Substrate heterogeneity promotes cancer cell dissemination

Zuzana Dunajova[1*], Saren Tasciyan[1*], Juraj Majek[1], Jack Merrin[1], Erik Sahai[2], Michael Sixt[1#], Edouard Hannezo[1#]

[1] Institute of Science and Technology Austria (IST Austria), am Campus 1, 3400 Klosterneuburg, Austria
[2] Tumour Cell Biology Laboratory, The Francis Crick Institute, London, UK

[*]Contributed equally, [#]contributed equally
[#]Correspondence: michael.sixt@ist.ac.at, edouard.hannezo@ist.ac.at

Abstract

While tumor malignancy has been extensively studied under the prism of genetic and epigenetic heterogeneity, tumor cell states also critically depend on reciprocal interactions with the microenvironment. This raises the hitherto untested possibility that heterogeneity of the untransformed tumor stroma can actively fuel malignant progression. As biological heterogeneity is inherently difficult to define and control, we here chose a highly reductionist bottom-up approach and let tumor cells invade into micro-engineered environments harboring obstacles with precision-controlled geometry. We find that not only the presence of obstacles, but more surprisingly their spatial disorder, causes a drastic shift from a collective to a single-cell mode of invasion. Combining live-imaging and perturbation experiments with minimal biophysical modeling, we demonstrate that cellular detachments result both from local geometrical constraints and a global temporal integration of the heterogeneity experienced during the invasion process. We show that different types of microenvironments map onto different universality classes of invasion dynamics, which can quantitatively explain our findings. We thus identify generic physical principles for how the mode of cancer cell invasion depends on environmental heterogeneity, with implications to understand tumor evolution *in vivo*.

Introduction

Cellular heterogeneity is at the core of tumor evolution. The canonical view suggests that new genetic variants produce new cancer cell phenotypes that physically invade extra-tumoral micro-milieus. If these cells experience more favorable growth conditions, they clonally expand and breed new heterogeneity. In an iterative process, the tumor ultimately reaches the fatal state of systemic dissemination[1–3]. While heritable genetic variations have long been recognized to play a crucial role, phenotypic variation can also be of epigenetic nature or even result from transient signaling states[4,5]. In any case, neither variation nor selection of cancer cells are autonomous to the tumor, but follow the metazoan principle that regulation occurs in a reciprocal crosstalk with the (untransformed) microenvironment [6–8]. Remarkably, in some tumor models even invasiveness itself is an active and transient response of rare cancer cells to signals released by the benign microenvironment[9–11]. This raises the possibility that more heterogeneous extra-tumoral microen-vironments act as drivers of malignancy[12], although this remains largely untested due to the diffi-culty of uncoupling substrate heterogeneity and malignancy *in vivo*.

In recent years, more reductionist approaches have gained increasing attention, where the behavior of cells emerges from first physical principles that do not even require genetic or epigenetic changes. For instance, bottom-up reconstitution of microenvironments have been suc-cessfully used to model cellular responses[13] and cancer dissemination in response to specific local topographies of the extracellular matrix (ECM)[14,15]. Cellular collectives can also shift between fluid and solid states as a simple function of cell-cell and cell-ECM adhesion, cell density or active migratory forces[16–22], with fluidized states favoring invasion[23–27]. Interestingly, such fluidized state can create large mechanical stresses on nuclei, which triggers mechano-adaptive responses and can thus feedback on cell malignancy to further promote invasiveness[28]. Finally, interface-dynamics of spreading tumor cell colonies display highly generic fractal-like features, that have been proposed to follow generic models for growing interfaces, such as the Kardar-Parisi-Zhang (KPZ) equation[29,30] or the Molecular Beam Epitaxy (MBE) equation[31], that have been validated in a number of non-living matter settings[32,33]. However, the above studies have all looked at migra-tion either in homogeneous or ordered environment, so that the influence of heterogeneity of the extracellular environment remains unexplored.

Here, we pursued a reductionist microengineering-based approach to test the response of tumor cells to ordered vs. disordered environments. We chose to specifically manipulate envi-ronmental geometry, while leaving all other aspects constant, which already demonstrated com-plex impacts on interfacial growth and tumor dispersion. In particular, we find that cellular detach-ments are both impacted by local features of substrate heterogeneity, as well as a much more

global effect of temporal integration of the heterogeneity experienced by the tumor front. This led

us to derive a quantitative description of the determinants of cellular dispersion, which we tested

by experiments in different environments with structured geometries.

Results

Environmental mechanics and heterogeneity are sufficient to drive dispersion *in vitro*

As a cellular model for tumor invasion, we employed A431 human carcinoma cells[34]. This

cell line is derived from a primary tumor of epithelial origin and forms cell-cell adhesions due to

the expression of E- and P-Cadherin[35]. When plated on fibronectin-coated 2D surfaces A431 cells

spread radially with a coherent cell front showing no cellular detachments from the collective

(Supplementary Fig. 1a). To test invasion into geometrically precisely controlled environments,

we engineered microfluidic devices where a centrally applied pool of cells spread into a fibron-

ectin-coated confined space of 9μm height. The two confining surfaces were intersected with

arbitrarily positionable 9μm diameter pillars, creating a 3D chamber that forces cells to invade as

a monolayer into the "pillar forest". The setup allowed free medium exchange, so that cells were

viable and could be imaged and tracked for more than a week using genetically encoded fluores-

cent reporters for cytoplasm (mCherry) and nuclei (GFP). We created microfluidic chips where

cells radially spread into 4 different sectors, allowing assessment of different pillar patterns under

otherwise identical conditions. To test the impact of ordered vs. disordered geometries we placed

pillars on grid-like patterns, creating regularly spaced obstacles that cells encounter while spread-

ing from the center to the periphery (Fig. 1a). We chose 9μm pillar spacing, where cells could

pass by only moderately deforming their ~12-18μm diameter nucleus. For disordered patterns we

kept the average pillar distance constant, but randomly shifted individual distances according to

a Laplace distribution (Fig. 1b). Tracking invasion over 8 days showed that cells invaded ordered

patterns with occasional detachments. Strikingly, in disordered patterns cells advanced with en-

hanced single cell detachments at the invasion front (Fig. 1c, Supplementary Movie 1). This was

in stark contrast to experiments on obstacle-free 2D surfaces, where A431 cells migrated in a

highly cohesive manner in the absence of any detachment (Fig. 1d, Supplementary Movie 2). To

quantify the magnitude of the dispersion effect in disordered patterns, we used A431 cells that

were genetically engineered to lack E- and P- cadherin (EPCad[-/-]). In collagen assays and pillar

patterns EPCad[-/-] tended to invade as single cells as they moved away from the bulk, irrespective

of environment geometry (Supplementary Fig. 1b-c, Supplementary Movie 3). A side-to-side com-

parison between wild type (WT) and EPCad[-/-] cells revealed that in disordered environments WT

cells showed cellular detachments comparable to EPCad$^{-/-}$ cells (Supplementary Fig. 1d). To-
gether, these data demonstrate that A431 cells undergo a drastic shift from collective to single
cell invasion when exposed to disordered geometries, with substrate heterogeneity having an
effect comparable to genetically inflicted loss of cell-cell adhesion.

Minimal biophysical simulations recapitulate the effect of environmental heterogeneity

Several mechanisms might explain the effect of environmental heterogeneity on single
cell detachments. Firstly, we reasoned that disordered pillar patterns contain regions of very nar-
row pore sizes, that might cause detachments by physically restricting the interfacial area of cell-
cell contacts. Such very small pores could also mechanically compress the nucleus, which can
trigger an active cellular response that might propel the cell forward, causing detachment from
the collective[36–41]. To test this 'pore size' hypothesis, we repeated the assay in ordered pillar pat-
terns with smaller spacing (6µm), requiring the nucleus to severely squeeze between obstacles.
We found that cells still moved collectively without increased detachment, arguing against a sim-
ple effect of pore size (Supplementary Fig. 1d). Next, we reasoned that gradients in pore sizes,
present only in disordered environments, might create detachments, as narrow constrictions fol-
lowed by free space might be permissive for cells at the leading edge to detach and move away
from the tissue bulk, which would stall in the constriction ('funnel' hypothesis). We tested this
option using two graded, yet ordered, pillar pattern designs, which change the pore size only in
the direction of cell invasion; either gradually (sinusoid design) or abruptly (stepwise design) (see
Supplementary Fig. 1e). Importantly, neither of these pore size variations triggered a significant
increase in the number of detachments (Supplementary Fig. 1f). Together, these data show that
cellular detachments do not simply arise from a single specific local geometry of obstacles. This
led us to explore other, more collective hypotheses, where detachments could result from a com-
plex series of transitions that the tissue experiences while passing through a disordered environ-
ment.

Fig. 1: *In vitro* experiments reveal a key role for substrate heterogeneity in driving tumor dispersal. **a.** Experimental setup of the microfluidic invasion assay, where surfaces are coated with fibronectin and filled with cell culture media before A431 cells are seeded into the center of the devices **b**. Illustration of the pillar patterns, where cells from the center invading into. **c.** A431 cell invasion imaged with time-lapse fluorescence microscopy over the days 3, 4 and 5.5 from left to right, where cells express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green). The segmentation of the cell nuclei in day 4 is shown on the right. **d.** Scratch assays performed as a control to compare the migration of A431 cells in homogeneous environment. **e.** Number of detached cells normalized by the radial interface length in the respective pillar patterns for A431 WT. Scale bars: 100 µm, box and whiskers plot displays median, 1st and 3rd quartiles (hinges) and bars display 1.5 interquartile range from the hinges, n=16 experiments for ordered and n=18 per disordered environment, followed over 8 days.

For this, we turned to minimal *in silico* simulations of invasion in complex environments.

Cells were modeled as soft adhesive and self-propelled particles[42–45], and we assumed repulsive

interactions with static pillars forming patterns of different geometries (Fig. 2a). Before seeking to

quantitatively fit the model parameters, we first performed a broad parameter screen, and found

that these minimal simulations robustly reproduced key qualitative features of the data, including

the fact that the number of detachments drastically increased with the presence of pillars, as well

as with pillar heterogeneity (Supplementary Fig. 2a). We then proceeded to constrain model pa-

rameters in a more precise manner, to understand quantitatively what causes detachments in the

simulations. As detailed in the Methods (see also Supplementary Table 1 for a summary of the

parameter fitting), parameters related to single-cell migration properties (e.g. migration speed and

bias, persistence time) were inferred based on the measured behavior of single detached cells in

pillar patterns. Cell-cell adhesion strength was constrained based on the lack of cell detachment

in homogeneous environment and infrequent cell-cell rearrangements, arguing for a glass-like

rheology with slow re-arrangements (Supplementary Fig. 2b, Supplementary Movie 4). Simula-

tions ran within this parameter regime confirmed the increase of detachments induced by pillar

heterogeneity (Fig. 2b, Supplementary Movie 5), as well as this effect diminishing when simulating

cells with lower cell-cell adhesion (Supplementary Fig. 2c,d, Supplementary Movie 6), similar to

the experiment (Fig. 1e).

With this parametrization, we started by running many repetitions of simulations with a

specific disordered pattern and calculating a probability distribution for detachment as a function

of position within the pillar pattern (the analysis was based on clustering with instantaneous cell

positions and tracks both transient and long-term detachments). Interestingly, the most striking

correlation which we found was not with local position, but instead with time since the start of

invasion. Indeed, this analysis revealed that the probability of detachments consistently increased

as the cells invaded in pillar patterns, with much steeper temporal increase in disordered patterns

(Fig. 2c).

Fig. 2: Minimal particle-based simulations reveal qualitatively different dynamics of interfacial growth and detachments in different environments. a. Simulation setup of cells (modeled in grey as soft attractive particles) invading a pillar pattern (modelled in black as soft repulsive particles), with schematic description of the relevant forces and interaction potentials. **b.** Comparison of the number of detached cells (at day 6) in simulations with patterns of different levels of disorder (heterogeneity values represent parameters of Laplace distribution, see Supplementary Table 1). Bars indicate mean number of detached cells and error bars show standard error (left bar - pore size 0.92 *L*, right bar pore size 1.06 *L*). **c.** Rate of detachments are increasing in time in simulations, with steeper increase with disordered patterns. **d.** Representative snapshots at different time points of the cell interface (magenta) in simulations for different environment geometries, side by side with the cell body segmentation from corresponding experiments (see Figure 1c,d). The interface gets rougher in time and in more heterogeneous environments. Cells are shown as grey and pillars as black beads, detached cells are highlighted in red. **e.** Interface roughness in different environment geometries in simulation, showing different exponents with and

without pillars, with the steepest increase for disordered patterns ($\beta_{homogeneous}$ = 0.33, $\beta_{ordered}$ = 0.46, β_{disor-}

$_{dered}$= 0.53). The markers represent the mean values, shaded area indicates the standard error and solid

line the linear fit. **f.** Experimental quantification of interfacial roughness over time and for different condi-

tions ($\beta_{homogeneous}$= 0.28, $\beta_{ordered}$ = 0.55, $\beta_{disordered}$ = 0.64), showing a good agreement with the theoretical

predictions. **g.** Correlation of interface width (day N) and detachment rate (day N+1) in all analyzed exper-

iments, showing strong overall correlation over different samples and conditions ($n_{ordered}$ = 16; n_{disor-}

$_{dered}$=18, $n_{homogeneous}$ = 6).

Substrate heterogeneity drives gradual roughening of the tissue interface

From a physical perspective, a classical result on the invasion dynamics of a fluid material

such as a tissue is that its interface should become rougher and rougher over time and that this

temporal increase follows a highly universal trend[29,46–49], which depends only on a few key char-

acteristics of the system. This led us to the idea that cell detachments could be connected to the

properties of the cell interface, and thus were sensitive to "time-integrated" heterogeneity, with

each pillar passed by the invading front increasing surface roughness by a small amount. To test

this, we computed in simulations both the overall perimeter of the front, which increases continu-

ously during cell invasion (Supplementary Fig. 3a), as well as the surface roughness (i.e. devia-

tions in the interface height with respect to its average, Fig. 2d,e). Interestingly, we found that the

interface roughness always increased in time, but with different growth exponents depending on

the nature of the geometrical environment. Fitting the increase to a power-law to get the growth

exponent β (following conventions from statistical physics, see Methods), we found that cells mi-

grating through homogeneous environments followed a roughness increase exponent of β=0.33.

This is what is expected from KPZ dynamics, a highly universal model of interfacial growth. Strik-

ingly, interfacial roughness in disordered patterns increased significantly faster, with a much larger

exponent of β=0.5-0.6. Although initially surprising, we reasoned that this was very close to pre-

vious theoretical predictions for the so-called KPZq universality class[50–53], which describes grow-

ing interfaces in the presence of quenched disorder (Fig. 2e). Computing these exponents in our

simulations for either homogeneous or disordered environments confirmed that we predict these

exponents in a wide range of parameter regimes (Supplementary Table 2), despite the fact that

our system is more complex than previous continuum theories, for instance due to the presence

of particle adhesion and active self-propulsion.

Strikingly, computing surface roughness and exponent β in the same manner in experi-

mental data revealed very good agreement with the data (best fit of 0.28 for homogeneous and

0.64 for disordered environments). Furthermore, in simulations of ordered pillar patterns, we

found that the predicted growth exponent depends on the average pore size as well as orientation

of the pattern relative to the direction of cell invasion (Supplementary Table 2, Supplementary

Fig. 3b,c), but is mostly found to be intermediate between homogeneous and disordered, with

systematically lower interface width compared to disordered environments. This was verified in

the experimental data, where we found a roughness exponent of 0.55 for ordered environments

with an interface width that was on average 21% lower in ordered compared to disordered envi-

ronments. Once we could rationalize the dynamics of interfacial growth in different environmental

conditions within this simple general model, we asked whether this could explain the different

trends of temporal increase in detachments across conditions. For this, we correlated at each

time point, for a given experimental replicate, the interface roughness at day N with the number

of detachments at day N+1. Strikingly, we found a strong correlation (R^2=0.65) across all data

points and different conditions, arguing that there might be a causal link between interfacial rough-

ening and cell detachments (Fig. 2f). To test further this theory, we went back to aforementioned

geometries which did not trigger detachments, such as ordered patterns with small pore size and

sinusoid or stepwise designs (Supplementary Fig. 3d, Supplementary Movie 7). Running simula-

tions in the same parameter regime, but corresponding pillar geometries, demonstrated good

agreement with experimental data, as we could predict both lower interfacial roughness and lower

number of detachments in all of these conditions compared to random pillars (Supplementary Fig.

3e).

Fig. 3: Predicting detachment dynamics from interfacial roughness across conditions. **a.** Details of the interface shape (magenta) analyzed in simulations. The cells are shown as grey and pillar as black beads, tips of interface protrusions are highlighted in orange, detached cells in red. **b.** Illustration of the measurements of position of detaching cell relative to interface shape, where the distance from the protrusion baseline and tip of protrusion is normalized. The measurement is performed in the last frame before detachment. Cell bodies were expressing cytoplasmic mCherry (blue) and nuclear H2B-GFP (green). Scale bar, 100µm. **c**. Relative position of the detachments compared to the finger baseline (0: at local minimum, 1: at the tip of the fingers). Middle line, upper and lower lines indicate the median, first and third quartiles, respectively (collection of n=100 detaching cells from n=7 experiments and n>700 from n=40 simulations for each condition). **d.** Local curvature of the interface in homogeneous, ordered and disordered environments, and local curvature of interface at detachment positions (n=6 experiments with homogeneous, n=16 for ordered and n=18 for disordered environment, n=40 simulations for each condition). Pillar patterns, especially the disordered pattern exhibit higher pool of interface positions with curvature high enough for the cells to detach. **e.** Aspect ratio of the interface protrusions (angle between tip of the

interface and "shoulders", ie. next local minima) in experiments. Disordered environment generates pat-

tern with higher number of long, narrow fingers. Same dataset was used as in d). **f-h.** Statistics of the dis-

tribution of interfacial protrusions (g) and detachments (h) across n=10 simulations with the same disor-

dered pillar pattern (f), confirming that the location of interface protrusion is highly predictive of detach-

ments (R^2=0.6).

Quantitative local and global determinants of cellular detachments

To follow up on this finding, we sought to study more precisely how a global property such

as interface roughness is linked to a local property such as cell detachments. We hypothesized

that a rougher invasion front leads to formation of interface regions with higher curvature and

fewer cell contacts, which might facilitate detachments[54–56]. Indeed, when we analyzed individual

detachment events in simulation and experiment, we found that the cells most likely to detach

were located at the tip of convex "finger-like" protrusions of the interface (Fig. 3a-c). Accordingly,

we analyzed local curvature of the interface shortly before the detachment (which negatively cor-

relates with the local number of cellular neighbors) (Fig. 3d), and we found that the majority of

cells detached from interface positions above a threshold curvature value. Interestingly, cells mi-

grating in pillar-free environments never reached such high interfacial curvature values, providing

a quantitative explanation for their lack of detachment. While there was no striking difference

between the curvature distribution of ordered and disordered patterns in experiment, which could

be due to the difficulty of segmenting extremely rough interfaces (Supplementary Fig. S4a), we

found that the aspect ratio of the fingers in disordered patterns was significantly higher than in

ordered patterns, and increased towards later time points (Fig. 3e, Supplementary Fig. 4b), which

again supports a close relationship between detachments and local interfacial curvature created

by the heterogeneous environment (Fig. 3f-h, Supplementary Fig. 4c). Overall, our findings sup-

port the idea that invasion into pillar patterns is a cumulative phenomenon. It results in a gradual

increase of interfacial roughness over time, with different exponents and slopes depending on

environmental geometry. The existence of a critical roughness which is permissive for detach-

ments explains that detachments do not occur at early time points in the presence of pillars and

never appear in the absence of pillars (at least within our experimental time).

To test this further via independent experiments, we designed new pillar geometries, char-

acterized by sinusoid patterns perpendicular to the direction of invasion. By making the sinusoidal

modulation relatively mild, we ensured invasion across the entire pillar pattern, but with a con-

trolled curvature modulation (Fig. 4a, Supplementary Movie 8). As predicted by the simulations,

this geometry promoted formation of periodic finger-like protrusions in the regions of maximal

pore width (Fig. 4b), as well as higher detachment probability with increasing frequency of sinus-
oid modulation, resulting in sharper fingers (Supplementary Fig. 4d). In both simulations and ex-
periments, the average probability of detachments scaled with the local interface shape (Fig. 4b-
c, e-f).

Finally, we asked whether there were additional local geometrical predictors for detach-
ment, on top of such global integrated heterogeneity. Indeed, as discussed above, simulations
repeated over the same disordered geometry revealed locations where more detachments oc-
curred. Given that we found that sinusoidal patterns parallel to the direction of invasion did not
yield detachments (Supplementary Fig. 1d-e, 3c-d), whereas sinusoidal patterns perpendicular to
the direction of invasion did (Fig. 4a-c), we specifically computed different metrics for local pillar
anisotropies in the front, back or sides of the invasion direction (Fig. 4g, Supplementary Fig. 4e-
f). Interestingly, we found in simulations that regions with locally anisotropic free space (i.e. less
pillars in the direction parallel to invasion compared to perpendicular) displayed a small but sig-
nificant correlation with detachments (R^2=0.1). By combining this metric of local pillar anisotropy
together with the long-term trend of roughness-induced increases in detachment, we could
achieve good predictive power for the spatiotemporal statistics of detachments in simulations
(R^2=0.64, Supplementary Fig. 4e). To confirm this, we followed individual cell trajectories in sim-
ulations, and found that increased pillar anisotropy was observed specifically at the time of de-
tachments (Fig. 4h). To test this experimentally, we computed the local anisotropy of pillar geom-
etry across n=94 cellular trajectories with detachments, and found indeed a statistically larger
number of pillars perpendicular compared to the parallel direction of movement at the time of
detachment (Fig. 4i). Together, our data show that a geometrically disordered environment pro-
moted detachments primarily via globally increased interfacial roughening, with finger-like regions
of high curvature, but that, in addition, local substrate anisotropy tunes the likelihood of cell de-
tachment.

Fig. 4: Predicting the statistics of detachments as a function of global and local heterogeneity. **a.** Simulation and corresponding experiments of cells in longitudinal sinusoid pillar pattern featuring only lateral heterogeneity of pore size. Cells invade with forming regularly spaced finger-like protrusions. Top panel: simulation at day 6, cells are represented as grey and pillar as black beads, finger maxima shown in orange, cell interface in magenta, detached cells in red. Bottom panel: experiment, cell bodies were expressing cytoplasmic mCherry (blue) and nuclear H2B-GFP (green). **b.** Interface profile in simulations (purple line, shaded area indicates standard error, n=10 simulations), averaged over all timepoints and multiple periods of the sinusoid pore size profile, showing that interface position correlates strongly with regions of low pillar densities. **c.** Instantaneous detachment positions in simulations collected from all timepoints and periods of sinusoid pore size profile. **d.** Interface profile in experiment averaged over all timepoints in spherical coordinates (shaded area indicates standard error, n=4 experiments). **e-f.** Averaged interface profile

(e, purple line, shaded area indicates standard error) and position of recently detached cells (f, red bars), which match well the prediction of simulations (see resp. panels b and c), n=4 experiments. **g.** Schematics for how local features of pillar geometry are quantified (see Methods). At each point in space, nearby pillars are divided into 4 quadrants – front, back, left and right, and we compute different statistics for how the pillar density in each quadrant correlates with detachments. **h.** Local pillar geometry along the trajectory of detaching cells in simulations (time=0 at detachment, n > 10000 detachment events from 40 simulations). As the cell is going to detach, the number of pillars in front of the cell decreases and increases on the sides of the cell (leading to detachments in locally anisotropic regions). **i.** Experimental number of pillars at the time point of detachment in different quadrants (dot plot and error bars display mean and standard error from the mean, collection of n=68 detaching cells from n=7 experiments, P 0.0045 (Back vs. Left/Right), 0.018 (Front vs. Left/Right) and 0.74 (Back vs. Front), Wilcoxon test), showing a statistically significant increase in side pillars (increased local anisotropic), as predicted experimentally.

Discussion

Our results demonstrate the importance of environmental heterogeneity in the dissemination of cancer cells and illustrate the crucial role of invasion interface shape in this process, an intrinsically collective tissue property. We find that substrate heterogeneity drives cell detachments from the bulk in a non-trivial way, acting via two complementary mechanisms. As cells go through a disordered environment, the free invasion interface gets gradually rougher, effectively integrating heterogeneity over time. We expect that this is a highly generic effect, as we show that we can map the dynamics of interfacial roughening to a generic universality class of active physical processes (KPZq). After a critical amount of time, the interface roughness is large enough that cells at the tip of invading fingers have few enough neighbors to detach. This permissive, integrated effect is then compounded by the local substrate geometry, which selects specific sites of local anisotropy where detachments are specifically promoted.

Although we have investigated this in an *in vitro* setting, where environmental geometry could be finely tuned, a next step would be to quantitatively compare our findings to more realistic *in vivo* situations[57]. For instance, different organs have different organization of extracellular matrix[58], which seem to sit between fully ordered and fully disordered states, although this has not been systematically quantified. Generalizing our simulations to tumor invasion in a 3D fibrous environment could lead to key insights to understand the influence of tissue-environment crosstalk in invasion dynamics. In line with that idea, quantitative imaging of human tumor samples revealed that spatially disordered or tilted stromal compartments can be negative prognostic factors for clinical outcome[59,60]. Another natural extension of the model would be to consider remodelling of the environment due to mechanical forces exerted by cell migration[61].

Using an experimentally controlled bottom-up approach, we introduce stromal heteroge-

neity as a potential driver of malignant progression. In analogy to organismal evolution, where

more heterogenous habitats drive genotypic diversity, a more heterogenous tumor stroma might

shift tumor cells into novel phenotypic states (in our case, the single cell as opposed to the col-

lective state). Despite being transient phenotypes without heritable traits, these states might later

become consolidated[28] by epigenetic and ultimately genetic accommodation.

Acknowledgments

The authors thank all the members of the Hannezo and Sixt groups for fruitful discussions and

Takuya Kato for advices on the experimental set-up. We are also thankful for support from the

Scientific Service Units (SSU) of IST Austria through resources provided by the Imaging and

Optics Facility (IOF) and the Lab Support Facility (LSF). Z.D. has received funding from the

Doctoral Programme of the Austrian Academy of Sciences (OeAW): grant agreement 26360.

This work was supported by the European Research Council (grant ERC-SyG 101071793 to

M.S).

Author contributions

S.T., Z.D., M.S., and E.H. designed the research. S.T. performed the experiments: S.T., Z. D.

and Ju.M. performed data and image analysis; Z.D. and Ju. M. performed simulations; Ja.M.

supported microfluidic engineering; E.S. provided cellular model systems. M.S. and E.H. super-

vised the study; S. T., Z.D., M.S. and E.H. wrote the manuscript, with input from all authors.

Methods

Experiments

Cell Culture

A431 cells expressing mCherry were grown in DMEM (Gibco, 10569010) containing phenol red,

high glucose, GlutaMAX, pyruvate, 10% fetal calf serum (FBS, Gibco, 10270106), non-essential

aminoacids (Gibco, 11140050), penicillin (50U/ml) and streptomycin (50µg/ml, Gibco 15070063)

in humidified incubators at 37°C with 5% CO_2. EPCad$^{-/-}$ A431 cells were made as described in

Labernadie et al. 2017[62]. Cells were regularly tested for mycoplasma contamination and pre-

served in growth medium containing 5% (v/v) DMSO in cryogenic conditions for longer periods.

Cells were transfected with lentivirus containing H2B-GFP construct and selected for blasticidin

resistance and then later sorted for GFP signal, regularly.

Device Fabrication

Patterns were generated with Python script in SVG format. Designs of photomasks were com-

pleted in (Inkscape, v0.92) in layers for each confinement. SVG files were converted to AutoCAD

DXF R14 files, further converted to Gerber files (LinkCAD) and finally printed on chrome photo-

masks at a resolution of 1 µm (JD Photo Data & Photo Tools). For high-resolution structures,

wafers were first coated with a sub-micrometre layer of SU-8 as a sticky substrate for later SU-8

structures. We used GM-1040 SU-8 photoresist (Gersteltec) at 5,000 rpm for 2 min with flood UV

exposure. The feature layer was then applied with SU8-2005 (Microchem). According to the

datasheet, a UV-absorbing glass plate and higher exposure were used to make sharp features.

PL-360-LP from Omega Optical and 550 mJ/cm^2 with an EVG mercury lampbased mask aligner

gave good results in the 3–9-µm-thickness range. The thick (25 µm) loading channels were made

with Microchem SU8-3025 using the EVG mask aligner according to the Microchem data sheet

without the PL-360-LP plate. The master wafers underwent vapour silanization for 1 h with (1H,

1H, 2H, 2H-perfluorooctyl) trichlorosilane. The final results were single hydrophobic master wa-

fers containing several aligned layers. Polydimethylsiloxane (PDMS) devices were made with Syl-

gard 184 Elastomer Kit (Dow Corning) using a ratio of 10:1. The PDMS was mixed in a Thinky

mixing machine and then poured on the wafer in a 3D printed tray and further degassed several

times in a vacuum chamber. The devices were baked at 80 °C overnight. The hardened devices

were carefully peeled off the mould and cut into single devices using a razor blade. Holes were

punched with a custom arbor press or with skin biopsy puncture tools (1 mm) for the central hole,

and sticky tape was used to remove debris. Glass coverslips were sonicated for 5 min in 100%

isopropanol and blown dry. Devices were bonded to the glass slides by oxygen plasma then

sealed together at 85 °C for 1 h to make the bond permanent.

Invasion Assays

Microfabricated devices were inserted into 12-well cell culture plates from the bottom side and

glued with molten paraffin. Devices were filled with fibronectin (F1141, Sigma) solution (diluted in

1:5 from 1mg/ml stock solution in PBS) using a syringe pump, sealed and incubated for 3h in cell

culture growth conditions. Devices were later flushed with 100 µl cell culture growth medium with

syringe pumps. Debris were removed from the center hole, either spheroids of cancer cells (gen-

erated with hanging drops) were dropped or a pipette tip filled with medium was inserted into the

hole, where 1.5 µl of the cell suspension was filled. Cells were incubated until they sank into the

hole and imaged at later days. Wells were checked for contamination under microscope.

2D Spread Assay

A431 cells were mixed with cytoplasmic mCherry reporter expressing A431 cells (10:1) and as-

sembled into 1000 cell spheroids in hanging drops overnight. Glass cover slips were coated with

fibronectin (F1141, Sigma, diluted in 1:40 from 1mg/ml stock solution in PBS) for 1h and then

washed with PBS. Subsequently, chambers were filled with cell culture media and spheroids were

placed to adhere on the substrate. Cells were observed with fluorescence and DIC microscopy.

Scratch Assay

In order to observe invasion on non-confined 2D surface with an initial straight interface, cells in

suspension were placed and grown to full confluence on fibronectin coated glass substrates (see

2D spread assay above). Then, approximately at the center of the cover slip, cells were scratched

with a rubber cell scraped and detached cells were aspirated and washed away with cell culture

media. Cells were observed with fluorescence and DIC microscopy.

Imaging

Cells were either imaged once a day at room conditions or in cell culture conditions for time-lapse

imaging (10 min interval) for several days (typically 3-4 days). Invasion was recorded with an

inverted wide-field Nikon Eclipse Ti-2 microscope equipped with a Lumencor light source (475

nm for GFP and 594 nm for mCherry) and an incubation chamber, heated stage and CO_2 mixer

(Ibidi). Generally, the fluorescence imaging was accompanied with light microscopy (DIC) imag-

ing. For long term imaging, phenol red color was checked and the temperature was monitored

with a temperature sensor. Custom made CNC milled box and a pair 3D printed plugs were used

to prevent CO_2 and humidity leak.

Collagen Assay

A collagen mixture was created with a final collagen concentration of 1.7 mg/ml by mixing bovine

collagen (PureCol, Advanced BioMatrix) in 1x minimum essential Eagle (Sigma) and 0.4% sodium

bicarbonate solution (Sigma Aldrich). Mixture was checked for a pale pink color to ensure for

correct pH for polymerization. 20 µl of the collagen mixture was mixed with 9 µl of the growth

medium and mixed gently. Subsequently, 1 µl of the spheroid containing growth medium was

added to the mixture and mixed by stirring with the tip. Final mixture was gently placed on a glass

coverslip, which was previously glued under a 60 mm dish with molten paraffin. The dish was
flipped to let the collagen drop hang from the glass coverslip, whereas the plastic lid was filled
with droplets of PBS to keep the dish humidified inside the incubator. Collagen drop was let to
polymerize in the incubator for 45-60 min under cell culture conditions. Before imaging, PBS was
removed and the dish was flipped again and medium was added to the chamber with the collagen
drops. Invasion into the collagen was imaged with light microscopy (DIC) over several days under
cell culture conditions.

Lentivirus Production

457 Lentivirus production was performed as described previously[63]. In brief, LX-293 HEK cells (Clon-
458 tech) were co-transfected with LentiCRISPRv1 or LentiCRISPRv2, packaging-ps-PAX2 (Addgene
459 no. 12260) and pCMV-VSV-G envelope plasmids using Lipofectamine 2000 (ThermoFisher Sci-
460 entific) as recommended by the manufacturer, and cells were resuspended in the growth medium
461 1 day before transfection. Supernatant was collected after 72 h and stored at −80 °C.

Image processing and experimental analysis

Image Analysis

FiJi was used for image processing, including using the TrackMate plugin for manual cell track-
ing[64,65]. Proprietary ND2 files from Nikon NIS Elements were converted using Bio-Formats plugin[66]
and the physically calibrated (pixel size and imaging interval). Images were aligned and pro-
cessed into TIFF and HDF files. Most of the analysis processes were automated with ImageJ
macros. Cell nuclei (H2B-GFP, green) and the remaining cell body (mCherry, blue) and nuclear
H2B-GFP (green) were initially segmented using Ilastik and then particle analysis feature was
used to quantify invasion[67]. Cells were tracked semi-automatically with TrackMate, tracks were
corrected manually. Dividing and dying cells were excluded, all gaps were filled with interpolation.
Final images were processed with automated scripts for display and exported into PNG. In im-
ages, cells were considered detached, when their cell bodies were not considered touching the
bulk. Finger measurements were performed manually by measuring the distance from the tip to
the finger baseline. Detachments were selected over a wide range of recorded experiments. The
pore size of the detaching cell during detachment vs. the pore size in front of the cell, which is
being detached from, was measured manually with an interactive macro.

Experimental data analysis

The cell interface in experiment was obtained by identifying the nuclei positions of cells at the

interface, for better comparison with simulations, which do not model the cell bodies (Supplemen-

tary Fig. 4a). Identification of cells at interface was done by first running Delaunay triangulation of

all cell nuclei in each timestep and subsequently removing the vertices, which cross through the

segmented interface of bulk cell bodies (blue line in Supplementary Fig. 4a). This way we could

efficiently find all cells, which are in contact with the very rough and irregular bulk interface. Before

using the interface positions for analyzing the roughness exponents, we had to remove the overall

tilts in the interface present in some experiments (see left panel in Fig. 1d, Supplementary Fig.

5a). This was done by removing long-wavelength Fourier modes (> 500 µm). The analysis of the

local pillar geometry around the detaching cells (Fig. 4i) was performed by identifying all close

pillar in a given radius around a the x,y position of the detaching cells and dividing them into 4

quadrants (front, back, left right, see Fig. 4g, the center of the front quadrant being the orientation

of the detaching cell). Multiple radii were tested, all resulting in qualitatively similar result (final

value: r_{max}= 27 µm, i.e three times the pore size). The identified pillars in each quadrant were

subsequently weighted by their distance from the reference position $(r_{max}-r)/r_{max}$ and summed.

Statistics and Data Visualization

All statistics for experimental data were performed in *R (4.0.3)* and plotted using *ggplot2* package

(R Core Team 2020; Wickham 2016:2). Tracking data was analyzed with *TraXpert*, an in-house

developed particle tracking analysis application. Directionality data was analyzed with *circular*

package and circular statistics[68,69]. Minor cosmetic adjustments were performed in *Inkscape*, with-

out affecting the data points or the interpretation of the plots. Figures were assembled in *Inkscape*

and exported into high resolution lossless PNG images.

Numerical model

We used two dimensional Brownian dynamics simulations as a minimalistic model of collective

cell migration[42–45] through complex environments (Fig. 2a). Each cell is represented by a soft self-

propelled particle (representing its nucleus[39]) on a substrate, so that the overdamped equation of

motion for each particle reads:

$$\gamma \frac{d\vec{r_i}}{dt} = \frac{dV_{pair}}{dr_{i,j}} + \vec{F_p}$$

507 where γ is the friction coefficient, $\vec{r_i}$ are the coordinates of the particles, V_{pair} is the pairwise inter-

508 action potential, and $r_{i,j}$ is the distance between particles. $\vec{F_p}$ is the active self-propulsion force

509 acting along the particle orientation $\vec{p_i} = (cos\theta_i, sin\theta_i)$ which in the absence of any interactions,

510 would result in an average cell speed of v_0, such that:

511
$$\vec{F_p} = \gamma v_0 \vec{p_i}$$

512 The polarity of migrating cells is affected by rotational diffusion and is additionally biased by an

513 external torque τ_{bias}, which represents the cues which drives cell migration away from the bulk

514 (see below for additional discussion of the source of this external cue):

515
$$\frac{\delta\theta_i}{\delta t} = \tau_{bias} + \sqrt{2D_r}\eta = -k_{bias}sin(\theta_i) + \sqrt{2D_r}\eta$$

Here, D_r is the rotational diffusion coefficient, η is a unit variance random variable, k_{bias} parameter

describing the strength of the biasing torque, where $\theta_i = 0$ represented the direction of the inva-

sion (right-hand side of all simulations shown here). To validate the numerical implementation,

we calculated the mean square displacement of a single active Brownian particle without the

biasing torque[70]. The detected crossover between ballistic and diffusive behavior was consistent

with the used value of D_r.

To describe cell-cell interactions, we assume a simple potential with short range repulsion and

middle range attraction[42,43,71] with the form

524
$$V_{pair} = \sum_i \sum_{j>i} v_{pair}(r_{i,j})$$

526
$$v_{pair}(r_{i,j}) = \begin{cases} 4\epsilon\left[\left(\frac{\sigma}{r_{i,j}}\right)^4 - \left(\frac{\sigma}{r_{i,j}}\right)^2\right] + \Delta v_{pair}(r_{i,j}), & r_{i,j} < r_{cut} \\ 0, & r_{i,j} \geq r_{cut} \end{cases}$$

where $r_{i,j}$ is the distance between the positions of particles i and j, ϵ is the depth of the potential

well, σ is the characteristic distance of interaction and r_{cut} is the cutoff distance of the potential.

The potential is shifted by a linear pair potential Δv_{pair}, to ensure that the potential and its derivative

smoothly goes to zero at the cutoff $\frac{dv_{pair}}{dr}(r_{cut}) = 0$.

Furthermore, to represent the complex environments that cells migrate in (pillar geometry), we

also positioned stationary particles into the simulation box, which contribute to the total pair po-

tential. The cell-pillar interaction is truncated at its minimum ($r_{cut} = \sqrt{2}\sigma$) to account only for repul-

sive interactions, given that we find no strong evidence of cell-pillar adhesion. An important as-
sumption of our minimal modeling approach is that cellular parameters are the same across con-
ditions, with differences only arising from substrate heterogeneity.

Details of the simulation setup

We used HOOMD-blue v3 library to run the numerical simulations[72]. The equations of motion
were numerically integrated using the Euler-Maruyama scheme with timestep $dt = 0.001\tau$. The
simulations are performed in two dimensions.

Simulation parameters and results are reported in dimensionless form. For convenient compari-
son to experiment the length is measured in units of cell-cell equilibrium distance L (defined as
the interaction minimum $1L = \sqrt{2}\sigma = 16$ um) and time $\tau = 10^3$ s. The simulation output is saved
every 600 frames, consistent with the smallest timestep in experiments (10 minutes).

Unless stated otherwise, the simulation parameters to model are summarized in the Supplemen-
tary Supplementary Table 1:

Supplementary Table 1: Summary of the used simulation parameters.

Parameter	Notation	Chosen value (simulation units)	Parameter fitting
Friction coefficient	γ	1	fixed parameter, non-dimensionalizes timescales
Attraction cutoff	r_{cut}	1.55 L (1 L for cell-pillar interaction)	fixed parameter (on the order of cell size, to avoid next neighbor interactions)
Self-propulsion speed	v_0	0.43 L/τ	fitted to correspond to mean single cell speed $6.9*10^{-3}$ μm/s inside the pillar pattern (Supplementary Fig. 5a,b)
Rotational diffusion	D_r	0.7 τ^{-1}	fitted according to MSAD of the experimental trajectories of detached cells (Supplementary Fig. 5c)
Bias torque	k_{bias}	0.75 rad/τ	fitted according to the drift of experimental trajectories of detached cells

			(Suplementary Fig. 5d)
Cell fluidity number	$\gamma v_0/\varepsilon$	0.52	constrained by confluency of cells, while keeping occasional cell rearrangements of bulk (Supplementary Fig. 2b, Supplementary Movie 4, Supplementary Table 3)
Pore size	L_p	0.88 -1.3 L	free parameter (on the order of cell size, lower bound is set that cells need to be able to pass the pores alone)
Pore heterogeneity	$\mathcal{H}$	$\mu = b = 0.1*L_p - 0.5*L_p$	defined by parameters of Laplace distribution, upper bound taken from disordered pattern in experiment

Before doing a systematic parameter search, we first constrained parameters to reasonable val-

ues and performed non-dimensional analysis to identify the key dimensionless ratios. The fluidity

of a tissue in the absence of any obstacles is governed by a dimensionless number relating self-

propulsion and cell-cell interaction strength ($v_0\gamma/\epsilon$)[44,73,74]. If the ratio is too low, cells behave as

solid with very few rearrangements of neighbors, while above a given value, the tissue fluidizes

as motility-driven re-arrangements become possible. However, when considering not only bulk

tissue, but also a cell interface, very large values of this ratio also cause spontaneous detach-

ments of cells in pillar-free environment as in a gas[75]. Importantly, experiments of monolayers

invading homogeneous environment support an intermediate value of the ratio, as we observed

virtually no detachments, but that cells could move relative to one another (Supplementary Fig.

1b, Supplementary Movie 4, Supplementary Table 3).

Furthermore, we can take advantage of the presence of single detached cells in experiments with

pillars to constrain a number of parameters related to single-cell migratory properties. In particular,

by calculating instantaneous (x,y) displacements and mean squared angular displacement

(MSAD)[76] of detached cells in pillar patterns in simulations and experiments, we could strongly

constrain the values of both self-propulsion speed and particle persistence length v_0/D_r (Supple-

mentary Fig. 5b-d). The value of external bias was fixed by analyzing the radial drift in the trajec-

tories of detached cells in experiment and compared to simulations (Supplementary Fig. 5e). Note

that we detected a strong biased migration even in single cells far from the leading edge, which

motivates the fact that this bias doesn't simply arise from pushing forces from bulk proliferation.

Given previous reports of self-generated gradients guiding long-range migration[77,78], such mech-
anisms would be compatible with this bias, which we chose to be agnostic about in the simula-
tions. Finally, we performed an additional sanity check by comparing mean squared displacement
(MSD) of single detached cells in simulation and experiment, allowing us to verify the chosen the
combination of parameters D_r, v_0, and k_{bias}. We achieved a reasonable fit, especially for long times
(Supplementary Fig. 5f).

The simulations of migrating cells used for comparative statistics in various pillar geometries and
parameters (Fig. 2b, Supplementary Fig. 2ab, Supplementary Fig. 5, Supplementary Table 2)
were initialized with a rectangular box of size $Lx=Ly=425$ L. The left part of the box was filled with
cells in hexagonal packing, while the right side contained the pillar geometry. Simulations used
for close temporal comparison with experiment (Fig. 2c-e, Fig. 3-4, Supplementary Fig. 3-4) were
initiated in a smaller box ($Lx=425$ L, $Ly=106$ L) with new cells being added on the left side of the
box during the simulation run to maintain a stable pressure in the bulk. Each combination of pa-
rameters was run with at least 10 repeats. The ordered pillar geometry was constructed as square
packing with pillar spacing being twice the effective pillar diameter. The reported results of ordered
pattern simulations (e. g. Fig. 2b,c,e and Fig. 3d) are averaged over 4 orientations of the square
spacing (0, 15, 30 and 45 degrees, see Supplementary Fig. 3b). The disordered pattern was
created from ordered patterns of all 4 tilts by shifting each pillar in a randomly chosen angle by
distance following Laplace distribution from the original positions (fully disordered pattern corre-
sponded to Laplace distribution parameters $\mu = b = 0.5 * L_p$).

Simulation analysis

The initial time needed for the cells to accumulate at the front of the pillar pattern and enter it was
discarded in the analysis. Unless stated otherwise, simulation frames were analyzed with fre-
quency 6 τ after the start of the invasion. The simulations were run until the bulk cells reached the
end of the pillar pattern. The simulation analysis was performed in Python.

The overall number of detached cells (Fig. 2b,d, Fig. 3a, Supplementary Fig. 2a,c, Supplementary
Fig. 3d) was analyzed by distance clustering of cells (r_{cut} was used as the natural clustering cutoff
from the physical point of view) and summing the number of cells inside of the pattern, which are
not connected to bulk in a given timepoint. The identification of the pattern locations with high
detachment probability and instantaneous detachments (Fig. 2c, Fig. 3c-d,f-g, Fig. 4c,h, Supple-
mentary Fig. 2d, Supplementary Fig. 3e, Supplementary Fig. 4c-e) was performed by first running

the same clustering analysis in each simulation frame. In the second step of the analysis, a cell

was considered to be detached if it was separated from the bulk (the biggest cluster) for at least

5 consecutive frames. For each detachment event we saved the x,y coordinates of detachment,

the cell ID, timestep to be used for further analysis (eg. analysis relating the detaching events to

interface properties in Fig. 3c,d, and local pillar positions Fig. 4d).

The shape of the cellular interface (eg. Fig. 2d,e, Fig. 3a, Fig. 4a) was analyzed by first computing

the Delaunay triangulation (convex hull) of cellular positions was in each simulation frame. The

Delaunay triangulation was then filtered using the alpha shape algorithm[79] (alphashape library in

Python) to determine the concave hull of the cellular interface (alpha = 0.6). The positions of

interface cells were subsequently used in the roughness scaling analysis (Fig. 2e, Supplementary

Fig. 3c,e) and their connectivity in the analysis of interface shape, perimeter and curvature (Fig.

3a,c,d,h, Fig. 4b, Supplementary Fig. 3a). The local maxima and minima of cell interface were

analyzed using the argrelextrema function in SciPy library[80], and the transient protrusions on the

order of cell size (protrusion length = average distance to next local minima < 1 L) were filtered

out.

The interface roughness (Fig. 2e, Supplementary Fig. S3c,e) was quantified using the height

function $h(y)$, i.e. the level of invasion of cells in different points of the interface (polar coordinates

were used in case of pillar pattern experiments). We analyze the increase of roughness in time

by computing interface width $w(l,t)$ measuring the root mean square deviations of an interface

about its mean value, defined as

629
$$w(l,t) = \left\{ \frac{1}{N} \sum_{i-1}^{N} \left[(h(y,t) - \bar{h}(t)) \right]^2 \right\}^{1/2}_{L_{int}}$$

where L_{int}=1000 μm is the length of the interface contour. These fluctuations grow in time in a

power law fashion $w(l,t) \sim t^{\beta}$, with a characteristic critical growth exponent β[31].

The analysis of local heterogeneity of pillars (Fig. 4h, Supplementary Fig. 4e,f) was performed by

identifying all close pillars around a given x,y position in disordered pattern (r_{max}= 3.18 L, i.e. 3

times average pore size) and dividing them into 4 quadrants (front, back, left right, Fig. 4g). The

identified pillars in each quadrant were subsequently weighted by their distance from the refer-

ence position (r_{max}-r)/r_{max} and summed.

Supplementary Figures

Supplementary Fig. 1: Invasion dynamics of cells in different environments a. 2D invasion assay of WT A431 cells with and without mCherry reporter with 1:10 mix respectively. **b.** Collagen invasion assay of WT and E-,P-Cad[-/-] spheroids imaged with brightfield microscopy. **c.** Detachments of the WT and E-,P-Cad[-/-] cells invading into ordered and disordered pillar patterns. White arrows indicate individual detached cells or cell clusters. **d.** Detached cell area in the respective pillar patterns for A431 WT and E-,P-CadKO cells. Scale bars: 100 μm, box and whiskers plot displays median, 1st and 3rd quartiles (hinges) and bars display 1.5 interquartile range from the hinges, n=3 experiments per genotype and pattern followed over 8 days. **e-f.** Cell invasion assays into smaller (6 μm) sized ordered pillar pattern (d) and step-

 wise or sinusoidal pattern (e). **g.** Quantified number of detachments for each pillar patterns, where box

and whiskers plot displays median, 1st and 3rd quartiles (hinges) and bars display 1.5 interquartile range

from the hinges, n=3 experiments per genotype and pattern followed over 8 days. All scale bars: 200µm.

Supplementary Figure 2: Phase diagram of invasion dynamics for different model parameters. a.

Temporal increase in detachment rate is a robust phenomenon present both in ordered and disordered

pillar patterns, as well as for different simulation parameters. The simulation snapshots show a representa-

tive state of the systems at day 6 (cells are shown as grey and pillar as black beads, detached cells are

highlighted in red), with the best-fit set of simulation parameters used in the main text (left column, see also

Supplementary Table 1). Each subsequent column shows the effect of changing a single simulation param-

eter (D_r = 1 τ^{-1}, D_r = 0.5 τ^{-1}, k_{bias} = 0.5 rad/τ, k_{bias} = 1 rad/τ, v_0 = 0.24 L/τ respectively). **b.** Snapshots of

simulation of cells in homogeneous environments (grey beads shown with smaller diameter for clarity, day

6), for different values of the fluidity number. High fluidity number (left) leads to numerous detachments,

something not observed in wild-type experiments, while low fluidity (right) leads to a solid-like rheology

without any re-arrangements on the time scale of the simulations (also not observed). **c.** Snapshots from

simulations of cells with lower cell-cell adhesion, corresponding to E-,P-Cad$^{-/-}$ cells (fluidity 0.87, cells are

shown as grey and pillar as black beads, detached cells are highlighted in red). This leads to detachments

even in the absence of any pillars, i.e. to a much weaker sensitivity to the environment, as seen experimen-

tally. **d.** Quantification for the increase of detachments when lowering the cell-cell adhesion, for either or-

dered or disordered pillar patterns, showing that the difference between ordered and disordered pattern

disappears compared to intermediate adhesion/fluidity.

Supplementary Figure 3: Sensitivity analysis of simulation outputs for different pillar patterns. a.
Temporal evolution of the interface perimeter, quantified in simulations (n=40 simulations, the shaded intervals indicate standard error). **b.** Differences in invasion behavior with different orientations of regular pattern (snapshots of day 4, cells are shown as grey and pillar as black beads, magenta line represents cell interface). We used the average results of the 4 shown orientations for comparing regular pattern with other geometries, given that invasion is radial in experiments. **c.** Interfacial roughness and roughness exponents in 4 orientations of regular patterns (corresponding to b) (n=10 simulations each, the shaded intervals correspond to standard error). **d.** Simulation snapshots of invasion in sinusoid and stepwise pillar patterns, exhibiting lower or comparable detachments, as in ordered pattern, as seen experimentally in Supplementary Fig 1e-g. Cells are represented as grey and pillars as black beads, detached cells are marked red, cell interface in magenta, day 4. **e.** Quantifications of interface width and detachment numbers in simulations of panel d, showing that it follows a stepwise trend corresponding to the pillar pattern, and that some cells tend to reattach to the bulk resulting in lower overall number of detachments.

Supplementary Figure 4: Detachments statistics in data and simulations. **a.** Analysis of interface roughness in experiments. The blue line represents the segmented outline of cell bodies and magenta line connects the cell nuclei along the interface, which are used for subsequent roughness and interface curvature analysis (see Methods). **b.** Difference between the finger angle (defined as the angle between the tip of finger and neighboring local minima, with small angles thus corresponding to sharp fingers, see Fig. 3e) of the protrusions in ordered and disordered patterns, which becomes more pronounced in later timepoints (n=16 experiments for regular and n=18 for disordered pattern). **c.** Distribution of detachment events in simulations with increasing disorder in pillar positions (color coded in red for increasing probability). **d.** Temporal increase of detachments in perpendicular sinusoid simulations with smaller wavelength (see Fig. 4a). **e.** Heat map of detachment positions compared to the local anisotropy of pillar positions in disordered pattern (Metric: Left + Right – Front, R^2 = 0.1) and local anisotropy of pillar positions multiplied by the mask, which takes into account the increase of detachments in the direction of invasion (R^2=0.63). detachments analyzed from n=10 simulations. **f.** Example of a different metric calculated using the same analysis (Back-Front), which exhibited lower spatial correlation with detachment positions.

Supplementary Figure 5: Fitting procedure for different model parameters. a) Correction of tilts in the detected experimental interface (blue line) by removing long Fourier modes (orange line). **b-c)** Fitting the active force v_0 based on displacements of single detached cells in a pillar patterns, respectively (experimental data shown in the first two columns of panel b, orange line in panel c represents the median of distribution of detached cells, n=91 trajectories from n=6 experiments). The displacements do not depend on D_r and grow linearly with v_0 (further columns in panel b show n=3 simulations of n=100 sparsely added single cells in ordered pillar patterns with pore size 1.06 L and different combinations of parameters). By

comparing the data to different values of v_0, we constrained v_0=0.43 L/τ as the best-fit parameter used in

the rest of the simulations unless explicitly mentioned **d)** Fitting of the rotational diffusion constant based

on MSAD of single detached cells in a regular pillar pattern (color coded according to value of v_0, black

dashed line represents MSAD of detached cells in experiment). As above, we compared the experimental

results from trajectories of detached cells to simulations with different rotational diffusion coefficients D_r.

We found that the best-fit occurred for 0.7 τ^{-1}, which we use for the rest of the paper unless explicitly

mentioned. **e)** Fitting of the k_{bias} based on the radial bias in trajectories of single detached cells in a regular

pillar pattern (same procedure as above). We found that k_{bias}=0.75 rad/τ reproduced best the data, which

we use for the rest of the paper unless explicitly mentioned. **f)** MSD comparing the detached cells in pillar

patterns in experiment and final simulation parameters D_r, v_0, and k_{bias}, chosen according to **b-e**.

Supplementary Table 2. Interface roughness of cells invading in ordered and disordered patterns in different parameter combinations (green – exponents from 0.25-0.44, blue – larger than 0.55).

Pore size	D_r [τ^{-1}]	Ordered pattern				Disordered pattern			
		fluidity				*fluidity*			
		0.44	0.52	0.60	0.76	0.44	0.52	0.60	0.76
0.88 *L*	0.5	0.27	0.33	0.41	0.50	0.55	0.70	0.70	0.57
	1	0.19	0.36	0.43	0.44	0.45	0.65	0.72	0.71
	2	0.37	0.25	0.23	0.37	0.44	0.36	0.39	0.49
1.06 *L*	0.5	0.48	0.57	0.62	0.59	0.54	0.62	0.78	0.57
	1	0.30	0.45	0.52	0.55	0.44	0.69	0.64	0.59
	2	0.34	0.23	0.28	0.42	0.35	0.32	0.40	0.43
1.24 *L*	0.5	0.32	0.32	0.38	0.38	0.57	0.66	0.80	0.49
	1	0.37	0.46	0.49	0.50	0.58	0.69	0.64	0.59
	2	0.33	0.27	0.31	0.41	0.56	0.33	0.40	0.43

Supplementary Table 3: Behavior of the cell bulk in homogeneous environment in simulations with different parameters (green – KPZ scaling, yellow – KPZ scaling with holes and detachments, red – jammed, collective movement; brown – jammed, no movement

Supplementary Movie Legends

Supplementary Movie 1: Invasion of A431 human skin carcinoma cells in ordered vs. disordered environments. Cells express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green).

Supplementary Movie 2: Invasion of A431 human skin carcinoma cells in free environments. Cells express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green).

Supplementary Movie 3: Invasion of EPCad[-/-] A431 human skin carcinoma cells in ordered vs. disordered environments. Cells express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green).

Supplementary Movie 4: Simulations of cells with different fluidity number (i.e. with varying cell-cell inter-
action strength) in homogeneous environment showing different level of cell rearrangements. Cells are
represented as grey beads.

Supplementary Movie 5: Simulations of cells migrating in different environments (homogeneous, ordered
and disordered. Cells are represented as grey and pillars as black beads.

Supplementary Movie 6: Simulations of EPCad$^{-/-}$ cells (lower cell-cell adhesion) invading different types
of environments. Simulations of cells migrating in different environments (homogeneous, ordered and dis-
ordered. Cells are represented as grey and pillars as black beads.

Supplementary Movie 7: Simulations of cell invasion in spatially structured environments which do not
increase detachment rate (regular with smaller pore, sinusoid, stepwise). Cells are represented as grey and
pillars as black beads.

Supplementary Movie 8: Simulation and experiment of cell invasion in sinusoid perpendicular pattern,
which creates localized interface protrusions and detachments. Cells in experiment express cytoplasmic
mCherry (blue) and nuclear H2B-GFP (green). Cells in simulations are represented as grey and pillars as
black beads.

References

1. Almendro, V., Marusyk, A. & Polyak, K. Cellular heterogeneity and molecular evolution in
cancer. *Annu. Rev. Pathol.* **8**, 277–302 (2013).

2. Altschuler, S. J. & Wu, L. F. Cellular Heterogeneity: Do Differences Make a Difference? *Cell*
141, 559–563 (2010).

3. McGranahan, N. & Swanton, C. Clonal Heterogeneity and Tumor Evolution: Past, Present,
and the Future. *Cell* **168**, 613–628 (2017).

4. Cheung, K. J. & Ewald, A. J. A collective route to metastasis: Seeding by tumor cell clusters.
Science **352**, 167–169 (2016).

5. Yang, R. *et al.* Distinct epigenetic features of tumor-reactive CD8+ T cells in colorectal can-
cer patients revealed by genome-wide DNA methylation analysis. *Genome Biol.* **21**, 2
(2019).

6. Ferguson, L. P., Diaz, E. & Reya, T. The Role of the Microenvironment and Immune System

in Regulating Stem Cell Fate in Cancer. *Trends Cancer* **7**, 624–634 (2021).

7. Friedl, P. & Alexander, S. Cancer Invasion and the Microenvironment: Plasticity and Reci-

procity. *Cell* **147**, 992–1009 (2011).

8. Spill, F., Reynolds, D. S., Kamm, R. D. & Zaman, M. H. Impact of the physical microenviron-

ment on tumor progression and metastasis. *Curr. Opin. Biotechnol.* **40**, 41–48 (2016).

9. Giampieri, S. *et al.* Localized and reversible TGFβ signalling switches breast cancer cells

from cohesive to single cell motility. *Nat. Cell Biol.* **11**, 1287–1296 (2009).

10. Brabletz, T. EMT and MET in Metastasis: Where Are the Cancer Stem Cells? *Cancer Cell*

22, 699–701 (2012).

11. Padmanaban, V. *et al.* E-cadherin is required for metastasis in multiple models of breast

cancer. *Nature* 1–6 (2019) doi:10.1038/s41586-019-1526-3.

12. Rezk, R. *et al.* Spatial heterogeneity of cell-matrix adhesive forces predicts human glioblas-

toma migration. *Neuro-Oncol. Adv.* **2**, vdaa081 (2020).

13. Flommersfeld, J., Stöberl, S., Shah, O., Rädler, J. O. & Broedersz, C. P. Geometry-Sensi-

tive Protrusion Growth Directs Confined Cell Migration. *Phys. Rev. Lett.* **132**, 098401

(2024).

14. Jouybar, M., Sleeboom, J. J. F., Vaezzadeh, E., Sahlgren, C. M. & den Toonder, J. M. J. An

in vitro model of cancer invasion with heterogeneous ECM created with droplet microfluid-

ics. *Front. Bioeng. Biotechnol.* **11**, (2023).

15. Wang, W., Law, R. A., Perez Ipiña, E., Konstantopoulos, K. & Camley, B. A. Confinement,

Jamming, and Adhesion in Cancer Cells Dissociating from a Collectively Invading Strand.

PRX Life **3**, 013012 (2025).

16. Bocanegra-Moreno, L., Singh, A., Hannezo, E., Zagorski, M. & Kicheva, A. Cell cycle dy-

namics controls fluidity of the developing mouse neuroepithelium. 2022.01.20.477048 Pre-

print at https://doi.org/10.1101/2022.01.20.477048 (2022).

17. Mongera, A. *et al.* A fluid-to-solid jamming transition underlies vertebrate body axis elonga-

tion. *Nature* (2018) doi:10.1038/s41586-018-0479-2.

18. Petridou, N. I., Corominas-Murtra, B., Heisenberg, C. P. & Hannezo, E. Rigidity percolation

uncovers a structural basis for embryonic tissue phase transitions. *Cell* **184**, 1914-1928.e19

(2021).

19. Pinheiro, D., Kardos, R., Hannezo, É. & Heisenberg, C.-P. Morphogen gradient orchestrates

pattern-preserving tissue morphogenesis via motility-driven unjamming. *Nat. Phys.* **18**,

1482–1493 (2022).

20. Bi, D., Lopez, J. H., Schwarz, J. M. & Manning, M. L. A density-independent rigidity transi-

tion in biological tissues. *Nat. Phys.* **11**, 1074–1079 (2015).

21. Mitchel, J. A. *et al.* In primary airway epithelial cells, the unjamming transition is distinct from

the epithelial-to-mesenchymal transition. *Nat. Commun.* **11**, 1–14 (2020).

22. Kato, T. *et al.* Interplay of adherens junctions and matrix proteolysis determines the invasive

pattern and growth of squamous cell carcinoma. *eLife* **12**, e76520 (2023).

23. Grosser, S. *et al.* Cell and Nucleus Shape as an Indicator of Tissue Fluidity in Carcinoma.

Phys. Rev. X **11**, (2021).

24. Ilina, O. *et al.* Cell–cell adhesion and 3D matrix confinement determine jamming transitions

in breast cancer invasion. *Nat. Cell Biol.* **22**, 1103–1115 (2020).

25. Malinverno, C. *et al.* Endocytic reawakening of motility in jammed epithelia. *Nat. Mater.* **16**,

587–596 (2017).

26. Oswald, L., Grosser, S., Smith, D. M. & Käs, J. A. Jamming transitions in cancer. *J. Phys.*

Appl. Phys. **50**, 483001 (2017).

27. Fuhs, T. *et al.* Rigid tumours contain soft cancer cells. *Nat. Phys.* **18**, 1510–1519 (2022).

28. Frittoli, E. *et al.* Tissue fluidification promotes a cGAS–STING cytosolic DNA response in in-

vasive breast cancer. *Nat. Mater.* (2022) doi:10.1038/s41563-022-01431-x.

29. Kardar, M., Parisi, G. & Zhang, Y.-C. Dynamic Scaling of Growing Interfaces. *Phys. Rev.*

Lett. **56**, 889–892 (1986).

30. Muzzio, N. E., Pasquale, M. A., González, P. H. & Arvia, A. J. Influence of individual cell

motility on the 2D front roughness dynamics of tumour cell colonies. *J. Biol. Phys.* **40**, 285–

308 (2014).

31. Brú, A., Albertos, S., Luis Subiza, J., García-Asenjo, J. L. & Brú, I. The Universal Dynamics

of Tumor Growth. *Biophys. J.* **85**, 2948–2961 (2003).

32. Maunuksela, J. *et al.* Kinetic Roughening in Slow Combustion of Paper. *Phys. Rev. Lett.* **79**,

1515–1518 (1997).

33. Takeuchi, K. A. & Sano, M. Evidence for Geometry-Dependent Universal Fluctuations of the

Kardar-Parisi-Zhang Interfaces in Liquid-Crystal Turbulence. *J. Stat. Phys.* **147**, 853–890

(2012).

34. Giard, D. J. *et al.* In vitro cultivation of human tumors: establishment of cell lines derived

from a series of solid tumors. *J. Natl. Cancer Inst.* **51**, 1417–1423 (1973).

35. Macpherson, I. R. *et al.* p120-catenin is required for the collective invasion of squamous cell

carcinoma cells via a phosphorylation-independent mechanism. *Oncogene* **26**, 1210334

(2007).

36. Bertillot, F., Miroshnikova, Y. A. & Wickström, S. A. SnapShot: Mechanotransduction in the

nucleus. *Cell* **185**, 3638-3638.e1 (2022).

37. Kalukula, Y., Stephens, A. D., Lammerding, J. & Gabriele, S. Mechanics and functional con-

sequences of nuclear deformations. *Nat. Rev. Mol. Cell Biol.* **23**, 583–602 (2022).

38. Venturini, V. *et al.* The nucleus measures shape changes for cellular proprioception to con-

trol dynamic cell behavior. *Science* **370**, eaba2644 (2020).

39. Venturini, V. *et al.* The nucleus as a mechanosensitive controller of cell behavior. *Biophys.*

J. **121**, 490a (2022).

40. Lomakin, A. J. *et al.* The nucleus acts as a ruler tailoring cell responses to spatial con-

straints. *Science* **370**, eaba2894 (2020).

41. Kidiyoor, G. R. *et al.* ATR is essential for preservation of cell mechanics and nuclear integ-

rity during interstitial migration. *Nat. Commun.* **11**, 4828 (2020).

42. Alert, R. & Trepat, X. Physical Models of Collective Cell Migration. *Annu. Rev. Condens.*

Matter Phys. **11**, 77–101 (2020).

43. Basan, M., Elgeti, J., Hannezo, E., Rappel, W.-J. & Levine, H. Alignment of cellular motility

forces with tissue flow as a mechanism for efficient wound healing. *Proc. Natl. Acad. Sci.*

110, 2452–2459 (2013).

44. Garcia, S. *et al.* Physics of active jamming during collective cellular motion in a monolayer.

Proc. Natl. Acad. Sci. **112**, 15314–15319 (2015).

45. Henkes, S., Fily, Y. & Marchetti, M. C. Active jamming: Self-propelled soft particles at high

density. *Phys. Rev. E* **84**, 040301 (2011).

46. Family, F. & Vicsek, T. Scaling of the active zone in the Eden process on percolation net-

works and the ballistic deposition model. *J. Phys. Math. Gen.* **18**, L75 (1985).

47. Chepizhko, O. *et al.* Bursts of activity in collective cell migration. *Proc. Natl. Acad. Sci.* **113**,

11408–11413 (2016).

48. Podewitz, N., Jülicher, F., Gompper, G. & Elgeti, J. Interface dynamics of competing tissues.

New J. Phys. **18**, 083020 (2016).

49. Rapin, G. *et al.* Roughness and dynamics of proliferating cell fronts as a probe of cell–cell

interactions. *Sci. Rep.* **11**, 8869 (2021).

50. Olami, Z., Procaccia, I. & Zeitak, R. Interface roughening in systems with quenched disor-

der. *Phys. Rev. E* **52**, 3402–3414 (1995).

51. Dias, C. S., Yunker, P. J., Yodh, A. G., Araújo, N. a. M. & Gama, M. M. T. da. Interaction an-

isotropy and the KPZ to KPZQ transition in particle deposition at the edges of drying drops.

Soft Matter **14**, 1903–1907 (2018).

52. Csahók, Z., Honda, K., Somfai, E., Vicsek, M. & Vicsek, T. Dynamics of surface roughening
in disordered media. *Phys. Stat. Mech. Its Appl.* **200**, 136–154 (1993).

53. Muzzio, N. E. *et al.* Spatio-temporal morphology changes in and quenching effects on the
2D spreading dynamics of cell colonies in both plain and methylcellulose-containing culture
media. *J. Biol. Phys.* **42**, 477–502 (2016).

54. Mark, S. *et al.* Physical Model of the Dynamic Instability in an Expanding Cell Culture. *Bio-*
phys. J. **98**, 361–370 (2010).

55. Tarle, V., Ravasio, A., Hakim, V. & Gov, N. S. Modeling the finger instability in an expanding
cell monolayer. *Integr. Biol.* **7**, 1218–1227 (2015).

56. Büscher, T., Diez, A. L., Gompper, G. & Elgeti, J. Instability and fingering of interfaces in
growing tissue. *New J. Phys.* **22**, 083005 (2020).

57. Dai, W. *et al.* Tissue topography steers migrating Drosophila border cells. *Science* **370**,
987–990 (2020).

58. Park, D. *et al.* Extracellular matrix anisotropy is determined by TFAP2C-dependent regula-
tion of cell collisions. *Nat. Mater.* 1–12 (2019) doi:10.1038/s41563-019-0504-3.

59. Conklin, M. W. *et al.* Aligned Collagen Is a Prognostic Signature for Survival in Human
Breast Carcinoma. *Am. J. Pathol.* **178**, 1221–1232 (2011).

60. Mukherjee, A., Jana, A., Koons, B. & Nain, A. Design of Fiber Networks for Studying Meta-
static Invasion. in *Biomechanics in Oncology* (eds. Dong, C., Zahir, N. & Konstantopoulos,
K.) 289–318 (Springer International Publishing, Cham, 2018). doi:10.1007/978-3-319-
95294-9_14.

61. Tsingos, E., Bakker, B. H., Keijzer, K. A. E., Hupkes, H. J. & Merks, R. M. H. Hybrid cellular
Potts and bead-spring modeling of cells in fibrous extracellular matrix. *Biophys. J.* **122**,
2609–2622 (2023).

62. Labernadie, A. *et al.* A mechanically active heterotypic E-cadherin/N-cadherin adhesion en-
ables fibroblasts to drive cancer cell invasion. *Nat. Cell Biol.* **19**, 224–237 (2017).

63. Leithner, A. *et al.* Diversified actin protrusions promote environmental exploration but are

dispensable for locomotion of leukocytes. *Nat. Cell Biol.* **18**, 1253–1259 (2016).

64. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods*

9, 676–682 (2012).

65. Tinevez, J.-Y. *et al.* TrackMate: An open and extensible platform for single-particle tracking.

Methods **115**, 80–90 (2017).

66. Linkert, M. *et al.* Metadata matters: access to image data in the real world. *J. Cell Biol.* **189**,

777–782 (2010).

67. Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nat. Methods* **16**,

1226–1232 (2019).

68. Agostinelli, C. & Lund, U. *R Package \texttt{circular: Circular Statistics (Version 0.4-95)*. (CA:

Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University,

Venice, Italy. UL: Department of Statistics, California Polytechnic State University, San Luis

Obispo, California, USA, 2022).

69. Pewsey, A., Neuhäuser, M. & Ruxton, G. D. *Circular Statistics in R*. (OUP Oxford, 2013).

70. Howse, J. R. *et al.* Self-Motile Colloidal Particles: From Directed Propulsion to Random

Walk. *Phys. Rev. Lett.* **99**, 048102 (2007).

71. Shaebani, M. R., Wysocki, A., Winkler, R. G., Gompper, G. & Rieger, H. Computational

models for active matter. *Nat. Rev. Phys.* **2**, 181–199 (2020).

72. Anderson, J. A., Glaser, J. & Glotzer, S. C. HOOMD-blue: A Python package for high-perfor-

mance molecular dynamics and hard particle Monte Carlo simulations. *Comput. Mater. Sci.*

173, 109363 (2020).

73. Mandal, R., Bhuyan, P. J., Chaudhuri, P., Dasgupta, C. & Rao, M. Extreme active matter at

high densities. *Nat. Commun.* **11**, 2581 (2020).

74. Angelini, T. E. *et al.* Glass-like dynamics of collective cell migration. *Proc. Natl. Acad. Sci.*

108, 4714–4719 (2011).

75. Kang, W. *et al.* A novel jamming phase diagram links tumor invasion to non-equilibrium

phase separation. *iScience* **24**, 103252 (2021).

76. Narinder, N., F. Bos, M., Abaurrea-Velasco, C., Graaf, J. de & Bechinger, C. Understanding

enhanced rotational dynamics of active probes in rod suspensions. *Soft Matter* **18**, 6246–

6253 (2022).

77. Yamashita, H. *et al.* Lysophosphatidic Acid Upregulates Laminin-332 Expression during

A431 Cell Colony Dispersal. *J. Oncol.* **2010**, 107075 (2010).

78. Tweedy, L. *et al.* Seeing around corners: Cells solve mazes and respond at a distance using

attractant breakdown. *Science* **369**, eaay9792 (2020).

79. Zhou, W. & Yan, H. Alpha shape and Delaunay triangulation in studies of protein-related in-

teractions. *Brief. Bioinform.* **15**, 54–64 (2014).

80. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat.*

Methods **17**, 261–272 (2020).