A Goldilocks principle of JAK/STAT signaling governs airway

epithelial homeostasis and stress adaptation in *Drosophila*

Xiao Niu[1, 2], Christine Fink[1], Kimberley Kallsen[3, 4], Leizhi Shi[1, 5], Viktoria Mincheva[1],

Sören Franzenburg[6], Ruben Prange[1], Jingjing He[1], Anita Bhandari[1], Iris Bruchhaus[7],

Holger Heine[2, 8], Judith Bossen[1, 8]*, Thomas Roeder[1, 8]*

[1]Kiel University, Zoology, Dept. of Molecular Physiology, Kiel, Germany

[2]Current address: Weifang Medical University, College of Life Science and

Technology, Weifang, China

[3]Research Center Borstel – Leibniz Lung Center, Div. of Innate Immunity, Borstel,

Germany

[4]Current address: Boehringer Ingelheim, Ingelheim, Germany

[5]Department of Thoracic Surgery, Linyi People's Hospital, Linyi, Shandong, 276000,

China

[6]Kiel University, IKMB, Kiel, Germany

[7]Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany

[8]German Center for Lung Research, Airway Research Center North, Kiel, Germany

*Correspondence: Judith Bossen, email: jbossen@zoologie.uni-kiel.de; Thomas

Roeder, email: troeder@zoologie.uni-kiel.de

Graphical abstract

Summary

Airway epithelia must maintain barrier integrity while continuously adapting to
changing environmental conditions. This requires key signaling pathways in airway
epithelia to operate within a tightly controlled functional range. Here, we show that
airway epithelial homeostasis in *Drosophila* depends on balanced JAK/STAT signaling.
Basal pathway activity is constitutively present in differentiated epithelial cells and is
necessary for cell survival, epithelial integrity and stress resistance. In accordance
with this role as a stress responsive regulator in epithelial biology, environmental
stressors, including hypoxia, cigarette smoke, and cold exposure, induce JAK/STAT
signaling. In contrast, sustained pathway activation is associated with airway
remodeling characterized by epithelial thickening, luminal narrowing, and altered
cellular organization. Transcriptomic analysis reveals that sustained activation is
associated with a coordinated epithelial stress program integrating immune
signaling, proteostasis, and metabolic adaptation. Cross-species comparisons with
murine and human datasets suggest that key aspects of this response are conserved.
Pharmacological inhibition demonstrates that remodeling depends on continued
pathway activity and can be partially reversed *in vivo*. Together, our findings support
a model in which airway epithelial homeostasis depends on maintaining JAK/STAT
signaling within a defined functional range. This "Goldilocks" principle provides a
conceptual framework for understanding how epithelial stress responses are
balanced under physiological and pathological conditions.

Introduction
Airway epithelia form the first barrier between the organism and the external
environment and are therefore continuously exposed to environmental stressors,
including pathogens, pollutants, and fluctuations in oxygen availability[1, 2].
Maintaining epithelial integrity under these conditions requires the coordinated
regulation of signaling pathways that govern epithelial survival, stress responses, and
tissue architecture[3, 4]. Dysregulation of epithelial signaling can lead to pathological
remodeling processes that contribute to chronic airway diseases, including asthma
and chronic obstructive pulmonary disease [5-7]. Understanding how epithelial
signaling pathways maintain airway homeostasis while enabling adaptive responses
to environmental stress, therefore, represents a fundamental challenge in lung
research.
Among the signaling pathways implicated in epithelial regulation, the JAK/STAT
signaling pathway plays a central role in epithelial stress responses and tissue
remodeling across metazoans[8, 9]. In mammalian airways, JAK/STAT signaling is
activated by inflammatory cytokines and has been implicated in epithelial
proliferation, immune responses, and airway remodeling during chronic lung
disease[9-11]. However, despite extensive work in mammalian systems, it remains
unclear how JAK/STAT signaling contributes to airway epithelial homeostasis under
physiological conditions and how different levels of pathway activity affect epithelial
structure and function[12-15][4].
The *Drosophila* tracheal system provides a powerful in vivo model to study airway
epithelial biology. The fly airway epithelium shares key structural and functional
features with mammalian airway epithelia while offering unparalleled genetic
accessibility to manipulate signaling pathways directly in epithelial cells[16-21].
Previous work has established that JAK/STAT signaling is important during tracheal
morphogenesis[22-24] and that it regulates epithelial responses to infection and
injury in multiple tissues in *Drosophila*[25, 26]. However, the role of this pathway in
maintaining airway epithelial homeostasis and coordinating responses to
environmental stress has not been systematically investigated.
Here, we use the genetically tractable *Drosophila* airway system to examine how
JAK/STAT signaling regulates epithelial homeostasis. We show that basal pathway
activity contributes to epithelial integrity, whereas sustained activation is associated
with structural remodeling. Furthermore, we demonstrate that environmental stress
dynamically modulates pathway activity and that chronic activation induces a
coordinated epithelial stress program. Based on these findings, we propose that
airway epithelia require a "Goldilocks" level of JAK/STAT signaling, in which balanced
pathway activity maintains homeostasis while insufficient or excessive signaling
disrupts tissue integrity.
89
90

Results

To investigate how JAK/STAT signaling regulates airway epithelial homeostasis, we combined genetic, environmental, and transcriptomic approaches in the *Drosophila* airway system. We first assessed basal pathway activity and its requirement for epithelial integrity. We then examined how environmental stress modulates pathway activity, followed by analysis of the structural and molecular consequences of sustained pathway activation. Finally, we tested whether pharmacological inhibition can reverse the observed remodeling phenotypes.

JAK/STAT Signaling Is Present throughout the Larval Tracheal System

To determine whether JAK/STAT signaling is active under physiological conditions, we examined pathway activity using a STAT-responsive transcriptional reporter (Fig. 1A). Under basal conditions, the trachea of third-instar *Drosophila* larvae (L3) displayed near-ubiquitous reporter activity across all tracheal compartments (Fig. 1B-E). However, marked regional differences in signal intensity were evident. Specifically, JAK/STAT signaling was enriched in defined zones of the larval tracheal system, including the dorsal trunks, spiracular branches (SB), and Tr2, regions previously associated with regenerative capacity[26,27] (Fig. 1D, E). In contrast, tracheal regions fated for apoptosis during metamorphosis (e.g., Tr10) exhibited only minimal reporter activity (Fig. 1C), highlighting a functional correlation between JAK/STAT activity and tissue maintenance.

In adult flies, reporter analysis revealed sustained pathway activity throughout the airway network (Fig. 1F), confirming a role for JAK/STAT signaling beyond developmental morphogenesis. Transcriptional mapping of *upd2* further localized ligand expression to SB and Tr2 tracheoblasts (Fig. S1), consistent with an epithelial-intrinsic autocrine regulatory circuit.

These observations indicate that JAK/STAT signaling is constitutively active in differentiated airway epithelial cells, suggesting a potential role in maintaining epithelial homeostasis. We next asked whether this basal pathway activity is functionally required for airway epithelial integrity.

Basal JAK/STAT signaling is required for airway epithelial homeostasis

To assess the functional relevance of basal pathway activity, we reduced pathway activity in airway epithelial cells. We selectively inhibited the pathway in airway epithelial cells using the TARGET system [27] to drive a dominant-negative *Domeless* receptor allele (Dome.DN) (Fig. 2A). Early (embryo) pathway blockade caused larval lethality (Fig. 2B) and led to severe architectural defects, including loss of dorsal trunks and segmental fusion (Fig. 2C). In larval trachea, inhibition of JAK/STAT signaling (here, driven by nach-Gal4) induced epithelial defects characterized by liquid-filled airways, a phenotype that is apparent as a lack of visible trachea in the animals, as liquid filling pushes away the air and thus the differences in refractive

indexes between the two media (Fig. 2D, yellow arrows). Moreover, the outcome of
the tracheal activation of JAK/STAT signaling critically depended on the time of
activation (Fig. S2). At the level of single epithelial cells (here, driven by the vvl-
coin system) strong changes in cell structure, reminiscent of apoptosis, occurred
(Fig. 2E), which was confirmed by cleaved Dcp-1 staining (Fig. 2F–G).
In adults, trachea-specific pathway inhibition reduced organismal lifespan and
sensitized animals to chronic CSE exposure, but not to hypoxia (Fig. 2H–J),
highlighting stress-specific functional dependencies. Here, CSE and reduced
JAK/STAT activity independently reduced lifespan. These findings demonstrate that
basal JAK/STAT signaling is required to maintain airway epithelial integrity and
organismal stress tolerance. Because airway epithelia constantly encounter
environmental challenges, we next investigated whether JAK/STAT signaling
responds dynamically to stress.
145
Environmental stress activates JAK/STAT signaling in airway epithelial cells
To determine whether environmental stress regulates epithelial JAK/STAT signaling,
we exposed animals to several physiologically relevant stress conditions. To evaluate
the responsiveness of airway epithelial cells to environmental insults, we employed a
GFP-based reporter driven by STAT92E activation in *Drosophila*. Exposure to airborne
stressors, including cold air, hypoxia, and cigarette smoke extract (CSE), triggered

robust and reproducible pathway activation in larval tracheae (Fig. 3A, B). Similar

results were observed in adult flies (Fig. 3C, D). Quantitative imaging confirmed a
significant increase in reporter fluorescence under all tested stress conditions.
Strikingly, pathway activation co-localized with elevated expression of the cytokine
upd3, and *upd2* was additionally induced by CSE, whereas other stressors induced
much smaller effects (Fig. S3). To test whether a simple intra-organ signaling system
is operative, we subjected flies expressing the STAT-reporter, concurrently with RNAi
targeting either upd2 or upd3, to hypoxia. Here, we found no rescue of the hypoxia-
induced STAT-activation by the RNAi against these most important ligands of the
pathway (Fig. 3E, F), which implies that this system is more complex, including other
sources of the unpaired ligands or plasticity among different unpaired ligands
replacing each other.
These results demonstrate that airway epithelial JAK/STAT signaling functions as a
stress-responsive signaling pathway activated by environmental challenges. We next
asked how sustained pathway activation affects the structure of the airway
epithelium.
168
Chronic JAK/STAT activation drives airway epithelial remodeling
To determine the consequences of sustained pathway activation, we chronically
activated JAK/STAT signaling in airway epithelial cells. Therefore, we overexpressed

either *Hop.CA* (constitutively active JAK) or the cytokine *upd3* (Fig. 4A). Both
perturbations induced at the embryonic state led to embryonic or early larval lethality
(Fig. 4B, C). The structural analysis of these perturbations revealed severe epithelial
disruption. Here, *upd3* overexpression yielded more severe epithelial disruption than
Hop.CA overexpression (Fig. 4D-E). Conditional activation of the pathway in larvae via
the TARGET system resulted in structural changes in the tracheal system (Fig. 4F, G).
Here, structural defects became apparent, including airway stenoses. A closer
analysis of the structural changes of the dorsal trunks of L3 larvae using a mosaic
system based vvl-coin revealed a narrowing of the airway lumen only in the affected
epithelial cells (Fig. 4F, G). A detailed analysis of the chitinous intima of the larval
trachea showed that the usually highly regular taenidial structure was strongly
impaired and characterized by irregularities (Fig. S4).
These findings indicate that excessive JAK/STAT signaling induces pathological airway
remodeling, suggesting that epithelial homeostasis requires tightly controlled
pathway activity. We next sought to determine the cellular mechanisms underlying
this remodeling phenotype.
188
JAK/STAT hyperactivation alters cell structure and junctional organization
In response to JAK/STAT activation, the tracheal epithelial cells exhibited pronounced
epithelial thickening. This thickening response was observed in response to *Hop.CA*
and *upd3* overexpression, with much higher thickening in response to *Hop.CA* (Fig.
4H, I). In addition to the other structural defects observed, the localization of
junctional proteins and epithelial polarity markers was analyzed in control and
Hop.CA-expressing airway epithelial cells. This phenotype was fully cell-autonomous,
as demonstrated by mosaic analysis (Fig. 4L), and driven by increased cell volume
rather than hyperplasia (Fig. S5). We observed locally occurring thickening of the
epithelium to strong CSE application (Fig. 4K), resembling reactions to infection in
the larval airway epithelium [28]. Furthermore, weaker expression drivers (*nach*-Gal4)
produced the same phenotype with reduced severity (Fig. S6). Epithelial thickening in
the trachea also occurs in response to activation of the IMD pathway (TNF-α receptor
equivalent, induced by *btl-Gal4*, *UAS-PGRP-LC*) [29]. To test, if JAK/STAT pathway
activation uses the same mechanisms, we blocked STAT expression while activating
the IMD pathway and found no phenotype rescue (Fig. S6E).
To investigate cellular mechanisms underlying epithelial remodeling, we examined
the localization of junctional proteins (Coracle (Cora) and Armadillo (Arm)). In control
tissues, these proteins were restricted to cell–cell boundaries. In contrast, sustained
JAK/STAT activation was associated with intracellular accumulation of junctional
components, suggesting altered protein trafficking or localization. Using mosaic
analysis, we observed intracellular accumulation of Arm and Cora, respectively, in
affected cells (Fig. 4M-P), which is not seen in control cells. To investigate the

molecular programs underlying these structural changes, we next performed
transcriptomic analysis of airway epithelial cells with activated JAK/STAT signaling.
214
Sustained JAK/STAT Activation induces a Coordinated Transcriptional Response
Transcriptomic analysis of airway epithelial cells with sustained JAK/STAT activation
identified widespread changes in gene expression. RNAseq data are deposited in the
GEO database (GSE212132). Upregulated genes included canonical pathway targets as
well as genes associated with immune responses, stress pathways, and metabolism.
Downregulated genes were enriched for functions related to tissue structure and
differentiation (Fig. 5A–B). A closer analysis revealed major upregulated genes (Table
1) and the KEGG pathways involved (Table S1). Transcription factor binding site
analyses revealed STAT92e as one of the hits (Table 2).
Comparison with previously described gene sets revealed partial overlap with
immune and stress-related transcriptional programs (Trabid-deficiency, tracheal
infection, and PGRP-LE overexpression signatures [29-31]). These findings suggest
that JAK/STAT activation induces a composite transcriptional state integrating
multiple regulatory pathways. The corrected Trabid gene set (n = 890) showed a
substantial overlap of 291 genes with the Hop dataset (32.7% of the Trabid set;
Jaccard index = 0.113). Hypergeometric testing revealed a >2-fold enrichment over
random expectation (expected ≈127 genes; observed = 291; p < 10^{-100}), indicating
strong biological convergence. The overlapping genes included canonical
antimicrobial peptides (e.g., *Drs*, *IM4*), heat shock proteins (*Hsp22*, *Hsp70Bb*),
cytochrome P450 enzymes, and lipid metabolic regulators, consistent with activation
of immune, stress, and metabolic remodeling programs. Comparisons with Infection
and PGRP-LE signatures revealed partial but not complete overlap, supporting the
interpretation that Hop activation induces an infection-like state that extends beyond
classical Imd pathway activation.
Together, these findings demonstrate that Hop signaling engages a composite
transcriptional program integrating immune activation, proteotoxic stress responses,
and metabolic reprogramming.
242
Cross-Species Comparison Reveals Conserved Features of Epithelial Stress
Responses
To assess the broader relevance of these transcriptional changes, we compared the
Drosophila dataset with gene expression profiles from murine asthma models and
cytokine-stimulated human airway epithelia. These analyses revealed shared
features, including activation of immune signaling, oxidative stress responses, and
alterations in epithelial transport and metabolism. To enable the direct comparison,
we generated lists of mouse homologs of the regulated *Drosophila* genes using the
DIOPT online tool [32]. The overlaps are highly significant (Fig. 5C, D), with more than
15-20% overlap with the mouse data. Constitutive Hop/JAK signaling induces a

transcriptional program characterized by activation of innate immune effectors,
antimicrobial peptides, stress responses, and secretory pathway components,
accompanied by extensive remodeling of metabolism and trafficking. We evaluated
this murine dataset by testing cytokine genes and selected genes involved in
vesicular transport by qRT-PCR (Fig. 5E). The direct comparison with the acute
mouse asthma model revealed a common signature for the upregulated genes,
including 1) antimicrobial/innate immune activation, 2) extensive cytokine signaling,
3) oxidative stress response, and 4) increased secretory activity. Downregulated are
especially programs associated with tissue differentiation and metabolic homeostasis
(Fig. 5F).
Furthermore, we compared the RNAseq data with those derived from human airways
subjected to a strong and specific JAK/STAT stimulus, namely IL-6 transsignaling [33].
Again, we observed an overlap in the DEGs with the *Drosophila* data transformed
using the DIOPT pipeline (Fig. 5G). Commonalities of the differentially expressed
genes comprise a coordinated transcriptional program characterized by activation of
ER stress, inflammatory signaling, epithelial transport, and metabolic pathways.
Genes involved in the unfolded protein response and ER proteostasis (*HSPA5,*
HSP90B1, HYOU1, ATF4, ERP44) and ER–Golgi trafficking (*SEC23B, SEC31B, COPB1*) were
enriched. Concurrently, key components of innate immune signaling and NF-κB
activation (*TLR1, IRAK1, IRAK4, RELA, RELB, BIRC3*) and oxidative stress regulators
(*NOX1, DUOX1, DUOX2, PRDX4, CAT*) were detected. The gene set also included
epithelial transport and barrier-associated genes such as *CFTR, SCNN1A/B/G, SLC26A9,*
and *AQP3/AQP4*, together with extracellular matrix and adhesion genes (*FBLN1,*
EFEMP1, LAMC2, ITGA family). Additionally, enzymes of the mevalonate and lipid
metabolism pathways (*HMGCR, HMGCS1, MVD, PLIN2/3/4/5*) were enriched, indicating
coordinated metabolic remodeling (Fig. 5H, I).
While these comparisons are based on orthology mapping and do not imply direct
equivalence, they suggest that key aspects of epithelial stress responses may be
conserved across species. Finally, we asked whether the airway remodeling
phenotype can be rescued by pharmacological intervention.
283
Pharmacological Inhibition of JAK/STAT Reverses Epithelial Remodeling
To determine whether airway remodeling depends on sustained pathway activation,
we tested whether pharmacological inhibition of JAK signaling could reverse
epithelial defects. Treatment of *Hop.CA*-expressing flies with a JAK inhibitors with a
JAK1-like profile (Oclacitinib, Baricitinib, and Filgotinib) reduced epithelial thickening.
Other JAK inhibitors with a different pharmacological profile (Tofacitinib, CYT387,
Ruxolitinib) failed to rescue the phenotype (Fig. 6A).
291

Summary
Together, these results demonstrate that airway epithelial homeostasis requires
JAK/STAT signaling to operate within a narrow functional range. Basal signaling
maintains epithelial integrity and stress resistance, whereas chronic hyperactivation
induces pathological airway remodeling through conserved epithelial stress
programs (Fig. 6C, D). Together, these findings demonstrate that airway epithelial
homeostasis requires JAK/STAT signaling to operate within a narrow functional
range. Basal pathway activity supports epithelial integrity and stress resistance,
whereas sustained hyperactivation disrupts epithelial architecture and induces
conserved airway remodeling programs.
302
303

Discussion

Basal Homeostasis and Epithelial Viability
Leveraging the genetically streamlined architecture of *Drosophila melanogaster*, this
study uncovers the bifunctional role of the JAK/STAT cascade in airway epithelia.
Because the system relies on a single receptor (Domeless), kinase (Hopscotch), and
transcription factor (STAT92E) [34], we circumvented vertebrate redundancy [35] to
dissect pathway dynamics at high resolution. Our data indicate that JAK/STAT
signaling operates within a narrow functional range in airway epithelial cells. Basal
pathway activity is required to maintain epithelial integrity and organismal stress
resistance, whereas sustained hyperactivation induces structural remodeling and
disrupts epithelial organization. Together, these findings support a "Goldilocks"
model in which balanced JAK/STAT signaling preserves airway homeostasis, while
both insufficient and excessive activity impairs epithelial function.
This requirement for basal activity perfectly aligns with recent mammalian findings
demonstrating that tonic JAK/STAT signaling is critical for maintaining cellular
homeostasis and identity across immune and epithelial compartments [36, 37].
Furthermore, our observation that early pathway disruption causes catastrophic
architectural defects during tracheal tube formation echoes vertebrate studies
establishing STAT3 as a fundamental requirement for lung organogenesis and
epithelial lineage maintenance [12, 38].
324
Stress Responsiveness and Dynamic Signaling
The airway epithelium represents the primary interface between the organism and
environmental stressors [39]. In this study, exposure to hypoxia, cold air, and
cigarette smoke extract (CSE) induced robust activation of JAK/STAT signaling in
airway epithelial cells. This response was associated with increased expression of
Upd ligands, suggesting that epithelial cells actively participate in amplifying stress

signals, but that they are not the only source of upd expression in response to
stressors.
These findings indicate that JAK/STAT signaling functions as a stress-responsive
pathway in airway epithelia. Importantly, our data suggest that both insufficient and
excessive pathway activity are detrimental: reduced signaling sensitizes the organism
to stress, whereas sustained activation is associated with structural remodeling.
Together, this supports a model in which airway epithelial homeostasis depends on
maintaining JAK/STAT signaling within a defined functional range.
339
A Goldilocks Principle of JAK/STAT Signaling in Airway Epithelia
Our findings support a "Goldilocks" principle of JAK/STAT signaling in airway
epithelia, in which both insufficient and excessive pathway activity compromise
tissue integrity. Basal signaling appears to support epithelial survival and stress
resistance, whereas chronic hyperactivation is associated with epithelial remodeling
and functional disruption.
This conceptual framework provides a unifying explanation for the dual roles of
JAK/STAT signaling observed in both physiological and pathological contexts. While
acute activation likely supports adaptive responses, sustained signaling may shift the
system toward a maladaptive state. This echoes the delicate balance of the JAK/STAT
pathway in airway epithelia [12]. If the pathway fails to activate, the organism is
heavily sensitized to chronic stress. Conversely, as seen in chronic lung conditions like
asthma and COPD where the JAK/STAT pathway remains persistently hyperactive [10,
40], the protective stress response transforms into pathology.
354
Stress-Induced Transcriptional Reprogramming
Transcriptomic profiling of airway epithelial cells with sustained JAK/STAT activation
revealed a coordinated transcriptional response characterized by upregulation of
innate immune effectors, stress response pathways, and metabolic regulators. These
findings suggest that chronic pathway activation induces a broad epithelial stress
program rather than a purely immune-specific response.
Comparison with previously described gene sets, including immune and stress-
related signatures, revealed significant overlap, indicating that JAK/STAT activation
engages conserved regulatory modules [30, 31]. The transcriptional reprogramming
induced by sustained JAK/STAT activation seems to fundamentally alter the cellular
identity, shifting it into a systemic, stress-adapted state that closely mimics the
inflammatory landscapes seen in idiopathic pulmonary fibrosis and acute lung injury
[41, 42]. However, these overlaps are based on orthology mapping and should be
interpreted cautiously, as species-specific differences in gene regulation are not fully
captured.
370
Molecular Mechanisms Underlying Airway Remodeling

Our data support a model in which sustained JAK/STAT activation induces a
coordinated stress response that increases secretory demand and proteostatic load
within epithelial cells. This imbalance is likely to affect vesicular trafficking and
protein localization, as suggested by the intracellular accumulation of junctional
proteins such as Coracle and Armadillo.
These observations are consistent with impaired trafficking and junctional
organization contributing to epithelial thickening and lumen narrowing. This
illustrates how the transcriptomic shift toward increased, yet defective, secretory
activity compromises the delivery of critical structural components, ultimately
leading to barrier failure and organismal lethality, a structural decline reminiscent to
the epithelial-to-mesenchymal transition (EMT) seen in progressive lung disease [43].
While the present study does not establish a direct causal sequence between these
processes, the data support a mechanistic link between sustained pathway
activation, cellular stress responses, and structural remodeling.
386

Conservation Across Species and Disease Contexts
Cross-species comparisons revealed overlap between the transcriptional signatures
induced by JAK/STAT activation in *Drosophila* and those observed in murine asthma
models and cytokine-stimulated human airway epithelia. These shared features
include activation of immune signaling pathways, stress responses, and alterations in
epithelial transport and metabolism.
While these comparisons rely on computational mapping and do not imply direct
equivalence, they suggest that core aspects of epithelial stress responses may be
conserved across species. This supports the use of the *Drosophila* airway system as a
model to investigate general principles of epithelial remodeling.
397

Pharmacological Modulation and Translational Implications
Pharmacological inhibition experiments demonstrated that airway remodeling
induced by sustained JAK/STAT activation can be partially reversed in vivo. Notably,
inhibitors with higher specificity toward JAK1-like activity showed greater efficacy in
reducing epithelial thickening. These results precisely mirror the specific kinase
binding profiles established in vertebrate models [44-46]. This functional divergence
suggests that the hyperactive *Drosophila* Hop.CA kinase shares a profound structural
or functional homology with vertebrate JAK1, establishing this in vivo model as a
highly specific screening platform for next-generation, JAK1-targeted respiratory
therapeutics [16, 17, 47]. Although inhibition of the JAK/STAT signaling pathway is
currently being actively explored as a therapeutic strategy, particularly for chronic
forms of asthma that are refractory to conventional treatments [48], it may interfere
with the regenerative capacity of the airway epithelium [13]. Thus, therapeutic
strategies will likely require careful modulation rather than complete inhibition.
412

Conclusion
In summary, our findings define a narrow functional window for JAK/STAT signaling
in airway epithelia, in which balanced pathway activity is required to maintain
epithelial integrity and stress resilience. Deviations from this balance—either through
insufficient or excessive signaling—lead to epithelial dysfunction and structural
remodeling. This "Goldilocks" principle provides a conceptual framework for
understanding how epithelial tissues integrate environmental signals while
preserving homeostasis and offers insights into conserved mechanisms underlying
airway disease.
422
423

Materials and Methods

Drosophila strains and husbandry
STAT92E-GFP was used to monitor the pathway's activation [49]; the Gal4-UAS
system [50] targeted ectopic expression to the tracheal system. The Gal4 drivers
included: *btl-Gal4, UAS-GFP* on the 3rd chromosome, and *btl-Gal4, UAS-GFP* on the 2nd
chromosome (obtained from the Leptin group, Heidelberg, Germany); *upd2-Gal4;*
upd3-Gal4 [19]; *nach-Gal4* [51]. The UAS responders consisted of: *UAS-LacZ.nls* (BDSC
3956), *UAS-domeΔcyt2.1* (*Dome.DN*), and *UAS-Hop.CA* (*UAS-hopTumL*), which were
obtained from N. Perrimon [22, 52]. The *UAS-upd3* was constructed in our lab. *TubP-*
Gal80[ts] (BDSC 7018) was obtained from the Bloomington stock center. Unless
otherwise stated, the flies were raised on standard medium at 25 °C with 50–60%
relative humidity under a 12:12 h light/dark cycle as described earlier [53].
436

Coin-FLP expression system
Vvl-FLP/CyO; btl-moe.mRFP (BDSC 64233), *tubP-Gal80[ts]* and *CoinFLP-Gal4, UAS-*
2xEGFP (BDSC 58751) were used to construct animals for the tracheal mosaic analysis.
Ventral veins lacking (*vvl*) was expressed in larval tracheal clones that covered
approximately 30 to 80% of the trachea [54, 55]. The genotype of the flies was *vvl-*
FLP, CoinFLP-Gal4, UAS-2xEGFP/CyO (*vvl-coin*) and *vvl-FLP, CoinFLP-Gal4, UAS-*
2xEGFP/CyO; tub-Gal80[ts] (*vvl-coin.ts*).
444

Developmental viability
For the developmental viability of eggs, they were collected overnight and not
physically handled in any way. The number of hatched eggs and pupae were counted
starting two days after collection. Each group had four replicates that included more
than 20 eggs. Tub-Gal80 [ts] was used to limit UAS responder expression during the
larval stage to study the developmental viability of larvae. Animals were raised at 18
°C to keep the UAS responder gene silent. Larvae at different instar stages were
transferred to a new medium at 29 °C. In this study, four replicates were typically

performed with 30 larvae each. The stage of the larvae was determined by the
appearance of the anterior spiracles.
455
Determination of epithelial thickness
The L2 and L3 larvae trachea were carefully dissected from the posterior side of the
body in PBS. The isolated trachea were immersed in 50% glycerol, and digital images
were captured within 15 min. L2 larvae were distinguished from L3 larvae by the
appearance of anterior spiracles. The relative ages of L3 larvae were inferred from
the animal's size. Thirty larvae were used in each group, and specimens were
analyzed by Image J.
463
Drug application in *Drosophila*
JAK inhibitors (Baricitinib #16707, Oclacitinib #18722, Filgotinib #17669 - Cayman
Chemicals, Michigan, USA) were diluted in DMSO [100 mM]. For later application, the
inhibitors were diluted 1:10 in 100% EtOH. We used 20 µl of each diluted inhibitor for 2
ml of concentrated medium (5% yeast extract, 5% corn flour, 5% sucrose, 1% low-melt
agarose, 1 ml of 10% propionic acid, and 3 ml of 10% Nipagin). The eggs of each
crossing have been applied to the modified medium and kept at 20 °C until the larvae
reach the L2 stage. Afterwards, they were incubated at 30 °C for 2 days to induce the
btl-Gal4, UAS-GFP; tub-Gal80[ts] (*btl.ts*)–driver. The trachea of L3 larvae were carefully
dissected from the posterior side of the body in PBS. The isolated trachea was
immersed in 50% glycerol, and digital images were captured every 15 minutes. 10
larvae were used for each group.
476
Experimental design for hypersensitivity pneumonitis in mice
Experimental design for hypersensitivity pneumonitis contains two processes,
sensitization and challenge of mice with ovalbumin (Sigma Aldrich, A5503). The
protocol refers to the protocol that [56] is described with a few modifications. 8-
week-old Balb/C mice were sensitized with intraperitoneal injections of 20 µg of OVA
emulsified in aluminum hydroxide in a total volume of 1 ml on days 7 and 14, followed
by three consecutive challenges each day by exposure to OVA or PBS aerosol for 30
min. Mice were sacrificed 24 h following the final challenge. The left lungs were
collected for histological analysis, and the superior lobes were dissected for RNA
analysis. Six mice per group were used in this experiment and four mice were
randomly selected in each group for analyses. Genesky performed total mRNA
sequencing, data processing, and statistical analysis (Shanghai, China). The
experimental protocols were approved by the Animal Care and Protection
Committee of Weifang Medical University (2021SDL418).
491
AB-PAS staining and immunofluorescence analysis of mouse lung

Mice were sacrificed in an excess of CO_2 gas. The lungs of euthanized mice were
inflated by intratracheal injection of cold 4% paraformaldehyde and then fixed and
embedded in paraffin as described in [57]. AB-PAS staining and Immunofluorescence
analysis were supported by Servicebio (Wuhan, China). All sections were
photographed using a microscope slide scanner (Pannoramic MIDI: 3Dhistech). The
materials and methods could be found at the following link
https://www.servicebio.cn/data-detail?id=3040&code=MYYGSYBG   and
https://www.servicebio.cn/data-detail?id=35_95&code=RSSYBG. We list the primary
reagents here: AB-PAS solution set (Servicebio, G1049), Anti-CD11b (Servicebio,
GB11058), and DAPI (Servicebio, G1012-10ML).

Time-lapse microscopy
All images were acquired using a ZEISS Axio Image Z1 and a ZEISS LSM 880
fluorescent microscope (INST 257/591-1 FUGG). Embryos were dechorionated in 3%
sodium hypochlorite and immersed in Halocarbon oil 700 (Sigma Aldrich, 9002-83-9,
Deisenhofen, Germany). Then the embryos were imaged after stage 15 when the
tracheal tree formed at 3-hour intervals. 40 embryos were investigated in each
group.
510
Cigarette smoke and hypoxic exposure
All cigarette smoke exposure experiments were carried out in a smoking chamber,
attached to a diaphragm pump. Common research 3R4F cigarettes (CTRP, Kentucky
University, Lexington, USA) were used for all experiments. The vials containing
animals were capped with a monitoring grid to allow the cigarette smoke to diffuse
into the vial. For long-term smoke experiments, L2 larvae were exposed to smoke
thrice a day for 30 minutes each, on two consecutive days. For heavy smoke
experiments, L3 larvae were exposed to the smoke of 2 cigarettes for 45 min, which
led to about 35-50% mortality. To study the effects of hypoxia on the activity of
JAK/STAT signaling pathway, larvae were exposed to long-term and short-term
hypoxia separately. For long-term hypoxia experiments, L2 larvae were exposed to
5% oxygen three times a day for 30 min each, on two consecutive days. L3 larvae were
exposed to 1% oxygen once for 5 hours for short-term hypoxia experiments. At least
20 animals in 4 vials were investigated in each group. Cigarette smoke and hypoxia
exposure of adults was performed as described earlier [19] .
526
527
Immunohistochemistry of *Drosophila* trachea
Larvae were dissected by ventral filleting and fixed in 4% paraformaldehyde for 30
min. Embryos were staged according to Campos-Ortega and Hartenstein [58] and
fixed in 4% formaldehyde for 30 min. Immunostaining was performed following
standard protocols as described earlier[59, 60]. GFP signals were amplified by
immunostaining with polyclonal rabbit anti-GFP (used at 1:500, Sigma-Aldrich, Merck

KGaA, Darmstadt, Germany, SAB4301138). 40-1a (used at 1:50, DSHB, Iowa City, USA)
was used to detect Beta-galactosidase. Coracle protein was detected with a
monoclonal mouse anti-coracle antibody (used at 1:200, DSHB, Iowa City, USA,
C566.9). Armadillo protein was detected with a monoclonal mouse anti-armadillo
antibody (DSHB, US, N2 7A1, used at 1:500). A monoclonal rabbit Cleaved Drosophila
Dcp1 (used at 1:200, Cell Signaling, Frankfurt/M, Germany, #9578) was used to detect
apoptotic cells. Secondary antibodies used were: Cy3-conjugated goat-anti-mouse,
Cy3-conjugated goat-anti-rabbit, Alexa488-conjugated goat-anti-mouse (used at
1:500, Jackson Immunoresearch, Dianova, Hamburg, Germany), Alexa488-conjugated
goat-anti-rabbit (used at 1:500, Cell signaling, Frankfurt/M, Germany, #9578). Tracheal
chitin was stained with 505 star conjugated chitin-binding probe (NEB, Frankfurt/M,
Germany, used at 1:300). Nuclei were stained with 4',6-Diamidino-2-Phenylindole,
Dihydrochloride (DAPI) (Roth, Karlsruhe, Germany, 6843). 30 specimens were
investigated in each group. Specimens were analyzed and digital images were
captured using either a confocal microscope (Zeiss LSM 880, Oberkochen, Germany)
or a conventional fluorescence microscope (ZEISS Axio Imager Z1, Zeiss, Oberkochen,
Germany).
551
RNA isolation and RNA sequencing
For the gene expression analysis of 3rd instar larvae trachea, animals were dissected
in cold PBS, and the isolated trachea were transferred to RNA Magic (BioBudget,
Krefeld, Germany) and processed essentially as described earlier [19] with slight
modifications. The tissue was homogenized using a Bead Ruptor 24 (BioLab
products, Bebensee, Germany), and the RNA was extracted with the PureLink RNA
Mini Kit (Thermo Fisher, Waltham, MA, USA) for phase separation with the RNA
Magic reagent. An additional DNase treatment was performed following the on-
column PureLink DNase treatment protocol (Thermo Fisher, Waltham, MA, USA).
Sequencing libraries were constructed using the TruSeq stranded mRNA kit (Illumina,
San Diego, USA) and 50 bp single-read sequencing was performed on an Illumina
HiSeq 4000 with 16 samples per lane. Resulting sequencing reads were trimmed for
low-quality bases and adapters using the fastq Illumina filter
(http://cancan.cshl.edu/labmembers/gordon /fastq_illumina_filter/) and cutadapt
(version 1.8.1)[61]. Transcriptomics analysis, including gene expression values and
differential expression analysis, was done using CLC Genomics Workbench. The
detailed protocols can be obtained from the CLC Web site (http://www.
clcbio.com/products/clc-genomics-workbench). *Drosophila melanogaster* reference
genome (Release 6)[62] was used for mapping in this research. RNAseq data are
deposited in the GEO database (GSE212132). Mouse and human orthologs of the
Drosophila DEGs were identified using the DIOPT program package[32].
The transcription factor binding site enrichment and the Gene Ontology enrichment
analyses of the differentially expressed genes were carried out using Pscan

(http://159.149.160.88/pscan/) and online GO enrichment analysis
(http://geneontology. org/), respectively. We chose -450- 50 bases of the annotated
transcription start site of the genes as the transcription factor binding sites for
enrichment analysis. The analysis used the TFBS matrices available in the JASPAR
databases (version Jaspar 2018_NR). Data were visualized using the Circos software
(http://circos.ca/software/).
581

Statistics and reproducibility
We did not use statistical methods to predetermine sample sizes, but the sample
sizes used in this study are similar to or higher than those used in previous studies
[29, 63]. Specific approaches to randomly allocate samples to groups were not used,
and the experiments were not performed in a blinded design. No data were excluded
from the analysis. Prism (GraphPad version 7) was used for statistical analyses, and
the corresponding tests used are listed in the figure legends.
589

Acknowledgments

We thank Britta Laubenstein and Christiane Sandberg for excellent technical
assistance, Maria Leptin and Norbert Perrimon for flies.

Funding

This work was funded by the DFG as part of CRC 1182 and by the CLSM (INST 257/591-1
FUGG). Xiao Niu received funds from the Chinese Scholarship Council and Weifang
Medical University, and Leizhi Shi received funds from the Linyi People's Hospital.
598

599 # Data availability

600 RNAseq data are deposited in the GEO database (GSE212132). Any additional
601 information required to reanalyze the data reported in this paper is available from
602 the lead contact upon request.
603

References

1.   Di YP, Kuhn JM, Mangoni ML: **Lung antimicrobial proteins and peptides: from host defense to**
therapeutic strategies. *Physiol Rev* 2024, **104**(4):1643–1677.

2.   Bao Y, Chen Q, Shi B, Zhu X: **The epithelial barrier: a Janus-faced regulator in allergic airway**
diseases from defense to inflammation. *Frontiers in Allergy* 2025, **Volume 6 - 2025**.
3.   Whitsett JA, Alenghat T: **Respiratory epithelial cells orchestrate pulmonary innate immunity**.
Nature immunology 2015, **16**(1):27–35.
4.   Hewitt RJ, Lloyd CM: **Regulation of immune responses by the airway epithelial cell landscape**.
Nature Reviews Immunology 2021, **21**(6):347–362.
5.   Brightling CE, Marone G, Aegerter H, Chanez P, Heffler E, Pavord ID, Rabe KF, Uller L, Dorscheid
D: **The epithelial era of asthma research: knowledge gaps and future direction for patient care**.
Eur Respir Rev 2024, **33**(174).
6.   Brightling CE, Bafadhel M, Han MK, Papon JF, Rabe KF, Rogliani P, Singh D: **A translational view**
of airway epithelial dysfunction in COPD. *Eur Respir Rev* 2025, **34**(178).
7.   He Q, Li P, Han L, Yang C, Jiang M, Wang Y, Han X, Cao Y, Liu X, Wu W: **Revisiting airway**
epithelial dysfunction and mechanisms in chronic obstructive pulmonary disease: the role of
mitochondrial damage. *Am J Physiol Lung Cell Mol Physiol* 2024, **326**(6):L754–l769.
8.   Kiu H, Nicholson SE: **Biology and significance of the JAK/STAT signalling pathways**. *Growth*
Factors 2012, **30**(2):88–106.
9.   Villarino AV, Kanno Y, O'Shea JJ: **Mechanisms and consequences of Jak-STAT signaling in the**
immune system. *Nat Immunol* 2017, **18**(4):374–384.
10. Yew-Booth L, Birrell MA, Lau MS, Baker K, Jones V, Kilty I, Belvisi MG: **JAK-STAT pathway**
activation in COPD. *Eur Respir J* 2015, **46**(3):843–845.
11. Hu X, Li J, Fu M, Zhao X, Wang W: **The JAK/STAT signaling pathway: from bench to clinic**. *Signal*
Transduct Target Ther 2021, **6**(1):402.
12. Tadokoro T, Wang Y, Barak LS, Bai Y, Randell SH, Hogan BLM: **IL-6/STAT3 promotes regeneration**
of airway ciliated cells from basal stem cells. *Proceedings of the National Academy of Sciences*
2014, **111**(35):E3641–E3649.
13. Major J, Crotta S, Llorian M, McCabe TM, Gad HH, Priestnall SL, Hartmann R, Wack A: **Type I and**
III interferons disrupt lung epithelial repair during recovery from viral infection. *Science* 2020,
369(6504):712–717.
14. Jin H, Ciechanowicz AK, Kaplan AR, Wang L, Zhang PX, Lu YC, Tobin RE, Tobin BA, Cohn L, Zeiss CJ
et al: **Surfactant protein C dampens inflammation by decreasing JAK/STAT activation during**
lung repair. *Am J Physiol Lung Cell Mol Physiol* 2018, **314**(5):L882–L892.
15. Makris S, Paulsen M, Johansson C: **Type I Interferons as Regulators of Lung Inflammation**. *Front*
Immunol 2017, **8**:259.
16. Pandey UB, Nichols CD: **Human disease models in Drosophila melanogaster and the role of the**
fly in therapeutic drug discovery. *Pharmacol Rev* 2011, **63**(2):411–436.
17. Roeder T, Isermann K, Kallsen K, Uliczka K, Wagner C: **A Drosophila asthma model - what the fly**
tells us about inflammatory diseases of the lung. *Adv Exp Med Biol* 2012, **710**:37–47.
18. Bossen J, Uliczka K, Steen L, Neugebauer P, Mai MM, Fink C, Stracke F, Heine H, Roeder T: **Driver**
mutations in major lung cancer oncogenes can be analyzed in Drosophila models. *ALTEX* 2021,
38(2):235–244.

19. Prange R, Thiedmann M, Bhandari A, Mishra N, Sinha A, Hasler R, Rosenstiel P, Uliczka K, Wagner
C, Yildirim AO *et al*: **A Drosophila model of cigarette smoke induced COPD identifies Nrf2**
signaling as an expedient target for intervention. *Aging (Albany NY)* 2018, **10**(8):2122–2135.
20. Levine BD, Cagan RL: **Drosophila Lung Cancer Models Identify Trametinib plus Statin as**
Candidate Therapeutic. *Cell Rep* 2016, **14**(6):1477–1487.
21. Roeder T, Isermann K, Kabesch M: **Drosophila in asthma research**. *Am J Respir Crit Care Med*
2009, **179**(11):979–983.
22. Brown S, Hu N, Hombria JC: **Identification of the first invertebrate interleukin JAK/STAT**
receptor, the Drosophila gene domeless. *Curr Biol* 2001, **11**(21):1700–1705.
23. Perrimon N, Mahowald AP: **l(1)hopscotch, A larval-pupal zygotic lethal with a specific maternal**
effect on segmentation in Drosophila. *Dev Biol* 1986, **118**(1):28–41.
24. Powers N, Srivastava A: **JAK/STAT signaling is involved in air sac primordium development of**
Drosophila melanogaster. *FEBS Lett* 2019, **593**(7):658–669.
25. Zeidler MP, Bausek N: **The Drosophila JAK-STAT pathway**. *JAKSTAT* 2013, **2**(3):e25353.
26. Arbouzova NI, Zeidler MP: **JAK/STAT signalling in Drosophila: insights into conserved regulatory**
and cellular functions. *Development* 2006, **133**(14):2605–2616.
27. McGuire SE, Mao Z, Davis RL: **Spatiotemporal gene expression targeting with the TARGET and**
gene-switch systems in Drosophila. *Sci STKE* 2004, **2004**(220):pl6.
28. Wagner C, Isermann K, Roeder T: **Infection induces a survival program and local remodeling in**
the airway epithelium of the fly. *FASEB J* 2009, **23**(7):2045–2054.
29. Wagner C, Uliczka K, Bossen J, Niu X, Fink C, Thiedmann M, Knop M, Vock C, Abdelsadik A, Zissler
UM *et al*: **Constitutive immune activity promotes JNK- and FoxO-dependent remodeling of**
Drosophila airways. *Cell Rep* 2021, **35**(1):108956.
30. Bossen J, Knop M, Niu X, Thiedmann M, Prange R, Tahanzadeh N, Franzenburg S, Bruchhaus I,
Heine H, Roeder T: **Chronic airway inflammation in Drosophila lacking the A20-like protein**
Trabid. *Front Immunol* 2025, **16**:1564386.
31. Wagner C, Isermann K, Fehrenbach H, Roeder T: **Molecular architecture of the fruit fly's airway**
epithelial immune system. *BMC Genomics* 2008, **9**:446.
32. Hu Y, Flockhart I, Vinayagam A, Bergwitz C, Berger B, Perrimon N, Mohr SE: **An integrative**
approach to ortholog prediction for disease-focused and other functional studies. *BMC*
Bioinformatics 2011, **12**:357.
33. El-Husseini ZW, Khalenkow D, Lan A, van der Molen T, Brightling C, Papi A, Rabe KF, Siddiqui S,
Singh D, Kraft M *et al*: **An epithelial gene signature of trans-IL-6 signaling defines a subgroup of**
type 2-low asthma. *Respiratory Research* 2023, **24**(1):308.
34. Herrera SC, Bach EA: **JAK/STAT signaling in stem cells and regeneration: from Drosophila to**
vertebrates. *Development* 2019, **146**(2):dev167643.
35. Liongue C, O'Sullivan LA, Trengove MC, Ward AC: **Evolution of JAK-STAT Pathway Components:**
Mechanisms and Role in Immune System Development. *PLOS ONE* 2012, **7**(3):e32777.
36. Fortelny N, Farlik M, Fife V, Gorki A-D, Lassnig C, Maurer B, Meissl K, Dolezal M, Boccuni L, Ravi
Sundar Jose Geetha A *et al*: **JAK-STAT signaling maintains homeostasis in T cells and**
macrophages. *Nature Immunology* 2024, **25**(5):847–859.

37.   D'Amico S, Shi J, Martin BL, Crawford HC, Petrenko O, Reich NC: **STAT3 is a master regulator of**
epithelial identity and KRAS-driven tumorigenesis. *Genes Dev* 2018, **32**(17-18):1175–1187.
38.   Piairo P, Moura RS, Baptista MJ, Correia-Pinto J, Nogueira-Silva C: **STATs in Lung Development:**
Distinct Early and Late Expression, Growth Modulation and Signaling Dysregulation in
Congenital Diaphragmatic Hernia. *Cell Physiol Biochem* 2018, **45**(1):1–14.
39.   Whitsett JA, Alenghat T: **Respiratory epithelial cells orchestrate pulmonary innate immunity**.
Nat Immunol 2015, **16**(1):27–35.
40.   Pernis AB, Rothman PB: **JAK-STAT signaling in asthma**. *J Clin Invest* 2002, **109**(10):1279–1283.
41.   Gao H, Guo RF, Speyer CL, Reuben J, Neff TA, Hoesel LM, Riedemann NC, McClintock SD, Sarma
JV, Van Rooijen N *et al*: **Stat3 activation in acute lung injury**. *J Immunol* 2004, **172**(12):7703–
7712.
42.   Milara J, Ballester B, Morell A, Ortiz JL, Escrivá J, Fernández E, Perez-Vizcaino F, Cogolludo A,
Pastor E, Artigues E *et al*: **JAK2 mediates lung fibrosis, pulmonary vascular remodelling and**
hypertension in idiopathic pulmonary fibrosis: an experimental study. *Thorax* 2018, **73**(6):519–
529.
43.   Liu RY, Zeng Y, Lei Z, Wang L, Yang H, Liu Z, Zhao J, Zhang HT: **JAK/STAT3 signaling is required for**
TGF-β-induced epithelial-mesenchymal transition in lung cancer cells. *Int J Oncol* 2014,
44(5):1643–1651.
44.   Traves PG, Murray B, Campigotto F, Galien R, Meng A, Di Paolo JA: **JAK selectivity and the**
implications for clinical inhibition of pharmacodynamic cytokine signalling by filgotinib,
upadacitinib, tofacitinib and baricitinib. *Ann Rheum Dis* 2021, **80**(7):865–875.
45.   Hu X, li J, Fu M, Zhao X, Wang W: **The JAK/STAT signaling pathway: from bench to clinic**. *Signal*
Transduction and Targeted Therapy 2021, **6**(1):402.
46.   Gadina M, Johnson C, Schwartz D, Bonelli M, Hasni S, Kanno Y, Changelian P, Laurence A, O'Shea
JJ: **Translational and clinical advances in JAK-STAT biology: The present and future of jakinibs**. *J*
Leukoc Biol 2018, **104**(3):499–514.
47.   Roeder T, Isermann K, Kabesch M: **Drosophila in Asthma Research**. *American Journal of*
Respiratory and Critical Care Medicine 2009, **179**(11):979–983.
48.   Georas SN, Donohue P, Connolly M, Wechsler ME: **JAK inhibitors for asthma**. *J Allergy Clin*
Immunol 2021, **148**(4):953–963.
49.   Bach EA, Ekas LA, Ayala-Camargo A, Flaherty MS, Lee H, Perrimon N, Baeg GH: **GFP reporters**
detect the activation of the Drosophila JAK/STAT pathway in vivo. *Gene Expr Patterns* 2007,
7(3):323–331.
50.   Brand AH, Perrimon N: **Targeted gene expression as a means of altering cell fates and**
generating dominant phenotypes. *Development* 1993, **118**(2):401–415.
51.   Liu L, Johnson WA, Welsh MJ: **Drosophila DEG/ENaC pickpocket genes are expressed in the**
tracheal system, where they may be involved in liquid clearance. *Proc Natl Acad Sci U S A* 2003,
100(4):2128–2133.
52.   Harrison DA, Binari R, Nahreini TS, Gilman M, Perrimon N: **Activation of a Drosophila Janus**
kinase (JAK) causes hematopoietic neoplasia and developmental defects. *EMBO J* 1995,
14(12):2857–2865.

53.   Fink C, Hoffmann J, Knop M, Li Y, Isermann K, Roeder T: **Intestinal FoxO signaling is required to**
733   **survive oral infection in Drosophila**. *Mucosal Immunol* 2016, **9**(4):927–936.
734   54.   Bosch JA, Tran NH, Hariharan IK: **CoinFLP: a system for efficient mosaic screening and for**
735   **visualizing clonal boundaries in Drosophila**. *Development* 2015, **142**(3):597–606.
736   55.   Chen F, Krasnow MA: **Progenitor outgrowth from the niche in Drosophila trachea is guided by**
737   **FGF from decaying branches**. *Science* 2014, **343**(6167):186–189.
738   56.   Daubeuf F, Frossard N: **Acute Asthma Models to Ovalbumin in the Mouse**. *Curr Protoc Mouse*
739   *Biol* 2013, **3**(1):31–37.
740   57.   Baligar P, Pokhrel S, Mukhopadhyay A: **Experimental Liver Fibrosis and Intrasplenic**
741   **Transplantation of CD45+ Bone Marrow Cells**. *Bio-protocol* 2016, **6**(20):e1972.
742   58.   Campos-Ortega JA, Hartenstein V: **The Embryonic Development of Drosophila melanogaster**:
743   Springer-Verlag Berlin Heidelberg; 1997.
744   59.   Jeon M, Nguyen H, Bahri S, Zinn K: **Redundancy and compensation in axon guidance: genetic**
745   **analysis of the Drosophila Ptp10D/Ptp4E receptor tyrosine phosphatase subfamily**. *Neural Dev*
746   2008, **3**:3.
747   60.   Levi BP, Ghabrial AS, Krasnow MA: **Drosophila talin and integrin genes are required for**
748   **maintenance of tracheal terminal branches and luminal organization**. *Development* 2006,
749   **133**(12):2383–2393.
750   61.   Martin M: **Cutadapt Removes Adapter Sequences From High-Throughput Sequencing Reads**.
751   *EMBnet.journal* 2011, **17**:10–12.
752   62.   Hoskins RA, Carlson JW, Wan KH, Park S, Mendez I, Galle SE, Booth BW, Pfeiffer BD, George RA,
753   Svirskas R *et al*: **The Release 6 reference sequence of the Drosophila melanogaster genome**.
754   *Genome Res* 2015, **25**(3):445–458.
755   63.   Proske A, Bossen J, von Frieling J, Roeder T: **Low-protein diet applied as part of combination**
756   **therapy or stand-alone normalizes lifespan and tumor proliferation in a model of intestinal**
757   **cancer**. *Aging (Albany NY)* 2021, **13**(21):24017–24036.

760   Table 1: Genes that were highly upregulated in response to *Hop.CA* overexpression.

Name	Maximum group mean	Fold change	FDR p-value
CG16772	11.85	135.44	0
hop	286.06	35.43	0
CG14570	42.21	35.37	0
CG14147	55.31	32.88	0
Hsp70Bb	161.51	29.99	0
Ubx	200.46	23.51	0
pre-mod(mdg4)-I	1.12	18.67	2.22648E-06
IM14	11.46	15.14	3.80132E-07
IM4	26.43	14.97	1.74645E-12

CG14569	1576.88	14.73	0
IM2	7.26	14.47	2.01792E-06
CG5791	1.79	13.80	2.90502E-05
CG9121	3.41	11.54	0
IM1	7.16	10.97	7.52122E-05
CG11413	35.70	10.64	3.44242E-07
CG16710	2.13	10.32	1.72352E-07
CG31960	267.26	10.05	0
CG16789	1.45	9.46	5.58576E-09
CG6034	6.92	8.47	4.7943E-11
Ace	2.06	8.19	0
CG13059	9.90	7.58	1.33893E-06
CG31809	8.64	7.39	0
CG7203	9.41	7.21	8.66989E-09
Vha14-2	2.51	7.05	2.31024E-07
wb	7.35	6.69	0
Drs	8166.47	6.64	0
CG5778	3.20	6.61	0.000294835
CG3457	2.69	6.34	2.38755E-06
Lcp65Ad	9.93	5.79	2.58249E-08
lectin-22C	29.06	5.67	6.88538E-06
fuss	3.97	5.67	0
CG43333	2.39	5.63	1.48392E-09
CG18336	2.96	5.49	5.52167E-05
CG32563	3.06	5.35	0.000696584
FASN2	1.65	5.34	2.95545E-10
Cyp4d21	45.70	5.31	0
Obp57d	1.80	5.28	0.000102926
CG17560	2.31	5.22	0.000766701
CG5888	68.16	5.20	0

761    $p < 1\times10^{-4}$, fold change > 5, maximum group mean > 1. The list is restricted to 39 genes.
762

763   Table 2: Transcription factor-binding site motifs enriched in 41 highly upregulated
764   genes.

Matrix ID	Matrix name	P-value
MA0023.1	dl (var.2)	0.000443261
MA0022.1	dl	0.00407113

MA0532.1	STAT92E	0.00794732
MA0450.1	hkb	0.00811969
MA0197.2	nub	0.0154201
MA0242.1	Bgb::run	0.0371696
MA0204.1	Six4	0.0443521
MA0444.1	CG34031	0.0488169

765   A total of 133 transcription factor profiles were used.
766
767
768

Figures

Figure 1: The activation of the JAK/STAT signaling pathway in the respiratory system
of *Drosophila*. The activation of the JAK/STAT signaling pathway was detected using
a *STAT92E-GFP* reporter (A), which includes *LacZ* expression under control of *btl-Gal4*
(*btl-Gal4, STAT92E-GFP; UAS-LacZ.nls*). (B-E) STAT92E activity pattern (green) in the
larval trachea. JAK/STAT activity is present in most tracheal parts (B). Only the
posterior part of the trachea (dorsal trunk (DT) of Tr 10, white dashed line in C)
showed decreased JAK/STAT signaling activity (and high *btl* expression; red)
compared to other tracheal epithelial cells. Conversely, some regions, like the
spiracular branch (SB; D, white arrow) and the dorsal branch (DB) in Tr 2 (E, white
arrow), showed increased JAK/STAT signaling activity compared to the tracheal
epithelial cells with *btl* expression. (F) STAT92E activity pattern in the adult trachea.
Scale bar: 50 μm.

783
Figure 2: Blocking JAK/STAT signaling induces epithelial cell apoptosis.
JAK/STAT signaling was blocked in the trachea by expressing *UAS-Dome.DN* under the
control of *btl-Gal4;UAS-GFP* (A). (B) The developmental viability in response to
JAK/STAT blockade is indicated by the percentage of embryos that hatched. (****p <
0.0001 by Student's t-test). (C) Blockade of JAK/STAT signaling induced major
impairments of the embryonic tracheal structure (arrows). (D) Surviving larvae
showed visible defects indicated by liquid-filled regions (arrows). (E) Blocking of the
JAK/STAT pathway by expressing *Dome.DN* in the trachea led to changes in cell
morphology (E' and E'', green arrows) and subsequent disintegration (E''', green

arrow). Scale bar: 20 μm. (F-G) Fluorescence micrographs of the tracheal cells of
control *vvl-coin.ts* larvae (F), and the tracheal cells of *vvl-coin.ts>Dome.DN* larvae (G)
stained for cleaved Dcp-1 (red, on day 2 after induction). Scale bar: 50 μm. (H)
Lifespan analysis under control conditions and cigarette smoke exposure (CS).
Median lifespan is indicated in brackets; n=76-103; Log-rank (Mantel-Cox) test. (I, J)
Short-term (H; n=6; unpaired t-test) and long-term (I; n=8-9; Mann-Whitney test)
hypoxia treatment. * p < 0.05, *** p < 0.001, **** p < 0.0001.
800
801
802

Figure 3: The JAK/STAT signaling pathway in the trachea was activated by external stimuli. (A, C) Fluorescence micrographs of the trachea from *STAT92E-GFP* larvae (A) and adults (C) under control conditions and exposed to external stimuli, which included hypoxia, cold air (cold), and cigarette smoke exposure (CSE). The external stimuli induced the activity of STAT92E, which could be observed and quantified through fluorescent intensity (A' and C'). (B, D) Fluorescence micrographs of the trachea from *upd3-Gal4; UAS-GFP* larvae (B) and adults (D) under control conditions and exposed to external stimuli. The expression of *upd3* induced by the external stimuli could be observed and quantified through fluorescent intensity (B' and D'). The fluorescence signal induced by hypoxia in larval trachea was quantified in flies

815   concurrently driving *upd2*-RNAi (E) or *upd3*-RNAi (F). ** $p < 0.01$, *** $p < 0.001$, ****
816    $p < 0.0001$ by Student's t-test. Scale bar: 100 μm.
817

Figure 4: Ectopic JAK/STAT signaling activation in the trachea induced structural

changes. Ectopic activation (*Hop.CA* or *upd3*) of epithelial JAK/STAT signaling was

achieved by ectopic activation of the pathway (A). (B, C) Signaling pathway activation

during development (*btl>Hop.CA* or *btl>upd3*) induced lethality. The percentage of

embryos that hatched (B) and the percentage of larvae that developed into pupae
(C) are shown. (D-E) Incomplete tracheal development during tracheal
morphogenesis. (F, G) Micrographs of DT8 of *vvl-FLP, CoinFLP-Gal4, UAS-EGFP* (*vvl-
coin*)>*Hop.CA* larvae. Scale bar: 20 μm. The thickening of the tracheal epithelium is
observed in those cells that express *Hop.CA* (arrows). A narrowing of the airspace
was observed in the affected epithelial cells (G; dotted lines indicate the airway
boundaries), whereas unaffected cells remained inconspicuous (F). (H) Ectopic
activation of the pathway either by upd3 or Hop.CA overexpression for 2 days or 4
days using the Target system by switching the temperature in the larval stage from
19°C to 30°C. Epithelial cells express GFP. Scale bar: 20 μm. (I) Quantification of the
epithelial thickness (DT8) in response to upd3-, Hop.CA-, or Dome.DN overexpression
for different times. (K) Quantitative evaluation of epithelial thicknesses of affected
regions of the trachea following repeated chronic smoke exposure. (L) Evaluation of
the thickening phenotype using the mosaic-inducing vvl system (green are affected
cells), unaffected cells remained normal. (M-P) Localization of the membrane-
associated proteins armadillo (arm) and Coracle (cora) in control epithelial cells and
those showing strong JAK/STAT activation. (M, N) Armadillo staining showed
accumulation in the cell of the immunoreactivity (arrow). (O, P) Similar localization
errors are shown for Coracle staining (arrow). ns = not significant, * $p < 0.05$, ** $p <$
0.01, *** $p < 0.001$, **** $p < 0.0001$; Student's t-test.
843
844

Figure 5: Analysis of genes differentially expressed between control trachea and
those ectopically activated by Hop.CA overexpression. (A) Schematic overview of
the DEGs that are up- or downregulated in response to Hop.CA expression. (B) Venn
diagram showing DEGs in response to Hop.CA overexpression compared with those
following Trabid-KO, tracheal infection, or ectopic activation of the IMD pathway
(PGRP-LC overexpression). (C) Using DIOPT to identify mouse orthologs of the
Hop.CA DEGs from (A), these orthologs were compared with DEGs from lungs
subjected to acute and chronic asthma models. (D) Key chemokines and chemokine
receptors induced by OVA aerosol are involved in the JAK/STAT signaling pathway.
(E) Analysis of the genes commonly regulated between Hop. CA-induced genes and
those from the acute and chronic asthma models. (F) DIOPT-assisted identification of
transformation of Hop.CA DEGs into human orthologs, which were compared with
genes differentially expressed in human airway epithelial cells exposed to IL-6 trans-
signaling. The common genes were analyzed regarding GO categories in the
biological processes group and through a string analysis of these data (G). *** $p <$
001; Fisher's exact test.
862
863

Figure 6: Pharmacological rescue of the epithelial thickening phenotype and concluding model of the JAK/STAT action in airway epithelial cells.

(A, B) Changes in tracheal epithelial thickness could be rescued by applying specific JAK inhibitors. (A) Microscopy images of the tracheal epithelium (3rd instar larvae) of Control (*btl.ts>w*^{1118}) compared to activated JAK/STAT signaling pathway (*btl.ts>Hop.CA*). Scale bar: 50 μm. (B) Quantification of epithelial thickness (highlighted in red) in JAK inhibitor treatment groups compared to the DMSO control. ** $p < 0.01$; Student's t-test. (C) Schematic characterization of JAK/STAT signaling under homeostatic conditions and in response to chronic stress leading to or prolonged overactivation (D).