Social interactions shape antiviral resistance outcomes in poliovirus via eco-evolutionary feedback

Alexander J. Robertson[1], Benjamin Kerr[2,*], Alison F. Feder[3,4,*]

[1] Molecular and Cellular Biology Graduate Program, University of Washington, Seattle, WA 98103

[2] Department of Biology, University of Washington, Seattle, WA 98103

[3] Department of Genome Sciences, University of Washington, Seattle, WA 98103

[4] Herbold Computational Biology Program, Fred Hutchinson Cancer Center, Seattle, WA 98109

[*] To whom correspondence should be addressed. E-mail: kerrb@uw.edu, affeder@uw.edu

Abstract

Antiviral resistance evolution poses a major obstacle for controlling viral infections. A promising strategy is to target shared viral proteins that allow drug susceptible viruses to sensitize resistant ones during cellular coinfection, muting selection for resistance. Pocapavir, a poliovirus capsid inhibitor, employs this sociovirological strategy. While susceptible viruses significantly suppressed resistance in the presence of pocapavir in cell culture, a pocapavir clinical trial observed widespread resistance evolution and limited improvements to clearance times. To reconcile these findings, we present an intra-host eco-evolutionary model of poliovirus in the presence of pocapavir, which reproduces both the potent interference observed *in vitro* and the resistance emergence seen in patients. In the short term, our model predicts that a high density of susceptible viruses sensitizes resistant ones to pocapavir, mirroring cell culture results. However, over multiple replication cycles, pocapavir's high potency collapses viral density, which reduces coinfection and allows resistance to evolve as observed in the clinical trial. Since coinfection is essential to suppress resistance, enabling greater survival of susceptible viruses could offer therapeutic advantages. Counterintuitively, we demonstrate that this can be achieved by *lessening* antiviral potency, which can limit resistance evolution while also maintaining a low viral load. These findings suggest that antivirals that rely on viral social interaction must balance immediate neutralization with the preservation of future coinfection, yielding more sustained inhibition. Explicitly considering the eco-evolutionary feedback encompassing viral density, social phenotypes and absolute fitness not only provides new insights into designing effective therapies but also illuminates viral evolutionary dynamics more broadly.

Introduction

Resistance evolution can undermine otherwise successful antimicrobial treatments if drug application permits microbes carrying resistance alleles to expand and prevent population extinction. Implicit in these dynamics is that selection for phenotypic resistance also selects genotypic resistance, which permits the trait to carry forward into future generations. While such an assumption is often valid in bacterial populations (**Figure 1A-C**) [1, 2], in viruses, the association between genotype and phenotype can be more complicated [3, 4, 5]. For example, both cellular coinfection or *de novo* mutation during viral replication can result in distinct genotypes occupying the same cell. These genotypes may share intracellular protein products, and as a result, a particular genotype may be associated, fully or partially, with the protein products of a different genotype (**Figure 1D&E**) [6].

This so-called "phenotypic mixing" has important implications for resistance evolution [5, 7]. First, the disassociation between genotype and phenotype can impair selection for resistant genotypes. Consider a cell that contains both treatment-resistant and susceptible genotypes. If resistant genotypes do not associate with resistant protein products, those genotypes may not survive drug application, limiting their contribution to future generations (**Figure 1F**). Second, phenotypic mixing can limit the creation of resistant phenotypes themselves. If a drug targets an oligomeric protein product (e.g. a capsid inhibitor that targets the capsid), susceptible proteins can act in

Figure 1: **Genotype-phenotype associations can limit resistance evolution.** In bacterial populations, resistance-conferring genotypes exist at low frequencies prior to treatment (**A**). Because genotypes are directly associated with the phenotypes they encode (**B**), drug application that selectively favors the survival of resistant phenotypes drives the expansion of resistant genotypes (**C**). In viruses, multiple genotypes (i.e., resistant and susceptible) may co-infect the same cell via mutation or superinfection and the protein products they encode can mix intracellularly (**D**). The resulting chimeric phenotypes can be associated with either genotype (**E**) and even those phenotypes that are partially composed of resistant proteins may be susceptible to drug neutralization (**F**). Throughout the figure, red and blue represent resistant and susceptible variants, respectively, and yellow triangles represent antimicrobials that can bind to susceptible proteins.

a dominant negative fashion to interfere with the phenotypic expression of resistance. Capsid inhibitors may neutralize chimeric capsids composed of both susceptible and resistant subunits if a sufficient number of susceptible subunits are available to bind the drug. If intracellular resistant genotypes are rare (such as after *de novo* mutation of a resistant mutant), cells may produce few or no phenotypically-resistant capsids. The presence of one or both of these factors could present an opportunity to treat viral infections while muting selection for resistance.

Exploiting phenotypic mixing to suppress resistance evolution while treating viral infections partially underlies the promise of the poliovirus capsid inhibitor pocapavir [6, 8]. Pocapavir strongly inhibits poliovirus, but mutations in the genes encoding capsid subunits VP1 or VP3 disrupt drug binding and confer resistance [9, 10]. When resistant viruses are co-cultured with susceptible ones and treated with pocapavir, resistant genomes are packaged in chimeric resistant-susceptible capsids and are neutralized by the drug [6]. As a result, co-cultured resistant viruses have significantly reduced viral titers compared to resistant viruses grown in isolation (**Figure 2A**). Promisingly, pocapavir was used to slow disease progression *in vivo* in poliovirus-infected mice with no detected resistance evolution [6]. However, in a larger placebo-matched human clinical trial where participants received the live attenuated polio vaccine and were treated with pocapavir, the drug both failed to significantly reduce time to viral clearance in three of four placebo-matched groups and resistance evolved in nearly half of pocapavir-treated participants (**Figure 2B**) [11]. While

Figure 2: **Pocapavir-treated poliovirus outcomes diverged between cell culture and clinical trial settings.** (**A**) *In vitro* experiments by Tanner et al. [6] demonstrated that coinfection of drug-resistant (res.) and susceptible (sus.) poliovirus strains suppresses the yield of resistant virus under pocapavir treatment (MOI—multiplicity of infection, the ratio of viruses to host cells, PFU– plaque forming units). (**B**) In the clinical trial reported by Collett et al. [11], pocapavir failed to significantly reduce time to infection clearance compared to a placebo in three of four matched groups of participants administered the live attenuated poliovirus vaccine and resistance was enriched in the pocapavir group. Points represent the clearance dates of individual trial participants and are colored by resistance status (resistance in red, susceptible in blue) and gray boxes indicate dates that were not sampled during the trial (DPI—days post infection).

a subset of pocapavir recipients did clear their virus early without apparent resistance evolution (and pocapavir therapy has been successfully employed in certain compassionate use cases [12, 13]), results from the clinic have been mixed, and, to our knowledge, have not led to additional trials. More generally, this study raises doubts about the therapeutic potential of exploiting phenotypic mixing.

To investigate these conflicting findings, we developed a dynamical model of poliovirus replication and evolution under drug treatment. Surprisingly, we find that a single model can reproduce the seemingly contradictory *in vitro* cell culture and clinical results via its behavior at different viral densities. At high viral density, susceptible viruses mask the phenotype of resistant ones and suppress selection for resistance, as observed in cell culture. However, as successful treatment drives the viral density down, limited intracellular viral interaction restores the standard genotype-phenotype association. At this point, resistant and susceptible genomes associate strongly with their own phenotype, and drugs can efficiently select for genotypic resistance. Counterintuitively, our model suggests that permitting more susceptible viruses to survive drug application can better suppress resistance evolution and lead to smaller viral population sizes over time. This study provides the first theoretical framework for evaluating viral evolutionary responses to therapies targeting socially-encoded resistance, serves as a guide for the development of novel antimicrobials and dosing strategies, and highlights the importance of emergent dynamical responses when exploiting virus-virus interactions in medicine.

Results

Poliovirus eco-evolutionary model

We developed a discrete-generation dynamical model that tracks poliovirus genotypes and phenotypes over multiple rounds of viral replication and analyzed the model using both deterministic and fully stochastic simulations (see Materials & Methods). Briefly, each generation consists of four steps (**Figure 3A**):

1. *Viral entry into host cells*: resistant and susceptible genomes enter cells as a function of their respective population sizes. Coinfection is more likely if viral population sizes are large relative to the host cell population.

2. *Viral genome replication and mutation*: intracellular viral genomes replicate up to the cell's burst size. Mutation can interconvert resistant and susceptible genotypes.

3. *Capsid formation*: the initial infecting genomes produce a pool of shared capsid subunits. For mixed infections, capsids are formed by randomly drawing 60 subunits from this pool and can be composed of both resistant and susceptible subunits.

4. *Capsid packaging*: newly replicated viral genomes are packaged into assembled capsids in proportion to their intracellular abundances.

Figure 3: **Discrete-time model of poliovirus replication, mutation and survival under pocapavir treatment.** (**A**) We simulate intracellular poliovirus dynamics in four stages: (1) viral entry into host cells, (2) genome replication with mutation, (3) production of capsid subunits, and (4) assembly and packaging of progeny virions. (**B**) Capsid-mediated survival is modeled by culling progeny virions according to their capsid composition. Capsid survival probability as a function of the number of resistant subunits (line) was fit from cell culture experimental data from Tanner et al. [6] (points). Resistant viral yield under different intensities of susceptible virus coinfection shows a density-dependent effect *in vitro* [6] (**C**) and *in silico* under our model (**D**). Throughout the figure, resistant variants are illustrated in red and susceptible variants are illustrated in blue.

Viruses then exit cells and pocapavir can bind and neutralize free virions according to their capsid phenotype. We parametrized viral neutralization rates based on the reduction in viral titers measured experimentally in pocapavir-treated populations of mixed resistant and susceptible cultures (see Materials & Methods—Parameter inference). Capsids composed solely of resistant subunits survive pocapavir application with probability 1, while those composed of fully susceptible subunits survive with probability 4×10^{-4} (**Figure 3B**). Our model recapitulates the observations in Tanner et al. [6] that titers of resistant viruses (specifically, viral genomes encoding resistant subunits) decrease when co-infected alongside susceptible viruses (specifically, viral genomes encoding susceptible subunits) when treated with pocapavir (**Figure 3C&D**).

Resistance suppression is dependent on susceptible virus density

We first assessed the conditions under which pocapavir resistance evolution is suppressed during a single round of replication with pocapavir treatment. Specifically, we measured the change in resistance frequency as a function of the total multiplicity of infection (MOI—a measure of viral density defined as the ratio of the total number of viruses to the total number of infectable host cells). We initialized simulations with the resistant genotype frequency set at $f_{\text{Res}} = 10^{-4}$, consistent

with levels observed in untreated poliovirus populations [6, 10].

At high MOIs (MOI $\approx 10^2$), genotypic resistance increased in frequency by less than 10^{-3} after a single round of replication in the presence of pocapavir (**Figure 4A**). These results are consistent with Tanner et al. [6] and the logic underlying phenotypic mixing. However, this resistance suppression did not extend to populations that were initialized at lower MOIs. At MOI ≤ 1, the resistant genotype frequency increased to more than 12% of the population in a single round of replication. These results can be understood in the context of the MOI controlling the degree of coinfection, and subsequently the strength of the genotype-phenotype association. At high MOIs, coinfection is ubiquitous and the association between phenotypes and rare genotypes breaks down, and rare resistant genomes are wrapped in susceptible capsids (**Figures 4B, S1**). However, at low MOIs, coinfection is rare and genotypes and phenotypes are directly associated (**Figure 4C**), allowing resistant genotypes to directly benefit from their associated phenotype without interference.

We next considered that infections are dynamic processes, and the degree of viral suppression or proliferation in one generation may determine viral density in subsequent generations. If pocapavir treatment drastically reduces the viral density (and consequently the total MOI), this may lead to conditions in which resistance can emerge. We therefore performed an *in silico* serial passaging experiment in which we inoculated cell populations at a high MOI (MOI = 100) with predominantly susceptible viruses (resistant genotype frequency $f_{Res} = 1 \times 10^{-4}$). We allowed viruses to replicate and be neutralized by pocapavir, and seeded the surviving viruses on fresh cell populations over multiple generations.

Consistent with single step experiments at high MOI, the viral population decreased in abundance after one round of replication in the presence of pocapavir (**Figure 4D**). As a result, the surviving viral progeny infected cells at significantly reduced MOI and resistant genomes increased in both frequency and abundance after a second and third round of passaging. At the passage at which the viral population size recovered enough for widespread coinfection, genotypic resistance was more than $100\times$ more common than genotypic susceptibility and susceptible subunits no longer effectively sensitized their resistant counterparts (**Figure S2**). These dynamics can be understood by tracking the change in MOI and resistance frequency on a stepwise phase diagram, initiating simulations across a range of initial total MOIs and resistance frequencies (**Figure 4E**). Initial conditions with low resistance frequencies and sufficiently high MOI lead first to a rapid reduction in the MOI, which then permits increases in the genotypic frequency of resistance. Note, one way to understand this sequence is that the low viral titer output of **Figure 4B** is the low MOI input of **Figure 4C**, enabling the spread of resistant viruses. Regardless of the initial conditions in our deterministic model (MOI > 0, $f_{Res} \in [0, 1]$), resistance becomes the dominant genotype over time. Despite phenotypic mixing effectively suppressing resistance at high MOIs, rapid population contraction in response to pocapavir ultimately undermines that suppression.

Figure 4: **Resistance suppression is MOI-dependent and resistance emerges if bottle-necks do not lead to extinction**. (**A**) Change in resistance frequency in a single generation (Δf_{Res}) depends on the MOI (initial $f_{Res} = 1 \times 10^{-4}$). (**B**) At high MOI, rare resistant genomes are encapsidated by phenotypically susceptible capsids, muting genotypic selection. Although nearly all capsids are phenotypically susceptible, some survive pocapavir administration (a fraction greatly exaggerated for this cartoon), as observed in cell culture. (**C**) At low MOI, viruses singly infect cells, and rare resistant genomes are encapsidated by phenotypically resistant capsids, enabling selection for resistance. (**D**) Over multiple generations, pocapavir treatment transiently reduces viral population size for both resistant and susceptible genomes but leads to viral rebound of a primarily genetically resistant population following low population density. (**E**) Discrete step phase diagram shows the joint change in genotypic resistance frequency and MOI from different initial conditions. Arrows are shortened by 80% to increase legibility. The trajectory from part (**D**) is overlaid in grey. (**F**) Clearance dates from the observed clinical trial and one simulated clinical trial of $n = 93$ viral populations treated with pocapavir. The dashed line indicates the date of the earliest placebo clearance. (**G**) Late and early clearers both experienced drops in resistant and susceptible viral population size with diverging outcomes following the population bottleneck. Throughout the figure, resistant variants are illustrated in red and susceptible variants are illustrated in blue.

Stochastic model replicates clinical trial outcomes

While our deterministic model can explain how resistance to pocapavir may have emerged in clinical trial participants despite phenotypic mixing, it can not recapture the clinical trial observation that a subset of pocapavir recipients clear their infections early with little to no resistance evolution (**Figure 2B**) [11]. To investigate the dynamics driving the clinical trial clearance times, we implemented a stochastic version of the model in which simulated viral elements are drawn from probability distributions at each replication step in a finite host cell population with an immune system that responds to infection (see Materials & Methods—Immune clearance). In brief, we modeled viral immune clearance via a non-specific, ramping innate immune response that removes viruses irrespective of capsid phenotype, and parameterized clearance rate and host cell population size based on the clearance dates in the clinical trial placebo group. We used the stochastic immune model to simulate an *in silico* pocapavir clinical trial by running 93 simulations (representing 93 trial participants) until viral extinction. The infections were initialized with one susceptible virus per host cell, and no resistant viruses. Pocapavir was administered after 24 hours (three rounds of replication, $n = 23$) or 72 hours (nine rounds of replication, $n = 70$), as in the clinical trial.

Our model broadly recaptures the clearance time and resistance evolution outcomes observed in Collett et al. [11]. Specifically, viral populations exhibit a bifurcation of outcomes, in which they clear shortly after pocapavir initiation (< 7 days post infection) with little genotypic resistance, or clear later (≥ 7 days post infection) with widespread genotypic resistance (**Figure 4F**). Analysis of the simulated population trajectories revealed that both early clearers and late clearers experienced sharp population bottlenecks and low MOIs shortly after pocapavir initiation (**Figures 4G, S3**). In late clearers, this bottleneck allowed for rapid, resistance emergence according to the dynamics explored above, while in early clearers, this bottleneck led to stochastic extinction. Our model suggests that the previously unexplained divergent participant outcomes among pocapavir recipients in Collett et al. [11] may be due to stochastic differences at low viral population sizes following pocapavir initiation.

Susceptible dominance has a limited effect on resistance suppression

We hypothesized that pocapavir resistance may evolve in poliovirus due in part to the particular relationship between the number of resistant subunits in a capsid and the virion's fitness. Specifically, when we parameterized the pocapavir neutralization curve using experimental measurements from [6], we found that the largest relative increases in capsid survival probability accompanied the incorporation of the first few resistant subunits (**Figure 3B**). We hypothesized that resistance could be successfully suppressed over longer timescales if more resistant subunits were required to increase the probability of a capsid surviving pocapavir (i.e., the susceptible subunits exerted a greater effect over phenotype than resistant subunits). We therefore repeated our experiments using a neutralization curve in which a virus needed to construct a fully resistant capsid (i.e., 60/60 resistant subunits) before it experienced any survival improvement in the presence of the drug (**Figure S4A**).

Intriguingly, under these conditions, dynamical outcomes were nearly identical to those under the neutralization curve of pocapavir, despite the seemingly more favorable properties. Again, resistance suppression was highly dependent on MOI (**Figure S4B**) and resistance evolved over multiple passages following a drug-induced reduction in viral population size (**Figure S4C**). More broadly, we found that as long as the survival probabilities of fully resistant and fully susceptible capsids were preserved (i.e. the end points of the fitness curve remained the same), a range of monotonically increasing functions describing the dominance interactions between susceptible and resistant subunits resulted in very similar clinical outcomes to pocapavir (**Figure S4D-F**). This consistent result emerges because the rapid contraction of the population following drug application results in most cells being infected by single viral genomes. At that point, interaction properties between resistant and susceptible subunits in the presence of the drug become secondary to the survival probabilities of fully resistant and susceptible virions. This suggests that efforts to find drug formulations in which resistance mutations behave more "recessively" are unlikely to yield substantively different results as long as drug application continues to produce singly-infected cells.

Reduced drug potency can enhance long-term control of resistance and lower viral burden

The critical liability of pocapavir identified above is that effective neutralization of the virus disrupts susceptible viruses' abilities to interfere with resistant phenotypes via coinfection. We hypothesized that increased survival of susceptible viruses would enhance resistance suppression by maintaining higher rates of coinfection over time. Furthermore, since much of the viral burden in a resistance-dominated infection is driven by high viral loads following the emergence of resistance, effectively suppressing resistance could reduce the viral load overall, even if more susceptible viruses survive. We therefore considered the effects of hypothetically less potent drugs (or alternatively, lower doses of pocapavir) that permitted greater degrees of survival by virions with susceptible or partially susceptible capsids (**Figure 5A**, see Materials & Methods—Variation in drug strength).

Reducing drug potency led to smaller increases in the frequency of genotypic resistance in a single round of replication than pocapavir, in line with evolutionary expectations for weaker selective pressures (**Figure S5A**). However, reducing drug efficacy also led to a less rapid decline in the absolute number of susceptible genomes. For example, a drug with a 100-fold reduced efficacy reduced the MOI after a single round of replication to approximately 10, whereas pocapavir reduced the MOI to substantially less than one virus per cell (**Figure 5B**). As a result of this sustained moderate MOI, genotypic resistance increased minimally (up to 0.39% of the population) over the next 6 passages while genotypic resistance under pocapavir reached near 100%. Because resistance remained suppressed, the population did not undergo full viral rebound and the total MOI after 6 passages was approximately 20 times smaller under the $100\times$ less potent drug than pocapavir. This suggests that reducing antiviral potency can, under some circumstances, improve multiple aspects of viral control.

To evaluate the potential clinical implications of reducing antiviral potency, we simulated *in*

Figure 5: **Drugs less potent than pocapavir can better suppress resistance and maintain lower viral loads.** (**A**) We modeled hypothetical drugs 10×, 100× and 1,000× weaker than pocapavir and plot the probability of virion survival as a function of the number of resistant capsid subunits under these drug conditions. (**B**) In deterministic serial passage experiments, the 100× weaker drug maintained a lower total viral population than pocapavir (resistant genomes, red; susceptible genomes, blue). We simulated clinical trials of 100 individuals treated with drugs 1×, 10×, 100×, and 1,000× weaker than pocapavir. Reducing drug potency delayed mean clearance time measured by days post infection (**C**) but had a non-monotonic effect on the sum viral load over the course of the infection (**D**). In both (**C**) and (**D**), dots represent individual simulations and are colored based on the frequency of resistance in the population over the course of the infection (scale in (**D**)). Error bars show mean (red triangle) ± variance. For an expanded range of drug potencies, see **Figure S5 B&C**.

silico clinical trials of 100 individuals treated with reduced potency drugs (**Figure 5C&D**, assessed across a greater range in **Figure S5B&C**). We found that pocapavir had an earlier mean clearance time compared to the less potent hypothetical drugs. Reduced drug potency weakened population bottlenecks, preventing stochastic extinction of viral populations at small population sizes when treated with drugs weaker than pocapavir.

In contrast, reducing drug potency had more complex effects on the frequency of resistance and the sum total viral population size over the course of infection (**Figure 5D**). Reducing drug potency by a factor of 10 resulted in near ubiquitous resistance evolution and high viral population size across trial participants. This reflects that a 10× reduced potency is a strong enough selective pressure to bring viral populations into a regime in which cells are singly infected and resistance can evolve, yet it is not strong enough to create population bottlenecks severe enough for stochastic extinction like pocapavir can. In contrast, reducing drug potency by a factor of 100× created a gradual enough decay during initial drug application that resistant-susceptible coinfection was

durably maintained. While this rate of viral decay was not strong enough to cause stochastic extinction during bottlenecks, the sustained coinfection reduced the rate of resistance evolution and subsequently the total viral load compared to pocapavir. Finally, a very strong reduction of drug potency by a factor of $1,000\times$ selected for almost no resistance evolution, but also did not restrict the viral population relative to an untreated control. This suggests that while reducing drug potency can improve multiple important clinical metrics, not all reductions in drug potency will have favorable effects and there is an optimal balance between preserving susceptible virus and limiting total infection burden.

Discussion

In this paper, we show that a single model of poliovirus population dynamics and genetics can reconcile seemingly divergent outcomes of pocapavir treatment in cell culture [6] and clinical trial settings [11]. Our key insight is that therapeutic strategies that rely on interaction between viruses must account for the demographic effects of therapeutic success. If the long-term efficacy of these therapies depends on the durability of social interactions, lowering viral density through successful treatment can decrease coinfection rates and thus the potential for therapeutically beneficial social interactions between viral genomes and their encoded proteins.

These considerations are important given the growing interest in exploiting social interactions between viral genomes in a therapeutic capacity. In most of these proposed cases, partially or fully-defective viral genomes serve as the therapeutic themselves rather than a sensitizing agent for a different small molecule as we consider here. Perhaps most promisingly, therapeutic interfering particles (TIPs) are replication-incompetent viral mutants that can interfere with their replication-competent counterparts when they co-infect the same cells [14, 15, 16]. Because TIPs expand faster than their targets upon coinfection, their interference capability grows rather than shrinks over time (as long as conditions favoring coinfection endure). Although the full therapeutic potential for TIPs is still an active area of investigation, Pitchai et al. [17] recently demonstrated an effective proof of concept for using TIPs to treat HIV over short timescales. The demographic feedback considered in our model highlights potential challenges with these approaches - over longer periods of time, if TIPs drive replication-competent viruses to near but not complete eradication, TIPs may lose their ability to self-renew and themselves become eliminated. This could ultimately lead to the unencumbered rebound of replication-competent viruses, especially in the case of viruses with a latent stage such as HIV where viruses can reactivate from a reservoir after TIPs have been eliminated.

If social interactions are crucial for the success of these treatments, how can they be maintained in the face of demographic collapse? In the case of TIPs, Weinberger et al. [15] found that *weaker* interference between TIPs with their wild-type counterparts favors the long-term success of TIP therapy [15]. Indeed, all successful demonstrations of TIPs in rodents [18, 19, 20] or mosquitoes [20] allow sufficient wild-type replication to maintain therapeutic pressure in subsequent generations.

In our model of poliovirus and pocapavir, we achieve sustained social interaction through a similar therapeutic intervention—decreasing drug intensity. Modifying drug intensity through dosage or administration frequency may allow more fine-tuned calibration of social interactions than identifying TIPs with the appropriate degree of interference. Similar strategies to prolong drug efficacy by exploiting social interactions are also being explored in non-virology settings. For example, in drug-treated bacterial [2, 1] and cancer [21, 22] populations, moderate-dose or pulsed strategies can prolong competition between susceptible and resistant cells, thereby lowering overall population size over time and delaying treatment failure.

Although these treatment strategies reckon with similar demographic dynamics across systems, they may require different medical and public health considerations. In our model, decreased drug potency can slow the emergence of resistance and lower viral load. However it simultaneously prolongs average infection clearance time by reducing stochastic early viral extinction. Similarly, in cancer adaptive therapy, alternating therapeutic pressure and drug holidays reduces tumor burden and increases the time to clinical progression, but sacrifices the possibility of eradication and ultimately only delays resistance evolution [21]. While both treatment strategies therefore affect individual patient clinical outcomes, infectious diseases carry the added risk of onward transmission. Decreased viral load from partial population control may result in reduced poliovirus infectivity as it does in SARS-CoV-2 [23, 24] and HIV [25], but slower clearance times may counterbalance this effect by increasing the number of contacts exposed. Resistance evolution also becomes a more acute concern than in cancer therapies, as resistant variants can transmit to other people and undermine future drug effectiveness. To determine if the decreased viral load and enhanced resistance suppression of a less potent drug outweighs the concomitant increase in the infectious period, an epidemiological model incorporating the intra-host dynamics we explore could prove valuable.

While intracellular interaction is a liability to viruses in our model, these same interactions may be beneficial or even critical to viruses in other settings. For example, multipartite viruses package their genome segments into separate capsids and therefore require cellular coinfection of virions carrying distinct segments to successfully replicate [26]. Coinfection may also be critical for influenza viruses, as individual virions often carry incomplete genomes [27, 28]. In less extreme cases, increased social interactions can allow viral populations to more effectively explore fitness landscapes [4, 3, 7] and exchange genes [29, 30]. When considering viral evolution in these settings, positive demographic feedback loops may need to be considered.

A central assumption in our model is that the number of viruses per host cell is the primary driver of coinfection rate. While this is a common assumption [31, 32, 33, 4], diverse viral behaviors can modulate coinfection. For example, the first virus infecting a cell can prevent others from entering in a process called superinfection exclusion [34]. Conversely, *en bloc* transmission, in which viruses are packaged and transmitted collectively, can enhance coinfection [35, 36]. To our knowledge, poliovirus does not have superinfection exclusion behavior [37, 38], but growing evidence suggests that *en bloc* transmission [35, 39, 36] and even shuttling of poliovirus by enteric bacteria [40] can be common during infection. These factors could elevate poliovirus coinfection rates beyond

what we consider in our model, or change which genotypes coinfect together.

Host and environmental factors could also affect the realized frequency of coinfection and subsequent evolutionary dynamics. We assume that cells are all equally susceptible to poliovirus infection and spatially well-mixed. In practice, expression of poliovirus' primary receptor CD155 varies considerably among cells [41] and at different stages of disease [42]. This could concentrate virus into a smaller number of cells, enhancing interference, or result in certain cells that can only be infected by few virions, potentially allowing greater expression of phenotypic resistance. The organization of host cells within a tissue also stands to impact coinfection dynamics [43]. Limited viral dispersal could increase coinfection and therefore interference, but it might also concentrate resistant genomes into the tissue sections in which they initially arose, limiting the degree to which susceptible genomes could interfere with resistant spread. The degree of this effect might also vary across organ systems. Notably, the limited resistance evolution observed in pocapavir-treated mice may relate to increased viral density in neuronal infections [6, 44] relative to the gut epithelial infections of the pocapavir clinical trial in which resistance evolution was common. Investigating the importance of intra-host spatial and environmental variation is an important future area of research.

A second potentially important form of spatial organization is intracellular. Many viruses, including poliovirus, form membrane-associated structures that can sequester viral components near their encoding genomes [45, 46]. This effect could limit protein diffusion and reduce phenotypic mixing. Poliovirus capsids are tethered to their membranes during virion assembly [47], so it is possible that some degree of intracellular segregation contributes to our observation that even very rare resistant genomes in a cell impart partial phenotypic resistance. Despite this, Tanner et al. [6] observed that phenotypically-distinct capsid proteins intermingle in single chimeric virions, so the exact extent of intracellular mixing of poliovirus capsid subunits remains unknown. Nevertheless, the intracellular organization of viruses within a cell is likely to be a key determinant of therapeutic strategies built around social resource sharing.

More broadly, our study contributes to a growing body of work framing viral coinfection through the lens of ploidy and drawing parallels to how multiple gene copies shape phenotype in cellular organisms [48, 7]. In classical diploid genetics, a single biallelic locus can produce up to three phenotypes, depending on the interactions between the alleles (i.e., the "dominance" of one allele over the other). Although the frequency of these phenotypes can shift between generations, the ploidy level itself remains fixed. Among viruses, the number of genomes that contribute to a phenotype can vary between host cells and dynamically over the course of an infection [7]. The association of several viral genomes in a host cell can lead to more complex and varied phenotypes than is possible in a diploid model [49]. The model that we explore here has a classical analog to incomplete dominance in a standard diploid framework, in which the addition of more resistant proteins always partially benefits a capsid in the presence of the drug. However, recent work has also described instances of apparent over- or underdominance in which coinfecting viruses have increased or decreased fitness relative to either non-mixed infection [50, 51, 52, 53].

Regardless of the exact form of intracellular "dominance," social phenotypes can clearly determine viral fitness. Because absolute fitness within a population governs viral density in the immediate future, it therefore impacts the degree of viral social interaction moving forward, and thus how new phenotypes are realized. Therefore, viral "ploidy" not only changes over time, but actually feeds back into itself (**Figure 6**). Although our model centers on poliovirus and pocapavir, its core principles—density-driven coinfection, transient genotype-phenotype associations, and demographic feedback with the environment —are likely to be broadly relevant for designing therapies that exploit the social lives of viruses, and viral evolutionary dynamics more broadly.

Figure 6: **Targeting social viral traits requires an understanding of eco-evolutionary feedback**. Viral density determines the degree of intracellular social interactions, which determines realized phenotypes, which determine absolute fitness, which feed back into viral density in the next generation.

Materials & Methods

We developed both stochastic and deterministic versions of a discrete-time dynamical model of poliovirus replication that integrates viral entry, genome replication, mutation, capsid formation, pocapavir neutralization, and immune clearance. We describe the stochastic version here, but the deterministic version follows the same probability distributions but accounts for each possible outcome (i.e., integrates over the probability distribution) instead of drawing specific values. Full details are provided in the Supplemental Materials & Methods. All simulations and analyses were performed in R, version 4.1.0 [54]. Data visualization was performed with the `ggplot2` package [55].

Viral entry

At time t, the total viral population, $v_{\text{tot},t}$, consists of $r_{\text{tot},t}$ resistant and $s_{\text{tot},t}$ susceptible genomes, yielding

$$v_{\text{tot},t} = r_{\text{tot},t} + s_{\text{tot},t}.$$

The multiplicity of infection (MOI, represented as m) is defined by the ratio of viruses to host cells (host cell population size represented as γ):

$$m = \frac{v_{\text{tot},t}}{\gamma}.$$

The number of virions entering a cell (a random variable, V_{inf}) is modeled by a Poisson distribution,

$$V_{\text{inf}} \sim \text{Pois}(m).$$

Thus, on average, a cell will be infected by m viruses.

Given that V_{inf} takes on the value v_{inf}, the number of resistant genomes per cell (a random

variable, R_{inf}) is binomially sampled from the total viral population

$$R_{\text{inf}} \sim \text{Bin}\left(v_{\text{inf}}, \frac{r_{\text{tot},t}}{v_{\text{tot},t}}\right).$$

Given that R_{inf} takes on the value r_{inf}, the number of susceptible genomes in a cell is given by $s_{\text{inf}} = v_{\text{inf}} - r_{\text{inf}}$.

Genome replication and mutation

For each infected cell, progeny production (V_{rep}) follows a Poisson distribution with a mean of the inferred average burst size, β (i.e., $V_{\text{rep}} \sim \text{Pois}(\beta)$ for $v_{\text{inf}} > 0$). Replicated resistant genomes (R_{rep}) are modeled as

$$R_{\text{rep}} \sim \text{Bin}\left(v_{\text{rep}}, \frac{r_{\text{inf}}}{v_{\text{inf}}}\right).$$

Given that R_{rep} takes on some value, r_{rep}, the number of newly replicated susceptible genomes in a cell, s_{rep}, is given by:

$$s_{\text{rep}} = v_{\text{rep}} - r_{\text{rep}}.$$

Mutation between genotypes occurs per replication event at rate $\mu = 2 \times 10^{-5}$ [9], so that resistant and susceptible mutants (the random variables R_{mut} and S_{mut}, respectively) are found by:

$$R_{\text{mut}} \sim \text{Bin}\left(s_{\text{rep}}, \mu\right) \quad \text{and} \quad S_{\text{mut}} \sim \text{Bin}\left(r_{\text{rep}}, \mu\right).$$

Given that R_{mut} and S_{mut} take on the values r_{mut} and s_{mut}, respectively, post-mutation genome counts per cell (r_{pool} and s_{pool}) are then found as follows:

$$r_{\text{pool}} = r_{\text{rep}} - s_{\text{mut}} + r_{\text{mut}}, \quad s_{\text{pool}} = s_{\text{rep}} - r_{\text{mut}} + s_{\text{mut}}.$$

Capsid formation

Let σ represent the number of subunits in a capsid. Each progeny genome is packaged into a capsid comprised of 60 subunits ($\sigma = 60$). The number of resistant subunits per virion (the random variable D_{cap}) is modeled binomially, where:

$$D_{\text{cap}} \sim \text{Bin}\left(\sigma, \frac{r_{\text{inf}}}{v_{\text{inf}}}\right),$$

assuming that both resistant and susceptible genomes contribute equally to the pool of capsid subunits. The probability that a capsid has i subunits is:

$$p_i = \Pr(D_{\text{cap}} = i \mid \{v_{\text{inf}}, r_{\text{inf}}\}),$$

where i takes on a discrete value between 0 and σ.

Genomes are assigned to capsids via a multinomial sampling process for each infected cell in the population. The number of replicated resistant genomes that are packaged into a capsid with i resistant subunits is the random variable $R_{\mathrm{pack},i}$. Each $R_{\mathrm{pack},i}$ can be collected into a vector R_{pack}, where:

$$R_{\mathrm{pack}} \sim \mathrm{Multinom}(r_{\mathrm{pool}}, p_0, \ldots, p_\sigma)$$

with analogous sampling for susceptible genomes. At this point, virions leave their cells and are pooled into groups according to their capsid subunit composition and genotype.

Accounting for non-infectious particles

To account for non-infectious particles produced during poliovirus infection, we cull the viral population based on the particle to plaque-forming unit ratio, ϕ. Specifically, for capsids with i resistant subunits, the number of infectious particles carrying resistant or susceptible genomes is modeled as

$$R_{\mathrm{pfu},i} \sim \mathrm{Bin}\left(r_{\mathrm{pack},i}, \frac{1}{\phi}\right) \quad \text{and} \quad S_{\mathrm{pfu},i} \sim \mathrm{Bin}\left(s_{\mathrm{pack},i}, \frac{1}{\phi}\right),$$

where $r_{\mathrm{pack},i}$ and $s_{\mathrm{pack},i}$ are the numbers of resistant and susceptible genomes packaged into capsids containing i resistant subunits, respectively. We assume that non-infectious particles are inert, do not interfere with infectious particles, and that their probability of generation is not affected by capsid subunit composition.

Pocapavir neutralization

Drug neutralization is modeled by assigning each virion a survival probability, $\omega(i, t)$, that depends on its capsid composition (number of resistant subunits i) and on the time of drug administration t_{poc}. For $t < t_{\mathrm{poc}}$, $\omega(i, t) = 1$ (that is, before drug application, virions are not affected by the drug). When $t \geq t_{\mathrm{poc}}$, survival is given by a scaled logistic function:

$$\omega(i, t) = y_0 + (y_\sigma - y_0) \cdot \frac{L(i) - L(0)}{L(\sigma) - L(0)},$$

where $L(i)$ is the standard form of the logistic function:

$$L(i) = \frac{1}{1 + e^{-k(i - i_0)}}.$$

The variables y_0 and y_σ represent the survival probabilities of fully susceptible and fully resistant capsids, respectively, and k and i_0 are inferred by fitting the function to cell culture survival probabilities from Tanner et al. [6]. Under drug pressure, the survival of virions carrying resistant genomes (the random variable $R_{\mathrm{surv},i}$) is then found by binomial sampling $r_{\mathrm{pfu},i}$:

$$R_{\mathrm{surv},i} \sim \mathrm{Bin}\left(r_{\mathrm{pfu},i}, \omega(i, t)\right),$$

for each capsid subunit state (and similarly for susceptible virions). $R_{\mathrm{surv},i}$ then takes on the specific values $r_{\mathrm{surv},i}$. After pocapavir neutralization, the total number of resistant genomes, r_{sum}, is calculated by summing across all $r_{\mathrm{surv},i}$:

$$r_{\mathrm{sum}} = \sum_{i=0}^{\sigma} r_{\mathrm{surv},i}.$$

A similar sum is used to calculate total number of sensitive genomes.

Immune clearance

Following pocapavir neutralization, immune clearance is applied after a specified number of replications, t_{imm}, via the survival function $d(t)$. For $t < t_{\mathrm{imm}}$, $d(t) = 1$. For $t \geq t_{\mathrm{imm}}$, survival is given by an exponential decay function,

$$d(t) = e^{(-I_{\mathrm{init}}(t - t_{\mathrm{imm}}))},$$

where the initial immune sensitivity, I_{init}, is drawn from a lognormal distribution,

$$I_{\mathrm{init}} \sim \mathrm{Lognormal}(\xi, \tau^2).$$

The parameters governing I_{init} (t_{imm}, ξ and τ) and the host cell population size, γ, were inferred by maximizing the log-likelihood of observing simulation clearance times, given matched placebo clearance times from the pocapavir clinical trial reported by Collett et al. [11].

The virions that survive pocapavir treatment and carry resistant genomes are represented by R_{imm}. This is found by binomially sampling r_{sum}:

$$R_{\mathrm{imm}} \sim \mathrm{Bin}(r_{\mathrm{sum}}, d(t)).$$

Note, immune clearance is considered in the stochastic model only.

Variation in phenotypic dominance

To explore the impact of susceptible dominance over the resistant phenotype, we changed the drug neutralization function to simulate different relationships between resistant subunit composition and virion survival. Using the standard logistic function as our base, we used a steepness coefficient of $k = 100$ to simulate a step-like function, and varied i_0 to set the inflection point, corresponding to the minimum number of resistant subunits needed to render a virion phenotype resistant to drug. Otherwise, simulations were initialized with the same parameters as the pocapavir simulations.

Variation in drug strength

To explore the impact of drug strength on clinical outcomes, we scaled the original pocapavir fitness function such that,

$$\omega'(i,t) = y_0' + \frac{(\omega(i,t) - y_0)(y_\sigma - y_0')}{y_\sigma - y_0},$$

where y_0 is the fitness of a fully susceptible capsid in the presence of pocapavir, y_σ is the fitness of a fully resistant capsid in the presence of pocapavir, and y_0' is the new survival probability of a virion composed entirely of susceptible subunits. Otherwise, simulations were initialized with the same parameters as the pocapavir simulations.

Parameter inference

Parameters were inferred by numerical optimization in `R` (version 4.1.0) [54]. The specific objective function varied by model and is described below.

Estimation of viral burst size and particle-to-plaque-forming-unit ratio

We inferred the viral burst size and particle-to-plaque-forming unit (P:PFU) ratio by fitting model outputs to mixed cell culture data from Tanner et al. [6], using matched initial multiplicities of infection (MOIs). We specifically compared our simulated data to reported results of pure Mahoney strain PV with the VP3-A24V mutation in cell culture. This mutation is one of the most commonly observed in experiments selecting for resistance [9, 10], and was the only strain/mutation pairing for which there was a negative control reported in Tanner et al. [6]. We evaluated model fit by minimizing the sum of squared differences between the log-values of the observed and simulated PFU after one round of replication. Specifically, we minimized:

$$\sum_l \Big(\log(\mathrm{real\_pfu}_l) - \log(\mathrm{sim\_pfu}_l) \Big)^2,$$

where l is the set of resistant MOI and susceptible MOI co-infected pairs reported by Tanner et al. [6]. Fit values and their empirically measured comparisons are reported in Table S1.

Estimation of immune clearance parameters

Parameters governing immune clearance and host cell population size were estimated by fitting a stochastic model of infection and clearance (assuming no drug effect; i.e., survival = 1 for all phenotypes at all time points) to placebo group clearance data from Collett et al. [11]. Since participants were not sampled daily, Collett et al. reported the clearance date as the first sample at which no virus was detected (**Figure 2B**). To replicate this, we rounded each simulated clearance time up to the next available sampling day, following the trial design.

We simulated 480 placebo recipients ($10\times$ the original sample size) and calculated the probability of clearance on each sampled day. These model-based probabilities were compared to the empirical distribution using a multinomial log-likelihood. Specifically, after adding a small pseudo-count to avoid log-zero issues, we computed:

$$\log L = \sum_x n_x \cdot \log p_x$$

where n_x is the number of placebo participants observed to clear on day x, and p_x is the model-derived probability of clearance on day x. If a simulated participant cleared after day 43 (the final sampling day), we could not compare this outcome to empirical data and instead returned a fixed log-likelihood value of 1000 to penalize these parameter settings. Fit values and their empirically measured comparisons are reported in Table S1.

We note that $\gamma \times \mu$ has an important function in shaping the probability of viral extinction. There is a two step requirement for virion survival: first, resistant viruses must *de novo* mutate and be infectious virions, and second, those virions must survive pocapavir treatment during the first round of infection, given that they were wrapped in susceptible capsids. This is mathematically represented as $P(\text{extinction}) = \beta \frac{1}{\phi} \gamma \mu y_0$. Using our estimated parameters reported in Table S1, we calculate that the inflection point in the $P(\text{extinction})$ curve (the point at which 50% of populations should go extinct) is at $\gamma \times \mu \approx 6.5$, a result reflected in model simulations (**Figure S6**).

A complete account of the model equations, parameter inference, and simulation details is provided in the Supplemental Materials & Methods.

Data, Materials, and Software Availability

Code for simulations, data analysis, and data visualization have been deposited on GitHub (https://github.com/albobson/polio-res-eco-evo).

Acknowledgments

We thank Feder and Kerr lab members for useful discussions and manuscript feedback. We thank Gaël Thebaud for the helpful advice on fitting the pocapavir fitness function. This work was possible by funding from NIH training grant T32-GM136534-02 supporting A. J. R and by the Environmental Biology Division from the National Science Foundation (grant number 2142718 to B.K.).

Author contributions

A.J.R, B.K and A.F.F. contributed to study conceptualization and design. A.J.R. performed the analysis. A.J.R, B.K and A.F.F. interpreted the results, wrote the original draft of the manuscript and contributed to review and editing of the manuscript.

Competing interests

The authors declare no competing interests.

References

[1] Elsa Hansen, Jason Karslake, Robert J. Woods, Andrew F. Read, and Kevin B. Wood. Antibiotics can be used to contain drug-resistant bacteria by maintaining sufficiently large sensitive populations. *PLOS Biology*, 18(5):e3000713, May 2020. ISSN 1545-7885. doi: 10.1371/journal.pbio.3000713.

[2] Caroline Colijn and Ted Cohen. How competition governs whether moderate or aggressive treatment minimizes antibiotic resistance. *eLife*, 4:e10559, September 2015. ISSN 2050-084X. doi: 10.7554/eLife.10559.

[3] Raul Andino and Esteban Domingo. Viral quasispecies. *Virology*, 479–480:46–51, May 2015. ISSN 0042-6822. doi: 10.1016/j.virol.2015.03.022.

[4] Claus O Wilke and Isabel S Novella. Phenotypic mixing and hiding may contribute to memory in viral quasispecies. *BMC Microbiology*, 3:11, June 2003. ISSN 1471-2180. doi: 10.1186/1471-2180-3-11.

[5] Claude Loverdo and James O. Lloyd-Smith. Inter-Generational Phenotypic Mixing in Viral Evolution. *Evolution; international journal of organic evolution*, 67(6):1815–1822, June 2013. ISSN 0014-3820. doi: 10.1111/evo.12048.

[6] Elizabeth J Tanner, Hong-mei Liu, M Steven Oberste, Mark Pallansch, Marc S Collett, and Karla Kirkegaard. Dominant drug targets suppress the emergence of antiviral resistance. *eLife*, 3:e03830, November 2014. ISSN 2050-084X. doi: 10.7554/eLife.03830.

[7] Asher Leeks, Lisa M. Bono, Elizabeth A. Ampolini, Lucas S. Souza, Thomas Höfler, Courtney L. Mattson, Anna E. Dye, and Samuel L. Díaz-Muñoz. Open questions in the social lives of viruses. *Journal of Evolutionary Biology*, 36(11):1551–1567, 2023. ISSN 1420-9101. doi: 10.1111/jeb.14203.

[8] Elizabeth J. Tanner, Karla A. Kirkegaard, and Leor S. Weinberger. Exploiting Genetic Interference for Antiviral Therapy. *PLOS Genetics*, 12(5):e1005986, May 2016. ISSN 1553-7404. doi: 10.1371/journal.pgen.1005986.

[9] Diana V Kouiavskaia, Eugenia M Dragunsky, Hong-Mei Liu, M Steven Oberste, Marc S Collett, and Konstantin M Chumakov. Immunological and Pathogenic Properties of Poliovirus Variants Selected for Resistance to Antiviral Drug V-073. *Antiviral Therapy*, 16(7):999–1004, October 2011. ISSN 1359-6535. doi: 10.3851/IMP1838.

[10] Hong-Mei Liu, Jason A. Roberts, Deborah Moore, Barbara Anderson, Mark A. Pallansch, Daniel C. Pevear, Marc S. Collett, and M. Steven Oberste. Characterization of Poliovirus Variants Selected for Resistance to the Antiviral Compound V-073. *Antimicrobial Agents and Chemotherapy*, 56(11):5568–5574, November 2012. doi: 10.1128/AAC.00539-12.

[11] Marc S. Collett, Jeffrey R. Hincks, Kimberley Benschop, Erwin Duizer, Harrie van der Avoort, Eric Rhoden, Hongmei Liu, M. Steven Oberste, Mark A. McKinlay, and Marianne Hartford. Antiviral Activity of Pocapavir in a Randomized, Blinded, Placebo-Controlled Human Oral Poliovirus Vaccine Challenge Model. *The Journal of Infectious Diseases*, 215(3):335–343, February 2017. ISSN 0022-1899. doi: 10.1093/infdis/jiw542.

[12] Julie Copelyn, Jeffrey R. Hincks, Jo M. Wilmshurst, Wonita Petersen, Wayne Howard, Sabelle Jallow, Shelina Moonsamy, Lerato Seakamela, Melinda Suchard, Marc S. Collett, and Brian Eley. Clearance of Immunodeficiency-associated Vaccine-derived Poliovirus Infection With Pocapavir. *The Pediatric Infectious Disease Journal*, 39(5):435, May 2020. ISSN 0891-3668. doi: 10.1097/INF.0000000000002584.

[13] Sanet Torres-Torres, Angela L. Myers, J. Michael Klatte, Eric E. Rhoden, M. Steven Oberste, Marc S. Collett, and Russell J. McCulloh. First Use of Investigational Antiviral Drug Pocapavir (V-073) for Treating Neonatal Enteroviral Sepsis. *The Pediatric Infectious Disease Journal*, 34(1):52, January 2015. ISSN 0891-3668. doi: 10.1097/INF.0000000000000497.

[14] Ashleigh S. Griffin and Asher Leeks. Exploiting social traits for clinical applications in bacteria and viruses. *npj Antimicrobials and Resistance*, 3(1):1–9, March 2025. ISSN 2731-8745. doi: 10.1038/s44259-025-00091-6.

[15] Leor S. Weinberger, David V. Schaffer, and Adam P. Arkin. Theoretical Design of a Gene Therapy To Prevent AIDS but Not Human Immunodeficiency Virus Type 1 Infection. *Journal of Virology*, 77(18):10028–10036, September 2003. ISSN 0022-538X. doi: 10.1128/JVI.77.18.10028-10036.2003.

[16] Vincent T. Metzger, James O. Lloyd-Smith, and Leor S. Weinberger. Autonomous Targeting of Infectious Superspreaders Using Engineered Transmissible Therapies. *PLoS Computational Biology*, 7(3):e1002015, March 2011. ISSN 1553-734X. doi: 10.1371/journal.pcbi.1002015.

[17] Fathima N Nagoor Pitchai, Elizabeth J Tanner, Neha Khetan, Gustavo Vasen, Clara Levrel, Arjun J Kumar, Shilpi Pandey, Tracy Ordonez, Philip Barnette, David Spencer, et al. Engineered deletions of HIV replicate conditionally to reduce disease in nonhuman primates. *Science*, 385(6709):eadn5866, 2024.

[18] Sonali Chaturvedi, Gustavo Vasen, Michael Pablo, Xinyue Chen, Nathan Beutler, Arjun Kumar, Elizabeth Tanner, Sylvia Illouz, Donna Rahgoshay, John Burnett, Leo Holguin, Pei-Yi Chen, Blaise Ndjamen, Melanie Ott, Robert Rodick, Thomas Rogers, Davey M. Smith, and

Leor S. Weinberger. Identification of a therapeutic interfering particle—A single-dose SARS-CoV-2 antiviral intervention with a high barrier to resistance. *Cell*, 184(25):6022–6036.e18, December 2021. ISSN 0092-8674, 1097-4172. doi: 10.1016/j.cell.2021.11.004.

[19] Sonali Chaturvedi, Nathan Beutler, Gustavo Vasen, Michael Pablo, Xinyue Chen, Giuliana Calia, Lauren Buie, Robert Rodick, Davey Smith, Thomas Rogers, and Leor S. Weinberger. A single-administration therapeutic interfering particle reduces SARS-CoV-2 viral shedding and pathogenesis in hamsters. *Proceedings of the National Academy of Sciences*, 119(39): e2204624119, September 2022. doi: 10.1073/pnas.2204624119.

[20] Veronica V. Rezelj, Lucía Carrau, Fernando Merwaiss, Laura I. Levi, Diana Erazo, Quang Dinh Tran, Annabelle Henrion-Lacritick, Valérie Gausson, Yasutsugu Suzuki, Djoshkun Shengjuler, Bjoern Meyer, Thomas Vallet, James Weger-Lucarelli, Veronika Bernhauerová, Avi Titievsky, Vadim Sharov, Stefano Pietropaoli, Marco A. Diaz-Salinas, Vincent Legros, Nathalie Pardigon, Giovanna Barba-Spaeth, Leonid Brodsky, Maria-Carla Saleh, and Marco Vignuzzi. Defective viral genomes as therapeutic interfering particles against flavivirus infection in mammalian and mosquito hosts. *Nature Communications*, 12(1):2290, April 2021. ISSN 2041-1723. doi: 10.1038/s41467-021-22341-7.

[21] Lei Zhang, Jianli Ma, Lei Liu, Guozheng Li, Hui Li, Yi Hao, Xin Zhang, Xin Ma, Yihai Chen, Jiale Wu, Xinheng Wang, Shuai Yang, and Shouping Xu. Adaptive therapy: a tumor therapy strategy based on Darwinian evolution theory. *Critical Reviews in Oncology/Hematology*, 192: 104192, December 2023. ISSN 1040-8428. doi: 10.1016/j.critrevonc.2023.104192.

[22] Robert A. Gatenby and Joel S. Brown. The Evolution and Ecology of Resistance in Cancer Therapy. *Cold Spring Harbor Perspectives in Medicine*, 10(11):a040972, November 2020. ISSN 2157-1422. doi: 10.1101/cshperspect.a040972.

[23] Olha Puhach, Benjamin Meyer, and Isabella Eckerle. SARS-CoV-2 viral load and shedding kinetics. *Nature Reviews Microbiology*, 21(3):147–161, March 2023. ISSN 1740-1534. doi: 10.1038/s41579-022-00822-w.

[24] Hitoshi Kawasuji, Yusuke Takegoshi, Makito Kaneda, Akitoshi Ueno, Yuki Miyajima, Koyomi Kawago, Yasutaka Fukui, Yoshihiro Yoshida, Miyuki Kimura, Hiroshi Yamada, Ippei Sakamaki, Hideki Tani, Yoshitomo Morinaga, and Yoshihiro Yamamoto. Transmissibility of COVID-19 depends on the viral load around onset in adult and symptomatic patients. *PLOS ONE*, 15(12):e0243597, December 2020. ISSN 1932-6203. doi: 10.1371/journal.pone.0243597.

[25] Robert W. Eisinger, Carl W. Dieffenbach, and Anthony S. Fauci. HIV Viral Load and Transmissibility of HIV Infection: Undetectable Equals Untransmittable. *JAMA*, 321(5):451–452, February 2019. ISSN 0098-7484. doi: 10.1001/jama.2018.21167.

[26] Anne Sicard, Yannis Michalakis, Serafín Gutiérrez, and Stéphane Blanc. The Strange Lifestyle of Multipartite Viruses. *PLOS Pathogens*, 12(11):e1005819, November 2016. ISSN 1553-7374. doi: 10.1371/journal.ppat.1005819.

[27] Christopher B. Brooke, William L. Ince, Jens Wrammert, Rafi Ahmed, Patrick C. Wilson, Jack R. Bennink, and Jonathan W. Yewdell. Most Influenza A Virions Fail To Express at Least One Essential Viral Protein. *Journal of Virology*, 87(6):3155–3162, March 2013. doi: 10.1128/jvi.02284-12.

[28] Alistair B Russell, Cole Trapnell, and Jesse D Bloom. Extreme heterogeneity of influenza virus infection in single cells. *Elife*, 7:e32303, 2018.

[29] D S Burke. Recombination in HIV: an important viral evolutionary strategy. *Emerging Infectious Diseases*, 3(3):253–259, 1997. ISSN 1080-6040. doi: 10.3201/eid0303.970301.

[30] Elena V Romero and Alison F Feder. Elevated HIV Viral Load is Associated with Higher Recombination Rate In Vivo. *Molecular Biology and Evolution*, 41(1):msad260, January 2024. ISSN 1537-1719. doi: 10.1093/molbev/msad260.

[31] Michael B Schulte, Jeremy A Draghi, Joshua B Plotkin, and Raul Andino. Experimentally guided models reveal replication principles that shape the mutation distribution of RNA viruses. *eLife*, 4:e03753, January 2015. ISSN 2050-084X. doi: 10.7554/eLife.03753.

[32] Michael B. Schulte and Raul Andino. Single-Cell Analysis Uncovers Extensive Biological Noise in Poliovirus Replication. *Journal of Virology*, 88(11):6205–6212, June 2014. ISSN 0022-538X, 1098-5514. doi: 10.1128/JVI.03539-13.

[33] Roberto Mateo, Claude M. Nagamine, and Karla Kirkegaard. Suppression of Drug Resistance in Dengue Virus. *mBio*, 6(6):10.1128/mbio.01960–15, December 2015. doi: 10.1128/mbio.01960-15.

[34] Michael Hunter and Diana Fusco. Superinfection exclusion: A viral strategy with short-term benefits and long-term drawbacks. *PLOS Computational Biology*, 18(5):e1010125, May 2022. ISSN 1553-7358. doi: 10.1371/journal.pcbi.1010125.

[35] Adeline Kerviel, Mengyang Zhang, and Nihal Altan-Bonnet. A New Infectious Unit: Extracellular Vesicles Carrying Virus Populations. *Annual Review of Cell and Developmental Biology*, 37(1):171–197, 2021. doi: 10.1146/annurev-cellbio-040621-032416.

[36] Ying-Han Chen, WenLi Du, Marne C. Hagemeijer, Peter M. Takvorian, Cyrilla Pau, Ann Cali, Christine A. Brantner, Erin S. Stempinski, Patricia S. Connelly, Hsin-Chieh Ma, Ping Jiang, Eckard Wimmer, Grégoire Altan-Bonnet, and Nihal Altan-Bonnet. Phosphatidylserine Vesicles Enable Efficient En Bloc Transmission of Enteroviruses. *Cell*, 160(4):619–630, February 2015. ISSN 0092-8674. doi: 10.1016/j.cell.2015.01.032.

[37] Denise Egger and Kurt Bienz. Intracellular location and translocation of silent and active poliovirus replication complexes. *Journal of General Virology*, 86(3):707–718, 2005. ISSN 1465-2099. doi: 10.1099/vir.0.80442-0.

[38] Purnell W. Choppin and Kathryn V. Holmes. Replication of SV5 RNA and the effects of superinfection with poliovirus. *Virology*, 33(3):442–451, November 1967. ISSN 0042-6822. doi: 10.1016/0042-6822(67)90119-5.

[39] Elizabeth R. Aguilera, Andrea K. Erickson, Palmy R. Jesudhasan, Christopher M. Robinson, and Julie K. Pfeiffer. Plaques Formed by Mutagenized Viral Populations Have Elevated Coinfection Frequencies. *mBio*, 8(2):10.1128/mbio.02020–16, March 2017. doi: 10.1128/mbio.02020-16.

[40] Andrea K. Erickson, Palmy R. Jesudhasan, Melinda J. Mayer, Arjan Narbad, Sebastian E. Winter, and Julie K. Pfeiffer. Bacteria Facilitate Enteric Virus Co-infection of Mammalian Cells and Promote Genetic Recombination. *Cell Host Microbe*, 23(1):77–88.e5, January 2018. ISSN 1931-3128. doi: 10.1016/j.chom.2017.11.007.

[41] Marion S. Freistadt, Gerardo Kaplan, and Vincent R. Racaniello. Heterogeneous Expression of Poliovirus Receptor-Related Proteins in Human Cells and Tissues. *Molecular and Cellular Biology*, 10(11):5700–5706, November 1990. ISSN null. doi: 10.1128/mcb.10.11.5700-5706.1990.

[42] Rosa Molfetta, Beatrice Zitti, Mario Lecce, Nadia Domenica Milito, Helena Stabile, Cinzia Fionda, Marco Cippitelli, Angela Gismondi, Angela Santoni, and Rossella Paolini. CD155: A Multi-Functional Molecule in Tumor Progression. *International Journal of Molecular Sciences*, 21(33):922, January 2020. ISSN 1422-0067. doi: 10.3390/ijms21030922.

[43] Alex Farrell, Tin Phan, Christopher B Brooke, Katia Koelle, and Ruian Ke. Semi-infectious particles contribute substantially to influenza virus within-host dynamics when infection is dominated by spatial structure. *Virus Evolution*, 9(1):vead020, March 2023. ISSN 2057-1577. doi: 10.1093/ve/vead020.

[44] P. J. Buontempo, S. Cox, J. Wright-Minogue, J. L. DeMartino, A. M. Skelton, E. Ferrari, R. Albin, E. J. Rozhon, V. Girijavallabhan, J. F. Modlin, and J. F. O'Connell. SCH 48973: a potent, broad-spectrum, antienterovirus compound. *Antimicrobial Agents and Chemotherapy*, 41(6):1220–1225, June 1997. ISSN 0066-4804. doi: 10.1128/AAC.41.6.1220.

[45] A Schlegel, T H Giddings, M S Ladinsky, and K Kirkegaard. Cellular origin and ultrastructure of membranes induced during poliovirus infection. *Journal of Virology*, 70(10):6576–6588, October 1996. doi: 10.1128/jvi.70.10.6576-6588.1996.

[46] Reyes R. Novoa, Gloria Calderita, Rocío Arranz, Juan Fontana, Harald Granzow, and Cristina Risco. Virus factories: associations of cell organelles for viral replication and morphogenesis. *Biology of the Cell*, 97(2):147–172, 2005. ISSN 1768-322X. doi: 10.1042/BC20040058.

[47] Selma Dahmane, Adeline Kerviel, Dustin R. Morado, Kasturika Shankar, Björn Ahlman, Michael Lazarou, Nihal Altan-Bonnet, and Lars-Anders Carlson. Membrane-assisted assembly and selective secretory autophagy of enteroviruses. *Nature Communications*, 13(1):5986, October 2022. ISSN 2041-1723. doi: 10.1038/s41467-022-33483-7.

[48] Claus O. Wilke. Quasispecies theory in the context of population genetics. *BMC Evolutionary Biology*, 5(1):44, August 2005. ISSN 1471-2148. doi: 10.1186/1471-2148-5-44.

[49] John P. DeLong, Maitham A. Al-Sammak, Zeina T. Al-Ameeli, David D. Dunigan, Kyle F. Edwards, Jeffry J. Fuhrmann, Jason P. Gleghorn, Hanqun Li, Kona Haramoto, Amelia O. Harrison, Marcia F. Marston, Ryan M. Moore, Shawn W. Polson, Barbra D. Ferrell, Miranda E. Salsbery, Christopher R. Schvarcz, Jasmine Shirazi, Grieg F. Steward, James L. Van Etten, and K. Eric Wommack. Towards an integrative view of virus phenotypes. *Nature Reviews Microbiology*, 20(2):83–94, February 2022. ISSN 1740-1534. doi: 10.1038/s41579-021-00612-w.

[50] Iris Yousaf, William W. Hannon, Ryan C. Donohue, Christian K. Pfaller, Kalpana Yadav, Ryan J. Dikdan, Sanjay Tyagi, Declan C. Schroeder, Wun-Ju Shieh, Paul A. Rota, Alison F. Feder, and Roberto Cattaneo. Brain tropism acquisition: The spatial dynamics and evolution of a measles virus collective infectious unit that drove lethal subacute sclerosing panencephalitis. *PLOS Pathogens*, 19(12):e1011817, December 2023. ISSN 1553-7374. doi: 10.1371/journal.ppat.1011817.

[51] Katherine S Xue, Kathryn A Hooper, Anja R Ollodart, Adam S Dingens, and Jesse D Bloom. Cooperation between distinct viral variants promotes growth of H3N2 influenza in cell culture. *Elife*, 5:e13974, 2016.

[52] Yuta Shirogane, Hidetaka Harada, Yuichi Hirai, Ryuichi Takemoto, Tateki Suzuki, Takao Hashiguchi, and Yusuke Yanagi. Collective fusion activity determines neurotropism of an en bloc transmitted enveloped virus. *Science Advances*, 9(4):eadf3731, 2023.

[53] Celia Perales, Roberto Mateo, Mauricio G. Mateu, and Esteban Domingo. Insights into RNA Virus Mutant Spectrum and Lethal Mutagenesis Events: Replicative Interference and Complementation by Multiple Point Mutants. *Journal of Molecular Biology*, 369(4):985–1000, June 2007. ISSN 0022-2836. doi: 10.1016/j.jmb.2007.03.074.

[54] R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2024. URL https://www.R-project.org/.

[55] Hadley Wickham. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York, 2016. ISBN 978-3-319-24277-4. URL https://ggplot2.tidyverse.org.

[56] Adi Stern, Ming Te Yeh, Tal Zinger, Matt Smith, Caroline Wright, Guy Ling, Rasmus Nielsen, Andrew Macadam, and Raul Andino. The Evolutionary Pathway to Virulence

of an RNA Virus. *Cell*, 169(1):35–46.e19, March 2017. ISSN 0092-8674, 1097-4172. doi: 10.1016/j.cell.2017.03.013.

[57] Nadège Néant, Guillaume Lingas, Quentin Le Hingrat, Jade Ghosn, Ilka Engelmann, Quentin Lepiller, Alexandre Gaymard, Virginie Ferré, Cédric Hartard, Jean-Christophe Plantier, Vincent Thibault, Julien Marlet, Brigitte Montes, Kevin Bouiller, François-Xavier Lescure, Jean-François Timsit, Emmanuel Faure, Julien Poissy, Christian Chidiac, François Raffi, Antoine Kimmoun, Manuel Etienne, Jean-Christophe Richard, Pierre Tattevin, Denis Garot, Vincent Le Moing, Delphine Bachelet, Coralie Tardivon, Xavier Duval, Yazdan Yazdanpanah, France Mentré, Cédric Laouénan, Benoit Visseaux, Jérémie Guedj, for the French COVID Cohort Investigators, and French Cohort Study groups. Modeling SARS-CoV-2 viral kinetics and association with mortality in hospitalized patients from the French COVID cohort. *Proceedings of the National Academy of Sciences*, 118(8):e2017962118, February 2021. doi: 10.1073/pnas.2017962118.

[58] Achim Zeileis and Gabor Grothendieck. zoo: S3 infrastructure for regular and irregular time series. *Journal of Statistical Software*, 14(6):1–27, 2005. doi: 10.18637/jss.v014.i06.

[59] Olen M. Kew, Roland W. Sutter, Esther M. de Gourville, Walter R. Dowdle, and Mark A. Pallansch. Vaccine-Derived Polioviruses and the Endgame Strategy for Global Polio Eradication. *Annual Review of Microbiology*, October 2005. doi: 10.1146/annurev.micro.58.030603.123625. URL https://zenodo.org/record/1235033.

[60] Carlton E. Schwerdt and Jørgen Fogh. The ratio of physical particles per infectious unit observed for poliomyelitis viruses. *Virology*, 4(1):41–52, August 1957. ISSN 0042-6822. doi: 10.1016/0042-6822(57)90042-9.

[61] Andreas Dotzauer and Leena Kraemer. Innate and adaptive immune responses against picornaviruses and their counteractions: An overview. *World Journal of Virology*, 1(3):91–107, June 2012. ISSN 2220-3249. doi: 10.5501/wjv.v1.i3.91.