Ecological constraints and evolutionary trade-offs shape nitrogen fixation across habitats

Morgan S. Sobol[1,2*], Aya S. Klos[1], Cécile Ané[3,4,] Katherine D. McMahon[5] and Betül Kaçar[1*]

1. Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA
2. Department of Biology, Texas State University, San Marcos, TX, USA
3. Department of Statistics, University of Wisconsin-Madison, Madison, WI, USA
4. Department of Botany, University of Wisconsin-Madison, Madison, WI, USA
5. Department of Civil and Environmental Engineering, University of Wisconsin–Madison, Madison, WI, USA

Correspondence: msobol@txstate.edu, bkacar@wisc.edu

Abstract

From its earliest beginnings, life's expansion into new habitats has been profoundly shaped by its reciprocal interactions with Earth's changing environments. Understanding how ancient metabolisms co-evolved with their environments requires uncovering the ecological and evolutionary processes that structured the functionally linked genes and networks underlying these metabolisms. Here, we focus on nitrogen (N_2) fixation, one of life's most critical metabolisms, and investigate the drivers of complexity in its associated gene machinery today. We used a large-scale comparative genomics framework to construct a comprehensive catalog of extant N_2 fixation-associated genes and assessed their distribution across diverse microbial genomes and environmental backgrounds. Genomes enriched in N_2 fixation genes generally have larger genome sizes, broader metabolic capabilities, wider habitat ranges, and are predominantly associated with mesophilic and aerobic lifestyles. Evolutionary reconstructions reveal a pattern of early gene gains in ancestral diazotrophs followed by lineage-specific gene losses in later diverging taxa, suggesting evolutionary trade-offs shaped by changing environments. These findings demonstrate that the evolution of N_2 fixation has been intertwined with the composition and organization of the genes supporting the overarching N_2 metabolism,

32   driven by feedback between genome evolution and shifting environmental and ecological

33   conditions.

Introduction

Biological nitrogen fixation ("N$_2$ fixation") is an ancient metabolism characterized by the

reduction of atmospheric N$_2$ to bioavailable ammonia [1, 2]. Nitrogenases are the only

known enzymes capable of catalyzing this energy-intensive reaction, giving them a crucial

historical role expanding the biosphere through nitrogen supply [3, 4].

Phylogenetic reconstructions support a shared ancestry among five major clades of

nitrogenases that have diversified over time [5]. These include three isozymes defined by

the metal in their active site cofactors, molybdenum (Mo), vanadium (V), or iron (Fe), and

the three Mo-nitrogenase clades that differ in host ecology and taxonomy [2, 6, 7].

Structural variations distinguish the Mo-nitrogenase clades and align with historic

transitions in environmental oxygen levels [6]. Despite this diversification, nitrogenase's

core structure [6] and catalytic mechanism [8, 9] have remained largely conserved over

~3 billion years of evolution [10, 11].

Outside the main nitrogenase complex (*nifHDK* for Mo-nitrogenases), N$_2$ fixation depends

on a myriad of other proteins that have coevolved with it throughout the history of this

metabolism [12, 13]. These include proteins involved in cofactor assembly (*nifBEN*) [14],

electron transfer (*nifFJ*), transcriptional regulation (*nifALI$_1$I$_2$*), and biosynthesis

(*nifXQUVYS*) [15, 16]. Beyond the *nif* cluster, numerous other gene families contribute to

N$_2$ fixation, including those involved in oxygen protection (*nod, ahp, coo, cow, sod*),

electron donation and transfer (*fix, rng, fdx*), transcriptional regulation (*draGT, fixJLK,*

regRS, ntc, ntr), metal transport (*mod, feo, cys*), iron-sulfur cluster assembly (*suf, isc, fdx*)

[17], as well as nitrogen metabolism (*gln, nir, nap, hox*) [18] (**Figure 1**). Additional genes

with potential roles in N$_2$ fixation remain functionally uncharacterized [19].

The composition of the complex genetic machinery responsible for N$_2$-fixation varies

widely among diazotrophs, with the simplest gene sets found in hyperthermophilic

methanogens and the most elaborate networks in aerobes [12]. Previous studies have

highlighted the diverse evolutionary trajectories of nitrogenase-associated genes [13] and
revealed a dynamic history of gene acquisitions and losses linked to the emergence of
aerobic metabolisms [12]. Nevertheless, a central challenge remains to reconstruct the
evolutionary history of N_2 fixation in deep time and relate it to the ecological diversification
of diazotrophs. Here, we address this challenge using large-scale comparative genomics
and show how the evolution of N_2 fixation genes enabled diazotrophs to expand into
diverse and increasingly oxygenated environments.

Figure 1. Structure of the major and minor molybdenum nitrogenase (Nif) clusters, vanadium nitrogenase
(Vnf) and iron nitrogenase (Anf) clusters, and the molybdenum acquisition (Mod) cluster in model diazotroph
Azotobacter vinelandii DJ. Genome positions and gene sizes for each cluster are based on KEGG genome
reference T00885 for *A. vinelandii* DJ.

Methods

Genome selection

The reference phylogeny Web of Life (WoL) [20] was selected as the basis of our study
because of its extensive genome coverage and phylogenetic reconstruction. For our
analysis, we kept WoL genomes with ≥90% completeness and <5% contamination,
following quality thresholds established by [21]. Protein fasta files associated with the
genome accessions (named "*_protein.faa.gz") were downloaded from NCBI with eutils

[22] (accessed July 17th, 2023). Genomes missing available NCBI protein fasta files were

excluded from analysis. After filtering for high-quality genomes and removing those

without annotated fasta files, the WoL genome dataset was reduced from 10,575

genomes to 7,458. Nucleotide sequences of noncoding RNA genes were downloaded if

they were available at GenBank or RefSeq and the 16S rRNA genes were extracted.

Nitrogen fixation gene catalog curation

Protein fasta files were functionally annotated using the Kyoto Encyclopedia of Genes

and Genomes (KEGG) database [23]. Specifically, the translated amino acid sequences

were annotated with KEGG orthology (KO) identifiers by profiling each genome against

KEGG hidden Markov models (KEGG release 106) for all KOs [24] (cutoff E-value = 1e-

5) using *hmmscan* in HMMER version 3.2.2 (http://hmmer.org/). Custom Python scripts

were used to parse through each genome's *hmmscan* output and generate the final KO

presence-absence matrix. The presence/absence matrix was subset to genes known to

be involved in N_2 fixation (and which had KO identifiers), which were collected through an

extensive literature search that included foundational studies with thorough analyses of

genes involved in N_2 fixation (**Supplemental Table 1**). Gene involvement was supported

by functional confirmation, for example by evidence from protein interactions with

nitrogenase, gene knockout phenotypes, or differential gene expression under N_2 fixation.

Species tree construction and visualization

The WoL species tree was subsetted to genomes annotated with *nifHDK* using the

drop.tip function from the APE package [25] in R. We additionally confirmed the presence

of nitrogenase by phylogenetic placement of concatenated *nifDK* sequences, as

described in reference [5], from each genome. The resulting species tree, the gene

presence-absence matrix, and trait data were plotted using the ggTree R package

(version 3.6.2) [26].

Trait data curation

Optimal growth temperature (OGT), oxygen, pH preference, and habitat preference trait
data were collected for each genome by matching the GenBank accession number to
BacDive [27], NCBI BioProject [28], or GOLD JGI [29] as well as matching the NCBI taxon
IDs to the Madin et al (2020) phenotypic trait database [30]. Trait data was also predicted
for all genomes. OGTs were predicted using the OGT_prediction tool [31]. The regression
model used included both Bacteria and Archaea, but excluded using information of the
16S rRNA gene and genome size (*superkingdom-all species-*
all features ex genome size and rRNA.txt). The *Oxyphen* package was used for
oxygen metabolism prediction based on the number of oxygen utilizing genes in each
genome [32]. Predictions for pH were done with the Ramoneda et al. (2023) prediction
model which is based on genes that influence pH tolerance [33]. Habitat preference and
coverage were determined with ProkAtlas [34] by mapping the 16S rRNA gene
sequences to the ProkAtlas metagenome database. Barrnap version 0.9 [35] was used
to extract 16S rRNA sequences for genomes with missing annotations. Using a custom
python script, the longest 16S rRNA sequence was extracted for each genome as its
representative 16S rRNA gene sequence to correct for errors in sequence length due to
sequencing and assembly bias. Twenty-one genomes without a 16S rRNA gene
sequence were excluded. A nucleotide identity of 97% and a query length match of at
least 150 base pairs were used as thresholds. Habitat preference, categorized into
freshwater, saline, terrestrial, plant-associated, non-plant host-associated, and
hydrothermal, was defined as the environment with the highest 16S abundance for each
genome. Habitat coverage was determined by summing up the total number of habitats
where a genome was considered present and dividing it by the total number of habitats
(n=100) investigated by ProkAtlas. The presence of KEGG modules was predicted using
the KEGG KO annotations with MetaPathPredict (version 0) [36]. Metabolic coverage was
calculated as the total number of KEGG modules present in a genome divided by the total
number of modules (n=194) searched for with MetaPathPredict. Data collected from
databases and predicted data were cross-referenced and manually annotated to provide
the final trait data for each genome.

Data analysis and statistics

We compared the mean number of N_2 fixation genes across specific traits (oxygen,
temperature, and habitat preferences) with non-parametric methods using the
stat_compare_means function in the ggpubr R package (version 0.6.0) [37]. For the
categorical traits habitat, oxygen, and temperature preference, we first applied a Kruskal-
Wallis test, followed by pairwise Wilcoxon rank-sum tests with Benjamini-Hochberg
correction. For temperature preference (categories: mesophilic and thermophilic) we
used the Wilcoxon rank-sum test directly. Compact letter display was used to report
pairwise groups whose means are discernibly different, based on a *P*-value of 0.05.

The gene presence-absence matrix was used in phylogenetic principal-component
analysis (PCA) performed with the *phyl.pca* function (method = "BM", mode ="corr") in
phytools R package [38]. This was done to visualize similarities and dissimilarities
between genomes regarding N_2 fixation gene content while accounting for phylogenetic
relationships. This approach reduces the risk of over-interpreting similarities simple due
to common ancestry among taxa. The first two components were retained for
visualization. Pearson correlation between the presence-absence of N_2 fixation genes
was calculated with the *cor* function in the stats R package [39] and plotted using corrplot
[40] with hierarchical clustering and default *P*-value of 0.05, to identify patterns of gene
co-occurrence.

Evolutionary trends analysis

In order to test whether certain traits shared a coevolutionary history with N_2 fixation gene
abundance, phylogenetic linear regression models were fit using the *phylolm* function in
the phylolm R package (version 2.62) [41], using Pagel's lambda model of phylogenetic
correlation. The lambda parameter (λ) ranges from 0 (no phylogenetic signal) to 1 (high
phylogenetic signal) in the response trait when the model has no covariates, or in the
residuals when covariates are included. When assessing phylogenetic signal within
individual clades, those represented by fewer than 20 genomes were not included in the
analysis, as statistical power to detect signal decreases substantially in trees with fewer
than 20 branches [42, 43]. Non-phylogenetic linear regressions were performed using the
lm function in the stats R package [39]. The *distRoot* function in the adephylo R package
[44] was used to calculate the root-to-tip patristic distances by measuring the total length

of branches (in terms of the number of substitutions per site) from the root of the species

tree to each tip.

Gene gain/loss was determined with Count [45] using a non-binary count matrix [46] for

all accessory (non-structural) *nif* genes. Wagner parsimony with a gain-to-loss penalty

ratio of 2 was applied to infer the most parsimonious scenario of gene presence at internal

nodes, as described previously [47]. This ratio is widely used because recurring

observations indicate that gene loss occurs at least twice as frequently than gene gain

[48–50]. Ancestral trait reconstructions were performed on an ultrametric and

dichotomous tree, which was made using the *multi2di* and *chronos* (model = "relaxed")

functions of the APE package [25] in R. The *fitdiscrete* function in geiger [51] was used

for discrete reconstructions of oxygen preference, which were mapped onto the

phylogeny with the *simmap* function in phytools [38]. The *fastAnc* function in phytools was

used for continuous variable reconstructions of metabolic and habitat diversity, which was

mapped onto the tree using the *contMap* function in phytools [38].

Sampling genomes from non-diazotrophs

To assess whether the observed co-evolutionary trends between traits and N_2 fixation

gene abundance were specific to diazotrophs, we analyzed a random sample of non-

diazotrophic genomes using the same phylogenetic linear regression methodology

described above. Among the 6,785 genomes missing one or more of the catalytic *nifHDK*

subunits, 700 were randomly selected with the base R *sample* function (version 4.4.2)

[39], for further analysis (**Supplemental Table 9**). Trait data (including OGT, oxygen

status, pH preference, and habitat preference) and gene presence/absence for N_2

fixation-related genes were collected as described for diazotrophs (**Supplemental**

Tables 9, 10). Fifty-one genomes without 16S rRNA gene sequences were excluded from

analyses involving habitat coverage.

<u>Results</u>

Distribution of nitrogenases across the tree of life

We used a large prokaryote reference phylogeny, "WoL" [20], containing over 7,000 high-
quality genomes [21] to first assess the representative diversity and distribution of
nitrogenase today. A total of 673 genomes (~9.0%) were found to contain nitrogenase
(**Supplemental Table 2**) based on the presence of the catalytic HDK subunits of
nitrogenase (**Figure 2A**). Our estimate of nitrogenase diversity falls within the range of
recent estimates (~5.9% to 16%) [7, 12, 52, 53]. Here, approximately 46.9% of these
genomes   belonged   to   Proteobacteria   (in   order   of   abundance:
Alpha>Delta>Gamma>Beta>Epsilon>Acidithiobacillia), while the remaining top three
phyla included Firmicutes (17.3%), Cyanobacteria (12.6%), and Euryarchaeota (11.4%).
Amongst the nitrogenase-containing genomes, molybdenum (Mo)-only nitrogenase made
up the majority (90.3%), followed by genomes with both Mo-nitrogenase and iron (Fe)
nitrogenase (5.9%), genomes with both Mo-nitrogenase and vanadium (V) nitrogenase
(2.2%), and lastly with 1.5% of genomes containing all three nitrogenases (**Figure 1**)
(**Figure 2B**). Genomes with V-nitrogenase were found in 6 phyla: Euryarchaeota,
Firmicutes,   Alphaproteobacteria,   Deltaproteobacteria,   Cyanobacteria,   and
Gammaproteobacteria. Fe-nitrogenases were found in 11 phyla: Firmicutes,
Epsilonproteobacteria, Alphaproteobacteria, Gammaproteobacteria, Verrucomicrobia,
Euryarchaeota,   Deltaproteobacteria,   Gammaproteobacteria,   Bacteroidetes,
Spirochaetes, and Chlorobi. Euryarchaeota represented 50% of the genomes with all
three nitrogenases; the remaining phyla included Alphaproteobacteria, Firmicutes, and
Gammaproteobacteria.

Figure 2. Diversity and phylogenetic distribution of nitrogenase. **A)** The overall distribution of nitrogenase across the microbial diversity from the Zhu et al (2019) Web of Life (WoL) phylogeny [20]. A total of 7,458 high-quality genomes were assessed. Of these, ~9.0% of all genomes (n=673) encoded genes for the full nitrogenase complex (*nifHDK*). Black bars on the outer ring represent the presence of nitrogenase. Numbers reflect the top 14 taxonomic groups in the dataset with the highest proportion of genomes with nitrogenase. The phylogeny was plotted with iTOL [54]. **B)** Proportions and taxonomic distributions of the different nitrogenase isozymes across WoL. Mo-nitrogenase constitutes the majority of nitrogenases identified, followed by Fe-nitrogenase, and then V-nitrogenase in abundance. Only ~1% of genomes contained all three nitrogenases, but ~50% of genomes with all three isozymes belonged to Euryarchaeota.

An N₂ fixation genomic and gene-centric ecological catalog

We screened all 673 genomes with the full *nifHDK* complement for 93 known genes ("machinery") involved in N_2 fixation (**Table 1, Supplemental Table 3, Supplemental Figure 1**). In Figure 3, genes are grouped by functional category in relation to N_2 fixation, and the relative abundance of genes within each category is shown. Excluding the structural genes *nifHDK*, the total number of N_2 fixation genes detected per genome ranged from 14 to 67 genes. The largest N_2 fixation gene repertoires were observed in

the Alphaproteobacteria *Rhodopseudomonas palustris CGA009* and the

Gammaproteobacteria *Azotobacter vinelandii* DJ, which contained 67 and 65 genes,

respectively. In contrast, the archaea taxa *Methanococcus aeolicus* Nankai-3,

Methanothermococcus okinawensis IH1 and *Methanocaldococcus infernus* ME each

possessed the smallest repertoires with 14 genes. Across the species phylogeny, the

number of N_2 fixation genes per genome scaled positively with branch lengths, indicating

lineages with greater cumulative evolutionary change tend to possess larger N_2 fixation

gene complements (**Supplemental Figure 2**).

Table 1. Functional groupings of N_2 fixation machinery genes.

Functional group	N_2 fixation machinery genes
Structural	*nifHDK*, *vnfHDGK*, *anfHDGK*
Cofactor synthesis and maturation	*nifBEMNPQVXZ*, *vnfE*, *iscAR*, *iscU/nifU*, *iscS/nifS*, *sufABCDES*
Electron donation and transfer	*fdx*, *fdxA*, *fer*, *fixABCX*, *fldA*, *fnr*, *nifJ*, *rnfABCDEG*
Metal acquisition	*feoABC*, *modABCDEF*, *mosAB*
Protection	*nodABCD*, *ahpCD*, *cooA*, *cowN*, *devAC*, *hghD*, *katG*, *nifW*, *oxyR*, *sod1,2*, *sodN*
Regulation	*draGT*, *fixJLK*, *nif/anf/vnfA*, *nifL*, *nifI₁I₂*, *nifR3*, *ntcA*, *ntrBC*, *regRS*
Unknown	*nifO*, *nifT*

Species Phylogeny   **Functional Group Matrix**   **Species Traits**

Euryarchaeota

Chloroflexi

Cyanobacteria

Firmicutes

Chlorobi

ε-proteo

δ-proteo

β-proteo

γ-proteo

α-proteo

0.1

A B C D E F G   1 2 3 4 5   Trait Frequency

Figure key

Functional Group Matrix
A. Structural
B. Cofactor synthesis/maturation
C. E- donation/transport
D. Metal aquisition
E. Protection
F. Regulation
G. Unknown

Trait Matrices
1. Genome size
2. Growth temperature
3. Oxygen preference
4. pH preference
5. Habitat preference

Gene Abundance
1.00
0.75
0.50
0.25
0.00

Genome Size (Mb)
12
9
6
3

Growth Temp (°C)
80
60
40
20

Oxygen Preference
Unknown
Aerobe
Anaerobe
Microaerophile
Facultative

pH
8
6
4
2

Habitat Preference
Saline
Freshwater
Hydrothermal
Terrestrial
Plant
Host-associated

Trait Frequency
N₂ fix genes
Metabolisms
Habitats

Figure 3. Distribution of N₂ fixation machinery mapped onto the phylogeny of nitrogenase-containing
genomes, highlighting associations with species-associated ecological (habitat preference, genome size,

coverage of metabolic pathways) and physiological traits (oxygen, temperature, and pH preferences). N_2
fixation gene abundance correlates with those that are mesophilic, prefer aerobic and/or facultative
lifestyles, possess larger genome sizes, broader metabolic capabilities, and occupy a wider range of
habitats. For this figure, N_2 fixation genes were organized into functional categories and the relative
abundance of genes for each category calculated. The full gene matrix visualization can be found in
Supplemental Figure 1 and the gene matrix and associated trait categories in Supplemental Table 3. Trait
frequencies were calculated as the relative abundances of N_2 fixation genes (blue line), metabolic pathways
(green line), and habitats (orange line) associated with genome hosts. The phylogeny and matrices were
plotted using the ggTree R package (version 3.6.2) [26].

To explore the physiological and ecological diversity of diazotrophs identified in this study,
we mapped their habitat preferences, oxygen tolerance, optimal growth pH and
temperature, metabolic versatility, and genome size onto a phylogenetic tree depicting
relationships among diazotroph lineages (**Figure 3**).

Amongst nitrogenase-containing genomes, 47.5% belonged anaerobes, 31.6% to
aerobes, 5.3% to facultative species, and 1.5% to microaerophiles. The remaining 14%
could not be classified due to missing data or uncertainty in predictions. Temperature
preferences further define nitrogenase distribution with most genomes (91.2%) classified
as mesophiles favoring growth between 18ºC and 45ºC. The remaining ~8.8%
nitrogenase-containing genomes belonged to thermophiles that grow best at
temperatures above 45ºC. Similarly, neutrophiles (pH 5.5-8.0) dominated the pH
preference dataset (87.2%), whereas acidophiles and alkaliphiles were comparatively
rare.

Habitat distributions varied across taxa. Freshwater and terrestrial environments were the
most common inferred habitats of preference, encompassing 32.5% and 26.2% of
genomes, respectively. Other environments included saline systems (14.3%), non-plant
host associations (11.4%), plant-associated niches (8.8%), and hydrothermal systems
(6.8%). These patterns likely both reflect genuine ecological preferences and uneven
environmental sampling, which may skew our understanding of where nitrogenase-
containing organisms are most abundant in nature.

Diazotrophs also differed in their metabolic and ecological breadth. Metabolic coverage

ranged from 9.8 to 46.4% with an average of 23.2%, and habitat coverage varied from 0

to 48% with an average of 12%, excluding the ~3% of genomes for which habitat data

were unavailable. Genome size spanned from 1.3 to 12.3 Mbp with a mean of 4.6 Mbps.

Figure 4. N_2 fixation gene abundance varies across traits. The number of genes was compared across **A)**
temperature**, B)** oxygen, and **C)** habitat preferences**.** Thermophiles and anaerobes have discernably fewer
N_2 fixation genes than mesophiles and all other oxygen phenotypes, respectively. Genomes associated
with hydrothermal environments have on average, the fewest number of N_2 fixation genes, whereas plant-
associated genomes have the highest average. Pairwise differences between means were calculated using
non-parametric Wilcoxon rank-sum tests. For each panel, the means of those that do not differ with

290   statistical evidence share the same letter, whereas those that do not share a letter are significantly different
291   (*P*<0.05). Boxplots were plotted with the ggpubr R package [37].

Physiological, metabolic, and habitat preferences vary with N_2 fixation genetic complexity

Next, we examined how the number of N_2 fixation genes per genome relates to
environmental and physiological traits, to identify the factors influencing the genetic
complexity of the N_2 fixation machinery. Here, genetic complexity is defined as the
proportion of N_2 fixation genes detected within a given genome.

We observed that genomes sharing similar temperature, oxygen, and habitat preferences
had comparable numbers of N_2 fixation genes (**Figure 3**). Specifically, thermophiles and
anaerobes contained significantly fewer genes than their mesophilic (p.adj=8.60×10^{-23})
and aerobic (p.adj=4.80×10^{-45}) counterparts (**Figure 4AB**). Consistently, taxa preferring
hydrothermal environments on average have fewer N_2 fixation genes than those from
other habitats (**Figure 4C**). No significant differences were detected among terrestrial and
freshwater, saline, or non-plant host-associated environments. Plant-associated taxa
exhibited the highest average of N_2 fixation gene counts, exceeding those from other
environments except non-plant host-associated taxa.

N_2 fixation genetic complexity as a predictor of co-evolutionary trait associations

To test the degree to which traits are phylogenetically conserved versus independently
evolved, we measured phylogenetic signal for gene abundance, habitat coverage,
metabolic coverage, genome size, temperature, pH, and oxygen preference across the
species phylogeny using Pagel's lambda (λ), as implemented in the *phylolm* R package
[41]. All traits had a moderate to high phylogenetic signal (λ≥0.69; **Supplemental Table
4**), indicating closely related genomes tend to share more similar trait values than those
drawn randomly from the phylogeny.

We further assessed phylogenetic signal within individual phyla to test for lineage-specific
evolutionary patterns and differences in selection pressures. This analysis clarified

whether the overall phylogenetic signal measured across the species phylogeny was
disproportionately driven by particular phyla. Because some phyla had too few genomes
(<20), phylum-specific analyses were restricted to Alpha-, Beta-, Delta-,
Gammaproteobacteria, as well as Cyanobacteria, Firmicutes, and Euryarchaeota.
Genome size, habitat coverage, and metabolic coverage all exhibited strong phylogenetic
signal across these major nitrogenase-containing phyla, ($\lambda \geq 0.8$, **Supplemental Table 6**),
indicating that these traits are broadly conserved across these lineages, rather than
confined to a single clade. N_2 fixation gene abundance likewise showed a high λ (>0.8)
within most phyla with the exception of Cyanobacteria ($\lambda = 0.34$), potentially due to limited
variability of N_2 fixation gene sets within this clade.

By contrast, no phylogenetic signal for temperature was detected in
Gammaproteobacteria ($\lambda = 0.001$), consistent with their largely mesophilic lifestyle,
whereas other phyla retained strong signals ($\lambda > 0.8$). Oxygen tolerance showed little
phylogenetic structure in either Gammaproteobacteria or Firmicutes. In contrast, pH
exhibited strong phylogenetic conservation in Cyanobacteria, Euryarcheaota, and
Gamma- and Betaproteobacteria, but low conservation in Deltaproteobacteria.

Next, we assessed whether certain traits shared an evolutionary history with N_2 fixation
gene abundance via phylogenetic linear regression models (**Figure 5, Supplemental**
Figure 3, Supplemental Tables 5, 11). Gene abundance (calculated as the proportion
of the ninety-three N_2 fixation machinery genes screened in this study, per genome) was
chosen as the independent variable to determine if the complexity of N_2 fixation genes in
a genome could be used to predict other trait values. Response (trait) variables were
transformed using either natural log or square root to stabilize variance prior to analysis.
Genome size (**Figure 5A, Supplemental Figure 3A**) had a positive, significant linear
relationship with N_2 fixation gene abundance for diazotrophs ($R^2=0.104$, $P=2.2\times10^{-16}$).
The same trend was observed screening a random sample of non-diazotrophs from our
dataset ($R^2=0.23$, $P=8.3\times10^{-42}$), suggesting N_2 fixation gene complexity may interact with
broader genomic expansion/subtraction forces, irrespective of an organism's ability to fix
nitrogen.

For temperature, we initially observed a significant, negative linear relationship with N_2
fixation gene abundance (**Figure 5B, Supplemental Figure 3B**). Because prior work has
shown that adaptation to high temperatures is often linked to genome reduction [55], we
repeated the regression using N_2 fixation gene abundance normalized to genome size
(**Supplemental Table 2**). When normalized, the negative relationship with temperature
disappeared, suggesting that the original pattern likely reflected genome size constraints
rather than a temperature specific effect on N_2 fixation. A similar pattern emerged for
metabolic and habitat coverage. Both traits showed positive relationships with N_2 fixation
gene number, significant for metabolic coverage (R^2=0.006, P=0.025) and marginally
significant for habitat coverage (R^2=0.004, P=0.066) (**Figures 5CD, Supplemental
Figures 3CD**). After accounting for genome size, these relationships disappeared,
suggesting that gene abundance scales broadly with genomic content rather than specific
ecological traits.

Figure 5. Phylogenetic linear regressions assessed whether the number of N_2 fixation genes in a genome could predict other traits, such as **A)** genome size, **B)** metabolic coverage, **C)** optimal growth temperature,

and **D)** habitat coverage. Positive relationships were observed between N_2 fixation gene abundance with
genome size, metabolic, and habitat coverage. Response variables were either transformed using natural
log or square root to stabilize variance prior analysis. Solid blue lines represent non-phylogenetic linear
models, dashed blue lines represent phylogenetic linear models. Correlation coefficients (R^2), associated
P-values for both regression models, and lambda values are shown for each plot. Scatter plots were plotted
using ggplot2 in R [56].

Differences in N_2 fixation machinery distribution

We used phylogenetic principal component analysis (PCA) to cluster genomes based on
the similarity of their N_2 fixation gene presence-absence patterns, while accounting for
shared evolutionary history. Phylogenetic relationships accounted for most of the
variation, as indicated by the low variance captured by the phylogenetic PCA
(**Supplemental Figure 10**). This finding suggests that clades are clustered primarily
because they share conserved gene sets inherited from common ancestors.

We then examined the key genes driving separation among genomes in the phylogeny-
corrected PCA and interpreted their distribution across taxa and oxygen preferences. The
ten genes with the strongest influence on each component of the phylogeny-corrected
PCA included the different regulation elements (*nifI₂*, *nifL*, *regRS*, *ntrB*), two different
electron-donating/transfer mechanisms (*rnfABCDEG, fixABCX*), nodulation and
heterocyst differentiation genes (*nodABCD*, *devC*), oxidative protection mechanisms
(*sodN*, *cooA*), cofactor synthesis and iron acquisition genes (*nifM*, *sufD*, *feoC2*)*,* and the
genes that make up both the Fe- and V-nitrogenases (*vnfHGDKE*, *anfHGDK*). The post-
translational *nifI₂* regulator is primarily found in anaerobes, such as Archaea and
Firmicutes [12, 13], while other regulatory mechanisms such as *regS*, *nifL*, and *ntrB* are
primarily found in aerobic/facultative Proteobacteria (**Supplemental Figure 1**). Pearson
correlations of gene co-occurrence (**Supplemental Figure 11**) confirmed that *nifI₂* and
nifI₁ had strong negative correlations with genes *nifTQWXZ*, *fnr*, *fdx*, *oxyR,* and *ntrBC*,
many of which were correlated with the ability to use oxygen in diazotrophs [12].

A preference for either *rnfACDEG* or *fixACX* electron donation systems was evident in
the PCA where the genes pointed opposite from one another (**Supplemental Figure**

10,11). Nodulation genes *nodABC* also pointed opposite to *rnfACDEG* and instead had a

positive correlation with *fixACX* (**Supplemental Figure 11**). This is confirmed by the fact

that diazotrophs forming symbioses with plants through nodulation prefer *fix* genes over

rnf genes for electron donation [57, 58]. None of the N_2 fixation genes were unique to

thermophiles and neither Fe- nor V- nitrogenases were found in genomes with optimal

growth temperatures greater than 41ºC.

Dynamic history of gene gain and loss

We estimated net gain or loss of N_2 fixation machinery genes at each node of the

nitrogenase-containing species tree using Wagner parsimony (see methods). Our results

show a dynamic history of gene gain and loss over time (**Figure 6**). Specifically, net gene

gain was predicted at many of the early branching nodes, whereas gene loss primarily

occurred closer to terminal tree branches. This suggests an initial expansion of genes

early in diazotroph evolution, followed by successive gene loss events, potentially

corresponding to an expansion of diazotroph lineages into new environments where

certain components of the N_2 fixation machinery became obsolete.

The nodes with the highest number of gene gains (+11) occurs at the most recent

common ancestor (MRCA) of Bacteria and at the MRCA of the Euryarchaeota clade

Methanosarcina (not shown). Some of the clades that experienced large gene losses

include Acidithiobacillia (-18), the Firmicute-Clostridia clade Thermoanaerobacterales (-

18) (not shown), Thermodesulfobacteria (-17) (not shown), Nitrospirae (-16), and the

MRCA of Aquificae and Epsilonproteobacteria (-11) (**Supplemental Tables 7-8**).

Acidothiobacillia's loss of genes *cowN*, *fixJL*, and the Suf and Rnf systems occurs in

tandem with a loss of habitat coverage (**Figure 6**). The loss of post-transcriptional

regulators *nifI$_1$I$_2$* unsurprisingly coincides with a gain in aerobic lifestyles at the MRCA of

Cyanobacteria, Actinobacteria and the MRCA of Gamma- and Alphaproteobacteria

(**Figure 6**) as *nifI$_1$I$_2$* is the main regulatory mechanism in anaerobes. The appearance of

transcriptional regulation by *nifA*, mostly used by aerobes, predates the loss of *nifI$_1$I$_2$* in

the MRCA of Gamma- and Alphaproteobacteria (**Figure 6**). The gain of genes involved

in protection against oxidants (*katG*, *oxyR*, and *cowN*) and oxygen response regulators

fixJL occurs early along the MRCA nodes of Verrucomicrobia, Chlorobi, Bacteroidetes,

and Proteobacteria (**Figure 6**). We noted two distinct gains of *rnfACDEG*, *nifTWZ*, *katG* and *fixCX* in distantly related diazotrophs, indicating they were either inherited from a common ancestor that does not have nitrogenase (and therefore would not be represented in this tree), and/or they were gained through horizontal gene transfer.

Figure 6. N_2 fixation gene gain (green triangles) and gene loss (red triangles) throughout the evolution of
diazotrophs. Events are mapped to the WoL [20] species tree which was pruned to include only tips
associated with diazotrophs identified in this study (n=673). Gene gain dominated the earliest nodes,
whereas gene loss was most common near terminal branches. No change (grey circle) represents no gain
or loss at that node, or an equal number or gains and losses. Asterisks highlight potential candidates for
HGT or genes derived from non-diazotroph ancestors not in this subsetted phylogeny. All gene events can
be found in Supplemental table 8. Key transitional events of gene abundance and habitat coverage can be
found in Supplemental Figure 12.

Discussion

Nitrogenase Distribution, Diversity, and Evolution Across Prokaryotic Lineages

Nitrogenase is found across a taxonomically diverse range of microbes with ecologies
spanning root-associated symbiosis in the rhizosphere [59] to deep-sea sediments
kilometers below the ocean surface [60–62]. The factors underlying this broad yet uneven
distribution remain unclear but likely reflect a complex evolutionary history shaped by
dynamic interplay between nitrogenase, its host cell, and the surrounding environment.

To investigate these patterns, we developed a comprehensive catalog of 673
nitrogenase-encoding genomes and their associated N_2 fixation machinery, tracing
historical genomic changes that shaped the evolution of this important metabolism
(**Figure 3**). In addition to the *nif* cluster and its accessory genes, we included 63 genes
outside of this cluster with reported roles in N_2 fixation, yielding a total of 93 genes
representing the N_2 fixation genetic machinery (**Table 1, Supplemental Table 1**). These
data were integrated with host-associated trait information (e.g., oxygen metabolism,
temperature, pH, genome size, habitat coverage), linking the evolutionary history of
nitrogenase and its supporting genes to the environmental diversity of N_2 fixation
observed today.

Oxygen inhibits nitrogenase activity [63], and modern diazotrophs employ a range of
strategies to protect the enzyme from this interference [19, 64, 65]. Consistent with
previous findings [12], we observed that on average, aerobic diazotrophs harbor more N_2
fixation-associated genes than anaerobes (**Figure 4B**), likely reflecting the additional

genetic components required for oxygen protection and regulation under aerobic
conditions. For example, we found that genes *katG*, *oxyR*, and *cowN*, which provide
protection against oxidants [66, 67], along with the oxygen response regulators *fixJL* [15]
were acquired along the lineages leading up to the divergence of Proteobacteria (**Figure
6**), a period that coincides with the historic accumulation of oxygen in the atmosphere
~2.45 billion years ago known as the Great Oxidation Event (GOE) [68]. This pattern
suggests these genes played a role enabling ancient diazotrophs to expand into
increasingly oxygenated environments. Recent studies further support the divergence of
the alternative nitrogenases during this transition to an oxic atmosphere [69],
underscoring how environmental transitions, particularly the historical rise of atmospheric
oxygen levels, reshaped the molecular evolution of N_2 fixation in deep time.

We recapitulate the relationship between N_2 fixation gene abundance and temperature
previously reported by Boyd et al. [12], that on average, thermophiles possess the
simplest N_2 fixation-related gene sets (**Figure 4AB, Figure 5B**). Our analyses indicate
that this pattern is in part due to genome reduction associated with long-term adaptation
to high temperatures [55]. Yet, if the earliest diazotrophs were indeed thermophilic [12],
then temperature's role shaping the evolution of N_2 fixation cannot be dismissed. For
example, some cold-adapted diazotrophs switch from Mo- to V-based nitrogenases [70–
72], indicating how shifts in temperature regimes may have shaped the functional
diversification of nitrogenase over evolutionary time.

Outside our set of nitrogenase-containing genomes, we identified genomes with partial
detection of the core N_2 fixation genes. Notably, several anaerobic thermophiles lack *nifE*,
nifN, or both (**Figure 3**). These genes are involved in FeMo cofactor assembly and are
part of a proposed minimal *nif* gene set (*nifHDKENB)* required for N_2 fixation [73].
Lineages exhibiting this partial gene set include Euryarcheaota (such as members of the
ANME groups [74]), Chloroflexi, and Firmicutes. Some have verified N_2 fixation activity ,
while others, like *Roseiflexus castenholzii*, do not [75–78]. Ancestral sequence
reconstructions suggest that *nifDK* may have originated from a duplicated maturase-like
nifEN ancestor [79], raising the possibility that some extant diazotrophs lacking *nifEN*
evolved alternative cofactor maturation strategies, perhaps as a way to maximize energy

conservation under thermophilic conditions [55]. Support for this idea comes from
Roseiflexus sp. RS-1, which lacks *nifEN* and appears to use *nifDK* to fulfill both the
cofactor maturase and catalytic roles [80]. Further work will be required to determine
whether this streamlined mechanism occurs in other diazotrophs missing *nifEN*.

Plant-associated genomes showed the highest N_2 fixation gene counts (**Figure 4C**),
consistent with the pivotal role of rhizobia in plant host nitrogen supply [81, 82]. Our
reconstructions capture the divergence of Actinobacteria *Frankia* from its ancestor,
marked by a transition to aerobic metabolism, accompanying genome and habitat
expansion, the loss of $nifI_1I_2$ regulators, and the gain of both nodulation (*nodC*) and
oxidative stress response (*sodN*) genes to support its dual free-living and symbiosis
states (**Figure 6**).

We observed that *Frankia* lacks key regulators such as *nifAL* [17]. However, we noted the
presence of the post-translational regulator *draG*, without its partner *draT*, indicating that
draG-draT regulation is likely non-functional and perhaps a novel, as-yet-unknown
mechanism for nitrogenase inactivation is present [83] – a possibility that warrants further
study.

In many plant symbioses, oxygen levels are tightly regulated by the host to activate the
FixL - FixJ pathway, ultimately driving nitrogenase expression [84], while *nod* genes
mediate nodulation [85]. Terrestrial habitats, being more variable than oceans [86],
impose fluctuating environmental conditions that demand dynamic nitrogenase regulation
[87–89]. Soil molybdenum scarcity likely favors the deployment of alternative nitrogenases
[90], contributing to the expansion of gene repertoire in terrestrial environments. Our
results illustrate this pattern. For example, several lichen-symbionts within the Nostocales
order of Cyanobacteria possess V-nitrogenase [91], likely as a way to cope with limited
Mo in high latitude boreal forests [92]. Together, these findings highlight how diazotroph
N_2 fixation gene repertoires have been shaped by environmental constraints and the
selective pressures of symbiotic lifestyles.

Eco-evolutionary history of nitrogenase-associated genes

Central to our framework is that N_2 fixation gene complexity is not an isolated trait, rather,
it is embedded within a larger evolutionary framework of genome expansion, metabolic
versatility, and habitat diversity (**Figures 4, 5**). Gene complexity can even serve as an
ecological marker distinguishing generalists, with broad habitats and large, versatile
genomes, from specialists with narrow niches and smaller genomes [93–95]. This
distinction is particularly relevant today as specialist diazotrophs may contribute
disproportionately to total soil N_2 fixation and thereby differentially influence nitrogen input
into these environments [96].

Our reconstructions show that as diazotrophs diversified, so did their N_2 fixation
machinery (**Figure 3**). Gene gain dominated early evolution, with clades such as Alpha-,
Gamma-, and Betaproteobacteria maintaining extensive N_2 fixation genes (**Figure 6**). In
contrast, later stages were marked by lineage-specific losses, particularly in terminal
branches, suggesting a shift towards ecological specialization and genome streamlining
in diazotrophs. Losses were most pronounced in Chloroflexi, Nitrospirae, Aquificae,
Acidithiobacillia, Thermodesulfobacterota, and Epsilonproteobacteria, groups often
associated with anaerobic, thermophilic, acidophilic, or host-adapted lifestyles
(**Supplemental Table 2**).

In some cases, N_2 fixation gene loss tracked closely with reduced habitat coverage, as in
Epsilonproteobacteria and Acidithiobacillia (**Figure 6**), consistent with prior work on
genome streamlining in these clades [97–99]. Adaptation to deep-sea hydrothermal vents
[100], shifts toward heterotrophy [97], and transitions to acidophily [101] all show how
niche transitions often coincide with simplification of N_2 fixation gene sets. There are
exceptions to this trend between genome size and the size of the N_2 fixation machinery.
For example, *Zymomonas mobilis* (Alphaproteobacteria) maintains the highest proportion
of N_2 fixation genes (~2-3% of all coding genes) despite having a compact ~2Mb genome
(**Supplemental Table 2**). Similarly, in symbiont systems, genome reduction often reflects
host compensation [102]. UCYN-A2, a marine symbiont recently identified as an early

stage "nitroplast" in *Braarudosphaera bigelowii* [103, 104], has lost CO_2 fixation and

photosystem II [105, 106], yet retains an intact *nif* cluster (*nifTZVBSHDKENXW)* [107].

Horizontal gene transfer (HGT) has further shaped N_2 fixation evolution [11, 108], likely

explaining several gene acquisitions we observed (e.g., *rnfACDEG*, *nifTWZ*, *katG* and

fixCX in **Figure 6**). Previous studies have documented entire *nif* clusters moving across

major clades – including a >20kb transfer from Gammaproteobacterium in to the diatom

symbiont "*Candidatus* Tectiglobus diatomicola" [86, 109]. In our dataset, *Coleofasciculus*

chthonoplastes PCC 7420 (formerly *Microcoleus chthonoplastes*) appears to have

acquired *nifHDKEN* from a Deltaproteobacterium [88, 110]. Together, these patterns

show that the complexity of N_2 fixation genes reflects a balance between the forces of

gene loss and gain, specialization, horizontal transfer - all embedded in the broader

trajectory of diazotroph genome evolution.

Limitations of comparative genomics

As with any comparative genomic approach, there are inherent limitations in identifying

the full complement of genes and traits within genomes. One potential limitation is that

our analysis focuses on genes already recognized to be playing a role in N_2 fixation,

meaning additional, yet-undiscovered genes may contribute to the process, particularly

in non-model diazotrophs, where many functional mechanisms remain uncharacterized.

Another caveat is our reliance on KEGG KO number. Not all genes could be assigned

KO numbers, and some KO identifiers fail to distinguish between genes with high

sequence similarity, such as *nifA*/*vnfA*/*anfA*. Continued improvements in gene functional

annotation and database curation will be essential for building a more complete

understanding of diazotroph capabilities [111].

Conclusion

Examining the distribution of diverse N_2 fixation machinery offers insight into which genes

were critical for enabling transitions into new environments throughout evolutionary

history. The rise of atmospheric oxygen, for example, fundamentally reshaped electron-

donor availability [112] and, together with changes in metal availability [113, 114], likely

exerted strong selective pressures on N_2 fixation systems, as demonstrated here.

Reconstructing historical patterns from modern genetic records is thus pivotal for
understanding the processes that govern evolution [115]. Global trends in genome
evolution not only reveal how organisms responded to past environmental fluctuations
but connects them to their present-day ecological niches.

In his book *Life's Engines*, Paul Falkowski likens nitrogenase and its core machinery to a
computer's hardware, while accessory genes function as the software, essential for
maintenance, optimization and updates [116]. This analogy is compelling, capturing the
deep interconnection between core biochemical processes and their regulatory networks.
These elements must evolve in concert to navigate the challenges posed by changing
environments, a dynamic reflected in the expansions and contractions within diazotroph
genomes which we have documented across both ancient and modern contexts.

Acknowledgements

We gratefully acknowledge support from the NASA Interdisciplinary Consortium for
Astrobiology Research (ICAR): Metal Utilization and Selection Across Eons, MUSE under
grant no 80NSSC17K0296, and the NASA Exobiology Program under grant no
NNH23ZDA001N. M.S. was supported by the NASA Postdoctoral Program, administered
by Oak Ridge Associated Universities under contract with NASA. We thank Patricia Tran,
Shaomei He, colleagues in the Kaçar Lab, members of NASA MUSE ICAR, and
colleagues at Texas State University for helpful feedback on manuscript; and the Center
for High Throughput Computing (CHTC) and the Department of Bacteriology at the
University of Wisconsin–Madison for providing computing resources. We would like to
finally thank the four anonymous reviewers for their constructive feedback.

Data availability

597 All custom scripts, tree files, raw data, and PDF versions of the figures are available at
598 https://github.com/kacarlab/Sobol_2025_NifGeneEcoEvo. All supplemental tables are
599 available in the uploaded datasets. Specifically, supplementary Tables 2 and 9 include
600 genome accession numbers, taxonomy, and trait data information.

References

603   1.   Canfield DE, Glazer AN, Falkowski PG. The Evolution and Future of Earth's Nitrogen

604   Cycle. *Science* 2010. https://doi.org/10.1126/science.1186120

605   2.   Raymond J et al. The Natural History of Nitrogen Fixation. *Mol Biol Evol* 2004;**21**:541–

606   554. https://doi.org/10.1093/molbev/msh047

607   3.   Navarro-González R, McKay CP, Mvondo DN. A possible nitrogen crisis for Archaean life

608   due to reduced nitrogen fixation by lightning. *Nature* 2001;**412**:61–64.

609   https://doi.org/10.1038/35083537

610   4.   Koehler MC et al. Transient surface ocean oxygenation recorded in the ~2.66-Ga

611   Jeerinah Formation, Australia. *Proc Natl Acad Sci* 2018;**115**:7711–7716.

612   https://doi.org/10.1073/pnas.1720820115

613   5.   Garcia AK et al. Reconstructing the evolutionary history of nitrogenases: Evidence for

614   ancestral molybdenum-cofactor utilization. *Geobiology* 2020;**18**:394–411.

615   https://doi.org/10.1111/gbi.12381

616   6.   Cuevas Zuviría B et al. Structural evolution of nitrogenase over 3 billion years. *eLife*

617   2025;**14**:RP105613. https://doi.org/10.7554/eLife.105613

618   7.   Koirala A, Brözel VS. Phylogeny of Nitrogenase Structural and Assembly Components

619   Reveals New Insights into the Origin and Distribution of Nitrogen Fixation across Bacteria

620   and Archaea. *Microorganisms* 2021;**9**:1662.

621   https://doi.org/10.3390/microorganisms9081662

622   8.   Garcia AK et al. Nitrogenase resurrection and the evolution of a singular enzymatic

623   mechanism. *eLife* 2023;**12**:e85003. https://doi.org/10.7554/eLife.85003

624   9.   Harris DF et al. Ancient nitrogenases are ATP dependent. *mBio* 2024;**15**:e01271-24.

625   https://doi.org/10.1128/mbio.01271-24

626   10.   Stüeken EE et al. Isotopic evidence for biological nitrogen fixation by molybdenum-

627   nitrogenase from 3.2 Gyr. *Nature* 2015;**520**:666–669. https://doi.org/10.1038/nature14180

628   11.   Parsons C et al. Radiation of nitrogen-metabolizing enzymes across the tree of life tracks

629   environmental transitions in Earth history. *Geobiology* 2021;**19**:18–34.

630   https://doi.org/10.1111/gbi.12419

631   12.   Boyd ES et al. Evolution of Molybdenum Nitrogenase during the Transition from

632   Anaerobic to Aerobic Metabolism. *J Bacteriol* 2015;**197**:1690–1699.

633   https://doi.org/10.1128/jb.02611-14

634   13.   Nichio BT de L et al. Exploring diazotrophic diversity: unveiling Nif core distribution and

635   evolutionary patterns in nitrogen-fixing organisms. *BMC Genomics* 2025;**26**:81.

636   https://doi.org/10.1186/s12864-024-10994-9

637   14.   Kaiser JT et al. Structure of Precursor-Bound NifEN: A Nitrogenase FeMo Cofactor

638   Maturase/Insertase. *Science* 2011;**331**:91–94. https://doi.org/10.1126/science.1196954

639   15.   Dixon R, Kahn D. Genetic regulation of biological nitrogen fixation. *Nat Rev Microbiol*

640   2004;**2**:621–631. https://doi.org/10.1038/nrmicro954

641   16.   Rubio LM, Ludden PW. Biosynthesis of the Iron-Molybdenum Cofactor of Nitrogenase.

642   *Annu Rev Microbiol* 2008;**62**:93–111.

643   https://doi.org/10.1146/annurev.micro.62.081307.162737

644   17.   Mus F et al. Exploring the alternatives of biological nitrogen fixation. *Metallomics*

645   2018;**10**:523–538. https://doi.org/10.1039/c8mt00038g

646   18.   Kleinschmidt JA, Kleiner D. Relationships between nitrogenase, glutamine synthetase,

647   glutamine, and energy charge in Azotobacter vinelandii. *Arch Microbiol* 1981;**128**:412–

648   415. https://doi.org/10.1007/BF00405923

649   19.   Nonaka A et al. Accessory Proteins of the Nitrogenase Assembly, NifW, NifX/NafY, and

650   NifZ, Are Essential for Diazotrophic Growth in the Nonheterocystous Cyanobacterium

651   Leptolyngbya boryana. *Front Microbiol* 2019;**10**. https://doi.org/10.3389/fmicb.2019.00495

652 20. Zhu Q et al. Phylogenomics of 10,575 genomes reveals evolutionary proximity between

653   domains Bacteria and Archaea. *Nat Commun* 2019;**10**:5477.

654   https://doi.org/10.1038/s41467-019-13443-4

655 21. Bowers RM et al. Minimum information about a single amplified genome (MISAG) and a

656   metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol*

657   2017;**35**:725–731. https://doi.org/10.1038/nbt.3893

658 22. Nadkarni PM, Parikh CR. An eUtils toolset and its use for creating a pipeline to link

659   genomics and proteomics analyses to domain-specific biomedical literature. *J Clin*

660   *Bioinforma* 2012;**2**:9. https://doi.org/10.1186/2043-9113-2-9

661 23. Kanehisa M et al. KEGG as a reference resource for gene and protein annotation.

662   *Nucleic Acids Res* 2016;**44**:D457–D462. https://doi.org/10.1093/nar/gkv1070

663 24. Aramaki T et al. KofamKOALA: KEGG Ortholog assignment based on profile HMM and

664   adaptive score threshold. *Bioinformatics* 2020;**36**:2251–2252.

665   https://doi.org/10.1093/bioinformatics/btz859

666 25. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary

667   analyses in R. *Bioinformatics* 2019;**35**:526–528.

668   https://doi.org/10.1093/bioinformatics/bty633

669 26. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr Protoc Bioinforma*

670   2020;**69**:e96. https://doi.org/10.1002/cpbi.96

671 27. Reimer LC et al. BacDive in 2022: the knowledge base for standardized bacterial and

672   archaeal data. *Nucleic Acids Res* 2022;**50**:D741–D746.

673   https://doi.org/10.1093/nar/gkab961

674 28. Barrett T et al. BioProject and BioSample databases at NCBI: facilitating capture and

675   organization of metadata. *Nucleic Acids Res* 2012;**40**:D57–D63.

676   https://doi.org/10.1093/nar/gkr1163

29.   Mukherjee S et al. Genomes OnLine Database (GOLD) v.10: new features and updates.

Nucleic Acids Res 2025;**53**:D989–D997. https://doi.org/10.1093/nar/gkae1000

30.   Madin JS et al. A synthesis of bacterial and archaeal phenotypic trait data. *Sci Data*

2020;**7**:170. https://doi.org/10.1038/s41597-020-0497-4

31.   Sauer DB, Wang D-N. Predicting the optimal growth temperatures of prokaryotes using

only genome derived features. *Bioinformatics* 2019;**35**:3224–3231.

https://doi.org/10.1093/bioinformatics/btz059

32.   Jabłońska J, Tawfik DS. The number and type of oxygen-utilizing enzymes indicates

aerobic vs. anaerobic phenotype. *Free Radic Biol Med* 2019;**140**:84–92.

https://doi.org/10.1016/j.freeradbiomed.2019.03.031

33.   Ramoneda J et al. Building a genome-based understanding of bacterial pH preferences.

Sci Adv 2023;**9**:eadf8998. https://doi.org/10.1126/sciadv.adf8998

34.   Mise K, Iwasaki W. Environmental Atlas of Prokaryotes Enables Powerful and Intuitive

Habitat-Based Analysis of Community Structures. *iScience* 2020;**23**:101624.

https://doi.org/10.1016/j.isci.2020.101624

35.   Seemann T. tseemann/barrnap. 2025. 2025.

36.   Geller-McGrath D et al. Predicting metabolic modules in incomplete bacterial genomes

with MetaPathPredict. *eLife* 2024;**13**:e85749. https://doi.org/10.7554/eLife.85749

37.   Alboukadel Kassambara. ggpubr: 'ggplot2' Based Publication Ready Plots. 2025. 2025.

38.   Revell LJ. phytools 2.0: an updated R ecosystem for phylogenetic comparative methods

(and other things). *PeerJ* 2024;**12**:e16505. https://doi.org/10.7717/peerj.16505

39.   R CORE TEAM (2014) R A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna. - References - Scientific Research

Publishing. https://www.scirp.org/reference/referencespapers?referenceid=2145566. .

40.   Taiyun Wei, Viliam Simko. R package 'corrplot': Visualization of a Correlation Matrix.

2024. 2024.

41.   Tung Ho L si, Ané C. A Linear-Time Algorithm for Gaussian and Non-Gaussian Trait

704   Evolution Models. *Syst Biol* 2014;**63**:397–408. https://doi.org/10.1093/sysbio/syu005

705   42.   Rezende EL, Diniz-Filho JAF. Phylogenetic Analyses: Comparing Species to Infer

706   Adaptations and Physiological Mechanisms. *Compr Physiol* 2012;**2**:639–674.

707   https://doi.org/10.1002/j.2040-4603.2012.tb00401.x

708   43.   Freckleton RP, Rees M. Comparative analysis of experimental data. *Methods Ecol Evol*

709   2019;**10**:1308–1321. https://doi.org/10.1111/2041-210X.13164

710   44.   Jombart T, Balloux F, Dray S. adephylo: new tools for investigating the phylogenetic

711   signal in biological traits. *Bioinformatics* 2010;**26**:1907–1909.

712   https://doi.org/10.1093/bioinformatics/btq292

713   45.   Csűrös M. Ancestral Reconstruction by Asymmetric Wagner Parsimony over Continuous

714   Characters and Squared Parsimony over Distributions. In: Nelson CE, Vialette S (eds.)

715   *Comp. Genomics*. 2008. Berlin, Heidelberg: Springer, 2008, pp 72–86.

716   46.   Kapust N et al. Failure to Recover Major Events of Gene Flux in Real Biological Data Due

717   to Method Misapplication. *Genome Biol Evol* 2018;**10**:1198–1209.

718   https://doi.org/10.1093/gbe/evy080

719   47.   Oyserman BO et al. Ancestral genome reconstruction identifies the evolutionary basis for

720   trait acquisition in polyphosphate accumulating bacteria. *ISME J* 2016;**10**:2931–2945.

721   https://doi.org/10.1038/ismej.2016.67

722   48.   Pál C, Papp B, Lercher MJ. Adaptive evolution of bacterial metabolic networks by

723   horizontal gene transfer. *Nat Genet* 2005;**37**:1372–1375. https://doi.org/10.1038/ng1686

724   49.   Puigbò P et al. Genomes in turmoil: quantification of genome dynamics in prokaryote

725   supergenomes. *BMC Biol* 2014;**12**:66. https://doi.org/10.1186/s12915-014-0066-4

726   50.   Albalat R, Cañestro C. Evolution by gene loss. *Nat Rev Genet* 2016;**17**:379–391.

727   https://doi.org/10.1038/nrg.2016.39

728   51.   Pennell MW et al. geiger v2.0: an expanded suite of methods for fitting macroevolutionary

729   models to phylogenetic trees. *Bioinforma Oxf Engl* 2014;**30**:2216–2218.

730   https://doi.org/10.1093/bioinformatics/btu181

731   52.   Pi H-W et al. Origin and Evolution of Nitrogen Fixation in Prokaryotes. *Mol Biol Evol*

732   2022;**39**:msac181. https://doi.org/10.1093/molbev/msac181

733   53.   Bellanger M et al. NFixDB (Nitrogen Fixation DataBase)-a comprehensive integrated

734   database for robust 'omics analysis of diazotrophs. *NAR Genomics Bioinforma*

735   2024;**6**:lqae063. https://doi.org/10.1093/nargab/lqae063

736   54.   Letunic I, Bork P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic

737   tree display and annotation tool. *Nucleic Acids Res* 2024;**52**:W78–W82.

738   https://doi.org/10.1093/nar/gkae268

739   55.   Sabath N et al. Growth Temperature and Genome Size in Bacteria Are Negatively

740   Correlated, Suggesting Genomic Streamlining During Thermal Adaptation. *Genome Biol*

741   *Evol* 2013;**5**:966–977. https://doi.org/10.1093/gbe/evt050

742   56.   Villanueva RAM, Chen ZJ. ggplot2: Elegant Graphics for Data Analysis (2nd ed.). *Meas*

743   *Interdiscip Res Perspect* 2019;**17**:160–167.

744   https://doi.org/10.1080/15366367.2019.1565254

745   57.   Earl CD, Ronson CW, Ausubel FM. Genetic and structural analysis of the Rhizobium

746   meliloti fixA, fixB, fixC, and fixX genes. *J Bacteriol* 1987;**169**:1127–1136.

747   https://doi.org/10.1128/jb.169.3.1127-1136.1987

748   58.   Gubler M, Hennecke H. Regulation of the fixA gene and fixBC operon in Bradyrhizobium

749   japonicum. *J Bacteriol* 1988;**170**:1205–1214. https://doi.org/10.1128/jb.170.3.1205-

750   1214.1988

751   59.   Smercina DN et al. To Fix or Not To Fix: Controls on Free-Living Nitrogen Fixation in the

752   Rhizosphere. *Appl Environ Microbiol* 2019;**85**:e02546-18.

753   https://doi.org/10.1128/AEM.02546-18

60.   Reese BK et al. Nitrogen Cycling of Active Bacteria within Oligotrophic Sediment of the

Mid-Atlantic Ridge Flank. *Geomicrobiol J* 2018;**35**:468–483.

https://doi.org/10.1080/01490451.2017.1392649

61.   Kapili BJ et al. Evidence for phylogenetically and catabolically diverse active diazotrophs

in deep-sea sediment. *ISME J* 2020;**14**:971–983. https://doi.org/10.1038/s41396-019-

0584-8

62.   Semler AC et al. Abundance, Identity, and Potential Diazotrophic Activity of nifH-

Containing Organisms at Marine Cold Seeps. *Environ Microbiol* 2025;**27**:e70058.

https://doi.org/10.1111/1462-2920.70058

63.   Gallon JR. Reconciling the incompatible: N2 fixation And O2. *New Phytol* 1992;**122**:571–

609. https://doi.org/10.1111/j.1469-8137.1992.tb00087.x

64.   Milligan AJ et al. Light-dependent oxygen consumption in nitrogen-fixing cyanobacteria

plays a key role in nitrogenase protection. *J Phycol* 2007;**43**:845–852.

https://doi.org/10.1111/j.1529-8817.2007.00395.x

65.   Maier RJ, Moshiri F. Role of the Azotobacter vinelandiiNitrogenase-Protective Shethna

Protein in Preventing Oxygen-Mediated Cell Death. *J Bacteriol* 2000;**182**:3854–3857.

https://doi.org/10.1128/jb.182.13.3854-3857.2000

66.   Medina MS et al. CowN sustains nitrogenase turnover in the presence of the inhibitor

carbon monoxide. *J Biol Chem* 2021;**296**:100501.

https://doi.org/10.1016/j.jbc.2021.100501

67.   Zhao Y et al. OxyR-regulated catalase activity is critical for oxidative stress resistance,

nodulation and nitrogen fixation in Azorhizobium caulinodans. *FEMS Microbiol Lett*

2016;**363**:fnw130. https://doi.org/10.1093/femsle/fnw130

68.   Poulton SW et al. A 200-million-year delay in permanent atmospheric oxygenation.

Nature 2021;**592**:232–236. https://doi.org/10.1038/s41586-021-03393-7

69.   Cuevas-Zuviría B et al. Emergence of an Orphan Nitrogenase Protein Following

780   Atmospheric Oxygenation. *Mol Biol Evol* 2024;**41**:msae067.

781   https://doi.org/10.1093/molbev/msae067

782   70.   Miller RW, Eady RR. Molybdenum and vanadium nitrogenases of Azotobacter

783   chroococcum. Low temperature favours N2 reduction by vanadium nitrogenase. *Biochem*

784   *J* 1988;**256**:429–432. https://doi.org/10.1042/bj2560429

785   71.   Walmsley J, Kennedy C. Temperature-Dependent Regulation by Molybdenum and

786   Vanadium of Expression of the Structural Genes Encoding Three Nitrogenases in

787   Azotobacter vinelandii. *Appl Environ Microbiol* 1991;**57**:622–624.

788   https://doi.org/10.1128/aem.57.2.622-624.1991

789   72.   Darnajoux R, Bradley R, Bellenger J-P. In Vivo Temperature Dependency of Molybdenum

790   and Vanadium Nitrogenase Activity in the Heterocystous Cyanobacteria Anabaena

791   variabilis. *Environ Sci Technol* 2022;**56**:2760–2769.

792   https://doi.org/10.1021/acs.est.1c05279

793   73.   Dos Santos PC et al. Distribution of nitrogen fixation and nitrogenase-like sequences

794   amongst microbial genomes. *BMC Genomics* 2012;**13**:162. https://doi.org/10.1186/1471-

795   2164-13-162

796   74.   Laso-Pérez R et al. Evolutionary diversification of methanotrophic ANME-1 archaea and

797   their expansive virome. *Nat Microbiol* 2023;**8**:231–245. https://doi.org/10.1038/s41564-

798   022-01297-4

799   75.   Mehta MP, Baross JA. Nitrogen fixation at 92 degrees C by a hydrothermal vent

800   archaeon. *Science* 2006;**314**:1783–1786. https://doi.org/10.1126/science.1134772

801   76.   Dekas AE, Poretsky RS, Orphan VJ. Deep-sea archaea fix and share nitrogen in

802   methane-consuming microbial consortia. *Science* 2009;**326**:422–426.

803   https://doi.org/10.1126/science.1178223

77.   Ivanovsky RN et al. Nitrogen Metabolism of an Anoxygenic Filamentous Phototrophic

Bacterium Oscillocholris trichoides Strain DG-6. *Microbiology* 2021;**90**:428–434.

https://doi.org/10.1134/S0026261721040068

78.   Chen Y, Nishihara A, Haruta S. Nitrogen-fixing Ability and Nitrogen Fixation-related

Genes of Thermophilic Fermentative Bacteria in the Genus Caldicellulosiruptor. *Microbes*

Environ 2021;**36**:ME21018. https://doi.org/10.1264/jsme2.ME21018

79.   Garcia AK, Kolaczkowski B, Kaçar B. Reconstruction of Nitrogenase Predecessors

Suggests Origin from Maturase-Like Proteins. *Genome Biol Evol* 2022;**14**:evac031.

https://doi.org/10.1093/gbe/evac031

80.   Payá-Tormo L et al. Iron-molybdenum cofactor synthesis by a thermophilic nitrogenase

devoid of the scaffold NifEN. *Proc Natl Acad Sci* 2024;**121**:e2406198121.

https://doi.org/10.1073/pnas.2406198121

81.   Franche C, Lindström K, Elmerich C. Nitrogen-fixing bacteria associated with leguminous

and non-leguminous plants. *Plant Soil* 2009;**321**:35–59. https://doi.org/10.1007/s11104-

008-9833-8

82.   Bennett AB, Pankievicz VCS, Ané J-M. A Model for Nitrogen Fixation in Cereal Crops.

Trends Plant Sci 2020;**25**:226–235. https://doi.org/10.1016/j.tplants.2019.12.004

83.   Bueno Batista M, Dixon R. Manipulating nitrogen regulation in diazotrophic bacteria for

agronomic benefit. *Biochem Soc Trans* 2019;**47**:603–614.

https://doi.org/10.1042/BST20180342

84.   Mus F et al. Symbiotic Nitrogen Fixation and the Challenges to Its Extension to

Nonlegumes. *Appl Environ Microbiol* 2016;**82**:3698–3710.

https://doi.org/10.1128/AEM.01055-16

85.   Ferguson BJ et al. Molecular analysis of legume nodule development and autoregulation.

J Integr Plant Biol 2010;**52**:61–76. https://doi.org/10.1111/j.1744-7909.2010.00899.x

86.   Steele JH, Brink KH, Scott BE. Comparison of marine and terrestrial ecosystems:

830   suggestions of an evolutionary perspective influenced by environmental variation. *ICES J*

831   *Mar Sci* 2019;**76**:50–59. https://doi.org/10.1093/icesjms/fsy149

832   87.   Zheng M et al. Effects of human disturbance activities and environmental change factors

833   on terrestrial nitrogen fixation. *Glob Change Biol* 2020;**26**:6203–6217.

834   https://doi.org/10.1111/gcb.15328

835   88.   Dynarski KA, Houlton BZ. Nutrient limitation of terrestrial free-living nitrogen fixation. *New*

836   *Phytol* 2018;**217**:1050–1061. https://doi.org/10.1111/nph.14905

837   89.   Yan G et al. Long-time precipitation reduction and nitrogen deposition increase alter soil

838   nitrogen dynamic by influencing soil bacterial communities and functional groups.

839   *Pedosphere* 2020;**30**:363–377. https://doi.org/10.1016/S1002-0160(19)60834-9

840   90.   Bellenger JP et al. Biological nitrogen fixation by alternative nitrogenases in terrestrial

841   ecosystems: a review. *Biogeochemistry* 2020;**149**:53–73. https://doi.org/10.1007/s10533-

842   020-00666-7

843   91.   Chen M-Y et al. Phylogenomics Uncovers Evolutionary Trajectory of Nitrogen Fixation in

844   Cyanobacteria. *Mol Biol Evol* 2022;**39**:msac171. https://doi.org/10.1093/molbev/msac171

845   92.   Darnajoux R et al. Is vanadium a biometal for boreal cyanolichens? *New Phytol*

846   2014;**202**:765–771. https://doi.org/10.1111/nph.12777

847   93.   Sriswasdi S, Yang C, Iwasaki W. Generalist species drive microbial dispersion and

848   evolution. *Nat Commun* 2017;**8**:1162. https://doi.org/10.1038/s41467-017-01265-1

849   94.   Chen Y-J et al. Metabolic flexibility allows bacterial habitat generalists to become

850   dominant in a frequently disturbed ecosystem. *ISME J* 2021;**15**:2986–3004.

851   https://doi.org/10.1038/s41396-021-00988-w

852   95.   Carbonero F, Oakley BB, Purdy KJ. Metabolic Flexibility as a Major Predictor of Spatial

853   Distribution in Microbial Communities. *PLOS ONE* 2014;**9**:e85105.

854   https://doi.org/10.1371/journal.pone.0085105

96.   Hu W et al. Biological nitrogen fixation and the role of soil diazotroph niche breadth in

representative terrestrial ecosystems. *Soil Biol Biochem* 2024;**189**:109261.

https://doi.org/10.1016/j.soilbio.2023.109261

97.   Waite DW et al. Comparative Genomic Analysis of the Class Epsilonproteobacteria and

Proposed Reclassification to Epsilonbacteraeota (phyl. nov.). *Front Microbiol* 2017;**8**.

https://doi.org/10.3389/fmicb.2017.00682

98.   Kloesges T et al. Networks of Gene Sharing among 329 Proteobacterial Genomes

Reveal Differences in Lateral Gene Transfer Frequency at Different Phylogenetic Depths.

Mol Biol Evol 2011;**28**:1057–1074. https://doi.org/10.1093/molbev/msq297

99.   Moya-Beltrán A et al. Genomic evolution of the class Acidithiobacillia: deep-branching

Proteobacteria living in extreme acidic conditions. *ISME J* 2021;**15**:3221–3238.

https://doi.org/10.1038/s41396-021-00995-x

100. Campbell BJ et al. Adaptations to Submarine Hydrothermal Environments Exemplified by

the Genome of Nautilia profundicola. *PLOS Genet* 2009;**5**:e1000362.

https://doi.org/10.1371/journal.pgen.1000362

101. González-Rosales C et al. Integrative Genomics Sheds Light on Evolutionary Forces

Shaping the Acidithiobacillia Class Acidophilic Lifestyle. *Front Microbiol* 2022;**12**.

https://doi.org/10.3389/fmicb.2021.822229

102. McCutcheon JP, Moran NA. Extreme genome reduction in symbiotic bacteria. *Nat Rev

Microbiol* 2011;**10**:13–26. https://doi.org/10.1038/nrmicro2670

103. Cornejo-Castillo FM et al. Cyanobacterial symbionts diverged in the late Cretaceous

towards lineage-specific nitrogen fixation factories in single-celled phytoplankton. *Nat

Commun* 2016;**7**:11071. https://doi.org/10.1038/ncomms11071

104. Coale TH et al. Nitrogen-fixing organelle in a marine alga. *Science* 2024;**384**:217–222.

https://doi.org/10.1126/science.adk1075

105. Tripp HJ et al. Metabolic streamlining in an open-ocean nitrogen-fixing cyanobacterium.

Nature 2010;**464**:90–94. https://doi.org/10.1038/nature08786

106. Zehr JP et al. Unusual marine unicellular symbiosis with the nitrogen-fixing

cyanobacterium UCYN-A. *Nat Microbiol* 2016;**2**:16214.

https://doi.org/10.1038/nmicrobiol.2016.214

107. Kantor EJH et al. Metagenomics reveals the genetic diversity between sublineages of

UCYN-A and their algal host plastids. *ISME Commun* 2024;**4**:ycae150.

https://doi.org/10.1093/ismeco/ycae150

108. Wang S et al. Evolutionary Timeline and Genomic Plasticity Underlying the Lifestyle

Diversity in Rhizobiales. *mSystems* 2020;**5**:10.1128/msystems.00438-20.

https://doi.org/10.1128/msystems.00438-20

109. Zehr JP, Capone DG. Changing perspectives in marine nitrogen fixation. *Science* 2020.

https://doi.org/10.1126/science.aay9514

110. Xie J-B et al. Comparative genomic analysis of N2-fixing and non-N2-fixing Paenibacillus

spp.: organization, evolution and expression of the nitrogen fixation genes. *PLoS Genet*

2014;**10**:e1004231. https://doi.org/10.1371/journal.pgen.1004231

111. de Crécy-Lagard V et al. A roadmap for the functional annotation of protein families: a

community perspective. *Database J Biol Databases Curation* 2022;**2022**:baac062.

https://doi.org/10.1093/database/baac062

112. Anbar AD, Knoll AH. Proterozoic Ocean Chemistry and Evolution: A Bioinorganic Bridge?

Science 2002;**297**:1137–1142. https://doi.org/10.1126/science.1069651

113. Rucker H, Kacar B. Enigmatic evolution of microbial nitrogen fixation: insights from

Earth's past. *Trends Microbiol* 2023. https://doi.org/10.1016/j.tim.2023.03.011

114. Rico KI et al. Mysteries of metallome evolution: Integrating insights from the Earth and life

sciences. *Treatise Geochem* 2024.

115.   Kaçar B. Reconstructing Early Microbial Life. *Annu Rev Microbiol* 2024.

906   https://doi.org/10.1146/annurev-micro-041522-103400

907   116.   Falkowski PG. Life's Engines: How Microbes Made Earth Habitable. Princeton University

908   Press, 2015.