Ecological constraints and evolutionary trade-offs shape nitrogen fixation across habitats

Morgan S. Sobol[1,2*], Cécile Ané[3,4,] Katherine D. McMahon[5] and Betül Kaçar[1*]

1. Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA
2. Department of Biology, Texas State University, San Marcos, TX, USA
3. Department of Statistics, University of Wisconsin-Madison, Madison, WI, USA
4. Department of Botany, University of Wisconsin-Madison, Madison, WI, USA
5. Department of Civil and Environmental Engineering, University of Wisconsin–Madison, Madison, WI, USA

Correspondence: msobol@txstate.edu, bkacar@wisc.edu

Abstract

Life's transition into new habitats has profoundly shaped its trajectory through a reciprocal relationship with Earth's changing environments. To understand how ancient metabolisms co-evolved with these environments, it is essential to identify the ecological and evolutionary processes that structured their underlying genetic networks. Here, we focus on nitrogen (N_2) fixation, one of life's most critical and ancient metabolisms, to investigate the drivers of complexity in its associated genetic network. We used a large-scale comparative genomics framework to construct a comprehensive catalog of today's N_2 fixation-associated genes and assessed their distribution across diverse microbial genomes and environmental backgrounds. Genomes enriched in N_2 fixation genes generally have larger genome sizes, broader metabolic capabilities, wider habitat ranges, and are predominantly associated with mesophilic and aerobic lifestyles. Evolutionary reconstructions reveal a pattern of early gene gains in ancestral diazotrophs followed by lineage specific gene losses in later diverging taxa, suggesting evolutionary trade-offs shaped by changing environments. These findings demonstrate that the evolution of N_2 fixation has been intertwined with the restructuring of its genetic network, driven by feedbacks between microbial evolution and changing environmental and ecological conditions.

Introduction

Biological nitrogen fixation ("N_2 fixation") is an ancient metabolism tightly coupled to
Earth's geosphere [1–6]. Nitrogenases, catalytically responsible for reducing N_2 to NH_3,
are the only known enzymes capable of this reaction found in microorganisms, collectively
known as diazotrophs [7]. By supplying essential nitrogen for life, nitrogenase has been
transformative for biological productivity and Earth's environment [1, 2, 8–10].

While the global significance of diazotrophs is well established through studies of marine
and freshwater cyanobacteria [11] [12], and plant-associated symbioses [13], the factors
shaping N_2 fixation in both historical and modern contexts remain unclear [14].

The broad distribution of diazotrophs and their habitats reflects their evolution under
diverse ecological and physiological demands. The origin of nitrogenase remains
unsolved [15], but its core structural and functional features have remained largely
conserved over its long evolutionary history [4, 16]. In contrast, numerous genes involved
in its function and regulation have diversified [3, 17]. The composition of this genetic
"machinery" varies widely across the tree of life, from a small gene set in some genomes,
to more than 80 nitrogenase-associated genes in others [7, 18]. This variability suggests
that the environmental context and physiological requirements govern the nitrogenase-
related genes distribution among taxa [3, 19].

An integrated synthesis that brings together the diversity of N_2 fixation and its associated
genes within a broader ecological and evolutionary framework is only beginning to take
shape. This integration is essential to connect how the molecular evolution of nitrogenase
has been shaped by the interplay of genes within its broader interaction network.

Here, we employed large-scale comparative genomics to investigate how organismal
lifestyles influence evolutionary trajectories, particularly in in relation to the varying N_2
fixation-associated gene content. Our findings demonstrate that the evolutionary
diversification of the N_2 fixation machinery has facilitated the expansion of diazotrophs
into increasingly broader ecological niches over time.

Figure 1. Structure of the major and minor molybdenum nitrogenase (Nif) clusters, vanadium nitrogenase (Vnf) and iron nitrogenase (Anf) clusters, and the molybdenum acquisition (Mod) cluster in model diazotroph *Azotobacter vinelandii* DJ. Genome positions and gene sizes for each cluster are based on KEGG genome reference T00885 for *A. vinelandii* DJ.

Methods

Genome selection

The reference phylogeny Web of Life (WoL) [20] was selected as the basis of our study because of its extensive genome coverage and phylogenetic reconstruction. For our analysis, we kept WoL genomes with ≥90% completeness and <5% contamination, following quality thresholds established by [21].

Protein fasta files associated with the genome accessions (named "*_protein.faa.gz") were downloaded from NCBI with eutils [22] (accessed July 17th, 2023). Genomes missing available NCBI protein fasta files were excluded from analysis. This reduced the WoL genome dataset from 10,575 genomes to 7,458. Nucleotide sequences of noncoding RNA genes were downloaded if they were available at GenBank or RefSeq and the 16S rRNA genes were extracted.

Curation of nitrogen fixation gene catalog

Protein fasta files were functionally annotated using the Kyoto Encyclopedia of Genes and Genomes (KEGG) database [23]. Proteomes were annotated with KEGG orthology (KO) identifiers by profiling each genome against KEGG hidden Markov models (KEGG release 106) for all KOs [24] (cutoff E-value = 1e-5) using *hmmscan* in HMMER version 3.2.2 (http://hmmer.org/). Custom Python scripts were used to parse through each genome's *hmmscan* output and generate the final KO presence-absence matrix. The presence/absence matrix was subset to genes known to be involved in N_2 fixation (and which had KO identifiers), which were collected through a literature search (**Supplemental Table 1**).

Trait data curation

Optimal growth temperature (OGT), oxygen, pH preference, and habitat preference trait data were collected for each genome by matching the Genbank accession number to BacDive [25], NCBI BioProject [26], or GOLD JGI [27] as well as matching the NCBI taxon IDs to the Madin et al (2020) phenotypic trait database [28]. Trait data was also predicted for all genomes. OGTs were predicted using the OGT_prediction tool [29]. The regression model used included both Bacteria and Archaea, but excluded using information of the 16S rRNA gene and genome size (*superkingdom-all species-all features ex genome size and rRNA.txt*). The *Oxyphen* package was used for oxygen metabolism prediction based on the number of oxygen utilizing genes in each genome [30]. Predictions for pH were done with the Ramoneda et al. (2023) prediction model which is based on genes that influence pH tolerance [31]. Habitat preference and coverage were determined with ProkAtlas [32] by mapping the 16S rRNA gene sequences to the ProkAtlas metagenome database. Barrnap version 0.9 [33] was used to extract 16S rRNA sequences for genomes with missing annotations. Using a custom python script, the longest 16S rRNA sequence was extracted for each genome as its representative 16S rRNA gene sequence to correct for errors in sequence length due to sequencing and assembly bias. Twenty-one genomes without a 16S rRNA gene sequence were excluded. A nucleotide identity of 97% and a query length match of at least 150 base pairs were used as thresholds. Habitat coverage was determined by

summing up the total number of habitats where a genome was considered present and
dividing it by the total number of habitats investigated by ProkAtlas. The presence of
KEGG modules was predicted using the KEGG KO annotations with MetaPathPredict
(version 0) [34]. Metabolic ~~breadth~~ coverage was calculated as the total number of KEGG
modules present in a genome divided by the total number of modules searched for with
MetaPathPredict. Data collected from databases and predicted data were cross-
referenced and manually annotated to provide the final trait data for each genome.

Data analysis and statistics

We compared the mean number of N_2 fixation genes across specific traits (oxygen,
temperature, and habitat preferences) with non-parametric methods using the
stat_compare_means function in the ggpubr R package (version 0.6.0) [35]. For multi-
group traits habit and oxygen preference, we first applied a Kruskal-Wallis test, followed
by pairwise Wilcoxon rank-sum tests with Benjamini-Hochberg correction. For
temperature preference (categories: mesophilic and thermophilic) we used the Wilcoxon
rank-sum test directly. Compact letter display was used to report pairwise groups whose
means are discernibly different, based on a p-value of 0.05. The *distRoot* function in the
adephylo R package [36] was used to calculate the root-to-tip patristic distances by
measuring the total length of branches (in terms of the number of substitutions per site)
from the root of the species tree to each tip. Non-phylogenetic linear regressions were
performed using the *lm* function in the stats R package [37].

The gene presence-absence matrix and trait data were mapped onto the species
phylogeny using the ggTree R package (version 3.6.2) [38]. This matrix was also used in
phylogenetic principal-component analysis (PCA) performed with the *phyl.pca* function
(method = "BM", mode ="corr") in phytools R package [39], to visualize similarities and
dissimilarities between genomes. The first two components were retained for
visualization.

Evolutionary trends analysis

Phylogenetic regression models were fit using the *phylolm* function in the phylolm R
package (version 2.62) [40], using Pagel's lambda model of phylogenetic correlation. The

lambda parameter (λ) ranges from 0 (no phylogenetic signal) to 1 (high phylogenetic
signal) in the response trait when the model has no covariates, or in the residuals when
covariates are included. Gene gain/loss was determined with Count [41] using a non-
binary count matrix [42] for all accessory (non-structural) Nif genes. Wagner parsimony
with a gain/loss penalty of 2 was applied to infer the most parsimonious scenario of genes
present at the internal nodes, as done previously [43]. Ancestral trait reconstructions were
performed on an ultrametric and dichotomous tree, which was made using the *multi2di*
and *chronos* (model = "relaxed") functions of the APE package [44] in R. The *fitdiscrete*
function in geiger [45] was used for discrete reconstructions of oxygen preference, which
were mapped onto the phylogeny with the *simmap* function in phytools [39]. The *fastAnc*
function in phytools was used for continuous variable reconstructions of metabolic and
habitat diversity, which was mapped onto the tree using the *contMap* function in phytools.

Limitations of comparative genomics

As with any comparative genomic approach, there are inherent limitations when working
to identify the genes and traits of genomes. One limitation of this study is the selection of
genes that have been identified as playing a role in N_2 fixation, which means there may
be numerous other genes involved in the process that remain undiscovered due to the
lack of comprehensive functional characterization of many diazotrophs. Additionally, the
use of KEGG KO numbers presents another caveat, as not all genes could be assigned
KO numbers, and some KO numbers cannot adequately distinguish between genes with
high sequence similarities, e.g. *nifA*/*vnfA*/*anfA*. Future improvements in functional
annotation of genes and database curation is required to obtain a complete view of the
capabilities of diazotrophs [46].

Data availability

162 Supplementary Table 2 contains genome accession, taxonomy, and trait data
163 information. Custom scripts, raw data, and PDF versions of the figures are available at
164 https://github.com/kacarlab/Sobol_2025_NifGeneEcoEvo.

Results

Distribution of nitrogenases across the tree of life

We used a large prokaryote reference phylogeny, "WoL" [20], containing over 7,000 high-quality genomes [21] to first assess the representative diversity and distribution of nitrogenase today. A total of 673 genomes (~6.4%) were found to contain nitrogenase (**Supplemental Table 2**) based on the presence of the catalytic HDK subunits of nitrogenase (**Figure 2A**). Our estimate of nitrogenase diversity falls within range of recent estimates (~5.9% to 16%) [19, 47–49]. Here, approximately 46.9% of these genomes belonged to Proteobacteria (in order of abundance: Alpha>Delta>Gamma>Beta>Epsilon>Acidithiobacillia), while the remaining top three phyla included Firmicutes (17.3%), Cyanobacteria (12.6%), and Euryarchaeota (11.4%). Amongst the nitrogenase-containing genomes, molybdenum (Mo)-only nitrogenase made up the majority (90.3%), followed by genomes with both Mo-nitrogenase and iron (Fe) nitrogenase (5.9%), genomes with both Mo-nitrogenase and vanadium (V) nitrogenase (2.2%), and lastly with 1.5% of genomes containing all three nitrogenases (Figure 1) (**Figure 2B**). Genomes with V-nitrogenase were found in 6 phyla: Euryarchaeota, Firmicutes, Alphaproteobacteria, Deltaproteobacteria, Cyanobacteria, and Gammaproteobacteria. Fe-nitrogenases were found in 11 phyla: Firmicutes, Epsilonproteobacteria, Alphaproteobacteria, Gammaproteobacteria, Verrucomicrobia, Euryarchaeota, Deltaproteobacteria, Gammaproteobacteria, Bacteroidetes, Spirochaetes, and Chlorobi. Euryarchaeota represented 50% of the genomes with all three nitrogenases; the remaining phyla included Alphaproteobacteria, Firmicutes, and Gammaproteobacteria.

Figure 2. Diversity and phylogenetic distribution of nitrogenase. **A)** The overall distribution of nitrogenase across the microbial diversity from the Zhu et al (2020) Web of Life (WoL) phylogeny. Black bars on the outer ring represent the presence of nitrogenase. Numbers reflect the top 14 taxonomic groups in the dataset with the highest proportion of genomes with nitrogenase. **B)** Proportions and taxonomic distributions of the different nitrogenase isozymes across WoL.

An Ecological and Genomic Survey of N₂ Fixation Genes

We screened all 673 genomes for 93 known genes ("machinery") involved in N_2 fixation (**Supplemental Table 3, Supplemental Figure 1**). In Figure 3, genes were organized into categories and the relative abundance of genes within each category was calculated. At the individual genome level, without considering *nifHDK*, the total number of N_2 fixation genes within a genome ranged from 14 genes to 67 genes. The highest number of genes were found in the Alphaproteobacteria *Rhodopseudomonas palustris CGA009,* followed closely by *Azotobacter vinelandii* DJ with 65 genes. Archaea *Methanococcus aeolicus* Nankai-3, *Methanothermococcus okinawensis* IH1 and *Methanocaldococcus infernus* ME all tied for the lowest number of genes (14). Throughout the entire species phylogeny, we observed a general trend of increasing number of genes scaled with increasing taxa branch lengths (**Supplemental Figure 2**).

Figure 3. Distribution of N₂ fixation machinery mapped onto the phylogeny of nitrogenase-containing genomes, highlighting associations with species-associated ecological and physiological traits. This pattern

208
209

also correlates with those that are mesophilic, prefer aerobic and/or facultative lifestyles, possess larger
genome sizes, broader metabolic capabilities, and occupy a wider range of habitats. For this figure, N_2
fixation genes were organized into functional categories and the relative frequency of genes for each
category calculated. The full gene matrix visualization can be found in Supplemental Figure 1 and the gene
matrix and associated categories in Supplemental Table 3.

For each genome, we gathered trait data for optimal growth temperature and genome
size, as well as preferences for oxygen, pH, and habitats (**Figure 3**). In addition, we
calculated the frequency of metabolic pathways encoded in each genome based on the
total number of pathways considered (i.e., metabolic coverage) and the proportion of
habitats a genome was assigned based on the total number of habitats surveyed (i.e.,
habitat coverage). To improve trait assignments, we combined data from multiple
databases and predicted traits based on genomic features (see methods).

Amongst the nitrogenase-containing genomes, 47.5% of the genomes were anaerobes,
31.6% were aerobes, 5.3% were facultative, and 1.5% were microaerophiles. The
remaining 14% could not be determined due to missing data or uncertainty in predictions.
Temperature preferences further define nitrogenase distribution, with most genomes
(91.2%) belonging to mesophiles that favor temperatures from 18ºC to 45ºC. The
remaining ~8.8% belonged to thermophiles that preferred temperatures above 45ºC.
Similarly, neutrophiles (pH 5.5-8.0) dominated the pH preferences (87.2%). Acidophiles
and alkaliphiles, though present, were not as common. Habitat preferences varied
amongst genomes. Here, freshwater and terrestrial environments were particularly
favored by 32.5% and 26.2% of genomes, respectively. The remaining genomes were
associated with saline environments (14.3%), non-plant host associations (11.4%),
associations with plants (8.8%), and hydrothermal systems (6.8%). In addition to these
traits, we assessed differences in metabolic and habitat coverage to highlight the
ecological versatility of diazotrophs. Metabolic coverage ranged from 9.8 to 46.4%, with
an average of 23.2%. Habitat coverage ranged from 0 to 48%, with an average of 12%,
when not accounting for the ~3% of genomes whose habitat coverage could not be
determined. We also assessed variations in genome size. Here, genome size ranged
from 1.3 to 12.3 Mbps, with the average being 4.6 Mbps.

Figure 4. N₂ fixation gene richness varies across traits. The number of genes was compared across **A)** temperature**, B)** oxygen, and **C)** habitat preferences**.** Pairwise differences between means were calculated using non-parametric Wilcoxon rank-sum tests. For each panel, the means of those that do not differ with statistical evidence share the same letter, whereas those that do not share a letter are discernibly different ($p < 0.05$).

Physiological, metabolic, and habitat preferences vary with N₂ fixation genetic complexity

Next, we looked at relationships between the number of N₂ fixation genes in a genome and their environmental and physiological traits to understand factors that have altered

the complexity of machinery needed for N_2 fixation. We observed that genomes with
similar preferences for temperature, oxygen, and habitat type had comparable numbers
of N_2 fixation genes (**Figure 3**). Specifically, thermophiles and anaerobes have discernibly
fewer genes compared to their mesophilic (p.adj=8.60×10^{-23}) and aerobic (p.adj=4.80×10^{-45})
counterparts (**Fig. 4A,B**). Accordingly, organisms who prefer hydrothermal
environments have discernibly fewer genes on average compared to all other preferred
environments (**Fig. 4C**). There were no discernible differences in gene richness between
terrestrial and freshwater, saline, or host-associated environments. Plant-associated taxa
had the highest average of N_2 fixation genes, which was discernibly higher than all
environments, except host-associated.

N_2 *fixation genetic complexity as a predictor of co-evolutionary trait associations*

To test the extent to which traits are evolutionarily conserved or have evolved
independently, we measured the phylogenetic signal for gene richness, habitat coverage,
and metabolic coverage, as well as genome size, temperature, pH, and oxygen
preference across the species phylogeny using Pagel's lambda (λ) with the *phylolm* R
package [40]. All traits had a moderate to high ($\lambda \geq 0.69$) phylogenetic signal
(**Supplemental Table 4**), meaning that closely related genomes demonstrated greater
similarity in trait values compared to genomes randomly selected from the phylogeny.

Next, we assessed whether certain traits shared an evolutionary history with gene
richness via phylogenetic linear regression models (**Figure 5, Supplemental Table 5**).
Gene richness was chosen as the independent variable to determine if the complexity of
N_2 fixation machinery in a genome could be used to predict other trait values. Response
(trait) variables were transformed using either natural log or square root to stabilize
variance prior to analysis. Genome size (**Figure 5A**) and metabolic coverage (**Figure 5B**)
had a positive, significant linear relationship with N_2 fixation gene richness. As for
temperature, a significant, negative linear relationship with gene richness was measured
(**Figure 5C**). Outliers (n=15) were identified following log transformation of the
temperature variable, but we confirmed that removing these outliers prior to the
regression did not influence the strength of statistical evidence, nor the direction of the
relationship. Habitat richness was marginally significantly dependent (p=0.054) on gene

number, and had a positive slope, indicating a general relationship between the two traits

(**Figure 5D**).

Figure 5. Phylogenetic linear regressions assessed whether the number of N_2 fixation genes in a genome
could predict other traits, such as **A)** genome size, **B)** metabolic coverage, **C)** optimal growth temperature,
and **D)** habitat coverage. Response variables were either transformed using natural log or square root to
stabilize variance prior analysis. Solid blue lines represent non-phylogenetic linear model coefficients,
dashed blue lines represent phylogenetic linear coefficients.

We assessed whether phylogenetic signal at the individual phylum level remained in order
to test for phylum-specific evolutionary histories and differing selection pressures. This
analysis helped to determine whether the overall phylogenetic signal measured across
the entire species phylogeny was driven by specific phyla more so than others. Due to
some phyla having too few genomes (<20), phyla-specific analysis was only carried out
on genomes assigned to Alpha-, Beta-, Delta-, Gammaproteobacteria, Cyanobacteria,
Firmicutes, and Euryarchaeota. Genome size, habitat coverage, and metabolic coverage
maintained strong phylogenetic signal across these major nitrogenase-containing phyla
($\lambda \geq 0.8$) (**Supplemental Table 6**). Gene richness had a high λ (>0.8) within most phyla,
except Cyanobacteria, where λ was 0.34. No phylogenetic signal for temperature was
observed in Gammaproteobacteria ($\lambda=0.001$) as they are all are mesophiles, whereas the
other phyla had high signal ($\lambda>0.8$). Oxygen did not have a strong signal in
Gammaproteobacteria nor Firmicutes. As for pH, the strength of phylogenetic
conservation was high for Cyanobacteria, Euryarcheaota, and Gamma- and
Betaproteobacteria, but was low for Deltaproteobacteria.

Differences in N_2 fixation machinery distribution

Phylogenetic principal component analysis (PCA) was used to cluster genomes together
based on similarity of N_2 fixation gene presence-absence patterns, while accounting for
phylogenetic relationships. Indeed, we found that phylogenetic relationships accounted
for most of the variation, indicated by the low explained variance in the phylogenetic PCA
(**Supplemental Figure 3**).

The top ten most influential genes from each component of the phylogeny-corrected PCA
(**Supplemental Figure 3**) included the different regulation elements (*nifI₂*, *nifL*, *regRS*,
ntrB), two different electron-donating/transfer mechanisms (*rnfABCDEG, fixABCX*),
nodulation and heterocyst differentiation genes (*nodABCD*, *devC*), oxidative protection
mechanisms (*sodN*, *cooA*), cofactor synthesis and iron acquisition genes (*nifM*, *sufD*,
feoC2)*, and the genes that make up both the Fe- and V-nitrogenase (*vnfHGDKE*,
anfHGDK). The post-translational *nifI₂* regulator is primarily found in anaerobes, such as
Archaea and Firmicutes, whereas *regS*, *nifL*, and *ntrB* are primarily found in aerobic
Proteobacteria (**Supplemental Figure 1**). The preference for nitrogenase regulation
mechanisms therefore drives part of the separation of aerobes and anaerobes in the PCA
(**Supplemental Figure 3**). Pearson correlation of gene co-occurrence (**Supplemental**
Figure 4) also showed that *nifI₂* and *nifI₁* had strong negative correlations with genes
nifTQWXZ, *fnr*, *fdx*, *oxyR,* and *ntrBC*, which are genes primarily found in aerobes.

A preference for either *rnfACDEG* or *fixACX* electron donation systems was evident in
the PCA where the genes pointed opposite from one another (**Supplemental Figure 3**).
This anti-correlation was also seen in **Supplemental Figure 2**. Nodulation genes
nodABC also pointed opposite to *rnfACDEG* (**Supplemental Figure 3**) and instead had
a positive correlation with *fixACX* (**Supplemental Figure 4**). This indicates that
diazotrophs forming symbioses with plants through nodulation may prefer Fix over Rnf
for electron donation. None of the N₂ fixation genes were unique to thermophiles and
neither Fe- nor V- nitrogenases were found in genomes with optimal growth temperatures
greater than 41ºC.

Dynamic history of gene gain and loss

Although previous studies have noted the potential dynamic nature of N₂ fixation gene
acquisitions and losses [19, 50], none have specifically analyzed these patterns in detail,
nor in relation to their traits. By observing gene gain/loss patterns across the phylogeny
and linking them to shifts in lifestyle changes, we can begin to observe the selective
pressures on the retention or loss of particular genes. We estimated net gain or loss at
each node of the nitrogenase-containing species tree using Wagner parsimony (see
methods) (**Figure 6**). Evolutionary trajectories of N₂ fixation machinery complexity indeed

indicated a dynamic history of gene gain and loss over time. Specifically, early branching

nodes imply net gene gain, suggesting an initial expansion of genes early in diazotroph

evolution, whereas gene loss primarily occurs closer to the terminal tree branches.

The nodes with the highest number of gene gains (+11) occurs at the most recent

common ancestor (MRCA) of Bacteria and at the MRCA of the Euryarchaeota clade

Methanosarcina (not shown). Some of the clades that experienced large gene losses

include Acidithiobacillia (-18), the Firmicute-Clostridia clade Thermoanaerobacterales (-

18) (not shown), Thermodesulfobacteria (-17) (not shown), Nitrospirae (-16), and the

MRCA of Aquificae and Epsilonproteobacteria (-11) (**Supplemental Tables 7-8**).

Acidothiobacillia's loss of genes *cowN*, *fixJL*, and the Suf and Rnf systems occurs in

tandem with a loss of habitat coverage (**Figure 6**). The loss of post-transcriptional

regulators *nifI$_1$I$_2$* unsurprisingly coincides with a gain in aerobic lifestyles at the MRCA of

Cyanobacteria, Actinobacteria and the MRCA of Gamma- and Alphaproteobacteria

(**Figure 6**). The appearance of transcriptional regulation by *nifA* predates the loss of *nifI$_1$I$_2$*

in the MRCA of Gamma- and Alphaproteobacteria (**Figure 6**). The gain of genes involved

in protection against oxidants (*katG*, *oxyR*, and *cowN*) and oxygen response regulators

fixJL occurs early along the MRCA nodes of Verrucomicrobia, Chlorobi, Bacteroidetes,

and Proteobacteria (**Figure 6**). We noted two distinct gains of *rnfACDEG*, *nifTWZ*, *katG*

and *fixCX* in distantly related diazotrophs, indicating they were either inherited from a

common ancestor that does not have nitrogenase (and therefore would not be

represented in this tree), and/or they were gained through horizontal gene transfer.

Figure 6. N_2 fixation gene gain (green triangles) and gene loss (red triangles) throughout the evolution of diazotrophs. No change (grey circle) represents no gain or loss at that node, or an equal number or gains and losses. Asterisks denote candidate genes for HGT, or genes potentially derived from non-diazotrophic ancestors not represented in this subsetted phylogeny. All gene events can be found in Supplemental table 8. Key transitional events of gene richness and habitat coverage can be found in Supplemental Figure 5.

Discussion

Nitrogenase Distribution, Diversity, and Evolution Across Prokaryotic Lineages

Nitrogenase is widespread across prokaryotes, appearing within phylogenetically and ecologically diverse Bacteria and Archaea (**Figures 2 and 3**). Mo-nitrogenase is the most prevalent form, reported everywhere from the rhizosphere [51] to deep-sea sediments kilometers below the ocean surface [52–54]. Although V- and Fe-nitrogenases occur less frequently, they also span across diverse taxa (**Figure 2**; [55]). However, focusing solely on nitrogenase's distribution overlooks the evolutionary and genetic complexity of N_2 fixation.

To address this, we developed a comprehensive catalog of nitrogenase-containing genomes and associated N_2 fixation machinery, mapped to environmental preferences and phylogenetic history (**Figure 3**). Expanding on prior work [19], our work incorporates ~2.5× more N_2 fixation genes and genomes, along with data on oxygen metabolism, temperature, pH, genome size, habitat range. Whereas previous comparative studies often targeted single clades or ecosystems [56, 57], our framework unifies nitrogenase and its supporting genes across the entire microbial tree.

Oxygen is a well-known inhibitor of nitrogenase [58], supporting its origin under anoxic conditions and later diversification following planetary oxygenation [5, 17]. Consistent with this, aerobic diazotrophs harbor significantly more N_2 fixation genes than anaerobes (**Figure 4B**), likely due to the need for oxygen protection and regulation under aerobic growth (**Figure 3**).

Temperature also shapes the complexity of N_2 fixation gene repertoires. Mesophiles harbor more N_2 fixation genes than thermophiles (**Figure 4A**) and hydrothermal organisms exhibit the lowest average counts (**Figure 4B**). These findings align with previous research showing temperature-driven influences on molecular evolution [59–61]. Cold-adapted diazotrophs often switch from Mo- to V-based nitrogenases [62–64]. In further support of this view, we detected a temperature-dependent trend in gene richness across evolutionary histories of nitrogenase-encoding organisms (**Figure 5C**).

Notably, several anaerobic thermophiles lack *nifE, nifN*, or both (**Figure 3**), genes
involved in FeMo cofactor assembly and part of the typical minimal nif genetic set
(*nifHDKENB)* required for active N_2 fixation [7]. These lineages include Euryarcheaota,
such as members of the ANME groups [65], Chloroflexi, and Firmicutes. Some have
verified N_2 fixation activity [66–69], while others, like *Roseiflexus castenholzii*, do not [70].
Interestingly, recent work suggests that *nifEN* itself may have originated from a duplicated
ancestral nitrogenase [15], raising the possibility that some extant diazotrophs lacking
nifEN represent primitive traits of early nitrogenase evolution or that they developed
genetic strategies to maximize energy conservation [71].

Plant-associated genomes showed the highest N_2 fixation gene counts (**Figure 4C**),
consistent with their pivotal role in host nitrogen supply [72, 73]. In symbiosis, plants
regulate oxygen to initiate the FixL and FixJ pathway, ultimately driving the expression of
nitrogenase [13], while *nod* genes mediate nodulation [74]. Terrestrial habitats, being
more variable than oceans [75], impose fluctuating environmental conditions that
necessitate dynamic nitrogenase regulation [76–78]. Molybdenum scarcity in soils also
favors alternative nitrogenases [79], likely contributing to gene expansion in terrestrial
environments.

Reconstructing the Eco-Evolutionary History of Nitrogenase-Associated Genes

Central to our framework is that N_2 fixation gene complexity is not an isolated trait, rather,
it fits into a larger evolutionary framework that includes genome expansion, metabolic
versatility, and habitat diversity (**Figure 5**). As such, gene complexity may serve as an
ecological marker—distinguishing generalists, which inhabit diverse environments and
possess large, versatile genomes [80–82], from specialist, which typically occupy
narrower niches and have smaller genomes [83, 84]. Distinguishing the ecological
strategies of diazotrophs may be especially important today, as specialist diazotrophs
may contribute disproportionately to total soil N_2 fixation [85].

We find that as diazotrophs diversified and specialized over time, so did their N_2 fixation
machinery (**Figure 3**). Early evolution of diazotrophs was dominated by gene gain; later
stages were marked by loss (**Figure 6**), particularly in terminal branches, pointing to

lineage specific streamlining or ecological specialization. We observed this pattern most
notably in Chloroflexi, Nitrospirae, Aquificae, Acidithiobacillia, Thermodesulfobacterota,
and Epsilonproteobacteria, groups often characterized with either anaerobic,
thermophilic, acidophilic, or host-adapted lifestyles (**Supplemental Table 2**). These traits
suggest niche-specialization from once generalist ancestors.

In Epsilonproteobacteria and Acidithiobacillia, gene loss correlates with reduced habitat
breath, consistent with previous findings [86–88]. Whether via adaptation to deep-sea
hydrothermal vents [89], shifting toward heterotrophy [87], or acidophilic transitions from
a neutrophile ancestor [90], these patterns indicate broader evolutionary trends. In
obligate symbioses, gene loss often reflects host compensation [91]. A well-known case
is UCYN-A, a marine symbiotic cyanobacterium with a reduced genome [92, 93], which
was recently identified as an early stage N_2-fixing organelle ("Nitroplast") in
Braarudosphaera bigelowii [94, 95]. This and other examples underscore the central
importance of gene loss in shaping diazotroph evolution.

Examining the distribution of diverse diazotroph genetic machinery offers insight into
which genes were critical for enabling transitions into new environments over evolutionary
time. The rise of atmospheric oxygen, for example, fundamentally reshaped electron-
donor availability [96] and, along with changes in in metal availability [14, 97], likely
exerted strong selective pressures on N_2 fixation machinery. Certain genes, such as the
G-subunit of alternative nitrogenases [17], appear to have emerged during this transition
to an oxic atmosphere, underscoring how environmental transitions, particularly the Great
Oxidation Event, shaped molecular evolution in deep time.

Reconstructing historical patterns from the modern genetic records is thus pivotal for
understanding the processes that govern evolutionary change [98]. Global trends in
genome evolution not only reveal how organisms responded to past environmental
fluctuations but connects them to their present-day ecological niches. In particular, N_2
fixation complexity may help infer a diazotroph's ecological role, highlighting how
environmental factors and cellular demands intersect to enable colonization of new
habitats.

In his book *Life's Engines*, Paul Falkowski likens nitrogenase and its core machinery to a computer's hardware, while accessory genes function as the software, essential for maintenance, optimization and updates [99]. This analogy is compelling, capturing the deep interconnection between core biochemical processes and their regulatory networks. These elements must evolve in concert to navigate the challenges posed by changing environments, a dynamic reflected in the expansions and contractions of diazotrophs which we have documented across both ancient and modern contexts.

Acknowledgements

We gratefully acknowledge support from the NASA Interdisciplinary Consortium for Astrobiology Research (ICAR): Metal Utilization and Selection Across Eons, MUSE under grant no 80NSSC17K0296, and the NASA Exobiology Program under grant no NNH23ZDA001N and a UW-Madison Research Forward Award to BK. M.S. was supported by the NASA Postdoctoral Program, administered by Oak Ridge Associated Universities under contract with NASA. We thank Patricia Tran, Shaomei He, colleagues in the Kaçar Lab, members of NASA MUSE ICAR, and colleagues at Texas State University for helpful feedback on manuscript; and the Center for High Throughput Computing (CHTC) and the Department of Bacteriology at the University of Wisconsin–Madison for providing computing resources.

References

1. Falkowski PG. Evolution of the nitrogen cycle and its influence on the biological sequestration of CO2 in the ocean. *Nature* 1997; **387**: 272–275.

2. Canfield DE, Glazer AN, Falkowski PG. The evolution and future of Earth's nitrogen cycle. *Science* 2010; **330**: 192–196.

3. Mus F, Colman DR, Peters JW, Boyd ES. Geobiological feedbacks, oxygen, and the evolution of nitrogenase. *Free Radic Biol Med* 2019; **140**: 250–259.

4. Garcia AK, McShea H, Kolaczkowski B, Kaçar B. Reconstructing the evolutionary history of nitrogenases: Evidence for ancestral molybdenum-cofactor utilization.

Geobiology 2020; **18**: 394–411.

5.   Parsons C, Stüeken EE, Rosen CJ, Mateos K, Anderson RE. Radiation of nitrogen-

metabolizing enzymes across the tree of life tracks environmental transitions in

Earth history. *Geobiology* 2021; **19**: 18–34.

6.   Lyons TW, Tino CJ, Fournier GP, Anderson RE, Leavitt WD, Konhauser KO, et al.

Co-evolution of early Earth environments and microbial life. *Nat Rev Microbiol* 2024;

22: 572–586.

7.   Dos Santos PC, Fang Z, Mason SW, Setubal JC, Dixon R. Distribution of nitrogen

fixation and nitrogenase-like sequences amongst microbial genomes. *BMC*

Genomics 2012; **13**: 162.

8.   Navarro-González R, McKay CP, Mvondo DN. A possible nitrogen crisis for

Archaean life due to reduced nitrogen fixation by lightning. *Nature* 2001; **412**: 61–

64.

9.   Raymond J, Siefert JL, Staples CR, Blankenship RE. The natural history of nitrogen

fixation. *Mol Biol Evol* 2004; **21**: 541–554.

10.   Allen JF, Thake B, Martin WF. Nitrogenase inhibition limited oxygenation of earth's

Proterozoic atmosphere. *Trends Plant Sci* 2019; **24**: 1022–1031.

11.   Zehr JP, Capone DG. Changing perspectives in marine nitrogen fixation. *Science*

2020; **368**: eaay9514.

12.   Marcarelli AM, Fulweiler RW, Scott JT. Nitrogen fixation: a poorly understood

process along the freshwater-marine continuum. *Limnol Oceanogr Lett* 2022; **7**: 1–

10.

13.   Mus F, Crook MB, Garcia K, Garcia Costas A, Geddes BA, Kouri ED, et al.

Symbiotic nitrogen fixation and the challenges to its extension to nonlegumes. *Appl*

Environ Microbiol 2016; **82**: 3698–3710.

14. Rucker HR, Kaçar B. Enigmatic evolution of microbial nitrogen fixation: insights

from Earth's past. *Trends Microbiol* 2024; **32**: 554–564.

15. Garcia AK, Kolaczkowski B, Kaçar B. Reconstruction of nitrogenase predecessors

suggests origin from maturase-like proteins. *Genome Biol Evol* 2022; **14**.

16. Harris DF, Rucker HR, Garcia AK, Yang Z-Y, Chang SD, Feinsilber H, et al. Ancient

nitrogenases are ATP dependent. *MBio* 2024; **15**: e0127124.

17. Cuevas-Zuviría B, Garcia AK, Rivier AJ, Rucker HR, Carruthers BM, Kaçar B.

Emergence of an orphan nitrogenase protein following atmospheric oxygenation.

Mol Biol Evol 2024; **41**.

18. Rubio LM, Ludden PW. Biosynthesis of the iron-molybdenum cofactor of

nitrogenase. *Annu Rev Microbiol* 2008; **62**: 93–111.

19. Boyd ES, Costas AMG, Hamilton TL, Mus F, Peters JW. Evolution of molybdenum

nitrogenase during the transition from anaerobic to aerobic metabolism. *J Bacteriol*

2015; **197**: 1690–1699.

20. Zhu Q, Mai U, Pfeiffer W, Janssen S, Asnicar F, Sanders JG, et al. Phylogenomics

of 10,575 genomes reveals evolutionary proximity between domains Bacteria and

Archaea. *Nat Commun* 2019; **10**: 5477.

21. Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy

TBK, et al. Minimum information about a single amplified genome (MISAG) and a

metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol*

2017; **35**: 725–731.

22. Nadkarni PM, Parikh CR. An eUtils toolset and its use for creating a pipeline to link

genomics and proteomics analyses to domain-specific biomedical literature. *J Clin*

Bioinforma 2012; **2**: 9.

23. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference

resource for gene and protein annotation. *Nucleic Acids Res* 2016; **44**: D457-62.

24. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al.

KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive

score threshold. *Bioinformatics* 2020; **36**: 2251–2252.

25. Reimer LC, Sardà Carbasse J, Koblitz J, Ebeling C, Podstawka A, Overmann J.

BacDive in 2022: the knowledge base for standardized bacterial and archaeal data.

Nucleic Acids Res 2022; **50**: D741–D746.

26. Barrett T, Clark K, Gevorgyan R, Gorelenkov V, Gribov E, Karsch-Mizrachi I, et al.

BioProject and BioSample databases at NCBI: facilitating capture and organization

of metadata. *Nucleic Acids Res* 2012; **40**: D57-63.

27. Mukherjee S, Stamatis D, Bertsch J, Ovchinnikova G, Sundaramurthi JC, Lee J, et

al. Genomes OnLine Database (GOLD) v.8: overview and updates. *Nucleic Acids*

Res 2021; **49**: D723–D733.

28. Madin JS, Nielsen DA, Brbic M, Corkrey R, Danko D, Edwards K, et al. A synthesis

of bacterial and archaeal phenotypic trait data. *Sci Data* 2020; **7**: 170.

29. Sauer DB, Wang D-N. Predicting the optimal growth temperatures of prokaryotes

using only genome derived features. *Bioinformatics* 2019; **35**: 3224–3231.

30. Jabłońska J, Tawfik DS. The number and type of oxygen-utilizing enzymes

indicates aerobic vs. anaerobic phenotype. *Free Radic Biol Med* 2019; **140**: 84–92.

31. Ramoneda J, Stallard-Olivera E, Hoffert M, Winfrey CC, Stadler M, Niño-García JP,

et al. Building a genome-based understanding of bacterial pH preferences. *bioRxiv* .

2023. , 2023.01.24.524446

32. Mise K, Iwasaki W. Environmental atlas of prokaryotes enables powerful and

intuitive habitat-based analysis of community structures. *iScience* 2020; **23**:

101624.

33. Seemann T. barrnap: Bacterial ribosomal RNA predictor.

https://github.com/tseemann/barrnap. Accessed 2 Dec 2024.

34. Geller-McGrath D, Konwar KM, Edgcomb VP, Pachiadaki M, Roddy JW, Wheeler

TJ, et al. Predicting metabolic modules in incomplete bacterial genomes with

MetaPathPredict. *Elife* 2024; **13**.

35. Kassambara A. ggpubr:'ggplot2'based publication ready plots. *R package version*

2018; 2.

36. Jombart T, Balloux F, Dray S. Adephylo: New tools for investigating the

phylogenetic signal in biological traits. *Bioinformatics* 2010; **26**: 1907–1909.

37. R Core Team. R: A Language and Environment for Statistical Computing. Vienna,

Austria: R Foundation for Statistical Computing; 2014. R Foundation for Statistical

Computing. 2015.

38. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr Protoc*

Bioinformatics 2020; **69**: e96.

39. Revell LJ. phytools 2.0: an updated R ecosystem for phylogenetic comparative

methods (and other things). *PeerJ* 2024; **12**: e16505.

40. Ho L si T, Ané C. A linear-time algorithm for Gaussian and non-Gaussian trait

evolution models. *Syst Biol* 2014; **63**: 397–408.

41. Csűrös M. Ancestral reconstruction by asymmetric Wagner parsimony over

continuous characters and squared parsimony over distributions. *Comparative*

Genomics. 2008. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 72–86.

42. Kapust N, Nelson-Sathi S, Schönfeld B, Hazkani-Covo E, Bryant D, Lockhart PJ, et

al. Failure to recover major events of gene flux in real biological data due to method

misapplication. *Genome Biol Evol* 2018; **10**: 1198–1209.

43. Oyserman BO, Moya F, Lawson CE, Garcia AL, Vogt M, Heffernen M, et al.

Ancestral genome reconstruction identifies the evolutionary basis for trait

acquisition in polyphosphate accumulating bacteria. *ISME J* 2016; **10**: 2931–2945.

44. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics* 2019; **35**: 526–528.

45. Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, FitzJohn RG, et al.

Geiger v2.0: An expanded suite of methods for fitting macroevolutionary models to

phylogenetic trees. *Bioinformatics* 2014; **30**: 2216–2218.

46. de Crécy-Lagard V, Amorin de Hegedus R, Arighi C, Babor J, Bateman A, Blaby I,

et al. A roadmap for the functional annotation of protein families: a community

perspective. *Database (Oxford)* 2022; **2022**: baac062.

47. Koirala A, Brözel V. Phylogeny of nitrogenase structural and assembly components

reveals new insights into the origin and distribution of nitrogen fixation across

bacteria and Archaea. *Microorganisms* 2021; **9**: 1662.

48. Pi H-W, Lin J-J, Chen C-A, Wang P-H, Chiang Y-R, Huang C-C, et al. Origin and

evolution of nitrogen fixation in prokaryotes. *Mol Biol Evol* 2022; **39**.

49. Bellanger M, Figueroa JL 3rd, Tiemann L, Friesen ML, Iii RAW. NFixDB (Nitrogen

Fixation DataBase)-a comprehensive integrated database for robust 'omics analysis

of diazotrophs. *NAR Genom Bioinform* 2024; **6**: lqae063.

50. Nichio BT de L, Chaves RBR, Pedrosa FO, Raittz RT. Exploring diazotrophic

diversity: Unveiling nif core distribution and evolutionary patterns in nitrogen-fixing

organisms. *Research Square* . 2024.

51. Smercina DN, Evans SE, Friesen ML, Tiemann LK. To fix or not to fix: Controls on

free-living nitrogen fixation in the rhizosphere. *Appl Environ Microbiol* 2019; **85**.

52. Reese BK, Zinke LA, Sobol MS, LaRowe DE, Orcutt BN, Zhang X, et al. Nitrogen

cycling of active bacteria within oligotrophic sediment of the mid-Atlantic ridge flank.

Geomicrobiol J 2018; **35**: 468–483.

53. Kapili BJ, Barnett SE, Buckley DH, Dekas AE. Evidence for phylogenetically and

catabolically diverse active diazotrophs in deep-sea sediment. *ISME J* 2020; **14**:

971–983.

54. Semler AC, Paris ER, Salvador M, Dekas AE. Abundance, identity, and potential

diazotrophic activity of nifH-containing organisms at marine cold seeps. *Environ*

Microbiol 2025; **27**: e70058.

55. McRose DL, Zhang X, Kraepiel AML, Morel FMM. Diversity and activity of

alternative nitrogenases in sequenced genomes and coastal environments. *Front*

Microbiol 2017; **8**: 267.

56. Xie J-B, Du Z, Bai L, Tian C, Zhang Y, Xie J-Y, et al. Comparative genomic analysis

of N2-fixing and non-N2-fixing Paenibacillus spp.: organization, evolution and

expression of the nitrogen fixation genes. *PLoS Genet* 2014; **10**: e1004231.

57. Higdon SM, Pozzo T, Kong N, Huang BC, Yang ML, Jeannotte R, et al. Genomic

characterization of a diazotrophic microbiota associated with maize aerial root

mucilage. *PLoS One* 2020; **15**: e0239677.

58. Gallon JR. Tansley review no. 44. Reconciling the incompatible: N_2 fixation and O_2.

New Phytol 1992; **122**: 571–609.

59. Friedman R, Drake JW, Hughes AL. Genome-wide patterns of nucleotide

substitution reveal stringent functional constraints on the protein sequences of

thermophiles. *Genetics* 2004; **167**: 1507–1512.

60. Drake JW. Avoiding dangerous missense: thermophiles display especially low

mutation rates. *PLoS Genet* 2009; **5**: e1000520.

61. Groussin M, Gouy M. Adaptation to environmental temperature is a major

determinant of molecular evolutionary rates in archaea. *Mol Biol Evol* 2011; **28**:

2661–2674.

62. Miller RW, Eady RR. Molybdenum and vanadium nitrogenases of Azotobacter

chroococcum. Low temperature favours N2 reduction by vanadium nitrogenase.

Biochem J 1988; **256**: 429–432.

63. Walmsley J, Kennedy C. Temperature-dependent regulation by molybdenum and

vanadium of expression of the structural genes encoding three nitrogenases in

Azotobacter vinelandii. *Appl Environ Microbiol* 1991; **57**: 622–624.

64. Darnajoux R, Bradley R, Bellenger J-P. In vivo temperature dependency of

molybdenum and vanadium nitrogenase activity in the heterocystous Cyanobacteria

Anabaena variabilis. *Environ Sci Technol* 2022; **56**: 2760–2769.

65. Laso-Pérez R, Wu F, Crémière A, Speth DR, Magyar JS, Zhao K, et al.

644 Evolutionary diversification of methanotrophic ANME-1 archaea and their expansive

645 virome. *Nat Microbiol* 2023; **8**: 231–245.

646 66. Mehta MP, Baross JA. Nitrogen fixation at 92 degrees C by a hydrothermal vent

647 archaeon. *Science* 2006; **314**: 1783–1786.

648 67. Dekas AE, Poretsky RS, Orphan VJ. Deep-sea archaea fix and share nitrogen in

649 methane-consuming microbial consortia. *Science* 2009; **326**: 422–426.

650 68. Ivanovsky RN, Lebedeva NV, Keppen OI, Tourova TP. Nitrogen metabolism of an

651 anoxygenic filamentous phototrophic bacterium oscillocholris trichoides strain DG-6.

652 *Microbiology* 2021; **90**: 428–434.

653 69. Chen Y, Nishihara A, Haruta S. Nitrogen-fixing ability and nitrogen fixation-related

654 genes of thermophilic fermentative bacteria in the genus Caldicellulosiruptor.

655 *Microbes Environ* 2021; **36**.

656 70. Thiel V, Wood JM, Olsen MT, Tank M, Klatt CG, Ward DM, et al. The dark side of

657 the Mushroom Spring microbial mat: Life in the shadow of chlorophototrophs. :I.

658 microbial diversity based on 16S RRNA gene amplicons and metagenomic

659 sequencing. 2016. CRC Press, Boca Raton, FL.

660 71. Sabath N, Ferrada E, Barve A, Wagner A. Growth temperature and genome size in

661 bacteria are negatively correlated, suggesting genomic streamlining during thermal

662 adaptation. *Genome Biol Evol* 2013; **5**: 966–977.

663 72. Franche C, Lindström K, Elmerich C. Nitrogen-fixing bacteria associated with

664 leguminous and non-leguminous plants. *Plant Soil* 2009; **321**: 35–59.

665 73. Bennett AB, Pankievicz VCS, Ané J-M. A model for nitrogen fixation in cereal crops.

666 *Trends Plant Sci* 2020; **25**: 226–235.

74. Ferguson BJ, Indrasumunar A, Hayashi S, Lin M-H, Lin Y-H, Reid DE, et al.

Molecular analysis of legume nodule development and autoregulation. *J Integr*

Plant Biol 2010; **52**: 61–76.

75. Steele JH, Brink KH, Scott BE. Comparison of marine and terrestrial ecosystems:

suggestions of an evolutionary perspective influenced by environmental variation.

ICES J Mar Sci 2019; **76**: 50–59.

76. Dynarski KA, Houlton BZ. Nutrient limitation of terrestrial free-living nitrogen fixation.

New Phytol 2018; **217**: 1050–1061.

77. Zheng M, Zhou Z, Zhao P, Luo Y, Ye Q, Zhang K, et al. Effects of human

disturbance activities and environmental change factors on terrestrial nitrogen

fixation. *Glob Chang Biol* 2020; **26**: 6203–6217.

78. Yan G, Xing Y, Han S, Zhang J, Wang Q, Mu C. Long-time precipitation reduction

and nitrogen deposition increase alter soil nitrogen dynamic by influencing soil

bacterial communities and functional groups. *Pedosphere* 2020; **30**: 363–377.

79. Bellenger JP, Darnajoux R, Zhang X, Kraepiel AML. Biological nitrogen fixation by

alternative nitrogenases in terrestrial ecosystems: a review. *Biogeochemistry* 2020;

149: 53–73.

80. Sriswasdi S, Yang C-C, Iwasaki W. Generalist species drive microbial dispersion

and evolution. *Nat Commun* 2017; **8**: 1162.

81. Chen Y-J, Leung PM, Wood JL, Bay SK, Hugenholtz P, Kessler AJ, et al. Metabolic

flexibility allows bacterial habitat generalists to become dominant in a frequently

disturbed ecosystem. *ISME J* 2021; **15**: 2986–3004.

82. Carbonero F, Oakley BB, Purdy KJ. Metabolic flexibility as a major predictor of

spatial distribution in microbial communities. *PLoS One* 2014; **9**: e85105.

83. Van Tienderen PH. Evolution of generalists and specialists in spatially

heterogeneous environments. *Evolution* 1991; **45**: 1317–1331.

84. Poisot T, Bever JD, Nemri A, Thrall PH, Hochberg ME. A conceptual framework for

the evolution of ecological specialisation: Evolution of ecological specialisation. *Ecol*

Lett 2011; **14**: 841–851.

85. Hu W, Wang X, Xu Y, Wang X, Guo Z, Pan X, et al. Biological nitrogen fixation and

the role of soil diazotroph niche breadth in representative terrestrial ecosystems.

Soil Biol Biochem 2024; **189**: 109261.

86. Kloesges T, Popa O, Martin W, Dagan T. Networks of gene sharing among 329

proteobacterial genomes reveal differences in lateral gene transfer frequency at

different phylogenetic depths. *Mol Biol Evol* 2011; **28**: 1057–1074.

87. Waite DW, Vanwonterghem I, Rinke C, Parks DH, Zhang Y, Takai K, et al.

Comparative Genomic Analysis of the Class Epsilonproteobacteria and Proposed

Reclassification to Epsilonbacteraeota (phyl. nov.). *Front Microbiol* 2017; **8**: 682.

88. Moya-Beltrán A, Beard S, Rojas-Villalobos C, Issotta F, Gallardo Y, Ulloa R, et al.

Genomic evolution of the class Acidithiobacillia: deep-branching Proteobacteria

living in extreme acidic conditions. *ISME J* 2021; **15**: 3221–3238.

89. Campbell BJ, Smith JL, Hanson TE, Klotz MG, Stein LY, Lee CK, et al. Adaptations

to submarine hydrothermal environments exemplified by the genome of Nautilia

profundicola. *PLoS Genet* 2009; **5**: e1000362.

90. González-Rosales C, Vergara E, Dopson M, Valdés JH, Holmes DS. Integrative

genomics sheds light on evolutionary forces shaping the Acidithiobacillia class

acidophilic lifestyle. *Front Microbiol* 2021; **12**: 822229.

91. McCutcheon JP, Moran NA. Extreme genome reduction in symbiotic bacteria. *Nat*

Rev Microbiol 2011; **10**: 13–26.

92. Tripp HJ, Bench SR, Turk KA, Foster RA, Desany BA, Niazi F, et al. Metabolic

streamlining in an open-ocean nitrogen-fixing cyanobacterium. *Nature* 2010; **464**:

90–94.

93. Zehr JP, Shilova IN, Farnelid HM, Muñoz-MarínCarmen MDC, Turk-Kubo KA.

Unusual marine unicellular symbiosis with the nitrogen-fixing cyanobacterium

UCYN-A. *Nat Microbiol* 2016; **2**: 16214.

94. Cornejo-Castillo FM, Inomura K, Zehr JP, Follows MJ. Metabolic trade-offs

constrain the cell size ratio in a nitrogen-fixing symbiosis. *Cell* 2024; **187**: 1762-

1768.e9.

95. Coale TH, Loconte V, Turk-Kubo KA, Vanslembrouck B, Mak WKE, Cheung S, et

al. Nitrogen-fixing organelle in a marine alga. *Science* 2024; **384**: 217–222.

96. Anbar AD, Knoll AH. Proterozoic ocean chemistry and evolution: a bioinorganic

bridge? *Science* 2002; **297**: 1137–1142.

97. Rico KI, Garcia AK, Saito MA, Kaçar B, Anbar AD. Mysteries of metallome

evolution: Integrating insights from the Earth and life sciences. In: Anbar A, Weis D

(eds). *Treatise on Geochemistry*. 2025. Elsevier, pp 337–364.

98. Kaçar B. Reconstructing early microbial life. *Annu Rev Microbiol* 2024; **78**: 463–

492.

99. Falkowski PG. Life's engines: How microbes made earth habitable. *Life's Engines:*

How Microbes Made Earth Habitable 2015.