***Burkholderia* Genes Involved in Extracellular Bioplastic Degradation Revealed by Tn-seq**

and CRISPR-Cas

Running title: Extracellular Bioplastic Degradation in *Burkholderia* by Tn-seq and CRISPR-Cas

Zhong Ling Yap[a], Gabriel G. Lucas[b], Mohamad Safklou[a], Ryan Darragh[a], Andrew M. Hogan[a],

Anna Motnenko[a], Warren Blunt[c], Richard Sparling[a], David B. Levin[c], Dario Fernández Do Porto[b],

Silvia T. Cardona[a,d*]

[a]Department of Microbiology, University of Manitoba, Winnipeg, Manitoba, Canada R3T

2N2

[b]Plataforma de Bioinformática Argentina, Instituto de Cálculo, Ciudad Universitaria,

Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires (UBA), Ciudad

Autónoma de Buenos Aires, Argentina

[c]Department of Biosystems Engineering, University of Manitoba, Winnipeg, Manitoba,

Canada R3T 5V6

[d]Department of Medical Microbiology and Infectious Diseases, University of Manitoba,

Winnipeg, Manitoba, Canada R3E 0J9

email: silvia.cardona@umanitoba.ca

Keywords: *Burkholderia*, *Burkholderia vietnamiensis*, Transposon mutagenesis, CAST, gene

insertion

Abstract

Bioplastics represent promising alternatives to petroleum-based plastics, yet their biodegradation

remains insufficiently understood. Identifying bacteria capable of degrading bioplastics

extracellularly could enhance end-of-life management practices. To investigate *Burkholderia*'s

capacities for the degradation of medium-chain-length polyhydroxyalkanoate (mcl-PHA), we

screened a panel of *Burkholderia* strains and identified such capacity in some strains of *B. gladioli*,

B. multivorans, and *B. vietnamiensis*. To elucidate the genetic basis of this activity, we performed

transposon mutagenesis followed by activity-based screening and Tn-seq on *B. vietnamiensis*

LMG 16232. Transposon insertions with negative phenotype were identified in genes encoding a

triacylglycerol lipase, a lipase chaperone, type II secretion system components, HTH-type

transcriptional activators, and a polyhydroxyalkanoate synthesis repressor. To validate the

involvement of these genetic elements in mcl-PHA degradation, we generated insertional mutants

using a CRISPR-associated transposase system (CAST). Complete loss or reduced extracellular

mcl-PHA depolymerase activity was observed in the CAST mutants, validating their involvement

in mcl-PHA degradation. In addition, the interruption of these genetic elements showed a loss of

lipase activity, suggesting that the enzymes responsible for mcl-PHA depolymerase are lipases

with substrate ambiguity; moreover, docking experiments supported these findings. Together, we

identify *B. vietnamiensis* as a source of enzymes capable of degrading extracellular mcl-PHA and

demonstrate the power of combining activity-based screening, Tn-seq, and CAST to rapidly

establish gene-to-function links.

Importance

Due to their versatile metabolism, *Burkholderia* strains play critical roles in degradation of multiple compounds in the environment. Here we show that several Burkholderia species can degrade medium-chain-length polyhydroxyalkanoates (mcl-PHAs), a promising class of bioplastics. By integrating transposon mutagenesis, Tn-seq, and CRISPR-associated transposase (CAST) technologies, we identify and validate key genetic determinants involved in mcl-PHA degradation in B. vietnamiensis. These genes encode a lipase, a secretion system component, and regulatory factors, underscoring the complexity and specificity of microbial bioplastic degradation pathways. The findings not only advance our understanding of PHA biodegradation but also identifies *B. vietnamiensis* as as a source of enzymes capable of degrading extracellular mcl-PHA.

Introduction

Environmental pollution from petroleum-based plastic has driven growing interest in bioplastics, which are bio-based, biodegradable substitutes. Among these, poly(3-hydroxyalkanoates) (PHAs) are biologically produced polyesters that have a wide-range of applications in various sectors, such as packaging, agriculture, and biomedical devices (1). PHAs are synthesized by many microorganisms for intracellular energy and carbon storage under conditions of nutrient imbalance, namely nitrogen and phosphorous limitation. Depending on the number of carbon atoms in each 3-hydroxyalkanoate monomer unit, PHAs can be broadly classified as either short-chain-length PHAs (scl-PHA), which consist of 3–5 carbons per monomer or medium-chain-length PHAs (mcl-PHA), which contain, 6–14 carbons per monomer, each with different physico-chemical properties (2). The monomer of the carbon side chain length of PHAs will influence the enzyme-substrate specificity for degradation (3).

PHAs are degraded by either intracellular or extracellular PHA depolymerases. Intracellular PHA depolymerases hydrolyze PHA located in carbonosomes, which are endogenous carbon reservoirs or in native amorphous PHA granules, which have a surface layer of proteins and phospholipids (4). Extracellular PHA depolymerases degrade exogenous denatured granules that are semi-crystalline and lack the surface layer (3). Both intracellular and extracellular PHA depolymerases are carboxylesterases and belong to the α/β-hydrolase fold family (5). PHA depolymerases have been isolated and biochemically characterized in Ascomycetes, Firmicutes, Proteobacteria, and Actinobacteria (6).

Many studies have reported microorganisms that degrade scl-PHA and have purified, and characterized the corresponding enzymes (5, 6). However, microorganisms that can degrade mcl-PHA are relatively rare compared to those that can degrade scl-PHA (6). Most of the extracellular

mcl-PHA depolymerases that have been purified and biochemically validated belong to

Pseudomonas sp. (7–9), the obligated predator *Bdellovibrio bacteriovorus* (10), *Thermus*

thermophilus (11) and the Actinobacteria *Streptomyces* sp (12–14). Amino acid sequence

similarity searches of metagenomes using biochemically validated extracellular mcl-PHA

depolymerases as queries, have identified candidate extracellular mcl-PHA depolymerases only in

the taxonomic groups Myxococcota, Proteobacteria, Actinobacteriota and Bdellovibrionota (6).

These studies suggest that mcl-PHA degradation is not widely distributed among Bacteria.

While all extracellular PHA depolymerases hydrolyze PHA, extracellular mcl-PHA

depolymerases show little overall sequence homology with scl-PHA despite both being involved

in the degradation of PHAs. However, most PHA depolymerases share structural and mechanistic

characteristics with lipases and other α/β-hydrolase enzymes (15). Both PHA depolymerases and

lipases have a catalytic triad consisting of serine, histidine, and aspartic acid at the active site. The

catalytic serine is located within the GxSxG sequence motive (also known as lipase box), also

found in other α/β-hydrolases. Furthermore, a conserved non-catalytic histidine near the oxyanion

hole is positioned and functions similarly to the corresponding residue in lipases (3, 5, 16),

underscoring putative functional promiscuity between PHA depolymerases and lipases. Indeed,

extracellular lipases were purified from various species, including *Bacillus subtilis, Pseudomonas*

aeruginosa, Pseudomonas alcaligenes, and *Pseudomonas fluorescens* (16). However, most of

these lipases were not able to hydrolyze poly3-hydroxyhexanoate or poly3-hydroxybutyrate.

While the extracellular lipases of *Pseudomonas chlororaphis* PA23 and *Acinetobacter lwoffii* can

hydrolyze various PHA polymers, including mcl-PHA: polyhydroxyoctanoate (PHO) and

polyhydroxydecanoate (17), it is not known if *Burkholderia* strains, which are known lipase

producers (18–21) possess the capacity to degrade extracellular mcl-PHA.

In this work, we used poly(3-hydroxyoctanoate) (PHO) to explore extracellular mcl-PHA

degradation in *B. vietnamiensis* LMG 16232 and identified the genetic elements involved in this

process. Transposon mutagenesis followed by activity-based screening and Tn-seq identification

of transposon insertions linked genes with loss of extracellular mcl-PHA degradation. The

identified genes encode a putative triacylglycerol lipase, a lipase chaperone, type II secretion

system components, an HTH-type transcriptional activator, and a polyhydroxyalkanoate synthesis

repressor. By combining activity-based screening with the power of random and site-directed

transposon mutagenesis to rapidly identify all genetic elements that contribute to a specific

function, we were able to uncover genes associated mcl-PHA degradation that would not have

been predicted based on gene annotation.

Results

Diversity of extracellular mcl-PHA degradation activity in Burkholderia strains

To identify mcl-PHA degradation activity in the *Burkholderia* genus, we performed an

initial screen in a solid minimal medium containing PHO and limiting amounts of phenylalanine,

supplemented with Nile red to enhance the visualization of PHO degradation as a clear halo around

isolated bacterial colonies. Phenylalanine was chosen as a secondary carbon source in the medium

because (i) it is known to be catabolized by *Burkholderia* species (22)*,* and (ii) its metabolism is

known to influence several bacterial pathways (23). The plates were incubated for 20 days to

ensure cells grew densely and produced sufficient enzyme to visually detect mcl-PHA degradation.

After dense growth was achieved, a clear halo was observed around some colonies, suggesting

extracellular PHO degradation.

Several strains of *B. gladioli*, *B. multivorans*, and *B. vietnamiensis,* were able to produce a

visible halo suggesting extracellular mcl-PHA degradation (**Table 1, Fig. S1**). Of the strains with

activity, *B. gladioli* strain VC19233, *B. multivorans* 21NH942533, and VC9825 exhibited weak

activity in degrading mcl-PHA; *B. stagnalis* strains MSMB618, MSMB642, and *B. vietnamiensis*

LMG 16232 had moderate activity in degrading mcl-PHA. There were two strains, *B. multivorans*

F9091687 and *B. gladioli* 132208, where the temperature affected the activity. No mcl-PHA

depolymerase activity was observed for *B. cenocepacia*, *B. contaminans,* and *B. dolosa* isolates

tested (**Table 1**).

Genetic basis of B. vietnamiensis LMG 16232 extracellular mcl-PHA degradation activity

To determine the gene elements involved in the extracellular mcl-PHA degradation activity

we chose *B. vietnamiensis* LMG 16232 as ongoing work in our lab suggested its genetic

manipulation amenability by a ShCAST-based genetic tool for targeted insertional mutagenesis

(24). After sequencing its genome (25) we performed transposon mutagenesis and activity-based

screening followed by identification of transposon sites with Illumina sequencing (Tn-seq) (**Fig.**

1). To achieve genomic coverage of transposon insertions in *B. vietnamiensis* LMG 16232, we

determined the estimated transposon insertion density using an analytical model based on a

Poisson distribution of transposon insertions (26) (see material and methods).

The primary screening of single isolated transconjugants identified 84 colonies that did not

produce a visible halo. These colonies were subjected to secondary screening, where each isolated

transconjugant clone was propagated and screened on solid medium containing mcl-PHA. We

observed that 66 transconjugants had lost the ability to degrade extracellular mcl-PHA, showing

no halo (Pool 1). In contrast, 18 transconjugants exhibited a reduced halo (Pool 2). The complete

list of transposon mutants from pool 1 and pool 2 are shown in Table 2.

Further examination of the putative functions of the genes identified by transposon

insertions identified a gene cluster that encodes a type II secretion system (**Fig. 2A**), and a lipase

gene cluster, encoding triacylglycerol lipases, (designated as *lip1* and *lip2* herein) and a lipase

chaperone, (designated as *lipO*) (**Fig. 2B**). Additionally, three transposons mapped to a *pha* gene

cluster. Transposons in this cluster were found in *phaR* (P4G95_13560), a polyhydroxyalkanoate

synthesis repressor and *phaA* (P4G95_13570), which encodes for an acetyl-CoA acetyltransferase

involved in the biosynthesis of polyhydroxybutyrate (**Fig. 2C**). In addition, a gene coding for a

A24 family peptidase (P4G95_07805) and four genes encoding transcriptional regulators were

identified, which are a TetR/AcrR family transcriptional regulator (P4G95_00865), a MarR family

winged helix-turn-helix transcriptional regulator CDS (P4G95_05570), *phaR* (P4G95_13560), the

polyhydroxyalkanoate synthesis repressor, and a helix-turn-helix domain-containing protein CDS

(P4G95_16325) (Table 2). Together these results suggest that *lip1* (P4G95_16805) might be the

enzyme responsible for extracellular mcl-PHA degradation. The lipase encoded by *lip1* contains a

signal peptide probably cleaved by the A24 family peptidase (P4G95_07805). In the periplasm,

the lipase chaperone, *lipO* (P4G95_16815), may be responsible for folding and activating the

lipase prior to secretion. Finally, the lipase may be then finally excreted to extracellular space via

a type II secretion system (P4G95_05485, P4G95_05490, P4G95_05495, P4G95_05500,

P4G95_05505, P4G95_05510, P4G95_05515, P4G95_05520, P4G95_05525, P4G95_05530,

P4G95_05535).

A few genetic elements obtained from the Tn-seq analysis had a less clear relation with

extracellular mcl-PHA degradation in *B. vietnamiensis* LMG 16232. These were genes encoding

for trehalose-6-phosphate phosphatase (P4G95_10885), hypothetical proteins (P4G95_20805 and

P4G95_20810), the N-acetylglucosamine-6-phosphate deacetylase, NagA (P4G95_03530), a

phenylacetate-CoA oxygenase/reductase subunit PaaK and PaaJ (P4G95_06510 and

P4G95_06515) and an acetyl-CoA acetyltransferase PhaA (P4G95_13570).

Characteristics of a mcl-PHA degrading enzyme in B. vietnamiensis LMG 16232 predicted via in

silico analysis

Bacterial lipases are known to be capable of degrading polymers, such as polycaprolactone,

polyethylene terephthalate, and mcl-PHA (16, 27). The genome of *B. vietnamiensis* LMG 16232,

contains three genes putatively associated with lipase function: P4G95_16805 and P4G95_16810,

both coding for a triacylglycerol lipase, and P4G95_14150, coding for a lipase family protein.

While these candidate genes could encode enzymes capable for mcl-PHA degradation, our

activity-based screening followed by Tn-seq analysis identified that clones that have lost the halo

of PHO degradation, had transposon insertions in the P4G95_16805 locus (*lip1*), but not in the

other genes with predicted lipase activity.

The *lip1*-encoded protein, GenBank: WHU94860.1 is a putative triacylglycerol lipase that

shows common features to other extracellular lipases and mcl-PHA depolymerase, such as

containing a signal peptide and a catalytic triad composed of a serine (embedded in a lipase box

motif), histidine and aspartic acid (5). This enzyme is predicted to have 364 amino acids, with a

calculated molecular weight of 37.6 kDa and a theoretical isoelectric point (pI) of 6.43, computed

with Expasy (28). The computed molecular mass and isoelectric point are in the range of currently

known mcl-PHA depolymerase, where the molecular range is approximately 25 to 41 kDa, and

the pI value is mainly in the acidic range (11). This enzyme was predicted via SignalP 5.0 to be

secreted via secretory signal peptides transported by the Sec translocon and cleaved by Signal

Peptidase I (Lep) with a probability of 0.83. Signal peptidase I is essential in many bacteria

functions, and it could be the reason why insertional mutants in *LepB* (P4G95_10750) were not

recovered. The cleavage site position was predicted between 43-44 with a probability of 0.5069.

The presence of the motive (G-H-S-Q-G) in the amino acid sequence encode by *lip1* aligns with

the lipase box consensus sequence. This sequence has also been found in a previously identified

novel subgroup of extracellular mcl-PHA depolymerases from Actinobacteria, specifically those

from *S. roseolus* and *S. venezuelae*, and *S. exfoliates* (12–14).

In contrast, extracellular mcl-PHA depolymerases identified from the genus *Pseudomonas*

(7–9), as well as in *Bdellovibrio* (10), exhibit a different lipase consensus sequence, which is (G-

I-S-S-G). This putative protein sequence (WHU94860.1) shows very little sequence similarity to

currently known enzymes which can degrade mcl-PHA from the genus *Streptomyces*,

Pseudomonas, and *Bdellovibrio,* where there is only 9 to 12 % sequence identity (**Fig. 3**). The

enzyme also has a putative catalytic amino acid (Ser131-Asp308-His330). In summary, while *in*

silico analysis shows that P4G95_16805 codes for a lipase, the low similarity of this protein with

other mcl-PHA depolymerases makes *in silico* prediction for PHA degradation activity difficult.

While Tn-seq analysis strongly suggests that the activity of the encoded protein is that of an

extracellular mcl-PHA depolymerase, further validation is required.

Genetic elements responsible for mcl-PHA degradation are confirmed by RhaCAST

To validate the genetic elements involved in extracellular mcl-PHA degradation activity

identified by Tn-seq, we created independent insertional mutants using RhaCAST (24) and

investigated their phenotypes. RhaCAST is a genetic tool we recently developed for targeted

insertional mutagenesis in *Burkholderia cepacia* complex and previously applied on *B.*

cenocepacia and *B. multivorans.* We applied RhaCAST in *B. vietnamiensis* LMG 16232

interrupting the coding genes for the putative lipase (*lip1*, P4G95_16805), A24 family peptidase

(*A24*, P4G95_07805), the TetR/AcrR family transcriptional regulator (*tetR/acrR*, P4G95_00865),

trehalose-phosphatase (*otsB,* P4G95_10885), and a gene from type II secretion system (*gspE*,

P4G95_05530) (**Fig. S2-S6**). These mutants did not produce a halo around the mutant strain in an

agar plate containing mcl-PHA (**Fig. 4**).

To further confirm that the halo of degradation observed resulted from the putative lipase

is mcl-PHA degradation, a plug of agar from underneath the inoculated mutant was taken,

lip1::CAST-Tp and wild type as control from the agar-based assay. We determined the absence

and presence of PHO in the agar with gas chromatography with flame ionization detection (GC-

FID). As expected, no PHO was detected in the agar beneath the WT strain, whereas PHO was

detected under the *lip1*::CAST-Tp strain, indicating that PHO was not being degraded (**Fig. 5**).

To determine if the *B. vietnamiensis* LMG 16232 putative lipase and the genes that we

chose to make the insertional mutant are involved in mcl-PHA degradation are also involved in

lipid degradation, we tested *B. vietnamiensis* LMG 16232 and its mutants for tributyrin

degradation. The wild type *B. vietnamiensis* LMG 16232 exhibited high activity against tributyrin.

The mutant *lip1::CAST-Tp* exhibited reduced activity in degrading tributyrin, suggesting other

lipases or esterases might be produced and secreted. By disrupting *gspE,* which is part of the gene

cluster encoding for the type II secretion system, the mutant (*gspE::CAST-Tp*) showed complete

abrogation of tributyrin degradation, indicating that the lipases or esterases might be produced and

secreted via type II secretion system. Disrupting the gene encoding for A24 family peptidase also

completely abolishes tributyrin degradation, suggesting that the lipases or esterases responsible for

tributyrin degradation are extracellular enzymes with a leader peptide. For the TetR/AcrR family

transcriptional regulator insertional mutant, the ability to degrade tributyrin was reduced to

residual activity (**Fig. 6**).

In silico analysis: structural properties of catalytic site and docking

In the gene cluster of *lip1,* there is another lipase-encoding gene, *lip2* located adjacent to

lip1 (**Fig. 2**). This lipase gene cluster comprises *lip1*, *lip2* and *lipO* (P4G95_16805, P4G95_16810,

and P4G95_16815). Transposon insertions located in clones with no halo of degradation that

mapped to this region were located in *lip1* and *lipO*, but not in the *lip2* gene, posing the question

of whether *lip2* codes for an active mcl-PHA depolymerase. To investigate this possibility, we

decided to analyze and compare the predicted structures of these proteins. To determine the

differences in the structural properties of the target proteins, we predicted the structures of Lip1

and Lip2 using AlphaFold 2 and analyzed multiple sequence alignments, AlphaFold predicted

structural confidence, and inter-residue error estimates. The sequence coverage analysis, which

combines alignment depth and sequence identity, illustrates the extent of conservation across

homologous sequences for both proteins (**Fig. S7A**). A well-conserved core region (residues ~50–

160) is evident in both cases, with lower coverage and diversity in the N- and C-terminal regions

suggesting flexible or functionally variable domains. To evaluate the reliability of the structural

predictions, we examined the Predicted Aligned Error (PAE) matrices of the five top-ranked

AlphaFold models (**Fig. S7B**). As expected, low predicted error values (blue) correspond to well-

defined domains, while higher values (red) are associated with flexible or ambiguous regions. The

consistent patterns observed across different models support a robust domain architecture with

potential inter-domain flexibility. Additionally, the predicted Local Distance Difference Test

(pLDDT) scores (**Fig. S7C**) further validate the confidence in structural predictions. The high

pLDDT values (>90) in the core region confirm a well-defined fold, whereas significant drops

around residues 50–100 and 200–250 suggest intrinsically disordered or flexible regions. These

trends are consistent across both Lip1 and Lip2, reinforcing the reliability of the predictions and

indicating a shared structural organization.

The superposition of the AlphaFold-predicted Lip1 and Lip2 (not shown) structures with

the 4LIP crystallographic template (**Fig. S8A**) revealed a high degree of structural similarity,

particularly in the conserved secondary structure elements. However, a notable difference was

observed in the predicted models of Lip1 and Lip2, where an alpha-helix appears to obstruct a

cavity present in 4LIP (**Fig. S8B**).

To evaluate the catalytic site and determine whether Lip1 and Lip2 possess enzymatic

activity, we performed a structural superposition with 4LIP. The overlay reveals a remarkable

similarity in the spatial arrangement of the catalytic triad (**Fig. S9**). The conserved positioning of

the key catalytic residues strongly suggests that Lip1 and Lip2 adopt a functional fold compatible

with lipase activity. This structural conservation provides additional support for the hypothesis

that both proteins share mechanistic features with known lipases.

Using 4LIP as a template, we guided the docking of LIP1 and LIP2 with two substrates: 3-

Hydroxyoctanoate (OCT) and Tributyrin (TRI). Both ligands, OCT and TRI, were energy-

minimized before docking, and a grid-based docking algorithm was used to position the ligands

within the predicted active sites of both proteins. Notably, all combinations of protein-ligand

docking yielded clusters with low energy, indicating favourable interactions (**Table S1**).

Docking simulations with Lip1 revealed a single predominant low-energy cluster for both

OCT and TRI (**Fig. 7A, B, E, Table S1**). This convergence, together with the consistently low

binding energies, strongly suggests a high affinity between Lip1 and both ligands. In contrast, the

docking of both ligands against Lip2 (**Fig. 7C, D, E, Table S1**) yielded three distinct low-energy

clusters for each substrate. This multiplicity of binding modes suggests a weaker or less specific

interaction, potentially due to a more flexible or heterogeneous binding pocket that compromises

optimal substrate accommodation. Structural analysis of the catalytic site revealed that Lip1 and

4LIP share a similar configuration, whereas Lip2 contains a different amino acid at position 73

compared to Lip1, which is GLN (Lip2) instead of TYR (Lip1) (**Fig. 8, Fig. S10**). This amino acid

difference may explain the activities observed between Lip1, which retains enzymatic function,

and Lip2, in which detectable activity could not be linked to the *lip1* gene by loss of function-

derived transposon mutagenesis.

Discussion

The production and use of bioplastic, including mcl-PHA, has increased recently. These

materials are considered alternatives to petroleum-based plastics because they can be derived from

renewable resources and have a wide range of applications (1). Despite growing interest,

microorganisms and enzymes capable of degrading mcl-PHA remain relatively understudied (6).

Extracellular mcl-PHA depolymerases have been found in various genera, including *Pseudomonas*

(7–9), *Bdellovibrio* (10), *Thermus* (11) and *Streptomyces* (12–14). However, to date, the

mechanism of action for extracellular mcl-PHA by microorganisms has not been thoroughly

explored. Here, we report that several *Burkholderia* strains, including *B. gladioli*, *B. multivorans*,

and *B. vietnamiensis,* can degrade mcl-PHA. Through transposon mutagenesis combined with Tn-

seq analysis, we identified genetic elements associated with mcl-PHA degradation in *B.*

vietnamiensis LMG 16232. We further validated some of these genetic elements by generating

insertional mutants using a CRISPR-associated transposase system (CAST) in extracellular mcl-

PHA degradation activity. Our findings indicate that the enzyme responsible for extracellular mcl-

PHA degradation is a lipase with substrate versatility.

A technique commonly used to enhance the visualization of mcl-PHA degradation involves

incorporating Nile red, a lipophilic fluorescent dye, into agar plates (29). Thus, we adopted this

visualization method in our screening medium to increase sensitivity. In comparison to other

studies, where the degradation of extracellular mcl-PHA can be observed within 2 to 3 days,

Burkholderia vietnamiensis LMG16232 exhibited a slower degradation process, requiring

approximately 7 days for observable activity. Some strains of *Burkholderia* from the initial

screening showed weak activity, with degradation taking up to 20 days, despite the use of Nile red

to enhance detection. Nevertheless, by implementing this visualization technique and alongside

transposon mutagenesis in *B. vietnamiensis* LMG 16232, we were able to screen 50,000 individual

colonies in a rapid manner. This approach enabled us to identify transconjugant colonies that had

lost the ability to degrade mcl-PHA, providing a sensitive and efficient screening method. This

method may be applicable to other polymers that can be stained with suitable dyes, enhancing

assay sensitivity.

According to the data on the PHA depolymerase engineering database, several species of

Burkholderia, including *Burkholderia thailandensis*, *Burkholderia ambifaria, Burkholderia*

graminis, *Burkholderia cenocepacia*, *Burkholderia dolosa*, *Burkholderia multivorans*, *B.*

vietnamiensis and *Burkholderia oklahomensis* have genes encoding proteins putatively capable of

degrading extracellular scl-PHA (5). While *B. cepacia* DP1 has been shown to degrade

extracellular scl-PHA (30, 31), there are no reports of *Burkholderia* species that degrade

extracellular mcl-PHA. We show evidence of *Burkholderia*'s capability to degrade extracellular

mcl-PHA, linking the enzymatic activity to the P4G95_16805 locus of *B. vietnamiensis* LMG

16232.

Burkholderia is a is known as a major lipase producer. Many strains can secrete

extracellular lipases under inducing conditions, and some of those lipases are capable of degrading

various compounds (32–35). For example, lipases from *Burkholderia* stains MC 16-3 (GenBank:

AY772173) and 99-2-1 (GenBank: AY772174) can degrade methyl(R)-N-(2,6-dimethyl-phenyl)alaninate (36). Additionally, lipase from *B. cepacia* PBSA-1 (GenBank: EF189714.1) has been shown to degrade poly(butylene succinate-co-butylene adipate) (37), while *B. cepacia* lipase can degrade Polycaprolactone (PCL) (16). These lipases, including the one identified in this work, share a conserved lipase consensus sequence of G-H-S-Q-G, resembling the oxyanion hole consensus sequence. Enzymes that degrade extracellular mcl-PHA exhibit similar characteristics to these extracellular lipases, including the presence of the lipase box and a conserved non-catalytic histidine near the oxyanion hole (3, 5, 16). Consequently, this study suggests that the mcl-PHA degrading enzyme from *Burkholderia* are lipases that also possess the capability to degrade mcl-PHA.

Besides the identification of the gene responsible for mcl-PHA degradation activity, transposon mutagenesis combined with Tn-seq identified other genetic elements involved in the degradation of mcl-PHA in *B. vietnamiensis* LMG 16232. These genes can be classified into three functional groups: a) lipase/chaperone gene cluster, b) components of type II secretion system, and c) regulatory elements. The genetic elements associated with the mcl-PHA degradation in *B. vietnamiensis* LMG 16232 are consistent with the findings in various *Burkholderia* lipases, particularly regarding the production and secretion of extracellular lipase. It is well established that *Burkholderia* extracellular lipases are secreted through the type II secretion system. In this process, the N-terminal signal peptide is transported across the inner membrane via the Sec secretion system. Once in the periplasm, a chaperone interacts with the lipase, facilitating the formation of the active form of the lipase (38, 39). This work also supports that the extracellular lipase of *Burkholderia* depends on a dedicated chaperone. In *B. glumae*, this chaperone (lipase-specific foldase) facilitates lipase folding, functioning similarly to the propeptide domains found

in bacterial proteases (39). Similar to the lipases in *B. cepacia* and *B. glumae*, the putative lipase

responsible for degrading mcl-PHA from *B. vietnamiensis* LMG 16232 is predicted to have an N-

terminal signal peptide. In addition, the gene encoding a chaperone is located downstream of the

putative lipase gene and is predicted to assist the correct folding and efficient secretion of this

putative lipase (40, 41). This putative enzyme (WHU94860.1), encoded by *lip1,* is likely secreted

via the general secretion pathway (GSP), also known as the type II secretion system, which relies

on a signal sequence for translocation. This mechanism is similarly predicted for the excretion of

the enzyme that degrades extracellular mcl-PHA in *B. bacteriovorus* (10).

One of the putative regulatory elements identified in this work is a TetR/AcrR family

transcriptional regulator CDS (P4G95_00865). This type of regulator is widely distributed among

bacteria and controls various cellular processes, including the metabolism of lipid compounds,

such as the synthesis and degradation of fatty acid and the storage of polymer (42). The

involvement of other regulatory elements, such as HTH-type transcriptional activators and a PHA

synthesis repressor, suggests that extracellular mcl-PHA degradation activity of *B. vietnamiensis*

LMG 16232 may be integrated with other metabolic pathways. While we did not identify a specific

genetic element directly responsible for the uptake of mcl-PHA degradation products, we propose

passive diffusion. This hypothesis is based on the structural similarity between mcl-PHA

monomers and fatty acids, which both contain a hydrophobic alkyl chain (43). Medium-chain fatty

acids (C_7-C_{11}) can enter bacteria through simple diffusion without the involvement of active

transport systems or an unknown mechanism (44, 45) (**Fig. 9**).

Although our Tn-seq experiment was expected to saturate the whole genome, we cannot

rule out the possibility that the lipase production cluster (**Fig. 2**) may constitute a cold spot for

transposon insertions. Thus, this could decrease the probability of obtaining clones with insertions

in *lip2* having insertions with a halo of degradation. Furthermore, since *lip2* and *lipO* probably form an operon, an insertion in *lip2* may not be informative about its function due to the potential polar effect on *lipO*. Therefore, we took an *in silico* approach to analyze and compare the predicted structures of Lip1 and Lip2. Our structural analysis of Lip1 and Lip2, based on AlphaFold predictions, highlights a conserved core architecture and variable N- and C-terminal regions, suggesting potential regulatory or accessory roles. The Predicted Aligned Error (PAE) matrices confirm a stable domain architecture with flexible inter-domain regions, while high pLDDT values reinforce confidence in the structural models. Superposition with 4LIP demonstrated overall structural similarity but also revealed an obstructive alpha-helix in Lip1 and Lip2, which is absent in 4LIP, which may influence enzyme activity. Further studies are needed to determine its functional relevance. Both proteins maintain a conserved catalytic triad, consistent with lipase activity. Docking experiments showed strong and specific binding for Lip1. In contrast, Lip2 exhibited multiple binding modes, indicating a more flexible binding pocket. A difference in the amino acid at position 73: Lip1 contains glutamine (GLN), while Lip2 has tyrosine (TYR), which may suggest reduced catalytic activity. Based on our experimental and docking results, Lip1 appears to function as an active lipase, while Lip2 may have altered activity due to structural differences. Additional biochemical and structural studies are needed to validate these conclusions.

Overall this study provided new insight onto the degradation of extracellular mcl-PHA by the genus *Burkholderia* thanks to a combination of activity screening, Tn-seq, and further validation with RhaCAST.

Material and methods

Strains, Selective Antibiotics, and Growth Conditions

Strains and plasmids are shown in **Table 3**. Primers used in this study are listed in **Table**

4. All strains were grown in LB-Lennox medium (Difco). Strains of *Burkholderia* and the strains

of *E. coli* were initially grown at 37 °C, except for *E. coli* Stbl2, which was grown at 30 °C. The

following selective antibiotics were used: chloramphenicol (Sigma; 20 µg/mL for *E. coli*),

trimethoprim (Sigma; 100 µg/mL for strains of *Burkholderia*, 50 µg/mL for *E. coli*), gentamicin

(Sigma; 50 µg/mL for *Burkholderia*).

Genomic and plasmid extraction

Genomic DNA of the *B. vietnamiensis* LMG 16232 mutants made with RhaCAST were

extracted using the PureLink microbiome DNA purification kit (Invitrogen). Genomic DNA of the

Tn-seq pools were extracted using standard isopropanol precipitation. Plasmid DNA was extracted

using the EZNA Plasmid DNA Mini Kit (Omega Bio-tek). Both genomic and plasmid DNA were

eluted or solubilized in Tris-HCl pH 8.5.

Substrate and media preparation

The Poly(3-hydroxyoctanoate) (PHO) colloidal suspension was created by dissolving 400

mg of PHO into 200 mL acetone (Polyferm Canada Inc., Canada). Once the PHO was dissolved,

40 mL of dH_2O was added. The acetone was evaporated off using a speed evaporator. The colloidal

suspension was stored at 4° C until needed.

The mcl-PHA containing medium consisted of an M9 minimal medium agar (BD Difco,

USA) with 2 mM phenylalanine, 2 mM $MgSO_4$, 10 mg/ mL of PHO and 0.3 µg/mL Nile red. PHO

was added to the medium by using a previously prepared colloidal suspension of PHO. The PHO,

Nile red, $MgSO_4$, and phenylalanine were added after autoclaving to the M9 agar whilst melted.

25 mL of medium was then poured into petri plates. After solidifying, the plates were stored at

4°C until needed.

Tributyrin medium was made using a minimal medium with 2 mM phenylalanine and 2

mM $MgSO_4$ with tributyrin added to a final concentration of 4% (v/v) to the medium according to

the Sigma-Aldrich instructions (Sigma, St. Louis, Missouri, USA). The tributyrin was added after

autoclaving whilst the agar medium was around 80°C. The medium with tributyrin was vigorously

mixed until ready to pour into petri plates. After solidifying, the plates were stored at 4°C until

needed.

Screening for mcl-PHA and tributyrin degradation

To determine the strain's ability to degrade mcl-PHA and tributyrin degradation. The

overnight cultures were grown at 30°C and washed twice with PBS. 20 µL of the culture was

inoculated after being adjusted to an OD_{600} of 1. For mcl-PHA degradation determination, the

plates were incubated for 7-20 days; For tributyrin degradation determination, the plates were

incubated for 3 days, both incubated at 30°C. The visualization of the halo degradation was

performed using AlphaImager EP (Cell Biosciences, Santa Clara, CA). For the mcl-PHA plates

containing Nile red, the image was taken under UV excitation. Colonies that appeared to have lost

the PHO degradation activity on the first round of screening were then subjected to a secondary

screening to rule out false positives. 66 and 18 transconjugants were isolated without or with weak

PHO degradation activity, respectively. *Transposon mutagenesis, Tn-seq library preparation and*

genome mapping

B. vietnamiensis LMG 16232 was subjected to transposon mutagenesis with

pRBrhaBoutgfp (46). A total of 50,000 single colonies were screened. This number is based on

the expected average transposon insertion density from the model, approximately eight transposon

insertions per gene. Assuming a genome with N = 6000 non-essential genes and a collection of N

= 50,000 transposon insertion mutants, the probability that a single gene will see no transposon

insertions is: $P(0, k, N) = e^{(-50,000/6,000)} = 2.4 \times 10^{-4}$; the total number of genes that will see no

insertions is: $N* e^{(-k/N)} = 6,000 \times 2.4 \times 10^{-4} = 1.4$ genes; and the total number of genes will be

found with this transposon density is n(insertion) $6,000* (1-e^{(-50,000/6,000)}) = 5,999$ genes.

Tn-seq was chosen to identify the transposon insertion sites. To enrich transposon-genome

junctions and to determine the insertion site, an adapted Tn-seq method, Tn-seq circle, developed

by Gallagher et al. (47), was performed as previously described with some minor modifications

(47, 48). Briefly, cultures of individual mutants were grown individually to approximately OD_{600}

of 3 and were combined into two separate pools, based on without (Pool 1) or with weak (Pool 2)

PHO degradation activity. The cultures were harvested, followed by the genomic DNA extraction.

The DNA was sheared to an average size of 300 bp via ultrasonication with a Covaris M220 (duty

factor of 20%, a peak incident power of 50W, 200 cycles per burst, and a duration of 75s), followed

by end repair with the NEBNext End Repair Module (NEB) and A-tailing. Next, the products were

ligated with adaptors, consisting of annealing oligos 683 and 684 with the Quick Ligation Kit

(NEB) and digested with *PacI* (NEB). Probe 682 in the presence of Ampligase (Lucigen) was used

to circularize fragments containing the transposon sequence. The exonuclease treatment was

carried out with a mix of T7 gene 6 exonuclease (ThermoFisher), Lambda exonuclease (NEB),

and Exonuclease I (NEB) to digest the linear fragments. The enrichment of the transposon–genome

junctions was confirmed by comparing the amplification of the transconjugant's pools with wild-

type *B. vietnamiensis* LMG 16232 using qPCR with iTaq Universal SYBR Green Supermix (Bio-

Rad). The number of PCR cycles corresponding to ~50% saturation by qPCR was chosen to

amplify transposon-genome junctions with iTaq DNA polymerases (Bio-Rad) for Illumina

sequencing. PCR primers 681, 690, 718, and 719 contain the Nextera indices and were used in

appropriate pairs (**Table 4**). The indexed samples were then cleaned-up with the SeraMag beads

(Cytiva), analysed on a TapeStation4150 (Agilent Technologies) and sequenced with an Illumina

MiSeq 2500. Raw reads were deposited in the NCBI Sequence Read Archive (SRA) repository

and will be publicly available after publication under BioProject: PRJNA1082765, accession:

SRR28216496 (Pool 1), SRR28216495 (Pool 2).

The sequenced reads of these mutants were mapped to the *B. vietnamensis* LMG 16232

genome with the Galaxy bioinformatics platform (https://usegalaxy.org). Using Cutadapt

(Galaxy v2.9+galaxy0) (49), the transposon sequence

5'-NNNNAATCTAGACTAGTGCGGCCGCACTTGTGTATAAGAGTCA-3' was removed

from the FASTQ file. Next, using Burrows-Wheeler transform (BWA) (Galaxy v0.7.17.4) (50),

the reads were mapped to the close genome of *B. vietnamiensis* LMG 16232 (25). The BWA's

BAM output was then converted to SAM using BAM-to-SAM (Galaxy 2.0.2) (51). The output

was visualized using Geneious 8.1.9.

Determination of the PHO content beneath inoculated culture

The inoculated culture from the phenotypic assay was scraped off from the agar. An agar

disk beneath the inoculated culture was cut using the back of the P1000 pipette tip. For control, a

disk was cut where no cell was inoculated. The samples were dried at 60°C for 16 hours, and the

PHO was extracted in chloroform using the acid-catalyzed methanolysis procedure as previously

described (52, 53). The determination of PHO in the agar was performed using an Agilent 7890A

gas chromatograph equipped with a DB-23 capillary column ($30\,m \times 250\,\mu m \times 0.25\,\mu m$) and a

flame ionization detector (Agilent Technologies, Santa Clara, CA). Method operating parameters

and peak quantification have been previously described (52, 54).

In silico analysis

The calculated molecular weight and theoretical isoelectric point (pI) of the putative enzyme were computed with Expasy (28). The putative catalytic amino acid was predicted with Phyre2 server (Protein Homology/analogY Recognition Engine V 2.0) (55). Signal peptides were predicted for predicted PHA depolymerases using Signal P 5.0 (https://services.healthtech.dtu.dk/service.php?SignalP) with default settings using both Gram-negative and Gram-positive predictions (56). The alignment of amino acid sequences was performed using Clustal Omega (https://www.ebi.ac.uk/jdispatcher/msa) (57). Protein conformation was predicted using ColabFold 2 (58). Molecular docking studies were conducted using Autodock 4 (59) to investigate the interaction between the enzyme and its potential substrates. The docking protocol was guided by the crystal structure of the reference protein (PDB ID: 4LIP). Visual representations of the predicted protein structures and docking interactions were generated using PyMOL version 3.0.3 (60).

Construction of B. vietnamiensis LMG 16232 insertional mutants

Genes were interrupted using the RhaCAST method reported previously (24). Briefly, 24 bp crRNAs were designed from the genes to be targeted, and the templates were ordered as single-stranded oligos and PCR-amplified using PCR primers 3392 and 3421 to create double-stranded DNA fragments. The PCR products were then introduced into pRhaCAST using Golden Gate cloning with SapI (NEB). The newly constructed plasmids were then transformed into *E. coli* Stbl2. pRhaCAST was then introduced to *B. vietnamiensis* LMG 16232 by tetra-parental mating (24). *E. coli* MM290/pRK2013 helper strain, *E. coli* Stbl2/pRhaCAST with corresponding crRNA, and *E. coli* PIR1/pDonorTp was conjugated into *B. vietnamiensis* LMG 16232 on LB agar plate containing 0.05% rhamnose. The plate was incubated at 37°C for 24 hours. Successful

exconjugants were selected on LB plates containing trimethoprim and gentamycin to select against

the *E. coli*. Colonies that appeared within 48 h at 37°C were screened by colony PCR. All

insertional mutants constructed with RhaCAST were verified by whole genome sequencing using

Oxford Nanopore Technologies platforms with v14 library prep chemistry and R10.4.1 flow cells

by Plasmidsaurus (Oregon, USA). The genomic sequence of the insertional mutants is available

after under BioProject: PRJNA1205771.

Author Contributions

STC and ZLY formulated the ideas and design of the project; GGL performed docking

experiment; ZLY and MS designed and cloned the plasmids and mutants; ZLY, MS, RD, and WB

performed phenotypic assays; ZLY and AMH performed library prep for Tn-seq. ZLY and AM

processed and analyzed the Tn-seq data; ZLY wrote the manuscript and edited together with STC,

GGL, MS, RD, AMH, AM, WB, RS, DBL, and DFDP; STC, DFDP, DBL, RS, and WB provided

financial support and supervised the work.

Acknowledgements and Funding Sources

This work was supported by a Natural Sciences and Engineering Research Council

(NSERC) of Canada Discovery Grant to SC and a University of Manitoba Collaborative Research

Program (UCRP) grant to SC and DBL. ZLY was supported by a Research Manitoba and

University of Manitoba Graduate Fellowship (UMGF); MS was supported by NSERC

Undergraduate Student Research Awards (USRA); AMH was supported by a Cystic Fibrosis

Canada studentship and a Canadian Institutes of Health Research (CIHR) Vanier award.

Conflict of Interest Statement

527 The authors declare no competing financial interests.

References

1. Muneer F, Rasul I, Azeem F, Siddique MH, Zubair M, Nadeem H. 2020. Microbial Polyhydroxyalkanoates (PHAs): Efficient Replacement of Synthetic Polymers. J Polym Environ 28:2301–2323.

2. Li Z, Yang J, Loh XJ. 2016. Polyhydroxyalkanoates: opening doors for a sustainable future. NPG Asia Mater 8:e265–e265.

3. Jendrossek D, Handrick R. 2002. Microbial degradation of polyhydroxyalkanoates. Annu Rev Microbiol 56:403–432.

4. Jendrossek D. 2009. Polyhydroxyalkanoate Granules Are Complex Subcellular Organelles (Carbonosomes). J Bacteriol 191:3195–3202.

5. Knoll M, Hamm TM, Wagner F, Martinez V, Pleiss J. 2009. The PHA Depolymerase Engineering Database: A systematic analysis tool for the diverse family of polyhydroxyalkanoate (PHA) depolymerases. BMC Bioinformatics 10:89.

6. Viljakainen VR, Hug LA. 2021. The phylogenetic and global distribution of bacterial polyhydroxyalkanoate bioplastic-degrading genes. Environ Microbiol 23:1717–1731.

7. Schirmer A, Jendrossek D. 1994. Molecular characterization of the extracellular poly(3-hydroxyoctanoic acid) [P(3HO)] depolymerase gene of Pseudomonas fluorescens GK13 and of its gene product. J Bacteriol 176:7065–7073.

8. Kim DY, Kim HC, Kim SY, Rhee YH. 2005. Molecular characterization of extracellular medium-chain-length poly(3-hydroxyalkanoate) depolymerase genes from Pseudomonas alcaligenes strains. J Microbiol Seoul Korea 43:285–294.

9.   Rhee YH, Kim YH, Shin K-S. 2006. Characterization of an extracellular poly (3-hydroxyoctanoate) depolymerase from the marine isolate, *Pseudomonas luteola* M13-4. Enzyme Microb Technol 38:529–535.

10.   Martínez V, de la Peña F, García-Hidalgo J, de la Mata I, García JL, Prieto MA. 2012. Identification and Biochemical Evidence of a Medium-Chain-Length Polyhydroxyalkanoate Depolymerase in the Bdellovibrio bacteriovorus Predatory Hydrolytic Arsenal. Appl Environ Microbiol 78:6017–6026.

11.   Papaneophytou CP, Velali EE, Pantazaki AA. 2011. Purification and characterization of an extracellular medium-chain length polyhydroxyalkanoate depolymerase from *Thermus thermophilus* HB8. Polym Degrad Stab 96:670–678.

12.   Martínez V, de Santos PG, García-Hidalgo J, Hormigo D, Prieto MA, Arroyo M, de la Mata I. 2015. Novel extracellular medium-chain-length polyhydroxyalkanoate depolymerase from Streptomyces exfoliatus K10 DSMZ 41693: a promising biocatalyst for the efficient degradation of natural and functionalized mcl-PHAs. Appl Microbiol Biotechnol 99:9605–9615.

13.   Gangoiti J, Santos M, Prieto MA, de la Mata I, Serra JL, Llama MJ. 2012. Characterization of a Novel Subgroup of Extracellular Medium-Chain-Length Polyhydroxyalkanoate Depolymerases from Actinobacteria. Appl Environ Microbiol 78:7229–7237.

14.   Santos M, Gangoiti J, Keul H, Möller M, Serra JL, Llama MJ. 2013. Polyester hydrolytic and synthetic activity catalyzed by the medium-chain-length poly(3-hydroxyalkanoate) depolymerase from *Streptomyces venezuelae* SO1. Appl Microbiol Biotechnol 97:211–222.

15.   Gupta R, Gupta N, Rathi P. 2004. Bacterial lipases: an overview of production, purification and biochemical properties. Appl Microbiol Biotechnol 64:763–781.

16.   Jaeger KE, Steinbüchel A, Jendrossek D. 1995. Substrate specificities of bacterial

polyhydroxyalkanoate depolymerases and lipases: bacterial lipases hydrolyze poly(omega-

hydroxyalkanoates). Appl Environ Microbiol 61:3113–3118.

17.   Sharma PK, Mohanan N, Sidhu R, Levin DB. 2019. Colonization and degradation of

polyhydroxyalkanoates by lipase-producing bacteria. Can J Microbiol 65:461–475.

18.   Rathi P, Saxena RK, Gupta R. 2001. A novel alkaline lipase from *Burkholderia cepacia* for

detergent formulation. Process Biochem 37:187–192.

19.   Oliveira BH, Santos RÉ, Loiola LEA, Nascimento VMG. 2015. Overproduction and properties of

lipase by a wild strain of Burkholderia lata LBBIO-BL02 using chicken fat. 2. Ann Microbiol

65:865–877.

20.   Shu Z, Wu J, Chen D, Cheng L, Zheng Y, Chen J, Huang J. 2012. Optimization of Burkholderia sp.

ZYB002 lipase production for pitch control in thermomechanical pulping (TMP) processes.

Holzforschung 66:341–348.

21.   Lo C-F, Yu C-Y, Kuan I-C, Lee S-L. 2012. Optimization of Lipase Production by Burkholderia sp.

Using Response Surface Methodology. Int J Mol Sci 13:14889.

22.   Yudistira H, McClarty L, Bloodworth RA, Hammond SA, Butcher H, Mark BL, Cardona ST. 2011.

Phenylalanine induces *Burkholderia cenocepacia* phenylacetic acid catabolism through degradation

to phenylacetyl-CoA in synthetic cystic fibrosis sputum medium. Microb Pathog 51:183–196.

23.   Tung TT, Quoc Thang N, Cao Huy N, Bao Phuong P, Ngoc Minh D, Hai Nam N, Nielsen J. 2024.

Identification of novel phenylalanine derivatives bearing a hydroxamic acid moiety as potent

quorum sensing inhibitors. RSC Med Chem 15:1320–1328.

24.   Yap ZL, Rahman ASMZ, Hogan AM, Levin DB, Cardona ST. 2024. A CRISPR-Cas-associated

592   transposon system for genome editing in Burkholderia cepacia complex species. Appl Environ

593   Microbiol 90:e0069924.

594   25.   Yap ZL, Motnenko A, Cardona S. 2023. Complete Genome Sequence of Burkholderia

595   vietnamiensis LMG16232. Microbiol Resour Announc 12:e0032823.

596   26.   Baym M, Shaket L, Anzai IA, Adesina O, Barstow B. 2016. Rapid construction of a whole-genome

597   transposon insertion collection for *Shewanella oneidensis* by Knockout Sudoku. Nat Commun 7:1–

598   13.

599   27.   Mohanan N, Montazer Z, Sharma PK, Levin DB. 2020. Microbial and Enzymatic Degradation of

600   Synthetic Plastics. Front Microbiol 11:580709.

601   28.   Gasteiger E, Hoogland C, Gattiker A, Duvaud S, Wilkins MR, Appel RD, Bairoch A. 2005. Protein

602   Identification and Analysis Tools on the ExPASy Server, p. 571–607. *In* Walker, JM (ed.), The

603   Proteomics Protocols Handbook. Humana Press, Totowa, NJ.

604   29.   Mohanan N, Wong CH, Budisa N, Levin DB. 2022. Characterization of Polymer Degrading

605   Lipases, LIP1 and LIP2 From Pseudomonas chlororaphis PA23. Front Bioeng Biotechnol

606   10:854298.

607   30.   Nor Azura Azam et al. 2018. Enhanced degradation of polyhydroxyalkanoates (PHAs) by newly

608   isolated *Burkholderia cepacia* DP1 with high depolymerase activity. 27 Apri 2017 75:11.

609   31.   Azura Azami N, Ira Aryani W, Aik-Hong T, Amirul AA. 2019. Purification and characterization of

610   new bio-plastic degrading enzyme from *Burkholderia cepacia* DP1. Protein Expr Purif 155:35–42.

611   32.   Sánchez DA, Tonetto GM, Ferreira ML. 2018. Burkholderia cepacia lipase: A versatile catalyst in

612   synthesis reactions. Biotechnol Bioeng 115:6–24.

33. Liu C-H, Huang C-C, Wang Y-W, Chang J-S. 2012. Optimizing lipase production from isolated

Burkholderia sp. J Taiwan Inst Chem Eng 43:511–516.

34. Li J, Shen W, Fan G, Li X. 2018. Screening, purification and characterization of lipase from

Burkholderia pyrrocinia B1213. 3 Biotech 8:387.

35. Alnoch RC, Cardoso RLA, Guizelini D, Balsanelli E, Tadra-Sfeir MZ, de Oliveira Pedrosa F,

Sassaki GL, Cruz LM, Mitchell DA, de Souza EM, Krieger N, Muller-Santos M. 2019. Genome

sequencing of Burkholderia contaminans LTEB11 reveals a lipolytic arsenal of biotechnological

interest. Braz J Microbiol 50:619–624.

36. Park O-J, Lee S-H. 2005. Stereoselective lipases from *Burkholderia* sp., cloning and their

application to preparation of methyl (*R*)-*N*-(2,6-dimethylphenyl)alaninate, a key intermediate for

(*R*)-Metalaxyl. J Biotechnol 120:174–182.

37. Lee S-H, Kim M-N. 2010. Isolation of bacteria degrading poly(butylene succinate-*co*-butylene

adipate) and their *lip A* gene. Int Biodeterior Biodegrad 64:184–190.

38. Knapp A, Voget S, Gao R, Zaburannyi N, Krysciak D, Breuer M, Hauer B, Streit WR, Müller R,

Daniel R, Jaeger K-E. 2015. Mutations improving production and secretion of extracellular lipase

by Burkholderia glumae PG1. Appl Microbiol Biotechnol 100:1265.

39. Khattabi ME, Gelder PV, Bitter W, Tommassen J. 2000. Role of the Lipase-specific Foldase of

Burkholderia glumae as a Steric Chaperone. J Biol Chem 275:26885–26891.

40. Wang X, Yu X, Xu Y. 2009. Homologous expression, purification and characterization of a novel

high-alkaline and thermal stable lipase from *Burkholderia cepacia* ATCC 25416. Enzyme Microb

Technol 45:94–102.

41.   Frenken LGJ, de Groot A, Tommassen J, Verrips CT. 1993. Role of the lipB gene product in the

635   folding of the secreted lipase of Pseudomonas glumae. Mol Microbiol 9:591–599.

636   42.   Cuthbertson L, Nodwell JR. 2013. The TetR Family of Regulators. Microbiol Mol Biol Rev

637   77:440–475.

638   43.   Kniewel R, Revelles Lopez O, Prieto MA. 2017. Biogenesis of Medium-Chain-Length

639   Polyhydroxyalkanoates, p. 1–25. *In* Geiger, O (ed.), Biogenesis of Fatty Acids, Lipids and

640   Membranes. Springer International Publishing, Cham.

641   44.   Black PN, DiRusso CC. 1994. Molecular and biochemical analyses of fatty acid transport,

642   metabolism, and gene regulation in *Escherichia coli*. Biochim Biophys Acta 1210:123–145.

643   45.   Jimenez-Diaz L, Caballero A, Segura A. 2017. Pathways for the Degradation of Fatty Acids in

644   Bacteria, p. 1–23. *In* Rojo, F (ed.), Aerobic Utilization of Hydrocarbons, Oils and Lipids. Springer

645   International Publishing, Cham.

646   46.   Cardona ST, Mueller C, Valvano MA. 2006. Identification of essential operons in *Burkholderia*

647   *cenocepacia* with a rhamnose inducible promoter. Appl Environ Microbiol 72:2547–2555.

648   47.   Gallagher LA, Shendure J, Manoil C. 2011. Genome-scale identification of resistance functions in

649   *Pseudomonas aeruginosa* using Tn-seq. mBio 2:e00315-10.

650   48.   Gislason AS, Turner K, Domaratzki M, Cardona ST. 2017. Comparative analysis of the

651   *Burkholderia cenocepacia* K56-2 essential genome reveals cell envelope functions that are uniquely

652   required for survival in species of the genus *Burkholderia*. Microb Genomics 3:e000140.

653   49.   Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. 1.

654   EMBnet.journal 17:10–12.

655   50.   Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler transform.

656   Bioinformatics 26:589–595.

657   51.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,

658   Subgroup 1000 Genome Project Data Processing. 2009. The Sequence Alignment/Map format and

659   SAMtools. Bioinformatics 25:2078–2079.

660   52.   Fu J, Sharma U, Sparling R, Cicek N, Levin DB. 2014. Evaluation of medium-chain-length

661   polyhydroxyalkanoate production by Pseudomonas putida LS46 using biodiesel by-product streams.

662   Can J Microbiol 60:461–468.

663   53.   Braunegg G, Sonnleitner B, Lafferty RM. 1978. A rapid gas chromatographic method for the

664   determination of polyhydroxybutyric acid in microbial biomass. Eur J Appl Microbiol Biotechnol

665   6:29–37.

666   54.   Blunt W, Lagassé A, Harvey J, Sparling R, Gapes D, Levin D, Cicek N. 2024. Semi-continuous

667   non-sterile production of medium chain-length polyhydroxyalkanoates from fatty acids. Can J

668   Chem Eng 102:3896–3908.

669   55.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web portal for

670   protein modeling, prediction and analysis. Nat Protoc 10:845–858.

671   56.   Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, von

672   Heijne G, Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using deep neural

673   networks. Nat Biotechnol 37:420–423.

674   57.   Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R, Butcher S.

675   2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024. Nucleic Acids

676   Res 52:W521–W525.

677   58.   Kim G, Lee S, Levy Karin E, Kim H, Moriwaki Y, Ovchinnikov S, Steinegger M, Mirdita M. 2025.

678   Easy and accurate protein structure prediction using ColabFold. Nat Protoc 20:620–642.

679   59.   Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS, Olson AJ. 2009.

680   AutoDock4 and AutoDockTools4: Automated docking with selective receptor flexibility. J Comput

681   Chem 30:2785–2791.

682   60.   DeLano WL. 2002. Pymol: An open-source molecular graphics tool. CCP4 Newsl Protein

683   Crystallogr.

684   61.   Darling P, Chan M, Cox AD, Sokol PA. 1998. Siderophore production by cystic fibrosis isolates of

685   *Burkholderia cepacia*. Infect Immun 66:874–877.

686   62.   Bloodworth RA, Gislason AS, Cardona ST. 2013. *Burkholderia cenocepacia* conditional growth

687   mutant library created by random promoter replacement of essential genes. MicrobiologyOpen

688   2:243–258.

689   63.   Figurski DH, Helinski DR. 1979. Replication of an origin-containing derivative of plasmid RK2

690   dependent on a plasmid function provided in trans. Proc Natl Acad Sci U S A 76:1648–1652.

Figures and Legends

Fig 1. Overview of transposon mutagenesis and Tn-seq workflow to identify genetic elements

involved in the extracellular mcl-PHA degradation activity in *B. vietnamiensis* LMG 16232. *B.*

vietnamiensis LMG 16232 was subjected to transposon mutagenesis, and 50,000 colonies were

screened for those with absent or reduced extracellular mcl-PHA degradation activity. Colonies

showing reduced or no degradation activity were selected for further analysis. Selected mutants

were grown individually and then grouped into two pools based on their mcl-PHA degradation

activity (Pool 1 for no activity, Pool 2 for weak activity). Genomic DNA (gDNA) was extracted

from each pool. The gDNA was prepared for Tn-Genome junction enrichment. The enriched DNA

fragments were sequenced using Illumina MiSeq. The sequencing data were analyzed by mapping

the reads to the *B. vietnamiensis* LMG 16232 genome.

(A) Type II secretion system

gspN *gspM* *gspL* *gspK* *gspJ* *gspI* *gspH* *gspG* *gspC* *gspF* *gspE* *gspD*

(B) Lipase production

lip1 *lip2* *lipO*

(C) Polyhydroxyalkanoate synthesis

phaR *phaB* *phaA* *phaC*

1 kb

Fig 2. Genetic Organization of the *B. vietnamiensis* LMG 16232 gene clusters that affect the

extracellular mcl-PHA degradation upon interruption by transposon insertion. The triangles

indicate transposon insertions that resulted in the complete loss (Pool 1, purple) or severe reduction

(Pool 2, green) of extracellular mcl-PHA degradation activity in *B. vietnamiensis* LMG 16232. (A)

The type II secretion system of *B. vietnamiensis* LMG 16232 arranges in a gene cluster containing

12 genes (P4G95_05480, P4G95_05485, P4G95_05490, P4G95_05495, P4G95_05500,

P4G95_05505, P4G95_05510, P4G95_05515, P4G95_05520, P4G95_05525, P4G95_05530,

P4G95_05535). All genes of the gene cluster are interrupted except for *gspN*. (B) lipase gene

cluster consists of three genes, *lip1, lip2* and *lipO* (P4G95_16805, P4G95_16810, and

P4G95_16815). Interruption of *lip1* and *lipO* by transposon abolishes *B. vietnamiensis* LMG

16232 ability to degrade extracellular mcl-PHA. (C) The polyhydroxyalkanoate synthesis gene

cluster of *B. vietnamiensis* LMG 16232 consists of four genes, *phaR, phaB, phaA, and phbC*

(P4G95_13560, P4G95_13565, P4G95_13570, and P4G95_13575). Transposon interruption on

phaR and *phaA* severely reduces *B. vietnamiensis* LMG 16232 to degrade extracellular mcl-PHA.

The genes are depicted to scale, as indicated by the bars on the right side of the panels. The putative

function of genes interrupted by transposon is provided in Table 2.

```
WHU94860.1    MVRSMRSRV-------------------------------VAAAVACAMSGAQFAGTTAVM    30
AFQ93688.1    ------------------------------------------------------------     0
AFQ93689.1    ------------------------------------------------------------     0
CAE81078.1    ----MKKLLAGVFGVVAMSLSAQAAKKASNCEVT----GLVDRMTCPYLEKL-VSGP-HL    50
AAA64538.1    --MPLRTLLCGLLLAVCLGQHALA---ASRCSERPRTLLRPAEVSCSYQSTWLDSGL-VG    54

WHU94860.1    TLATTHAATAATAATDDYAATRYPVILVH-GL-TGTDKYA---GVL-EYW--YGIQEDLQ    82
AFQ93688.1    ----VHTF-YH-PRAMGASGERHPVVIW-----------GNGTGAVPGIY--SSLLRHWA    41
AFQ93689.1    ----VHTF-YH-PRSMGASGERHPVVIW-----------GNGTGAVPGIY--SSLLRHWA    41
CAE81078.1    ---TRHVK-YSLPKGKTPKA-GWPTVILYQGSLFPVE-FSRSSLMIAGGYNEIRLIQTLL    104
AAA64538.1    ---QRKII-YQTPLGTPPAG-GWPVVLIYQGSFFPLNDFSYHSNLPFGGYYEGKLVQNLL    109
                       :         .    *.::              .          :    : .

WHU94860.1    RHGATVY----------VANLSGFQSD----DGPNGRGEQLLAYVKTVLAATGATKVNLI    128
AFQ93688.1    SHGFIVAAANTPTSNFALS----------MRAGIDVLERRNADPGSEYFGRVDLARIGSA    91
AFQ93689.1    SHGIIVAAANTPTSNFAIS----------MRAGIDVLERRNADPGSEYFGRVDLEHIGSA    91
CAE81078.1    DSGFAVIAPPAIEGVAWMTNIVGI--DYDTSEDFYFVEELLVAMGNGEFGKLNMDRLYAT    162
AAA64538.1    DHGYAVIAPSAPADLFWQTNIPGLAQAYELSTDYDFLGNVLAAIASGHFGPLNAQRQYAT    169
               *  *              :             .         .     . :.  . :

WHU94860.1    GHSQGGLTSRYVAAVAPDLVASVTTIGTPHRGSEFADFVQSVLAYDPTGLSSSVIATFFN    188
AFQ93688.1    GHSQGGAAAVNAAVDAR-VDTAVPIQ-----PG--------PL-----TDP---------    123
AFQ93689.1    GHSQGGAAAVNAAIDAR-VDTAVPIQ-----PG--------PL-----TDP---------    123
CAE81078.1    GISSGGYHSSRMAVAFPGVFKALAVH-----SASYADCGG-PMCFVPAQVP---------    207
AAA64538.1    GISSGGYNTSRMAVSFPGKFRALAVQ-----SGSYATCSG-PLCVVPDQLP---------    214
              * *.** :   *        ::        .              :

WHU94860.1    VFGILTSSSNNTNQDALAALKTLTTSQAAAYNQNYPSAGLGAPGSCQTGAPTETVGGNTH    248
AFQ93688.1    ------------------------------------------------DLTGVPVFY------    132
AFQ93689.1    ------------------------------------------------DLTDVPMFY------    132
CAE81078.1    ------------------------------------------------ENHPPTIF------    215
AAA64538.1    ------------------------------------------------ADHPPTLF------    222
                                                                    *

WHU94860.1    LLYSWAGTAIQPTVSMFGVTGAKDVSTIPVIDPANALDLSTLALLGTGTVMINRGAGQND    308
AFQ93688.1    ------------------LAGERDLTVWPAL----VKALYRDSD---------------    154
AFQ93689.1    ------------------LAGQRDLTVWPAL----VKALYRDSS---------------    154
CAE81078.1    ------------------LHGRLDPVV-PVR----TMYPYHETL---------------    236
AAA64538.1    ------------------LHGFVDAVV-PWW----SMDLYYDRL---------------    243
                               : *   *   . *

WHU94860.1    GLVSKCSALYGQVLGTRYKWNHLDEINQLLGVRGAYAEDPVAVIRTHANRLKLAGV    364
AFQ93688.1    ----HLPAVYGEVRGAGHLS--------SIGDGGDFRAPT-TA-------------    184
AFQ93689.1    ----HLPAVYGEVRGAGHLS--------SIGDGGDFRAPT-T--------------    183
CAE81078.1    ----KNQG----VETEMFVS--------PWARHEWLEEAPELITNWFINHK-----    271
AAA64538.1    ----LHQG----IETARYTE--------PLGGHEWFAASPGKVLAWFNAHP-----    278
                      .    :    .             .
```

Fig 3. Amino acid sequence alignment of the putative lipase that is able to degrade extracellular

mcl-PHA depolymerase of *B. vietnamiensis* LMG 16232 (protein ID: WHU94860.1) with other

mcl-PHA depolymerase sequences. The sequences analyzed include those from *Streptomyces*

roseolus (GenBank: AFQ93688.1), *Streptomyces venezuelae* (GenBank: AFQ93689.1),

Bdellovibrio bacteriovorus HD100 (GenBank: CAE81078.1)*, Pseudomonas fluorescens*

(GenBank: AAA64538.1). The lipase box consists of the pentapeptide sequence (G-X1-S-X2-G),

which is indicated in the red box. The signal peptide of *B. vietnamiensis* LMG 16232 was predicted

by the SignalP (version 5.0) program (purple box). The main putative catalytic amino acids

Ser131-Asp308-His330 of *B. vietnamiensis* LMG 16232, predicted with Phyre 2, are marked with

a dark blue box.

Fig 4. Targeting genes involved in mcl-PHA degradation in *B. vietnamiensis* LMG 16232 abolish

the strain's ability to degrade mcl-PHA. (A) Schematic model of RNA-guided DNA transposition

by RhaCAST applied to *B. vietnamiensis* LMG 16232. The two plasmids of the RhaCAST system,

pRhaCAST and pDonorTp, encode all the necessary components, tnsB, tnsC, tniQ, Cas12k,

crRNA scaffold, and the cargo gene (TpR) that is flanked by the transposon left end (LE) and right

end (RE). (B) The ability to degrade mcl-PHA was ablated by CAST targeting triacylglycerol

lipase CDS (*lip1*), A24 family peptidase CDS (*A24*), *otsB, gspE,* TetR/AcrR family transcriptional

740   regulator CDS (*tetR/acrR*) in LMG 16232 Wild-type (WT). This figure shows a representative

741   plate of the mcl-PHA degradation assay of three biological replicates in duplicates.

Fig 5. Targeting the putative *lipase* gene (locus tag: P4G95_16805) in *B. vietnamiensis* LMG 16232 abolishes the ability to degrade mcl-PHA. (A) Strains were grown on an agar plate containing Nile red and PHO and visualized under UV irradiation. Inoculation of wild type (WT) results in a clear halo around the strain, whereas the inoculation of the mutant *lip1::CAST-Tp* does not have a halo around the strain. (B) The absence or presence of PHO in the agar after strain inoculation was determined with gas chromatography with flame ionization detection (GC-FID). Samples were obtained by cutting an agar disk under the strain. For control, a disk was cut at an area of the agar where no cell was inoculated. The peak at ~9.3 minutes retention time corresponds to PHO, which can be observed at the no cell control (Top). The absence of a peak corresponds to PHO in WT, and the presence of PHO in *lip1::CAST-Tp* supports the phenotypic test.

Fig 6. Targeting genes involved in mcl-PHA degradation in *B. vietnamiensis* LMG 16232 affects

the ability to degrade tributyrin. A) Representative plate of tributyrin degradation assay; B)

Diameter of the halos degradation in (A). The ability to degrade tributyrin was ablated by CAST

targeting triacylglycerol lipase CDS (*lip1*), A24 family peptidase CDS (*A24*), *otsB, gspE,*

TetR/AcrR family transcriptional regulator CDS (*tetR/acrR*) in LMG 16232 wild-type (WT). Data

is representative of two biological replicates repeated in technical triplicate.

Fig. 7 Docking analysis of Lip1 and Lip2-substrate complexes. A) Docking poses of Lip1 with 3-

Hydroxyoctanoate (OCT). B) Docking poses of Lip1 with Tributyrin (TRI). C) Docking poses of

Lip2 with 3-Hydroxyoctanoate (OCT). D) Docking poses of Lip2 with Tributyrin (TRI). The

images illustrate the single predominant low-energy binding cluster obtained for both substrates,

highlighting the catalytic triad in violet. (E) Binding energy and cluster distribution for protein-

ligand complexes. The plot shows the lowest binding energy for Lip1 and Lip2 docking with 3-

Hydroxyoctanoate (OCT) and Tributyrin (TRI). The size of the circle represents the cluster size

(number of poses), while colours differentiate protein-ligand pairs: Lip1-OCT (blue), Lip1-TRI

(orange), Lip2-OCT (green), and Lip2-TRI (red). Lip1 exhibits a higher affinity for both ligands,

characterized by a single predominant low-energy cluster, whereas Lip2 displays multiple binding

modes for both substrates.

Fig 8. Structural superposition of Lip1 and Lip2 at amino acid at position 73. Superposition of

Lip1 (green) and Lip2 (blue), highlighting the catalytic triad (violet) and the difference in amino

acid in position 73: Lip1 (TYR) and Lip2 (GLN), illustrating its potential impact on enzyme

structure and function.

Monomer ← PHA polymer

Simple diffusion

• Triacylglycerol lipase (P4G95_16805)

Type II secretion system, and signal peptide processing:

OM

• Lipase secretion chaperone (P4G95_16815)

Periplasm

gspD (P4G95_05535)
gspK (P4G95_05495)
gspJ (P4G95_05500)
gspI (P4G95_05505)
gspH (P4G95_05510)
gspC (P4G95_05520)
gspM (P4G95_05485)
gspG (P4G95_05515)
gspF (P4G95_05525)
gspL (P4G95_05490)

IM

Cytosol

Sec

gspE (P4G95_05530)

• A24 family peptidase (P4G95_07805)

+/− ?

Transcriptional regulator:
• TetR/AcrR family transcriptional regulator (P4G95_00865)
• MarR family winged helix-turn-helix transcriptional regulator (P4G95_05570)
• Polyhydroxyalkanoate synthesis repressor PhaR (P4G95_13560)
• Helix-turn-helix domain-containing protein (P4G95_16325)

Unknown relationship with extracellular mcl-PHA depolymerase activity
• Trehalose-phosphatase otsB (P4G95_10885)
• N-acetylglucosamine-6-phosphate deacetylase NagA (P4G95_03530)
• Phenylacetate-CoA oxygenase/reductase subunit PaaK and PaaJ (P4G95_06510 and P4G95_06515)
• Acetyl-CoA acetyltransferase (P4G95_13570)
• Hypothetical proteins (P4G95_20805 and P4G95_20810)

Fig 9. Proposed model of components involved in *B. vietnamiensis* LMG 16232 extracellular mcl-PHA degradation activity. The elements in this model were deduced from the transconjugants that either lost or showed severely reduced extracellular mcl-PHA depolymerase activity due to transposon insertion. The components where the text is in red have been verified via the creation of insertional mutagenesis using RhaCAST. The regulatory elements influencing this activity are TetR/AcrR family transcriptional regulator (P4G95_00865), Polyhydroxyalkanoate synthesis repressor (P4G95_13560) and HTH-type regulators (P4G95_05570 and P4G95_16325). The Sec system is responsible for transporting the unfolded protein into the periplasmic, as the protein is predicted to contain a sec tag. The putative lipase chaperone (P4G95_16815) is responsible for folding the putative lipase into a mature protein and excreted to the outside of the cell via the type II secretion system. *B. vietnamiensis* LMG 16232 extracellular mcl-PHA degradation activity is also affected by the transposon interruption in genetic elements related to polyhydroxyalkanoate production and several genes involved in metabolism.

Table 1. *Burkholderia* isolates that exhibit extracellular mcl-PHO degradation activity

Species	Strain	Ability to degrade PHO after 20 days	
		30 ˚C	37 ˚C
Burkholderia cenocepacia	K56-2	-	-
	F9141525	-	-
	F3071909	-	-
	G1181568-1	-	-
	F4200989-3	-	-
	VC14488	-	-
	VC14543	-	-
	VC10414	-	-
	VC14761	-	-
	130723	-	-
	125007	-	-
Burkholderia contaminans	21NG595759	-	-
	21NG609698	-	-
Burkholderia dolosa	AU6940	-	-
	AU2167	-	-
	LM018941	-	-
	AU0645	-	-
Burkholderia gladioli	VC19233	+	+
	VC14812	-	-
	132208	+	-
	113611	-	-
Burkholderia multivorans	21NH942533	+	+
	21NJ503211	+	-
	V1120157	-	-
	LM013010	-	-
	F9091687	+	-
	VC19694	-	-
	VC15555	-	-
	VC9825	+	+
	131077	-	-
Burkholderia stagnalis	MSMB618	++	++
	MSMB642	++	++
Burkholderia vietnamiensis	LMG 22486	-	-
	LMG 10929	-	-
	LMG 16232	++	++
	VC11431	-	-
	VC18984	-	-

794 mcl-PHA degradation was ranked based on (–) no clearing, (+) weak clearing, (++) moderate
795 clearing. Fig. S1 is the source of the data interpretation.

Table 2. Transposon mutant of *B. vietnamiensis* LMG 16232 with abolished (Pool 1) or reduced (Pool 2) extracellular mcl-PHA activity

Chromosome location	Locus tag	Gene name	Putative function
Pool 1			
1	P4G95_00865	TetR/AcrR family transcriptional regulator CDS	TetR/AcrR family transcriptional regulator
1	P4G95_05485	gspM	type II secretion system protein M
1	P4G95_05490	gspL	type II secretion system protein GspL
1	P4G95_05495	gspK	type II secretion system minor pseudopilin GspK
1	P4G95_05500	gspJ/prepilin-type N-terminal cleavage/methylation domain-containing protein CDS	prepilin-type N-terminal cleavage/methylation domain-containing protein
1	P4G95_05505	gspI	type II secretion system minor pseudopilin GspI
1	P4G95_05510	GspH/FimT family pseudopilin CDS	GspH/FimT family pseudopilin
1	P4G95_05515	gspG	type II secretion system major pseudopilin GspG
1	P4G95_05520	general secretion pathway protein GspC CDS	general secretion pathway protein GspC CDS
1	P4G95_05525	gspF	type II secretion system inner membrane protein GspF
1	P4G95_05530	gspE	type II secretion system ATPase GspE
1	P4G95_05535	gspD	type II secretion system secretin GspD
1	P4G95_07805	A24 family peptidase CDS	A24 family peptidase
1	P4G95_10885	otsB CDS	trehalose-phosphatase
2	P4G95_16805	lip1/triacylglycerol lipase CDS	triacylglycerol lipase
2	P4G95_16815	lipO/lipase secretion chaperone CDS	lipase secretion chaperone
2	P4G95_20805	hypothetical protein CDS	hypothetical protein
3	P4G95_28980	gspD CDS	type II secretion system secretion GspD
Pool 2			
1	P4G95_03530	nagA	N-acetylglucosamine-6-phosphate deacetylase
1	P4G95_05495	gspK	type II secretion system minor pseudopilin GspK
1	P4G95_05505	gspI	type II secretion system minor pseudopilin GspI

1	Between P4G95_05555 and P4G95_05560	Between mnmC and cation:proton antiporter CDS	cation:proton antiporter
1	P4G95_05570	MarR family winged helix-turn-helix transcriptional regulator CDS	MarR family winged helix-turn-helix transcriptional regulator
1	P4G95_06510 and P4G95_06515	paaK and paaJ CDS	phenylacetate-CoA oxygenase/reductase subunit PaaK and phenylacetate-CoA oxygenase subunit PaaJ
1	P4G95_13560	phaR	polyhydroxyalkanoate synthesis repressor PhaR
1	P4G95_13570	acetyl-CoA C-acetyltransferase CDS	acetyl-CoA C-acetyltransferase
1	Between P4G95_15285 and P4G95_15290	Between cupin domain-containing protein CDS and hypothetical protein	
2	P4G95_16325	helix-turn-helix domain-containing protein CDS	helix-turn-helix domain-containing protein
2	P4G95_20810	hypothetical protein CDS	hypothetical protein
3	Between P4G95_29770 and P4G95_29775	Between porin CDS and helix-turn-helix domain-containing protein CDS	

798   **Table 3.** List of bacterial strains and plasmids used in this work.

Strain or plasmid	Features	Source
Burkholderia cenocepacia K56-2	ET12 lineage cystic fibrosis clinical isolate from Toronto	(61)
Burkholderia cenocepacia F9141525	Cystic fibrosis clinical isolate from Toronto	SickKids, Toronto
Burkholderia cenocepacia F3071909	Cystic fibrosis clinical isolate from Toronto	SickKids, Toronto
Burkholderia cenocepacia G1181568-1	Cystic fibrosis clinical isolate from Toronto	SickKids, Toronto
Burkholderia cenocepacia F4200989-3	Cystic fibrosis clinical isolate from Toronto	SickKids, Toronto
Burkholderia cenocepacia VC14488	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia cenocepacia VC14543	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia cenocepacia VC10414	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia cenocepacia VC14761	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia cenocepacia 130723	Clinical respiratory isolated collected in Halifax	CANWARD/CARA database
Burkholderia cenocepacia 125007	Clinical respiratory isolated collected in Vancouver	CANWARD/CARA database
Burkholderia contaminans 21NG595759	Canadian cystic fibrosis isolate	H. Adam (Shared Health MB)
Burkholderia contaminans 21NG609698	Canadian cystic fibrosis isolate	H. Adam (Shared Health MB)
Burkholderia dolosa AU6940	Cystic fibrosis clinical isolate	SickKids, Toronto
Burkholderia dolosa AU2167	Cystic fibrosis clinical isolate	SickKids, Toronto
Burkholderia dolosa LM018941	Cystic fibrosis clinical isolate	SickKids, Toronto
Burkholderia dolosa AU0645	Cystic fibrosis clinical isolate	SickKids, Toronto
Burkholderia gladioli VC19233	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia gladioli VC14812	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia gladioli 132208	Clinical respiratory isolated collected in Montreal	CANWARD/CARA database

Burkholderia gladioli 113611	Clinical respiratory isolated collected in Saskatoon	CANWARD/CARA database
Burkholderia multivorans 21NH942533	Canadian cystic fibrosis isolate	H. Adam (Shared Health MB)
Burkholderia multivorans 21NJ503211	Canadian cystic fibrosis isolate	H. Adam (Shared Health MB)
Burkholderia multivorans V1120157	Cystic fibrosis clinical isolate	SickKids, Toronto
Burkholderia multivorans LM013010	Cystic fibrosis clinical isolate	SickKids, Toronto
Burkholderia multivorans F9091687	Cystic fibrosis clinical isolate	SickKids, Toronto
Burkholderia multivorans VC19694	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia multivorans VC15555	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia multivorans VC9825	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia multivorans 131077	Clinical respiratory isolated collected in Toronto	CANWARD/CARA database
Burkholderia stagnalis MSMB618	N/A	NAU-PMI
Burkholderia stagnalis MSMB642	N/A	NAU-PMI
Burkholderia vietnamiensis LMG 22486	Wastewater isolate	BCCM
Burkholderia vietnamiensis LMG 10929	Oryza sativa, rhizosphere soil isolate	BCCM
Burkholderia vietnamiensis LMG 16232	Clinical isolate from cystic fibrosis patient	BCCM
Burkholderia vietnamiensis VC11431	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia vietnamiensis VC18984	Canadian cystic fibrosis isolate	CBCCRRR
Burkholderia vietnamiensis LMG 16232 *lip1*::CAST-Tp	Derived from LMG 16232; Cargo containing Tp^r integrated into triacylglycerol lipase CDS (P4G95_16805)	This study
Burkholderia vietnamiensis LMG 16232 A24 family peptidase CDS:: CAST-Tp	Derived from LMG 16232; Cargo containing Tp^r integrated into A24 family peptidase CDS (P4G95_07805)	This study
Burkholderia vietnamiensis LMG 16232 *otsB*::CAST-Tp	Derived from LMG 16232; Cargo containing Tp^r integrated into *otsB* (P4G95_10885)	This study

Burkholderia vietnamiensis LMG 16232 *gspE*::CAST-Tp	Derived from LMG 16232; Cargo containing Tp^r integrated into *gspE* (P4G95_05530)	This study
Burkholderia vietnamiensis LMG 16232 TetR/AcrR family transcriptional regulator CDS::CAST-Tp	Derived from LMG 16232; Cargo containing Tp^r integrated into TetR/AcrR family transcriptional regulator CDS (P4G95_00865)	This study
E. coli Stbl2	*F- mcrA Δ(mcrBC-hsdRMS-mrr) recA1 endA1lon gyrA96 thi supE44 relA1 λ- Δ(lac-proAB)*	Invitrogen
pRBrhaBoutgfp	pSCrhaBoutgfp derivative, ori$_{R6K}$, *rhaR rhaS* PrhaB e-gfp	(62)
pRK2013	ori$_{colE1}$ RK2 derivative *Kan^r mob$^+$ tra$^+$*	(63)
pDonorTp	pDonor_ShCAST_kanR with modification of Km^R to Tp^R	(24)
pRhaCAST	*Sh-cas12K-tnsBC-tniQ* drives by the rhamnose inducible promoter	(24)
pRhaCAST-3668	Modified from pRhaCAST. crRNA targeting *B. vietnamiensis* LMG 16232 triacylglycerol lipase CDS (P4G95_16805)	This study
pRhaCAST-3952	Modified from pRhaCAST. crRNA targeting *B. vietnamiensis* LMG 16232 A24 family peptidase CDS (P4G95_07805)	This study
pRhaCAST-3956	Modified from pRhaCAST. crRNA targeting *B. vietnamiensis* LMG 16232 *otsB* (P4G95_10885)	This study
pRhaCAST-3996	Modified from pRhaCAST. crRNA targeting *B. vietnamiensis* LMG 16232 *gspE* (P4G95_05530)	This study
pRhaCAST-4000	Modified from pRhaCAST. crRNA targeting *B. vietnamiensis* LMG 16232 TetR/AcrR family transcriptional regulator CDS (P4G95_00865)	This study

800 **Table 4.** Primers/templates used in this study

Number	Sequence	Description
Oligonucleotides for Tn-seq		
683	CCGTAGTGAGTTCTTCGTCCGAGCCACTCGGAGATGTGTATAAGAGACAGT	Tn-seq adapter (top strand)
684	CTGTCTCTTATACACATCTCCGAGTGGCTCGGACGAAGAACTCACTACGG	Tn-seq adapter (bottom strand)

682	CACAAGTGCGGCCGCACTAGTCTAG ATTTAAATTACCGTAGTGAGTTCTTC GTCCGAGCCAC	Tn-seq oligo probe
681	CAAGCAGAAGACGGCATACGAGATT CGCCTTAGTCTCGTGGGCTCGGAGAT GTGTATAAGAGACAG	Illumina Nextera index N701 (Pool 1)
690	AATGATACGGCGACCACCGAGATCT ACACTAGATCGCTCGTCGGCAGCGTC AGATGTGTATAAGAGACAGNNNNNAA TCTAGACTAGTGCGGCC	Illumina Nextera index N501 (Pool 1)
718	CAAGCAGAAGACGGCATACGAGATA GGAGTCCGTCTCGTGGGCTCGGAGAT GTGTATAAGAGACAG	Illumina Nextera index N705 (Pool 2)
719	AATGATACGGCGACCACCGAGATCT ACACCTCTCTATTCGTCGGCAGCGTC AGATGTGTATAAGAGACAGNNNNNAA TCTAGACTAGTGCGGCC	Illumina Nextera index N502 (Pool 2)

Oligonucleotides for cloning crRNA into pRhaCAST (crRNA is in bold, *sapI* restriction site is in italics)		
3392	ACTAACGGCTCTTCGAG	Reverse primer to amplify single stranded nucleotide template containing the crRNA
3421	CATTGGCTCTTCGAAAG	Forward primer to amplify single stranded nucleotide template containing the crRNA
3668	CATT*GGCTCTTC*GAAAG**ATCCTCGTG CACGGCCTTACCGGC**GGCTC*GAAGA GC*CGTTAGT	Targeting triacylglycerol lipase CDS (P4G95_16805) of LMG 16232
3952	CATT*GGCTCTTC*GAAAG**GTGATCGG TCTGTGCGTCGGCAGC**GGCTC*GAAG AGC*CGTTAGT	Targeting A24 family peptidase CDS (P4G95_07805) of LMG 16232
3956	CATT*GGCTCTTC*GAAAG**CCTGACCG ATACCGCGTTCTTTTT**GGCTC*GAAG AGC*CGTTAGT	Targeting *otsB* (P4G95_10885) of LMG 16232
3996	CATT*GGCTCTTC*GAAAG**TGCAAGGA AGAGCGACACGAGGAC**GGCTC*GAA GAGC*CGTTAGT	Targeting *gspE* (P4G95_05530) of LMG 16232
4000	CATT*GGCTCTTC*GAAAG**CCACCATCG ACGAGACCTGGTCGT**GGCTC*GAAGA GC*CGTTAGT	Targeting TetR/AcrR family transcriptional regulator CDS (P4G95_00865) of LMG 16232

Oligonucleotides for colony PCR to verify genomic insertion		
3001	GGTGGGTTGAAAGCAAGTCC	left end of the cargo insertion
3670	CTATCCGGTCATCCTCGTGC	Colony PCR for triacylglycerol lipase CDS (P4G95_16805) of LMG 16232
3953	GTGATCGGTCTGTGCGTCG	Colony PCR for A24 family peptidase CDS (P4G95_07805) of LMG 16232
3957	CTGACCGATACCGCGTTCTT	Colony PCR for *otsB* (P4G95_10885) of LMG 16232

3997	GCAAGGAAGAGCGACACGAG	Colony PCR for *gspE* (P4G95_05530) of LMG 16232
4001	CCACCATCGACGAGACCTG	Colony PCR for TetR/AcrR family transcriptional regulator CDS (P4G95_00865) of LMG 16232

801