PIWIL2 downregulation in colon cancer promotes transposon activity and pro-tumorigenic phenotypes

Authors:

Alyssa Risner[1], Joyce Nair-Menon[1], Abhinav Cheedipudi[2], Joe R Delaney[3], Vamsi Gangaraju[3], Antonis Kourtidis[1]

Affiliations

[1]Department of Regenerative Medicine and Cell Biology, Medical University South Carolina, Charleston, SC

[2]University of South Carolina, Columbia, SC.

[3]Department of Biochemistry and Molecular Biology, Medical University South Carolina, Charleston, SC

Corresponding author: Antonis Kourtidis, kourtidi@musc.edu

ABSTRACT

Reactivation of transposable elements (TEs) in somatic tissues, particularly of LINE-1, is associated with disease by causing gene mutations and DNA damage. Previous work has shown that the PIWI pathway is crucial for TE suppression in the germline. However, the status and function of this pathway is not well characterized in differentiated somatic cells and there is lack of consensus on the role of the pathway in somatic tumorigenesis. To shed light on this conundrum, we examined the PIWI pathway in colon cancer through combining bioinformatic analyses and cell-based assays. Shifted Weighted Annotation Network (SWAN) analysis revealed that the pathway experiences significant allelic losses in colon cancer and that PIWIL2, the main catalytic component of the pathway responsible for TE silencing, experiences the highest percent deletions. PIWIL2 is downregulated in colon tumors of advanced stage, nodal metastasis, and in certain subtypes, correlating with poor survival, while it is also downregulated in ulcerative colitis, an inflammatory bowel disease that predisposes to colon cancer. PIWIL2 depletion in colon epithelial Caco2 cells leads to increased anchorage-independent growth, loss of transposon-targeting - non-canonical - piRNAs, increased LINE-1 levels and activity, and in DNA damage, altogether highlighting a tumor-suppressing role of PIWIL2 in the colon.

SUMMARY STATEMENT

This study investigates the PIWI-piRNA pathway in colon cancer using a nuanced bioinformatic and cell-based approach, linking the downregulation of PIWIL2 to disease progression, transposable element activation and DNA damage.

INTRODUCTION

Transposons, or transposable elements (TEs), are mobile DNA elements that make up approximately 45% of the human genome. Although they have been primarily studied in the germline and are overall considered inactive in differentiated somatic epithelial tissues, their reactivation has been shown to lead to a variety of diseases through the introduction of regulatory or protein coding changes when integrated into a new site (Beck et al., 2011, Elbarbary et al., 2016, Hancks and Kazazian, 2016). Additionally, recent studies have demonstrated that genomic instability, oncogene expression, and high mutation rates in approximately 50% of tumors are linked to increased TE activity (Tubio et al., 2014, Jang et al., 2019).

Among the several TE families existing in the human genome, the Long Interspersed Nuclear Element-1 (LINE-1) is the most abundant and the only autonomously mobilizing TE family in humans(Beck et al., 2011). While most copies of the LINE-1 TE are inactive due to truncations and mutations, 100-150 copies are functionally active in disease(Wang et al., 2024, Beck et al., 2010, Tubio et al., 2014, Rodriguez-Martin et al., 2020). The functional LINE-1 copies contain protein-coding open reading frames (ORFs), ORF1 and ORF2, that encode an RNA-binding protein and reverse transcriptase, respectively, to facilitate reverse transcription of the LINE1 RNA transcript(Beck et al., 2011, Elbarbary et al., 2016, Hancks and Kazazian, 2016). Reactivation of LINE-1 has been frequently observed in somatic tissues and particularly in the gastrointestinal tract, where it has been associated with colon cancer, the second most frequent and third deadliest cancer type in both sexes(Pitkanen et al., 2014, Solyom et al., 2012, Ewing et al., 2015, Tubio et al., 2014, Zhuo et al., 2015, Nam et al., 2023, Siegel et al., 2024). Indeed, recurrent LINE-1 insertions have been identified to cause mutations in known oncogenes or tumor suppressors such as APC, a key initiating event in colorectal tumorigenesis, indicating that LINE-1 retro-transposition events have the ability to serve as tumor-initiating events in colon cancer(Cajuso et al., 2019, Scott et al., 2016, Miki et al., 1992). In fact, colon tumors are significantly enriched with somatic retro-transpositions(Rodriguez-Martin et al., 2020). However, the reasons for increased TE activity in these tumors are still poorly understood.

Mechanisms of TE regulation previously described in the germline include DNA methylation and histone modifications of LINE-1 loci, as well as RNA interference by small RNAs through the PIWI pathway(Kuramochi-Miyagawa et al., 2008, Sigurdsson et al., 2012, Ariumi, 2016, Berrens et al., 2017, Walter et al., 2016). The PIWI pathway was first described in *Drosophila* to suppress TEs in the germline and to be required for spermatogenesis(Ku and Lin, 2014). PIWI proteins are members of the Argonaute family of endoribonucleases that bind PIWI interacting RNAs (piRNAs), a distinct class of small non-coding RNAs that are 24-32 nucleotides in length, altogether forming the piRNA-induced silencing complex (piRISC)(Siomi et al., 2011). Most piRNAs are derived from piRNA clusters within the genome,

which are also locations of a large number of truncated TEs (Siomi et al., 2011). piRISC recognizes piRNA - complementary RNA transcripts of TEs in their cytosolic phase and cleaves them, subsequently preventing retro-transposition and DNA mutagenesis(Ramat and Simonelig, 2021, Watanabe and Lin, 2014, Weick and Miska, 2014).

The PIWI-piRNA mechanism is conserved throughout evolution, including in mammals, where it has been found to be required for germline maintenance in mice(Unhavaithaya et al., 2009, Sun et al., 2022, Kuramochi-Miyagawa et al., 2008, Li et al., 2021). In humans, there are four PIWI protein members, namely PIWIL1, PIWIL2, PIWIL3 and PIWIL4(Zeng et al., 2020). In addition to PIWI proteins, the TE targeting function of piRISC is enabled by a number of additional protein components, including the Tudor family of proteins that act as protein scaffolds, as well as by RNA helicases, such as MOV10L1 and DDX4(Siomi et al., 2010, Mathioudakis et al., 2012, Reuter et al., 2009, Gao et al., 2024, Frost et al., 2010, Vourekas et al., 2015, Fu et al., 2019, Li et al., 2021, Wenda et al., 2017). Although it was previously thought that PIWI proteins were germline - specific in humans, a recent study that sought to characterize their expression patterns across multiple tissue types showed that PIWI proteins, namely PIWIL2 and PIWIL4, are also expressed substantially in somatic tissues(Meseure et al., 2020).

In addition to their expression in normal somatic tissues, a number of studies have suggested that PIWI proteins are overexpressed in somatic tumors and act as oncogenes, in a similar fashion to the overall scheme of re-expression of stem cell and germline markers in tumors; however, other studies have shown that PIWI proteins in fact act as tumor suppressors, altogether exposing a lack of consensus on the function of PIWI proteins in somatic tissues and tumors (Lee et al., 2006, Lu et al., 2012, Wang et al., 2015a, Zeng et al., 2017, Eckstein et al., 2018, Meseure et al., 2020, Kishani Farahani et al., 2023, Shi et al., 2020). This is particularly the case for PIWIL2, which is the key endoribonuclease that binds primary piRNAs in piRISC to target TE RNA transcripts for degradation, including LINE-1(Siomi and Siomi, 2015, De Fazio et al., 2011). Still, the function of PIWIL2 in differentiated epithelial tissues and its role in tumorigenesis has not been well characterized(Yan et al., 2011, Peng and Lin, 2013, Martinez et al., 2015, Jacobs et al., 2016, Gao, 2008, He et al., 2010, Liu et al., 2010, Ross et al., 2014). Previous *in vitro* work studying PIWIL2 has been primarily conducted in transformed cancer cell lines, or in commonly used cell lines such as HEK-293 cells, none of which are representative of the normal differentiated epithelium. Here, we aimed to elucidate the expression and function of the PIWI complex in colon cancer, through performing bioinformatic analyses and functional assays in colon epithelial cells.

RESULTS

SWAN analysis identifies dysregulation of the PIWI pathway in colon cancer

We first confirmed high expression levels of PIWIL2 in normal tissues, particularly of gastrointestinal origin, including colon and rectal, through querying PIWIL2 expression levels both through NCBI and through The Cancer Genome Atlas (TCGA), using the University of Alabama at Birmingham CANcer data analysis portal (UALCAN) (**Fig. S1**). Then, to interrogate the status and potential role of the PIWI-piRNA pathway in colon cancer, we initially used the Shifted Weighted Annotation Network (SWAN) algorithm, which has the capability to interrogate pathway and copy-number alterations (CNAs) within the TCGA data set(Bowers et al., 2022). Networks in the algorithm are scored based on the number of interactions within the pathway and if haploinsufficiency data are available for the dataset. The Single-pathway SWAN algorithm was used to perform Gene Ontology Biological Process Analysis on the PIWI-piRNA metabolic pathway in the TCGA colon adenocarcinoma (COAD) dataset (**Fig. 1**). The pathway was scored by the SWAN algorithm as overall haploinsufficient **(Fig. 1A)**. Allelic losses or gains were then mapped to each respective chromosome by the algorithm (**Fig. 1B**). This analysis showed that there is not a single "hot spot" across the genome where all the observed allelic losses or gains are concentrated, but that these are dispersed throughout the genome. Notably, PIWIL2, which is the main catalytic component of the primary piRNA pathway involved in TE regulation, experienced the highest percent deletion and allelic losses of the pathway queried (**Fig. 1C**). PIWIL1 and PIWIL4 were not significantly enriched in the SWAN algorithm with gains or losses, and PIWIL3 was excluded from the pathway by the algorithm. Proteins required for both primary and secondary piRNA processing experienced allelic losses, including the nuclease EXD1; RNA helicase DDX4 (Vasa in *Drosophila*; MVH in mice); PLD6 (Zucchini in *Drosophila*; mitoPLD in mice) that generates the 5' end of primary piRNAs; RNA helicase MOV10L1 that is required for stabilizing the primary piRNA transcript; as well as TDRD1 and TDRD9 that are required for stabilization of the complex and piRNA loading(Yang et al., 2016, Wenda et al., 2017, Nishimasu et al., 2012, Han et al., 2015a, Vourekas et al., 2015, Zhang et al., 2019, Mathioudakis et al., 2012). Interestingly, four components of the pathway experienced high percentages of allelic amplifications and allelic gains in the COAD dataset, including TDRD12, TDRKH, ASZ1 and FKBP6. TDRD12 is a binding partner of EXD1 while TDRKH is required for 3'end processing of piRNAs(Pandey et al., 2013, Saxe et al., 2013). ASZ1 is required for phased piRNA biogenesis through the stabilization of PIWIL2 and PIWIL4, while FKBP6 is a co-chaperone to facilitate piRNA loading (Ma et al., 2009, Ikeda et al., 2022, Saxe et al., 2013, Xiol et al., 2012). Furthermore, we queried whether PIWIL2 allelic loss correlates with patient survival using the cBioportal copy number alteration (CNA) tool; indeed, allelic loss of PIWIL2 correlates with poorer patient survival compared to patients without an allelic loss (**Fig. 1D**). The results from the SWAN and cBioportal analyses highlight the overall loss of regulation of the entire piRNA biogenesis pathway in colon cancer and identify PIWIL2 as a top candidate for the loss of the pathway's regulation.

PIWIL2 is downregulated in colon cancer in correlation with disease progression

Then, to examine whether PIWIL2's allelic losses are also accompanied with downregulation of PIWIL2 in colon cancer, we interrogated the COAD (colon adenocarcinoma) dataset of TCGA using UALCAN, to examine PIWIL2 mRNA expression stratified by normal vs: primary tumor; tumor stages; nodal metastasis; p53 status; histological subtype (adenocarcinoma, mucinous adenocarcinoma); microsatellite stability status; patient age; weight; race; and biological sex (**Fig. 2A-J**). Comparisons between normal and primary tumors showed decreased PIWIL2 levels in tumors, as well as when tumors were stratified by stage I-IV (**Fig. 2A,B**). Due to high variance in expression, this downregulation of PIWIL2 only reached statistical significance in the transition to stage IV (**Fig. 2B**). We also did not find any significant downregulation of PIWIL2 in adenomas through interrogating Gene Expression Omnibus (GEO) repository dataset profiles (**Fig. S2**). However, downregulation of PIWIL2 in tumors was significant when the tumors were stratified by lymph node metastasis, particularly in N0 tumors compared to normal and between N0 and N2 tumors, denoting correlation of PIWIL2 both with early transformation at N0 and with tumor aggressiveness at N2 (**Fig. 2C**). Interestingly, PIWIL2 downregulation does not correlate with p53 status (**Fig. 2D**); however, PIWIL2 was significantly downregulated in adenocarcinomas compared to normal tissues, and even more downregulated in mucinous adenocarcinomas (**Fig. 2E**). Mucinous adenocarcinomas possess a high degree of microsatellite instability, which is a feature that is used to categorize colon tumors in MSI-H (microsatellite instability - high), MSI-L (microsatellite instability - low), and MSS (microsatellite stable). Since the UALCAN database does not include the option of stratifying tumors based on microsatellite instability, we used GEPIA2(Tang et al., 2019) to examine expression of PIWIL2 in these sub-categories of tumors. Interestingly, PIWIL2 is downregulated in MSI-L tumors and even more significantly in MSI-H tumors, compared to MSS tumors (**Fig. 2F**). This result is in agreement with the downregulation of PIWIL2 in the mucinous adenocarcinomas, which are microsatellite unstable. PIWIL2 also seems to be progressively downregulated by age, where MSI-H tumors are also more prevalent (Guidoboni et al., 2001) (**Fig. 2G**). Finally, there were no statistically significant differences in PIWIL2 expression by weight, race, or biological sex (**Fig. 2H,I,J**). Low PIWIL2 expression in colon tumors was also correlated with poor patient survival among all tumor stages, when stratified by stages 1-3 combined, and even more pronounced in stage 4, using the Kaplan-Meier Plotter web-based tool (**Fig. 2K-M**), which is in agreement with the CNA data (**Fig. 1D**).

Furthermore, examination of the methylation levels of the PIWIL2 promoter using UALCAN showed that this was significantly elevated in colon tumors overall, indicated by a higher methylation beta-value, as well as when these tumors were stratified by stage, nodal metastasis; p53 status; histological

subtype; age: biological sex; weight; and race (**Fig. 3**). These findings indicate that PIWIL2 expression is suppressed across colon cancer possibly due to promoter methylation and associates with microsatellite instability.

PIWIL2 is also downregulated in ulcerative colitis

Since we identified downregulation of PIWIL2 in colon tumors beginning from early forms of the disease, such as in N0 tumors, we then sought to examine the status of PIWIL2 in precancerous conditions in the colon, such as in inflammatory bowel disease that significantly increases the risk for colon cancer(Kim and Chang, 2014, Gordon et al., 2018). To do this, we examined the expression levels of members of the PIWI pathway that we identified using the SWAN analysis, using expression data from patients with ulcerative colitis, one of the main manifestations of inflammatory bowel disease, available through the PreMedIBD database(Linggi et al., 2021). This analysis showed that out of all the members of the PIWI-piRNA pathway that were queried, PIWIL2 was the gene that was more predominantly downregulated in ulcerative colitis (**Fig. 4**). This further denotes PIWIL2 as the member of the pathway that is potentially most critical for tumor initiation. Altogether, the above analyses demonstrate that the PIWI pathway is dysregulated in colon cancer, particularly its main component PIWIL2, and its downregulation correlates with disease initiation and progression.

PIWIL2 depletion in colon epithelial cells promotes pro-tumorigenic behavior

Given that PIWIL2 exhibits the most allelic losses in the entire PIWI pathway and is also downregulated in ulcerative colitis and colon cancer correlating with disease progression (**Fig. 1C, Fig. 2, Fig. 3, Fig. 4**), we sought to investigate whether loss of PIWIL2 may indeed promote pro-tumorigenic phenotypes. To do this, we used Caco2 colon epithelial cells, a broadly used in vitro model of the well-differentiated colonic epithelium(Grasset et al., 1984, Hidalgo et al., 1989, Sambuy et al., 2005). We first confirmed that PIWI proteins were expressed in these cells, by examining our publicly available RNA-sequencing data(Daulagala et al., 2024). The data confirmed expression of PIWIL2 and PIWIL4 in Caco2 cells, but not of PIWIL1 and PIWIL3 (**Table S1**). We then generated a PIWIL2 CRISPR/Cas9 mediated knockout (KO) in Caco2 cells (**Fig. S3**). The PIWIL2 KO was overall not well tolerated by these cells, since most PIWIL2 CRISPR/Cas9 – transfected cells did not survive selection; however, we isolated a clone (**Fig. S3**) where we confirmed 50% downregulation of PIWIL2 by qRT-PCR (**Fig. 5A**) and by western blot, when assessing the full length PIWIL2 at the predicted molecular weight of 109 kDa and when quantified over 3 biological replicates (**Fig. 5B, C**). We confirmed the specificity of the loss of the full length PIWIL2 through introducing an LZRS-based retroviral expression vector with full length PIWIL2 FLAG-tagged. (**Fig. 5D**).

We then sought to assess the potential phenotypic effects of PIWIL2 depletion in these cells on cell proliferation, since increased proliferation is an indication of pro-tumorigenic behavior. To do this, we used the xCELLigence Real-Time Cell Analysis (RTCA) system, to monitor cell proliferation rates in standard 2-dimensional conditions and in real time (**Fig. 5E**). Interestingly, the PIWIL2 KO cells had a significantly lower proliferation rate indicated by a slower growth curve, than the Caco2 WT control cells. Although we observed a similarly slower proliferation rate by partially depleting PIWIL2 using a shRNA construct (**Fig. S4**), introduction of the PIWIL2-FLAG construct in the PIWIL2 KO cells not only did not rescue the proliferation decrease, but it actually exacerbated it (**Fig. 5E**). This suggests that although the full length PIWIL2 is responsible for proliferation suppression, there may be other isoforms of PIWIL2 that act opposingly in promoting proliferation.

We then examined the ability of the PIWIL2 KO cells to grow in anchorage-independent conditions, which is a hallmark of pro-tumorigenic transformation. To do this, we cultured cells using a low attachment plate assay to determine if the PIWIL2 KO cells could form spheroids in these conditions (**Fig. 5F-J**). Caco2 WT cells were able to form only small and very few spheroids after 7 days, which is expected since these cells are well-differentiated epithelial cells and not expected to grow in anchorage-independent conditions(Kourtidis et al., 2015). However, the PIWIL2 KO cells grew into much larger spheroids that exhibited a significantly larger area, perimeter and diameter, and were relatively symmetrical, when considering there were no significant changes in circularity. Introduction of the PIWIL2-FLAG construct in the PIWIL2 KO cells reversed the increased anchorage-independent growth and all the observed spheroid phenotypes, demonstrating the specificity of the results (**Fig 5. F-J**). These data demonstrate that the full-length PIWIL2 suppresses proliferation and cell growth in an anchorage independent manner, indicating a tumor-suppressing role of PIWIL2.

PIWIL2 depletion results in loss of transposon-targeting piRNAs and in upregulation of the LINE-1 transposable element

Since PIWIL2 is the main enzymatic component of the PIWI pathway responsible for TE targeting through piRNAs, we then investigated whether PIWIL2 depletion would indeed affect TE targeting. To do this, we first performed RNA sequencing of Caco2 WT and PIWIL2 KO cells for short RNAs to identify any effects on piRNAs upon PIWIL2 depletion. Intriguingly, our sequencing and subsequent database search for canonical piRNAs using both the piRNA and the repBase databases did not yield any hits, whereas the peak denoting the piRNA size at 25-31 nt was absent (**Fig. 6A**), both indicating that canonical piRNAs are not expressed in Caco2 cells. However, we did detect a peak at the ~16 nt size mark, which has been reported to be a product of piRNA processing and transposon targeting (**Fig. 6A**)(Wenda et al., 2017, Wang et al., 2015b, Xiol et al., 2012). These short RNAs are also enriched for

T/U at their 5' end (**Fig. 6B**), which further corroborates that this peak represents processed piRNAs

and it is not a product of random degradation. Indeed, when we aligned those short RNA sequences to

TE sequences, we identified a significant population of those to align with TEs (**Table S2**). More

specifically, when we examined the "antisense" piRNA-like sequences, meaning the ones

complementary to TEs that would indicate the piRNAs targeting those TEs, we found that these are

almost entirely absent in the PIWIL2 KO cells, compared to the WT control cells, including many piRNAs

that target LINE-1 elements (**Fig. 6C**). Additionally, many of the "sense" short RNA fragments that would

represent the cleavage products of the same LINE-1 elements are also absent in the PIWIL2 KO cells

(**Fig. 6D**). This indicates that these LINE-1 elements are not targeted for degradation by piRNAs in the

PIWIL2 KO cells, which would result in increased overall levels of LINE-1. Indeed, qRT-PCR of the

L1RE1 RNA transcript, which includes the first ORF of the LINE-1 mRNA that encodes the ORF1p RNA

binding protein required for retro-transposition(Beck et al., 2011, Elbarbary et al., 2016), confirmed that

LINE-1 levels were increased in the PIWIL2 KO cells (**Fig. 6E**). Altogether, these results demonstrate

that PIWIL2 depletion impairs piRNA-mediated targeting of TEs, especially of LINE-1 elements,

resulting in increased LINE-1 levels in these cells.

PIWIL2 depletion increases activity of the LINE-1 transposable element

Then, to examine whether the increased LINE-1 levels upon PIWIL2 depletion were also accompanied

by increased retro-transposition activity, we utilized a retro-transposition LINE1-GFP reporter

assay(Moran et al., 1996, Ostertag et al., 2000, Garcia-Perez et al., 2010, Kopera et al., 2016). In this

assay, a plasmid with a full-length retro-transposition competent LINE-1 element produces GFP only

when retro-transposition occurs and is an indicator of TE activity within the cell (denoted LINE1-GFP).

Additionally, the assay uses a negative control plasmid with an inactive LINE-1 element that contains a

defective ORF1p protein (LINE1 inactive (-)) that is unable to retro-transpose and produce GFP. Caco2

WT or PIWIL2 KO cells were transfected with either a constitutively active GFP reporter plasmid

(pCEP4+) to determine transfection efficiency, the LINE1-GFP retro-transposition reporter, or the LINE1

inactive (-) plasmid as a negative control. After 48 hours, transfected Caco2 WT and PIWIL2 KO cells

were placed in puromycin selection for 4 days and then imaged by fluorescence microscopy to identify

GFP positive cells (**Fig. 7A**). The average percentage of positive GFP expressing cells was normalized

to the transfection efficiency of each cell line for comparison (**Fig. 7B**). Caco2 WT + LINE1-GFP

fluorescence was minimal and not significantly different compared to Caco2 WT LINE-1 inactive (-)

cells, indicating there is minimal LINE-1 activity in the Caco2 WT cells. However, PIWIL2 KO + LINE1-

GFP exhibited significantly higher LINE-1 activity compared to the PIWIL2 KO + LINE1 inactive (-) cells,

indicating specific LINE-1 activation in the PIWIL2 KO cells; this increased activity was also significantly

higher compared to Caco2 WT + LINE1-GFP cells, demonstrating that PIWL2 depletion not only increases LINE-1 mRNA transcript levels, but also LINE1 activity in colon epithelial Caco2 cells.

PIWIL2 depletion promotes DNA damage

The DNA insertional activity of TEs and especially of LINE-1 elements can cause DNA damage, which is a hallmark of cancer (Gasior et al., 2006, Hanahan and Weinberg, 2011). Therefore, we examined whether the increased LINE-1 activity upon PIWIL2 depletion in Caco2 cells (**Fig. 7A,B**) also results in DNA damage in these cells (**Fig. 8**). Immunofluorescent staining followed by confocal microscopy of Caco2 WT and PIWIL2 KO cells for γH2AX, a phosphorylated histone marker of DNA double stranded breaks(Kuo and Yang, 2008), revealed that there was a significant increase in the number of γH2AX foci per cell (**Fig. 8A,B**). We also confirmed increased levels of γH2AX compared to total H2AX by immunoblotting (**Fig. 8C**). Furthermore, examination of γH2AX foci together with the LINE1-GFP retro-transposition assay showed substantial overlap of γH2AX foci with transposon activity (**Fig. 8D,E**). These results demonstrate that PIWIL2 depletion in somatic, well-differentiated colon epithelial Caco2 cells results in up regulation of transposon levels and activity, as well as in subsequent increased DNA damage, which are all promoters of pro-tumorigenic transformation. These findings, taken together with the dysregulation of the PIWI pathway and downregulation of PIWIL2 in colon tumors, as well as with the significantly increased ability of the PIWIL2 KO cells to grow in anchorage-independent conditions, underscore a tumor-suppressing role of PIWIL2 in the colon.

DISCUSSION

Prior studies have shown that the activation of TEs in tumors, including colon cancer, was frequent, although little is known regarding the potential mechanisms of TE activation (Pitkanen et al., 2014, Ewing et al., 2015, Solyom et al., 2012). Recently, the PIWI pathway, which is well-established to suppress transposons in the germline, has also been shown to be expressed in the soma and be implicated in cancer(Ross et al., 2014, Siomi et al., 2011). However, reports regarding the presence of PIWI proteins in somatic tissues and the role of this pathway in tumorigenesis have been conflicting, with some supporting a tumor-suppressing and others a tumor-promoting role of the pathway. Our bioinformatic analysis, as well as our RNAseq, qRT-PCR, and protein expression data, confirm expression of PIWIL2, the key regulator of the pathway, in the colon and in the well-differentiated colon epithelial Caco2 cells (**Fig. 5, Fig. S1, Table S1**). Our results also show that the pathway experiences significant allelic losses and its expression is downregulated in colon adenocarcinoma, with PIWIL2 having the highest percentage of deletions and allelic losses within the pathway (**Fig. 1**). Both PIWIL2 allelic loses and downregulation also correlate with poor patient survival (**Fig. 1, 2**). Furthermore, knockout of PIWIL2 in Caco2 cells resulted in increased expression and activity of the LINE-1 TE, the

most abundant and only autonomous TE in humans, as well as in DNA damage and in anchorage-independent growth, all hallmarks of pro-tumorigenic transformation (**Fig. 5,6,7,8**). Altogether, the data support the idea that PIWIL2 acts as a tumor suppressor in the colon.

Furthermore, when we interrogated PIWIL2 expression levels in the TCGA dataset by UALCAN analysis, PIWIL2 loss was more prevalent in the mucinous adenocarcinoma subtype of colon cancer (**Fig. 2E**), which is found in 10%-20% of colon cancer patients and is characterized by ≥50% of the tumor volume consisting of extracellular mucins(Fleming et al., 2012, Luo et al., 2019, Huang et al., 2021). Studies have also shown that mucinous adenocarcinoma tumors exhibit a higher frequency of microsatellite instability, CpG island methylation, and lower frequency of p53 mutations(Luo et al., 2019, Huang et al., 2021). Indeed, PIWIL2 downregulation in colon tumors seems to correlate with microsatellite instability, independently of p53 status (**Fig. 2D,F**). These findings further suggest the involvement of TE mobility in tumorigenesis, since retrotransposons like LINE-1 are known to give rise to de novo microsatellites during retro-transposition events due to internal proto-microsatellite sequences and poly-A mononucleotide repeats (Grandi and An, 2013). Since PIWIL2 downregulation seems to be independent of p53 status, which is key for maintaining genome integrity, it is likely that PIWIL2 dysregulation in microsatellite tumors is a potential mechanism of LINE-1 activation which would be interesting to be further interrogated.

Our database examination also showed that PIWL2 is downregulated in samples from patients with ulcerative colitis, which highlights a gap in current research, since the PIWI pathway has not been previously studied in colitis and in inflammatory bowel disease (**Fig. 4**). A few studies have indicated that there is upregulation of LINE-1 elements in inflammatory bowel disease, particularly in Crohn's Disease; however, the information on transposon expression and their regulation in these diseases is still very limited(Kanke et al., 2022, Wang et al., 2017). Given the above and that colitis and inflammatory bowel disease overall strongly predispose for colon cancer, it would be interesting to examine whether this PIWIL2 downregulation is a contributing factor to this predisposition, through upregulation of LINE elements in these diseases.

Interestingly, our *in vitro* studies showed that only a 50% knockout of PIWIL2 is tolerated by Caco2 cells, which correlates with the haploinsufficiency identified by the SWAN algorithm (**Fig. 1, 5A-C, Fig. S3**). In fact, this low tolerance for PIWIL2 loss may be due to the ensuing DNA damage caused by PIWIL2 downregulation (**Fig. 8**). However, we found that even a 50% depletion of PIWIL2 was able to significantly promote anchorage independent growth of Caco2 cells, which are otherwise well-differentiated and don't grow efficiently under these conditions (**Fig. 5F-J**). Acquisition of anchorage independent growth capability is a key event in epithelial cell pro-tumorigenic transformation. Furthermore, the increased DNA damage upon depletion of PIWIL2 was accompanied by increased

levels and activity of the LINE-1 TE, indicating that TE regulation and silencing is disrupted upon PIWIL2 depletion (**Fig. 7**). Both TE activation and DNA damage are driver events in tumorigenesis (Cajuso et al., 2019, Scott et al., 2016, Miki et al., 1992, Gasior et al., 2006). Therefore, these findings, taken together with our current results showing early downregulation of PIWIL2 in colon tumors or even in pre-cancerous IBD, underscore a potential role of PIWIL2 in early colon tumorigenesis, which deems further investigation.

Although our results in this study overall support a tumor suppressing role for PIWIL2 in colon tumorigenesis, there have been conflicting reports regarding the role of PIWIL2 in cancer (Lee et al., 2006, Lu et al., 2012, Wang et al., 2015a, Zeng et al., 2017, Eckstein et al., 2018, Meseure et al., 2020, Kishani Farahani et al., 2023, Shi et al., 2020). Along these lines, both our CRISPR/Cas9-generated PIWIL2 KO cells, as well as our PIWIL2 knockdown cells using a shRNA, exhibit slower proliferation rate, compared to the control cells, which would imply a tumor-promoting role for PIWIL2 (**Fig. 5E, Fig. S4**). However, unlike the anchorage-independent growth phenotype, we were not able to rescue the effect of PIWIL2 depletion on proliferation, using our full length PIWIL2-FLAG construct, which actually suppressed proliferation even more, further supporting the tumor suppressing role of PIWIL2 (**Fig. 5E**). It is likely that this discrepancy is due to the existence of additional PIWIL2 isoforms, which may have opposing roles to the full length PIWIL2 protein. Such isoforms have been reported in the literature and have been involved in tumorigenesis, or have been associated with certain cancer cells and tumor types (Gainetdinov et al., 2014, Ye et al., 2010). These isoforms apparently lack parts of the PAZ or PIWI domains, which are essential for piRNA processing and TE targeting, and seem to act through downstream pathways other than the canonical piRNA-TE targeting pathway (Gainetdinov et al., 2014, Ye et al., 2010). Although our PIWIL2 examination in this study shows an overall downregulation in colon cancer and supports a tumor-suppressing role of the protein through piRNA-mediated TE targeting, considering these additional isoforms in different cancer types could resolve any potentially conflicting findings regarding the role of PIWIL2 in cancer.

PIWIL2 is part of a larger protein complex (piRISC) and the SWAN analysis identified additional members of the PIWI-piRNA pathway and of piRISC as also significantly downregulated. For example, other top targets from the SWAN analysis include the exonuclease EXD1 found in pi-bodies, the RNA helicases DDX4 and MOV10L1, PLD6, and several members of the Tudor family of proteins that are key for the functionality of piRISC. These proteins have also been identified to be dysregulated in other cancer types, mainly breast and ovarian cancers, but they have not been studied in the colonic epithelium, since this is still a novel and ongoing field of investigation(Olotu et al., 2024, Kim et al., 2014, Lee et al., 2020). Therefore, it would be of particular interest to further investigate these additional PIWI pathway components and their roles in suppressing TEs in colonic diseases. In addition, our

finding of the expression of non-canonical piRNAs in Caco2 cells is intriguing and warrants further investigation: it is likely that the PIWI pathway in somatic tissues acts through and processes different sets of short RNAs to target transposons, as opposed to the well-characterized piRNA substrates of the pathway in the germline. In summary, our work, through using a nuanced bioinformatic approach and by performing cell - based assays, sheds light into the currently conflicting views about the role of the PIWI pathway in somatic tissues and in cancer, opening future directions of investigation that will fully assess the tumor suppressing potential of this pathway.

METHODS

Bioinformatic Analysis

Multiple bioinformatic tools were used throughout this study to investigate changes in the PIWI protein pathway using data from The Cancer Genome Atlas (TCGA). The analytical tools are detailed below.

Shifted Weighted Annotation Network (SWAN): SWAN is an analytical tool used to interrogate pathway and copy-number alterations (CNAs) within the TCGA data set and can be found at the following webpage: https://www.delaneyapps.com/. The Single-pathway SWAN tool was used to perform Gene Ontology Biological Process Analysis on the piRNA metabolic pathway in COAD with a 0.001 significance threshold over 1000 permutations of the algorithm. RNA data was integrated with CNA data for this analysis and a Wilcoxon rank test was calculated as described (Bowers et al., 2022).

cBioportal: cBioportal is another open-source web-based tool to visualize multidimensional cancer genomics data sets, including data from TCGA Pan Cancer Atlas (Cerami et al., 2012, Gao et al., 2013, de Bruijn et al., 2023). This tool can be found at the following webpage: https://www.cbioportal.org. The TCGA COAD data set was selected and then queried for samples with shallow deletion CNAs (loss of one allele; HETLOSS) of PIWIL2 and a plot of Overall Survival was generated using the Comparison/Survival Tool.

UALCAN: UALCAN is a web-based tool from the University of Alabama at Birmingham that allows users to stratify TCGA data for more in-depth analysis of TCGA expression profiles. This tool can be found at the following webpage: https://ualcan.path.uab.edu/. A pan-tissue analysis of PIWIL2 expression was performed across multiple tissuetypes to compare overall expression in normalsamples. PIWIL2 expression and methylation was then queried in the TCGA colon adenocarcinoma (COAD) data set and stratified by normal vs tumor, tumor stage, nodal metastasis, p53 status, histological subtype, biological sex, age, weight, and race. The median,interquartile range, minimum and maximum values

are displayed, and statistical significance was calculated by the UALCAN algorithm using a Welch's T-test as previously described (Chandrashekar et al., 2017, Chandrashekar et al., 2022). Results were exported from UALCAN, and graphical displays were made in Prism 10 (Graphpad).

GEPIA2: GEPIA2 is an interactive web-based tool for analyzing RNA-seq data from the TCGA and GTEx projects that can be found at the following webpage: http://gepia2.cancer-pku.cn/#index. The single gene analysis search for PIWIL2 was used to plot box plots stratified by cancer subtype (MSS, MSI-L, MSI-H) of gene expression. For this study, the median expression of colon adenocarcinoma (COAD) was plotted from TCGA data available with the upper and lower quartiles. The GEPIA2 program calculated statistical significance using a one-way ANOVA. A log_2-fold-change of 0.5 and p-value of 0.5 were set as thresholds for the analysis(Tang et al., 2019).

Kaplan-Meier Plotter: This web-based tool is capable of correlating mRNA expression data and survival data of colon cancer datasets from the Gene Expression Omibus (Gyorffy, 2024). A hazard ratio with 95% confidence intervals and logrank P value are calculated by the program. PIWIL2 was interrogated to determine Regression Free Survival and stratified by all tumor stages, stages 1-3 combined and stage 4 alone. This web-based tool can be found at the following webpage https://www.kmplot.com (Gyorffy, 2024).

PreMedIBD: PreMedIBD is a web-based data mining tool for meta-analysis of ulcerative colitis tissue gene expression. This tool can be found at the following webpage: https://premedibd.com/genes.html. PIWIL2, PIWIL4, DDX4, PLD6, EXD1, TDRD1, TDRD9, TDRD12 and MOV10L1 were queried and the log2-Fold gene expression changes between ulcerative colitis patients and non-IBD controls were determined for each gene. The meta-analysis was calculated as previously described using a random effects model and shows the meta-log2-Fold change and 95% confidence interval(Linggi et al., 2021). A log_2-fold-change of 0.5 was used as a threshold for significance. Datasets used are available in the Gene Expression Omnibus: GSE13367, GSE9452, GSE53306, GSE38713, GSE47908, GSE73661, GSE114527, GSE87466.

Cell Culture

Caco2 colon epithelial cells (ATCC, #HTB-37) were cultured in MEM (Corning, #10-010-CV) with 10% fetal bovine serum (FBS) (Gibco,Fetal Bovine Serum Qualified One Shot #A31605-02), 1 mM sodium pyruvate (Gibco, #11360-070), and 1x non-essential amino-acids (NEAA) solution (Gibco, #11140-050) and were maintained at 37°C with 5% CO_2. Caco2 cells were authenticated by the University of Arizona Genetics Core (via Science Exchange) and checked for misidentified, cross contaminated, or

genetically drifted cells and tested negative for mycoplasma contamination (LookOut Mycoplasma PCR Detection Kit, Sigma Aldrich). The Caco2 PIWIL2 knockout (KO) cell line was generated using CRISPR/Cas9 technology in collaboration with the Cell Models Core as part of the Medical University of South Carolina COBRE in Digestive and Liver Disease following the protocol by Ran et al.(Ran et al., 2013) The sgRNA sequence was designed to target exon 14 of PIWIL2 (sense:CACCGCCAATGAACTGATGCGTTGG/antisense: AAACCCAACGCATCAGTTCATTGGC) using the online Synthego tool (www.synthego.com) and synthesized by Integrated DNA Technologies, IDT. This PIWIL2-targeting sgRNA was cloned into the PX459 pSPCas9(BB)-2A-Puro vector and Caco2 cells were transfected with 30 µg the sgRNA-containing vector using Lipofectamine 3000 (Invitrogen) according to the manufacturer's protocol. After 24 hours, cells were selected with 5 µg/ml puromycin for 48 h. Successfully transfected cells were seeded to obtain single colonies, which were picked and expanded for screening by western blot and qPCR. PCR using primers designed to amplify the target region was performed (Forward1: CTCGAACACCTGGGCTCAAGCG and Reverse1: ACCACGCACAGAGGTTCATACC and Forward2: ATCACATTCACCTTCCTAAGAGA and Reverse2: GACCACGCACAGAGGTTCAT) and sequenced by Sanger sequencing (Eurofins Genomics). Alignment of the PCR products was performed to characterize the indel of the PIWIL2 KO.

RNA isolation and qPCR

RNA was isolated by adding 1ml of Trizol (Invitrogen # 15596018), incubating the sample at RT for 5 minutes then using the PureLink RNA mini kit (Invitrogen #12183018A) for total RNA, using the whole transcriptome protocol of the kit. Final RNA concentrations and purity were determined by using a NanoDrop spectrophotometer. RNA was converted to cDNA using the High Capacity cDNA Reverse Transcriptase Kit (Applied Biosystems #4368814). qPCR reactions were performed using the Taqman FAST Universal PCR master mix (Applied Biosystems #4352042), in a BioRad CFX96 Touch/Connect real time quantitative PCR machine. Data were analyzed and normalized to 18S ribosomal RNA by ΔΔCt calculations to calculate fold change compared to control samples.

RNA sequencing

RNA from Caco2 WT and PIWIL2 KO cells was extracted as above and short RNA sequencing was performed using the NovaSeq SE50 strategy. Sequenced reads were first processed using Cutadapt (v1.15) to trim the 3′ adapter sequence (5′-AGATCGGAAGAGCACACGTCT-3′). Adapter-trimmed reads were analyzed using the piPipes small RNA pipeline to quantify transposon-mapped reads and normalize sense and antisense counts as counts per million (CPM) uniquely mapped to the hg19 genome(Han et al., 2015b). In parallel, reads were aligned to Repbase and known piRNA clusters using Bowtie2(Langmead and Salzberg, 2012, Langmead et al., 2019) and only successfully aligned reads

were retained. These reads were then assessed for size distribution and 5′ nucleotide signature using the TBr2_basicanalyses.pl script (Rosenkranz et al., 2015). Results were summarized and plotted as shown in **Fig. 6.** The detailed list of hits is shown in **Table S2**.

The Caco2 total mRNA sequencing that was used to extract the PIWI1-4 expression data shown in **Table S1** have been previously published and are publicly available in the GEO public database, under the accession number GSE156860(Daulagala et al., 2024).

Immunoblotting

Whole-cell protein lysates were obtained using RIPA buffer (50 mM Tris, pH 7.4, 150 mM NaCl, 1% NP-40, 0.5% deoxycholic acid, and 0.1% SDS) supplemented with protease inhibitor (RPI Research Products, #P50750-1) and phosphatase inhibitor (Thermo Fisher Scientific, Halt Phosphatase inhibitor cocktail, #1862495) at 1:100 dilution. Cell culture plates were scraped with a cell scraper, lysates were homogenized through a syringe and then cleared by centrifugation at 15,000 rpm for 5 minutes. Protein quantification was performed using a Pierce BCA Protein Assay (Thermo Fisher Scientific, #23227). Protein extracts were mixed with Laemmli sample buffer and separated by SDS–PAGE using 4–20% Tris-Glycine eXtended (TGX) gels (Bio-Rad) and transferred to 0.2 μm nitrocellulose membranes (Bio-Rad) with the Trans-Blot Turbo Transfer System (Bio-Rad). Transfer efficiency was determined using a Ponceau S stain (Thermo Fisher Scientific, #A40000279). Membranes were blocked in 3% milk for 45 minutes and incubated overnight with primary antibody rotating at 4°C. The membranes were washed with 1x TBST, incubated with secondary antibody for 45 min at 4°C, followed by room temperature for 1 hour, and then washed with 1x TBST. Signals were detected by luminescence using Pierce ECL (Thermo Fisher Scientific, #32209) and a ChemiDoc Imaging System (Bio-Rad). Blots were quantified using the Analyze gels program in FIJI/ImageJ.

PIWIL2-FLAG Expression Constructs

For PIWIL2 expression and rescue experiments, a C-terminal PIWIL2-3xFLAG tagged construct was designed and synthesized using the LZRS-MS-NEO retroviral backbone with the PIWIL2-3xFLAG (PIWIL2-FLAG) CDNA sequence inserted at the EcoRI site (Genscript). The retrovirus was generated using HEK Pheonix cells according to standard protocols; and Caco2 wild type (WT) and PIWIL2 KO cells were infected with either the empty vector (LZRS) or LZRS-PIWIL2-FLAG. Retroviral infection was followed by antibiotic selection using 250 μg/ml G418 disulfate (Thermo Fisher Scientific, #J63871.AD) for 12 days, then maintaining the cells in 125 μg/ml G418 disulfate. PIWIL2 and FLAG expression was confirmed through immunoblotting.

shRNA Knockdown

shRNAs were derived from the pLKO.1-based TRC1 (Sigma-Aldrich/RNAi Consortium) shRNA library (pLKO.1-puro Non-Target shRNA Control, SHC016; PIWIL2 #7882 TRCN0000007882). Caco2 cells were transfected with 1.5 μg of the shRNA-containing vector using Lipofectamine 3000 (Invitrogen) according to the manufacturer's protocol and cell culture media was changed 16 hours post-transfection. 48 hours post transfection, protein and RNA was isolated, and cells were used for the xCelligence proliferation assay in Supplemental Figure 4.

xCELLigence Proliferation Assay

16 well E-plates (Agilent, # 300 600 890) were used for the experiment on an Agilent xCELLigence Real Time Cell Analysis (RTCA) Dual Purpose (DP) machine and RTCA Basic software (Agilent). Pre-warmed media was added to each well and used to take a background measurement for 1 minute. Cells were grown until 80% confluency, washed with 1xPBS, trypsinized, centrifuged at 1000 rpm for 5 min and then resuspended in cell culture media for a manual cell count using trypan blue. 20,000 cells/well were then plated for each cell line in a volume of 100 μl/well for a total volume of 150 μl/well. Each plate included blank wells with 150 μl of media for a negative control. Cells were allowed to settle and attach to the plate at room temperature in the cell culture hood for 30 minutes, after which the plate was inserted into the machine. The Xcelligence machine was set to take a measurement every hour for 50 hours. The experiment was completed with n = 3 biological replicates and 4 technical replicates per cell line that were averaged to determine the growth curve. Results were exported from the RTCA software and graphical displays were made in Prism 10 (Graphpad). The cell index from each run was normalized to the final end point of the Caco2 WT cell line to determine fold change and a two-way repeated measures ANOVA with Greenhouse–Geisser correction and Bonferroni correction.

Low Attachment (Anchorage-Independent Growth) Assay

Cells were trypisinzed with 0.25% Trypsin-EDTA (Gibco, #25200-056) and counted using a manual cell count and trypan blue (Corning, #25-900-CI). Cells were then suspended in culture media appropriate for the cell line that contained 2.5% basement membrane substrate (Cultrex Basement Membrane Extract, Pathclear, #3432-001-01) and plated on an ultra-low attachment 96-well plate (Corning, #3474) at a seeding density of 1000 cells per well. Plates were cultured for a total of 7 days, adding 50 ul of media to the wells every other day and were fixed using 4% paraformaldehyde (PFA) (Electron Microscopy Sciences, # 157-4) diluted to a final concentration of 1% PFA. The experiment was completed with n = 3 biological replicates and 9 technical replicates per cell line. Following fixation, plates were imaged on a Keyence BZ-X810 microscope at 2x magnification. Images were analyzed in

FIJI/ImageJ by thresholding and using the analyze particles function to measure the area, perimeter,
Feret's diameter, and circularity of the spheroids.

LINE1 Reporter Assay
The LINE1 reporter constructs used in this study were a gift from Eline Luning Prak. The plasmids used
in this study contain the indicated modifications to the pCEP4 vector backbone as previously
described(Moran et al., 1996, Ostertag et al., 2000, Garcia-Perez et al., 2010, Kopera et al., 2016). In
detail:

pCEP4puroeGFP (pCEP4+) The vector backbone is the pCEP4 plasmid with the hygromycin
resistance gene replaced with puromycin for selection. It contains the coding sequence for enhanced
GFP (eGFP) and constitutively expresses eGFP with puromycin selection. This is the positive control
for the retrotransposition assay and is used to determine the transfection efficiency (Addgene
#177922).

pJM111L1eGFP (LINE1 inactive) The vector backbone is the pCEP4 plasmid with the hygromycin
resistance gene replaced with puromycin for selection. This vector contains a retro-transposition
defective allele of LINE1 that contains two missense mutations ($RR_{261-262}AA$) in the LINE1 ORF1-
encoded protein (ORF1p) and also contains the eGFP indicator cassette. This plasmid cannot express
eGFP and is a negative control for the retro-transposition assay (Addgene #42941).

p99LINE1RP-GFP (LINE1-GFP reporter) The vector backbone is the pCEP4 plasmid with the
hygromycin resistance gene replaced with puromycin for selection. This vector lacks the CMV promoter
and contains a full-length retro-transposition competent LINE1 element and the eGFP indicator
cassette subcloned into the 3'UTR of LINE1 (Addgene #42940).

Caco2 wild type or PIWIL2 KO cells were seeded onto 18mm circular glass coverslips in 12 well plates
for immunofluorescence staining and allowed to grow until 70% confluency. Cells were transfected
using Lipofectamine 3000 (Thermo Fisher #L3000-001) according to manufacturer's instructions with
1µg of pCEP4puroeGFP, or 1µg pJM111L1eGFP or 1µg p99LINE1RP-GFP(Day 0). 16 hours post
transfection, the media was changed with fresh cell culture media (Day 1). At 48 hours post-transfection,
cells were placed in puromycin selection at a concentration of 5 µg/ml (Day 2). 48 hours later, media
was changed with fresh, and cells were kept in puromycin selection (Day 4). After 4 days of puromycin
selection (Day 6 post-transfection), the cells were then fixed with 4% paraformaldehyde, washed with
1xPBS with 10mM glycine followed by permeabilization with 0.02% Triton-X 100 and stained with DAPI.
For the co-localization experiments, the coverslips were fixed and stained for EGFP (Abcam
ab184601,1:100 dilution) and γH2AX (Cell Signaling 20 E3 (9718T), 1:400 dilution) and DAPI. Three
fields per condition were captured and the experiment was completed with n = 4 biological replicates
for Figure 7 and n = 3 biological replicates for Figure 8 and imaged using a Keyence BZ-X810

microscope. Images were analyzed by FIJI/ImageJ by thresholding and using the analyze particles function. The average number of retro-transposition events was divided by the transfection efficiency of the pCEP4+ positive control plasmid for each cell line.

Immunofluorescence

Cells were grown in 12-well plates on 18 mm circular sterile glass coverslips (Fisher Scientific #72222-01) until they reached 80% confluency. Cells were washed once with 1x PBS and fixed with 100% methanol (Thermo Fisher Scientific) at –20°C for 7 min followed by 1x PBS wash. Coverslips were then blocked with serum free Protein Block reagent (Dako, #X0909) at RT for 1hr and stained with primary antibodies diluted in Antibody Diluent (Dako, #S3022) as described in the antibodies table overnight at 4°C. Cells were then washed with 1x PBS and stained with the fluorescent-labeled secondary antibodies for 1 hour at room temperature. Coverslips were then washed with 1x PBS, co-stained with DAPI (Sigma-Aldrich, #D8417) and mounted with Aqua-Poly/Mount (Polysciences, #18606-20).

Confocal Imaging and Image processing

Confocal images of γH2AX foci were acquired using a Leica SP8 confocal microscope with a 63x Plan-Apochromat 1.4NA DIC oil immersion objective (Leica) and 405 nm and 633 nm lasers. Image acquisition was completed using the Leica Application Suite software at 2048 × 2048 resolution and with 0.3 μm intervals along the z-axis. The same imaging parameters were used across conditions for each set of experiments to allow for comparisons. Z-stacks of all samples were analyzed using the FIJI/ImageJ analysis software and a max projection of each image was used to visualize the nucleus and account for uneven cell thickness over the entire field. Three fields of each sample were captured, and the experiment was performed in n = 3 biological replicates. Total γH2AX foci were quantified by thresholding the images and using the analyze particles function, then normalized to the total number of cells in the field. Nuclei on the borders were excluded from the cell count calculation in FIJI.

Analysis and Statistics

For all measurements, sample size and related statistics are indicated in the respected figure legends and were analyzed and graphed using Prism 10 (GraphPad). All experiments were performed in at least three independent experiments, unless otherwise noted, with alpha = 0.05 and representative images are shown. Statistical tests are noted in each figure legend along with p-values. For tests that required multiple comparisons, the multiplicity adjusted p-value is reported.

Antibodies

Primary Antibodies

Antibody Name	Company	Catalogue Number	Animal	WB dilution	IF dilution
PIWIL2*	Sigma Aldrich	SAB3500749	Rb	1 to 1000	-
β-actin (Actin)	Cell Signaling	4967L	Rb	1 to 2000	-
Anti-FLAG M2	Sigma Aldrich	F1804	Ms	1 to 2000	-
H2AX total	Santa Cruz	sc517336	Ms	1 to 1000	-
γH2AX	Cell Signaling	20 E3 (9718T)	Rb	1 to 1000	1 to 400 (MeOH and PFA Fixed)
EGFP	Abcam	ab184601	Ms	-	1:100 (PFA Fixed)

623 * The PIWL2 antibody was validated by using the PIWIL2-FLAG construct in Fig 5D. All other antibodies

624 are standard, commercially available and vendor-validated antibodies.

626 <u>Secondary Antibodies</u>

Antibody Name	Company	Catalogue Number	Animal	WB dilution	IF dilution
HRP-anti-rabbit	Jackson ImmunoResearch	711-035-152	Rb	1 to 2000	-
HRP-anti-mouse	Jackson ImmunoResearch	715-035-150	Ms	1 to 2000	-
AlexaFluor 647 anti-mouse	Invitrogen	A21236	Ms	-	1 to 500 (MeOH and PFA Fixed)
AlexaFluor 488 anti-rabbit	Invitrogen	A11034	Rb	-	1 to 500 (PFA Fixed)

628 **qPCR Probes**

Probe name	Company	Assay ID
18S	Thermo Fisher	Hs99999901_s1
PIWIL2	Thermo Fisher	Hs01032719_m1
L1RE1(APFVNJ2)	Thermo Fisher	custom sequence

ACKNOWLEDGMENTS

We would like to thank Drs. Christiana Kappler and Stephen Duncan, Cell Models Core, Medical University of South Carolina Center of Biomedical Research Excellence (COBRE) in Digestive & Liver Disease (CDLD; National Institutes of Health P20 GM130457) for support with CRISPR/Cas9 reagents.

COMPETING INTERESTS

No competing interests declared.

FUNDING

This work was supported by National Institute of Health grants R01 DK124553, R01 DK136658, R21 CA246233, P20 GM130457 (COBRE in Digestive & Liver Disease, MUSC), P30 DK123704 (Digestive Disease Research Center, MUSC), to AK; R01 GM160815 to VG; and DP2 CA280626 to JRD. A.R. was supported by National Institutes of Health training grants TL1 TR001451, UL1 TR001450, T32 5T32DK124191, and 1F31DK138780-01.

DATA AND RESOURCE AVAILABILITY

All data are included within the manuscript, Figures, Supplementary Figures and Tables. The piRNA sequencing raw data shown in Fig 6 and Table S2 are being deposited to GEO and an accession number is pending and will be provided upon acceptance of the manuscript. The PIWIL2-FLAG construct will be deposited on Addgene and will be available upon acceptance of the manuscript.

REFERENCES

ARIUMI, Y. 2016. Guardian of the Human Genome: Host Defense Mechanisms against LINE-1 Retrotransposition. *Front Chem,* 4**,** 28.

BECK, C. R., COLLIER, P., MACFARLANE, C., MALIG, M., KIDD, J. M., EICHLER, E. E., BADGE, R. M. & MORAN, J. V. 2010. LINE-1 retrotransposition activity in human genomes. *Cell,* 141**,** 1159-70.

BECK, C. R., GARCIA-PEREZ, J. L., BADGE, R. M. & MORAN, J. V. 2011. LINE-1 elements in structural variation and disease. *Annu Rev Genomics Hum Genet,* 12**,** 187-215.

BERRENS, R. V., ANDREWS, S., SPENSBERGER, D., SANTOS, F., DEAN, W., GOULD, P., SHARIF, J., OLOVA, N., CHANDRA, T., KOSEKI, H., VON MEYENN, F. & REIK, W. 2017. An endosiRNA-Based Repression Mechanism Counteracts Transposon Activation during Global DNA Demethylation in Embryonic Stem Cells. *Cell Stem Cell,* 21**,** 694-703 e7.

BOWERS, R. R., JONES, C. M., PAZ, E. A., BARROWS, J. K., ARMESON, K. E., LONG, D. T. & DELANEY, J. R. 2022. SWAN pathway-network identification of common aneuploidy-based oncogenic drivers. *Nucleic Acids Res,* 50**,** 3673-3692.

CAJUSO, T., SULO, P., TANSKANEN, T., KATAINEN, R., TAIRA, A., HANNINEN, U. A., KONDELIN, J., FORSSTROM, L., VALIMAKI, N., AAVIKKO, M., KAASINEN, E., RISTIMAKI, A., KOSKENSALO, S., LEPISTO, A., RENKONEN-SINISALO, L., SEPPALA, T., KUOPIO, T., BOHM, J., MECKLIN, J. P., KILPIVAARA, O., PITKANEN, E., PALIN, K. & AALTONEN, L. A. 2019. Retrotransposon insertions can initiate colorectal cancer and are associated with poor survival. *Nat Commun,* 10**,** 4022.

CERAMI, E., GAO, J., DOGRUSOZ, U., GROSS, B. E., SUMER, S. O., AKSOY, B. A., JACOBSEN, A., BYRNE, C. J., HEUER, M. L., LARSSON, E., ANTIPIN, Y., REVA, B., GOLDBERG, A. P., SANDER, C. & SCHULTZ, N. 2012. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov,* 2**,** 401-4.

CHANDRASHEKAR, D. S., BASHEL, B., BALASUBRAMANYA, S. A. H., CREIGHTON, C. J., PONCE-RODRIGUEZ, I., CHAKRAVARTHI, B. & VARAMBALLY, S. 2017. UALCAN: A Portal for Facilitating Tumor Subgroup Gene Expression and Survival Analyses. *Neoplasia,* 19**,** 649-658.

CHANDRASHEKAR, D. S., KARTHIKEYAN, S. K., KORLA, P. K., PATEL, H., SHOVON, A. R., ATHAR, M., NETTO, G. J., QIN, Z. S., KUMAR, S., MANNE, U., CREIGHTON, C. J. & VARAMBALLY, S. 2022. UALCAN: An update to the integrated cancer data analysis platform. *Neoplasia,* 25**,** 18-27.

DAULAGALA, A. C., CETIN, M., NAIR-MENON, J., JIMENEZ, D. W., BRIDGES, M. C., BRADSHAW, A. D., SAHIN, O. & KOURTIDIS, A. 2024. The epithelial adherens junction component PLEKHA7 regulates ECM remodeling and cell behavior through miRNA-mediated regulation of MMP1 and LOX. *bioRxiv 10.1101/2024.05.28.596237.*

DE BRUIJN, I., KUNDRA, R., MASTROGIACOMO, B., TRAN, T. N., SIKINA, L., MAZOR, T., LI, X., OCHOA, A., ZHAO, G., LAI, B., ABESHOUSE, A., BAICEANU, D., CIFTCI, E., DOGRUSOZ, U., DUFILIE, A., ERKOC, Z., GARCIA LARA, E., FU, Z., GROSS, B., HAYNES, C., HEATH, A., HIGGINS, D., JAGANNATHAN, P., KALLETLA, K., KUMARI, P., LINDSAY, J., LISMAN, A., LEENKNEGT, B., LUKASSE, P., MADELA, D., MADUPURI, R., VAN NIEROP, P., PLANTALECH, O., QUACH, J., RESNICK, A. C., RODENBURG, S. Y. A., SATRAVADA, B. A., SCHAEFFER, F., SHERIDAN, R., SINGH, J., SIROHI, R., SUMER, S. O., VAN HAGEN, S., WANG, A., WILSON, M., ZHANG, H., ZHU, K., RUSK, N., BROWN, S., LAVERY, J. A., PANAGEAS, K. S., RUDOLPH, J. E., LENOUE-NEWTON, M. L., WARNER, J. L., GUO, X., HUNTER-ZINCK, H., YU, T. V., PILAI, S., NICHOLS, C., GARDOS, S. M., PHILIP, J., AACR PROJECT GENIE BPC CORE TEAM, A. P. G. C., KEHL, K. L., RIELY, G. J., SCHRAG, D., LEE, J., FIANDALO, M. V., SWEENEY, S. M., PUGH, T. J., SANDER, C., CERAMI, E., GAO, J. & SCHULTZ, N. 2023. Analysis and Visualization of Longitudinal Genomic and Clinical Data from the AACR Project GENIE Biopharma Collaborative in cBioPortal. *Cancer Res,* 83**,** 3861-3867.

DE FAZIO, S., BARTONICEK, N., DI GIACOMO, M., ABREU-GOODGER, C., SANKAR, A., FUNAYA, C., ANTONY, C., MOREIRA, P. N., ENRIGHT, A. J. & O'CARROLL, D. 2011. The endonuclease activity of Mili fuels piRNA amplification that silences LINE1 elements. *Nature,* 480**,** 259-63.

ECKSTEIN, M., JUNG, R., WEIGELT, K., SIKIC, D., STOHR, R., GEPPERT, C., AGAIMY, A., LIEB, V., HARTMANN, A., WULLICH, B., WACH, S. & TAUBERT, H. 2018. Piwi-like 1 and -2 protein expression levels are prognostic factors for muscle invasive urothelial bladder cancer patients. *Sci Rep,* 8**,** 17693.

ELBARBARY, R. A., LUCAS, B. A. & MAQUAT, L. E. 2016. Retrotransposons as regulators of gene expression. *Science,* 351**,** aac7247.

EWING, A. D., GACITA, A., WOOD, L. D., MA, F., XING, D., KIM, M. S., MANDA, S. S., ABRIL, G., PEREIRA, G., MAKOHON-MOORE, A., LOOIJENGA, L. H., GILLIS, A. J., HRUBAN, R. H., ANDERS, R. A., ROMANS, K. E., PANDEY, A., IACOBUZIO-DONAHUE, C. A., VOGELSTEIN, B., KINZLER, K. W., KAZAZIAN, H. H., JR. & SOLYOM, S. 2015. Widespread somatic L1 retrotransposition occurs early during gastrointestinal cancer evolution. *Genome Res,* 25**,** 1536-45.

FLEMING, M., RAVULA, S., TATISHCHEV, S. F. & WANG, H. L. 2012. Colorectal carcinoma: Pathologic aspects. *J Gastrointest Oncol,* 3**,** 153-73.

FROST, R. J., HAMRA, F. K., RICHARDSON, J. A., QI, X., BASSEL-DUBY, R. & OLSON, E. N. 2010. MOV10L1 is necessary for protection of spermatocytes against retrotransposons by Piwi-interacting RNAs. *Proc Natl Acad Sci U S A,* 107**,** 11847-52.

FU, K., TIAN, S., TAN, H., WANG, C., WANG, H., WANG, M., WANG, Y., CHEN, Z., WANG, Y., YUE, Q., XU, Q., ZHANG, S., LI, H., XIE, J., LIN, M., LUO, M., CHEN, F., YE, L. & ZHENG, K. 2019. Biological and RNA regulatory function of MOV10 in mammalian germ cells. *BMC Biol,* 17**,** 39.

GAINETDINOV, I. V., SKVORTSOVA, Y. V., STUKACHEVA, E. A., BYCHENKO, O. S., KONDRATIEVA, S. A., ZINOVIEVA, M. V. & AZHIKINA, T. L. 2014. Expression profiles of PIWIL2 short isoforms differ in testicular germ cell tumors of various differentiation subtypes. *PLoS One,* 9**,** e112528.

GAO, J., AKSOY, B. A., DOGRUSOZ, U., DRESDNER, G., GROSS, B., SUMER, S. O., SUN, Y., JACOBSEN, A., SINHA, R., LARSSON, E., CERAMI, E., SANDER, C. & SCHULTZ, N. 2013. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal,* 6**,** pl1.

GAO, J., JING, J., SHANG, G., CHEN, C., DUAN, M., YU, W., WANG, K., LUO, J., SONG, M., CHEN, K., CHEN, C., ZHANG, T. & DING, D. 2024. TDRD1 phase separation drives intermitochondrial cement assembly to promote piRNA biogenesis and fertility. *Dev Cell,* 59**,** 2704-2718 e6.

GAO, J. X. 2008. Cancer stem cells: the lessons from pre-cancerous stem cells. *J Cell Mol Med,* 12**,** 67-96.

GARCIA-PEREZ, J. L., MORELL, M., SCHEYS, J. O., KULPA, D. A., MORELL, S., CARTER, C. C., HAMMER, G. D., COLLINS, K. L., O'SHEA, K. S., MENENDEZ, P. & MORAN, J. V. 2010. Epigenetic silencing of engineered L1 retrotransposition events in human embryonic carcinoma cells. *Nature,* 466**,** 769-73.

GASIOR, S. L., WAKEMAN, T. P., XU, B. & DEININGER, P. L. 2006. The human LINE-1 retrotransposon creates DNA double-strand breaks. *J Mol Biol,* 357**,** 1383-93.

GORDON, I. O., AGRAWAL, N., WILLIS, E., GOLDBLUM, J. R., LOPEZ, R., ALLENDE, D., LIU, X., PATIL, D. Y., YERIAN, L., EL-KHIDER, F., FIOCCHI, C. & RIEDER, F. 2018. Fibrosis in ulcerative colitis is directly linked to severity and chronicity of mucosal inflammation. *Aliment Pharmacol Ther,* 47**,** 922-939.

GRANDI, F. C. & AN, W. 2013. Non-LTR retrotransposons and microsatellites: Partners in genomic variation. *Mob Genet Elements,* 3**,** e25674.

GRASSET, E., PINTO, M., DUSSAULX, E., ZWEIBAUM, A. & DESJEUX, J. F. 1984. Epithelial properties of human colonic carcinoma cell line Caco-2: electrical parameters. *Am J Physiol,* 247**,** C260-7.

GUIDOBONI, M., GAFA, R., VIEL, A., DOGLIONI, C., RUSSO, A., SANTINI, A., DEL TIN, L., MACRI, E., LANZA, G., BOIOCCHI, M. & DOLCETTI, R. 2001. Microsatellite instability and high content of activated cytotoxic lymphocytes identify colon cancer patients with a favorable prognosis. *Am J Pathol,* 159**,** 297-304.

GYORFFY, B. 2024. Integrated analysis of public datasets for the discovery and validation of survival-associated genes in solid tumors. *Innovation (Camb),* 5**,** 100625.

HAN, B. W., WANG, W., LI, C., WENG, Z. & ZAMORE, P. D. 2015a. Noncoding RNA. piRNA-guided transposon cleavage initiates Zucchini-dependent, phased piRNA production. *Science,* 348**,** 817-21.

HAN, B. W., WANG, W., ZAMORE, P. D. & WENG, Z. 2015b. piPipes: a set of pipelines for piRNA and transposon analysis via small RNA-seq, RNA-seq, degradome- and CAGE-seq, ChIP-seq and genomic DNA sequencing. *Bioinformatics,* 31**,** 593-5.

HANAHAN, D. & WEINBERG, R. A. 2011. Hallmarks of cancer: the next generation. *Cell,* 144**,** 646-74.

HANCKS, D. C. & KAZAZIAN, H. H., JR. 2016. Roles for retrotransposon insertions in human disease. *Mob DNA,* 7**,** 9.

HE, G., CHEN, L., YE, Y., XIAO, Y., HUA, K., JARJOURA, D., NAKANO, T., BARSKY, S. H., SHEN, R. & GAO, J. X. 2010. Piwil2 expressed in various stages of cervical neoplasia is a potential complementary marker for p16. *Am J Transl Res,* 2**,** 156-69.

HIDALGO, I. J., RAUB, T. J. & BORCHARDT, R. T. 1989. Characterization of the human colon carcinoma cell line (Caco-2) as a model system for intestinal epithelial permeability. *Gastroenterology,* 96**,** 736-49.

755 HUANG, A., YANG, Y., SHI, J. Y., LI, Y. K., XU, J. X., CHENG, Y. & GU, J. 2021. Mucinous adenocarcinoma: A unique
756   clinicopathological subtype in colorectal cancer. *World J Gastrointest Surg,* 13**,** 1567-1583.
757 IKEDA, S., TANAKA, K., OHTANI, R., KANDA, A., SOTOMARU, Y., KONO, T. & OBATA, Y. 2022. Disruption of piRNA
758   machinery by deletion of ASZ1/GASZ results in the expression of aberrant chimeric transcripts in
759   gonocytes. *J Reprod Dev,* 68**,** 125-136.
760 JACOBS, D. I., QIN, Q., LERRO, M. C., FU, A., DUBROW, R., CLAUS, E. B., DEWAN, A. T., WANG, G., LIN, H. & ZHU, Y.
761   2016. PIWI-Interacting RNAs in Gliomagenesis: Evidence from Post-GWAS and Functional Analyses.
762   *Cancer Epidemiol Biomarkers Prev,* 25**,** 1073-80.
763 JANG, H. S., SHAH, N. M., DU, A. Y., DAILEY, Z. Z., PEHRSSON, E. C., GODOY, P. M., ZHANG, D., LI, D., XING, X., KIM,
764   S., O'DONNELL, D., GORDON, J. I. & WANG, T. 2019. Transposable elements drive widespread expression
765   of oncogenes in human cancers. *Nat Genet,* 51**,** 611-617.
766 KANKE, M., KENNEDY NG, M. M., CONNELLY, S., SINGH, M., SCHANER, M., SHANAHAN, M. T., WOLBER, E. A.,
767   BEASLEY, C., LIAN, G., JAIN, A., LONG, M. D., BARNES, E. L., HERFARTH, H. H., ISAACS, K. L., HANSEN, J. J.,
768   KAPADIA, M., GUILLEM, J. G., FESCHOTTE, C., FUREY, T. S., SHEIKH, S. Z. & SETHUPATHY, P. 2022. Single-
769   Cell Analysis Reveals Unexpected Cellular Changes and Transposon Expression Signatures in the Colonic
770   Epithelium of Treatment-Naive Adult Crohn's Disease Patients. *Cell Mol Gastroenterol Hepatol,* 13**,** 1717-
771   1740.
772 KIM, E. R. & CHANG, D. K. 2014. Colorectal cancer in inflammatory bowel disease: the risk, pathogenesis,
773   prevention and diagnosis. *World J Gastroenterol,* 20**,** 9872-81.
774 KIM, K. H., KANG, Y. J., JO, J. O., OCK, M. S., MOON, S. H., SUH, D. S., YOON, M. S., PARK, E. S., JEONG, N., EO, W.
775   K., KIM, H. Y. & CHA, H. J. 2014. DDX4 (DEAD box polypeptide 4) colocalizes with cancer stem cell marker
776   CD133 in ovarian cancers. *Biochem Biophys Res Commun,* 447**,** 315-22.
777 KISHANI FARAHANI, R., SOLEIMANPOUR, S., GOLMOHAMMADI, M. & SOLEIMANPOUR-LICHAEI, H. R. 2023.
778   PIWIL2 Regulates the Proliferation, Apoptosis and Colony Formation of Colorectal Cancer Cell Line. *Iran J
779   Biotechnol,* 21**,** e3176.
780 KOPERA, H. C., LARSON, P. A., MOLDOVAN, J. B., RICHARDSON, S. R., LIU, Y. & MORAN, J. V. 2016. LINE-1 Cultured
781   Cell Retrotransposition Assay. *Methods Mol Biol,* 1400**,** 139-56.
782 KOURTIDIS, A., NGOK, S. P., PULIMENO, P., FEATHERS, R. W., CARPIO, L. R., BAKER, T. R., CARR, J. M., YAN, I. K.,
783   BORGES, S., PEREZ, E. A., STORZ, P., COPLAND, J. A., PATEL, T., THOMPSON, E. A., CITI, S. &
784   ANASTASIADIS, P. Z. 2015. Distinct E-cadherin-based complexes regulate cell behaviour through miRNA
785   processing or Src and p120 catenin activity. *Nat Cell Biol,* 17**,** 1145-57.
786 KU, H. Y. & LIN, H. 2014. PIWI proteins and their interactors in piRNA biogenesis, germline development and gene
787   expression. *Natl Sci Rev,* 1**,** 205-218.
788 KUO, L. J. & YANG, L. X. 2008. Gamma-H2AX - a novel biomarker for DNA double-strand breaks. *In Vivo,* 22**,** 305-9.
789 KURAMOCHI-MIYAGAWA, S., WATANABE, T., GOTOH, K., TOTOKI, Y., TOYODA, A., IKAWA, M., ASADA, N., KOJIMA,
790   K., YAMAGUCHI, Y., IJIRI, T. W., HATA, K., LI, E., MATSUDA, Y., KIMURA, T., OKABE, M., SAKAKI, Y., SASAKI, H.
791   & NAKANO, T. 2008. DNA methylation of retrotransposon genes is regulated by Piwi family members MILI
792   and MIWI2 in murine fetal testes. *Genes Dev,* 22**,** 908-17.
793 LANGMEAD, B. & SALZBERG, S. L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods,* 9**,** 357-9.
794 LANGMEAD, B., WILKS, C., ANTONESCU, V. & CHARLES, R. 2019. Scaling read aligners to hundreds of threads on
795   general-purpose processors. *Bioinformatics,* 35**,** 421-432.
796 LEE, E., LOKMAN, N. A., OEHLER, M. K., RICCIARDELLI, C. & GRUTZNER, F. 2020. A Comprehensive Molecular and
797   Clinical Analysis of the piRNA Pathway Genes in Ovarian Cancer. *Cancers (Basel),* 13.
798 LEE, J. H., SCHUTTE, D., WULF, G., FUZESI, L., RADZUN, H. J., SCHWEYER, S., ENGEL, W. & NAYERNIA, K. 2006.
799   Stem-cell protein Piwil2 is widely expressed in tumors and inhibits apoptosis through activation of
800   Stat3/Bcl-XL pathway. *Hum Mol Genet,* 15**,** 201-11.
801 LI, Y., ZHANG, Y. & LIU, M. 2021. Knockout Gene-Based Evidence for PIWI-Interacting RNA Pathway in Mammals.
802   *Front Cell Dev Biol,* 9**,** 681188.
803 LINGGI, B., JAIRATH, V., ZOU, G., SHACKELTON, L. M., MCGOVERN, D. P. B., SALAS, A., VERSTOCKT, B.,
804   SILVERBERG, M. S., NAYERI, S., FEAGAN, B. G. & VANDE CASTEELE, N. 2021. Meta-analysis of gene
805   expression disease signatures in colonic biopsy tissue from patients with ulcerative colitis. *Sci Rep,* 11**,**
806   18243.

LIU, J. J., SHEN, R., CHEN, L., YE, Y., HE, G., HUA, K., JARJOURA, D., NAKANO, T., RAMESH, G. K., SHAPIRO, C. L., BARSKY, S. H. & GAO, J. X. 2010. Piwil2 is expressed in various stages of breast cancers and has the potential to be used as a novel biomarker. *Int J Clin Exp Pathol,* 3**,** 328-37.

LU, Y., ZHANG, K., LI, C., YAO, Y., TAO, D., LIU, Y., ZHANG, S. & MA, Y. 2012. Piwil2 suppresses p53 by inducing phosphorylation of signal transducer and activator of transcription 3 in tumor cells. *PLoS One,* 7**,** e30999.

LUO, C., CEN, S., DING, G. & WU, W. 2019. Mucinous colorectal adenocarcinoma: clinical pathology and treatment options. *Cancer Commun (Lond),* 39**,** 13.

MA, L., BUCHOLD, G. M., GREENBAUM, M. P., ROY, A., BURNS, K. H., ZHU, H., HAN, D. Y., HARRIS, R. A., COARFA, C., GUNARATNE, P. H., YAN, W. & MATZUK, M. M. 2009. GASZ is essential for male meiosis and suppression of retrotransposon expression in the male germline. *PLoS Genet,* 5**,** e1000635.

MARTINEZ, V. D., VUCIC, E. A., THU, K. L., HUBAUX, R., ENFIELD, K. S., PIKOR, L. A., BECKER-SANTOS, D. D., BROWN, C. J., LAM, S. & LAM, W. L. 2015. Unique somatic and malignant expression patterns implicate PIWI-interacting RNAs in cancer-type specific biology. *Sci Rep,* 5**,** 10423.

MATHIOUDAKIS, N., PALENCIA, A., KADLEC, J., ROUND, A., TRIPSIANES, K., SATTLER, M., PILLAI, R. S. & CUSACK, S. 2012. The multiple Tudor domain-containing protein TDRD1 is a molecular scaffold for mouse Piwi proteins and piRNA biogenesis factors. *RNA,* 18**,** 2056-72.

MESEURE, D., VACHER, S., BOUDJEMAA, S., LAE, M., NICOLAS, A., LECLERE, R., CHEMLALI, W., CHAMPENOIS, G., SCHNITZLER, A., LESAGE, L., DUBOIS, T. & BIECHE, I. 2020. Biopathological Significance of PIWI-piRNA Pathway Deregulation in Invasive Breast Carcinomas. *Cancers (Basel),* 12.

MIKI, Y., NISHISHO, I., HORII, A., MIYOSHI, Y., UTSUNOMIYA, J., KINZLER, K. W., VOGELSTEIN, B. & NAKAMURA, Y. 1992. Disruption of the APC gene by a retrotransposal insertion of L1 sequence in a colon cancer. *Cancer Res,* 52**,** 643-5.

MORAN, J. V., HOLMES, S. E., NAAS, T. P., DEBERARDINIS, R. J., BOEKE, J. D. & KAZAZIAN, H. H., JR. 1996. High frequency retrotransposition in cultured mammalian cells. *Cell,* 87**,** 917-27.

NAM, C. H., YOUK, J., KIM, J. Y., LIM, J., PARK, J. W., OH, S. A., LEE, H. J., PARK, J. W., WON, H., LEE, Y., JEONG, S. Y., LEE, D. S., OH, J. W., HAN, J., LEE, J., KWON, H. W., KIM, M. J. & JU, Y. S. 2023. Widespread somatic L1 retrotransposition in normal colorectal epithelium. *Nature,* 617**,** 540-547.

NISHIMASU, H., ISHIZU, H., SAITO, K., FUKUHARA, S., KAMATANI, M. K., BONNEFOND, L., MATSUMOTO, N., NISHIZAWA, T., NAKANAGA, K., AOKI, J., ISHITANI, R., SIOMI, H., SIOMI, M. C. & NUREKI, O. 2012. Structure and function of Zucchini endoribonuclease in piRNA biogenesis. *Nature,* 491**,** 284-7.

OLOTU, O., KOSKENNIEMI, A. R., MA, L., PARAMONOV, V., LAASANEN, S., LOURAMO, E., BOURGERY, M., LEHTINIEMI, T., LAASANEN, S., RIVERO-MULLER, A., LOYTTYNIEMI, E., SAHLGREN, C., WESTERMARCK, J., VENTELA, S., VISAKORPI, T., POUTANEN, M., VAINIO, P., MAKELA, J. A. & KOTAJA, N. 2024. Germline-specific RNA helicase DDX4 forms cytoplasmic granules in cancer cells and promotes tumor growth. *Cell Rep,* 43**,** 114430.

OSTERTAG, E. M., PRAK, E. T., DEBERARDINIS, R. J., MORAN, J. V. & KAZAZIAN, H. H., JR. 2000. Determination of L1 retrotransposition kinetics in cultured cells. *Nucleic Acids Res,* 28**,** 1418-23.

PANDEY, R. R., TOKUZAWA, Y., YANG, Z., HAYASHI, E., ICHISAKA, T., KAJITA, S., ASANO, Y., KUNIEDA, T., SACHIDANANDAM, R., CHUMA, S., YAMANAKA, S. & PILLAI, R. S. 2013. Tudor domain containing 12 (TDRD12) is essential for secondary PIWI interacting RNA biogenesis in mice. *Proc Natl Acad Sci U S A,* 110**,** 16492-7.

PENG, J. C. & LIN, H. 2013. Beyond transposons: the epigenetic and somatic functions of the Piwi-piRNA mechanism. *Curr Opin Cell Biol,* 25**,** 190-4.

PITKANEN, E., CAJUSO, T., KATAINEN, R., KAASINEN, E., VALIMAKI, N., PALIN, K., TAIPALE, J., AALTONEN, L. A. & KILPIVAARA, O. 2014. Frequent L1 retrotranspositions originating from TTC28 in colorectal cancer. *Oncotarget,* 5**,** 853-9.

RAMAT, A. & SIMONELIG, M. 2021. Functions of PIWI Proteins in Gene Regulation: New Arrows Added to the piRNA Quiver. *Trends Genet,* 37**,** 188-200.

RAN, F. A., HSU, P. D., WRIGHT, J., AGARWALA, V., SCOTT, D. A. & ZHANG, F. 2013. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc,* 8**,** 2281-2308.

REUTER, M., CHUMA, S., TANAKA, T., FRANZ, T., STARK, A. & PILLAI, R. S. 2009. Loss of the Mili-interacting Tudor domain-containing protein-1 activates transposons and alters the Mili-associated small RNA profile. *Nat Struct Mol Biol,* 16**,** 639-46.

RODRIGUEZ-MARTIN, B., ALVAREZ, E. G., BAEZ-ORTEGA, A., ZAMORA, J., SUPEK, F., DEMEULEMEESTER, J.,
SANTAMARINA, M., JU, Y. S., TEMES, J., GARCIA-SOUTO, D., DETERING, H., LI, Y., RODRIGUEZ-CASTRO, J.,
DUESO-BARROSO, A., BRUZOS, A. L., DENTRO, S. C., BLANCO, M. G., CONTINO, G., ARDELJAN, D., TOJO,
M., ROBERTS, N. D., ZUMALAVE, S., EDWARDS, P. A., WEISCHENFELDT, J., PUIGGROS, M., CHONG, Z.,
CHEN, K., LEE, E. A., WALA, J. A., RAINE, K. M., BUTLER, A., WASZAK, S. M., NAVARRO, F. C. P.,
SCHUMACHER, S. E., MONLONG, J., MAURA, F., BOLLI, N., BOURQUE, G., GERSTEIN, M., PARK, P. J.,
WEDGE, D. C., BEROUKHIM, R., TORRENTS, D., KORBEL, J. O., MARTINCORENA, I., FITZGERALD, R. C., VAN
LOO, P., KAZAZIAN, H. H., BURNS, K. H., GROUP, P. S. V. W., CAMPBELL, P. J., TUBIO, J. M. C. &
CONSORTIUM, P. 2020. Pan-cancer analysis of whole genomes identifies driver rearrangements promoted
by LINE-1 retrotransposition. *Nat Genet,* 52**,** 306-319.
ROSENKRANZ, D., HAN, C. T., ROOVERS, E. F., ZISCHLER, H. & KETTING, R. F. 2015. Piwi proteins and piRNAs in
mammalian oocytes and early embryos: From sample to sequence. *Genom Data,* 5**,** 309-13.
ROSS, R. J., WEINER, M. M. & LIN, H. 2014. PIWI proteins and PIWI-interacting RNAs in the soma. *Nature,* 505**,** 353-
359.
SAMBUY, Y., DE ANGELIS, I., RANALDI, G., SCARINO, M. L., STAMMATI, A. & ZUCCO, F. 2005. The Caco-2 cell line as
a model of the intestinal barrier: influence of cell and culture-related factors on Caco-2 cell functional
characteristics. *Cell Biol Toxicol,* 21**,** 1-26.
SAXE, J. P., CHEN, M., ZHAO, H. & LIN, H. 2013. Tdrkh is essential for spermatogenesis and participates in primary
piRNA biogenesis in the germline. *EMBO J,* 32**,** 1869-85.
SCOTT, E. C., GARDNER, E. J., MASOOD, A., CHUANG, N. T., VERTINO, P. M. & DEVINE, S. E. 2016. A hot L1
retrotransposon evades somatic repression and initiates human colorectal cancer. *Genome Res,* 26**,** 745-
55.
SHI, S., YANG, Z. Z., LIU, S., YANG, F. & LIN, H. 2020. PIWIL1 promotes gastric cancer via a piRNA-independent
mechanism. *Proc Natl Acad Sci U S A,* 117**,** 22390-22401.
SIEGEL, R. L., GIAQUINTO, A. N. & JEMAL, A. 2024. Cancer statistics, 2024. *CA Cancer J Clin,* 74**,** 12-49.
SIGURDSSON, M. I., SMITH, A. V., BJORNSSON, H. T. & JONSSON, J. J. 2012. The distribution of a germline
methylation marker suggests a regional mechanism of LINE-1 silencing by the piRNA-PIWI system. *BMC
Genet,* 13**,** 31.
SIOMI, H. & SIOMI, M. C. 2015. RNA. Phased piRNAs tackle transposons. *Science,* 348**,** 756-7.
SIOMI, M. C., MANNEN, T. & SIOMI, H. 2010. How does the royal family of Tudor rule the PIWI-interacting RNA
pathway? *Genes Dev,* 24**,** 636-46.
SIOMI, M. C., SATO, K., PEZIC, D. & ARAVIN, A. A. 2011. PIWI-interacting small RNAs: the vanguard of genome
defence. *Nat Rev Mol Cell Biol,* 12**,** 246-58.
SOLYOM, S., EWING, A. D., RAHRMANN, E. P., DOUCET, T., NELSON, H. H., BURNS, M. B., HARRIS, R. S., SIGMON,
D. F., CASELLA, A., ERLANGER, B., WHEELAN, S., UPTON, K. R., SHUKLA, R., FAULKNER, G. J.,
LARGAESPADA, D. A. & KAZAZIAN, H. H., JR. 2012. Extensive somatic L1 retrotransposition in colorectal
tumors. *Genome Res,* 22**,** 2328-38.
SUN, Y. H., LEE, B. & LI, X. Z. 2022. The birth of piRNAs: how mammalian piRNAs are produced, originated, and
evolved. *Mamm Genome,* 33**,** 293-311.
TANG, Z., KANG, B., LI, C., CHEN, T. & ZHANG, Z. 2019. GEPIA2: an enhanced web server for large-scale expression
profiling and interactive analysis. *Nucleic Acids Res,* 47**,** W556-W560.
TUBIO, J. M., LI, Y., JU, Y. S., MARTINCORENA, I., COOKE, S. L., TOJO, M., GUNDEM, G., PIPINIKAS, C. P., ZAMORA,
J., RAINE, K., MENZIES, A., ROMAN-GARCIA, P., FULLAM, A., GERSTUNG, M., SHLIEN, A., TARPEY, P. S.,
PAPAEMMANUIL, E., KNAPPSKOG, S., VAN LOO, P., RAMAKRISHNA, M., DAVIES, H. R., MARSHALL, J.,
WEDGE, D. C., TEAGUE, J. W., BUTLER, A. P., NIK-ZAINAL, S., ALEXANDROV, L., BEHJATI, S., YATES, L. R.,
BOLLI, N., MUDIE, L., HARDY, C., MARTIN, S., MCLAREN, S., O'MEARA, S., ANDERSON, E., MADDISON, M.,
GAMBLE, S., GROUP, I. B. C., GROUP, I. B. C., GROUP, I. P. C., FOSTER, C., WARREN, A. Y., WHITAKER, H.,
BREWER, D., EELES, R., COOPER, C., NEAL, D., LYNCH, A. G., VISAKORPI, T., ISAACS, W. B., VAN'T VEER,
L., CALDAS, C., DESMEDT, C., SOTIRIOU, C., APARICIO, S., FOEKENS, J. A., EYFJORD, J. E., LAKHANI, S. R.,
THOMAS, G., MYKLEBOST, O., SPAN, P. N., BORRESEN-DALE, A. L., RICHARDSON, A. L., VAN DE VIJVER,
M., VINCENT-SALOMON, A., VAN DEN EYNDEN, G. G., FLANAGAN, A. M., FUTREAL, P. A., JANES, S. M.,
BOVA, G. S., STRATTON, M. R., MCDERMOTT, U. & CAMPBELL, P. J. 2014. Mobile DNA in cancer. Extensive

transduction of nonrepetitive DNA mediated by L1 retrotransposition in cancer genomes. *Science,* 345**,** 1251343.

UNHAVAITHAYA, Y., HAO, Y., BEYRET, E., YIN, H., KURAMOCHI-MIYAGAWA, S., NAKANO, T. & LIN, H. 2009. MILI, a PIWI-interacting RNA-binding protein, is required for germ line stem cell self-renewal and appears to positively regulate translation. *J Biol Chem,* 284**,** 6507-19.

VOUREKAS, A., ZHENG, K., FU, Q., MARAGKAKIS, M., ALEXIOU, P., MA, J., PILLAI, R. S., MOURELATOS, Z. & WANG, P. J. 2015. The RNA helicase MOV10L1 binds piRNA precursors to initiate piRNA processing. *Genes Dev,* 29**,** 617-29.

WALTER, M., TEISSANDIER, A., PEREZ-PALACIOS, R. & BOURC'HIS, D. 2016. An epigenetic switch ensures transposon repression upon dynamic loss of DNA methylation in embryonic stem cells. *Elife,* 5.

WANG, H. L., CHEN, B. B., CAO, X. G., WANG, J., HU, X. F., MU, X. Q. & CHEN, X. B. 2015a. The clinical significances of the abnormal expressions of Piwil1 and Piwil2 in colonic adenoma and adenocarcinoma. *Onco Targets Ther,* 8**,** 1259-64.

WANG, L., NORRIS, E. T. & JORDAN, I. K. 2017. Human Retrotransposon Insertion Polymorphisms Are Associated with Health and Disease via Gene Regulatory Phenotypes. *Front Microbiol,* 8**,** 1418.

WANG, W., HAN, B. W., TIPPING, C., GE, D. T., ZHANG, Z., WENG, Z. & ZAMORE, P. D. 2015b. Slicing and Binding by Ago3 or Aub Trigger Piwi-Bound piRNA Production by Distinct Mechanisms. *Mol Cell,* 59**,** 819-30.

WANG, Z. Y., GE, L. P., OUYANG, Y., JIN, X. & JIANG, Y. Z. 2024. Targeting transposable elements in cancer: developments and opportunities. *Biochim Biophys Acta Rev Cancer,* 1879**,** 189143.

WATANABE, T. & LIN, H. 2014. Posttranscriptional regulation of gene expression by Piwi proteins and piRNAs. *Mol Cell,* 56**,** 18-27.

WEICK, E. M. & MISKA, E. A. 2014. piRNAs: from biogenesis to function. *Development,* 141**,** 3458-71.

WENDA, J. M., HOMOLKA, D., YANG, Z., SPINELLI, P., SACHIDANANDAM, R., PANDEY, R. R. & PILLAI, R. S. 2017. Distinct Roles of RNA Helicases MVH and TDRD9 in PIWI Slicing-Triggered Mammalian piRNA Biogenesis and Function. *Dev Cell,* 41**,** 623-637 e9.

XIOL, J., CORA, E., KOGLGRUBER, R., CHUMA, S., SUBRAMANIAN, S., HOSOKAWA, M., REUTER, M., YANG, Z., BERNINGER, P., PALENCIA, A., BENES, V., PENNINGER, J., SACHIDANANDAM, R. & PILLAI, R. S. 2012. A role for Fkbp6 and the chaperone machinery in piRNA amplification and transposon silencing. *Mol Cell,* 47**,** 970-9.

YAN, Z., HU, H. Y., JIANG, X., MAIERHOFER, V., NEB, E., HE, L., HU, Y., HU, H., LI, N., CHEN, W. & KHAITOVICH, P. 2011. Widespread expression of piRNA-like molecules in somatic tissues. *Nucleic Acids Res,* 39**,** 6596-607.

YANG, Z., CHEN, K. M., PANDEY, R. R., HOMOLKA, D., REUTER, M., JANEIRO, B. K., SACHIDANANDAM, R., FAUVARQUE, M. O., MCCARTHY, A. A. & PILLAI, R. S. 2016. PIWI Slicing and EXD1 Drive Biogenesis of Nuclear piRNAs from Cytosolic Targets of the Mouse piRNA Pathway. *Mol Cell,* 61**,** 138-52.

YE, Y., YIN, D. T., CHEN, L., ZHOU, Q., SHEN, R., HE, G., YAN, Q., TONG, Z., ISSEKUTZ, A. C., SHAPIRO, C. L., BARSKY, S. H., LIN, H., LI, J. J. & GAO, J. X. 2010. Identification of Piwil2-like (PL2L) proteins that promote tumorigenesis. *PLoS One,* 5**,** e13406.

ZENG, G., ZHANG, D., LIU, X., KANG, Q., FU, Y., TANG, B., GUO, W., ZHANG, Y., WEI, G. & HE, D. 2017. Co-expression of Piwil2/Piwil4 in nucleus indicates poor prognosis of hepatocellular carcinoma. *Oncotarget,* 8**,** 4607-4617.

ZENG, Q., WAN, H., ZHAO, S., XU, H., TANG, T., OWARE, K. A. & QU, S. 2020. Role of PIWI-interacting RNAs on cell survival: Proliferation, apoptosis, and cycle. *IUBMB Life,* 72**,** 1870-1878.

ZHANG, X., YU, L., YE, S., XIE, J., HUANG, X., ZHENG, K. & SUN, B. 2019. MOV10L1 Binds RNA G-Quadruplex in a Structure-Specific Manner and Resolves It More Efficiently Than MOV10. *iScience,* 17**,** 36-48.

ZHUO, C., LI, Q., WU, Y., LI, Y., NIE, J., LI, D., PENG, J., LIAN, P., LI, B., CAI, G., LI, X. & CAI, S. 2015. LINE-1 hypomethylation in normal colon mucosa is associated with poor survival in Chinese patients with sporadic colon cancer. *Oncotarget,* 6**,** 23820-36.

    FIGURE LEGENDS

    Figure 1. The PIWI-piRNA pathway experiences copy number alterations (CNAs) in colon cancer.
    Shifted Weighted Annotation Network (SWAN) analysis was performed to interrogate pathway and
    CNAs within The Cancer Genome Atlas (TCGA) dataset. The Single-pathway SWAN tool was used to
    perform Gene Ontology Biological Process Analysis on the piRNA metabolic pathway in colon
    adenocarcinoma (COAD). A) SWAN pathway scoring for each member of the piRNA pathway where
    blue represents negatively scored allelic losses and red represents positively scored copy number
    gains within the network with an overall trend of haploinsufficiency as indicated by the grey line
    (Wilcoxon rank sum p < 1.7E-10). B) Circos plot mapping the frequency of allelic losses (blue) or gains
    (red) to each respective chromosome labeled on the outside ring. Gene labels are included if a score
    was beyond one standard deviation from a zero-change value. C) Table of the percentages of CNAs
    for each member of the piRNA pathway ranked by percent biallelic deletions. D) The cBioportal
    database was used to interrogate TCGA COAD dataset for Overall survival of PIWIL2 allelic loss
    compared to unaltered tumors. Red indicates PIWIL2 shallow deletions while blue indicates samples
    with no CNAs. (Logrank Test p = 0.0204)

    Figure 2. PIWIL2 is differentially expressed in colon tumors of advanced stage and nodal
    metastasis, histological subtype, and age. PIWIL2 expression was queried in the TCGA colon
    adenocarcinoma (COAD) data set and stratified by comparing normal colon tissue vs A) primary tumor,
    B) tumor stage, C) tumor nodal metastasis, D) tumor p53 status, E) tumor histological subtype, F) tumor
    microsatellite stability status, as well as G) age, H) weight, I) race and J) biological sex of colon cancer
    patients. Normal tissues were defined as adjacent non-cancerous tissue. For panels A-E and G-I the
    UALCAN web-based tool was used to stratify TCGA RNA-seq data for more in-depth analysis of
    expression profiles. The box and whisker plots display the median, interquartile range, minimum and
    maximum values of PIWIL2 expression in transcript per million (TPM) for each dataset, and *p<0.05,
    **p<0.01, ***p<0.001. For panel F The GEPIA2 tool was utilized to further stratify PIWIL2 expression
    in the TCGA COAD dataset by: micro-satellite stable (MSS), micro-satellite instable low (MSI-L), and
    micro-satellite instable high (MSI-H). The GEPIA2 program calculated statistical significance using one-
    way ANOVA (*p<0.05). K-M) The Kaplan-Meier Plotter tool was used to determine if PIWIL2 expression
    impacted patient regression free survival for all stages combined (K), Stages 1+2+3 (L) and Stage 4(M).

    Figure 3. The promoter of PIWIL2 is highly methylated in colon cancer. PIWIL2 promoter
    methylation was queried in the TCGA colon adenocarcinoma (COAD) data set and stratified by
    comparing normal colon tissue vs A) primary tumor, B) tumor stage, C) tumor nodal metastasis, D)
    tumor p53 status, E) tumor histological subtype, as well as F) biological sex, G) age, H) weight, and I)

race of colon cancer patients. Normal tissues were defined as adjacent non-cancerous tissue. The

median, interquartile range, minimum and maximum values of the methylation beta value is displayed

and *p <0.05, **p<0.01, ***p<0.001. A beta value closer to 1 indicates higher levels of methylation.

Figure 4. PIWIL2 is downregulated in ulcerative colitis patients compared to healthy, non-IBD

patients. A meta-analysis of ulcerative colitis tissue gene expression was performed using the

PreMedIBD Gene Tool. A) PIWIL2, B) PIWIL4, C) MOV10L1, D) TDRD1, E) TDRD9, F) TDRD12, G)

DDX4, H) PLD6 and I) EXD1 were queried and the $\log_2$-fold gene expression changes between

ulcerative colitis patients and non-IBD controls were determined for each gene. The meta-analysis was

calculated using a random effects model and shows the meta-log2-Fold change and 95% confidence

interval. Ulcerative Colitis datasets used are available in the Gene Expression Omnibus: GSE13367,

GSE9452, GSE53306, GSE38713, GSE47908, GSE73661, GSE114527, GSE87466 and are

represented by grey dots. The summary fold change is displayed with a blue line indicating

downregulation, while a red line indicates upregulation. A $\log_2$-fold-change of 0.5 was used as a

threshold for significance.

Figure 5. PIWIL2 suppresses pro-tumorigenic phenotypes. A) qPCR of PIWIL2 comparing PIWIL2

KO to Caco2 wild type (WT). Data were analyzed and normalized to 18S ribosomal RNA by ΔΔCt

calculations to calculate fold change. (n=3 biological replicates, mean ± SE, t-test *p=0.0309). B)

Quantification of the upper band of PIWIL2 by western blot to confirm the knockout (n=3 biological

replicates, mean ± SE, t-test **p = 0.0025). C) Representative western blot of total PIWIL2 and β-actin

(Actin) loading control in Caco2 WT and PIWIL2 KO cells. The band at the expected molecular weight

of full length PIWIL2 of 109 kDa was quantified and indicated by a red arrow. D) Caco2 WT and PIWIL2

KO cells were infected with an LZRS empty vector (LZRS), and PIWIL2 KO cells were infected with a

FLAG-tagged PIWIL2 construct (PIWIL2-FLAG) to show rescue of the upper band of PIWIL2 by western

blot (red arrow), FLAG, and β-actin (Actin) E) xCELLigence cellular impedance assay to examine the

proliferation rates of Caco2+LZRS, PIWIL2 KO+LZRS and PIWIL2 KO+PIWIL2-FLAG cells in 2-

dimensions (2D) (n=3 biological replicates, two-way repeated measures ANOVA with Geisser-

Greenhouse correction and Bonferroni correction for multiple comparisons (Caco2+LZRS vs PIWIL2

KO+LZRS **p<0.01, Caco2+LZRS vs PIWIL2 KO+PIWIL2-FLAG *p<0.05, PIWIL2 KO+LZRS vs

PIWIL2 KO+PIWIL2-FLAG n.s). F) Representative images of Caco2+LZRS, PIWIL2 KO+LZRS and

PIWIL2 KO+PIWIL2-FLAG cells grown in anchorage independent conditions after 7 days. G) Area, H)

Perimeter I) Diameter and J) Circularity of Caco2+LZRS, PIWIL2 KO+LZRS and PIWIL2 KO+PIWIL2-

FLAG spheroids (*p<0.05). For the low attachment assay n=3 biological replicates, mean ± SE is

displayed, and a one-way ANOVA with a Bonferroni correction for multiple comparisons was used for
statistical analysis between groups.

Figure 6. RNA sequencing analysis shows that PIWIL2 regulates a set of non-canonical piRNAs
targeting transposons. A) Size distribution of RNA reads obtained through short RNA sequencing of
Caco2 WT and PIWIL2 KO cells, after aligning reads to both the piRNA cluster and the repBase
databases. Nucleotide lengths is shown on the x axis, whereas the fraction of the respected reads
against the total sequencing reads are shown on the y axis. B) Nucleotide enrichment of the 5' end of
the short RNA reads identified in the piRNA cluster and the repBase databases. C-D) Scatter plots of
the antisense (meaning complementary to the respected TEs indicated in the graph) and of the sense
(meaning the same as the respected TEs indicated in the graph) short RNA (piRNA-like) sequence
reads as counts per million (CPM) from Caco2 WT (x axis) and PIWIL2 KO (y axis) cells. In red are
examples of specific LINE-1 element targets that are present in both the antisense and sense piRNA
sequences in Caco2 WT cells but absent in PIWIL2 KO cells. All sequencing results are normalized
average reads from n=3 biological replicates. Source data can be found in Table S2. E) qPCR of L1RE1
comparing PIWIL2 KO to Caco2 WT. Data were analyzed and normalized to 18S ribosomal RNA by
$\Delta\Delta$Ct calculations to calculate fold change. (n=3 biological replicates, mean ± SE, t-test *p = 0.0267)

Figure 7. PIWIL2 depletion increases LINE1 activity. A) Representative fluorescent images of Caco2
WT or PIWIL2 KO cells that were transfected with either a constitutively active GFP reporter (pCEP4+),
a LINE1 retro-transposition defective reporter (LINE1 inactive (-)) or a LINE1-GFP reporter plasmid that
expresses GFP only when LINE1 retro-transposition occurs. B) Quantification of the average % positive
GFP expressing cells normalized to transfection efficiency for each cell line. (n = 4 biological replicates
with 3 fields taken per condition and averaged, mean ± SE, two-way ANOVA with Bonferroni correction
for multiple comparisons, PIWIL2 KO LINE1 inactive (-) vs. PIWIL2 KO LINE1 GFP *p = 0.0212,
Caco2 WT LINE1 inactive (-) vs. PIWIL2 KO LINE1 GFP **p = 0.0076, Caco2 WT LINE1 GFP vs.
PIWIL2 KO LINE1 GFP **p = 0.0091).

Figure 8. PIWIL2 depletion promotes DNA damage via LINE-1 transposon activation. A) γH2AX
increases in PIWIL2 KO cells compared to Caco2 WT cells by immunofluorescence. Representative
confocal images shown B) Quantification of γH2AX foci per cell (n = 9 fields (3 biological replicates, 3
fields per replicate), mean ± SE, t-test *p = 0.0151) C) Representative western blot of total H2AX,
phosphorylated γH2AX, and β-actin (Actin) loading control in Caco2 and PIWIL2 KO cells. D) LINE-1
GFP reporter assay staining for eGFP and γH2AX co-localization as indicated by white arrows. E)
Quantification of the average % positive GFP and γH2AX expressing cells normalized to transfection

efficiency for each cell line. (n = 3 biological replicates with 3 fields taken per condition and averaged,

mean ± SE, two-way ANOVA with Bonferroni correction for multiple comparisons, Caco2 WT LINE1

inactive (-) vs. PIWIL2 KO LINE1 GFP *$p = 0.0351$.

Figure S1. PIWIL2 is highly expressed in normal somatic tissues particularly of gastrointestinal

origin. A) RNA-seq data of normal tissues (project PRJEB4337) for PIWIL2, derived from NCBI and

from the related PIWIL2 entry (Gene ID: 55124). Mean RPKMs (Reads Per Kilobase of exon per Million

mapped reads) and error bars are shown from n=2 (duodenum, ovary, pancreas, urinary bladder), n=3

(adrenal, appendix, brain, endometrium, esophagus, fat, gall bladder, liver, salivary gland, skin,

stomach), n=4 (bone marrow, heart, kidney, placenta, prostate, small intestine, spleen, thyroid), n=5

(colon, lung, lymph node) samples per indicated tissue. B) The UALCAN web-based tool was used to

perform analysis of TCGA RNA-seq data to determine PIWIL2 expression across multiple normal tissue

samples. Box plots show median log2TPM (transcripts per kilobase million), upper-lower quartile, and

maximum-minimum distribution. Tumor types and number of samples queried: bladder urothelial

carcinoma (BLCA; n=19), breast invasive carcinoma (BRCA; n=114), cervical squamous cell carcinoma

and endocervical adenocarcinoma (CESC; n=3), cholangiocarcinoma (CHOL; n=9), colon

adenocarcinoma (COAD; n=41), esophageal carcinoma (ESCA; n=11), glioblastoma multiforme (GBM;

n=5), head and neck squamous cell carcinoma (HNSC; n=44), kidney chromophobe (KICH; n=25),

kidney renal clear cell carcinoma (KIRC; n=72), kidney renal papillary cell carcinoma (KIRP; n=32),

liver hepatocellular carcinoma (LIHC; n=50), lung adenocarcinoma (LUAD; n=59), lung squamous cell

carcinoma (LUSC; n=52), pancreatic adenocarcinoma (PAAD; n=4), prostate adenocarcinoma (PRAD;

n=52), pheochromocytoma and paraganglioma (PCPG, n=3), rectum adenocarcinoma (READ; n=10),

sarcoma (SARC; n=2), skin cutaneous melanoma (SKCM; n=1), thyroid carcinoma (THCA; n=59),

thymoma (THYM; n=2), stomach adenocarcinoma (STAD; n=34), and uterine corpus endometrial

carcinoma (UCEC; n=35). Tissues showing the highest expression in both datasets are highlighted in

red boxes and are of gastrointestinal origin.

Figure S2. PIWIL2 is not differentially expressed in colon adenomas compared to normal

samples. The colon adenoma datasets used are available from the Gene Expression Omnibus (GEO):

GDS2947/239303_at, GDS2947/220686_s_at and GDS2947/217421_at. N=32 samples normal and

adenoma were included in all cases; a Welch's t-test was used to compare normal vs colon adenoma

samples within a dataset. The box and whisker plots show the median, interquartile range, minimum

and maximum values of PIWIL2 expression for each dataset, displayed in arbitrary units between

datasets.

Figure S3. PIWIL2 KO clone selection and mutation characterization. A) Representative western blots of the screening process to identify PIWIL2 knockout clones, after transfection of Caco2 cells with CRISPR/Cas9 and a gRNA targeting PIWIL2. Clone 11 (indicated by *) was selected for subsequent studies as it was the only clone to survive selection and maintenance in cell culture. B) Schematic of the genomic region including exon 14 of PIWIL2 that was targeted by the gRNA, where highlighted are the genomic alterations that we identified after DNA sequencing of clone 11: a nucleotide substitution (G to A), altering the splicing target site and a large deletion within the intron. The PCR primers used to amplify and sequence the specific region are also indicated. C) Alignment of the DNA sequencing results between the WT cells and the PIWIL2 KO cells, where we identified both an unaffected allele (allele 2), as well as the actual modified allele (allele 1); modifications are indicated in red boxes.

Figure S4. Depletion of PIWIL2 by transient shRNA knockdown exhibits slowed growth in 2-dimensions (2D). A) qPCR of PIWIL2 comparing Caco2 cells treated with a Non-Target shRNA or shRNA targeting PIWIL2. Expression was normalized to 18S ribosomal RNA and fold change was calculated by ΔΔCt calculations. (n=3 biological replicates, mean ± SE, t-test *p=0.0243). B) Representative western blot of Caco2 cells treated with a Non-Target shRNA or shRNA targeting PIWIL2. C) xCELLigence cellular impedance assay to examine the proliferation rates of Caco2 cells treated with a Non-Target shRNA or shRNA targeting PIWIL2 (n=3 biological replicates, two-way repeated measures ANOVA with a Bonferroni correction for multiple comparisons, *p=0.0247).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

B

Figure S1

Figure S2

Figure S3

Figure S4