Xist Repeat A coordinates an assembly of SR proteins to recruit SPEN and induce gene silencing

Jackson B. Trotman[1,2,3], Alessandro Porrello[2,3], Scott A. Schactler[7,8,10], Lorenzo E. DeLeon[7,8,10], Quinn E. Eberhard[1,2,3,5], Samuel P. Boyson[1,2,3,4], Zhiyue Zhang[1,2,3,11], David M. Lee[1,2,3,4,12], Sarah E. Kirik[1,6,13], Nadia Sultana[14,15,16], Son N. Nguyen[14,15], Michael-Claude G. Beltejar[7,9], Min Kyung Shinn[17], Shao-En Ong[7,9], Maria P. Gonzalez-Perez[14,15], Scott A. Shaffer[14,15], Daniel Dominguez[1,2,3], David M. Shechner[7,8,10], J. Mauro Calabrese[1,2,3,18]

[1] Department of Pharmacology,

[2] RNA Discovery Center,

[3] Lineberger Comprehensive Cancer Center,

[4] Curriculum in Genetics and Molecular Biology,

[5] Curriculum in Bioinformatics and Computational Biology,

[6] Department of Biochemistry and Biophysics,

University of North Carolina, Chapel Hill, NC 27599

[7] Department of Pharmacology,

[8] Institute of Stem Cell and Regenerative Medicine,

[9] Department of Genome Sciences,

University of Washington, Seattle, WA 98195

[10] Brotman Baty Institute for Precision Medicine, Seattle, WA 98195

[11] Present address: Computer Science Graduate Program, Princeton University, Princeton, NJ 08544

[12] Present address: Dechert LLP, Boston, MA 02110

[13] Present address: Department of Integrative Immunobiology, Duke University, Durham, NC 27710

[14] University of Massachusetts Chan Medical School, Mass Spectrometry Facility, Shrewsbury, MA 01545

15 University of Massachusetts Chan Medical School, Department of Biochemistry and Molecular Biotechnology, Worcester, MA 01605

16 Present address: AbbVie Bioresearch Center, Worcester, MA 01605

17 Department of Biomedical Engineering and Center for Biomolecular Condensates, Washington University in St. Louis, St. Louis, MO 63130

18 Lead contact; jmcalabr@med.unc.edu

ABSTRACT

The lncRNA *Xist* silences gene expression by recruiting the protein SPEN through its 5′-proximal Repeat A domain. How Repeat A recruits SPEN, how SPEN enacts silencing, and why Repeat A is required for *Xist* biogenesis in addition to silencing remain unclear. We find that sequences in Repeat A critical for SPEN recruitment, silencing, and *Xist* biogenesis directly bind SR-rich splicing factors and not SPEN. SRSF1, one such factor, is necessary and sufficient for SPEN recruitment to Repeat A. SPEN and SR protein-binding motifs in Repeat A enable *Xist*'s association with many proteins, including those involved in m^6A methylation, RNA turnover, and transcriptional elongation. Our results reveal an unexpectedly essential role for splicing factors in coordinating silencing by SPEN and suggest a unifying model for the origin of Repeat A, the function of SPEN, and their roles in *Xist* biogenesis and silencing.

Keywords: *Xist* RNA, X chromosome inactivation, long noncoding RNA, gene silencing, transcription regulation, RNA splicing, RNA methylation, RNA binding protein, SR protein, nuclear speckle

INTRODUCTION

Xist is an 18-kilobase (kb) long noncoding RNA (lncRNA) that orchestrates X-chromosome inactivation, a regulatory process that silences transcription across an entire chromosome early during the development of female eutherian mammals. *Xist* and many of the genes it regulates are essential for health and homeostasis, and their dysregulation can cause human disease. In the decades since its discovery, *Xist* has served as an exemplary model to dissect mechanisms of lncRNA-mediated gene regulation.[1]

Xist is distinguished by a unique sequence organization that contains several conserved tandem repeat domains embedded within exons of exceptional length (~10 kb).[2-6] The most intriguing of these may be Repeat A, a 450-nucleotide (nt) domain located near the 5′ end of *Xist*. Repeat A is comprised of eight tandemly arrayed copies of a ~50-nt monomer that itself is made up of a degenerate, structurally flexible U-rich spacer followed by a GC-rich "core" that is linearly conserved across eutherians.[7,8] Repeat A is essential for *Xist*-induced gene silencing, a function that depends on its interaction with the RNA-binding protein SPEN.[9-12] In turn, interactions between Repeat A-bound SPEN, histone deacetylates, and other histone-modifying enzymes have been proposed to mediate *Xist*-induced silencing.[10,13]

However, while deletion of SPEN or Repeat A results in a near-complete loss of *Xist*-induced silencing, inhibition or deletion of SPEN-associated histone deacetylates, including HDAC3, have only minor effects.[13-15] This suggests that Repeat A and SPEN may engage yet-to-be-defined proteins to induce silencing.[16] Indeed, although SPEN binds Repeat A through its U-rich spacers,[17-19] chemical probing suggests that sequence motifs across Repeat A exist in single-stranded states and bind proteins (Figure S1A).[6,8] Moreover, previous work has shown that in the context of a synthetic, consensus Repeat A sequence embedded in an *Xist* cDNA transgene, mutations of the GC-rich cores – but not mutations or truncations of the U-rich spacers – impair silencing as measured indirectly by a cell-survival assay.[20,21] Perplexingly, the same GC-rich core mutations qualitatively had no impact on SPEN binding *in vitro*.[11,16]

In addition to or perhaps conflated with its role in gene silencing, Repeat A also harbors silencing-independent functions. Most notably, Repeat A deletion reduces *Xist* abundance in a manner that depends at least partly on SPEN.[22-29] Relatedly, in the context of a 2-kb transgene, Repeat A enables readthrough of downstream polyadenylation signals, an activity that depends on its GC-rich cores and not SPEN.[26] Lastly, Repeat A associates with serine-and-arginine-rich (SR) proteins *in vivo* and *in vitro*, and its deletion can alter *Xist* splicing, linking Repeat A to the

splicing machinery.[2,5,26-28,30] These data highlight the need to study Repeat A in quantitative contexts that can report on all aspects of *Xist* biology. In turn, studying Repeat A might provide insights into the function of SPEN, a deeply conserved protein that is recurrently mutated in cancers and neurodevelopmental disorders yet whose molecular functions remain incompletely characterized.[16,31-40]

We sought to reconcile the different functions attributed to Repeat A by mutationally dissecting its endogenous sequence in a full-length *Xist* transgene including all endogenous exons and introns. We demonstrate that Repeat A strongly suppresses aberrant splicing within *Xist* exon 1 and that SR protein-binding motifs in the GC-rich core and at least one SR protein, SRSF1, are critical for recruitment of SPEN and *Xist*-induced gene silencing. We likewise find that the N-terminal region of SPEN is an SR protein sensor that facilitates its recruitment to both Repeat A and SR protein-binding motifs transcriptome-wide. Both SPEN and SR protein-binding motifs synergistically enable *Xist* to associate with the m^6A methylation, RNA turnover, and transcriptional elongation machineries. Our data uncover an unexpected connection between canonical splicing factors and silencing by SPEN and suggest a unifying model to explain the dual roles of SPEN and Repeat A in *Xist* biogenesis and silencing. The sensing of SR protein-rich protein assemblies may be a general mechanism by which SPEN is targeted to RNA.

RESULTS

Sequences critical for *Xist*-induced silencing, *Xist* production, and *Xist* splicing fidelity are located within the GC-rich cores of Repeat A

To clarify how sequence elements throughout Repeat A impact *Xist* biogenesis and function, we sought to evaluate the effects of new and previously investigated[11,20,21] mutations in the context of a full-length doxycycline (dox)-inducible *Xist* transgene that includes its exons and introns. *Xist* transgenes were inserted as a single copy into the C57BL/6 (B6) allele of the *Rosa26* locus of an F1-hybrid mouse embryonic stem cell (ESC) line.[26] In these ESCs, *Xist* expression represses genes on the B6 copy of chromosome 6 (chr6) without affecting expression on the CAST/EiJ (CAST) chromosome (Figure 1A).[26,41]

Using this system, we made two separate excisions of Repeat A and ten smaller mutations or deletions within each monomer of Repeat A (Figure 1B; Figure S2). Additionally, we made a deletion as well as an insertion downstream of Repeat A (Figure 1B). The deletion encompassed

a 742-nt region immediately downstream of Repeat A that is predicted to form at least three stable RNA secondary structures ("ss234")[8,42] and harbors the highest N6-methyladenosine (m^6A) density within *Xist*.[25] This region also contains the only transcribed CpG island in *Xist*, perhaps related to its requirement for readthrough of proximal polyadenylation signals in a 2-kb *Xist* transgene.[26,43,44] Separately, we inserted a synthetic intron[45] just downstream of Repeat F. Previous work suggested that ectopic splicing of an intron downstream of Repeat A may attenuate *Xist*-induced silencing,[46] and we sought to test that hypothesis. Each alteration was confirmed by whole-plasmid sequencing before inserting into *Rosa26*. Four independent wild-type (WT) *Xist* clonal lines and two independent clonal lines for each *Xist* mutant were selected and rendered dox-inducible via piggyBac-mediated insertion of the reverse tetracycline-controlled transactivator (rtTA).[26] In parallel, we prepared four independent clonal lines of empty-cargo control ESCs containing the same dox-inducible promoter but no lncRNA sequence inserted.

We first defined a high-confidence set of *Xist*-repressed genes on chr6. We performed RNA-seq for each wild-type *Xist* and empty-cargo cell line treated with dox for 3 days (Figure 1A). Comparing allelic expression between the two genotypes, we identified 204 chr6 genes significantly repressed by *Xist* (adjusted *p*-value < 0.001).[47] As expected, *Xist*-repressed genes were located along the length of chr6, and were, on average, repressed by about 50% relative to empty-cargo control (Figure S1C). Replicates across genotypes were highly reproducible (Figure S1D,E).

We next performed RNA-seq for our panel of mutants to assess effects on *Xist*-induced gene silencing, *Xist* production, and *Xist* splicing. This revealed a broad range of phenotypes, many of which were unanticipated. As expected, Repeat A deletion led to a near-complete loss of silencing and a ~50% reduction in *Xist* abundance (Figure 1C,D, Figure S1F). Repeat A deletion also caused previously reported aberrant splicing events that excise the bulk of *Xist* exon 1; an estimated ~15-20% of *Xist* transcripts in Repeat A deletion mutants lacked a functional exon 1 (Figure 1H,I).[28,48] Surprisingly, deletion of the U-rich spacers – Repeat A's presumed SPEN-binding region – only intermediately affected silencing, did not alter *Xist* abundance, and minimally altered *Xist* splicing, whereas deletion of the GC-rich cores phenocopied deletion of Repeat A in all respects (Figure 1C,D,I). Mutations predicted to preserve or disrupt possible secondary structure within the GC-rich cores[20] did not impact *Xist* abundance and minimally impacted splicing but did have moderate ("preserve" mutant) to severe ("disrupt" mutant) effects on silencing (Figure 1C,D,I). Strikingly, mutation of a predicted single-stranded AWCG motif previously shown to bind splicing factor SRSF1,[27] deletion of a GG dinucleotide downstream of

the AWCG, or a combination of both mutations in the same *Xist* gene phenocopied Repeat A deletion in terms of reduced *Xist* abundance and silencing. These mutations also altered exon 1 splicing but to a much lesser degree than Repeat A or GC-rich core deletion (Figure 1C,D,I). Mutation of U19 or deletion of the CCUGC motif had severe defects on silencing without reducing *Xist* abundance; U19 mutation altered splicing to a similar degree as the AWCG mutation, whereas CCUGC deletion did not (Figure 1C,D,I). Mutation of U23 weakly reduced silencing, weakly affected splicing, and did not alter *Xist* abundance (Figure 1C,D,I). Deletion of ss234 affected both *Xist* abundance and silencing and had no effect on splicing (Figure 1C,D,I). Insertion of the ectopic intron phenocopied wild-type *Xist* in all respects (Figure 1C,D,I); *Xist*-induced silencing was unaffected by variably efficient ectopic intron splicing (Figure S1D,G).

Leveraging our panel of mutants, we sought to determine the extent to which reductions in silencing activity could be attributed to concomitant reductions in *Xist* abundance. We performed RNA-seq in ESCs expressing lower levels of wild-type *Xist* and demonstrated a linear relationship between silencing and *Xist* abundance (Figure 1E-G, Figure S1E), consistent with expectations set by prior works.[41,49,50] Next, plotting silencing activity as a function of *Xist* RNA levels for all genotypes and expression conditions revealed that every mutant except two (Δss234 and the intron-insertion) affected silencing to a greater degree than could be attributed solely to reduced *Xist* abundance (Figure 1G). These experiments confirm a role for Repeat A in silencing beyond its role in promoting *Xist* production.

Taken together, we observed a striking non-uniformity of phenotypes upon deletion or mutation of Repeat A. Repeat A was critical for *Xist*-induced gene silencing and for its production. However, different alterations disrupted silencing and *Xist* abundance to different extents, with silencing affected to a greater degree than would have been expected from reduced *Xist* levels. Almost every alteration also caused at least some level of aberrant splicing-out of *Xist* exon 1, indicating that, in addition to its roles in silencing and promoting *Xist* abundance, Repeat A is a potent local suppressor of splicing. We also observed that the ss234 region downstream of Repeat A is essential for promoting wild-type levels of *Xist* and thus wild-type levels of silencing. The sequence elements most critical for *Xist*-induced silencing, *Xist* production, and suppression of aberrant splicing are located in the GC-rich cores of Repeat A and not its U-rich spacers.

SPEN association with Repeat A *in vivo* depends on sequences that it does not bind *in vitro*

SPEN is recruited by Repeat A and is required for *Xist*-induced silencing.[9-11,13] However, *in vitro*, SPEN directly binds the U-rich spacers of Repeat A and not the GC-rich cores critical for silencing.[11,17,18] To address this apparent discrepancy, we performed RNA immunoprecipitation from formaldehyde-crosslinked cells, followed by quantitative PCR (RIP-qPCR),[41] to measure the extent to which SPEN associates with each *Xist* mutant *in vivo* (Figure 2A). Because formaldehyde induces RNA-protein and protein-protein crosslinks, RIP enables detection of both direct and indirect (i.e., protein-bridged) RNA-protein interactions (Figure 2A). We confirmed that our SPEN antibody retrieved Repeat A in a SPEN-dependent manner by performing RIP-qPCR with wild-type and *Spen*-knockout ESCs (Figure 2B; Figure S3A,B).[51] In parallel, we examined SPEN association across our panel of mutants, normalizing signal by the amount of *Xist* RNA in each corresponding input (Figure 2B). Consistent with the necessity of SPEN for *Xist*-induced silencing, we observed a remarkably strong correlation between SPEN RIP recovery and extent of *Xist*-induced gene silencing across all mutants (Figure 2C; Pearson's r = -0.91; p = 5 x 10^{-6}). The mutants associating with the least amount of SPEN *in vivo* were ΔRepA; ΔGC-core; ΔGG16,17; and the AWCG+ΔGG mutant (Figure 2B). Recovery of a non-specific control region was uniform across genotypes (Figure 2B). Therefore, the sequences most essential for SPEN-Repeat A association *in vivo* are not U-rich but are instead contained within its GC-rich cores.

To determine whether GC-rich sequences can direct SPEN association with RNAs other than *Xist*, we sequenced RNA recovered by SPEN RIPs from wild-type and *Spen*-knockout SM33 ESCs (Figure 2B). We identified 69,045 genomic regions ("peaks") enriched in SPEN association. This revealed that Repeat A is by far the most-recovered RNA target of SPEN and also the most specifically recovered from wild-type versus *Spen*-knockout cells (Figure 2D). Nonetheless, we identified thousands of RNA peaks whose wild-type recovery was at least two times greater than knockout across two independent experiments. Analysis of peaks with STREME[52] revealed enrichments for multiple CUG- and GA-rich motifs (Figure 2E, Figure S3C). A single poly-U motif, ranked 13th, was also enriched. These results suggest that, while SPEN can associate with U-rich sequences *in vivo*, it more frequently associates with G- and C-rich sequences, consistent with our *Xist*-mutant RIP results.

Next, we examined how the different mutations impact the relative binding affinity between Repeat A RNA and SPEN *in vitro*. SPEN directly binds RNA through its RNA-recognition motifs 2 through 4 (RRM2-4), located near its N-terminus.[19] We therefore used a modified version of the RNA-Bind-n-qPCR assay (RBNQ) to measure the equilibrium binding affinity between bacterially purified SPEN RRM2-4 and *in vitro* transcribed Repeat A RNA (Figure 2F; Figure S3D,E).[53] SPEN

RRM2-4 bound to wild-type Repeat A RNA cooperatively and with a K_d of 190 nM, consistent with prior studies (Figure 2G).[18,19,54] We next performed RBNQ across our panel of mutant RNAs using a concentration of SPEN RRM2-4 that yielded ~50% recovery of wild-type RNA, enabling detection of both increases and decreases in binding (Figure 2F). Relative to wild-type Repeat A, U-rich-spacer deletion abrogated binding, consistent with expectations from prior *in vitro* experiments (Figure 2H;[11,17]). However, GC-rich-core deletion increased binding, and mutations within the cores either increased binding, made no difference, or reduced binding slightly but not to the same degree as deletion of the U-rich spacers (Figure 2H). Together, these findings demonstrate that the sequences within Repeat A that directly bind SPEN *in vitro* (its U-rich spacers) are not the ones most critical for SPEN association *in vivo* (its GC-rich cores).

Motif-dependent association between Repeat A and SR proteins *in vitro* and *in vivo*

To rationalize the sharp discordance between SPEN's preferred motifs *in vitro* and *in vivo*, we hypothesized that recruitment of SPEN to Repeat A *in vivo* might require proteins that interact with the GC-rich cores. To identify candidate interacting proteins, we modified a previous approach[2] in which internally biotinylated wild-type or mutant Repeat A RNA was incubated with ESC nuclear extracts, crosslinked with UV light, and streptavidin-purified prior to being analyzed by mass-spectrometry-based proteomics (Figure 3A, Figure S4A,B). We also used proteomics to quantify the proteins detectable in our nuclear extracts. In total, we identified 644 proteins reproducibly recovered with Repeat A RNA but not no-RNA control (Figure S4B; Figure S4C; Table S1). SPEN was not recovered, presumably due to its low abundance in ESCs (2766[th] out of 2826 detected nuclear extract proteins; Table S1).

Of the 644 Repeat A-recovered proteins, 124 were significantly enriched with wild-type RNA relative to ΔGC-core RNA ($p < 0.05$, two-sided t-test; Figure 3B). Gene ontology analysis of the GC-core-interacting proteins revealed an enrichment in RNA processing and splicing factors, including components of the spliceosome and nuclear speckles (Figure 3C). We noted distinct motif-dependent enrichment patterns of SR proteins, a protein family distinguished by disordered serine- and arginine-rich domains that plays central roles in transcription and RNA processing (Figure 3D).[55,56] Indeed, relative to wild-type Repeat A, several SR proteins displayed reduced recovery upon deletion of the GC-rich cores and enhanced recovery by RNA containing only GC-rich cores (the ΔU-spacer mutant), strongly suggesting interactions with the GC-rich cores (Figure 3D, cluster 2).

We next sought to determine whether the proteins identified to interact with Repeat A *in vitro* associate with Repeat A *in vivo*. Using RIP-seq, we analyzed a panel of 22 RNA processing factors, including components of the spliceosome and nuclear speckle, with a focus on SR proteins. We analyzed SPEN as a positive control, and nonspecific IgG and anti-FLAG antibodies as negative controls. Repeat A association was enriched at least two-fold over negative controls for five SR proteins: SRSF1, SRSF2, ZC3H13, ZC3H18, and SCAF4 (Figure 3E,F; Figure S4D). RIPs for other proteins more closely resembled negative controls, indicating poor antibody performance or non-specific association with *Xist*. We conclude that *in vivo*, Repeat A associates with SR proteins that *in vitro* interact with Repeat A in a GC core-dependent manner.

SPEN associates with nuclear speckle components and other SR proteins via its N-terminus

Next, we investigated whether SPEN interacts with SR proteins *in vivo*. We first deleted *Spen* from RMCE-*Xist* WT ESCs and stably re-expressed SPEN from a cDNA harboring an N-terminal Halo tag and a C-terminal V5 tag, enabling immunoprecipitation and western blot detection (Figure S5A,B). Consistent with previous observations,[25] *Spen* deletion caused a 50% decrease in *Xist* levels (Figure 4A) and a loss of *Xist* silencing activity beyond that expected for comparable lower levels of wild-type *Xist* (Figure 1G, Figure 6G, Figure S5C). While ~18% of *Xist* transcripts displayed aberrant splicing-out of *Xist* exon 1 upon Repeat A deletion (Figure 1H,I), only 4% did so upon *Spen* deletion, indicating that SPEN plays a minor role in preserving *Xist* splicing fidelity (Figure S5D). Expressing Halo-SPEN-V5 in *Spen* knockout cells restored *Xist* abundance and splicing fidelity to wild-type levels and rescued silencing activity, albeit only partially, possibly due to the production of SPEN at lower-than-endogenous levels (Figure 4A; Figure S5C,D).

To identify SPEN-interacting partners, we immunoprecipitated Halo-SPEN-V5 from formaldehyde-crosslinked cells with RIP wash conditions and analyzed recovered proteins via mass spectrometry. Parallel immunoprecipitations from *Spen* knockout cells expressing no Halo-tagged construct ("no rescue") served as a specificity control. We identified 37 proteins whose signal was enriched at least two-fold over no-rescue control in both replicates, while also observing that SPEN peptides were only detected from its N-terminal half (Figure 4B, Table S2; not shown). Gene ontology analysis revealed significant enrichments with many of the same terms as our Repeat A affinity proteomics, including those related to splicing, RNA processing, and nuclear speckles (Figure 3C, Figure 4C). The enrichment of nuclear speckle components and

SR proteins was especially striking, with 8 of the top 19 SPEN-interacting proteins belonging to one or both categories (Figure 4B). One such protein, SRSF1, is an SR and nuclear speckle protein previously found to interact with SPEN,[55-58] which we confirmed via co-immunoprecipitation (Figure 4D).

We next examined how individual parts of SPEN contribute to its interaction with Repeat A and other proteins. SPEN is a ~400-kilodalton protein with 4 N-terminal RRMs and a C-terminal SPOC domain thought to mediate interactions with protein effectors (Figure 4E).[13] The rest of SPEN primarily consists of 14 intrinsically disordered regions (IDRs) defined by MobiDB (Figure 4E).[59] Reminiscent of canonical SR proteins, SPEN contains a bona fide RS domain within its RRM-adjacent IDR2.[55,60] IDR4, downstream of RRM4, is also arginine-rich, with many RE/ER dipeptides that can mimic phosphorylated RS dipeptides (Figure S5E,F).[61] These regions were of particular interest due to the roles that RS domains play in mediating interactions among SR proteins.[55,56] We constructed a panel of 12 Halo-and-V5-tagged SPEN deletion mutants and analyzed association with Repeat A via RIP-qPCR (Figure 4E). As expected, wild-type SPEN recovered Repeat A robustly relative to no-rescue and EGFP-Halo controls (Figure 4F). Deletion of RRM1-RRM4 completely abrogated SPEN's association with Repeat A, as did deletion of RRM2-4, consistent with previous results.[13,50,62] In contrast, deletion of the RRM1-IDR2 region had no impact on Repeat A association, nor did deletion of IDR2, the smaller RS domain, IDR4, or a combined deletion of IDR2 and IDR4. Intriguingly, deletion of the SPOC domain caused a ~50% increase in SPEN-Repeat A association. Lastly, we found that RRM2-4 alone was sufficient to associate with Repeat A but at a level ~3-fold lower than the larger RRM1-4 construct. This difference was mostly attributable to RRM1, although the addition of IDR2 to RRM2-4 also caused a minor increase in Repeat A association (Figure 4F). These results confirm that RRMs 2-4 are essential for Repeat A recognition and demonstrate that SPEN's N-terminus (RRM1-IDR2) promotes Repeat A association in a truncated version of SPEN and likely in combination with downstream IDRs in the context of full-length SPEN.

Previous work demonstrated no difference in Repeat A binding affinity between RRM1-4 and RRM2-4, implying that SPEN's N-terminal region functions outside of RNA-binding.[19] To investigate, we used Halo immunoprecipitation and proteomics to compare protein association between the RRM1-4 and RRM2-4 constructs. Likely due to higher expression of these constructs than full-length SPEN (Figure S5B), we identified 146 and 106 proteins enriched with RRM1-4 and RRM2-4, respectively, at least two-fold over no-rescue control in two independent replicates (Table S2). GO analysis of the seventy-five proteins enriched with RRM1-4 at least two-fold over

RRM2-4 revealed strong enrichments for terms related to splicing and nuclear speckles (Figure S5G). Indeed, compared to RRM2-4, RRM1-4 recovered significantly higher levels of nuclear speckle and SR proteins – including known Repeat A interactors RBM15, U2AF1, U2AF2, SRSF1, and SRSF2 – but not paraspeckle or nucleolar proteins (Figure 4G,H).[63]

In parallel, we performed RIP-seq to determine motifs enriched in the top non-*Xist* peaks for Halo-tagged full-length SPEN, RRM1-4, and RRM2-4 transcriptome-wide. Like endogenous SPEN (Figure 2E), full-length Halo-SPEN-V5 associated most strongly with GA- and GC-rich, SR protein-like binding motifs (Figure 4I). Notably, while RRM1-4 similarly enriched for SR protein-like binding motifs, RRM2-4 enriched most strongly for poly-U motifs, consistent with the latter's direct RNA-binding preference (Figure 2H, Figure 4I). We further identified the top RNA targets of each SPEN variant by totaling peak signal per gene, revealing that RRM1-4 uniquely associated with the nuclear speckle-associated *7SK* and *MALAT1* RNAs (ranked 1st and 3rd, respectively)[64-66] and, curiously, mRNAs encoding SR-, nuclear speckle-, and splicing-related proteins (e.g., *Srsf1* and *Son*, ranked 11th and 16th, respectively; Table S3). Therefore, relative to RRM2-4, RRM1-4 associates more strongly with SR proteins, SR protein-binding motifs and RNAs, and Repeat A itself.

Lastly, we investigated the proteins genetically associated with SPEN via an orthogonal method, performing "guilt-by-association" analysis of data from the Genotype-Tissue Expression (GTEx) project.[67-70] We identified 129 genes whose expression was significantly correlated with *SPEN* across 49 GTEx tissues (FDR < 0.1, see Methods), implying that they may function within the same or related pathways (Figure 4J, Table S4). Additional guilt-by-association analyses of normal and tumor tissue data from The Cancer Genome Atlas (TCGA) identified many of the same genes and pathways (Figure 4K, Figure S5H, Table S4), including *MAML1* and *KMT2D*, genes previously linked to SPEN in the context of NOTCH signaling.[71,72] Genes encoding SPEN-interacting proteins (merged list of WT, RRM1-4, and SPOC domain interactors; Table S2)[13] were significantly enriched in *SPEN*-correlated genes (*p* = 0.0004, two-tailed binomial test), further underscoring our ability to identify meaningful connections to SPEN via guilt-by-association. Consistent with our findings above, we observed strong enrichment for GO terms associated with splicing and nuclear speckles, along with terms and proteins related to transcriptional activation and repression (Figure 4J-L, Figure S5H). These include the top hit, SCAF4, an RS-domain-containing transcriptional antiterminator enriched over Repeat A (Figure 3E);[73] BRD4, a critical transcriptional elongation factor;[74] TRIM28 and NuRD (GATAD2B, MTA2), epigenetic modifiers previously linked to SPEN- and Repeat A-dependent silencing;[13,75] ZC3H18, an SR protein and

RNA exosome adapter enriched over Repeat A (Figure 3E);[76,77] the SR proteins ACIN1, SRSF4, and U2AF2; and YTHDC1, a (D/E)R-rich, nuclear speckle-associated m^6A-binding protein and exosome adapter, which is also enriched over Repeat A (Figure 3E,F, Figure 4H, Figure S7A).[78-81]

Thus, proteomic, genomic, and guilt-by-association analyses connect SPEN with multiple aspects of the transcriptional process and nominate the N-terminal region of SPEN as an SR protein sensor.

SRSF1 directly binds a conserved Repeat A motif to recruit SPEN to *Xist in vivo*

To address the possibility that SR-rich splicing factors play a role in recruiting SPEN to Repeat A, and for the following reasons, we focused on SRSF1: (i) SRSF1 interacts with Repeat A *in vitro* and *in vivo*, and, after SPEN, was the most Repeat-A-enriched protein we observed by RIP-seq (Figure 3, Figure S4D);[2,5,27,63,82] (ii) by proteomics, SRSF1 binding was reduced in the silencing-defective ΔGC-core and AWCG+ΔGG mutants (Figure 1D, Figure 3D); (iii) the GA-rich motifs enriched transcriptome-wide in SPEN RIP-seq peaks closely resemble canonical SRSF1 binding motifs (Figure 2E, Figure 4I);[56,83,84] (iv) we and others detect physical interactions between SRSF1 and SPEN (Figure 4A,C,I);[57,58] and (v) Repeat A's CAWCGGG motif resembles a consensus SRSF1 motif determined by SELEX.[30]

We first performed SRSF1 RIP-qPCR across our panel of *Xist* mutants (Figure 5A). We included wild-type and *Spen*-knockout SM33 cells, which revealed that SRSF1 associates with Repeat A two to three times more strongly in the absence of SPEN (Figure 5A). We observed that SRSF1 association was lost or reduced upon deletion of Repeat A, deletion of the GC-rich cores, or mutations within nucleotides 1 and 17 of the GC-rich cores (Figure 1B, Figure 5A). Conversely, SRSF1 association was unchanged by deletion of the U-rich spacers and increased with mutations between GC-rich-core nucleotides 19 and 25 (Figure 1B, Figure 5A). Comparing levels of SRSF1 and SPEN association for each mutant revealed a moderate positive correlation (Pearson's r = 0.47; p = 0.09); however, excluding mutations within GC-rich-core nucleotides 19 and 25 yielded a strong, highly significant correlation (Pearson's r = 0.84; p = 6 x 10^{-4}; Figure 5B). The CCUG(21-24) sequence of the GC-rich cores is a canonical SRSF2-binding motif;[85,86] however, deletion of this sequence did not impact SRSF2-Repeat A association (Figure S6A). Together, these results are consistent with a role for SRSF1 in recruiting SPEN to Repeat A and

suggest an SRSF1- and SRSF2-independent role for the 3' end of the GC-rich cores in recruiting SPEN.

We next determined the extent to which SRSF1 directly binds Repeat A *in vitro* by performing equilibrium RBNQ with SRSF1's RNA-binding domains (RRM1-2; Figure S6B). Like SPEN, SRSF1 bound Repeat A cooperatively, yet with a four-fold stronger K_d than that of SPEN, despite being ~1000-fold more abundant in ESCs (Figure 2G, Figure 5C).[87] Holding the concentration of SRSF1 RRM1-2 constant in binding reactions with Repeat A mutants, we observed patterns similar to *in vivo* SRSF1 association (Figure 5A), with the ΔGC-core, AWCG, and AWCG+ΔGG mutants exhibiting loss of SRSF1 binding (Figure 5D). Interestingly, the ΔGG16,17 mutant, which reduced SRSF1 association by RIP (Figure 5A) and reduced *in vitro* binding in another study,[27] did not reduce binding by RBNQ (Figure 5D). Our results demonstrate that the direct interaction between SRSF1 and Repeat A is strong and dependent on the single-stranded AWCG5-8 motif present in each GC-rich core.

To directly test whether SRSF1 is necessary for recruiting SPEN to *Xist*, we tagged the 5′ end of both *Srsf1* alleles in RMCE-*Xist* WT cells with *FKBP1A^F36V* to enable its degradation upon addition of the small molecule dTAG^V-1 (Figure S6C).[88-90] After 18 hours of dTAG^V-1 treatment, the FKBP1A^{F36V}-SRSF1 fusion protein was no longer detectable by western blot (Figure 5E). To determine the impact of SRSF1 depletion on SPEN-Repeat A recruitment, we treated *FKBP1A^F36V*-*Srsf1* cells with dTAG^V-1 for 18 hours before dox-inducing *Xist* expression for 26 hours and harvesting. Using SPEN RIP-qPCR, we observed that SRSF1 depletion caused a 50% decrease in SPEN-Repeat A association relative to wild-type cells and a 30% decrease relative to untreated *FKBP1A^F36V*-tagged controls, indicating that SRSF1 is necessary for optimal SPEN recruitment to Repeat A (Figure 5F). By RNA-seq, acute depletion of SRSF1 caused a mild ~20% decrease in *Xist* RNA levels and significantly weakened *Xist* silencing activity (Figure 5G,H). While a previous report suggested a role for SRSF1 in maintaining proper *Xist* splicing,[27] we observed no aberrant *Xist* splicing events upon SRSF1 depletion (Figure S6D and not shown). These results reveal SRSF1 is necessary to recruit SPEN to *Xist*.

To determine whether SRSF1 is sufficient to recruit SPEN to Repeat A, we devised an *in vivo* tethering assay in which SRSF1 or other proteins were fused to the MS2 bacteriophage coat protein (MCP), and within *Xist*, Repeat A was replaced with a synthetic sequence consisting of six units, each with a natural U-rich spacer followed by three copies of the MCP-cognate MS2 stem-loop (6x(U-spacer-3xMS2); Figure 5I). Initially, we MCP-tagged full length SRSF1 and compared effects to cells expressing untagged EGFP (Figure S6E). Using SRSF1 RIP-qPCR, we

confirmed successful tethering of the MCP-SRSF1 fusion construct to the 6x(U-spacer-3xMS2) sequence but did not observe a significant increase in SPEN association relative to EGFP control (p = 0.12; Figure 5J, Figure S6F). We speculated that full-length SRSF1 may not robustly recruit SPEN in this system due to steric effects and/or loss of heterotypic cooperativity due to presumed binding sites for multiple proteins in the 25-nt GC-rich core being replaced with monovalent, 95-nt 3xMS2 sequences. We therefore expressed a fusion construct consisting of MCP, an SV40 nuclear localization signal, a 3xFLAG tag, and four concatenated copies of SRSF1's RS domain, reasoning that the 4x-RS domain might better mimic cooperative cofactor recruitment by the GC-rich core (Figure 5F).[56] We additionally tested analogously tagged constructs of EGFP, three copies of the HNRNPK arginine-glycine-rich (RGG) domain, one copy of the 121-amino-acid SRSF2 RS domain, or alternating copies of the SRSF1 and SRSF2 RS domains (Figure S6E). We confirmed tethering of these constructs to *Xist* by FLAG RIP-qPCR (Figure S6G). Strikingly, relative to the other constructs, 4xSRSF1 RS recruited two-to-four times more SPEN to synthetic Repeat A (p = 1 x 10^{-4}; Figure 5J). The 4xSRSF1 RS construct nonetheless recruited four-fold less SPEN than wild-type *Xist* and did not induce gene silencing (Figure 5L, Figure S6H). However, 4xSRSF1 RS tethering also caused a 33% increase in levels of *Xist* RNA over EGFP control, likely a consequence of increased SPEN recruitment (Figure 4A, Figure 5K). Consistent with this, across Repeat A mutants, we observed that *Xist* RNA abundance significantly correlated with levels of association with both SRSF1 and SPEN (Figure S6H,I). In summary, using a reconstructed system, we find that the RS domain of SRSF1 is sufficient to increase levels of *Xist* RNA and recruit SPEN to a version of *Xist* lacking its GC-rich cores. Altogether, our results reveal that SRSF1 is a core component of the Repeat A ribonucleoprotein complex that recruits SPEN to *Xist*.

SPEN and SR protein-binding motifs promote interactions between Repeat A and the m⁶A, RNA turnover, and transcriptional elongation machineries

Our own and previous proteomic analyses detected interactions between SPEN and proteins involved in transcription, splicing, and m⁶A RNA methylation, the last of which has been implicated in the degradation and transcriptional silencing of chromatin-associated RNAs.[13,91,92] Likewise, SR proteins can interact with RNA polymerase II, the splicing and m⁶A machineries, and the exosome adapters ZC3H18 and YTHDC1, which themselves exhibit association with Repeat A (Figure 3E, Figure S7A).[64,76,80,93-95] Our observations that Repeat A and the N-terminus of SPEN are SR-protein-binding modules raised the possibility that Repeat A may use SR proteins and

SPEN to engage with the transcriptional and RNA processing machineries upstream of or in parallel to inducing transcriptional silencing by chromatin modification.

To investigate this possibility, we used *Xist* oligonucleotide-mediated proximity-interactome mapping (O-MAP) to assess how *Spen* deletion impacts the proteins present within the *Xist* microenvironment (Figure 6A).[96] We identified 1802 *Xist*-proximal proteins significantly enriched relative to no-probe controls in either RMCE-*Xist* WT or Δ*Spen* cells: 810 proteins exhibited reduced recovery upon *Spen* deletion, 34 exhibited enhanced recovery, and 958 were unchanged ($p < 0.05$, two-sided t-test; Figure 6B, Table S5). The first group can be considered "SPEN-dependent" for inclusion in the *Xist* microenvironment, whereas the latter two can be considered "SPEN-independent." The large number of SPEN-dependent proteins may reflect, in part, reduced *Xist* spreading and abundance in the absence of SPEN (Figure 4A).[23,25] Consistent with expectations, the well-characterized SPEN-binding protein RBPJ was SPEN-dependent, whereas proteins that bind *Xist* outside Repeat A, including HNRNPK, HNRNPU, PTBP1, and MATR3, were SPEN-independent (Table S5).[97-99] Strikingly, consistent with the notion that Repeat A binds SR proteins upstream of SPEN (Figure 5A), 40 of the 51 SR proteins detected as *Xist*-proximal were SPEN-independent,[55] including SRSF1/3/5/6/7/9, TRA2A/B, U2AF1/2, SCAF1/4/8, SNRNP70, and ACIN1 (Table S5). Conversely, the previously identified Repeat A-interacting proteins RNF20/40 were SPEN-dependent, as were nearly all components of the m^6A methyltransferase complex (METTL3/14, RBM15, WTAP, ZC3H13, VIRMA; Figure 6B).[9,79] Additionally, several components of the RNA exosome (EXOSC2/7/8, MTREX) and RNA polymerase II (POLR2A/C/E/J/H) were SPEN-dependent, along with many repressive and activating transcription factors and epigenetic modifiers (Figure 6B, Table S5). In sum, these data suggest that *Xist*'s association with SR proteins is largely SPEN-independent, whereas *Xist*'s association with the m^6A, RNA exosome, and transcriptional machineries is more SPEN-dependent.

To probe these observations further, we used RIP-qPCR to examine the extent to which proteins involved in RNA metabolism and transcription physically associate with Repeat A in a SPEN- or SR protein-binding motif-dependent fashion. As observed previously by RIP and O-MAP (Figure 5A, Figure 6B), the association of SRSF1 with Repeat A was SPEN-independent but lost with the *Xist* AWCG+ΔGG mutant, a pattern similar for the SR protein ACIN1 (Figure 6C). Repeat A association with the SR protein and spliceosome adapter U2AF2 was independent of both SPEN and the AWCG+ΔGG mutation, consistent with its preference for the U-rich spacers (Figure 3D, Figure 6C). Remarkably, for most of the other proteins we profiled, associations with

Repeat A were dually dependent on SPEN and Repeat A's SR protein-binding motifs. These included HDAC3 and TRIM28, factors previously implicated in silencing by *Xist* and SPEN;[10,13,14,75] m^{6}A methyltransferase complex components RBM15 and ZC3H13; nuclear exosome adapters YTHDC1 and ZC3H18; the transcription-elongation and antitermination factor SCAF4; and elongation-associated, serine-2-phosphorylated RNA polymerase II itself (Figure 6C). In contrast, transcription initiation-associated, serine-5-phosphorylated RNA polymerase II; the transcription elongation factor SUPT6H; and components of the CPSF/CFIm and Integrator complexes did not readily associate with Repeat A beyond IgG controls (Figure 6D).

Several of the proteins we found to associate with Repeat A in a SPEN- or SR protein motif-dependent manner had not previously been linked to *Xist* biology. We therefore performed a targeted siRNA screen to determine whether their depletion would alter *Xist* abundance or *Xist*-induced silencing. Knockdowns were confirmed by western blot, with efficiencies ranging from 63% to 94% (Figure S7B). Knockdowns of four proteins, HDAC3, TRIM28, ZC3H13, and YTHDC1, caused mild but significant reductions in the silencing of *Xist* target genes (Figure 6F); however, excepting HDAC3, weaker silencing could be attributed to reductions in *Xist* abundance (Figure 6E,G). Curiously, knockdown of ACIN1, a component of the ASAP complex linked to SPEN-induced silencing in plants,[100] caused an apparent increase in the abundance of *Xist* without potentiating its silencing function (Figure 6E-G). None of the knockdowns affected *Xist*-mediated silencing to same level as SPEN deletion (Figure 6G, Figure S5C). Together, our data demonstrate that SPEN and SR protein-binding motifs enable Repeat A to associate with multiple factors involved in transcription and RNA processing; however, no single factor we tested appears to be as important for *Xist*-mediated silencing as SPEN.

DISCUSSION

We have resolved major discrepancies in how *Xist* recruits SPEN by demonstrating that the sequences in Repeat A most essential for gene silencing do so by binding canonical splicing factors such as SRSF1, which in turn recruit SPEN. These same sequences are potent local suppressors of splicing and, together with SPEN, enable Repeat A to associate with multiple proteins and protein complexes, including m^{6}A writers and readers, nuclear exosome adapters, a transcriptional antiterminator, and elongating RNA polymerase II. Indeed, several factors previously implicated in *Xist* function – HDAC3, TRIM28, RBM15, and YTHDC1[10,12,75,79,101] – depend on SPEN for Repeat A association (Figure 7A).

Our findings suggest that Repeat A is a splicing factor-binding module that recruits SPEN predominantly through bridged interactions with a subset of SR proteins. While SRSF1 is one such protein, our data suggest that yet-to-be-identified proteins also bind the AWCG and downstream motifs in Repeat A, and that these proteins can substitute for or work in concert with SRSF1 to recruit SPEN. Many SR proteins have overlapping binding specificities and are several orders of magnitude more abundant than SPEN in cells.[56,58,73,102] Thus, it stands to reason that SR proteins assemble on Repeat A prior to recruiting SPEN, likely as *Xist* is transcribed. Moreover, both SR proteins and SPEN interact with elongating RNA polymerase II,[57,64,93,94,103] findings we corroborated by RIP. We speculate that in the context of nascent *Xist*, interactions among SR proteins, SPEN, and the transcriptional machinery are partly if not fully responsible for the role of Repeat A in promoting *Xist* production. Concurrently, the periphery of nuclear speckles and other foci rich in speckle-related proteins are hubs of transcription and splicing.[93,104-107] Thus, after mature *Xist* is processed and released from its own locus, SR protein- and SPEN-driven interactions might grant the *Xist* RNP access to actively transcribed regions (Figure 7B).[13]

Once near an actively transcribed region, our data suggest that *Xist* is capable of silencing via pathways parallel to or independent of chromatin modification, including by inducing premature transcriptional termination and/or nascent RNA turnover. By O-MAP, the presence of several proteins (HDAC3, TRIM28, ZC3H18, SCAF4) within the *Xist* microenvironment was SPEN-independent; however, by RIP-qPCR, their physical associations with Repeat A were strongly dependent on SPEN. Moreover, for all factors investigated, the magnitude of SPEN dependence was greater by RIP-qPCR than by O-MAP, suggesting that SPEN strengthens and integrates interactions between *Xist* and multiple specific nearby factors (Figure 7A). Indeed, at the levels of knockdown we achieved, none of the Repeat-A-associated proteins we profiled – including HDAC3 and TRIM28 – appeared to be on their own responsible for transcriptional silencing by *Xist*, suggesting that SPEN-dependent silencing relies on multiple or yet-to-be tested factors, or that SPEN primarily silences transcription through direct engagement with RNA polymerase II. Our observation that Repeat A associates with elongating but not initiating RNA polymerase II, along with prior observations that the chromatin-bound fraction of RNA polymerase II is destabilized in a SPEN-dependent manner at the onset of X-chromosome inactivation[108] support the possibility that SPEN represses transcription by destabilizing elongating RNA polymerase II. Whether the *Xist* RNP might also destabilize nascent RNA remains unclear. While we did not observe association between Repeat A and CPSF/CFIm or Integrator, we did detect SPEN- and splicing factor motif-dependent associations between Repeat A and the m^{6}A machinery and nuclear exosome adapters, both which are involved in degradation of chromatin-

associated RNAs.[76-78,91,92,109] Moreover, in plants, the SPEN homologue FPA induces premature cleavage and degradation of nascent RNAs subsequent to transcriptional silencing, raising the possibility that SPEN-family proteins have long been involved in clearance of nascent RNA.[110-112] Elucidating the mechanisms by which SPEN silences transcription is a critical next step.

SPEN and SR proteins are deeply conserved, predating the emergence of *Xist* by several hundred million years.[16,113] We surmise that SPEN's ability to recognize SR protein-rich assemblies also predates *Xist*. In contexts outside XCI, SPEN may recognize SR-rich assemblies to maintain homeostatic gene expression through transcriptional dampening or silence transcription of nascent RNAs upon which the spliceosome has misassembled or stalled. The notion that SPEN may dampen transcription through SR-rich assembly recognition is supported by our co-expression and O-MAP analyses, which identified associations between SPEN and multiple transcriptional activators and elongation factors, and by our proteomic and RIP-seq analyses, which demonstrate that the N-terminus of SPEN associates with protein and RNA components of nuclear speckles and the spliceosome. Conversely, the notion that SPEN may silence nascent RNAs bound by stalled spliceosomes or spliceosome-recruiting complexes could rationalize its role in the silencing of transposons, particularly if they lack efficient splicing signals.[18,112]

We speculate that the recognition of stalled splicing machinery by SPEN was the evolutionary force that gave rise to Repeat A. Repeat A is replete with GC-rich consensus SR protein-binding motifs and U-rich spacers that resemble canonical polypyrimidine tracts, which each associate with spliceosome-recruiting proteins, including SRSF1, SRSF2, U2AF1, and U2AF2. Yet, Repeat A lacks consensus 5' and 3' splice sites, does not appreciably associate with core spliceosome components (e.g., SNRNP70, SNRPA, or SNRNP200; Figure 3E), and is not spliced itself. Rather, it has a local suppressive effect on splicing. Thus, disruption of splicing near the 5' end of proto-*Xist*,[114] either by chance mutation or transposon insertion,[18,115] followed by expansion of a proto-Repeat A monomer, may represent the path by which SPEN recruitment and transcriptional silencing ability arose in present-day *Xist*. Evolutionary pressure has since kept the sequence of Repeat A well-conserved, as our results demonstrate that even minor sequence perturbations – down to a single nucleotide – can inflict major penalties in SPEN recruitment and *Xist*-induced silencing.

Lastly, our discoveries provide new perspectives on the roles of SPEN in development and disease. Haploinsufficiency of *SPEN*, located in the rearrangement-prone 1p36 region of the human genome, causes neurodevelopmental disorders.[31] *SPEN* is also recurrently mutated in

cancers, harboring a mutational profile generally consistent with that of a tumor suppressor,[32-40] and regulates NOTCH-, WNT-, growth factor-, and hormone-dependent gene expression programs.[40,98,116-125] In these contexts and others, the possibility that SPEN represses transcription through the sensing of SR protein-rich assemblies should now be evaluated.

ACKNOWLEDGMENTS

We thank Brian Golitz, Noah Sciaky, and Andrew Snipes of the UNC CRISPR Screening Facility for sequencing. This work was supported by NIH grants R01GM136819, R01GM121806, and R35GM153293; NSF grant DBI-2228805; and the Yang Family Biomedical Scholar Fund (J.M.C.); NIH grants R01GM138799 (to S.A.S., L.E.D., and D.M.S.), R21CA288806 (to S.E.O), R35GM142864 (to D.D.), R01HL160825 (to D.M.S.), and a Chan-Zuckerberg Institute Collaborative Pairs project award (to D.M.S.). J.B.T. was supported by T32CA217824 and a Marzluff Postdoctoral Fellowship from the RNA Discovery Center. S.B.P. was supported by T32GM135128. S.A.S. was supported by T32GM007750 and the ISCRM Fellows Program. M.K.S was supported by K99GM152778. Table S4 results are in part based upon data generated by TCGA (https://www.cancer.gov/tcga). This work used an EASY-nLC1200 UHPLC and Thermo Scientific Orbitrap Fusion Lumos Tribrid mass spectrometer purchased with funding from NIH-SIG grant S10OD021502 to S.E.O.

AUTHOR CONTRIBUTIONS

J.B.T., A.P., D.M.S., and J.M.C. conceived the study; J.B.T., A.P., S.A.S, L.E.D., Q.E.E., S.P.B., Z.Z., D.M.L., N.S., S.N.N., and M.-C.G.B. performed the experiments and/or data analysis; S.E.H., M.K.S., S.-E.O., M.P.G.-P., S.A.S., D.D., and D.M.S. provided reagents and advice; J.B.T., D.M.S., and J.M.C. acquired funding; and J.B.T. and J.M.C. wrote the paper with feedback and appropriate methods contributed by co-authors.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURES AND LEGENDS

Figure 1. Sequences critical for *Xist*-induced silencing, *Xist* production, and *Xist* splicing fidelity are located within the GC-rich cores of Repeat A. (A) *Xist* gene insertion into *Rosa26*.[41] **(B)** Summary of *Xist* mutants, including mutation positions in each Repeat A monomer. **(C, E)** *Xist* abundance relative to WT by RNA-seq across mutants (C) or WT dox titration (E). Dots, independent clonal cell lines. **(D, F)** *Xist* silencing activity by RNA-seq for *Xist* mutants (D) or WT dox titration (F). Dots, sample-averaged expression of B6 chr6 *Xist*-repressed genes relative to empty-cargo control. Black bars, median. **(G)** Median silencing vs. *Xist* RNA abundance. Dashed line, silencing expected for corresponding levels of WT *Xist*. **(H)** Sashimi plots illustrating normal (WT) and aberrant (ΔRepA-XB) *Xist* splicing. **(I)** Quantification of aberrant splicing across

genotypes (dots, independent clonal cell lines). SJ, splice junction. For all panels vs. WT (two-sided t-test): (*), $p < 0.05$; (**), $p < 0.01$. See Figures S1, S2.

Figure 2. SPEN association with Repeat A *in vivo* depends on sequences that it does not bind *in vitro*. (A) RIP schematic.[41] **(B)** SPEN-Repeat A association across genotypes by input-normalized RIP-qPCR. SM33 WT vs. Δ*Spen* ESCs confirm specificity of SPEN antibody.[51] *Srsf2* mRNA, control. Dots, triplicate qPCR values from two independent experiments. **(C)** Median silencing vs. SPEN-Repeat A association across genotypes. *r*, *p*: data fit to linear regression. **(D)** Specificity vs. signal for SPEN RIP-seq peaks in SM33 ESCs. **(E)** Motifs from top 1500 non-*Xist*, non-*Spen* SPEN RIP-seq peaks. **(F)** Overview of equilibrium RBNQ. SBP, streptavidin-binding peptide. **(G)** RBNQ with SPEN RRM2-4 and Repeat A. Dots, values from two independent experiments. Error bars, standard deviation of qPCR triplicates. CI, confidence interval. **(H)** RBNQ across Repeat A RNAs using fixed SPEN RRM2-4 concentration. Dots, qPCR triplicates from ≥2 independent experiments. For all panels vs. WT (two-sided t-test): (*), $p < 0.05$; (**), $p < 0.01$. See Figure S3.

Figure 3. Motif-dependent association between Repeat A and SR proteins *in vitro* and *in vivo*. (A) Strategy to identify Repeat A-interacting proteins *in vitro*. **(B)** Proteins enriched by WT over ΔGC-core Repeat A ($p < 0.05$, two-sided t-test, n=3). **(C)** GO terms of GC-core-specific proteins (top 20). **(D)** Relative recovery of SR proteins across Repeat A RNAs. Columns, independent experiments. **(E)** RMCE-*Xist* WT RIP-seq with antibodies against controls, SPEN,

and SR and/or RNA-processing proteins. **(F)** RIP-seq signal within Repeat A relative to total *Xist*. Dots, independent experiments. See Figure S4.

Figure 4. SPEN associates with nuclear speckle components and other SR proteins via its N-terminus. (A) *Xist* abundance across genotypes (RNA-seq). Dots, independent clonal lines. **(B)** Halo-SPEN-V5-interacting proteins ranked by proportion of peptides recovered. **(C)** GO terms of SPEN-interacting proteins (top 10). **(D)** V5 IP, Halo and SRSF1 western blots (no-rescue vs. Halo-SPEN-V5 ESCs). **(E)** SPEN domains and deletion mutants. IDRs, MobiDB.[126] **(F)** SPEN-Repeat A association for WT vs. mutants by input-normalized Halo RIP-qPCR. Dots, qPCR triplicates (≥2 independent experiments). Asterisks, significantly different from WT. Crosses, significantly different from RRM1-4. **(G)** Enrichment of proteins, by class, in RRM1-4 vs. RRM2-4 IP-MS/MS. "(0)," no recovery with RRM2-4. **(H)** Peptide counts of SR and/or speckle proteins in

RRM1-4 or RRM2-4 IP-MS/MS. Dots, independent experiments. **(I)** Motifs from top 1500 non-*Xist*, non-*Spen* Halo RIP-seq peaks across genotypes. **(J)** Significant *SPEN*-correlated genes from GTEx (guilt-by-association; $p < 0.01$ and FDR < 0.1, see Methods). **(K)** Overlap of significant *SPEN*-correlated genes (FDR < 0.1) from the GTEx, TCGA Tumor (non-chr1), and TCGA Normal datasets. **(L)** GO terms of *SPEN*-correlated genes from GTEx (top 10 and 14th). Excepting (J), *p*-values, two-sided t-tests: (*/†), $p < 0.05$; (**), $p < 0.01$. See Figure S5.

Figure 5. SRSF1 directly binds a conserved Repeat A motif to recruit SPEN to *Xist in vivo*.
(A) SRSF1-Repeat A association in SM33 or RMCE-*Xist* ESCs by input-normalized RIP-qPCR.
Srsf2 mRNA, control. Dots, qPCR triplicates from two independent experiments. **(B)** SPEN- vs.
SRSF1-Repeat A association (RIP-qPCR). *r*, *p*: data fit to linear regression. **(C)** RBNQ with
SRSF1 RRM1-2 and Repeat A. Dots, values from two independent experiments. Error bars,
standard deviation of qPCR triplicates. **(D)** RBNQ across Repeat A RNAs using fixed SRSF1
RRM1-2 concentration. Dots, qPCR triplicates values, ≥2 independent experiments. **(E)**
FKBP1A^{F36V}-SRSF1 degradation by dTAG^V-1 in cells derived from RMCE-*Xist* WT. **(F)** SPEN-
Repeat A association in WT or *FKBP1A^{F36V}-Srsf1* cells by input-normalized RIP-qPCR. Dots,

qPCR triplicates from independent clonal lines. **(G)** *Xist* abundance (RNA-seq) in *FKBP1A^F36V^-Srsf1* cells, with/without dox and with/without dTAG^V^-1. Dots, independent lines, relative to dox(+), dTAG^V^-1(-) condition. **(H)** *Xist* silencing activity as in Figure 1D,F, for *FKBP1A^F36V^-Srsf1* cells treated as in (G). **(I)** MCP-MS2 tethering to assess SPEN recruitment to *Xist* ΔRepA-XB-6x(U-spacer-3xMS2). **(J)** SPEN-6x(U-spacer-3xMS2) association across MCP genotypes by input-normalized RIP-qPCR. Dots, qPCR triplicates from ≥2 independent experiments. **(K)** *Xist* abundance (RNA-seq) across genotypes. Dots, independent experiments. **(L)** *Xist* silencing activity as in Figure 1D,F across MCP genotypes. "n.s.," not significantly different. *P*-values, two-sided t-tests relative to WT or EGFP unless specified by black bars: (*), $p < 0.05$; (**), $p < 0.01$. See Figure S6.

Figure 6. SPEN and SR protein-binding motifs promote interactions between Repeat A and the m⁶A, RNA turnover, and transcriptional elongation machineries. (A) O-MAP schematic; HRP, horseradish peroxidase.[96] **(B)** Proteins differentially recovered by O-MAP from WT and Δ*Spen* cells (*p* < 0.05, two-sided t-test, n=4). **(C)** Repeat A association (input-normalized RIP-qPCR) in non-targeting-control and Δ*Spen* ESCs derived from RMCE-*Xist* WT, and RMCE-*Xist* AWCG+ΔGG cells. Values relative to non-targeting controls. Dots, qPCR triplicates from ≥2 independent experiments. Values above non-targeting-control bars, experiment-averaged IP/input relative to IgG control. **(D)** RIP-qPCR as in (C), shown relative to IgG. **(E)** *Xist* abundance (RNA-seq) in RMCE-*Xist* WT ESCs treated with non-targeting control siRNA (siNT) and 1000, 25, or 10 ng/mL dox, or protein-targeting siRNAs and 1000 ng/mL dox. Values relative to siNT-1000. Dots, independent experiments. **(F)** *Xist* silencing activity as in Figure 1D,F, for siRNA-treated

cells. **(G)** Median silencing activity vs. *Xist* abundance. Δ*Spen* data, Figures 4A, S5C. Dashed line, silencing expected for corresponding levels of *Xist* in siNT ESCs. *P*-values, two-sided t-tests relative to WT or non-targeting controls: (*) indicates $p < 0.05$; (**) indicates $p < 0.01$. See Figure S7.

Figure 7. Models of Repeat A RNP composition and function. (A) Repeats 6 and 7 depicted according to structural model from Lu et al.[17,127] SPEN is recruited to Repeat A predominantly by proteins that bind motifs in the GC-rich core, whereas the U-rich spacers play minor roles. SPEN and GC-rich-core-bound proteins enable association with factors involved in transcriptional elongation and RNA processing. **(B)** Sensing of SR protein-rich transcriptional hubs by Repeat A and SPEN.

SUPPLEMENTARY FIGURES AND LEGENDS

Figure S1. Related to Figure 1.

(A) Sequences of human (top) and mouse (bottom) Repeat A, with each repeat monomer centered on its conserved GC-rich core sequence. Protein-interacting nucleotides detected by RNP-MaP[6] are highlighted in white and tallied for each position (grey bars above and below).

(B) Per-nucleotide ex-vivo SHAPE reactivity for Repeat A in mouse trophoblast stem (TS) cells.[8]

(C) *Xist*-induced silencing of genes along the B6 allele of chr6, determined by RNA-seq. The average expression level of each expressed gene in RMCE-*Xist* WT cells (grey and pink dots) is plotted relative to the average expression level in RMCE-empty cells (magenta dots). The 204 genes with significantly altered expression ($p_{adj} < 0.001$; DESeq2)[41] are depicted by pink dots and designated as "*Xist*-repressed genes."

(D and E) Silencing activity for *Xist*-repressed genes in individual clonal cell lines. Average values for each gene in each genotype/condition are shown in Figure 1D,F. Black bars, median values.

(F) *Xist* RNA stability across WT and Repeat A mutants, using RT-qPCR to measure differences following treatment for 4 h with actinomycin D (ActD). Dots, average values from three independent experiments.

(G) Semiquantitative PCR using intron-flanking primers to estimate splicing efficiency of the ectopic intron inserted downstream of Repeat A in independent cell lines. Spliced and unspliced controls used RMCE-*Xist* WT and intron plasmids, respectively, as PCR templates.

Figure S2. Related to Figure 1.

Repeat A mutant sequences, with deletions shown as dashes and substitutions shown in white. Each repeat monomer sequence is aligned and centered on its GC-rich core (flanked by black lines), with position of each GC-rich core nucleotide numbered above.

Figure S3. Related to Figure 2.

(A) Genotyping PCR for independent clonal lines of SM33 Δ*Spen* cells generated by CRISPR-Cas9. PCR1 was performed using primers flanking either side of the deletion; PCR2 and PCR3 were performed using primers flanking the upstream and downstream gRNA target sites.

(B) RT-qPCR measuring *Spen* mRNA in SM33 WT and Δ*Spen* cells. Values are relative to SM33 WT. Dots represent average qPCR-replicate values for independent cell lines.

(C) Top 15 most significant sequence motifs enriched in the 1500 most abundant non-*Xist*, non-*Spen* SPEN-specific peaks from SPEN RIP-seq, determined by STREME.

(D) Promega Diamond-stained urea-PAGE gels of in-vitro-transcribed Repeat A RNAs used for equilibrium RBNQ experiments.

(E) Coomassie blue-stained SDS-PAGE gel of two independent preps of SBP-tagged SPEN RRM2-4 used for replicates of equilibrium RBNQ experiments.

A *In-vitro*-transcribed, internally biotinylated RNAs
Urea-PAGE, Promega Diamond stain

B Biotinylated RNA pulldown for proteomics, 15% of eluates
SDS-PAGE, silver stain

*not used for pulldown experiments

C Biotinylated RNA pulldown, western blot

D

Figure S4. Related to Figure 3.

(A) Promega Diamond-stained urea-PAGE gels of in-vitro-transcribed, internally biotinylated Repeat A RNAs used for biotinylated RNA pulldown experiments.

(B) Representative silver-stained SDS-PAGE gel of proteins recovered by biotinylated Repeat A mutant pulldowns or no-RNA control pulldown. The remaining 85% of each sample was analyzed by quantitative proteomics.

(C) Validation of quantitative proteomics results by biotinylated RNA pulldown and western blotting for the proteins RBM15, U2AF2, and SRSF1.

(D) Quantification of the proportion of RIP-seq signal in Figure 3E within Repeat A relative to total reads per million. Dots, values from independent experiments.

Figure S5. Related to Figure 4.

(A) Genotyping PCR for independent clonal lines of RMCE-*Xist* WT Δ*Spen* cells or non-targeting gRNA-treated control cells generated by CRISPR-Cas9. PCR1 was performed using primers

flanking either side of the deletion; PCR2 and PCR3 were performed using primers flanking the upstream and downstream *Spen* gRNA target sites.

(B) Confirmation of expression of WT and mutant Halo-SPEN-V5 proteins by IP-WB or WB.

(C) *Xist* silencing activity determined by RNA-seq, as in Figure 1D,F, for non-targeting control cells and Δ*Spen* cells with and without WT Halo-SPEN-V5 expression. Asterisks, significant differences by two-sided t-test.

(D) Quantification of exon 1 aberrant splicing for each RMCE-*Xist* cell line (dots), as done in Figure 1I. Asterisks, values significantly different from WT by two-sided t-test.

(E and F) Quantification of RS/SR and RE/ER dipeptide content in each IDR of SPEN by percentage of residues (E) or overall residue count (F).

(G) Gene ontology analysis of the 75 proteins recovered with SPEN RRM1-4 two-fold more than with RRM2-4 in two independent experiments, showing the 10 most significantly enriched terms.

(H) Gene ontology analysis of the 44 non-chromosome 1 (chr1) genes with expression significantly correlated with *SPEN* across tissue types in the TCGA Tumor dataset (FDR < 0.1). An additional 59 chr1 genes were significantly correlated with *SPEN* in this dataset but were omitted from this analysis because their correlations with *SPEN*, a gene on chr1, cannot be decoupled from cancer-associated copy-number variations.

A FLAG RIP, Repeat A qPCR

B Purified, SBP-tagged SRSF1 RRM1-2 for RBNQ SDS-PAGE, Coomassie blue stain

C *FKBP1A^{F36V}-Srsf1* genotyping PCR RMCE-*Xist* WT cells

D *Xist* exon 1 aberrant splicing

E MCP-X-NLS-3xFLAG

*not used for RIP-qPCR experiments

F SRSF1 RIP, 6x(U-spacer-3xMS2) qPCR

FLAG RIP, 6x(U-spacer-3xMS2) qPCR

G SPEN RIP, Repeat A or 6x(U-spacer-3xMS2) qPCR

H *Xist* RNA abundance vs. SPEN-Repeat A association

I *Xist* RNA abundance vs. SRSF1-Repeat A association

Figure S6. Related to Figure 5.

(A) Levels of Repeat A recovery by input-normalized FLAG RIP-qPCR from RMCE-*Xist* WT cells expressing no FLAG-tagged construct and RMCE-*Xist* WT and ΔCCUGC cells expressing SRSF2-3xFLAG. Dots, qPCR triplicate values from one experiment.

(B) Coomassie blue-stained SDS-PAGE gel of two independent preps of SBP-tagged SRSF1 RRM1-2 used for replicates of equilibrium RBNQ experiments.

(C) Genotyping PCR for parental RMCE-*Xist* WT control cells or independent clonal lines of *FKBP1A^F36V^-Srsf1* cells generated by CRISPR-Cas9 and homology-directed repair. PCR was performed using primers flanking either side of the *FKBP1A^F36V^* degron tag insertion.

(D) Quantification of exon 1 aberrant splicing for each RMCE-*Xist* cell line (dots), as done in Figure 1I. Asterisks, values significantly different from WT by two-sided t-test.

(E) Western blot confirmation of expression of untagged EGFP or various MCP-tagged proteins in RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) cells.

(F) Confirmation of MCP-tagged protein tethering by input-normalized RIP-qPCR using antibodies against SRSF1 (left) or FLAG (right). Cells expressing untagged EGFP served as a negative control. Dots, qPCR triplicate values from two independent experiments. Asterisks, significantly different from untagged EGFP control by two-sided t-test.

(G) Levels of SPEN association with WT, mutant, or synthetic Repeat A in RMCE-*Xist* WT, ΔU-spacer, or ΔGC-core cells or RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) cells, determined by input-normalized RIP-qPCR. Dots, qPCR triplicate values from two independent experiments (one that included ΔGC-core). Asterisk, significantly different from WT by two-sided t-test (two-sided one-sample t-test for ΔGC-core).

(H and I) Plots of *Xist* RNA abundance as a function of SPEN-Repeat A association level (H) or SRSF1-Repeat A association level (I). Values of *r* and *p* reflect fit to a linear regression models with the indicated data. The ss234 deletion reduces *Xist* abundance independent of Repeat A and was thus omitted.

Figure S7. Related to Figure 6.

(A) RIP-seq from RMCE-*Xist* non-targeting gRNA control cells, with antibodies used in Figure 6C. IgG (grey) and SPEN (blue) RIP-seq tracks from Figure 3E are included as controls. Note that for some RIPs, enrichment over Repeat A relative to the rest of *Xist* was low; nevertheless, Repeat A was recovered at levels significantly higher than IgG control in input-normalized RIP-qPCR (Figure 6C).

(B) Assessment of siRNA-mediated protein knockdowns by western blot. RMCE-*Xist* non-targeting control and Δ*Spen* cells were included to confirm that differences in Repeat A association observed in Figure 6C are not attributable to differences in RIP target protein expression.

SUPPLEMENTARY TABLES

Table S1. Quantitative proteomics of biotinylated Repeat A RNA pulldowns. Related to Figure 3.

Table S2. Halo-SPEN IP proteomics. Related to Figure 4.

Table S3. Halo RIP-seq target genes and GO analyses. Related to Figure 4.

Table S4. *SPEN*-correlated genes by guilt-by-association. Related to Figure 4.

Table S5. WT vs. Δ*Spen* O-MAP proteomics. Related to Figure 6.

Table S6. Oligonucleotides and antibodies used in this study. Related to Figures 1-6.

MATERIALS AND METHODS

Cell culture

Male E14 mouse embryonic stem cells (mESCs, kind gift of D. Ciavatta), male B6/CAST F1-hybrid *Rosa26*-RMCE mESCs,[26] and male SM33 mESCs (with dox-inducible promoter at the endogenous *Xist* locus; kind gift of K. Plath; [52]), were grown on gelatin-coated plastic dishes in a humidified Thermo Fisher Forma Series II water-jacketed incubator at 37 °C and under 5% CO_2. Cells were cultured in growth medium consisting of DMEM high glucose plus sodium pyruvate (Gibco 11995-065), 15% ESC-qualified fetal bovine serum (Gibco 26140-079), 0.1 mM non-essential amino acids (Gibco 11140-050), 100 U/mL penicillin-streptomycin (Gibco 15140-122), 2 mM L-glutamine (Gibco 25030-081), 0.1 mM β-mercaptoethanol (Sigma-Aldrich 63689), and 1:500 LIF conditioned media produced from Lif-1Cα (COS) cells (kind gift of N. Hathaway).

Fresh growth medium was supplied daily, and cells were split every two days. To do so, cells were washed once with 1X phosphate-buffered saline (PBS; Corning 46-013-CM), dissociated by adding pre-warmed (37 °C) 0.125% trypsin-EDTA (Gibco 2520072; diluted from 0.25% with 1X PBS) for ~3-5 min, and quenched by adding two volumes of pre-warmed "standard" quenching medium (DMEM high glucose plus sodium pyruvate containing 10% fetal bovine serum, 100 U/mL penicillin-streptomycin, 2 mM L-glutamine and, 0.1 mM β-mercaptoethanol). Cells were gently dissociated to single-cell suspension using a P1000 pipet before centrifuging the desired number of cells (most typically a 1:6 split) in a conical tube at 200 x g for 5 min at room temp. Supernatants were aspirated, and cells were resuspended in pre-warmed growth medium via serological pipet before adding to tissue-culture-treated plastic dishes (Genesee Scientific) pre-treated with 0.1% (w/v) gelatin (Sigma G9391).

Plasmid construction

Repeat A mutants were first generated in plasmid pTETRISv1 [26] as follows. ~1.2-kb DNA sequences were synthesized (Genewiz) corresponding to the 5′ end of *Mus musculus Xist* and containing the desired Repeat A mutations (sequences shown in Figure S2). These sequences were inserted into pTETRISv1-*Xist*-2kb WT [26] digested with KpnI and NdeI. The insert sequences of the resulting plasmids were verified by Sanger sequencing.

To generate pCARGO-RMCE-*Xist*, ΔU-spacer, ΔGC-core, disrupt, AWCG, AWCG+ΔGG, U19C, U23C, and ΔCCUGC, mutant sequences were PCR-amplified using pTETRISv1-*Xist*-2kb mutants as template and inserted into pCARGO-RMCE-*Xist* digested with NotI and PmlI (cloned by Genewiz).

To generate pCARGO-RMCE-*Xist* ΔRepA (deletion of *Xist* nt 307-732), preserve, ΔGG(16,17), and Δss234 (deletion of *Xist* nt 731-1472), mutant sequences were PCR-amplified from corresponding pTETRISv1-*Xist*-2kb mutants using Q5 High-Fidelity DNA Polymerase (NEB M0491) and primers JT351 and JT352 (Table S6). PCR products (~0.02 pmol) were inserted into 0.01 pmol NotI-and-PmlI-digested pCARGO-RMCE-*Xist* using NEBuilder HiFi DNA Assembly (NEB E2621). Assembly products were diluted 4-fold with water and transformed into MAX Efficiency Stbl2 cells (Invitrogen 10268019), which were grown overnight at 30 °C with ampicillin selection.

To generate pCARGO-RMCE-*Xist* ΔRepA-XB, pGEM-*Xist*-2kb [42] was first modified via site-directed mutagenesis to simultaneously delete Repeat A (*Xist* nt 309-736) and replace it with XmaI and BsiWI restriction sites, generating pGEM-*Xist*-2kb ΔRepA-XB. This was done using NEB Q5 High-Fidelity DNA Polymerase, primers JT353 and JT354, and KLD enzyme mix (NEB M0554). The mutated *Xist* sequence was then PCR-amplified using NEB Q5 High-Fidelity DNA Polymerase and primers JT351 and JT352 and inserted into NotI-and-PmlI-digested pCARGO-RMCE-*Xist* using NEBuilder HiFi DNA Assembly as described above.

To generate pCARGO-RMCE-*Xist* intron, a 328-bp dsDNA gBlock (IDT, JT355, Table S6) was synthesized corresponding to *Xist* sequence containing a 133-bp chimeric β-globin/immunoglobulin intron [45] inserted between *Xist* bases 1919 and 1920 (NCBI NR_001463.3). This dsDNA fragment was inserted into PmlI-and-BsaBI-digested pCARGO-RMCE-*Xist* using NEBuilder HiFi DNA Assembly as described above.

To generate pCARGO-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2), the following 6x(U-spacer-3xMS2) sequence was first synthesized by Genewiz and inserted into pUC-GW-Kan, including the flanking 5′ XmaI and 3′ BsiWI sites (uppercase, not bold):

CCCGGG**TTTCTTTCATTGTTTATATATTCTT**cgggtggaggatcaccccacccgacacttcacaatcaggg
tggaggaacaccccaccctccagacacattacacagaagcaccatcagggcttctg**CTTTTCATTCTTTTTTTTTCTTA
TTATTTTTTTTTCTAAACTT**cgggaggacgatcacgcctcccgaatatcggcatgttgcgtggagcatcagcccacgca
gccaatcagagtccgcgaagagcatcagccttcgcg**TTTTATTCTTTTTTTTCTTCTCCTTA**caggaacagcatcagc
gttcctgcccagtacccaaccccatgcagcatcagcgcatgggccccaagaatacatccgagcaccaacagggctcgga**GTTT
AGTCTTTTTTTCCCAT**tccatggaccatcaggccatggactctcaccaacaatcgaagcagcatcagcgcttcgaaacact
cgagcattgggtggacgatcacgccacccaTGTTATTATTTTTTTTTTCTTTTTCTTTTgacgagcaccaccagggc
tcgtcgttccacgtccaacgggatcacgatcacggatcccgcagctacatcactagcatgcacgatcacggcatgct**TTTAAATT
TTTTTTTTCACG**tgggttcagcaccagcgaacccactcctacctcaaaggcaagcaggatcaccgcttgcccattccaacata
cctggtacagcatcagcgtaccagCGTACG.

This sequence contains, in order, the endogenous mouse U-rich spacers that precede the GC-rich cores of Repeat A repeats 1, 3, 4, 6, 7, and 8 (bold, uppercase). 3xMS2 sequences, synonymized to prevent recombination (lowercase, underlined), and linkers (lowercase, not underlined) between each U-rich spacer were derived from Addgene plasmid #84561 [128]. The 6x(U-spacer-3xMS2) sequence was liberated with XmaI and BsiWI and ligated, via NEB Quick Ligase (NEB M2200S), into XmaI-and-BsiWI-digested pCARGO-RMCE-*Xist* ΔRepA-XB.

For expression and purification of proteins N-terminally tagged with glutathione-S-transferase (GST; cleavable) and streptavidin-binding peptide (SBP), sequences coding for SPEN and SRSF1 RRMs were cloned into pDD001 (lab of D. Dominguez) as follows. To generate pDD001-SPEN-RRM2-4, an *Escherichia coli*-optimized coding sequence corresponding to *Mus musculus* SPEN amino acids 335-620 was first synthesized by Genewiz and then PCR-amplified with NEB Q5 High-Fidelity DNA Polymerase and primers JT396 and JT397 and inserted into NcoI-and-NotI-digested pDD001 using NEBuilder HiFi DNA Assembly. To generate pDD001-SRSF1-RRM1-2, SRSF1 RRM1-2 sequence (*Mus musculus* SRSF1 amino acids 1-197) was PCR-amplified from template plasmid pGEX6P-1-SF2 (Addgene #99020, a gift from Honglin Chen, [129]) using NEB Q5 High-Fidelity DNA Polymerase and primers JT388 and JT390 and inserted into NcoI-and-NotI-digested pDD001 using NEBuilder HiFi DNA Assembly. The sequences of the pDD001 plasmids were confirmed by diagnostic digestion with NcoI and NotI and by Sanger sequencing of the full insert sequences.

The mammalian expression plasmid pCL116(PuroR) – which enables piggyBac-mediated insertion, puromycin-based selection, and doxycycline-induced expression of genes of interest (EGFP in the parental plasmid) – was generated by removing the *HygroR* sequence in pCL116 ("pb-miRE-tre," from which Addgene plasmid #196087 was derived[130]) with NheI and SnaBI and replacing it with *PuroR* sequence (from pTETRISv1, Addgene plasmid #126033 [42]) generated by DNA synthesis (Genewiz).

To generate pCL116(PuroR)-SRSF2-3xFLAG, DNA was synthesized (Genewiz) containing mouse-optimized (IDT Codon Optimization Tool, https://www.idtdna.com/pages/tools/codon-optimization-tool) SRSF2 coding sequence followed by a 3xFLAG tag (MDYKDHDGDYKDHDIDYKDDDDK), and this sequence was inserted into pCL116(PuroR)-HA-MCP-SRSF1_v2 (described below) digested with AgeI and SalI (to remove HA-MCP-SRSF1).

To generate pCL116(PuroR)-Halo-SPEN-V5 WT, the Halo-SPEN-V5 sequence from PyPP-CAG-Halo-full-length-Spen-V5 (kind gift of K. Plath [62]) was PCR-amplified using NEB Q5 High-Fidelity DNA Polymerase and primers JT418 and JT419 and inserted into SalI-and-NcoI-digested (to remove EGFP) pCL116(PuroR) using NEBuilder HiFi DNA Assembly. The RS domain was

deleted by Genewiz. To generate the remaining SPEN deletion mutants in pCL116(PuroR)-Halo-SPEN-V5, a strategy was used in which two unique restriction enzymes were used to remove sequence from pCL116(PuroR)-Halo-SPEN-V5 encompassing the desired deletion(s), and NEBuilder HiFi DNA Assembly was used to reconstitute the removed region such that it contained the desired deletion(s). To generate the plasmid backbones for NEBuilder HiFi DNA Assembly, pCL116(PuroR)-Halo-SPEN-V5 was digested with the following pairs of restriction enzymes: NotI and MfeI for RRM1-4, RRM2-4, IDR2-RRM4, and RRM1-4 ΔIDR2; KpnI and MfeI for ΔSPOC; and NotI and PmlI for all others except IDR2-RRM4 and RRM1-4 ΔIDR2, which were generated using NotI-and-MfeI-digested pCL116(PuroR)-Halo-SPEN(RRM2-4)-V5. PCR fragments for NEBuilder HiFi DNA Assembly were generated using with NEB Q5 High-Fidelity DNA Polymerase, pCL116(PuroR)-Halo-SPEN-V5 as template, and primer pairs detailed in Table S6. The following PCR fragments (Table S6) were used to construct each mutant: ΔRRM1-4, fragment 8; ΔRRM2-4, fragments 9 and 10; ΔRRM1-IDR2, fragment 14; ΔIDR2, fragments 1 and 2; ΔIDR4, fragments 3 and 4; ΔIDR2 ΔIDR4, fragments 1, 4, and 5; ΔSPOC, fragments 6 and 7; RRM1-4, fragments 11 and 12; RRM2-4, fragments 12 and 13; IDR2-RRM4, fragment 15; RRM1-4 ΔIDR2, fragments 1 and 16. Assembly reaction products were transformed into NEB high-efficiency 5-alpha competent cells (NEB C2987H), which were grown in LB containing ampicillin.

The SPEN mutations by amino acid (aa) residue are as follows: ΔRRM1-4, Δaa1-590; ΔRRM2-4, Δaa335-590; ΔRRM1-IDR2, Δaa1-334; ΔIDR2, Δaa99-330; ΔRS, Δ186-310; ΔIDR4, Δaa694-900; ΔIDR2 ΔIDR4, both Δaa99-330 and Δaa694-900; ΔSPOC, Δaa3478-3644*; RRM1-4, aa1-620 only; RRM2-4, aa335-620 only; IDR2-RRM4, aa99-620 only; RRM1-4 ΔIDR2, aa1-620 with Δaa99-330 (*note that the T2201-A2202 dipeptide in the Uniprot mouse SPEN entry is missing in this plasmid; as a result, SPOC is technically aa 3476-3642 in this plasmid).

To modify the endogenous *Srsf1* loci for dTAG-mediated degradation, two plasmids were constructed, the first containing Cas9 and a guide RNA (gRNA) targeted to the immediate 5' end of *Srsf1* (gRNA sequence determined with Benchling CRISPR Guide RNA Design tool; https://www.benchling.com/crispr), and the second containing a homology-directed repair (HDR) donor with the PuroR-P2A-2xHA-FKBP1A^{F36V} tag. The first plasmid (PX459_v2-Srsf1-N1-gRNA) was constructed by annealing gRNA-sequence oligos JT459 and JT460 and inserting this sticky-ended dsDNA into BbsI-digested PX459 V2.0 (Addgene #62988, a gift from Feng Zhang,[131]) via NEB Quick Ligase (NEB M2200S). The second plasmid (pGW-Amp-Srsf1-N_term-dTAG-HDR) was created by DNA synthesis (Genewiz) in pGW-Amp, containing the ~1 kb PuroR-P2A-2xHA-FKBP1A^{F36V} sequence from Addgene plasmid #91793 [89], flanked on either side with ~1 kb of

sequence homologous to the endogenous *Srsf1* sequence upstream and downstream of its start codon [90].

To generate pCL116(PuroR)-MCP-SRSF1_v2, a coding sequence containing mouse-optimized (IDT Codon Optimization Tool) HA-MCP-SRSF1 was synthesized by Genewiz and inserted into AgeI-and-SalI-digested pCL116(PuroR). The 5' HA-MCP sequence was taken from Addgene plasmid #31230 [132]. An XbaI restriction site was included between MCP and SRSF1 to enable cloning of other MCP-tagged constructs.

To generate pCL116(PuroR)-MCP-X-NLS-3xFLAG constructs, pCL116(PuroR)-MCP-SRSF1_v2 was digested with XbaI and SalI to remove SRSF1, which was replaced with synthesized (Genewiz) DNA fragments containing mouse-optimized (IDT Codon Optimization Tool) X-NLS-3xFLAG sequence, where NLS is the SV40 nuclear localization signal (PKKKRKV), 3xFLAG is three copies of a FLAG tag (MDYKDHDGDYKDHDIDYKDDDDK), and X is one of the following: EGFP (sequence from Addgene plasmid #13031), three copies of the mouse HNRNPK RGG domain (aa 247-330), four copies of the mouse SRSF1 RS domain (aa 196-248), one copy of the mouse SRSF2 RS domain (aa 93-221), or one copy of the SRSF1 RS domain followed by one copy of the SRSF2 RS domain followed by one copy of the SRSF1 RS domain.

Large-scale preparations of all pCARGO-RMCE-*Xist* plasmids were purified from MAX Efficiency Stbl2 cells (Invitrogen 10268019) grown overnight at 30 °C with ampicillin selection (100 µg/mL ampicillin sodium salt, Fisher Scientific BP1760), using the NucleoBond BAC100 Maxiprep Kit (Macherey-Nagel 740579). To limit any potential physical shearing during handling, these large (~29 kb) plasmids below were mixed only by flicking and inverting tubes, taking care to avoid vortexing or extensively mixing by pipet. Pelleted plasmids were resuspended in 120-150 µL of Invitrogen TE Buffer before analyzing purity (A260/A280 > 1.8) with a NanoDrop Lite and DNA concentration with a Qubit 2.0 Fluorometer (dsDNA high-sensitivity or broad-range kit, Thermo Fisher Q32854 and Q32853). Integrity and sequence composition of all pCARGO-RMCE-*Xist* large-scale plasmid preparations were confirmed by diagnostic digestion with KpnI and XhoI, by whole-plasmid Oxford Nanopore sequencing (Plasmidsaurus), and by Sanger sequencing of the regions surrounding Repeat A, Repeat B, and the junction between intron 3 and exon 4, where whole-plasmid sequencing erroneously called an AG-to-AA mutation at the 3′ splice site for several plasmids. Large-scale preparations of all other plasmids were purified from High-Efficiency 5-alpha competent cells (NEB C2987H) or Subcloning Efficiency DH5α competent cells (Invitrogen 18265017) using the PureLink HiPure Plasmid Midiprep Kit (Invitrogen K210004), quantified via Qubit 2.0 fluorometer (dsDNA broad-range kit, Thermo Fisher Q32853), and cloned sequences were confirmed by whole-plasmid Oxford Nanopore sequencing (Plasmidsaurus).

For a complete list of oligonucleotides used in this study, see Table S6.

Generation of RMCE cell lines

Prior to electroporation, plasmid mixtures were prepared containing 6 µg PB-TRE-Cas9 (for transient hygromycin resistance; Addgene #126029 [133]), pOG-Cre (for RMCE-cargo integration; Addgene #233265; [26]; 2.2 µg for mixtures with pCARGO-RMCE-*Xist* and 4.1 µg for mixtures with pCARGO-RMCE-empty for a molar ratio of 1:2.4 for pOG-Cre:pCARGO-RMCE), and 19 µg pCARGO-RMCE-*Xist* (wild-type or mutant) or 6.25 µg pCARGO-RMCE-empty. Each plasmid mixture was ethanol-precipitated, air-dried, and thoroughly resuspended in 10 µL of TE Buffer (Invitrogen 12090015). The 10-µL plasmid mixtures were mixed with 1 million B6/CAST F1-hybrid *Rosa26*-RMCE mESCs (trypsin-dissociated single-cell suspension in 100 µL Neon Buffer R, from Invitrogen MPK10025) and electroporated using a Neon Transfection System (Invitrogen) with one 40-ms pulse of 1000 V before seeding sparsely onto a confluent 10-cm dish of gamma-irradiated drug-resistant (DR4) MEF feeder cells (ATCC SCRC-1045) in growth medium lacking penicillin-streptomycin. Growth medium was not changed the day after electroporation but was replaced daily thereafter. Two days after electroporation, growth medium was replaced with normal growth medium (containing penicillin-streptomycin). Starting three days after electroporation and lasting for four days, cells were selected in growth medium containing 150 µg/mL hygromycin B (Roche 10843555001). Starting seven days after electroporation and lasting for four days, cells were selected in growth medium containing 150 µg/mL hygromycin B and 3 µM ganciclovir (InvivoGen sud-gcv). Nine to eleven days after electroporation, individual colonies (now clearly visible by eye) were picked with aid of a light microscope fitted with a 4X objective, dissociated with 0.125% trypsin-EDTA, and each transferred to a 24-well plate well pre-seeded confluently with gamma-irradiated DR4 MEF feeder cells. Clonal lines of cells were grown until sufficiently confluent (>50%), at which point they were passaged for subsequent genotyping (see "Genotyping PCR" section), preparing cryogenic freeze-down stocks, and introducing the reverse tetracycline-controlled transactivator (rtTA) via plasmid transfection. Every colony screened by genotyping PCR (66 for this study) was positive for having received the cargo cassette.

For introduction of rtTA, four independent RMCE-empty-cargo cell lines, four independent RMCE-*Xist* wild-type cell lines, and two independent RMCE-*Xist* cell lines of each mutant were each seeded at a density of 750,000 cells per well of a 6-well plate. The following day, each cell line was given fresh growth medium and transfected with a plasmid mixture containing 1 µg PB-rtTA (Addgene #126034; [133]) and 1 µg pUC19-piggyBac transposase [133] using Lipofectamine 3000 (Thermo Fisher L3000001). Starting around 24 h after transfection and lasting at least 10

days, cells were selected in growth medium containing 200 µg/mL G418 sulfate (Thermo Fisher 10131035).

Generation of SM33 Δ*Spen* cell lines

To generate SM33 Δ*Spen* cells, 400,000 SM33 cells were seeded in a 6-well plate well and transfected the following day, via Lipofectamine 3000, with a plasmid mixture containing 0.2 µg of PB-TRE-Cas9 (for transient hygromycin resistance; Addgene #126029; [133]) and 0.8 µg of a 1:1:1:1 mixture of pX330 plasmids (Addgene #42230; [134]) expressing Cas9 and 4 different *Spen*-targeting gRNAs.[26] One day after transfection, cells began 3-day treatment with 150 µg/mL hygromycin B to select for transfected cells. Three days after transfection, cells were split and seeded sparsely (500, 1500, 3000, or 6000 cells) onto confluent 10-cm dishes of gamma-irradiated DR4 MEF feeder cells (ATCC SCRC-1045). Ten days after transfection, individual colonies of cells were picked and subsequently expanded and genotyped (see "Genotyping PCR" section) as done above. Of 48 colonies screened by genotyping PCR for *Spen* deletion, eight contained homozygously deleted alleles.

Generation of RMCE-*Xist* WT and ΔCCUGC SRSF2-3xFLAG cell lines

To generate cells with doxycycline-inducible expression of SRSF2-3xFLAG, RMCE-*Xist* WT (clone 4) cells and RMCE-*Xist* ΔCCUGC (clone 1) cells already containing rtTA were seeded at a density of 1.5 million cells in a 6-cm dish. The following day, cells were given fresh growth medium and transfected with a plasmid mixture containing 3 µg pCL116(PuroR)-SRSF2-3xFLAG and 1 µg pUC19-piggyBac transposase [133] using Lipofectamine 3000 (Thermo Fisher L3000001). Starting around 24 h after transfection and lasting at least 5 days, cells were selected in growth medium containing 2 µg/mL puromycin dihydrochloride (Corning 61-385-RA) and 200 µg/mL G418 sulfate (Thermo Fisher 10131035). These cells were cultured in 2 µg/mL puromycin dihydrochloride and 200 µg/mL G418 sulfate to maintain selective pressure.

Generation of RMCE-*Xist* Δ*Spen*, non-targeting gRNA control, and Halo-SPEN-V5 rescue cell lines

To generate RMCE-*Xist* Δ*Spen* cells, 500,000 RMCE-*Xist* Δ*HygroR* cells (chosen to enable addition of expression constructs with hygromycin selection; [130]) were transfected, via Lipofectamine 3000, with a plasmid mixture containing 0.4 µg of pCL116(PuroR) [for transient puromycin resistance] and 1.6 µg of a 1:1:1:1 mixture of pX330 plasmids (Addgene #42230; [134]) expressing Cas9 and 4 different *Spen*-targeting gRNAs [26]. In parallel, RMCE-*Xist* non-targeting

gRNA control cells were generated by transfecting 500,000 RMCE-*Xist* Δ*HygroR* cells, via Lipofectamine 3000, with a plasmid mixture containing 0.4 µg of pCL116(PuroR) [for transient puromycin resistance], 0.8 µg of PB-TRE-Cas9 (Addgene #126029; [133]), and 0.8 µg of PB-rtTA(BsmBI)-non-targeting-gRNA (gRNA sequence from TrueGuide sgRNA Negative Control, non-targeting 1 [Invitrogen A35526]; [135]). Cells were seeded to 6-well plate wells. One day after transfection, cells began 3-day treatment with 2 µg/mL puromycin to select for transfected cells; non-targeting control cells were additionally treated for 1 day with 1 µg/mL dox for transient expression of Cas9. Two days after transfection, cells were split and seeded sparsely (500 or 2000 cells) onto confluent 10-cm dishes of gamma-irradiated DR4 MEF feeder cells (ATCC SCRC-1045). Eight days after transfection, individual colonies of cells were picked and subsequently expanded and genotyped (see "Genotyping PCR" section) as done above. Of 16 colonies screened by genotyping PCR for *Spen* deletion, three contained homozygously deleted alleles.

For introduction of rtTA, two independent RMCE-*Xist* Δ*Spen* cell lines and one RMCE-*Xist* non-targeting gRNA control cell line were each seeded at a density of 500,000 cells per well of a 6-well plate. The following day, each cell line was given fresh growth medium and transfected with a plasmid mixture containing 1 µg PB-rtTA (Addgene #126034; [42]) and 1 µg pUC19-piggyBac transposase [42] using Lipofectamine 3000 (Thermo Fisher L3000001). Starting around 24 h after transfection and lasting at least 10 days, cells were selected in growth medium containing 200 µg/mL G418 sulfate (Thermo Fisher 10131035). These cells and those derived from them were cultured in 200 µg/mL G418 sulfate to maintain selective pressure.

To generate EGFP-Halo or Halo-SPEN-V5 rescue cell lines, RMCE-*Xist* Δ*Spen* (clone 27) cells already containing rtTA were seeded at a density of 400,000 cells per well of a 6-well plate. The following day, cells were given fresh growth medium and transfected with a plasmid mixture containing 1.5 µg pCL116-EGFP-Halo or pCL116(PuroR)-Halo-SPEN-V5 (WT or SPEN mutant) and 0.5 µg pUC19-piggyBac transposase [42] using Lipofectamine 3000 (Thermo Fisher L3000001). Starting around 24 h after transfection and lasting at least 5 days, cells were selected in growth medium containing 150 µg/mL hygromycin B (for EGFP-Halo; Roche 10843555001) or 2 µg/mL puromycin dihydrochloride (for Halo-SPEN-V5; Corning 61-385-RA). These cells were cultured in 150 µg/mL hygromycin B or 2 µg/mL puromycin dihydrochloride (as appropriate) and 200 µg/mL G418 sulfate to maintain selective pressure.

Generation of RMCE-*Xist* FKBP1A^{F36V}-Srsf1 cell lines

To generate RMCE-*Xist FKBP1A^F36V^-Srsf1* cell lines, RMCE-*Xist* WT (clone 4) cells already containing rtTA were seeded at a density of 1.5 million cells in a 6-cm dish. The following day, cells were given fresh growth medium and transfected with a plasmid mixture containing 2 µg PX459_v2-Srsf1-N1-gRNA and 2 µg pGW-Amp-Srsf1-N_term-dTAG-HDR using Lipofectamine 3000 (Thermo Fisher L3000001). One day after transfection, all cells were split and seeded onto a confluent 10-cm dish of gamma-irradiated DR4 MEF feeder cells (ATCC SCRC-1045). Two days after transfection, cells began selection with 2 µg/mL puromycin dihydrochloride (Corning 61-385-RA) and 200 µg/mL G418 sulfate (Thermo Fisher 10131035), which lasted at least 10 days. Seven days after transfection, individual colonies were picked and subsequently expanded and genotyped (see "Genotyping PCR" section) as done above. Of 10 colonies screened by genotyping PCR, two contained homozygously edited *Srsf1* alleles.

Note that at the time of picking colonies, many more colonies – at least several hundred – had survived than the anticipated ~50 to 100 [90], so care was taken to pick individual colonies before further growth would have made their isolation infeasible. This deviation from expectation may be attributable to Lipofectamine-mediated transfection being more efficient than the electroporation method in [90]; as a result, any efforts to repeat this procedure with Lipofectamine-mediated transfection are advised to seed cells at a range of lower densities.

Parallel work to generate C-terminally-tagged *Srsf1-FKBP1A^F36V^* cell lines yielded no edited colonies (0/12 screened), possibly a consequence of poor gRNA (GCTCTCGTACATAAGATGAT) performance or cell inviability upon allele editing. Likewise, work to N-terminally tag *Srsf1* using a different gRNA (GCCGCCGCCGCCATGTCGGG) yielded no edited colonies (0/10 screened).

Generation of RMCE-*Xist* cells for MS2-MCP tethering assay

To generate cells for MS2-MCP tethering, RMCE-*Xist* WT ΔRepA-XB-6x(U-spacer-3xMS2) cells (clone 1 for EGFP and SRSF1-FL; clone 4 for all others) were seeded at a density of 500,000 (clone 1) or 700,000 cells (clone 4) per well of a 6-well plate. The following day, each cell line was given fresh growth medium and transfected with a plasmid mixture containing 0.8 µg of the appropriate pCL116(PuroR) expression plasmid (parental plasmid for EGFP expression), 0.8 µg of PB-rtTA (Addgene #126034; [42]), and 0.4 µg of pUC19-piggyBac transposase [42]), using Lipofectamine 3000 (Thermo Fisher L3000001). Starting around 24 h after transfection and lasting at least 10 days, cells were selected in growth medium containing 2 µg/mL puromycin dihydrochloride (Corning 61-385-RA) and 200 µg/mL G418 sulfate (Thermo Fisher 10131035).

Genotyping PCR

Genomic DNA from individual clonal lines was prepared by first incubating 50-100% confluent 24-well-plate wells of cells (washed once with PBS) with 488 µL of a mixture containing 400 µL gDNA Lysis Buffer (100 mM Tris-HCl pH 8.0, 200 mM NaCl, 5 mM EDTA-NaOH pH 8.0, 0.2% (w/v) SDS), 80 µL of 20 mg/mL Proteinase K solution (Meridian Bioscience BIO-37084), and 8 µL of 5 mg/mL linear acrylamide (Thermo Fisher AM9520). Plates were wrapped in Parafilm and incubated at 55 °C (hybridization oven) for 1-2 h before mixing the contents of each well by pipet, transferring to 1.7-mL centrifuge tubes, and continuing to incubate at 55 °C overnight. The next day, samples were incubated at 100 °C for 1 h before placing on ice, adding 960 µL ice-cold 100% ethanol, inverting several times to mix, and incubating on ice 30 min with occasional inversion. Tubes were then centrifuged at 16,100 x g for 10 min at 4 °C to pellet genomic DNA. DNA samples were washed once with 1 mL ice-cold 80% ethanol, centrifuged again (16,100 x g for 5 min at 4 °C), and DNA pellets were air-dried before resuspending in 100 µL ice-cold TE Buffer (Invitrogen 8019005).

Genotyping PCR was performed using the Apex Taq polymerase kit (Genesee Scientific 42-801B2), in 25-µL reactions containing 1.5 µL genomic DNA, 1x Ammonium Buffer, 1 µM genotyping primers (Table S6), 0.4 mM dNTPs, 1.5 mM $MgCl_2$, and 0.1 U/µL (0.5 µL of 5 U/µL stock) Apex Taq polymerase. Reactions were run on a thermocycler with the following protocol: 95 °C for 3 min, 30 cycles of (95 °C for 30 s, 60 °C for 30 s, 72 °C for 1 min), 72 °C for 5 min. Reaction products were run on a 1% agarose-TAE gel containing ethidium bromide and visualized with a Bio-Rad ChemiDoc MP imager (Ethidium Bromide setting).

RNA-seq

Prior to harvesting RMCE cell lines (with rtTA) were treated with 10-1000 ng/mL doxycycline for 72 hours, with the exception of $FKBP1A^{F36V}$-Srsf1 cells, which were treated for 18 hours with 500 nM dTAG^V-1 (Fisher Scientific 69-145) or equivalent volume of DMSO (Sigma-Aldrich D2650) as a vehicle control, followed by 26 hours with 1000 ng/mL doxycycline hyclate (Sigma-Aldrich D9891-1G) while maintaining the dTAG^V-1 or DMSO treatment. Each 6-cm dish of cells was washed twice with 3 mL ice-cold PBS and harvested with 1 mL TRIzol (Thermo Fisher 15596026). Total RNA was prepared via standard TRIzol-chloroform extraction, per manufacturer's instructions, and resuspended in RNase-free water. RNA concentrations were measured via Qubit 2.0 fluorometer (RNA high-sensitivity or broad-range kit, Thermo Fisher Q32852 and Q10211), and integrity of RNA was confirmed by visualizing rRNA bands on an agarose gel. Single-read, strand-specific (reverse-stranded) libraries for RNA sequencing were prepared using the KAPA RNA HyperPrep Kit with RiboErase (HMR) (Roche 08098140702) and KAPA Unique

Dual-Indexed Adapters (Roche 08861919702) or TruSeq adapters, with each library's input consisting of 9 µL 100 ng/µL total RNA (900 ng) mixed with 1 µL of a 1:250 dilution of ERCC RNA spike-in controls (Thermo Fisher 4456740). RNA fragmentation was performed at 94 °C for 6 min, diluted adapter stocks were 7 µM, and libraries were amplified using 11 cycles of PCR. Libraries were quantified using a Qubit 2.0 fluorometer (dsDNA broad sensitivity kit, Thermo Fisher Q32850), and average fragment sizes were determined by agarose gel electrophoresis. Libraries were pooled and sequenced on an Illumina NextSeq500 platform with a 75-cycle NextSeq 500/550 High Output Kit v2.5 (Illumina 20024906) or Illumina NextSeq1000 platform with a 100-cycle NextSeq 1000/2000 P2 Reagent Kit v3 (Illumina 20046811) or 100-cycle NextSeq 1000/2000 P2 XLEAP-SBS Reagent Kit (Illumina 20100987).

Quantification of *Xist* levels by RNA-seq

For each sample, RNA-seq reads (fastq files) were aligned with STAR (v2.7.9a, [136]) to the mouse GRCm38 (mm10) genome (with reference to GTF file gencode.vM25.basic.annotation.gtf from https://www.gencodegenes.org/mouse/release_M25.html [137], using multi-sample two-pass alignment that incorporated novel splice junctions discovered among all samples in a given experiment [136]. STAR was run with the option "--outFilterMultimapNmax 1" to consider only uniquely mapping reads. For each sample, reads mapping to *Xist* were counted with featureCounts (Subread v2.0.4, [138]) and normalized by dividing by the total number of reads uniquely mapped by STAR.

Allelic RNA-seq, differential gene expression analysis, and measurement of *Xist* silencing activity

Analysis was performed as previously described [41]. RNA-seq reads were aligned to mm10, and independently, also aligned to a version of mm10 that had been modified to incorporate CAST single-nucleotide polymorphisms (SNPs) downloaded from the Sanger Mouse Genomes Project on 7/30/2020 [139]. Alignment was performed with STAR (v2.7.9a, [136]), using multi-sample two-pass mapping and the option "--outFilterMultimapNmax 1" to consider only uniquely mapping reads. Using a custom Perl script (intersect_reads_snps16.pl) and a file containing a list of CAST SNPs (downloaded from the Sanger Mouse Genomes Project on 7/30/2020, [139]), aligned reads in the B6 and CAST SAM files were parsed to identify reads clearly originating from either the B6 or CAST allele (i.e. reads that overlap at least 1 B6/CAST SNP). Reads marked as either B6 or CAST were then assigned to genes using a custom Perl script (ase_analyzer10.pl) and the GTF

file gencode.vM25.basic.annotation.gtf [137]. The number of B6-specific reads for each gene from each sample were then compiled.

DESeq2 (v1.32.0, [47]) was used to identify the set of B6 chr6 genes significantly repressed by wild-type *Xist*. Prior to running DESeq2, a pre-filtering step was used in which genes with fewer than an average of five B6 reads per sample (among four wild-type *Xist* and four empty-cargo control samples) were excluded. Using R, DESeq2 was run to compare the wild-type *Xist* samples against the empty-cargo control samples. Default settings were used except selecting a significance threshold of $p_{adj} < 0.001$, which yielded 204 chr6 genes significantly repressed by WT *Xist*.

To assess *Xist* silencing activity for each sample, B6-specific reads were compiled for each gene as described above. Using Python, the B6-specific reads for each gene were normalized by dividing by the average number of empty-cargo B6-specific reads for that gene; therefore, a value of 1 represents no silencing, and a value of 0 represents complete silencing. The distributions of empty-cargo-normalized expression values for the 204 wild-type *Xist*-repressed genes were visualized by generating swarmplots with the Seaborn package in Python [140].

Xist splicing analysis by RNA-seq

RNA-seq reads were aligned to mm10 using STAR (v2.7.9a, [136]), using multi-sample two-pass mapping and the option "--outFilterMultimapNmax 1" to consider only uniquely mapping reads. The option "--outSAMtype BAM SortedByCoordinate" were included to output alignments in the BAM format. For memory purposes, Samtools (v1.17, [141]) was used to generate smaller BAM files containing only chrX reads as well as corresponding index (.bai) files.

ChrX BAM files and corresponding index files were loaded into IGV (v2.8.9, [142]), and the viewer was navigated to the *Xist* locus. Sashimi plots were generated to visualize strand-specific (reverse-stranded, with "forward" option) junction-spanning reads, setting "Junction coverage minimum" to a value of 2 to remove spurious junction-spanning reads. Extents of aberrant splicing for each sample were calculated by dividing the number of aberrant, exon-1-removing junction-spanning reads by the total number of junction-spanning reads within exon 1 and those spanning exons 1 and 2.

RNA immunoprecipitation (RIP)

After treatment with 1000 ng/mL doxycycline hyclate (24 h for SM33 cells, 72 h for all RMCE cells except 26 h for *FKBP1A^{F36V}*-*Srsf1* cells following 18 h pre-treatment with 500 nM dTAG^V-1), cells were crosslinked with formaldehyde in 10- or 15-cm dishes as follows. Each 10- (15)-cm

dish was washed twice with 6 (10) mL ice-cold PBS before adding 8 (12) mL ice-cold, freshly prepared 0.3% (w/v) formaldehyde (Thermo Scientific Pierce 28906) in PBS and incubating at 4 °C for 30 min. Dishes were then moved to room temp and quenched by adding 1 (1.5) mL 2 M glycine to each dish, gently tilting to mix completely, and incubating for 5 min. Each dish was washed twice with 6 (10) mL ice-cold PBS before harvesting cells by scraping into 5 mL ice-cold 0.1% Tween 20 (Fisher Scientific BP337) in PBS and transferring the cell suspensions to pre-chilled conical tubes on ice. Cells were pelleted by centrifugation (1200 x g, 5 min, 4 °C), supernatants were carefully aspirated, and cell pellets were gently resuspended by pipet in a volume of ice-cold PBS to yield a solution of approximately 10 million cells per mL. (At this point, the sizes of all cell pellets within an experiment were judged relative to one another to ensure that an equivalent number of cells would be used in downstream analyses. For reference, 90% confluent 15-cm dishes routinely yielded approximately 75 million cells.) To pre-chilled 1.7-mL tubes on ice, 1 mL (10-million-cell) aliquots of frequently, gently mixed cell suspension was added. Cells were pelleted by centrifugation (1200 x g, 5 min, 4 °C), supernatants were carefully aspirated, and tubes were snap-frozen in liquid nitrogen before storing at -80 °C.

For each immunoprecipitation, 25 µL well-mixed slurry volume of Protein A/G PLUS-Agarose beads (Santa Cruz sc-2003) was washed three times with 1 mL room-temp 0.5% (w/v) bovine serum albumin (BSA; Fisher Scientific BP1600) in PBS. For each wash, 1.7-mL tubes were gently inverted three times and pelleted by centrifugation (2000 x g, 1 min, room temp), and supernatants were removed carefully, leaving ~10-20 µL to ensure no loss of beads. To washed beads, 300 µL 0.5% (w/v) BSA in PBS was added, followed by an amount of antibody specified in Table S6. Tubes were incubated overnight at 4 °C with end-over-end rotation.

The following day, formaldehyde-crosslinked cell pellets (10 million cells) were thawed on ice and resuspended with 10 gentle up-down strokes of a P1000 pipet in 508 µL ice-cold RIPA Buffer (50 mM Tris-HCl pH 8.0, 150 mM KCl, 5 mM EDTA-NaOH pH 8.0, 1% [v/v] Triton X-100 [Fisher Scientific BP151], 0.5% [w/v] sodium deoxycholate, 0.1% [w/v] sodium dodecylsulfate) containing the following components added immediately before use: 0.5 µL per mL 1 M dithiothreitol (DTT), 10 µL per mL protease inhibitor cocktail (Sigma P8340), and 5 µL per mL 20 U/µL SUPERase-In RNase inhibitor (Invitrogen AM2696). Keeping tubes on ice, each cell suspension was sonicated twice for 30 s with a Sonics Vibra-Cell 130-watt VCX 130 probe-tip sonicator (30% amplitude output), with a 1-min rest on ice between sonication pulses. Tubes were centrifuged (15000 x g, 15 min, 4 °C), and 510 µL of each supernatant (cleared cell lysate) was transferred to a new pre-chilled tube on ice and mixed with 510 µL ice-cold fRIP Buffer (25 mM Tris-HCl pH 7.5, 150 mL KCl, 5 mM EDTA, 0.5% [v/v] IGEPAL CA-630 [Fisher Scientific ICN19859650]) containing the

following components added immediately before use: 0.5 µL per mL 1 M DTT, 10 µL per mL Sigma protease inhibitor cocktail, and 5 µL per mL 20 U/µL SUPERase-In. A 25-µL aliquot of each 1020-µL cleared cell lysate mixture was taken and stored at -20 °C to be processed later as "5% input" samples.

The antibody-bound beads from the day before were washed twice with 1 mL ice-cold fRIP Buffer; for each wash, 1.7-mL tubes were gently inverted three times and pelleted by centrifugation (2000 x g, 1 min, 4 °C), and supernatants were carefully removed, leaving ~10-20 µL to ensure no loss of beads. To each tube of washed beads, 490 µL of an ice-cold 1:1 mixture of RIPA Buffer and fRIP Buffer containing freshly added 1 M DTT (0.5 µL per mL), Sigma protease inhibitor cocktail (10 µL per mL), and 20 U/µL SUPERase-In (5 µL per mL) was added. To each tube, 490 µL of appropriate cleared cell lysate mixture (equivalent to ~5 million cells) was added, and tubes were incubated overnight at 4 °C with end-over-end rotation. In the case of the SPEN RIP-qPCR experiments in Figure 2B, the remaining ~500 µL cleared cell lysate mixtures were snap-frozen in liquid nitrogen and stored at -80 °C to use later in the SRSF1 RIP-qPCR experiments in Figure 5A, using the same sample-matched input samples for normalization.

The next day, beads were pelleted by centrifugation (2000 x g, 1 min, 4 °C), and supernatants were carefully removed. The beads were washed once with 1 mL ice-cold fRIP Buffer and resuspended in 1 mL ice-cold Pol II ChIP Buffer (50 mM Tris-HCl pH 7.5, 140 mM NaCl, 1 mM EDTA-NaOH pH 8.0, 1 mM EGTA-NaOH pH 8.0, 1% [v/v] Triton X-100, 0.1% [w/v] sodium deoxycholate, 0.1% [w/v] sodium dodecylsulfate) before carefully transferring to a new 1.7-mL tube, ensuring no loss of beads. Samples were rotated end-over-end at 4 °C for five min, pelleted by centrifugation (2000 x g, 1 min, 4 °C), and supernatants were carefully removed. Samples were washed twice more with 1 mL Pol II ChIP Buffer, once with 1 mL High-Salt Pol II ChIP Buffer (50 mM Tris-HCl pH 7.5, 500 mM NaCl, 1 mM EDTA-NaOH pH 8.0, 1 mM EGTA-NaOH pH 8.0, 1% [v/v] Triton X-100, 0.1% [w/v] sodium deoxycholate, 0.1% [w/v] sodium dodecylsulfate), and once with 1 mL LiCl Buffer (20 mM Tris pH 8.0, 1 mM EDTA-NaOH pH 8.0, 250 mM LiCl, 0.5% [v/v] IGEPAL CA-630, 0.5% [w/v] sodium deoxycholate); each wash included a 5-min end-over-end rotation at 4 °C. In the final wash, samples were transferred to a new 1.7-mL tube. The final wash was carefully removed, first with a P1000 pipet and then with a P20 pipet, to ensure no loss of beads while leaving a volume of ~10 µL LiCl Buffer or less on the beads.

Input-sample Reverse Crosslinking buffer was prepared on ice by mixing, for each sample, 33 µL 3X Reverse Crosslinking Buffer (3x PBS, 6% [w/v] N-lauroylsarcosine, 30 mM EDTA-NaOH pH 8.0), 41 µL RNase-free water, 5 µL 100 mM DTT, 1 µL 20 U/µL SUPERase-In, and 10 µL 20 mg/mL proteinase K (Thermo Fisher 25530049). Bead-sample Reverse Crosslinking buffer was

prepared on ice by mixing, for each sample, 33 µL 3X Reverse Crosslinking Buffer (3x PBS, 6% [w/v] N-lauroylsarcosine, 30 mM EDTA-NaOH pH 8.0), 66 µL RNase-free water, 5 µL 100 mM DTT, 1 µL 20 U/µL SUPERase-In, and 10 µL 20 mg/mL proteinase K. The 25-µL 5% input samples were thawed on ice and mixed thoroughly by pipet with 90 µL Input-sample Reverse Crosslinking buffer. To washed beads, 115 µL bead-sample reverse crosslinking buffer was added forcefully to resuspend the beads, but the mixture was not otherwise mixed. To reverse crosslinks, samples were incubated for 1 hr at 42 °C, 1 hr at 55 °C, and 30 minutes at 65 °C. During these incubations, bead-containing samples had their beads resuspended every 15 to 30 min by forcefully ejecting ~100 µL of the supernatant onto the settled beads, a practice done to avoid losing beads to the inside of pipet tips. After heating, 1 mL TRIzol was added to each sample, and samples were inverted several times to mix thoroughly and stored at -80 °C.

The TRIzol-containing samples were thawed at room temp, 200 µL chloroform was added to each, and samples were inverted and vortexed for 3 to 5 s (or alternatively, shaken vigorously all together in a "tube rack sandwich" for 15 s), and centrifuged at 15,000 x g for 15 min at 4 °C. A 580-µL volume of the aqueous phase was mixed with 580 µL 100% ethanol, and samples were inverted, vortexed, and pulsed down before applying (in two steps to accommodate the entire volume) to Zymo-Spin IC Columns (from RNA Clean & Concentrator-5 Kit, Zymo R1013) and centrifuging (16,100 x g, 30 s, room temp). A 400-µL volume of RNA Wash Buffer (Zymo R1013) was added to each column and centrifuged (16,100 x g, 30 s, room temp). To each column, a 40-µL volume of a mixture containing 5 µL DNase I (1 U/µL) and 35 µL DNA Digestion Buffer (Zymo R1013) was added; samples were incubated at room temp for 20 min. A 400-µL volume of RNA Prep Buffer (Zymo R1013) was then added, and columns were centrifuged (16,100 x g, 30 s, room temp). Columns were washed again as before with 700 µL RNA Wash Buffer and then with 400 µL RNA Wash Buffer. The flowthrough was discarded and columns spun again for 2 minutes to remove all traces of wash buffer. Columns were transferred to a clean 1.7-mL tube, 15 µL RNase-free water was added to each, and after incubating at room temp for 5 min, samples were centrifuged (16,100 x g, 1 min, room temp). Eluted RNA was stored at -80 °C for subsequent analysis by RT-qPCR (RIP-qPCR) or sequencing (RIP-seq).

Replicate RNA immunoprecipitation experiments used cell pellets that were crosslinked on different days, and the immunoprecipitation, RNA isolation, and RT-qPCR steps were repeated on different days. The exception to this was the SPEN RIP-qPCR experiment in Figure 5F, in which two independent clonal lines of *FKBP1A^F36V^*-*Srsf1* cells were crosslinked and analyzed on the same days.

Reverse transcription and quantitative PCR analysis of RIPs (RIP-qPCR)

Reverse transcription was performed with the Applied Biosystems High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher 4368814) according to manufacturer instructions: 20-µL randomer-primed reverse transcription reactions were prepared in 0.2-mL PCR strip-tubes on ice, each containing 0.5 µL 40 U/µL RNaseOUT RNase Inhibitor (Thermo Fisher 10777019) and 5 µL of column-purified RNAs from 5% input or 100% RIP samples. Reverse transcription was performed with the following thermal cycler program: 25 °C for 10 min, 37 °C for 120 min, 85 °C for 5 min, hold at 4 °C. All reverse transcription products were first diluted 2-fold by thoroughly mixing with 20 µL nuclease-free water. For robust quantification via qPCR, standard curves were constructed by preparing 4-fold serial dilutions from each 5% input sample (10 µL previous standard mixed with 30 µL nuclease-free water), generating a total of 6 standards per standard curve.

To set up qPCR, a master mix was generated that contained, for each 10-µL qPCR reaction: 5 µL iTaq Universal SYBR Green Supermix (Bio-Rad 1725124), 3 µL nuclease-free water, 0.5 µL 10 µM forward primer, and 0.5 µL 10 µM reverse primer (see Table S6 for primer sequences). In Hard-Shell 96-Well PCR Plates (Bio-Rad HSP9601), qPCR reactions were prepared in technical triplicate by combining 9 µL qPCR master mix and 1 µL of standard-curve or RIP reverse transcription product. Standard-curve qPCR reactions from each 5% input and qPCR reactions for each 5% input's corresponding RIP(s) were always run on the same plate. Plates were firmly sealed with Microseal 'B' PCR Plate Sealing Film (Bio-Rad MSB1001), mixed several times by inversion and low intensity vortexing, and briefly pulsed down in a centrifuge (up to 3000 x g) to collect the volumes and remove bubbles. Reactions were analyzed on a Bio-Rad C1000 Touch Thermal Cycler equipped with a CFX96 Real-Time System with the following standard program: 95 °C for 10 min, 40 cycles of (95 °C for 15 s, 60 °C for 30 s [annealing], 72 °C for 30 s, plateread), followed by melt curve analysis: 95 °C for 10 s, 65 °C for 31 s, 61 cycles of (65 °C for 5 s +0.5 °C/cycle Ramp 0.5 °C/s, plateread). A variant of this standard program in which the annealing step was set to 56 °C was used for Repeat A-targeting qPCR primers to accommodate their lower annealing temperatures (see Table S6). All qPCR primer pairs were confirmed via agarose gel electrophoresis to specifically amplify a single product of the expected size. Data were analyzed using Bio-Rad CFX Manager software (v3.1), first setting "Cq Determination Mode" to "Regression" and then using each standard curve (Cq vs. % input, with the average value from the most concentrated standard set to 5% input) to convert Cq values to "% input" values for each technical replicate of each RIP sample. Each qPCR technical triplicate value was plotted as a dot in GraphPad Prism (v9.5.0), with the average values being represented by bars. Significance was

determined via two-tailed t-tests comparing the triplicate-averaged values from independent biological replicates.

RIP-seq library preparation and sequencing

Single-read, strand-specific (reverse-stranded) libraries for RNA sequencing were prepared using the KAPA RNA HyperPrep Kit with RiboErase (HMR) (Roche 08098140702) and KAPA Unique Dual-Indexed Adapters (Roche 08861919702) or TruSeq adapters, with each library's input consisting of 9 µL column-purified RNA from RIP and 1 µL of a 1:250 dilution of ERCC RNA spike-in controls (Thermo Fisher 4456740). RNA fragmentation was performed at 85 °C for 5 min, diluted adapter stocks were 1.5 µM, and libraries were amplified using 13 to 21 cycles of PCR depending on the sample, with the number of cycles determined by qPCR to compare against positive-control pre-amplification libraries that had been previously amplified, quantified, and sequenced. Libraries were quantified using a Qubit 2.0 fluorometer (dsDNA broad sensitivity kit, Thermo Fisher Q32850), and average fragment sizes were determined by agarose gel electrophoresis. Libraries were pooled and sequenced on an Illumina NextSeq500 platform with a 75-cycle NextSeq 500/550 High Output Kit v2.5 (Illumina 20024906) or Illumina NextSeq1000 platform with a 100-cycle NextSeq 1000/2000 P2 Reagent Kit v3 (Illumina 20046811) or 100-cycle NextSeq 1000/2000 P2 XLEAP-SBS Reagent Kit (Illumina 20100987).

RIP-seq wiggle track generation

RIP-seq reads were aligned to mm10 using STAR (v2.7.9a, [136]), using single-pass mapping and the option "--outFilterMultimapNmax 1" to consider only uniquely mapping reads. Output SAM files from each alignment were parsed by strand using Samtools (v1.15 or v1.20): reads from Samtools view -F 0x10 were retained as negative-stranded reads, and reads from Samtools view -f 0x10 were retained as positive-stranded reads. For each strand-specific SAM file, wiggle files were generated using the custom Perl script bigsam_to_wig_mm10_wcigar2.pl, which binned reads into 50-nt bins. Wiggle files were uploaded to the UCSC Genome Browser to visualize data. For viewing reads over *Xist*, browser tracks were configured such that "Data view" was set to "auto-scale to data view" and "Windowing function" was set to "mean." Views over *Xist* (which is on the mm10 negative strand) were exported in PDF file format and flipped horizontally in Adobe Illustrator to display data from 5′ to 3′. Track heights were RPM-normalized by dividing the auto-scaled heights (reads per 50-nt bin) by the total number of reads (in millions) uniquely mapped by STAR for that sample.

To quantify reads over Repeat A or the entirety of *Xist*, featureCounts (Subread v2.0.4, [138]) was used. For Repeat A reads, a SAF file was generated with a region defined as chrX:103482476-103482926, and featureCounts was run with options "-s 2 -F SAF -a Repeat_A.saf." For *Xist* reads, featureCounts was run with options "-t exon -g gene_id -a mm10_genes.gtf," and *Xist* reads were collected with the Unix command grep 'Xist'.

RIP-seq peak calling

SPEN and Halo RIP-seq FASTQ files were aligned with STAR (2.7.10a, [136]) to a STAR genomic index built from the primary assembly of the GRCm38 (mm10) genome with reference to the GENCODE vM25 basic annotation GTF. Additional flags of "--outMultimapperOrder Random" and "--outSAMmultNmax 1" were used. Alignments were then filtered to remove multimapping reads (MAPQ scores less than 30) with Samtools (v1.21) view -q 30. For all RIP-seq experiments, the filtered alignments were split by strand, reads from Samtools view -F 0x10 were retained as negative-stranded reads, and reads from Samtools view -f 0x10 were retained as positive-stranded reads. The strand information was then randomly re-assigned through a custom Perl script (macs_strand_rand_sam.pl),[130] and these strand-randomized alignments were converted to BAM files and provided as input to the MACS2 (v2.2.7.1; [143]) callpeak function with the flags "--keep-dup all", "--broad", "--broad-cutoff 0.3", and "--max-gap 76". To identify the number of reads under each peak, the MACS2 peaks for each RIP-seq experiment were converted from broadPeak to SAF file formats and then provided as input to featureCounts (Subread v2.0.6, [138]) with the additional flags "-F SAF and -s 2," as the data is reversely stranded. Additionally, the same peaks were also given to featureCounts with the appropriate control sample (SM33 Δ*Spen* or RMCE-*Xist* Δ*Spen* no-rescue) alignment. The number of reads under each peak was converted to RPM and the peak set was filtered to retain only peaks where the RPM of each replicate was greater than two times the RPM value of the control in that same peak region.

RIP-seq motif analysis

To identify motif enrichment that excluded RIP signal from *Spen* and *Xist* transcripts, the coordinates for both genes were used to make a miniature custom GTF file that was used in BEDTools intersect (v2.29, [144]) as -b input. The -v flag was also used to exclude intersecting peaks, resulting in a new final set of peaks in BED file format. Next, scaled signal over control for each peak region was calculated by squaring the RIP RPM value and dividing that by the sum of the RIP RPM and the control-sample RIP RPM. Peaks were then sorted by decreasing numerical

scaled signal values, and the top 1500 peaks were retained. The sequences for the top 1500 ranked peaks were extracted from the GRCm38 primary assembly genome FASTA by BEDTools getfasta command with the -s flag. A paired set of length-matched control sequences was generated for the top 1500 peak sequences of each RIP experiment by calculating the nucleotide probabilities from the nucleotide content of the genome, which weighted random nucleotide selection in random sequence creation. Finally, enriched motifs were established with STREME (MEME Suite v5.2.0, [145]) for the top 1500 peak sequences for each RIP experiment individually, with the corresponding length-matched control sequences provided with the -n flag, a maximum window of --maxw 12, minimum window of --minw 4, --nmotifs 20, and the --rna and -p flags.

RIP-seq target ranking by gene

To determine which genes had transcripts most associated with SPEN in the WT, RRM1-4, and RRM2-4 Halo RIPs, the RIP peaks over the *Spen* locus were removed in the same fashion as the *Spen* and *Xist* peaks from the motif methods above. Next, a limited version of the GENCODE vM25 basic annotation GTF was created that removed all non-gene feature lines and this limited GTF was provided to BEDTools intersect with the non-*Spen* peaks and the -wo and -s flags. The resultant output file contained one line for every peak that intersected a gene, the information of both the gene and the peak coordinates, and the number of nucleotides in that intersection. A custom Python script (rank_genes_by_peak_signal_3_31_25.py) was used to calculate the scaled signal over control for each peak as was calculated in the motif methods previously (the RIP RPM value squared and divided by the sum of the RIP RPM and the control-sample RIP RPM). For each gene, the scaled signals for all peaks over the annotated gene bounds were fractionally summed based on the percentage of the peak overlapping the gene bounds. Genes were sorted by their representative summed signal value and reported in Table S3.

Protein purification

To 400 mL pre-warmed lysogeny broth (LB; Research Products International L24041) containing 100 µg/mL ampicillin sodium salt (Fisher Scientific BP1760-25) and 34 µg/mL chloramphenicol (Acros Organics 227920250) in a 1-L flask, 4 mL of overnight culture of Rosetta *Escherichia coli* cells (MilliporeSigma 70953) transformed with pDD001-SPEN-RRM2-4 or pDD001-SRSF1 RRM1-2 was added. Cultures were grown at 37 °C with 180 rpm shaking until OD_{600} reached 0.4-0.6, after which cultures were cooled to 17 °C, and then isopropyl β-D-1-thiogalactopyranoside (IPTG; Sigma I6758) was added to a final concentration of 0.5 mM.

After ~16-h induction at 17 °C with 180 rpm shaking, cells were pelleted by centrifugation (5000 x g, 10 min, 4 °C) and gently resuspended by swirling in 20 mL GST Lysis Buffer (20 mM Tris-HCl pH 7.5, 20 mM NaCl, 1% [v/v] Triton X-100; containing freshly added SigmaFast EDTA-free protease inhibitor cocktail [1 tablet per 100 mL]) with freshly added 2 mM DTT and 2 mM phenylmethylsulfonyl fluoride (PMSF). Resuspended cells were lysed on ice, with 6 30-s pulses of a Sonics Vibra-Cell 130-watt VCX 130 probe-tip sonicator (50% amplitude output) and a 30-s rest on ice between each pulse. Lysates were cleared by centrifugation (36635 x g, 30 min, 4 °C) and then added to 1 mL slurry volume of Pierce Glutathione Agarose Beads (Thermo Fisher 16100) pre-equilibrated in GST Lysis Buffer in a 50-mL conical tube on ice. Lysates were incubated with beads for 30 min at 4 °C with end-over-end rotation and then gently pelleted by centrifugation (500 x g, 1 min, room temp). Unbound lysate supernatants were carefully removed from the beads, and then the beads were washed 5 min per wash at 4 °C with end-over-end rotation and pelleted as before, using 10 mL of the following buffers per wash: twice with GST Wash Buffer A (20 mM Tris-HCl pH 7.5, 300 mM NaCl), twice with GST Wash Buffer B (20 mM Tris-HCl pH 7.5, 1 M NaCl), and then twice again with GST Wash Buffer A. All but ~500 µL the final wash buffer was removed from the beads, the beads were resuspended in the remaining buffer and transferred to a 1.7-mL tube, and the remaining buffer was carefully removed after pelleting by centrifugation (1000 x g, 2 min, 4 °C). To the beads, 500 µL Cleavage Buffer (20 mM Tris-HCl pH 7.5, 100 mM NaCl, 0.01% [v/v] Triton X-100, 10% [v/v] glycerol, 5 mM DTT [freshly added]) was added, followed by 10 µL PreScission Protease (Cytiva 27-0843-01). SBP-tagged proteins were cleaved from the bead-bound GST by incubation at 4 °C with end-over-end rotation for ~18 h.

Beads were pelleted by centrifugation (2000 x g, 2 min, 4 °C), and supernatants were carefully collected and transferred to a new pre-chilled tube on ice. To clear residual beads from the supernatants, tubes were centrifuged again (16100 x g, 5 min, 4 °C), and supernatants (final protein preparations) were transferred to pre-chilled tubes on ice, aliquoted to small volumes, snap-frozen with liquid nitrogen, and stored at -80 °C. To assess purity and measure concentration, each protein preparation was run alongside a standard curve of known amounts of BSA (Fisher Scientific BP1600) on a pre-cast Bio-Rad TGX SDS-PAGE gel in Tris-glycine-SDS running buffer (2.5 mM Tris, 19.2 mM glycine, 0.01% [w/v] SDS, pH 8.3) at 150 V. The gel was stained with GelCode Blue (Thermo Fisher 24590), destained with water, and imaged with a Bio-Rad ChemiDoc MP imager (Coomassie Blue setting). Densitometry analysis was performed with Bio-Rad Image Lab v6.1 software to determine band intensities of SBP-tagged proteins and the

sums of band intensities in each BSA standard; using Microsoft Excel software, SBP-tagged protein concentrations were calculated by fitting to a best-fit line of the BSA standards.

In vitro transcription (IVT) of biotinylated and unlabeled RNA

Double-stranded DNA templates for IVT were generated by PCR using Q5 High-Fidelity DNA Polymerase (NEB M0491), pTETRISv1-*Xist*-2kb wild-type/mutant as template, and primers specific to *Xist* that add the SP6 promoter sequence upstream of Repeat A (Table S6). DNA templates were run on an agarose-TAE gel containing ethidium bromide, and excised DNA bands were purified using the GenCatch Gel Extraction Kit (Epoch Life Science 2260250). DNA concentration was measured with a NanoDrop Lite.

To generate internally biotinylated RNA, IVT was performed overnight (~16-24 h) at 37 °C using the MEGAscript SP6 Transcription Kit (Thermo Fisher AM1330) following manufacturer instructions, with the addition of biotinylated UTP (MilliporeSigma 11388908910 or TriLink Biotechnologies N-5005-1) to a final concentration of 0.5 mM (such that the molar ratio of biotinylated UTP to UTP was 1:10) and 0.5 pmol DNA template per 20 µL of reaction. To generate large-scale amounts of RNA, reaction volumes were frequently scaled up to 100 µL. IVT products were treated with TURBO DNase (0.8 µL per 20 µL reaction, 15 min at 37 °C) and purified via standard TRIzol-chloroform extraction, per manufacturer instructions (Thermo Fisher 15596026). RNA concentration was measured with a NanoDrop Lite, and purity was assessed by urea-PAGE with Promega Diamond (Promega H1181) staining. RNA was stored in small aliquots at -80 °C.

To generate unlabeled RNA for RBNQ, the same procedure was performed with the following differences: biotinylated UTP was omitted, RNA was purified using the MEGAclear Transcription Clean-Up Kit (Thermo Fisher AM1908) following manufacturer instructions except that elution was performed at 37 °C to preserve native RNA folding, and RNA concentration was measured with a Qubit 2.0 fluorometer (RNA high-sensitivity or broad-range kit, Thermo Fisher Q32852 and Q10211).

Equilibrium RNA bind-n-qPCR (RBNQ)

The following protocol is an adaptation of one published previously [53], modified to enable equilibrium assessment of binding and accurate comparison to input. SBP-tagged proteins were first diluted into room-temp RBNQ Binding Buffer with Random RNA (3 mM $MgCl_2$, 25 mM Tris-HCl pH 8.05, 150 mM KCl, 1 mg/mL BSA [Thermo Fisher AM2616], 2 U/mL SUPERase-In [Thermo Fisher Scientific AM2694], 50 nM random 16-nt RNA [IDT, bases randomized by machine-mixing, purified by standard desalting]) to a range of different protein concentrations (~1-

1400 nM). Using a room-temp magnetic separator for buffer exchanges, a volume of 250 µL well-mixed Dynabeads MyOne Streptavidin T1 (Thermo Fisher 65602) was washed once with 250 µL RNase-free water, twice with 250 µL RBNQ Blocking Buffer (3 mM MgCl$_2$, 25 mM Tris-HCl pH 8.05, 150 mM KCl, 1 mg/mL BSA [Thermo Fisher AM2616], 2 U/mL SUPERase-In [Thermo Fisher AM2694], 1 mg/mL yeast tRNA [Thermo Fisher AM7119]), and twice with 250 µL RBNQ Binding Buffer with Random RNA. Beads were then resuspended in 125 µL RBNQ Binding Buffer with Random RNA, and 10 µL of the well-mixed slurry was aliquoted to each of 12 1.7-mL tubes. (Volumes of beads and washes were scaled proportionately for experiments with more than 12 binding reactions.) To each tube, 30 µL of SBP-tagged protein (at various concentrations) was added and gently mixed by pipet, and tubes were incubated for 10 min (in a time-staggered manner from this point forward to ensure consistent handling) at room temp with end-over-end rotation for proteins to bind to the beads. Tubes were placed on the magnetic separator; supernatants were removed; beads were carefully resuspended in 30 µL of 0.1 nM in-vitro-transcribed, unlabeled Repeat A RNA in RBNQ Binding Buffer with Random RNA; and tubes were incubated for 30 min at room temp with end-over-end rotation to enable RNA-protein binding reactions to equilibrate. Meanwhile, 30 µL of each 0.1 nM in-vitro-transcribed, unlabeled Repeat A RNA was transferred to prechilled tubes on ice for "input RNA" samples. Keeping uniform, time-staggered treatment, tubes were placed on the magnetic separator, unbound supernatants were carefully and completely removed, beads were resuspended in 100 µL of Elution Buffer (25 mM Tris-HCl pH 8.05, 4 mM D-biotin [Thermo Fisher B1595], 8% [v/v] dimethyl sulfoxide [DMSO]), and tubes were incubated for 30 min at 37 °C. Tubes were placed on the magnetic separator, and "eluate RNA" supernatants were transferred to pre-chilled tubes on ice. "Input RNA" samples were brought up to 100 µL by adding 70 µL Elution Buffer and mixing well.

For RBNQ assays examining differences in protein binding to wild-type versus mutant Repeat A RNA, the same protocol was followed except that the volumes of Dynabeads and bead washes were cut in half (i.e., 10 µL Dynabead slurry was used per binding reaction and washes were with 125 µL) and that a single concentration of protein was used (near the pre-determined K$_d$ for wild-type RNA). Replicate RBNQ experiments used protein preps that were purified on separate days; protein concentrations of the independent preps were measured at the same time to ensure accuracy and precision of RBNQ data.

Reverse transcription was performed with the Applied Biosystems High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher 4368814) according to manufacturer instructions: 20-µL randomer-primed reverse transcription reactions were prepared in 0.2-mL PCR strip-tubes on ice, each containing 0.5 µL 40 U/µL RNaseOUT RNase Inhibitor (Thermo Fisher 10777019) and

5 µL of input or eluate RNA. Reverse transcription was performed with the following thermal cycler program: 25 °C for 10 min, 37 °C for 120 min, 85 °C for 5 min, hold at 4 °C. All reverse transcription products were first diluted 2-fold by thoroughly mixing with 20 µL nuclease-free water. For robust quantification of binding-curve RBNQ experiments via qPCR, standard curves were constructed by preparing 2-fold serial dilutions from each input sample (20 µL previous standard mixed with 20 µL nuclease-free water), generating a total of 8 standards per standard curve. For fixed-concentration RBNQ comparing WT and mutant Repeat A, 6-standard standard curves were constructed via 2.5-fold serial dilutions from each input sample.

To set up qPCR, a master mix was generated that contained, for each 10-µL qPCR reaction: 5 µL iTaq Universal SYBR Green Supermix (Bio-Rad 1725124), 3 µL nuclease-free water, 0.5 µL 10 µM forward primer, and 0.5 µL 10 µM reverse primer (see Table S6 for oligonucleotide sequences). In Hard-Shell 96-Well PCR Plates (Bio-Rad HSP9601), qPCR reactions were prepared in technical triplicate (technical duplicate for standard-curve samples in fixed-concentration WT vs. mutant Repeat A RBNQ experiments) by combining 9 µL qPCR master mix and 1 µL of input or eluate reverse transcription product. Standard-curve qPCR reactions from each input and corresponding eluate(s) were always run on the same plate. Plates were firmly sealed with Microseal 'B' PCR Plate Sealing Film (Bio-Rad MSB1001), mixed several times by inversion and low intensity vortexing, and briefly pulsed down in a centrifuge (up to 3000 x g) to collect the volumes and remove bubbles. Reactions were analyzed on a Bio-Rad C1000 Touch Thermal Cycler equipped with a CFX96 Real-Time System with the following standard program: 95 °C for 10 min, 40 cycles of (95 °C for 15 s, 56 °C for 30 s, 72 °C for 30 s, plateread), followed by melt curve analysis: 95 °C for 10 s, 65 °C for 31 s, 61 cycles of (65 °C for 5 s +0.5 °C/cycle Ramp 0.5 °C/s, plateread). All qPCR primer pairs were confirmed via agarose gel electrophoresis to specifically amplify a single product of the expected size. Data were analyzed using Bio-Rad CFX Manager software (v3.1), first setting "Cq Determination Mode" to "Regression" and then using each standard curve (Cq vs. % input, with the average value from the most concentrated standard set to 100% input) to convert Cq values to "% input" values for each technical replicate of each eluate sample. For binding-curve RBNQ experiments, input-normalized values from each of two experimental replicates were plotted together in GraphPad Prism (v9.5.0), with the average values and standard deviation of qPCR technical triplicates being represented by dots and error bars, respectively. The equilibrium dissociation constant K_d and Hill slope h were determined in GraphPad Prism by fitting to the equation $Y=B_{max}*X^h/(K_d^h + X^h)$ with Analyze > XY analyses > nonlinear regression (curve fit) > Binding - Saturation > Specific binding with Hill Slope. For fixed-concentration WT vs. mutant Repeat A RBNQ experiments, each qPCR technical triplicate value

was plotted as a dot in GraphPad Prism (v9.5.0), with the average values being represented by bars. Significance was determined via two-tailed t-tests comparing the triplicate-averaged values from independent biological replicates.

Preparation of nuclear extracts for biotinylated RNA pulldown

Between 14 and 20 75-85% confluent 15-cm dishes of E14 mESCs were each washed three times with 10 mL ice-cold PBS before harvesting by scraping into 1.5 mL ice-cold PBS. Cells were transferred to pre-chilled conical tubes and pelleted by centrifugation (1136 x g, 5 min, 4 °C), supernatants were removed, and cells were gently resuspended by serological pipet collectively in 7 mL ice-cold PBS and consolidated in one 15-mL conical tube before pelleting by centrifugation (1136 x g, 5 min, 4 °C). The volume of the cell pellet was estimated (~1.2-1.5 mL), and cells were resuspended by serological pipet in 2 volumes of ice-cold NE Buffer A (10 mM HEPES-KOH pH 7.9, 1.5 mM MgCl$_2$, 10 mM KCl, with the following added fresh: 0.5 mM DTT, 0.5 mM PMSF, 1:200 protease inhibitor cocktail [Sigma P8340]). IGEPAL CA-630 was added to a final concentration of 0.1% (v/v), and the cell suspension was gently inverted to mix before incubating on ice for 5 min. The cell suspension was placed in a pre-chilled Dounce homogenizer and homogenized with 10 up-down strokes of a tight-fitting pestle, taking care avoid introducing bubbles. The mixture was pelleted by centrifugation (2148 x g, 4 min, 4 °C), and the supernatant (cytoplasmic extract) was removed. To prepare nuclear extract, the pellet of nuclei was resuspended by serological pipet in 1.5 original-cell-pellet-volumes of ice-cold NE Buffer C (20 mM HEPES-KOH pH 7.9, 25% [v/v] glycerol, 420 mM KCl, 1.5 mM MgCl$_2$, 0.2 mM EDTA-NaOH pH 8.0, with the following added fresh: 0.5 mM DTT, 0.5 mM PMSF, 1:200 protease inhibitor cocktail [Sigma P8340]) and incubated end-over-end at 4 °C for 30 min. The mixture was pelleted by centrifugation (5000 x g, 5 min, 4 °C), and supernatant (nuclear extract) was transferred to a pre-chilled tube on ice. From the pellet of nuclei and nuclear debris, the extraction process was repeated with another 1.5 original-cell-pellet-volumes of ice-cold NE Buffer C. The nuclear extract samples were cleared by centrifugation (16,100 x g, 15 min, 4 °C), and the supernatants were pooled together to a volume of ~3-6 mL. Using Slide-A-Lyzer 10-kDa MWCO devices (Thermo Fisher 88404), cleared nuclear extracts were dialyzed twice at 4 °C into NE Buffer D (20 mM HEPES-KOH pH 7.9, 20% [v/v] glycerol, 100 mM KCl, 0.2 mM EDTA-NaOH pH 8.0, with 0.5 mM DTT added fresh); one dialysis step lasted 2-3 hours and the other lasted ~18 hours. To remove precipitated insoluble material, dialyzed nuclear extracts were pelleted by centrifugation (16,100 x g, 15 min, 4 °C), and supernatants (final nuclear extracts) were pooled, measured by Bradford assay (Bio-Rad 5000006) against a standard curve of BSA to determine protein concentration

(~2.5-3 µg/µL), and snap-frozen in liquid nitrogen before storing at -80 °C. Aliquots were shipped on dry ice for proteomics analysis of nuclear extracts (biotinylated RNA pulldown inputs).

Biotinylated RNA pulldown

The following protocol is a modification of one published previously [2], modified to increase the amount of RNA and protein, to clarify crosslinking conditions, and to detail the elution of proteins with RNase A. In a pre-chilled tube on ice for each pulldown sample, a volume of nuclear extract containing 1100 µg protein was brought to 500 µL with NE Buffer D, after which 1 mL ice cold Incubation Buffer (150 mM NaCl, 15 mM Tris-HCl pH 7.5, 0.5 mM EDTA-NaOH pH 8.0, 0.025% [v/v] IGEPAL CA-630, with the following added fresh: 1 mM DTT, 1:100 protease inhibitor cocktail [Sigma P8340], 100 U/mL SUPERase-In RNase inhibitor [Thermo Fisher AM2696]) and 11 µg internally biotinylated, in-vitro-transcribed RNA were added. Tubes were incubated end-over-end at 4 °C overnight.

The next day, for each pulldown, 25 µL slurry volume of Dynabeads M-280 streptavidin beads (Thermo Fisher 11205D) was washed twice with 500 µL ice-cold Incubation Buffer (without DTT, protease inhibitor cocktail, and SUPERase-In) with aid of a magnetic separator, keeping the beads in the second wash on ice until ready for addition of crosslinked RNA-protein mixtures. The RNA-protein mixtures were transferred to individual wells of a pre-chilled 6-well cell culture plate (Genesee Scientific 25-105) on ice. To induce crosslinks with UV light, the plate on ice (with lid removed) was placed in a Stratagene UV Stratalinker 1800 fitted with 254-nm bulbs and irradiated with the auto-crosslink setting (2500 x 100 µJ/cm^2; value of 2500 entered on the digital screen). Washed beads were resuspended in crosslinked RNA-protein mixtures and incubated end-over-end at 4 °C overnight.

The next day, the beads were washed (5 min per wash, end-over-end at 4 °C) with 500 µL of the following buffers per wash: twice with ice-cold PD Wash Buffer A (150 mM NaCl, 15 mM Tris-HCl pH 7.5, 0.5 mM EDTA-NaOH pH 8.0, 1 mM DTT, 1:100 protease inhibitor cocktail [Sigma P8340]), twice with room-temp (to prevent SDS precipitation) PD Wash Buffer B (450 mM NaCl, 15 mM Tris-HCl pH 7.5, 0.5 mM EDTA-NaOH pH 8.0, 0.025% [v/v] IGEPAL CA-630, 0.05% [w/v] SDS, 0.1 mM DTT, 1:100 protease inhibitor cocktail [Sigma P8340]), and twice with ice-cold PD Wash Buffer C (150 mM NaCl, 15 mM Tris-HCl pH 7.5, 0.5 mM EDTA-NaOH pH 8.0, 1:100 protease inhibitor cocktail [Sigma P8340]). Mixtures were transferred to new pre-chilled tubes on ice before each of the final two washes (resuspending the beads in PD Wash Buffer C before transferring). After complete removal of the final wash, beads were resuspended in 20 µL of PD Wash Buffer C containing 0.005 µg/µL RNase A (Thermo Fisher EN0531, a 1:20,000 dilution).

Tubes were incubated at 37 °C for 30 min, gently resuspending every 5 min to keep beads well suspended. Supernatants (first eluates) were transferred to pre-chilled tubes on ice. To collect residual proteins, beads were again resuspended in 15 µL of PD Wash Buffer C, placed on the magnetic separator, and supernatants were added to the first eluates and mixed for a total eluate volume of ~35 µL. A volume of 5 µL of each eluate was analyzed by SDS-PAGE on a 4-20% polyacrylamide gel (Bio-Rad 4561094) and silver staining with the Pierce Silver Stain Kit (Thermo Fisher 24612). The remaining eluates were snap-frozen in liquid nitrogen and stored at -80 °C prior to shipping on dry ice for proteomics analysis.

Repeat A proteomics sample analysis

Nuclear extract (input) samples: For in-solution Suspension Trap (S-Trap) digestion, 11.5 µL (~2 µg/µL) of each nuclear extract sample (inputs for RNA pulldowns) was mixed with in 11.5 µL of 10% SDS solution for 23 µL of initial-volume S-Trap protocol (ProtiFi K02-micro-10). After overnight trypsin digestion, peptides were reconstituted with 25 µL of 5% acetonitrile and 0.1% formic acid, and absorbances were measured. Due to higher absorbance, they were further diluted 2 times, and 0.6 pmole of yeast alcohol dehydrogenase (ADH) was spiked in for quantitative sample-to-sample comparison, retention time reproducibility, and mass accuracy throughout the runs. A 3.8-µL portion of each sample (independent replicate input samples, each analyzed three times) was injected into a to Waters nanoACQUITY UPLC coupled to a Thermo Scientific Orbitrap Fusion Lumos Tribrid mass spectrometer. The proteins were searched against the SwissProt Mouse database (V9) in Mascot Search Engine in Proteome Discoverer v2.1, followed by export to Scaffold v4 for false-discovery-rate (FDR) and intensity-based absolute quantitation (iBAQ) analysis.

RNA pulldown samples: In-solution S-Trap digestion was performed with 30-µL initial volume for each sample. After digesting with trypsin, samples were reconstituted with 20 µL 5% acetonitrile, 0.1% (v/v) formic acid in water following 0.2 pmole ADH spike-in. A 3.8 µL portion of each sample (independent replicate RNA pulldown samples, each analyzed once) was then injected to a Waters nanoAcquity UPLC coupled to a Thermo Scientific Orbitrap Fusion Lumos Tribrid mass spectrometer. After data acquisition, raw files were loaded to Proteome Discoverer software, and database SwissProt Mouse was used for the Mascot Search Engine followed by Scaffold analysis. The Scaffold analysis was carried out at 1% protein FDR.

Repeat A proteomics data analysis

Nuclear extract (input) samples: iBAQ data were exported from Scaffold to Microsoft Excel for processing. All iBAQ values of 0 or -1 were considered missing values and converted to values of 0. Proteins deemed as sample contaminants (yeast ADH, all keratins, and spurious entry "744") were removed, and proteins with multiple entries (Hnrnpa3 [2 entries], Hnrnpk [2 entries], Tmpo [2 entries], Hnrnpc [3 entries], Cpsf4 [2 entries]) were combined into one entry, summing their iBAQ values within each sample replicate. Within each sample replicate, all protein iBAQ values were normalized to total signal by dividing each protein's iBAQ value by the sum of all iBAQ values within the sample replicate, then multiplying by an arbitrary scaling factor of 1 million. To ensure robust estimation of relative protein abundance in the nuclear extracts, 149 proteins were removed for which 5 or more of the 9 normalized iBAQ values were missing. For the 2826 proteins remaining, nonmissing (nonzero) normalized iBAQ values were averaged, and the base-2 logarithm of this average was calculated to compare values to those from biotinylated RNA pulldowns on the same logarithmic scale (below).

RNA pulldown samples: iBAQ data were exported from Scaffold to Microsoft Excel and processed using Microsoft Excel and Python. All iBAQ values of 0 or -1 were considered missing values and converted to values of 0. Proteins deemed as sample contaminants were removed: yeast ADH, all keratins, and the 8 proteins with nonmissing iBAQ values in the "no RNA" control. Proteins with multiple entries (Hnrnpa3 [2 entries], Hnrnpc [2 entries]) were combined into one entry, summing their iBAQ values within each sample replicate. Within each sample replicate, all protein iBAQ values were normalized to total signal by dividing each protein's iBAQ value by the sum of all iBAQ values within the sample replicate, then multiplying by an arbitrary scaling factor of 1 million. (Due to intra-group variability of total-sample iBAQ values [e.g., within the WT samples], normalization to total-sample iBAQ signal was deemed more appropriate than normalization to the yeast ADH spike-in. One of four WT replicates and one of three ΔCCUGC replicates were removed from the analysis due to substantially poorer total protein detection.) To consider only proteins recovered reproducibly in the RNA pulldowns, the 844 proteins with iBAQ signal in at least one sample were filtered to a list of 644 proteins that fit at least one of two criteria: (1) at least 2 of the 3 WT values were nonmissing [614 proteins], or (2) all of the values for at least one mutant group were nonmissing [30 additional proteins]. Each nonmissing value was replaced with its base-2 logarithm to transform the data to a roughly normal distribution, and missing values were replaced according to the nature of their missingness: when only 1 value of a sample group was missing, it was assumed to be missing-at-random and was replaced with a not-a-number (NaN) to omit it from averages and statistical tests; when 2 or 3 values of a sample group were missing, these were assumed to be missing-not-at-random and underwent "Perseus-

style" imputation, replacing them with values generated at random from a normal distribution centered at (iBAQ_log2_mean) – 2*(iBAQ_log2_sd) with a standard deviation of 0.3*[(iBAQ_log2_mean) – 2*(iBAQ_log2_sd)], where iBAQ_log2_mean and iBAQ_log2_sd were the average and standard deviation, respectively, of all nonmissing values (total-normalized and log2-converted) in the dataset.

Two-sample t-tests were performed comparing the log2-transformed replicate values for WT and ΔGC-core pulldowns from three independent experiments, and the proteins recovered more strongly with WT and with $p < 0.05$ were considered the set of GC-rich core-interacting proteins highlighted in Figure 3B.

The heatmap in Figure 3D was generated by first filtering rows of proteins for those classified as SR proteins [55] and then using Morpheus (https://software.broadinstitute.org/morpheus), keeping column order fixed but enabling hierarchical clustering of rows with "metric" set to "one minus Pearson correlation" and "linkage method" set to "average."

Gene ontology analysis

Gene ontology of gene or protein sets were analyzed with the Molecular Signatures Database (MSigDB) at https://www.gsea-msigdb.org/gsea/msigdb [146,147], using mouse collections for all sets except for the set of *SPEN*-correlated genes from GTEx, which used the human collections. The top overlapping gene sets in GO biological process (GO:BP) and GO cellular component (GO:CC) were determined and reported with their enrichment p-values. Gene/protein classifications were defined by the following sources: SR proteins [55], nuclear speckles (MSigDB GO term GOCC_NUCLEAR_SPECK.v2023.2.Mm for mouse and MSigDB GO term GOCC_NUCLEAR_SPECK.v2024.1.Hs for human), paraspeckles [148], and nucleolus (MSigDB GO term GOCC_NUCLEOUS.v2023.2.Mm).

Halo-SPEN immunoprecipitation for proteomics analysis

Immunoprecipitation for proteomics analysis generally followed the RIP protocol above but with modifications made to accommodate larger inputs and a different method to reverse formaldehyde-induced crosslinks.

For each immunoprecipitation, well-mixed slurry volume of 50 μL Protein A/G PLUS-Agarose beads (Santa Cruz sc-2003) was washed three times with 1 mL room-temp 0.5% (w/v) bovine serum albumin (BSA; Fisher Scientific BP1600) in PBS. For each wash, 1.7-mL tubes were gently inverted three times and pelleted by centrifugation (2000 x g, 1 min, room temp), and supernatants were removed carefully, leaving ~10-20 μL to ensure no loss of beads. To washed beads, 300 μL

0.5% (w/v) BSA in PBS was added, followed by 15 µL 1 µg/µL anti-Halo antibody (Promega G9281). Tubes were incubated overnight at 4 °C with end-over-end rotation.

The following day, formaldehyde-crosslinked 10-million-cell pellets (7 pellets for Halo-SPEN-V5 WT, ΔIDR2 ΔIDR4, ΔSPOC, and no-rescue control IPs; 3 pellets for Halo-SPEN-V5 RRM1-4 and RRM2-4 IPs; see "RNA Immunoprecipitation" section) were thawed on ice and resuspended with 10 gentle up-down strokes of a P1000 pipet in 500 µL ice-cold RIPA Buffer (50 mM Tris-HCl pH 8.0, 150 mM KCl, 5 mM EDTA-NaOH pH 8.0, 1% [v/v] Triton X-100 [Fisher Scientific BP151], 0.5% [w/v] sodium deoxycholate, 0.1% [w/v] sodium dodecylsulfate) containing the following components added immediately before use: 0.5 µL per mL 1 M dithiothreitol (DTT), 10 µL per mL protease inhibitor cocktail (Sigma P8340), and 5 µL per mL 20 U/µL SUPERase-In RNase inhibitor (Invitrogen AM2696). Keeping tubes on ice, each cell suspension was sonicated twice for 30 s with a Sonics Vibra-Cell 130-watt VCX 130 probe-tip sonicator (30% amplitude output), with a 1-min rest on ice between sonication pulses. Tubes were centrifuged (15000 x g, 15 min, 4 °C), and 505 µL of each supernatant (cleared cell lysate) was transferred to a new pre-chilled 15-mL conical tube, pooling lysates for each genotype.

The antibody-bound beads from the day before were washed twice with 1 mL ice-cold fRIP Buffer; for each wash, 1.7-mL tubes were gently inverted three times and pelleted by centrifugation (2000 x g, 1 min, 4 °C), and supernatants were carefully removed, leaving ~10-20 µL to ensure no loss of beads. A 1-mL portion of complete fRIP Buffer (25 mM Tris-HCl pH 7.5, 150 mL KCl, 5 mM EDTA-NaOH pH 8.0, 0.5% [v/v] IGEPAL CA-630 [Fisher Scientific ICN19859650]; containing the following components added immediately before use: 0.5 µL per mL 1 M DTT, 10 µL per mL Sigma protease inhibitor cocktail, and 5 µL per mL 20 U/µL SUPERase-In) was used to completely collect washed antibody-bound beads and transfer to the 15-mL tubes containing lysates. A volume of fRIP Buffer was then added to each tube such that the total volume of complete fRIP Buffer was the same as the volume of pooled lysates in complete RIPA Buffer.

The next day, beads were pelleted by centrifugation (2000 x g, 1 min, 4 °C), and supernatants were carefully removed. The beads were washed once with 1 mL ice-cold fRIP Buffer, which was used to carefully transfer them to new 1.7-mL tubes. Beads were then washed three times with 1 mL ice-cold Pol II ChIP Buffer (50 mM Tris-HCl pH 7.5, 140 mM NaCl, 1 mM EDTA-NaOH pH 8.0, 1 mM EGTA-NaOH pH 8.0, 1% [v/v] Triton X-100, 0.1% [w/v] sodium deoxycholate, 0.1% [w/v] sodium dodecylsulfate), once with ice-cold 1 mL High-Salt Pol II ChIP Buffer (50 mM Tris-HCl pH 7.5, 500 mM NaCl, 1 mM EDTA-NaOH pH 8.0, 1 mM EGTA-NaOH pH 8.0, 1% [v/v] Triton X-100, 0.1% [w/v] sodium deoxycholate, 0.1% [w/v] sodium dodecylsulfate), and once with 1 mL

LiCl Buffer (20 mM Tris pH 8.0, 1 mM EDTA-NaOH pH 8.0, 250 mM LiCl, 0.5% [v/v] IGEPAL CA-630, 0.5% [w/v] sodium deoxycholate); each wash included a 5-min end-over-end rotation at 4 °C. In the final wash, samples were transferred to a new 1.7-mL tube. The final wash was carefully removed, first with a P1000 pipet and then with a P20 pipet, to ensure no loss of beads while leaving a volume of ~5 μL LiCl Buffer on the beads. A volume of 50 μL 2% (w/v) sodium dodecylsulfate was forcefully added to the beads to suspend them, and samples were incubated at 95 °C for 40 min to reverse crosslinks. A volume of 200 μL water was added, samples were vortexed at maximum power for 10 s, and centrifuged to pellet beads (2000 x g, 1 min, room temp). Supernatants were transferred to pre-chilled tubes on ice, snap-frozen in liquid nitrogen, and stored at -80 °C prior to shipping on dry ice for proteomics analysis.

Halo-SPEN IP proteomics sample analysis

Samples were flash frozen on dry ice and subsequently dried using a Speed-Vac. Next, 23 μL of 1x lysis buffer (5% SDS, 50 mM TEAB pH 8.5) was added to the protein pellet, initiating the Suspension trap (S-Trap) protocol (ProtiFi K02-micro-10). The proteins underwent reduction by 10 mM TCEP and alkylation by 20 mM iodoacetamide. Following an overnight digestion at 37 °C, 40 μL of 50 mM TEAB elution buffer was added and spun at 4000 rpm for 1 minute. This was followed by a 1-min spin with 40 μL of 0.2% formic acid (FA) and, lastly, 40 μL of 50% acetonitrile (ACN). The elution was then pooled together and dried using a Speed-Vac.

The digested samples were reconstituted with 20 μL of 0.1% formic acid in 5% ACN, followed by centrifugation at 16000 x g for 15 minutes. Subsequently, 18 μL of the sample was transferred to non-binding HPLC vials.

The samples (2 μL injection) were analyzed on a TimsTOF Pro2 (Bruker) mass spectrometer, which was coupled to a nanoElute LC system from Bruker. Peptides were loaded and separated on a house-made 75-μm I.D. fused silica analytical column packed with 25 cm ReproSil-Pur C18-AQ (Dr. Maisch, GmbH, 120 Å pore size, 3 μm particle size) particles to a gravity-pulled tip. A 30-min gradient was employed for DDA-PASEF with a flow rate set at 500 nL/min. The captive nano-electrospray voltage was maintained at 1600 V, using one column configuration (no trap) and a solvent composition of Solvent A: 0.1% FA in water and Solvent B: 0.1% FA in ACN. The analytical column was changed between the two batches of sample runs: the first batch consisted of the first biological replicates of no-rescue and Halo-SPEN-V5 WT IPs and only replicates of ΔIDR2 ΔIDR4 and ΔSPOC IPs, and the second batch contained all other IP samples (Table S2). Therefore, comparisons to no-rescue control were made within each batch.

DDA-PASEF method: The method consists of 10 MS/MS PASEF scans per topN acquisition cycle, with ramp and accumulation times of 100 ms, covering an m/z range from 100 to 1700 and an ion mobility range (1/K0) from 0.70 to 1.30 V s/cm^2. Collision energy settings followed a linear function of ion mobility, ranging from 20 eV at 0.6 V s/cm^2 to 59 eV at 1.6 V s/cm^2, using default parameters. Calibration of the instrument was performed using three ions from the ESI-L Tuning Mix (Agilent) [m/z 622, 922, 1222].

Halo-SPEN IP proteomics data analysis

The custom workflow 'LFQ-MBR' from FragPipe version 21.1 was used [149], including database search with MSFragger (version 4.0), deep-learning prediction rescoring with MSBooster, Percolator and ProteinProphet (Philosopher version 5.1.0) for PSM validation and protein inference. The raw .d files from Bruker were searched against *Mus musculus* reference proteome FATSA file UP000000589 from Uniprot (downloaded on 11/20/2023), appended with common contaminant proteins, containing in total 25483 sequence entries. Decoy reversed sequences were appended to the search database. The default MSFragger search parameters were used, except precursor and fragment mass tolerances were set to 20 ppm; enzyme: trypsin (not cutting before P); peptide length 7-30. Variable modifications were set to oxidation of Met and acetylation of protein N-t, and fixed modifications to carbamidomethylation of Cys. MSBooster, Percolator and ProteinProphet default options were used, and results were filtered by 1% FDR at protein level. The searched results were imported to Scaffold Q/S 5.2.2. Total spectrum counts were then exported to Microsoft Excel to calculate enrichments of proteins relative to no-rescue control (and between RRM1-4 and RRM2-4 samples) within each experimental replicate (Table S2). In Excel, all entries containing "keratin", "albumin", "Ig gamma", "Ig kappa", or "REVERSE" were considered contaminants and removed from the analysis. Data were visualized with GraphPad Prism (v9.5.0).

Immunoprecipitations for analysis by western blot

Halo-SPEN-V5 WT and near-full-length mutant constructs expressed at low levels, and repeated attempts to visualize them (or endogenous SPEN) via western blot of lysate samples (up to 60-100 µg per lane) were unsuccessful. Therefore, an immunoprecipitation step was used prior to western blot analysis in order to enrich constructs from larger amounts of lysate (1000 µg) and to reduce nonspecific signal on the membrane.

For each immunoprecipitation, 25 µL well-mixed slurry volume of Protein A/G PLUS-Agarose beads (Santa Cruz sc-2003) was washed three times with 1 mL room-temp 0.5% (w/v) bovine

serum albumin (BSA; Fisher Scientific BP1600) in PBS. For each wash, 1.7-mL tubes were gently inverted three times and pelleted by centrifugation (2000 x g, 1 min, room temp), and supernatants were removed carefully, leaving ~10-20 µL to ensure no loss of beads. To washed beads, 300 µL 0.5% (w/v) BSA in PBS was added, followed by 5 µg anti-V5 antibody (Thermo Fisher R960-25). Tubes were incubated overnight at 4 °C with end-over-end rotation.

After 3 d of treatment with 1000 ng/mL doxycycline hyclate, approximately 70-90% confluent 15-cm dishes of Halo-SPEN-V5-expressing RMCE-*Xist* Δ*Spen* cells were washed twice with 15 mL ice-cold PBS and harvested by scraping into 0.8 mL ice-cold PBS before transferring to a 1.7-mL tube. Cells were pelleted (1200 x g, 5 min, 4 °C) before removing supernatants and storing at -20 °C for later processing. Cell pellets were thawed on ice and resuspended with 10 up-down strokes of a P1000 pipet in 500 µL ice-cold RIPA Lysis Buffer (10 mM Tris-HCl pH 7.5, 140 mM NaCl, 1 mM EDTA-NaOH pH 8.0, 0.5 mM EGTA-NaOH pH 8.0, 1% [v/v] IGEPAL CA-630, 0.1% [w/v] sodium deoxycholate, 0.1% [w/v] SDS; 1 mM PMSF and 1:100 protease inhibitor cocktail [Sigma P8340-5ML] added fresh). Using a Sonics Vibracell 130-watt VCX 130 probe-tip sonicator (30% amplitude output), mixtures were sonicated on ice, twice for 10 s with a 1-min rest in between before pelleting cell debris (16,100 x g, 15 min, 4 °C). Cleared lysates (supernatants) were transferred to new, pre-chilled tubes on ice, and protein concentrations were measured against a standard curve of BSA (Fisher Scientific BP1600-100) via Bradford assay using 1X Bio-Rad Protein Assay Dye Reagent (Bio-Rad 5000006). Antibody-bound beads from the day before were washed twice with 1 mL ice-cold RIPA Lysis Buffer (not containing PMSF or protease inhibitor cocktail) before adding 1000 µg of lysate, brought to a total volume of 500 µL with the addition of RIPA Lysis Buffer. Tubes were incubated overnight at 4 °C with end-over-end rotation.

The next day, beads were washed three times with 1 mL ice-cold RIPA Lysis Buffer (not containing PMSF or protease inhibitor cocktail). The final wash was removed, leaving approximately 10-20 µL residual volume on top of the beads, to which 6 µL 4X NuPAGE LDS Sample Buffer (Thermo Fisher NP0007) containing 200 mM DTT was added and mixed by gentle flicking. Samples were heated at 95 °C for 5 min, allowed to cool to room temp, and pulsed down in a microcentrifuge to collect condensation prior to analyzing by SDS-PAGE and western blot.

Western blots

For western blots not requiring enrichment by immunoprecipitation, lysates were prepared and measured via Bradford assay as described above. Samples for SDS-PAGE were prepared by adding one volume 4X Laemmli sample buffer (Bio-Rad 1610747) containing 10% (v/v) β-mercaptoethanol to three volumes lysate. The exception to this was the right-most western blot

in Figure S5B, in which samples were instead prepared with 4X NuPAGE LDS Sample Buffer (Thermo Fisher NP0007) containing 200 mM DTT. All samples were heated at 95 °C for 5 min, allowed to cool to room temp, and pulsed down in a microcentrifuge to collect condensation prior to analyzing by SDS-PAGE and western blot.

For western blots to detect WT or near-full-length SPEN, protein samples in 1X NuPAGE LDS Sample Buffer (Thermo Fisher NP0007) containing 50 mM DTT were run on NuPAGE 3-8% Tris-Acetate pre-cast SDS-PAGE gels (Thermo Fisher EA0375BOX) alongside HiMark Pre-stained Protein Standard (Thermo Fisher LC5699) in 1X NuPAGE Tris-Acetate SDS Running Buffer (Thermo Fisher LA0041) at 150 V and at room temp. Proteins were wet-transferred to methanol-activated Immobilon-P PVDF membranes (MilliporeSigma IPVH00010) in cold 1X NuPAGE Transfer Buffer (Thermo Fisher NP0006) containing 20% methanol, for 1 h 45 min at 110 V and at 4 °C.

For all other western blots, protein samples in 1X Laemmli sample buffer (Bio-Rad 1610747) containing 2.5% (v/v) β-mercaptoethanol were run on TGX pre-cast SDS-PAGE gels (e.g., Bio-Rad 4561084) alongside Precision Plus all-blue protein ladder (Bio-Rad 1610373) in Tris-glycine-SDS running buffer (2.5 mM Tris, 19.2 mM glycine, 0.01% [w/v] SDS, pH 8.3) at 150 V and at room temp. Proteins were wet-transferred to methanol-activated Immobilon-P PVDF membranes (MilliporeSigma IPVH00010) in cold Tris-glycine-methanol-SDS transfer buffer (2.5 mM Tris, 19.2 mM glycine, 20% [v/v] methanol, 0.001% [w/v] SDS, pH 8.3) for 60 min at 100 V and at 4 °C.

After transfer, membranes were frequently cut horizontally with a razor blade to enable parallel detection of loading controls or co-immunoprecipitated proteins (Figure 4D). Membranes were blocked for at least 30 min on a room-temperature orbital shaker in 3% BSA (Fisher Scientific BP1600-100) or 5% nonfat milk (Bio-Rad 1706404) in tris-buffered saline (TBS; 20 mM Tris-HCl, 137 mM NaCl, pH 7.6) before adding primary antibody (see Table S6 for antibodies and concentrations) and incubating overnight on a 4 °C rocking platform. Membranes were washed with a quick rinse of TBS-T and then three times for 10 min in TBS-T on a room-temperature orbital shaker. An appropriate secondary antibody (see Table S6 for antibodies and concentrations) in 3% BSA or 5% milk in TBS was added and incubated for 30 min on a room-temperature orbital shaker. Membranes were washed as before with TBS-T. For detection of proteins probed with HRP-conjugated secondary antibodies, membranes were developed for 5 min with Clarity ECL substrate (Bio-Rad 1705061) or 3 min with SuperSignal West Femto Maximum Sensitivity ECL substrate (Thermo Fisher 34094) before placing in a clear plastic sheet protector to prevent drying while imaging. Membranes were imaged with a Bio-Rad ChemiDoc MP imager in chemiluminescence, DyLight 680, or DyLight 800 mode as appropriate. Protein

markers were visualized by simultaneously imaging a second channel in DyLight 680 or AlexaFluor 680 mode. Densitometry analysis (Figure S7B) was performed with Bio-Rad Image Lab 6.1 software.

SPEN "guilt-by-association" co-expression analysis

To enhance prediction accuracy, we took a conservative and multi-tissue/multi-tumor type approach, relying on experimental data from major public datasets and selecting high-quality results, as described below. We used three datasets: Genotype-Tissue Expression (GTEx), The Cancer Genome Atlas (TCGA) Tumor, and TCGA Normal. The first dataset includes normal tissues taken from the entire human body.[150] The second and third datasets include tumor samples or normal tissues from cancer patients (https://www.cancer.gov/tcga). For the sake of homogeneity and reproducibility, the shared source of data for these datasets was the Gene Expression Profiling Interactive Analysis (GEPIA) database.[151] This computational resource relies on recomputed data from RNA-seq samples, which were processed through the University of California, Santa Cruz (UCSC) Xena project pipeline.[152]

Co-expression was assessed using the Pearson correlation coefficient of non-log transformed data between *SPEN* and other genes across available samples. Notably, for GTEx, we included only tissues present both in the GEPIA list and in the GTEx V8 version and, at the same time, which are associated, in aggregate, in the GEPIA GTEx list, with > 10 samples. For TCGA Normal and Tumor, the inclusion criterion was to have > 10 samples associated. Due to these criteria, we used 49 different tissues for GTEx, 33 tumor types for TCGA Tumor, and 15 normal tissues for TCGA Normal (see Table S4). We selected, for each admitted tissue of GTEx and of TCGA Normal and for each cancer type of TCGA Tumor, the top 100 most correlated genes with *SPEN* (largest Pearson correlation coefficients, based on the GEPIA data). In case of isoform entries (distinct Ensembl ID) pointing to the same gene symbol, we removed these occasional duplicates and extended the analysis up to reaching the top 100 genes for each tissue, cell or cancer type.

We aggregated these gene entries in three tables having as many genes as the distinct correlated genes found across the tissues/cancers (i.e., union of all genes of the dataset having at least one top correlation) of that dataset and as many columns as the included tissues/cancers (one table for each dataset, tabs in Table S4). This procedure was separately followed for every dataset. While doing so, data were binarized such that the only numerical elements of each column of this table were 1 (gene belonging to the top 100 correlated genes for the analyzed tissue or tumor type) and 0 (otherwise). Then, we calculated the sum of gene hits across all tissues/cancers of that dataset (sums by rows) and used each total as the score for the

corresponding gene in that dataset. The motivation for this approach is that, once a correlation is ranked at the very top for a tissue/cancer, the second variable to account for is how frequently we observe it across the dataset. Our assumption was that the more a correlation is common in a dataset, the more likely it is that that correlation is (a) numerically reliable, (b) biologically relevant, and (c) cell- or tissue-independent, and therefore potentially involved in *SPEN* functions in multiple human cell types.

To assess the statistical significance of the score of each gene, separately, for each table, we simulated the outcome of having 100 hits for each column, but randomly distributed in it, and calculated how many times each score was reached, in a simulated row, in a sequence of 1 million (M) simulated matrixes. Therefore, the simulated matrixes had as many rows as those of the three datasets (1,780, 1,144, and 907, respectively) and as many columns as the tissues/cancers of the datasets (see above). The randomization algorithm was applied so that only valid rows were used in the simulation (rows having scores of 0 were not allowed). We used the simulation results, through the following two formulas, to calculate the statistical significance: (a) for all gene scores ≥ maximum score reached in the 1 M randomizations (MS1M), p-value = times MS1M is reached across 1 M randomizations / 1 M; (b) for all gene scores < MS1M, p-value = (times in which scores equaling that score are reached in the 1 M randomizations + times in which scores greater than that score are reached in the 1 M randomizations) / 1 M [153]. These *p*-values were reassessed in terms of multiple hypothesis testing by using the Benjamini Hochberg false discovery rate (FDR) method [154], as implemented in MATLAB [155].

The described method to assess the statistical significance of gene scores can be regarded as conservative, since in these simulations we only considered genes having at least one hit (i.e., at least one tissue type in which the gene was a top-100 *SPEN* correlate), instead of the full set of genes for which expression data were available.

In the TCGA Tumor dataset, of the 103 genes significantly correlated with *SPEN*, 59 reside on chr1. These were omitted from comparisons to the GTEx and TCGA Normal datasets and GO analysis because their correlations with *SPEN*, a gene on chr1, cannot be decoupled from cancer-associated chr1 copy-number variations that presumably have a strong influence in this dataset.

Overlaps of genes among the three datasets and depictions via area-proportional Venn diagram were generated using DeepVenn (https://www.deepvenn.com/, [156]).

To assess whether this method significantly enriches for genes that encode SPEN-interacting proteins, we first created a merged list of our 37 WT SPEN interactors (Table S2), our 146 SPEN RRM1-4 interactors (Table S2), and 207 SPOC domain interactors ("Ratio" value >2, [13]). We removed all duplicate entries, resulting in a list of 347 genes. We matched these 347 genes, by

gene symbol, with all genes that had collected at least one hit across the 49 GTEx tissue/cell types of our analysis. To prevent bias when matching gene symbols from distinct sources and of mixed nature (due to the presence of coding as well as non-coding genes in the GTEx dataset), this procedure was performed with respect to a pre-defined common set, including only coding genes recognized by the Human Genome Organization (HUGO, https://www.genenames.org/). We assessed if the matching genes were enriched or depleted for genes having the highest scores in the GTEx analysis (those with FDR < 0.10). For this task, we compared the proportion of HUGO genes with FDR < 0.10 in the matched list and in the full GTEx list (used as baseline). Those two proportions were compared through a two-tailed binomial test using R.

siRNA knockdown

Lyophilized siGLO Green Transfection Indicator (Dharmacon D-001630-01-05), ON-TARGETplus non-targeting siRNA #1 (Dharmacon D-001810-01-05), and factor-targeting ON-TARGETplus SMARTpools of four individual siRNAs (Dharmacon; see Key Resources Table) were resuspended to 20 μM in siRNA Resuspension Buffer (60 mM KCl, 6 mM Tris-HCl pH 7.5, 0.2 mM $MgCl_2$) before preparing small aliquots and storing at -80 °C.

RMCE-*Xist* WT (clone 4) cells expressing rtTA were transfected once with siRNAs at t = 0, transfected with siRNAs a second time at 48 h, induced by addition of 10-1000 ng/mL doxycycline hyclate starting at 72 h, and harvested at 96 h to prepare RNA samples for RNA-seq and protein samples for western blots. Transfections were performed with DharmaFECT 1 reagent (Dharmacon T-2001-02), following manufacturer's instructions with 50 nM working concentration of siRNA. First transfections were performed with 1 million trypsin-dissociated cells and second transfections were performed with 1.5 million cells, from which 0.5 million were plated for collecting RNA samples and 1 million were plated for collecting protein samples.

Xist O-MAP

For *Xist* O-MAP, RMCE-*Xist* WT and Δ*Spen* mESCs were cultured as above. *Xist* expression was induced with 1 μg/mL doxycycline in all of 4 replicates for 72 h prior to O-MAP fixation. Cells were plated into 6-well plates (3.5×10^5 cells per well; 6 wells per replicate) 24 hours prior to O-MAP. *Xist* O-MAP labeling was conducted in parallel on all replicates, as previously described [96,157], using *Xist*-targeting primary probes, and the SABER1–HRP O-MAP secondary probe described previously.[96] Briefly, cells were fixed (2% paraformaldehyde, 10 min), permeabilized (0.5% Triton X-10, 10 min), and endogenous peroxidases were inactivated (0.5% H_2O_2, 10 min). Primary *Xist* O-MAP probes [96] were hybridized for 8 hours at 42 °C, in 40% formamide

Hybridization Buffer.[96] Cells were washed to remove excess unbound probes and hybridized to the HRP-conjugated O-MAP oligo (1 hour at 37 °C; 30% formamide). Samples were extensively washed, and *in situ* biotinylated (8 µM biotin tyramide; 1 mM H_2O_2; 10 min at 23 °C). Labeling reactions were quenched with sodium azide and ascorbate, and cells were harvested by scraping into 1x Dulbecco's Phosphate Buffered Saline (dPBS), pooled, and stored at -80°C before use.

Xist O-MAP-MS sample preparation

Xist O-MAP labeled cells (4.2×10^6 cells per replicate; 4 biological replicates per experimental condition) were lysed, processed for streptavidin pulldown, and prepared for mass spectrometry using the same manipulations and reagents described previously.[157] Briefly, cells were gently lysed (0.5% NP-40, 10 min) and centrifuged to remove supernatant (3000 x g, 5min at 4 °C, 2 times) for nuclear enrichment. Nuclei were more extensively lysed (0.1% SDS, 10 min) and sonicated with a Branson Probe Sonicator (4 cycles of 30 s at 0.7/1.3 seconds on/off intervals) in an ice block. Samples were reverse-crosslinked by boiling (1% SDS, 1 hour, 800 rpm mixing), cooled to 50°C, then sonicated for one additional cycle on ice, using the same parameters. Samples were centrifuged (15,000 x g, 10 min at 4 °C), and clarified lysates were gently mixed with Pierce Magnetic streptavidin beads (ThermoFisher 88817; 75 µL per sample) for 2 h, with end-over-end rocking, at 23 °C. Beads were washed twice with (Tris-HCl, 0.1% SDS, sodium azide, NaCl) and samples were moved to PCR strip tubes to improve sample recovery. Samples were then washed with the following series of buffers, with one-minute, off-magnet resuspensions between each washing step: (Tris-HCl, 0.1% SDS, sodium azide; two washes), 1x PBS (two washes), 1 M KCl Buffer (two washes), 2 M Urea Buffer (four washes), EPPS Buffer (two washes). Beads were then resuspended in new PCR strip tubes in 50 µL EPPS buffer. Samples were reduced (10 mM TCEP, 60 min at 700 rpm mixing), alkylated (10 mM IAM, 60 min at 700 rpm in the dark) and quenched with DTT (5 mM, 15 min at 700 rpm). Samples were then digested with LysC (1 µg per 100 µg total protein; 3 hours at 37 °C, 700 rpm mixing) and then trypsin (1 µg per 100 µg total protein; overnight at 37 °C, 700 rpm mixing). Beads were removed magnetically and samples were moved to fresh 0.2 mL PCR-strip tubes. Peptides were stage-tipped and analyzed by mass-spectrometry as described.[157]

RNA stability assay

RMCE-*Xist* WT and mutant cells in wells of a 6-well plate were treated for 4 h with 5 μg/mL actinomycin D or an equivalent volume of DMSO vehicle control ("0 h") before washing once with 1X PBS and harvesting with 1 mL TRIzol. RNA was prepared via standard TRIzol-chloroform extraction, according to manufacturer's instructions, and resuspended in RNase-free water. 400 ng RNA was reverse-transcribed with random primers using the Applied Biosystems High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher 4368814; see above) and analyzed by qPCR (see above) to measure *Xist* and *Gapdh* mRNA abundance, fitting to a serially-diluted standard curve of a representative sample for each primer pair to incorporate primer efficiencies (see Table S6 for primer sequences). *Gapdh*-normalized *Xist* abundance data were calculated relative to 0-h control, and results from three independent experiments were with GraphPad Prism (v9.5.0).

Semiquantitative PCR analysis of *Xist* ectopic intron splicing

One microgram of total RNA from RMCE-*Xist* WT or intron mutant cells (see RNA-seq section) was reverse-transcribed with random primers using the Applied Biosystems High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher 4368814; see above), and products were diluted 10-fold with nuclease-free water. PCR reactions (10 μL) were prepared containing 5 μL iTaq Universal SYBR Green Supermix (Bio-Rad 1725124), 3 μL nuclease-free water, 0.5 μL 10 μM forward primer JT328, and 0.5 μL 10 μM reverse primer JT329 (see Table S6 for primer sequences). For PCR templates, 1 μL 1:10-diluted reverse transcription product was added, or in the case of the "spliced" and "unspliced" controls, 1 μL 0.5 ng/μL pTETRISv1-*Xist*-2kb or pTETRISv1-*Xist*-2kb-β-globin/IgG-intron,[26] respectively. PCR was run on a thermocycler with the following program: 95 °C for 10 min, 24 cycles of (95 °C for 15 s, 60 °C for 30 s, 72 °C for 30 s, 12 °C hold. Twenty-four cycles was chosen due to this being in the "linear range" of amplification curves from qPCR analyzed previously; parallel semiquantitative PCRs were run with 22 and 26 cycles, which yielded similar results. PCR products were mixed with Purple Gel Loading Dye (NEB B7024S) to 1X concentration and run on a 2% agarose-TAE gel containing ethidium bromide and visualized with a Bio-Rad ChemiDoc MP imager (Ethidium Bromide setting) alongside 1 Kb Plus DNA Ladder (Thermo Fisher MPK10025). Band intensities were determined by densitometry with Bio-Rad Image Lab 6.1 software. For each sample, splicing efficiency was determined as the ratio of spliced band intensity to the sum of spliced and unspliced band intensities.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
SPEN	Novus	NBP1-82952
IgG	Invitrogen	02-6102
FLAG	Sigma-Aldrich	F1804
CHERP	Bethyl	A304-621A
CPSF6	Bethyl	A301-356A
DDX46	Proteintech	16927-1-AP
LUC7L3	Bethyl	A301-498A
RBM39	Proteintech	21339-1-AP
SCAF4	Bethyl	A303-951A
SNRNP70	Santa Cruz	sc-390899
SNRNP200	Bethyl	A303-453A
SNRPA	Proteintech	10212-1-AP
SON	Abcam	ab121759
SRRM1	Proteintech	12822-1-AP
SRRM2	Invitrogen	PA5-66827
SRRT	Bethyl	A304-550A
SRSF1	Santa Cruz	sc-33652
SRSF3	MBL International	RN080PW
SRSF5	MBL International	RN082PW
SRSF6	Bethyl	A303-669A
SRSF9	MBL International	RN081PW
SRSF10	Santa Cruz	sc-101132
ZC3H13	Bethyl	A300-748A
ZC3H18	Bethyl	A304-682A
Halo (for IP and RIP)	Promega	G9281
Halo (for WB)	Promega	G9211
V5	Thermo Fisher	R960-25
U2AF2	Santa Cruz	sc-53942
ACIN1	Bethyl	A300-999A
HDAC3	GeneTex	GTX113303
TRIM28	Abcam	AB10483
RBM15	Proteintech	10587-1-AP
YTHDC1	Proteintech	29441-1-AP
Pol II S2P	Abcam	ab5095
Pol II S5P	Active Motif	61986
SUPT6H	Cell Signaling	15616
CPSF3	Bethyl	A301-091A
CPSF7	Bethyl	A301-360A
INTS1	Bethyl	A300-361A
INTS11	Sigma-Aldrich	HPA029025
EGFP	Proteintech	50420-2-AP
Vinculin	Santa Cruz	sc-73614
GAPDH	Proteintech	600004-1-Ig
POLR2D	Proteintech	16093-1-AP
Anti-mouse secondary antibody, HRP	Invitrogen	A16072
Anti-rabbit secondary antibody, HRP	Santa Cruz	sc-2357

Anti-mouse secondary antibody, DyLight 680	Invitrogen	35518
Anti-rabbit secondary antibody, DyLight 800	Invitrogen	SA5-10036
Bacterial and virus strains		
MAX Efficiency Stbl2 *E. coli* cells	Invitrogen	10268019
High-Efficiency 5-alpha competent *E. coli* cells	NEB	C2987H
Subcloning Efficiency DH5α competent *E. coli* cells	Invitrogen	18265017
Rosetta *E. coli* cells	MilliporeSigma	70953
Chemicals, peptides, and recombinant proteins		
DMEM high glucose plus sodium pyruvate	Gibco	11995-065
ESC-qualified fetal bovine serum	Gibco	26140-079
Non-essential amino acids	Gibco	11140-050
Penicillin-streptomycin	Gibco	15140-122
L-glutamine	Gibco	25030-081
β-mercaptoethanol	Sigma-Aldrich	63689
Trypsin-EDTA, phenol red	Gibco	2520072
10X Phosphate-Buffered Saline (PBS)	Corning	46-013-CM
Gelatin	Sigma	G9391
Hygromycin B	Roche	10843555001
Ganciclovir	InvivoGen	sud-gcv
G418 sulfate	Thermo Fisher	10131035
Puromycin dihydrochloride	Corning	61-385-RA
Q5 High-Fidelity DNA Polymerase	NEB	M0491
NEBuilder HiFi DNA Assembly Master Mix	NEB	E2621
KLD Enzyme Mix	NEB	M0554
Quick Ligase	NEB	M2200S
TE Buffer	Invitrogen	12090015
Lipofectamine 3000	Thermo Fisher	L3000001
Tris	Fisher Scientific	AC327360010
Hydrochloric acid (HCl)	Fisher Scientific	A142-212
Sodium chloride (NaCl)	Fisher Scientific	BP358-1
Ethylenediaminetetraacetic acid (EDTA)	Fisher Scientific	BP120-500
Sodium hydroxide (NaOH)	Fisher Scientific	BP359-500
Sodium dodecylsulfate (SDS)	Fisher Scientific	BP166-100
Proteinase K (for genotyping)	Meridian Bioscience	BIO-37084
Linear acrylamide	Thermo Fisher	AM9520
Ethanol	Decon Labs	2701
Agarose	Genesee Scientific	20-102QD
Acetic acid, glacial	J.T. Baker	9508-05
Ethidium bromide (1% solution)	Fisher Scientific	BP1302-10

Dimethyl sulfoxide (DMSO)	Sigma-Aldrich	D2650
Tocris dTAG^V-1 (1-mM stock prepared in DMSO)	Fisher Scientific	69-145
Doxycycline hyclate (1 µg/mL stock prepared in sterile water)	Sigma-Aldrich	D9891-1G
TRIzol	Thermo Fisher	15596026
Chloroform	Fisher Scientific	BP1145-1
ERCC RNA spike-in controls	Thermo Fisher	4456740
Dithiothreitol (DTT)	Thermo Fisher	15508013
Protease inhibitor cocktail	Sigma	P8340
SUPERase-In RNase inhibitor	Invitrogen	AM2696
Pierce 16% Formaldehyde (w/v), Methanol-free	Thermo Fisher	28906
Glycine	Fisher Scientific	BP381-1
Tween 20	Fisher Scientific	BP337
Protein A/G PLUS-Agarose beads	Santa Cruz Biotechnology	sc-2003
Bovine serum albumin (BSA)	Fisher Scientific	BP1600
Potassium chloride (KCl)	Fisher Scientific	BP366-1
Triton X-100	Fisher Scientific	BP151
Sodium deoxycholate	Sigma-Aldrich	D6750-100G
IGEPAL CA-630	Fisher Scientific	ICN19859650
Ethylene glycol-bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA)	MilliporeSigma	324626-25GM
Lithium chloride (LiCl)	Sigma-Aldrich	L4408-100G
N-lauroylsarcosine sodium salt	Fisher Scientific	AC434371000
Proteinase K (for RIP)	Thermo Fisher	25530049
RNaseOUT RNase Inhibitor	Thermo Fisher	10777019
iTaq Universal SYBR Green Supermix	Bio-Rad	1725124
Lysogeny broth (LB)	Research Products International	L24041
Ampicillin sodium salt	Fisher Scientific	BP1760-25
Chloramphenicol	Acros Organics	227920250
Isopropyl β-D-1-thiogalactopyranoside (IPTG)	Sigma-Aldrich	I6758
SigmaFast EDTA-free protease inhibitor cocktail	Sigma-Aldrich	S8830
Phenylmethylsulfonyl fluoride (PMSF)	Thermo Fisher	36978B
Pierce Glutathione Agarose Beads	Thermo Fisher	16100
Glycerol	Thermo Fisher	327255000
PreScission Protease	Cytiva	27-0843-01
GelCode Blue	Thermo Fisher	24590
Biotinylated UTP	MilliporeSigma	11388908910
Biotinylated UTP	TriLink Biotechnologies	N-5005-1
SequaGel UreaGel 29:1 Concentrate	National Diagnostics	EC-828
SequaGel UreaGel Diluent	National Diagnostics	EC-840
SequaGel UreaGel Buffer	National Diagnostics	EC-835

5x TBE buffer	National Diagnostics	EC-861
Diamond Nucleic Acid Dye	Promega	H1181
Magnesium chloride (MgCl$_2$)	Thermo Fisher	223210010
Dynabeads MyOne Streptavidin T1 beads	Thermo Fisher	65602
Yeast tRNA	Thermo Fisher	AM7119
D-biotin (50 mM stock prepared in DMSO)	Thermo Fisher	B1595
N-(2-hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid) [HEPES]	Sigma-Aldrich	H4034-500G
Protein Assay Dye Reagent Concentrate (Bradford)	Bio-Rad	5000006
Dynabeads M-280 Streptavidin beads	Thermo Fisher	11205D
RNase A	Thermo Fisher	EN0531
Mini-PROTEAN 4-20% TGX SDS-PAGE gels	Bio-Rad	4561094
Mini-PROTEAN 4-15% TGX SDS-PAGE gels	Bio-Rad	4561084
4X NuPAGE LDS Sample Buffer	Thermo Fisher	NP0007
4X Laemmli sample buffer	Bio-Rad	1610747
NuPAGE 3-8% Tris-Acetate SDS-PAGE gels	Thermo Fisher	EA0375BOX
HiMark Pre-stained Protein Standard	Thermo Fisher	LC5699
NuPAGE Tris-Acetate SDS Running Buffer	Thermo Fisher	LA0041
Immobilon-P PVDF membrane	MilliporeSigma	IPVH00010
NuPAGE Transfer Buffer	Thermo Fisher	NP0006
Methanol	Sigma-Aldrich	646377-4L
Precision Plus all-blue protein ladder	Bio-Rad	1610373
Blotting-grade nonfat milk blocker	Bio-Rad	1706404
Clarity ECL substrate	Bio-Rad	1705061
SuperSignal West Femto Maximum Sensitivity ECL substrate	Thermo Fisher	34094
siGLO Green Transfection Indicator	Dharmacon (Horizon Discovery)	D-001630-01-05
ON-TARGETplus Non-targeting siRNA #1	Dharmacon (Horizon Discovery)	D-001810-01-05
ON-TARGETplus Mouse *Acin1* siRNA - SMARTPool	Dharmacon (Horizon Discovery)	L-065307-00-0005
ON-TARGETplus Mouse *Hdac3* siRNA - SMARTPool	Dharmacon (Horizon Discovery)	L-043553-02-0005
ON-TARGETplus Mouse *Trim28* siRNA - SMARTPool	Dharmacon (Horizon Discovery)	L-040800-01-0005
ON-TARGETplus Mouse *Zc3h13* siRNA - SMARTPool	Dharmacon (Horizon Discovery)	L-050497-01-0005
ON-TARGETplus Mouse *Ythdc1* siRNA - SMARTPool	Dharmacon (Horizon Discovery)	L-167076-00-0005
ON-TARGETplus Mouse *Zc3h18* siRNA - SMARTPool	Dharmacon (Horizon Discovery)	L-066066-01-0005

ON-TARGETplus Mouse *Scaf4* siRNA - SMARTPool	Dharmacon (Horizon Discovery)	L-167104-00-0005
DharmaFECT 1 Transfection Reagent	Dharmacon (Horizon Discovery)	T-2001-02
Actinomycin D, working stock dissolved in DMSO	Sigma	A9415-5MG
1 Kb Plus DNA Ladder	Thermo Fisher	MPK10025
Critical commercial assays		
Q5 High-Fidelity DNA Polymerase	NEB	M0491
NEBuilder HiFi DNA Assembly Master Mix	NEB	E2621
KLD Enzyme Mix	NEB	M0554
Quick Ligase	NEB	M2200S
NucleoBond BAC100 Maxiprep Kit	Macherey-Nagel	740579
Qubit dsDNA High-Sensitivity Kit	Thermo Fisher	Q32854
Qubit dsDNA Broad-Range Kit	Thermo Fisher	Q32853
PureLink HiPure Plasmid Midiprep Kit	Invitrogen	K210004
Neon Transfection System 100 µL Kit	Invitrogen	MPK10025
Apex Taq Polymerase Kit	Genesee Scientific	42-801B2
RNA High-Sensitivity Kit	Thermo Fisher	Q32852
RNA Broad-Range kit	Thermo Fisher	Q10211
KAPA RNA HyperPrep Kit with RiboErase (HMR)	Roche	08098140702
KAPA Unique Dual-Indexed Adapters	Roche	08861919702
RNA Clean & Concentrator-5 Kit	Zymo	R1013
Applied Biosystems High-Capacity cDNA Reverse Transcription Kit	Thermo Fisher	4368814
GenCatch Gel Extraction Kit	Epoch Life Science	2260250
MEGAscript SP6 Transcription Kit	Thermo Fisher	AM1330
MEGAclear Transcription Clean-Up Kit	Thermo Fisher	AM1908
Pierce Silver Stain Kit	Thermo Fisher	24612
S-Trap Proteomics Sample Prep Kit	ProtiFi	K02-micro-10
Deposited data		
GENCODE Release M25 (GRCm38.p6) *Mus musculus* reference genome	https://www.gencodegenes.org/ mouse/release_M25.html	gencode.vM25.basic.annotation.gtf
Sanger VCF SNP file	https://www.mousegenomes.org/	k_mgp.v5.vcf.parsed
Uniprot *Mus musculus* reference proteome	https://www.uniprot.org/proteomes/UP000000589	UP000000589
Gene Expression Profiling Interactive Analysis (GEPIA) database	Tang et al.[151]	
All RNA- and RIP-seq from this study	NCBI GEO	GEO: In progress
Gel and western blot images from this study	Mendeley	In progress

Original code from this study	Zenodo/GitHub	In progress
Proteomics data from this study	ProteomeXchange	In progress
Experimental models: Cell lines		
DR4 irMEF feeder cells	ATCC	SCRC-1045
B6/CAST F1-hybrid *Rosa26*-RMCE mouse ESCs	Trotman et al.[26]	R26-RMCE-i69-P19-E114
B6/CAST F1-hybrid *Rosa26*-RMCE-empty mouse ESCs	Trotman et al.[41]	empty_clone_1
B6/CAST F1-hybrid *Rosa26*-RMCE-empty mouse ESCs	Trotman et al.[41]	empty_clone_2
B6/CAST F1-hybrid *Rosa26*-RMCE-empty mouse ESCs	Trotman et al.[41]	empty_clone_3
B6/CAST F1-hybrid *Rosa26*-RMCE-empty mouse ESCs	Trotman et al.[41]	empty_clone_4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* WT mouse ESCs	Trotman et al.[41]	WT_clone_1
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* WT mouse ESCs	Trotman et al.[41]	WT_clone_3
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* WT mouse ESCs	Trotman et al.[41]	WT_clone_4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* WT mouse ESCs	Trotman et al.[41]	WT_clone_5
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA mouse ESCs	This study	ΔRepA_clone_4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA mouse ESCs	This study	ΔRepA_clone_5
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB mouse ESCs	This study	ΔRepA-XB_clone_3
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB mouse ESCs	This study	ΔRepA-XB_clone_6
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔU-spacer mouse ESCs	This study	ΔU-spacer_clone_1
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔU-spacer mouse ESCs	This study	ΔU-spacer_clone_3
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔGC-core mouse ESCs	This study	ΔGC-core _clone_4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔGC-core mouse ESCs	This study	ΔGC-core _clone_8
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* preserve mouse ESCs	This study	preserve_clone_2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* preserve mouse ESCs	This study	preserve_clone_4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* disrupt mouse ESCs	This study	disrupt_clone_1
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* disrupt mouse ESCs	This study	disrupt_clone_2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* AWCG mouse ESCs	This study	AWCG_clone_1

B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* AWCG mouse ESCs	This study	AWCG_clone_2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔGG16,17 mouse ESCs	This study	ΔGG16,17_clone_2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔGG16,17 mouse ESCs	This study	ΔGG16,17_clone_3
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* AWCG+ΔGG mouse ESCs	This study	AWCG+ΔGG_clone_5
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* AWCG+ΔGG mouse ESCs	This study	AWCG+ΔGG_clone_8
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* U19C mouse ESCs	This study	U19C_clone_1
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* U19C mouse ESCs	This study	U19C_clone_3
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* U23C mouse ESCs	This study	U23C_clone_1
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* U23C mouse ESCs	This study	U23C_clone_3
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔCCUGC mouse ESCs	This study	ΔCCUGC_clone_1
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔCCUGC mouse ESCs	This study	ΔCCUGC_clone_2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δss234 mouse ESCs	This study	Δss234_clone_2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δss234 mouse ESCs	This study	Δss234_clone_4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* intron mouse ESCs	This study	intron_clone_3
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* intron mouse ESCs	This study	intron_clone_4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) mouse ESCs	This study	ΔRepA-XB-6x(U-spacer-3xMS2)_clone_1
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) mouse ESCs	This study	ΔRepA-XB-6x(U-spacer-3xMS2)_clone_4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* WT SRSF2-3xFLAG mouse ESCs	This study	WT_SRSF2-3xFLAG
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔCCUGC SRSF2-3xFLAG mouse ESCs	This study	ΔCCUGC_SRSF2-3xFLAG
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* non-targeting gRNA control mouse ESCs	This study	NTG_clone_6
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* mouse ESCs	This study	ΔSpen_clone_27
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* mouse ESCs	This study	ΔSpen_clone_32

B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* EGFP-Halo WT mouse ESCs	This study	EGFP-Halo
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 WT mouse ESCs	This study	Halo-SPEN-V5_WT
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 ΔRRM1-4 mouse ESCs	This study	Halo-SPEN-V5_ ΔRRM1-4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 ΔRRM2-4 mouse ESCs	This study	Halo-SPEN-V5_ΔRRM2-4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 ΔRRM1-IDR2 mouse ESCs	This study	Halo-SPEN-V5_ΔRRM1-IDR2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 ΔIDR2 mouse ESCs	This study	Halo-SPEN-V5_ΔIDR2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 ΔRS mouse ESCs	This study	Halo-SPEN-V5_ΔRS
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 ΔIDR4 mouse ESCs	This study	Halo-SPEN-V5_ΔIDR4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 ΔIDR2 ΔIDR4 mouse ESCs	This study	Halo-SPEN-V5_ΔIDR2_ΔIDR4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 ΔSPOC mouse ESCs	This study	Halo-SPEN-V5_ΔSPOC
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 RRM1-4 mouse ESCs	This study	Halo-SPEN-V5_RRM1-4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 RRM2-4 mouse ESCs	This study	Halo-SPEN-V5_RRM2-4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 IDR2-RRM4 mouse ESCs	This study	Halo-SPEN-V5_IDR2-RRM4
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* Δ*Spen* Halo-SPEN-V5 RRM1-4 ΔIDR2 mouse ESCs	This study	Halo-SPEN-V5_RRM1-4_ΔIDR2
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist FKBP1A^{F36V}-Srsf1* mouse ESCs	This study	FKBP1A-Srsf1_clone_N1-9
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist FKBP1A^{F36V}-Srsf1* mouse ESCs	This study	FKBP1A-Srsf1_clone_N1-13
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) EGFP mouse ESCs	This study	EGFP

B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) MCP-SRSF1 FL mouse ESCs	This study	SRSF1_FL
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) MCP-EGFP FL mouse ESCs	This study	MCP-EGFP
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) MCP-3xHNRNPK RGG mouse ESCs	This study	3xHNRNPK_RGG
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) MCP-4xSRSF1 RS mouse ESCs	This study	4xSRSF1_RS
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) MCP-1xSRSF2 RS mouse ESCs	This study	1xSRSF2_RS
B6/CAST F1-hybrid *Rosa26*-RMCE-*Xist* ΔRepA-XB-6x(U-spacer-3xMS2) MCP-SRSF1-2-1 RS mouse ESCs	This study	SRSF1-2-1_RS
SM33 mouse ESCs	Engreitz et al.[51]	pSM33
SM33 Δ*Spen* mouse ESCs	This study	SM33_ΔSpen_clone_A1
SM33 Δ*Spen* mouse ESCs	This study	SM33_ΔSpen_clone_A3
E14 mouse ESCs (ES-E14TG2a)	ATCC	CRL-1821
Oligonucleotides		
Table S6		
Recombinant DNA		
pCARGO-RMCE-empty	Trotman et al.;[41] Addgene	233264
pCARGO-RMCE-Xist WT	Trotman et al.;[41] Addgene	233268
pCARGO-RMCE-Xist ΔRepA	This study; Addgene	
pCARGO-RMCE-Xist ΔRepA-XB	This study; Addgene	
pCARGO-RMCE-Xist ΔU-spacer	This study; Addgene	
pCARGO-RMCE-Xist ΔGC-core	This study; Addgene	
pCARGO-RMCE-Xist preserve	This study; Addgene	
pCARGO-RMCE-Xist disrupt	This study; Addgene	
pCARGO-RMCE-Xist AWCG	This study; Addgene	
pCARGO-RMCE-Xist ΔGG16,17	This study; Addgene	
pCARGO-RMCE-Xist AWCG+ΔGG	This study; Addgene	
pCARGO-RMCE-Xist U19C	This study; Addgene	
pCARGO-RMCE-Xist U23C	This study; Addgene	
pCARGO-RMCE-Xist ΔCCUGC	This study; Addgene	
pCARGO-RMCE-Xist Δss234	This study; Addgene	
pCARGO-RMCE-Xist intron	This study; Addgene	
pCARGO-RMCE-Xist ΔRepA-XB-6x(U-spacer-3xMS2)	This study; Addgene	

pDD001-SPEN-RRM2-4	This study; Addgene	
pDD001-SRSF1-RRM1-2	This study; Addgene	
pCL116(PuroR)	This study; Addgene	
pCL116(PuroR)-SRSF2-3xFLAG	This study; Addgene	
pCL116-EGFP-Halo	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 WT	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 ΔRRM1-4	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 ΔRRM2-4	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 ΔRRM1-IDR2	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 ΔIDR2	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 ΔRS	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 ΔIDR4	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 ΔIDR2 ΔIDR4	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 ΔSPOC	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 RRM1-4	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 RRM2-4	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 IDR2-RRM4	This study; Addgene	
pCL116(PuroR)-Halo-SPEN-V5 RRM1-4 ΔIDR2	This study; Addgene	
PX459_v2-Srsf1-N1-gRNA	This study; Addgene	
pGW-Amp-Srsf1-N_term-dTAG-HDR	This study; Addgene	
pCL116(PuroR)-MCP-SRSF1_v2	This study; Addgene	
pCL116(PuroR)-MCP-EGFP-NLS-3xFLAG	This study; Addgene	
pCL116(PuroR)-MCP-3xHNRNPK_RGG-NLS-3xFLAG	This study; Addgene	
pCL116(PuroR)-MCP-4xSRSF1_RS-NLS-3xFLAG	This study; Addgene	
pCL116(PuroR)-MCP-1xSRSF2_RS-NLS-3xFLAG	This study; Addgene	
pCL116(PuroR)-MCP-SRSF1-2-1_RS-NLS-3xFLAG	This study; Addgene	
PB-TRE-Cas9	Schertzer et al.;[133] Addgene	126029
pOG-Cre	Trotman et al.;[41] Addgene	233265
PB-rtTA	Schertzer et al.;[133] Addgene	126034
pUC19-piggyBac transposase	Schertzer et al.[133]	N/A

PB-rtTA(BsmBI)-non-targeting-gRNA	Braceros et al.;[135] Addgene	
pX330-Spen-sbdel-gRNA1	Trotman et al.;[26] Addgene	
pX330-Spen-sbdel-gRNA2	Trotman et al.;[26] Addgene	
pX330-Spen-sbdel-gRNA3	Trotman et al.;[26] Addgene	
pX330-Spen-sbdel-gRNA4	Trotman et al.;[26] Addgene	
Software and algorithms		
CRISPR Guide RNA Design Tool	Benchling	https://www.benchling.com/crispr
Codon Optimization Tool	IDT	https://www.idtdna.com/pages/tools/codon-optimization-tool
STAR	Dobin et al.[136]	v2.7.9a, v2.7.10a
featureCounts (Subread)	Liao et al. [138]	v2.0.4, v2.0.6
DESeq2	Love et al.[47]	v1.32.0
Samtools	Li et al. [141]	v1.15, v1.20
IGV	Robinson et al. [142]	v2.8.9
Seaborn	Waskom [140]	
Prism	GraphPad	v9.5.0
UCSC Genome Browser	UCSC	https://genome.ucsc.edu/
MACS2	Zhang et al. [143]	v2.2.7.1
BEDTools	Quinlan and Hall [144]	v2.29
CFX Manager	Bio-Rad	v3.1
STREME (MEME Suite)	Bailey [52]	v5.2.0
Image Lab	Bio-Rad	v6.1
Proteome Discoverer	Thermo Fisher	v2.1
Scaffold	Proteome Software	v4.0, v5.2.2
Molecular Signatures Database	Castanza et al. and Subramanian et al.[146,147]	https://www.gsea-msigdb.org/gsea/msigdb
LFQ-MBR workflow (FragPipe)	Kong et al. [149]	v21.1
MSFragger	Kong et al. [149]	v4.0
Philosopher	da Veiga Leprevost et al. [158]	v5.1.0
MobiDB	Potenza et al. [59]	v2.0
DeepVenn	Hulsen [158]	https://www.deepvenn.com/
RNA2Drawer	Johnson et al. [127]	https://rna2drawer.app/
Other		
Forma Series II Water-Jacketed CO_2 Incubator	Thermo Fisher	3110
NanoDrop Lite spectrophotometer	Thermo Fisher	840281400
Qubit 2.0 Fluorometer	Thermo Fisher	Q32866

Neon Transfection System	Thermo Fisher	10431915
Parafilm M	VWR	52858-032
Multidizer Hybridization Oven	UVP	95-0340-01
T100 Thermal Cycler	Bio-Rad	1861096
ChemiDoc MP Imaging System	Bio-Rad	12003154
NextSeq500 sequencing platform	Illumina	SY-415-1001
NextSeq1000 sequencing platform	Illumina	20038898
75-cycle NextSeq 500/550 High Output Kit v2.5	Illumina	20024906
100-cycle NextSeq 1000/2000 P2 Reagent Kit v3	Illumina	20046811
100-cycle NextSeq 1000/2000 P2 XLEAP-SBS Reagent Kit	Illumina	20100987
Vibra-Cell 130-W probe-tip sonicator	Sonics	VCX 130
Hard-Shell 96-Well PCR Plates	Bio-Rad	HSP9601
Microseal 'B' PCR Plate Sealing Film	Bio-Rad	MSB1001
C1000 Touch Thermal Cycler	Bio-Rad	1851196
CFX96 Real-Time System	Bio-Rad	1845097
BioPhotometer Plus spectrophotometer	Eppendorf	6132
Dounce homogenizer with tight pestle	DWK Life Sciences	357542
Slide-A-Lyzer 10-kDa MWCO devices	Thermo Fisher	88404
6-well cell culture plate	Genesee Scientific	25-105
Stratalinker 1800 UV crosslinker	Stratagene	400071
nanoACQUITY UPLC system	Waters	71500097503
Orbitrap Fusion Lumos Tribrid mass spectrometer	Thermo Scientific	FETD2-10001
nanoElute LC system	Bruker	
TimsTOF Pro2 mass spectrometer	Bruker	
ReproSil-Pur C18-AQ	Dr. Maisch	r13.aq.

REFERENCES

1.   Loda, A., Collombet, S., and Heard, E. (2022). Gene regulation in time and space during X-chromosome inactivation. Nat Rev Mol Cell Biol *23*, 231-249. 10.1038/s41580-021-00438-7.

2.   Pintacuda, G., Wei, G., Roustan, C., Kirmizitas, B.A., Solcan, N., Cerase, A., Castello, A., Mohammed, S., Moindrot, B., Nesterova, T.B., and Brockdorff, N. (2017). hnRNPK Recruits PCGF3/5-PRC1 to the Xist RNA B-Repeat to Establish Polycomb-Mediated Chromosomal Silencing. Mol Cell *68*, 955-969 e910. 10.1016/j.molcel.2017.11.013.

3.   Pandya-Jones, A., Markaki, Y., Serizay, J., Chitiashvili, T., Mancia Leon, W.R., Damianov, A., Chronis, C., Papp, B., Chen, C.K., McKee, R., et al. (2020). A protein assembly mediates Xist localization and gene silencing. Nature *587*, 145-151. 10.1038/s41586-020-2703-0.

4.   Lu, Z., Guo, J.K., Wei, Y., Dou, D.R., Zarnegar, B., Ma, Q., Li, R., Zhao, Y., Liu, F., Choudhry, H., et al. (2020). Structural modularity of the XIST ribonucleoprotein complex. Nature communications *11*, 6163. 10.1038/s41467-020-20040-3.

5.   Graindorge, A., Pinheiro, I., Nawrocka, A., Mallory, A.C., Tsvetkov, P., Gil, N., Carolis, C., Buchholz, F., Ulitsky, I., Heard, E., et al. (2019). In-cell identification and measurement of RNA-protein interactions. Nat Commun *10*, 5317. 10.1038/s41467-019-13235-w.

6.   Weidmann, C.A., Mustoe, A.M., Jariwala, P.B., Calabrese, J.M., and Weeks, K.M. (2021). Analysis of RNA-protein networks with RNP-MaP defines functional hubs on RNA. Nat Biotechnol *39*, 347-356. 10.1038/s41587-020-0709-7.

7.   Minks, J., Baldry, S.E., Yang, C., Cotton, A.M., and Brown, C.J. (2013). XIST-induced silencing of flanking genes is achieved by additive action of repeat a monomers in human somatic cells. Epigenetics Chromatin *6*, 23. 1756-8935-6-23 [pii]

10.1186/1756-8935-6-23.

8.   Smola, M.J., Christy, T.W., Inoue, K., Nicholson, C.O., Friedersdorf, M., Keene, J.D., Lee, D.M., Calabrese, J.M., and Weeks, K.M. (2016). SHAPE reveals transcript-wide interactions, complex structural domains, and protein interactions across the Xist lncRNA in living cells. Proc Natl Acad Sci U S A *113*, 10322-10327. 10.1073/pnas.1600008113.

9.   Chu, C., Zhang, Q.C., da Rocha, S.T., Flynn, R.A., Bharadwaj, M., Calabrese, J.M., Magnuson, T., Heard, E., and Chang, H.Y. (2015). Systematic discovery of Xist RNA binding proteins. Cell *161*, 404-416. 10.1016/j.cell.2015.03.025.

10.   McHugh, C.A., Chen, C.K., Chow, A., Surka, C.F., Tran, C., McDonel, P., Pandya-Jones, A., Blanco, M., Burghard, C., Moradian, A., et al. (2015). The Xist lncRNA interacts directly with SHARP to silence transcription through HDAC3. Nature *521*, 232-236. 10.1038/nature14443.

11.   Monfort, A., Di Minin, G., Postlmayr, A., Freimann, R., Arieti, F., Thore, S., and Wutz, A. (2015). Identification of Spen as a Crucial Factor for Xist Function through Forward Genetic Screening in Haploid Embryonic Stem Cells. Cell Rep *12*, 554-561. 10.1016/j.celrep.2015.06.067.

12.   Moindrot, B., Cerase, A., Coker, H., Masui, O., Grijzenhout, A., Pintacuda, G., Schermelleh, L., Nesterova, T.B., and Brockdorff, N. (2015). A Pooled shRNA Screen Identifies Rbm15, Spen, and Wtap as Factors Required for Xist RNA-Mediated Silencing. Cell Rep *12*, 562-572. 10.1016/j.celrep.2015.06.053.

13.   Dossin, F., Pinheiro, I., Zylicz, J.J., Roensch, J., Collombet, S., Le Saux, A., Chelmicki, T., Attia, M., Kapoor, V., Zhan, Y., et al. (2020). SPEN integrates transcriptional and epigenetic control of X-inactivation. Nature *578*, 455-460. 10.1038/s41586-020-1974-9.

14.   Zylicz, J.J., Bousard, A., Zumer, K., Dossin, F., Mohammad, E., da Rocha, S.T., Schwalb, B., Syx, L., Dingli, F., Loew, D., et al. (2019). The Implication of Early Chromatin Changes in X Chromosome Inactivation. Cell *176*, 182-197 e123. 10.1016/j.cell.2018.11.041.

15.   Dixon-McDougall, T., and Brown, C.J. (2022). Multiple distinct domains of human XIST are required to coordinate gene silencing and subsequent heterochromatin formation. Epigenetics Chromatin *15*, 6. 10.1186/s13072-022-00438-7.

16.   Kaufmann, C., and Wutz, A. (2023). IndiSPENsable for X Chromosome Inactivation and Gene Silencing. Epigenomes *7*. 10.3390/epigenomes7040028.

17.   Lu, Z., Zhang, Q.C., Lee, B., Flynn, R.A., Smith, M.A., Robinson, J.T., Davidovich, C., Gooding, A.R., Goodrich, K.J., Mattick, J.S., et al. (2016). RNA Duplex Map in Living Cells Reveals Higher-Order Transcriptome Structure. Cell *165*, 1267-1279. 10.1016/j.cell.2016.04.028.

18.   Carter, A.C., Xu, J., Nakamoto, M.Y., Wei, Y., Zarnegar, B.J., Shi, Q., Broughton, J.P., Ransom, R.C., Salhotra, A., Nagaraja, S.D., et al. (2020). Spen links RNA-mediated endogenous retrovirus silencing and X chromosome inactivation. Elife *9*. 10.7554/eLife.54508.

19.   Button, A.C., Hall, S.D., Ashley, E.L., and McHugh, C.A. (2024). Dissection of protein and RNA regions required for SPEN binding to XIST A-repeat RNA. RNA *30*, 240-255. 10.1261/rna.079713.123.

20.   Wutz, A., Rasmussen, T.P., and Jaenisch, R. (2002). Chromosomal silencing and localization are mediated by different domains of Xist RNA. Nat Genet *30*, 167-174. 10.1038/ng820

ng820 [pii].

21.   Duszczyk, M.M., Wutz, A., Rybin, V., and Sattler, M. (2011). The Xist RNA A-repeat comprises a novel AUCG tetraloop fold and a platform for multimerization. RNA *17*, 1973-1982. 10.1261/rna.2747411.

22.   Wang, Y., Zhong, Y., Zhou, Y., Tanaseichuk, O., Li, Z., and Zhao, J.C. (2019). Identification of a Xist silencing domain by Tiling CRISPR. Sci Rep *9*, 2408. 10.1038/s41598-018-36750-0.

23.   Rodermund, L., Coker, H., Oldenkamp, R., Wei, G., Bowness, J., Rajkumar, B., Nesterova, T., Susano Pinto, D.M., Schermelleh, L., and Brockdorff, N. (2021). Time-resolved structured illumination microscopy reveals key principles of Xist RNA spreading. Science *372*. 10.1126/science.abe7500.

24.   Hoki, Y., Kimura, N., Kanbayashi, M., Amakawa, Y., Ohhata, T., Sasaki, H., and Sado, T. (2009). A proximal conserved repeat in the Xist gene is essential as a genomic element for X-inactivation in mouse. Development *136*, 139-146. dev.026427 [pii]

10.1242/dev.026427.

25.   Nesterova, T.B., Wei, G., Coker, H., Pintacuda, G., Bowness, J.S., Zhang, T., Almeida, M., Bloechl, B., Moindrot, B., Carter, E.J., et al. (2019). Systematic allelic analysis defines the interplay of key pathways in X chromosome inactivation. Nature communications *10*, 3129. 10.1038/s41467-019-11171-3.

26.   Trotman, J.B., Lee, D.M., Cherney, R.E., Kim, S.O., Inoue, K., Schertzer, M.D., Bischoff, S.R., Cowley, D.O., and Calabrese, J.M. (2020). Elements at the 5' end of Xist harbor SPEN-independent transcriptional antiterminator activity. Nucleic Acids Res *48*, 10500-10517. 10.1093/nar/gkaa789.

27.   Royce-Tolland, M.E., Andersen, A.A., Koyfman, H.R., Talbot, D.J., Wutz, A., Tonks, I.D., Kay, G.F., and Panning, B. (2010). The A-repeat links ASF/SF2-dependent Xist RNA processing with random choice during X inactivation. Nat Struct Mol Biol *17*, 948-954. nsmb.1877 [pii]

10.1038/nsmb.1877.

28.   Colognori, D., Sunwoo, H., Kriz, A.J., Wang, C.Y., and Lee, J.T. (2019). Xist Deletional Analysis Reveals an Interdependency between Xist RNA and Polycomb Complexes for Spreading along the Inactive X. Mol Cell *74*, 101-117 e110. 10.1016/j.molcel.2019.01.015.

29.   Sakata, Y., Nagao, K., Hoki, Y., Sasaki, H., Obuse, C., and Sado, T. (2017). Defects in dosage compensation impact global gene regulation in the mouse trophoblast. Development *144*, 2784-2797. 10.1242/dev.149138.

30.   Cartegni, L., Wang, J., Zhu, Z., Zhang, M.Q., and Krainer, A.R. (2003). ESEfinder: A web resource to identify exonic splicing enhancers. Nucleic Acids Res *31*, 3568-3571. 10.1093/nar/gkg616.

31.   Radio, F.C., Pang, K., Ciolfi, A., Levy, M.A., Hernandez-Garcia, A., Pedace, L., Pantaleoni, F., Liu, Z., de Boer, E., Jackson, A., et al. (2021). SPEN haploinsufficiency causes a neurodevelopmental disorder overlapping proximal 1p36 deletion syndrome with an episignature of X chromosomes in females. Am J Hum Genet *108*, 502-516. 10.1016/j.ajhg.2021.01.015.

32.   Damrauer, J.S., Beckabir, W., Klomp, J., Zhou, M., Plimack, E.R., Galsky, M.D., Grivas, P., Hahn, N.M., O'Donnell, P.H., Iyer, G., et al. (2022). Collaborative study from the Bladder Cancer Advocacy Network for the genomic analysis of metastatic urothelial cancer. Nature communications *13*, 6658. 10.1038/s41467-022-33980-9.

33.   Heath, A.P., Ferretti, V., Agrawal, S., An, M., Angelakos, J.C., Arya, R., Bajari, R., Baqar, B., Barnowski, J.H.B., Burt, J., et al. (2021). The NCI Genomic Data Commons. Nat Genet *53*, 257-262. 10.1038/s41588-021-00791-5.

34.   Mendiratta, G., Ke, E., Aziz, M., Liarakos, D., Tong, M., and Stites, E.C. (2021). Cancer gene mutation frequencies for the U.S. population. Nature communications *12*, 5961. 10.1038/s41467-021-26213-y.

35.   Reddy, A., Zhang, J., Davis, N.S., Moffitt, A.B., Love, C.L., Waldrop, A., Leppa, S., Pasanen, A., Meriranta, L., Karjalainen-Lindsberg, M.L., et al. (2017). Genetic and Functional Drivers of Diffuse Large B Cell Lymphoma. Cell *171*, 481-494 e415. 10.1016/j.cell.2017.09.027.

36.   Sarkozy, C., Hung, S.S., Chavez, E.A., Duns, G., Takata, K., Chong, L.C., Aoki, T., Jiang, A., Miyata-Takata, T., Telenius, A., et al. (2021). Mutational landscape of gray zone lymphoma. Blood *137*, 1765-1776. 10.1182/blood.2020007507.

37.   Schmitz, R., Wright, G.W., Huang, D.W., Johnson, C.A., Phelan, J.D., Wang, J.Q., Roulland, S., Kasbekar, M., Young, R.M., Shaffer, A.L., et al. (2018). Genetics and Pathogenesis of Diffuse Large B-Cell Lymphoma. N Engl J Med *378*, 1396-1407. 10.1056/NEJMoa1801445.

38.   Seiler, M., Peng, S., Agrawal, A.A., Palacino, J., Teng, T., Zhu, P., Smith, P.G., Cancer Genome Atlas Research, N., Buonamici, S., and Yu, L. (2018). Somatic Mutational Landscape of Splicing Factor Genes and Their Functional Consequences across 33 Cancer Types. Cell Rep *23*, 282-296 e284. 10.1016/j.celrep.2018.01.088.

39.   Tyner, J.W., Tognon, C.E., Bottomly, D., Wilmot, B., Kurtz, S.E., Savage, S.L., Long, N., Schultz, A.R., Traer, E., Abel, M., et al. (2018). Functional genomic landscape of acute myeloid leukaemia. Nature *562*, 526-531. 10.1038/s41586-018-0623-z.

40.   Legare, S., Cavallone, L., Mamo, A., Chabot, C., Sirois, I., Magliocco, A., Klimowicz, A., Tonin, P.N., Buchanan, M., Keilty, D., et al. (2015). The Estrogen Receptor Cofactor SPEN Functions as a Tumor Suppressor and Candidate Biomarker of Drug Responsiveness in Hormone-Dependent Breast Cancers. Cancer Res *75*, 4351-4363. 10.1158/0008-5472.CAN-14-3475.

41.   Trotman, J.B., Abrash, E.W., Murvin, M.M., Braceros, A.K., Li, S., Boyson, S.P., Salcido, R.T., Cherney, R.E., Bischoff, S.R., Kaufmann, K., et al. (2025). Isogenic comparison of Airn and Xist reveals core principles of Polycomb recruitment by lncRNAs. Mol Cell *85*, 1117-1133 e1114. 10.1016/j.molcel.2025.02.014.

42.   Kirk, J.M., Kim, S.O., Inoue, K., Smola, M.J., Lee, D.M., Schertzer, M.D., Wooten, J.S., Baker, A.R., Sprague, D., Collins, D.W., et al. (2018). Functional classification of long non-coding RNAs by k-mer content. Nat Genet *50*, 1474-1482. 10.1038/s41588-018-0207-8.

43.   Hughes, A.L., Szczurek, A.T., Kelley, J.R., Lastuvkova, A., Turberfield, A.H., Dimitrova, E., Blackledge, N.P., and Klose, R.J. (2023). A CpG island-encoded mechanism protects genes from premature transcription termination. Nature communications *14*, 726. 10.1038/s41467-023-36236-2.

44.   Koerner, M.V., Pauler, F.M., Hudson, Q.J., Santoro, F., Sawicka, A., Guenzl, P.M., Stricker, S.H., Schichl, Y.M., Latos, P.A., Klement, R.M., et al. (2012). A downstream CpG island controls transcript initiation and elongation and the methylation state of the imprinted Airn macro ncRNA promoter. PLoS Genet *8*, e1002540. 10.1371/journal.pgen.1002540.

45.   Younis, I., Berg, M., Kaida, D., Dittmar, K., Wang, C., and Dreyfuss, G. (2010). Rapid-response splicing reporter screens identify differential regulators of constitutive and alternative splicing. Mol Cell Biol *30*, 1718-1728. 10.1128/MCB.01301-09.

46.   Matsuura, R., Nakajima, T., Ichihara, S., and Sado, T. (2021). Ectopic Splicing Disturbs the Function of Xist RNA to Establish the Stable Heterochromatin State. Front Cell Dev Biol *9*, 751154. 10.3389/fcell.2021.751154.

47.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-0550-8.

48.   Colognori, D., Sunwoo, H., Wang, D., Wang, C.Y., and Lee, J.T. (2020). Xist Repeats A and B Account for Two Distinct Phases of X Inactivation Establishment. Dev Cell *54*, 21-32 e25. 10.1016/j.devcel.2020.05.021.

49.   Schertzer, M.D., Braceros, K.C.A., Starmer, J., Cherney, R.E., Lee, D.M., Salazar, G., Justice, M., Bischoff, S.R., Cowley, D.O., Ariel, P., et al. (2019). lncRNA-Induced Spread of Polycomb Controlled by Genome Architecture, RNA Abundance, and CpG Island DNA. Mol Cell *75*, 523-537 e510. 10.1016/j.molcel.2019.05.028.

50.   Jachowicz, J.W., Strehle, M., Banerjee, A.K., Blanco, M.R., Thai, J., and Guttman, M. (2022). Xist spatially amplifies SHARP/SPEN recruitment to balance chromosome-wide silencing and specificity to the X chromosome. Nat Struct Mol Biol *29*, 239-249. 10.1038/s41594-022-00739-1.

51.   Engreitz, J.M., Pandya-Jones, A., McDonel, P., Shishkin, A., Sirokman, K., Surka, C., Kadri, S., Xing, J., Goren, A., Lander, E.S., et al. (2013). The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. Science *341*, 1237973. science.1237973 [pii]

10.1126/science.1237973.

52.   Bailey, T.L. (2021). STREME: Accurate and versatile sequence motif discovery. Bioinformatics. 10.1093/bioinformatics/btab203.

53.   Hinkle, E.R., Wiedner, H.J., Torres, E.V., Jackson, M., Black, A.J., Blue, R.E., Harris, S.E., Guzman, B.B., Gentile, G.M., Lee, E.Y., et al. (2022). Alternative splicing regulation of membrane trafficking genes during myogenesis. RNA *28*, 523-540. 10.1261/rna.078993.121.

54.   Aguilar, R., Spencer, K.B., Kesner, B., Rizvi, N.F., Badmalia, M.D., Mrozowich, T., Mortison, J.D., Rivera, C., Smith, G.F., Burchard, J., et al. (2022). Targeting Xist with compounds that disrupt RNA structure and X inactivation. Nature *604*, 160-166. 10.1038/s41586-022-04537-z.

55.   Cascarina, S.M., and Ross, E.D. (2022). Expansion and functional analysis of the SR-related protein family across the domains of life. RNA *28*, 1298-1314. 10.1261/rna.079170.122.

56.   Sliskovic, I., Eich, H., and Muller-McNicoll, M. (2022). Exploring the multifunctionality of SR proteins. Biochemical Society transactions *50*, 187-198. 10.1042/BST20210325.

57.   Malovannaya, A., Lanz, R.B., Jung, S.Y., Bulynko, Y., Le, N.T., Chan, D.W., Ding, C., Shi, Y., Yucer, N., Krenciute, G., et al. (2011). Analysis of the human endogenous coregulator complexome. Cell *145*, 787-799. 10.1016/j.cell.2011.05.006.

58.   Hein, M.Y., Hubner, N.C., Poser, I., Cox, J., Nagaraj, N., Toyoda, Y., Gak, I.A., Weisswange, I., Mansfeld, J., Buchholz, F., et al. (2015). A human interactome in three quantitative dimensions organized by stoichiometries and abundances. Cell *163*, 712-723. 10.1016/j.cell.2015.09.053.

59.   Potenza, E., Di Domenico, T., Walsh, I., and Tosatto, S.C. (2015). MobiDB 2.0: an improved database of intrinsically disordered and mobile proteins. Nucleic Acids Res *43*, D315-320. 10.1093/nar/gku982.

60.   Manley, J.L., and Krainer, A.R. (2010). A rational nomenclature for serine/arginine-rich protein splicing factors (SR proteins). Genes Dev *24*, 1073-1074. 10.1101/gad.1934910.

61.   Cazalla, D., Zhu, J., Manche, L., Huber, E., Krainer, A.R., and Caceres, J.F. (2002). Nuclear export and retention signals in the RS domain of SR proteins. Mol Cell Biol *22*, 6871-6882. 10.1128/MCB.22.19.6871-6882.2002.

62.   Markaki, Y., Gan Chong, J., Wang, Y., Jacobson, E.C., Luong, C., Tan, S.Y.X., Jachowicz, J.W., Strehle, M., Maestrini, D., Banerjee, A.K., et al. (2021). Xist nucleates local protein gradients to propagate silencing across the X chromosome. Cell *184*, 6174-6192 e6132. 10.1016/j.cell.2021.10.022.

63.   Van Nostrand, E.L., Pratt, G.A., Shishkin, A.A., Gelboin-Burkhart, C., Fang, M.Y., Sundararaman, B., Blue, S.M., Nguyen, T.B., Surka, C., Elkins, K., et al. (2016). Robust transcriptome-wide discovery of RNA-binding protein binding sites with enhanced CLIP (eCLIP). Nat Methods *13*, 508-514. 10.1038/nmeth.3810.

64.   Ji, X., Zhou, Y., Pandit, S., Huang, J., Li, H., Lin, C.Y., Xiao, R., Burge, C.B., and Fu, X.D. (2013). SR proteins collaborate with 7SK and promoter-associated nascent RNA to release paused polymerase. Cell *153*, 855-868. 10.1016/j.cell.2013.04.028.

65.   Chaturvedi, P., and Belmont, A.S. (2024). Nuclear speckle biology: At the cross-roads of discovery and functional analysis. Curr Opin Cell Biol *91*, 102438. 10.1016/j.ceb.2024.102438.

66.   Prasanth, K.V., Camiolo, M., Chan, G., Tripathi, V., Denis, L., Nakamura, T., Hubner, M.R., and Spector, D.L. (2010). Nuclear organization and dynamics of 7SK RNA in regulating gene expression. Mol Biol Cell *21*, 4184-4196. 10.1091/mbc.E10-02-0105.

67.   Consortium, G. (2013). The Genotype-Tissue Expression (GTEx) project. Nat Genet *45*, 580-585. 10.1038/ng.2653.

68.   van Dam, S., Craig, T., and de Magalhaes, J.P. (2015). GeneFriends: a human RNA-seq-based gene and transcript co-expression database. Nucleic Acids Res *43*, D1124-1132. 10.1093/nar/gku1042.

69.   Stuart, J.M., Segal, E., Koller, D., and Kim, S.K. (2003). A gene-coexpression network for global discovery of conserved genetic modules. Science *302*, 249-255. 10.1126/science.1087447.

70.   Quackenbush, J. (2003). Genomics. Microarrays--guilt by association. Science *302*, 240-241. 10.1126/science.1090887.

71.   Yuan, Z., VanderWielen, B.D., Giaimo, B.D., Pan, L., Collins, C.E., Turkiewicz, A., Hein, K., Oswald, F., Borggrefe, T., and Kovall, R.A. (2019). Structural and Functional Studies of the RBPJ-SHARP Complex Reveal a Conserved Corepressor Binding Site. Cell Rep *26*, 845-854 e846. 10.1016/j.celrep.2018.12.097.

72.   Oswald, F., Rodriguez, P., Giaimo, B.D., Antonello, Z.A., Mira, L., Mittler, G., Thiel, V.N., Collins, K.J., Tabaja, N., Cizelsky, W., et al. (2016). A phospho-dependent mechanism involving NCoR and KMT2D controls a permissive chromatin state at Notch target genes. Nucleic Acids Res *44*, 4703-4720. 10.1093/nar/gkw105.

73.   Gregersen, L.H., Mitter, R., Ugalde, A.P., Nojima, T., Proudfoot, N.J., Agami, R., Stewart, A., and Svejstrup, J.Q. (2019). SCAF4 and SCAF8, mRNA Anti-Terminator Proteins. Cell *177*, 1797-1813 e1718. 10.1016/j.cell.2019.04.038.

74.   Altendorfer, E., Mochalova, Y., and Mayer, A. (2022). BRD4: a general regulator of transcription elongation. Transcription *13*, 70-81. 10.1080/21541264.2022.2108302.

75.   Yu, B., Qi, Y., Li, R., Shi, Q., Satpathy, A.T., and Chang, H.Y. (2021). B cell-specific XIST complex enforces X-inactivation and restrains atypical B cells. Cell *184*, 1790-1803 e1717. 10.1016/j.cell.2021.02.015.

76.   Andersen, P.R., Domanski, M., Kristiansen, M.S., Storvall, H., Ntini, E., Verheggen, C., Schein, A., Bunkenborg, J., Poser, I., Hallais, M., et al. (2013). The human cap-binding complex is functionally connected to the nuclear RNA exosome. Nat Struct Mol Biol *20*, 1367-1376. 10.1038/nsmb.2703.

77.   Polak, P., Garland, W., Rathore, O., Schmid, M., Salerno-Kochan, A., Jakobsen, L., Gockert, M., Gerlach, P., Silla, T., Andersen, J.S., et al. (2023). Dual agonistic and antagonistic roles of ZC3H18 provide for co-activation of distinct nuclear RNA decay pathways. Cell Rep *42*, 113325. 10.1016/j.celrep.2023.113325.

78.   Liu, J., Dou, X., Chen, C., Chen, C., Liu, C., Xu, M.M., Zhao, S., Shen, B., Gao, Y., Han, D., and He, C. (2020). N (6)-methyladenosine of chromosome-associated regulatory

RNA regulates chromatin state and transcription. Science *367*, 580-586. 10.1126/science.aay6018.

79.   Patil, D.P., Chen, C.K., Pickering, B.F., Chow, A., Jackson, C., Guttman, M., and Jaffrey, S.R. (2016). m(6)A RNA methylation promotes XIST-mediated transcriptional repression. Nature *537*, 369-+. 10.1038/nature19342.

80.   Xiao, W., Adhikari, S., Dahal, U., Chen, Y.S., Hao, Y.J., Sun, B.F., Sun, H.Y., Li, A., Ping, X.L., Lai, W.Y., et al. (2016). Nuclear m(6)A Reader YTHDC1 Regulates mRNA Splicing. Mol Cell *61*, 507-519. 10.1016/j.molcel.2016.01.012.

81.   Nayler, O., Hartmann, A.M., and Stamm, S. (2000). The ER repeat protein YT521-B localizes to a novel subnuclear compartment. J Cell Biol *150*, 949-962. 10.1083/jcb.150.5.949.

82.   Bousard, A., Raposo, A.C., Zylicz, J.J., Picard, C., Pires, V.B., Qi, Y., Gil, C., Syx, L., Chang, H.Y., Heard, E., and da Rocha, S.T. (2019). The role of Xist-mediated Polycomb recruitment in the initiation of X-chromosome inactivation. EMBO Rep *20*, e48019. 10.15252/embr.201948019.

83.   Pandit, S., Zhou, Y., Shiue, L., Coutinho-Mansfield, G., Li, H., Qiu, J., Huang, J., Yeo, G.W., Ares, M., Jr., and Fu, X.D. (2013). Genome-wide analysis reveals SR protein cooperation and competition in regulated splicing. Mol Cell *50*, 223-235. 10.1016/j.molcel.2013.03.001.

84.   Feng, H., Bao, S., Rahman, M.A., Weyn-Vanhentenryck, S.M., Khan, A., Wong, J., Shah, A., Flynn, E.D., Krainer, A.R., and Zhang, C. (2019). Modeling RNA-Binding Protein Specificity In Vivo by Precisely Registering Protein-RNA Crosslink Sites. Mol Cell *74*, 1189-1204 e1186. 10.1016/j.molcel.2019.02.002.

85.   Daubner, G.M., Clery, A., Jayne, S., Stevenin, J., and Allain, F.H. (2012). A syn-anti conformational difference allows SRSF2 to recognize guanines and cytosines equally well. EMBO J *31*, 162-174. 10.1038/emboj.2011.367.

86.   Wheeler, E.C., Vora, S., Mayer, D., Kotini, A.G., Olszewska, M., Park, S.S., Guccione, E., Teruya-Feldstein, J., Silverman, L., Sunahara, R.K., et al. (2022). Integrative RNA-omics Discovers GNAS Alternative Splicing as a Phenotypic Driver of Splicing Factor-Mutant Neoplasms. Cancer Discov *12*, 836-855. 10.1158/2159-8290.CD-21-0508.

87.   Zhang, X., Smits, A.H., van Tilburg, G.B., Jansen, P.W., Makowski, M.M., Ovaa, H., and Vermeulen, M. (2017). An Interaction Landscape of Ubiquitin Signaling. Mol Cell *65*, 941-955 e948. 10.1016/j.molcel.2017.01.004.

88.   Nabet, B., Ferguson, F.M., Seong, B.K.A., Kuljanin, M., Leggett, A.L., Mohardt, M.L., Robichaud, A., Conway, A.S., Buckley, D.L., Mancias, J.D., et al. (2020). Rapid and direct control of target protein levels with VHL-recruiting dTAG molecules. Nat Commun *11*, 4687. 10.1038/s41467-020-18377-w.

89.   Nabet, B., Roberts, J.M., Buckley, D.L., Paulk, J., Dastjerdi, S., Yang, A., Leggett, A.L., Erb, M.A., Lawlor, M.A., Souza, A., et al. (2018). The dTAG system for immediate and

target-specific protein degradation. Nat Chem Biol *14*, 431-441. 10.1038/s41589-018-0021-8.

90.   Abuhashem, A., and Hadjantonakis, A.K. (2022). Generation of knock-in degron tags for endogenous proteins in mice using the dTAG system. STAR Protoc *3*, 101660. 10.1016/j.xpro.2022.101660.

91.   Chen, C., Liu, W., Guo, J., Liu, Y., Liu, X., Liu, J., Dou, X., Le, R., Huang, Y., Li, C., et al. (2021). Nuclear m(6)A reader YTHDC1 regulates the scaffold function of LINE1 RNA in mouse ESCs and early embryos. Protein Cell *12*, 455-474. 10.1007/s13238-021-00837-8.

92.   Xu, W., Li, J., He, C., Wen, J., Ma, H., Rong, B., Diao, J., Wang, L., Wang, J., Wu, F., et al. (2021). METTL3 regulates heterochromatin in mouse embryonic stem cells. Nature *591*, 317-321. 10.1038/s41586-021-03210-1.

93.   Guo, Y.E., Manteiga, J.C., Henninger, J.E., Sabari, B.R., Dall'Agnese, A., Hannett, N.M., Spille, J.H., Afeyan, L.K., Zamudio, A.V., Shrinivas, K., et al. (2019). Pol II phosphorylation regulates a switch between transcriptional and splicing condensates. Nature *572*, 543-548. 10.1038/s41586-019-1464-0.

94.   Das, R., Yu, J., Zhang, Z., Gygi, M.P., Krainer, A.R., Gygi, S.P., and Reed, R. (2007). SR proteins function in coupling RNAP II transcription to pre-mRNA splicing. Mol Cell *26*, 867-881. 10.1016/j.molcel.2007.05.036.

95.   Horiuchi, K., Kawamura, T., Iwanari, H., Ohashi, R., Naito, M., Kodama, T., and Hamakubo, T. (2013). Identification of Wilms' tumor 1-associating protein complex and its role in alternative splicing and the cell cycle. J Biol Chem *288*, 33292-33302. 10.1074/jbc.M113.500397.

96.   Tsue, A.F., Kania, E.E., Lei, D.Q., Fields, R., McGann, C.D., Marciniak, D.M., Hershberg, E.A., Deng, X., Kihiu, M., Ong, S.E., et al. (2024). Multiomic characterization of RNA microenvironments by oligonucleotide-mediated proximity-interactome mapping. Nat Methods *21*, 2058-2071. 10.1038/s41592-024-02457-6.

97.   Huttlin, E.L., Ting, L., Bruckner, R.J., Gebreab, F., Gygi, M.P., Szpyt, J., Tam, S., Zarraga, G., Colby, G., Baltier, K., et al. (2015). The BioPlex Network: A Systematic Exploration of the Human Interactome. Cell *162*, 425-440. 10.1016/j.cell.2015.06.043.

98.   Oswald, F., Kostezka, U., Astrahantseff, K., Bourteele, S., Dillinger, K., Zechner, U., Ludwig, L., Wilda, M., Hameister, H., Knochel, W., et al. (2002). SHARP is a novel component of the Notch/RBP-Jkappa signalling pathway. EMBO J *21*, 5417-5426.

99.   Trotman, J.B., Braceros, K.C.A., Cherney, R.E., Murvin, M.M., and Calabrese, J.M. (2021). The control of polycomb repressive complexes by long noncoding RNAs. Wiley Interdiscip Rev RNA *12*, e1657. 10.1002/wrna.1657.

100.   Chen, S.L., Rooney, T.J., Hu, A.R., Beard, H.S., Garrett, W.M., Mangalath, L.M., Powers, J.J., Cooper, B., and Zhang, X.N. (2019). Quantitative Proteomics Reveals a Role for SERINE/ARGININE-Rich 45 in Regulating RNA Metabolism and Modulating

Transcriptional Suppression via the ASAP Complex in Arabidopsis thaliana. Front Plant Sci *10*, 1116. 10.3389/fpls.2019.01116.

101.   Wei, G., H., C., Rodermund, L., Almeida, M., Roach, H.L., Nesterova, T., and Brockdorff, N. (2025). N6-methyladenosine and the NEXT complex direct Xist RNA turnover and X inactivation dynamics. bioRxiv. https://doi.org/10.1101/2025.01.09.632176.

102.   Cho, N.H., Cheveralls, K.C., Brunner, A.D., Kim, K., Michaelis, A.C., Raghavan, P., Kobayashi, H., Savy, L., Li, J.Y., Canaj, H., et al. (2022). OpenCell: Endogenous tagging for the cartography of human cellular organization. Science *375*, eabi6983. 10.1126/science.abi6983.

103.   Appel, L.M., Franke, V., Benedum, J., Grishkovskaya, I., Strobl, X., Polyansky, A., Ammann, G., Platzer, S., Neudolt, A., Wunder, A., et al. (2023). The SPOC domain is a phosphoserine binding module that bridges transcription machinery with co- and post-transcriptional regulators. Nature communications *14*, 166. 10.1038/s41467-023-35853-1.

104.   Zhang, L., Zhang, Y., Chen, Y., Gholamalamdari, O., Wang, Y., Ma, J., and Belmont, A.S. (2020). TSA-seq reveals a largely conserved genome organization relative to nuclear speckles with small position changes tightly correlated with gene expression changes. Genome Res *31*, 251-264. 10.1101/gr.266239.120.

105.   Bhat, P., Chow, A., Emert, B., Ettlin, O., Quinodoz, S.A., Strehle, M., Takei, Y., Burr, A., Goronzy, I.N., Chen, A.W., et al. (2024). Genome organization around nuclear speckles drives mRNA splicing efficiency. Nature *629*, 1165-1173. 10.1038/s41586-024-07429-6.

106.   Shopland, L.S., Johnson, C.V., Byron, M., McNeil, J., and Lawrence, J.B. (2003). Clustering of multiple specific genes and gene-rich R-bands around SC-35 domains: evidence for local euchromatic neighborhoods. J Cell Biol *162*, 981-990. 10.1083/jcb.200303131.

107.   Yoon, Y., Bournique, E., Soles, L.V., Yin, H., Chu, H.F., Yin, C., Zhuang, Y., Liu, X., Liu, L., Jeong, J., et al. (2025). RBBP6 anchors pre-mRNA 3' end processing to nuclear speckles for efficient gene expression. Mol Cell *85*, 555-570 e558. 10.1016/j.molcel.2024.12.016.

108.   Collombet, S., Rall, I., Dugast-Darzacq, C., Heckert, A., Halavatyi, A., Le Saux, A., Dailey, G., Darzacq, X., and Heard, E. (2023). RNA polymerase II depletion from the inactive X chromosome territory is not mediated by physical compartmentalization. Nat Struct Mol Biol *30*, 1216-1223. 10.1038/s41594-023-01008-5.

109.   Rouviere, J.O., Salerno-Kochan, A., Lykke-Andersen, S., Garland, W., Dou, Y., Rathore, O., Molska, E.S., Wu, G., Schmid, M., Bugai, A., et al. (2023). ARS2 instructs early transcription termination-coupled RNA decay by recruiting ZC3H4 to nascent transcripts. Mol Cell *83*, 2240-2257 e2246. 10.1016/j.molcel.2023.05.028.

110.   Parker, M.T., Knop, K., Zacharaki, V., Sherwood, A.V., Tome, D., Yu, X., Martin, P.G., Beynon, J., Michaels, S.D., Barton, G.J., and Simpson, G.G. (2021). Widespread premature transcription termination of Arabidopsis thaliana NLR genes by the spen protein FPA. eLife *10*. 10.7554/eLife.65537.

111. Hornyik, C., Terzi, L.C., and Simpson, G.G. (2010). The spen family protein FPA controls alternative cleavage and polyadenylation of RNA. Dev Cell *18*, 203-213. 10.1016/j.devcel.2009.12.009.

112. Baurle, I., Smith, L., Baulcombe, D.C., and Dean, C. (2007). Widespread role for the flowering-time regulators FCA and FPA in RNA-mediated chromatin silencing. Science *318*, 109-112. 10.1126/science.1146565.

113. Busch, A., and Hertel, K.J. (2012). Evolution of SR protein and hnRNP splicing regulatory factors. Wiley Interdiscip Rev RNA *3*, 1-12. 10.1002/wrna.100.

114. Duret, L., Chureau, C., Samain, S., Weissenbach, J., and Avner, P. (2006). The Xist RNA gene evolved in eutherians by pseudogenization of a protein-coding gene. Science *312*, 1653-1655. 10.1126/science.1126316.

115. Elisaphenko, E.A., Kolesnikov, N.N., Shevchenko, A.I., Rogozin, I.B., Nesterova, T.B., Brockdorff, N., and Zakian, S.M. (2008). A dual origin of the Xist gene from a protein-coding gene and a set of transposable elements. PLoS One *3*, e2521. 10.1371/journal.pone.0002521.

116. Kuroda, K., Han, H., Tani, S., Tanigaki, K., Tun, T., Furukawa, T., Taniguchi, Y., Kurooka, H., Hamada, Y., Toyokuni, S., and Honjo, T. (2003). Regulation of marginal zone B cell development by MINT, a suppressor of Notch/RBP-J signaling pathway. Immunity *18*, 301-312. 10.1016/s1074-7613(03)00029-3.

117. Oswald, F., Winkler, M., Cao, Y., Astrahantseff, K., Bourteele, S., Knochel, W., and Borggrefe, T. (2005). RBP-Jkappa/SHARP recruits CtIP/CtBP corepressors to silence Notch target genes. Mol Cell Biol *25*, 10379-10390. 10.1128/MCB.25.23.10379-10390.2005.

118. Shi, Y., Downes, M., Xie, W., Kao, H.Y., Ordentlich, P., Tsai, C.C., Hon, M., and Evans, R.M. (2001). Sharp, an inducible cofactor that integrates nuclear receptor repression and activation. Genes Dev *15*, 1140-1151. 10.1101/gad.871201.

119. Tsuji, M., Shinkura, R., Kuroda, K., Yabe, D., and Honjo, T. (2007). Msx2-interacting nuclear target protein (Mint) deficiency reveals negative regulation of early thymocyte differentiation by Notch/RBP-J signaling. Proc Natl Acad Sci U S A *104*, 1610-1615. 10.1073/pnas.0610520104.

120. Ariyoshi, M., and Schwabe, J.W. (2003). A conserved structural motif reveals the essential transcriptional repression function of Spen proteins and their role in developmental signaling. Genes Dev *17*, 1909-1920. 10.1101/gad.266203.

121. Chang, J.L., Lin, H.V., Blauwkamp, T.A., and Cadigan, K.M. (2008). Spenito and Split ends act redundantly to promote Wingless signaling. Dev Biol *314*, 100-111. 10.1016/j.ydbio.2007.11.023.

122. Chen, F., and Rebay, I. (2000). split ends, a new component of the Drosophila EGF receptor pathway, regulates development of midline glial cells. Curr Biol *10*, 943-946. 10.1016/s0960-9822(00)00625-4.

123. Doroquez, D.B., Orr-Weaver, T.L., and Rebay, I. (2007). Split ends antagonizes the Notch and potentiates the EGFR signaling pathways during Drosophila eye development. Mech Dev *124*, 792-806. 10.1016/j.mod.2007.05.002.

124. Feng, Y., Bommer, G.T., Zhai, Y., Akyol, A., Hinoi, T., Winer, I., Lin, H.V., Cadigan, K.M., Cho, K.R., and Fearon, E.R. (2007). Drosophila split ends homologue SHARP functions as a positive regulator of Wnt/beta-catenin/T-cell factor signaling in neoplastic transformation. Cancer Res *67*, 482-491. 10.1158/0008-5472.CAN-06-2314.

125. Kuang, B., Wu, S.C., Shin, Y., Luo, L., and Kolodziej, P. (2000). split ends encodes large nuclear proteins that regulate neuronal cell fate and axon extension in the Drosophila embryo. Development *127*, 1517-1529. 10.1242/dev.127.7.1517.

126. Piovesan, D., Del Conte, A., Mehdiabadi, M., Aspromonte, M.C., Blum, M., Tesei, G., von Bulow, S., Lindorff-Larsen, K., and Tosatto, S.C.E. (2025). MOBIDB in 2025: integrating ensemble properties and function annotations for intrinsically disordered proteins. Nucleic Acids Res *53*, D495-D503. 10.1093/nar/gkae969.

127. Johnson, P.Z., Kasprzak, W.K., Shapiro, B.A., and Simon, A.E. (2019). RNA2Drawer: geometrically strict drawing of nucleic acid structures with graphical structure editing and highlighting of complementary subsequences. RNA Biol *16*, 1667-1671. 10.1080/15476286.2019.1659081.

128. Wu, B., Eliscovich, C., Yoon, Y.J., and Singer, R.H. (2016). Translation dynamics of single mRNAs in live cells and neurons. Science *352*, 1430-1435. 10.1126/science.aaf1084.

129. Huang, X., Zheng, M., Wang, P., Mok, B.W., Liu, S., Lau, S.Y., Chen, P., Liu, Y.C., Liu, H., Chen, Y., et al. (2017). An NS-segment exonic splicing enhancer regulates influenza A virus replication in mammalian cells. Nature communications *8*, 14751. 10.1038/ncomms14751.

130. Cherney, R.E., Eberhard, Q.E., Giri, G., Mills, C.A., Porrello, A., Zhang, Z., White, D., Trotman, J.B., Herring, L.E., Dominguez, D., and Calabrese, J.M. (2023). SAFB associates with nascent RNAs and can promote gene expression in mouse embryonic stem cells. RNA *29*, 1535-1556. 10.1261/rna.079569.122.

131. Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F. (2013). Genome engineering using the CRISPR-Cas9 system. Nat Protoc *8*, 2281-2308. 10.1038/nprot.2013.143.

132. Grunwald, D., and Singer, R.H. (2010). In vivo imaging of labelled endogenous beta-actin mRNA during nucleocytoplasmic transport. Nature *467*, 604-607. 10.1038/nature09438.

133. Schertzer, M.D., Thulson, E., Braceros, K.C.A., Lee, D.M., Hinkle, E.R., Murphy, R.M., Kim, S.O., Vitucci, E.C.M., and Calabrese, J.M. (2019). A piggyBac-based toolkit for inducible genome editing in mammalian cells. RNA *25*, 1047-1058. 10.1261/rna.068932.118.

134. Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W., Marraffini, L.A., and Zhang, F. (2013). Multiplex genome engineering using CRISPR/Cas systems. Science *339*, 819-823. 10.1126/science.1231143.

135. Braceros, A.K., Schertzer, M.D., Omer, A., Trotman, J.B., Davis, E.S., Dowen, J.M., Phanstiel, D.H., Aiden, E.L., and Calabrese, J.M. (2023). Proximity-dependent recruitment of Polycomb repressive complexes by the lncRNA Airn. Cell Rep *42*, 112803. 10.1016/j.celrep.2023.112803.

136. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.

137. Mudge, J.M., Carbonell-Sala, S., Diekhans, M., Martinez, J.G., Hunt, T., Jungreis, I., Loveland, J.E., Arnan, C., Barnes, I., Bennett, R., et al. (2025). GENCODE 2025: reference gene annotation for human and mouse. Nucleic Acids Res *53*, D966-D975. 10.1093/nar/gkae1078.

138. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930. 10.1093/bioinformatics/btt656.

139. Keane, T.M., Goodstadt, L., Danecek, P., White, M.A., Wong, K., Yalcin, B., Heger, A., Agam, A., Slater, G., Goodson, M., et al. (2011). Mouse genomic variation and its effect on phenotypes and gene regulation. Nature *477*, 289-294. nature10413 [pii]

10.1038/nature10413.

140. Waskom, M. (2021). seaborn: statistical data visualization. Journal of Open Source Software *6*. 10.21105/joss.03021.

141. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079. 10.1093/bioinformatics/btp352.

142. Robinson, J.T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and Mesirov, J.P. (2011). Integrative genomics viewer. Nat Biotechnol *29*, 24-26. 10.1038/nbt.1754.

143. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-based analysis of ChIP-Seq (MACS). Genome Biol *9*, R137. gb-2008-9-9-r137 [pii]

10.1186/gb-2008-9-9-r137.

144. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics *26*, 841-842. 10.1093/bioinformatics/btq033.

145. Bailey, T.L. (2021). STREME: accurate and versatile sequence motif discovery. Bioinformatics *37*, 2834-2840. 10.1093/bioinformatics/btab203.

146. Castanza, A.S., Recla, J.M., Eby, D., Thorvaldsdottir, H., Bult, C.J., and Mesirov, J.P. (2023). Extending support for mouse data in the Molecular Signatures Database (MSigDB). Nat Methods *20*, 1619-1620. 10.1038/s41592-023-02014-7.

147. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A *102*, 15545-15550. 10.1073/pnas.0506580102.

148. Yamazaki, T., and Hirose, T. (2015). The building process of the functional paraspeckle with long non-coding RNAs. Front Biosci (Elite Ed) *7*, 1-41. 10.2741/715.

149. Kong, A.T., Leprevost, F.V., Avtonomov, D.M., Mellacheruvu, D., and Nesvizhskii, A.I. (2017). MSFragger: ultrafast and comprehensive peptide identification in mass spectrometry-based proteomics. Nat Methods *14*, 513-520. 10.1038/nmeth.4256.

150. Consortium, G.T. (2020). The GTEx Consortium atlas of genetic regulatory effects across human tissues. Science *369*, 1318-1330. 10.1126/science.aaz1776.

151. Tang, Z., Li, C., Kang, B., Gao, G., Li, C., and Zhang, Z. (2017). GEPIA: a web server for cancer and normal gene expression profiling and interactive analyses. Nucleic Acids Res *45*, W98-W102. 10.1093/nar/gkx247.

152. Goldman, M., Craft, B., Swatloski, T., Cline, M., Morozova, O., Diekhans, M., Haussler, D., and Zhu, J. (2015). The UCSC Cancer Genomics Browser: update 2015. Nucleic Acids Res *43*, D812-817. 10.1093/nar/gku1073.

153. Mielke, P.W., and Berry, K.J. (2001). Permutation methods : a distance function approach (Springer).

154. Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple hypothesis testing. Journal of the Royal Statistical Society *57*, 289-300. https://www.jstor.org/stable/2346101.

155. MathWorks. MATLAB version: 9.13.0 (R2022b). https://www.mathworks.com.

156. Hulsen, T. (2022). DeepVenn -- a web application for the creation of area-proportional Venn diagrams using the deep learning framework Tensorflow.js. arXiv, arXiv:2210.04597. 10.48550/arXiv.2210.04597.

157. Kania, E.E., Fenix, A., Marciniak, D.M., Lin, Q., Bianchi, S., Hristov, B., Li, S., Camplisson, C.K., Fields, R., Beliveau, B.J., et al. (2024). Nascent transcript O-MAP reveals the molecular architecture of a single-locus subnuclear compartment built by RBM20 and the TTN RNA. bioRxiv. 10.1101/2024.11.05.622011.

158. da Veiga Leprevost, F., Haynes, S.E., Avtonomov, D.M., Chang, H.Y., Shanmugam, A.K., Mellacheruvu, D., Kong, A.T., and Nesvizhskii, A.I. (2020). Philosopher: a versatile toolkit for shotgun proteomics data analysis. Nat Methods *17*, 869-870. 10.1038/s41592-020-0912-y.