A Conserved Mycobacterial Nucleomodulin Hijacks the Host COMPASS Complex to Reprogram Pro-Inflammatory Transcription and Promote Intracellular Survival

Liu Chen[1,2], Baojie Duan[1], Pingping Chen[1], Qiang Jiang[1], Yifan Wang[1], Lu Lu[1], Yingyu Chen[1,3,4], Changmin Hu[1,3], Lei Zhang[1,3,4]* and Aizhen Guo[1,2,3,4]*

[1]   National Key Laboratory of Agricultural Microbiology, College of Veterinary Medicine, Huazhong Agricultural University, Wuhan 430070, China

[2]   Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan 430070, China

[3]   National Professional Laboratory for Animal Tuberculosis, Ministry of Agriculture and Rural Affairs, Huazhong Agricultural University, Wuhan 430070, China

[4]   Hubei Jiangxia Laboratory, Wuhan 430207, China

*Address correspondence to: Lei Zhang, zhanglei2023@mail.hzau.edu.cn; Aizhen Guo, aizhen@mail.hzau.edu.cn

ABSTRACT

Nucleomodulins are a class of effector proteins secreted by bacterial pathogens that

translocate into the host cell nucleus to modulate nuclear processes. However, their target

proteins and underlying molecular mechanisms remain poorly understood in mycobacteria.

Herein, we identified a conserved hypothetical protein Rv1075c, designated MgdE, as a

nucleomodulin that enhances mycobacterial intracellular survival. MgdE undergoes nuclear

translocation via two nuclear localization signals, $KRIR^{108-111}$ and $RLRRPR^{300-305}$, and

interacts with ASH2L and WDR5, two subunits of the host histone methyltransferase

COMPASS complex. This interaction suppresses histone H3 lysine 4(H3K4) methylation-

mediated transcription of pro-inflammatory genes, including *il6* and *il1β*, thereby promoting

mycobacterial survival in both macrophages and mice. Our study provides the first

experimental evidence that a bacterial nucleomodulin facilitates intracellular survival by

directly targeting the host COMPASS complex. These findings advance our understanding of

mycobacterial pathogenesis by revealing a novel mechanism that contributes to its

intracellular survival strategy.

Keywords Mycobacteria; Nucleomodulin; Histone methyltransferase; Inflammatory genes;

Pathogenesis.

INTRODUCTION

Nucleomodulins constitute a distinct class of bacterial proteins capable of translocating into

the nuclei of eukaryotic cells, where they strategically modulate nuclear processes to

promote pathogen survival and persistence (Radoshevich et al., 2018; Hanford et al., 2021).

These evolutionarily diverse effector proteins, secreted by distinct pathogens, exhibit

remarkable adaptability in infiltrating host nuclei and reprogramming the transcriptional and

epigenetic regulatory networks (Bierne et al., 2012; Bierne et al., 2020). Upon nuclear

localization, nucleomodulins employ diverse molecular mechanisms to subvert host cellular

processes, including but not limited to: (i) direct interactions with genomic DNA or nuclear

proteins (Rennoll-Bankert et al., 2015; Sun et al., 2016; Prokop et al., 2017), (ii) hijacking

post-translational modification systems (Lebreton et al., 2014; Rolando et al., 2023), (iii)

dysregulating signaling molecule activity or localization (Zhao et al., 2019; Chambers et al.,

2020; Evans et al., 2018), and (iv) destabilizing nuclear architecture (Pourpre et al., 2022; Fu

et al., 2021). Specific examples include AnkA, a bacterial protein from *Anaplasma*

phagocytophilum, which binds to specific A/T-rich sequences in the host cell, alters

chromatin architecture, and modulates gene transcription (Rennoll-Bankert et al., 2015). In

Listeria monocytogenes, LntA interacts with the proline-rich domain of BAHD1, to disrupt

BAHD1-mediated gene silencing and promote histone H3 acetylation (Lebreton et al., 2014).

Another example is OspF from *Shigella flexneri*, which exploits its phosphothreonine lyase

activity to irreversibly dephosphorylate MAPKs, thereby preventing the transcriptional

activation of NF-κB-regulated genes (Chambers et al., 2020). Unlike these effectors, *L.*

monocytogenes InlP takes a different approach by hijacking the splicing factor RBM5. This

interaction not only suppresses apoptosis but also reorganizes nuclear speckles through

SC35 redistribution and RBM5-dependent granule formation (Pourpre et al., 2022). These

examples illustrate the diverse and sophisticated mechanisms used by nucleomodulins to

reprogram host nuclear functions, ultimately manipulating the intracellular environment to

their advantage.

The evolutionary refinement of nucleomodulins is exemplified by their ability to exploit

chromatin-based regulatory systems, with histone methylation emerging as a pivotal axis in

host-pathogen interactions (Bierne et al., 2020; Khan et al., 2021). As a central epigenetic

modification, histone methylation dynamically controls chromatin accessibility and

transcriptional outcomes (Yu et al., 2024; Wang et al., 2025). By precisely targeting specific

histone residues, nucleomodulins reprogram chromatin states to either suppress host

immune responses or promote cellular pathways conducive to pathogen survival (Lebreton

et al., 2014; Schator et al., 2023; Rolando et al., Denzer et al., 2020). A paradigm of this

molecular strategy is demonstrated by the *Chlamydia trachomatis* nuclear effector NUE,

which utilizes its conserved SET domain to catalyze the methylation of histones H2B, H3,

and H4, establishing a repressive chromatin state that downregulates antimicrobial gene

expression (Fol et al., 2020). Similarly, RomA uses its ankyrin and SET domains to bind and

methylate histone H3, thereby altering the host chromatin to enhance *Legionella* intracellular

survival (Schator et al., 2023; Rolando et al., 2013). *Legionella* LegAS4 catalyzes H3K4

methylation to promote an open chromatin state, enhancing ribosomal RNA transcription and

facilitating bacterial intracellular replication (Denzer et al., 2020). These cases emphasize

how pathogens exploit histone methylation as a molecular lever to hijack the host epigenetic

machinery.

Mycobacterium tuberculosis (Mtb), the causative agent of tuberculosis, is a highly

successful pathogen because of its ability to survive within macrophages, evade the immune

system, and persist in the host, ultimately leading to a chronic disease state (Chandra et al.,

2022; Chai et al., 2020). Central to its pathogenic strategy is the secretion of a diverse array

of effector proteins (Chai et al., 2022; Bates et al., 2024; Qiang et al., 2023), including

nucleomodulins that subvert host nuclear processes. The tyrosine phosphatase PtpA

exemplifies this strategy by directly binding to the promoter regions of immune-related

genes, including *GADD45A*, to suppress innate immune responses (Wang et al., 2017). The

methyltransferase Rv1988 mediates an unconventional histone modification by methylating

histone H3 at arginine 42 (H3R42me), thereby silencing antimicrobial defense mechanisms

(Yaseen et al., 2015). Similarly, the acetyltransferase Rv3423.1 modulates chromatin

accessibility through acetylation of histone H3 at lysine 9 and 14 (H3K9/K14ac), attenuating

pro-inflammatory signaling pathways (Jose et al., 2016). Rv2966c represents a unique

bacterial DNA methyltransferase that establishes non-canonical CpG methylation patterns to

epigenetically silence host immune-related gene expression (Sharma et al., 2015). Despite

these molecular insights, the comprehensive landscape of the nuclear effector systems of

Mtb, particularly the mechanistic details of nucleomodulin interactions with host chromatin-

modifying complexes and their spatiotemporal regulation during infection, remains poorly

defined. This critical knowledge gap impedes our understanding of how Mtb coordinates

system-level transcriptional rewiring in the host cellular environment.

In this study, we utilized bioinformatics analysis combined with fluorescence imaging to

rapidly screen for potential nucleomodulins in pathogenic mycobacterial species. Using this

approach, the hypothetical protein Rv1075c (designated as MgdE) was identified and

subsequently validated. MgdE undergoes nuclear translocation mediated by two nuclear

localization signals, KRIR[108-111] and RLRRPR[300-305], and interacts with the histone

methyltransferase COMPASS complex subunits ASH2L and WDR5. This interaction

suppresses H3K4me3-mediated inflammatory gene expression, thus dampening host

immune responses and enhancing bacterial survival in macrophages and murine infection

models. Our findings identify MgdE as a nucleomodulin that targets the COMPASS complex,

revealing a novel epigenetic strategy exploited by mycobacteria for intracellular survival.

RESULTS

Functional screening of potential nucleomodulins in pathogenic mycobacterial

species

To identify conserved nucleomodulins in pathogenic mycobacteria, we first conducted a pan-

genomic comparative analysis across four relevant species, including *M. tuberculosis* H37Rv

and H37Ra, *M. bovis* BCG, *M. marinum*, and *M. avium*, to categorize the core genes.

Subsequently, a dual-threshold screening approach was employed to predict the classical

and non-canonical secreted proteins encoded by these shared genes using SignalP 5.0 and

SecretomeP 2.0, with thresholds set at a D-score ≥0.5 (Hammond et al., 2018) and a neural

network (NN) score ≥0.9 (Bendtsen et al., 2005) (Table S1). This analysis yielded 135 high

confidence secreted protein candidates (Table S2). Further analysis of these candidates for

nuclear localization signal (NLS) motifs using cNLS Mapper (NLS score ≥2) revealed that 56

proteins harbored at least one putative NLS. Following systematic classification based on

secretion machinery dependencies, the candidate proteins were categorized into four distinct

pathways: 26 Sec/SPI (signal peptide-driven), 7 Tat/SPI (twin-arginine translocase pathway),

18 Sec/SPII (lipoprotein-targeted), and eight unclassified candidates lacking canonical

secretion motifs (Figure 1A and B, Figure S1A and B). Functional validation of nuclear

trafficking was conducted through the heterologous expression of enhanced green

fluorescent protein (EGFP)-tagged constructs in HEK293T cells, followed by quantitative

profiling of the subcellular distribution using laser confocal fluorescence microscopy. This

orthogonal validation confirmed the nuclear enrichment of 23 effectors, including six Tat/SPI-

associated candidates. Among these, Rv2577 exhibited the most pronounced nuclear

enrichment, displaying a nuclear localization capacity stronger than that of the previously

described nuclear modulator protein Rv1988 (Yaseen et al., 2015). Notably, MgdE not only

localized to the host cell nucleus but also exhibited a distinct punctate distribution within the

nucleus (Figure 1C). Nucleomodulins are effector proteins secreted by pathogens that

translocate into the host cell nucleus to modulate host functions. To assess whether MgdE is

secreted by mycobacteria, recombinant *M. bovis* BCG strains expressing C-terminally

2×Flag-tagged Ag85B (positive control) and MgdE were constructed. Immunoblot analysis

confirmed high levels of Ag85B-2×Flag and MgdE-2×Flag in bacterial lysates. Notably, both

proteins were also detected in the culture supernatants (Figure S1C), indicating that MgdE is

secreted. These findings identify MgdE as a nucleomodulin in mycobacteria and led us to

further investigate the mechanisms underlying its nuclear translocation and functional role

during infection.

MgdE translocates into the host nucleus through its dual NLS

MgdE is a putative GDSL-like lipase characterized by conserved esterase sequence motifs,

with the GDSx motif containing a nucleophilic Ser residue as part of block I (equivalent to the

classical G-x-S-x-G motif of lipases/esterases), and the consensus amino acids Gly, Asn,

and His located in blocks II, III, and V, respectively (Yang et al., 2019). Phylogenetic analysis

across the mycobacterial species revealed strict evolutionary conservation of MgdE, with

most strains encoding two predicted NLS: $KRIR^{108-111}$ (NLS1) and $RLRRPR^{300-305}$ (NLS2)

(Figure 2A and B). To further investigate the nuclear translocation capability of MgdE, MgdE-

EGFP was transfected into HEK293T cells, and its subcellular localization was analyzed at

various time points post-transfection. Confocal microscopy revealed a time-dependent

decrease in the cytoplasmic levels of MgdE, accompanied by its progressive accumulation in

the host cell nucleus. Notably, MgdE displayed a distinct punctate distribution pattern within

the nucleus, with nearly all the protein localizing to the nucleus approximately 36 h post-

transfection (Figure S2). To quantitatively validate these observations, the temporal

dynamics of MgdE expression in HEK293T cells were monitored, and nuclear enrichment

was further assessed through cell fractionation. The results demonstrated a time-dependent

increase in MgdE expression levels, which closely correlated with its gradual and sustained

nuclear accumulation (Figure 2C). Collectively, these findings indicate that MgdE is an

evolutionarily conserved mycobacterial nucleomodulin with nuclear translocation capacity.

To dissect the functional contributions of the predicted NLS motifs, EGFP-tagged NLS

deletion mutants including $MgdE^{\Delta NLS1}$, $MgdE^{\Delta NLS2}$, and $MgdE^{\Delta NLS1-2}$ were constructed and

transfected into HEK293T cells. Fluorescence intensity profiling revealed that individual

deletion of either NLS1 or NLS2 partially reduced the nuclear targeting efficiency, whereas

simultaneous deletion of both the NLS motifs ($MgdE^{\Delta NLS1-2}$) completely abrogated its nuclear

localization (Figure 2D and E). These observations were further confirmed by immunoblot

analysis of nuclear fractions, which showed that at 36 h post-transfection, MgdE-EGFP and

its mutant variants were consistently detected in the cytoplasmic compartment, and nuclear

accumulation of MgdE$^{\Delta NLS1-2}$ was nearly undetectable compared to that of MgdE, MgdE$^{\Delta NLS1}$,

and MgdE$^{\Delta NLS2}$ (Figure 2F). Together, these findings demonstrate that NLS1 and NLS2 are

both required for efficient nuclear translocation of MgdE.

Nuclear localization of MgdE enhances mycobacterial intracellular survival within

macrophages

Given that MgdE exhibits nuclear translocation ability, we sought to determine whether this

ability plays a role in bacterial survival during infection. We generated various recombinant

M. bovis BCG strains, including wild-type BCG (WT), a *mgdE* deletion mutant (ΔMgdE)

(Figure S3A and B), ΔMgdE complemented with wild-type *mgdE* (Comp-MgdE), and ΔMgdE

complemented with NLS-deleted variant (Comp-MgdE$^{\Delta NLS1-2}$). All strains exhibited similar

growth under standard culture conditions (Figure S3C), and subsequently used them to

infect THP-1 macrophages. As shown in Figure 3A, the ΔMgdE mutant strain exhibited

significantly reduced survival compared to the WT strain, with accelerated bacterial

clearance over time. This defect was rescued in the Comp-MgdE strain, with intracellular

survival levels restored to that in the WT strain. Notably, the Comp-MgdE$^{\Delta NLS1-2}$ showed

bacterial clearance equivalent to that in the ΔMgdE strains. These results were consistent

with the infection experiments using RAW264.7 macrophage (Figure 3B). Furthermore,

macrophages infected with the ΔMgdE mutant strains exhibited significant upregulation of

inflammatory cytokines, including *il1β, il6*, *il10,* and the colony-stimulating factors *csf1-csf3*

compared to WT-infected controls (Figure 3C-H). Collectively, these results demonstrated

that MgdE enhances mycobacterial intracellular survival and suppresses host inflammatory

responses during infection, with both NLS1 and NLS2 motifs being essential for its nuclear

trafficking-dependent function.

MgdE physically interacts with the host COMPASS complex subunits ASH2L and

WDR5

To identify the potential host target proteins of MgdE, we employed AlphaFold to predict the

interactions between MgdE and majority of the nuclear proteins in the host. The prediction

results showed that MgdE binds to WDR5, a core component of the histone

methyltransferase COMPASS complex, with high confidence (pLDDT = 0.77) (Figure S4A

and B). To experimentally validate this prediction, we performed yeast two-hybrid (Y2H)

assays. Strain co-expressing MgdE with either ASH2L or WDR5 exhibited robust growth on

selective SD/-Trp-Leu-His-Ade medium, whereas strain co-expressing MgdE with DPY30 or

RbBP5 showed no detectable growth, confirming specific interactions between MgdE and

ASH2L/WDR5 (Figure 4A). Further, these interactions were validated by co-

immunoprecipitation (Co-IP) assays. HEK293T cells were co-transfected with Flag-tagged

MgdE and HA-tagged ASH2L, WDR5, or RbBP5. The results revealed that HA-tagged

ASH2L and WDR5, but not RbBP5, co-precipitated with Flag-tagged MgdE (Figure 4B).

Conversely, Flag-tagged MgdE precipitated HA-tagged ASH2L and WDR5, but not RbBP5

(Figure 4C). Taken together, these results demonstrate that MgdE specifically interacts with

the COMPASS complex subunits ASH2L and WDR5.

The conserved residues D^{224} and H^{247} are critical for mediating the binding of MgdE to

WDR5

To further confirm the interactions between MgdE and WDR5/ASH2L, the key amino acids of

MgdE (S[80], D[244], and H[247]) were individually replaced with alanine (A) (Yang et al., 2019).

The Y2H assays showed that single-residue substitutions of MgdE did not affect its

interaction with ASH2L or WDR5. However, the D[244]A/H[247]A double mutation specifically

abrogated its interaction with WDR5 but retained its ASH2L binding ability (Figure 5A).

These findings were further confirmed by Co-IP assays in HEK293T cells. Flag-tagged

MgdE precipitated HA-tagged ASH2L and WDR5 but not RbBP5, with reciprocal pull-downs

confirming bidirectional binding specificity. The D[244]A/H[247]A double mutant maintained its

interaction with ASH2L but failed to bind WDR5, suggesting that these interactions are

mediated through distinct molecular interfaces (Figure 5B, Figure S5). To further investigate

the functional consequences of these mutations, we analyzed the subcellular localization of

the MgdE mutants in transfected HEK293T cells using fluorescence co-localization assays.

Wild-type MgdE exhibited progressive accumulation of punctate nuclear foci over time, a

phenomenon that was abrogated following transfection with the D[244]A/H[247]A mutant (Figure

5C), suggesting that the double mutation impairs MgdE activity.

Given that MgdE interacts with WDR5, a subunit of COMPASS complex, which is a key

regulator of H3K4me3 methylation (Deng et al., 2024), we sought to determine whether this

interaction contributes to an epigenetic regulatory mechanism that facilitates bacterial

persistence within host cells. We transfected HEK293T cells with the EGFP-tagged WT

MgdE and mutated MgdE-D[244]A/H[247]A, and monitored H3K4me3 modifications over time.

Strikingly, immunoblotting of nuclear extracts collected at defined time points revealed that

cells expressing WT MgdE exhibited approximately 2-fold lower H3K4me3 levels compared

to those expressing the MgdE-D^{244}A/H^{247}A mutant, an effect detectible within 4–12 h post-

transfection (Figure 5D). These findings indicate that MgdE inhibits COMPASS-dependent

methyltransferase activity via its functional interaction with WDR5.

Together, these data indicate that the residues D^{244} and H^{247} are essential for the MgdE-

WDR5 interaction, which enables MgdE to attenuate H3K4me3 deposition in the host nuclei

by subverting COMPASS complex function, a mechanism critical for bacterial survival and

immune evasion.

MgdE suppresses inflammatory responses mediated by the COMPASS complex via

inhibition of H3K4 methylation

As H3K4me3, a hallmark of transcriptionally active promoters, facilitates transcriptional

machinery assembly and RNA polymerase II recruitment (Wang et al., 2023), we further

investigated the transcriptional consequences of MgdE-mediated H3K4me3 suppression via

its interaction with the COMPASS complex. To assess the genome-wide effects of MgdE on

host gene expression, we performed RNA-seq profiling of THP-1 macrophages infected with

either WT BCG (BCG/wt) or MgdE-deleted (Ko*mgdE*) strains. Transcriptomic analysis at 24

h post-infection revealed that 271 genes showed significant differential expression in the

cells infected with Ko*mgdE* strain compared to that in the cells infected with WT BCG strain,

with 222 genes being transcriptionally upregulated in the absence of MgdE (Figure 6A).

Gene ontology (GO) enrichment analysis indicated that these upregulated genes were

primarily associated with biological processes involved in immune responses, with a notable

enrichment in "Positive regulation of cytokine production" (GO: 0001819), a critical pathway

for amplifying inflammatory signaling (Figure 6B). Furthermore, the Kyoto encyclopedia of

genes and genomes (KEGG) pathway enrichment analysis highlighted a significant

activation of the "Cytokine-cytokine receptor interaction" and "JAK-STAT signaling pathway"

in THP-1 cells infected with the Ko*mgdE* strain (Figure 6C, Figure S6A). Heatmap analysis

further confirmed the marked upregulation of inflammation-associated transcripts, including

inflammatory cytokines (*il6*, *il10,* and *il12b*), chemokines (*ccl1-ccl5*), and colony-stimulating

factors (*csf1-csf3*), in Ko*mgdE*-infected cells (Figure 6D, Figure S6B). Protein-protein

interaction network analysis of the upregulated genes revealed tightly interconnected

clusters within inflammatory networks, particularly those orchestrated by the central

inflammatory mediator IL-6 (Figure S6C). To validate this result, we infected macrophages

with the *M. bovis BCG* strains, including WT, ΔMgdE, Comp-MgdE, and Comp-MgdE$^{\Delta NLS1-2}$,

under identical conditions for 24 h and examined the expression of relevant inflammatory

factors. Compared to cells infected with the WT and Comp-MgdE strains, those infected with

the ΔMgdE and Comp-MgdE$^{\Delta NLS1-2}$ strains exhibited significantly increased expression levels

of inflammatory factors (*il6, il10,* and *il12b*) (Figure 6E-G). Collectively, these results

demonstrate that MgdE effectively inhibits the production of inflammatory factors during BCG

infection and highlights the critical role of its NLSs in this process. Through its interaction

with the COMPASS complex, MgdE suppresses H3K4me3 levels, thereby attenuating the

transcriptional activation of inflammation-related genes.

Nuclear localization of MgdE is essential for mycobacterial survival in mice

Our experimental data demonstrated that MgdE enhances bacterial survival in macrophages.

To further investigate the *in vivo* functional significance of MgdE in host-pathogen interplay,

we infected C57BL/6 mice with different *M. bovis* BCG strains, including WT, ΔMgdE, Comp-

MgdE, and Comp-MgdE$^{\Delta NLS1-2}$, and comprehensively evaluated bacterial survival, pulmonary

inflammation, and cytokine dynamics. Mice infected with ΔMgdE exhibited a significantly

reduced lung bacterial burden compared to those infected with WT or Comp-MgdE strains.

Moreover, mice infected with Comp-MgdE$^{\Delta NLS1-2}$ exhibited a significantly lower pulmonary

bacterial load than those infected with Comp-MgdE (Figure 7A). These findings collectively

establish that MgdE is essential for optimal bacterial survival during infection, with the NLS

being critical for its full functionality. Furthermore, histopathological assessment of the lung

tissues corroborated these observations. Hematoxylin-eosin (H&E) staining revealed more

pronounced inflammatory pathology in the lungs of mice infected with WT or Comp-MgdE

strain compared to those infected with ΔMgdE or Comp-MgdE$^{\Delta NLS1-2}$ (Figure 7B),

underscoring the necessity of MgdE and its NLSs in sustaining inflammation during infection.

To further investigate the systemic immune modulation mediated by MgdE, we quantified

splenic bacterial colonization and the inflammatory cytokine profiles. Consistent with the

attenuated bacterial survival observed in the lungs, mice infected with ΔMgdE and Comp-

MgdE$^{\Delta NLS1-2}$ exhibited a significantly reduced splenic bacterial load compared to those

infected with WT or omp-MgdE (Figure S7). Despite diminished bacterial dissemination,

infections with ΔMgdE and Comp-MgdE$^{\Delta NLS1-2}$ strains triggered elevated expression of

inflammatory cytokines (*il-1β, il-6*) relative to infection with the WT and Comp-MgdE strains

(Figure 7C and D). Collectively, these findings indicate that MgdE orchestrates

mycobacterial survival within host tissues by suppressing inflammatory responses to evade

pathogen clearance, with the NLS playing a pivotal role in mediating these host-pathogen

interactions.

DISCUSSION

MgdE functions as a critical nucleomodulin that contributes to mycobacterial pathogenesis. Its evolutionary conservation across mycobacterial species, coupled with nuclear translocation via dual N-terminal and C-terminal NLSs, supports its classification as a nucleomodulin. Previous investigations have described MgdE as a secreted effector protein (Penn et al., 2018), and our experimental results independently confirm its secretion (Figure S1C). Furthermore, our findings demonstrate that MgdE's nuclear localization plays a critical role during infection. Disruption of MgdE expression or its NLS motifs significantly attenuated *M. bovis* BCG viability within macrophages (Figure 3A and B) and reduced bacterial burden in mice (Figure 7A and Figure S7), supporting the notion that nuclear targeting contributes to bacterial virulence. This phenotype aligns with the observed upregulation of MgdE during Mtb infection of alveolar macrophages (Pisu et al., 2020), suggesting that nuclear activity of MgdE may contribute to an adaptive persistence mechanism during infection. Although MgdE was previously characterized as a lipase (Yang et al., 2019), our findings reveal a novel functional duality wherein nuclear targeting coordinates bacterial survival through epigenetic interference while preserving its metabolic roles. This functional pleiotropy positions MgdE as a master virulence regulator, exploiting host-pathogen interface plasticity to sustain infection.

MgdE directly subverts the COMPASS complex to reprogram host epigenetics. To our knowledge, MgdE is the first identified bacterial effector that directly targets and subverts the COMPASS complex, a central regulator of H3K4 methylation. Nuclear-localized MgdE specifically interacts with the core COMPASS subunits ASH2L and WDR5, disrupting

H3K4me3 deposition through structural interference. The COMPASS complex facilitates

H3K4 methylation via a conserved assembly mechanism in which WDR5 scaffolds

methyltransferases such as MLL1 through its WIN motif and stabilizes the ASH2L–RbBP5

dimer (Guarnaccia et al., 2021; Hsu et al., 2018). This dimer, in turn, positions the SET

domain for efficient substrate recognition (Xue et al., 2019). MgdE appears to interfere with

this assembly process, resembling strategies employed by host regulatory proteins.

Site-directed mutagenesis studies have identified critical residues (D^{224} and H^{247}) within

MgdE that are essential for WDR5 binding. Mutating these residues abolished both WDR5

interaction and H3K4me3 suppression (Figure 5C, D), confirming their functional relevance.

Moreover, nuclear MgdE accumulates in punctate subnuclear foci (Figure 1C, Figure 2D,

Figure 5C), reminiscent of *L. monocytogenes* InlP, which modulates nuclear speckles to

inhibit apoptosis (Pourpre et al., 2022). This interaction silences the expression of

antimicrobial effector genes, thereby facilitating immune evasion. Similar to *Legionella*

pneumophila RomA, which methylates H3K14 to repress innate immune responses (Schator

et al., 2023; Rolando et al., 2013), or *M. tuberculosis* Rv2067c, a structural mimic of host

DOT1L that catalyzes non-nucleosomal H3K79 trimethylation to subvert pro-inflammatory

signaling (Singh et al., 2023), MgdE exemplifies a broader pathogen strategy of hijacking the

host histone methylation machinery.

MgdE-mediated epigenetic silencing suppresses pro-inflammatory responses.

Transcriptome analysis showed that the *mgdE*-deleted BCG strain induced hyperactivation

of cytokine-cytokine receptor interactions and the JAK-STAT signaling pathway, along with

significantly elevated production of pro-inflammatory cytokines, including *il6* and *il1β*, in THP-

1 macrophages compared to the WT BCG strain (Figure 3C-F, Figure 6E-F). This profile is

consistent with the functional restoration of the COMPASS complex (Wang et al., 2025; Yu et

al., 2017). These findings support the hypothesis that MgdE suppresses host inflammatory

responses by disrupting COMPASS-mediated H3K4me3 deposition at active promoters.

H3K4me3 is a well-established histone mark associated with transcriptional activation, in

part through the recruitment of PHD domain-containing reader proteins (Hyun et al., 2017).

For instance, SET1-dependent H3K4me3 accumulation at NF-κB target promoters such as

il6 and *tnfα* enhances pro-inflammatory gene expression (Bhattacharya et al., 2023). In

addition, WDR5 has been implicated in upregulating immunosuppressive cytokines,

including *il6* and *tgfβ* (Deng et al., 2024). In transfected HEK293T cells, WT MgdE

significantly reduced global H3K4me3 levels, whereas the catalytically inactive $D^{224}A/H^{247}A$

mutant failed to do so (Figure 5D), supporting the notion that MgdE-mediated suppression of

H3K4me3 depends on its ability to interact with WDR5. Furthermore, during infection, the

MgdE-deleted BCG strain markedly activated the expression of inflammatory factors

compared to the WT BCG strain (Figure 6C-G). These results suggest that MgdE attenuates

the expression of inflammatory mediators such as *il6* and *il1β* by reducing host H3K4me3

levels during Mycobacterium infection. This mechanism is similar to those observed in other

pathogens, such as spirochete-derived factors that suppress inflammation by reducing

H3K4me3 levels at the *tnfα* and *il-6* promoters (Chauhan et al., 2015).

In summary, our study identified MgdE as a critical mycobacterial nucleomodulin and

uncovered a novel paradigm of pathogen-mediated epigenetic regulation through the "MgdE-

COMPASS complex-H3K4me3-cytokine" axis (Figure 7F). Mechanistically, MgdE specifically

interacts with ASH2L and WDR5 to destabilize the COMPASS assembly, resulting in

reduced H3K4me3 levels at the promoters of pro-inflammatory genes. This epigenetic

suppression downregulates cytokine transcription (e.g., *il6 and il1β*) and enhances bacterial

survival in macrophages and in mice. Our findings reveal the COMPASS complex as a

previously unrecognized target of bacterial effectors and offers mechanistic insights into

immune evasion and host-pathogen epigenetic interplay.

MATERIALS AND METHODS

Bacterial strains and culture conditions

The bacterial strains used in this study are detailed in Table S4. *Escherichia coli* DH5α was

cultured in Luria-Bertani medium under standard conditions. *Mycobacterium bovis BCG*

strains were grown in Middlebrook 7H9 broth (BD Biosciences), containing 1× OADC (oleic

acid, albumin, dextrose, catalase), 0.05% Tween-80, and 2% glycerol. Antibiotics were

added as required at the following final concentrations: 50 µg/mL hygromycin (for

mycobacteria) and 50 µg/mL kanamycin (for both mycobacteria and *E. coli*). The Y2HGold

yeast strain (Takara Bio) was maintained in YPDA (yeast extract, peptone, dextrose, and

adenine sulfate) medium. Transformants were selected on SD/-Trp medium, which lacks

tryptophan, to select for plasmid uptake. Protein-protein interactions were assessed in SD/-

Trp/-Leu (lacking tryptophan and leucine) and SD/-Trp/-Leu/-Ade/-His (lacking tryptophan,

leucine, adenine, and histidine) media, with growth on higher stringency media indicating

positive interactions.

Cell culture

THP-1 cells were cultured in Roswell Park memorial institute 1640 (RPMI 1640) medium

supplemented with 10% (v/v) heat-inactivated fetal bovine serum (FBS), 1 mM sodium

pyruvate, 2 mM L-glutamine, 10 mM HEPES buffer (pH 7.2–7.5), and 50 µM 2-

mercaptoethanol. For differentiation into macrophages, THP-1 cells were treated with 200

nM phorbol 12-myristate 13-acetate (PMA) for 24 h, followed by an additional 48-h

incubation in fresh PMA-free medium to allow complete differentiation prior to their use in

experiments.

HEK293T and RAW264.7 monocytes were cultured in Dulbecco's modified Eagle's medium

(DMEM) supplemented with 10% FBS and 50 µg/mL penicillin-streptomycin. All cells were

maintained at 37 °C in a humidified atmosphere with 5% CO_2.

Pan-genomic nucleomodulin screening

To systematically identify evolutionarily conserved nucleomodulins in pathogenic

mycobacteria, a pan-genomic comparative analysis was performed across four

phylogenetically representative mycobacterium strains: *Mycobacterium tuberculosis* H37Rv

(NCBI RefSeq assembly GCF_000195955.2), *Mycobacterium tuberculosis* H37Ra (NCBI

RefSeq assembly GCF_001938725.1), *Mycobacterium bovis* BCG (NCBI RefSeq assembly

GCF_000009445.1), *Mycobacterium marinum* (NCBI RefSeq assembly GCF_000018345.1),

and *Mycobacterium avium* subsp. (NCBI RefSeq assembly GCF_009741445.1). Complete

genome sequences and annotations were obtained from NCBI RefSeq with strict inclusion

criteria requiring "Complete Genome" status and fewer than 50 contigs to ensure the

assembly integrity. Plasmid sequences, as well as low-complexity and repetitive regions,

were filtered using the NCBI Genome Data Viewer toolkit. Genome quality control was

rigorously conducted using CheckM v1.2.2 with a *Mycobacteriaceae*-specific marker set

(including essential housekeeping genes such as *dnaA* and *rpoB*), confirming that all the

selected genomes exhibited >99% completeness and <1% contamination.

Ortholog clustering of protein-coding sequences (CDS) was executed through OrthoFinder

v2.5.4, employing Diamond alignment in ultra-sensitive mode (--ultra-sensitive), an E-value

cutoff of ≤1e-5 (--blastp_evalue 1e-5), and a minimum sequence identity threshold of ≥50%

(--min_percent_identity 50). The Markov cluster algorithm (MCL) was applied with an

inflation parameter of 1.5 (--mcl_inflation 1.5) to define orthologous groups, balancing

sensitivity, and specificity. Core orthologs were identified as clusters present in all four

strains, with additional filtering to retain only single-copy genes in ≥90% of the strains,

minimizing paralog-related artifacts.

Functional annotation of the core genes was conducted using InterProScan v5.62, which

integrates domain predictions from the Pfam, TIGRFAM, and SUPERFAMILY databases. To

prioritize candidate nucleomodulins, a sequential screening pipeline was applied, beginning

with the prediction of classical secretory proteins using SignalP v5.0 (D-score ≥0.5), followed

by the identification of non-classically secreted proteins via SecretomeP v2.0 (NN score

≥0.9), and culminating in NLS detection using cNLS Mapper (NLS score ≥2.0), which

predicts both monopartite and bipartite NLSs. This integrated strategy enabled the

systematic identification of conserved candidates that exhibit both secretory potential and

nuclear targeting capacity.

Recombinant expression and secretion of MgdE

Recombinant *M. bovis* BCG strains expressing C-terminally 2×Flag-tagged Ag85B (positive

control) and MgdE were constructed to assess protein secretion under standard in vitro

culture conditions. The strains were cultured in 200 mL of Middlebrook 7H9 broth

supplemented with albumin-dextrose-catalase (ADC), glycerol, and 0.05% Tween-80 at

37 °C until reaching mid-log phase (OD_{600} = 0.6). Bacterial cultures were centrifuged at

4,000 × g for 60 minutes to separate cells from supernatants. Pellets were washed and lysed

by bead beating in PBS supplemented with protease inhibitors. Culture supernatants were

passed through 0.22 µm filters to remove residual bacteria and subsequently concentrated to

2 mL using Amicon Ultra centrifugal filters (10 kDa cutoff). Both bacterial lysates and

concentrated supernatants were subjected to SDS-PAGE followed by immunoblotting with

anti-Flag antibodies to detect the expression and secretion of 2×Flag-tagged proteins.

Phylogenetic and sequence analysis

To investigate the evolutionary conservation and sequence features of MgdE homologs,

phylogenetic reconstruction was performed using MEGA 12.0 with a neighbor-joining

algorithm. The parameters included pairwise deletion for gap treatment, a Poisson

substitution model (selected for its suitability in analyzing closely related bacterial

sequences), and 1,000 bootstrap replicates to assess node support. Nodes with ≥50%

bootstrap values were retained, and no additional pruning was performed.

The analysis incorporated MgdE homologs from 12 Mycobacterium strains spanning clinical,

environmental, and attenuated lineages. These included Rv1075c (*M. tuberculosis* H37Rv),

BCG_1133c (*M. bovis* BCG), MRA_1085 (*M. tuberculosis* H37Ra), MAF_10880 (*M.*

africanum), MT1105 (*M. tuberculosis* CDC1551), TBMG_02912 (*M. tuberculosis*

GM041182), MSHI_02760 (*M. shinjukuense*), MMAR_4392 (*M. marinum*), MULP_04607 (*M.*

lupini), MAV_1199 (*M. avium*), MMAN_54310 (*M. manitobense*), and MIP_01781 (*M.*

intracellulare). Genomic sequences were retrieved from the NCBI RefSeq database, filtered

for completeness, and aligned using Clustal Omega v1.2.4 with default parameters (gap

opening penalty = 10, gap extension = 0.2, and iterative refinement enabled). Sequence

conservation was visualized using ESPript 3.0, highlighting residues with ≥90% identity

across all homologs in red.

Computational prediction of sequence and structural features

AlphaFold v2.2.0 was used to predict the interaction between MgdE and COMPASS

complex subunits, following the methodology established in a previous study (Gómez

Borrego et al., 2024). The specific parameters used for the prediction were as follows:

model_preset = multimer; db_preset = full_dbs. A FASTA file containing the sequences of

both MgdE and COMPASS complex subunit proteins was input into the AlphaFold software.

The resulting models were evaluated and ranked based on the AF-score, which is a linear

combination of the interface score (ipTM) and the predicted TM-score (pTM). The top-ranked

model (rank 1) was selected for further analysis.

Cell transfection and confocal microscopy

HEK293T cells were seeded at 70% confluency in poly-L-lysine-coated 35-mm glass-bottom

dishes and maintained in DMEM supplemented with 10% FBS at 37 °C under 5% CO_2.

Transfection was performed using the Hieff Trans™ Liposomal Transfection Reagent

(Yeasen) according to the manufacturer's protocol. Briefly, 0.5 μg of DNA and 1.5 μL of

transfection reagent were mixed with 50 μL of Opti-MEM and incubated for 20 min at room

temperature. The mixture was added dropwise to each chamber of 35-mm six-chamber

glass-bottom dishes containing cells at 40–60% confluency.

At 24–48 h post-transfection, the cells were washed with phosphate-buffered saline (PBS),

fixed in 4% paraformaldehyde for 15 min, and permeabilized with 0.1% Triton X-100 for 10

min. Nuclei were counterstained with 1 µg/mL DAPI (4′,6-diamidino-2-phenylindole) for 10

min. Confocal imaging was performed using an Olympus FV1000 Confocal Laser Scanning

Microscope equipped with a 100×/1.40 NA oil immersion objective. EGFP and DAPI signals

were detected with 488 nm and 405 nm lasers, respectively. Spectral unmixing was applied

using the Olympus FV10-ASW software (v4.2) to minimize signal cross-talk, and Z-stacks

were acquired in sequential mode by applying a step size of 0.5 µm across all samples to

ensure comparability.

Immunoblot analysis

For immunoblotting, the cells were lysed in radioimmunoprecipitation assay (RIPA) buffer

supplemented with 1.5% protease inhibitor cocktail (Boster). Protein concentrations were

quantified using the BCA assay (Thermo Fisher Scientific). Equal amounts of protein (20–40

µg) were resolved by sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-

PAGE) and transferred onto polyvinylidene difluoride (PVDF) membranes (Millipore). The

membranes were blocked for 1 h at room temperature with 5% non-fat milk in TBST (25 mM

Tris, 150 mM NaCl, 2 mM KCl, and 0.2% Tween-20, pH 7.4) and incubated overnight at 4 °C

with primary antibodies. After three washes with TBST (10 min each), the membranes were

incubated with horseradish peroxidase (HRP)-conjugated secondary antibodies at room

temperature for 1 h. Following three additional TBST washes, protein signals were detected

using an enhanced chemiluminescence (ECL) substrate and imaged on a ChemiDoc MP

system (Bio-Rad). β-actin, GAPDH, or histone H3 served as housekeeping proteins for

normalization. The antibody information is listed in Table S3.

Cell fractionation was performed using a Nuclear and Cytoplasmic Protein Extraction Kit

(Beyotime) according to the manufacturer's protocol. Briefly, after treatment, the cells were

washed and harvested in cold PBS. Cytoplasmic and nuclear fractions were separated

according to the manufacturer's instructions. Protein concentrations were quantified using a

BCA protein assay kit, and equal amounts of protein were subjected to immunoblotting, as

described above.

Co-IP assays were performed to analyze protein-protein interactions. HEK293T cells were

washed twice with ice-cold PBS and lysed in lysis buffer (50 mM Tris-HCl, pH 7.4, 150 mM

NaCl, 1% NP-40) supplemented with 1.5% protease inhibitor cocktail (Boster) for 30 min on

ice. Crude lysates were centrifuged at 12,000 ×g for 15 min at 4 °C to remove the debris. For

immunoprecipitation, 1 mg of total protein was incubated with 5 µg of primary antibody

overnight at 4 °C with gentle agitation. Protein A/G magnetic beads (MedChemExpress)

were added to the antibody-protein complexes and incubated for 6 h at 4 °C. The beads

were washed three times with PBS-T buffer (PBS containing 0.5% Tween-20) to remove

nonspecific interactions. Proteins were eluted by boiling the beads in 1× SDS sample buffer

at 95 °C for 5 min. The eluted proteins were resolved by SDS-PAGE and analyzed by

immunoblotting.

Y2H assay

The Yeast Two-Hybrid System (Clontech) was used to assay protein-protein interactions

between MgdE (WT and mutants) and subunits of the COMPASS complex (ASH2L, WDR5,

RbBP5, and DPY30). Full-length cDNAs encoding WT MgdE and its mutants (see Table S5)

were cloned into the bait vector pGBKT7, whereas genes encoding the COMPASS subunits

were cloned into the prey vector pGADT7. The yeast strain Y2HGold was co-transformed

with bait and prey plasmids using the lithium acetate/polyethylene glycol method, and the

transformed colonies were initially selected on SD agar plates lacking leucine and

tryptophan (-Leu/-Trp) to confirm plasmid retention. For interaction screening, colonies were

replica-plated onto selective SD agar plates lacking leucine, tryptophan, adenine, and

histidine (-Leu/-Trp/-Ade/-His) and supplemented with 200 ng/µL aureobasidin A (AbA,

Sigma-Aldrich). The plates were incubated at 30 °C, and colony growth was monitored daily,

with final assessments performed after 5 d of incubation. Interaction strength was

qualitatively assessed by comparing colony density and growth rates with those of the

controls. The positive controls included the co-transformation of pGBKT7-p53 with pGADT7-

T, which produced robust growth on -Leu/-Trp/-Ade/-His plates. Negative controls included

co-transformation of pGBKT7-Lam (human lamin C, a non-interacting bait) with pGADT7-T,

as well as individual transformations with empty pGBKT7 or pGADT7 vectors to assess the

background growth. The yeast strains used in this study are listed in Table S4. All

experiments were performed in triplicate with biologically independent yeast cultures. The

results were consistent across replicates, confirming the reliability of the observed

interactions.

qRT-PCR assay

THP-1 cells were infected with the *M. bovis* BCG strain and harvested 12 h post-infection for

RNA isolation. Total RNA was extracted using 1 mL of TRIzol® reagent (Aidlab) according to

the manufacturer's instructions. To remove genomic DNA contamination, the RNA samples

were treated with DNase I (Thermo Fisher Scientific) for 30 min at 37 °C, followed by heat

inactivation at 65 °C for 10 min. RNA concentration and purity were assessed using a

NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific). For cDNA synthesis, 1 µg of

total RNA was reverse transcribed using the HiScript® III cDNA Reverse Transcription Kit

(Vazyme) with random hexamers, according to the manufacturer's protocol. qRT-PCR was

performed using ChamQ SYBR® Green RT-PCR Master Mix (Vazyme). The relative mRNA

expression of different genes was calculated by comparing their cycle threshold (Ct) values

to that of a control gene *hprt* using the $2^{-\Delta\Delta Ct}$ method. Real-time qRT-PCR experiments were

performed in triplicate, with at least three independent biological replicates and two technical

replicates per condition. Primer sequences used for qRT-PCR are listed in Table S6.

RNA-seq analysis

THP-1 cells were seeded at a density of 1×10^6 cells/well in 6-well plates and differentiated

into macrophages by treatment with 100 nM PMA for 48 h. Following differentiation, the cells

were washed with PBS and infected with *M. bovis* BCG WT or *KomgdE* strains at a

multiplicity of infection (MOI) of 10:1. Following 12 h of infection, total RNA was extracted

using TRIzol® reagent (Thermo Fisher Scientific) according to the manufacturer's

instructions. High-quality reads were then mapped to the human reference genome

(GRCh38) using HISAT2 (v2.2.1). FeatureCounts (v2.0.3) was used to quantify transcript

abundance, and differential gene expression analysis was conducted using DESeq2

(v1.38.3) in R. Genes with an adjusted P-value <0.05 and an absolute $\log_2$-foldchange ≥ 1

was considered significantly differentially expressed. RNA-seq data were visualized using

GraphPad Prism 8.0, and volcano plots, heatmaps, and other visualizations were generated

to summarize the results.

CFU assay

For bacterial preparation, mid-log phase cultures (OD_{600} = 0.8) of *M. bovis* BCG were

washed with PBS and resuspended in fresh medium. THP-1 cells were seeded at a density

of 5×10^5 cells/well in 24-well plates and subsequently infected at a multiplicity of infection

(MOI) of 1:10. Following 4 h of infection, extracellular bacteria were eliminated by treatment

with penicillin-streptomycin (100 µg/mL) for 2 h. The cells were then washed three times with

PBS and maintained in an antibiotic-free medium until harvest. For bacterial recovery, the

macrophages were lysed with 0.25% SDS for 10 min at room temperature, followed by serial

dilution of the lysates in PBS. Aliquots (100 µL) of each dilution were plated on Middlebrook

7H10 agar supplemented with 10% OADC and kanamycin (50 µg/mL). Plates were

incubated at 37 °C for 14 d, and CFUs were calculated by applying dilution factors to the

colony counts. *M. bovis* BCG strains used in this study are listed in Table S4.

Animal experiments

Specific pathogen-free (SPF) female C57BL/6 mice (6–8 weeks old) were obtained from

Chang-sheng Bio (Liaoning, China). All animals were housed under SPF conditions under

controlled temperature, humidity, and a 12-hour light/dark cycle, with free access to food and

water.

Mice infection

Mice were infected as described previously, with slight modifications. Mid-log phase *M. bovis* BCG cultures were washed twice in PBS containing 0.05% Tween 80 and sonicated to disrupt clumps. Female C57BL/6 mice (6–8 weeks old, 16–18g) were anesthetized using isoflurane and intranasally administered 1×10^7 CFUs of *M. bovis* BCG in 40 µL of PBS. At 48 h post-infection, the bacterial load in the lungs of five mice was assessed to confirm the inoculation dose. The lungs were homogenized in 1 mL of PBS using a tissue homogenizer, and serial dilutions of the homogenates were plated onto Middlebrook 7H10 agar with 10% OADC. The plates were then incubated at 37 °C with 5% CO_2 for 14 d prior to colony counting.

Histological analysis

Lung tissues from *M. bovis* BCG-infected mice were fixed in 4% phosphate-buffered formalin at room temperature for 24 h and embedded in paraffin wax. Paraffin-embedded tissue samples were sectioned into 2–3 µm thick slices using a microtome. The sections were deparaffinized, rehydrated with graded ethanol, and stained with hematoxylin and eosin (H&E). Stained slides were visualized using an Olympus BX53 light microscope, and CaseViewer version 2.0 (3DHISTECH) was used to view and annotate the scanned slides.

Ethics approval

All animal experiments were conducted in accordance with the protocols approved by the Animal Ethics Committee of the Huazhong Agricultural University.

Statistical analyses

Data are presented as mean ± standard deviation (SD). For comparisons between the two

groups, statistical significance was determined using two-tailed unpaired Student's *t*-tests

after confirming normality. Statistical significance was set at $P < 0.05$. Significance levels are

denoted in the figures as follows: $P > 0.05$ (ns, not significant), $P < 0.05$ (*), $P < 0.01$ (**),

and $P < 0.001$ (***). All statistical analyses were performed using GraphPad Prism (version

8.0). Data were collected from at least three independent biological replicates, each with two

or more technical replicates.

ACKNOWLEDGMENTS

This work was supported by the National Key Research and Development Program of China

(#2021YFD1800403), Fundamental Research Funds for the Central Universities

(#2662023DKQD001), Talent Start-up Funds of Huazhong Agricultural University

(#11042310008), National Natural Science Foundation of China (#32473123), China

Agriculture Research System of MOF and MARA (#CARS-37).

AUTHOR CONTRIBUTIONS

Conceptualization: L. C., A. G., and L. Z.; Methodology: L. C.; Validation: L. C., B. D., and Q.

J.; Formal analysis: L. C., Y. W.; Investigation: L. C., B. D., P. C., Q. J., Y. W., and L. L.;

Resources: A. G., L. Z., and Y. C.; Data curation: L. C., Q. J., Y. W., and L. L.; Writing -

original draft: L. C., L. Z., and A. G.; Writing - review & editing: L. C., L. Z., and A. G.;

Visualization: L. C., Q. J., Y. C., C. H., L. Z., and A. G.; Supervision: A. G., L. Z., Y. C., and C.

H.; Project administration: A. G., L. Z., Y. C., and C. H.; Funding acquisition: A. G., L. Z., and

Y. C.

DISCLOSURE AND COMPETING INTEREST STATEMENT

The authors declare no competing interests.

REFERENCES

Bates TA, Trank-Greene M, Nguyenla X, Anastas A, Gurmessa SK, Merutka IR, Dixon SD,

Shumate A, Groncki AR, Parson MAH (2024) **ESAT-6 undergoes self-association at**

phagosomal pH and an ESAT-6-specific nanobody restricts *M. tuberculosis* growth in

macrophages *Elife* **12**:RP91930.

Bendtsen JD, Kiemer L, Fausbøll A, Brunak S (2005) **Non-classical protein secretion in**

bacteria *BMC Microbiol* **5**:58.

Bhattacharya A, Fushimi A, Wang K, Yamashita N, Morimoto Y, Ishikawa S, Daimon T, Liu T,

Liu S, Long MD (2023) **MUC1-C intersects chronic inflammation with epigenetic**

reprogramming by regulating the set1a compass complex in cancer progression

Commun Biol **6**:1030.

Bierne H, Cossart P (2012) **When bacteria target the nucleus: the emerging family of**

nucleomodulins *Cell Microbiol* **14**:622-633.

Bierne H, Pourpre R (2020) **Bacterial factors targeting the nucleus: the growing family**

of nucleomodulins *Toxins* (Basel) **12**:220.

Chai Q, Wang L, Liu CH, Ge B (2020) **New insights into the evasion of host innate**

immunity by *Mycobacterium tuberculosis* *Cell Mol Immunol* **17**:901-913.

Chai Q, Yu S, Zhong Y, Lu Z, Qiu C, Yu Y, Zhang X, Zhang Y, Lei Z, Qiang L (2022) **A**

bacterial phospholipid phosphatase inhibits host pyroptosis by hijacking ubiquitin

Science **378**:eabq0132.

Chambers KA, Scheck RA (2020) **Bacterial virulence mediated by orthogonal post-**

translational modification *Nat Chem Biol* **16**:1043-1051.

Chandra P, Grigsby SJ, Philips JA (2022) **Immune evasion and provocation by**

Mycobacterium tuberculosis *Nat Rev Microbiol* **20**:750-766.

Chauhan A, Quenum FZ, Abbas A, Bradley DS, Nechaev S, Singh BB, Sharma J, Mishra BB

(2015) **Epigenetic modulation of microglial inflammatory gene loci in helminth-**

induced immune suppression: implications for immune regulation in

neurocysticercosis *ASN Neuro* **7**:1759091415592126.

Deng K, Liang L, Yang Y, Wu Y, Li Y, Zhang R, Tian Y, Lu C (2024) **The Wdr5-H3K4me3**

epigenetic axis regulates pancreatic tumor immunogenicity and immune suppression

Int J Mol Sci **25**:8773.

Denzer L, Schroten H, Schwerk C (2020) **From gene to protein-how bacterial virulence**

factors manipulate host gene expression during infection *Int J Mol Sci* **21**:3730.

Evans SM, Rodino KG, Adcox HE, Carlyon JA (2018) ***Orientia tsutsugamushi* uses two**

Ank effectors to modulate NF-κB p65 nuclear transport and inhibit NF-κB

transcriptional activation *PLoS Pathog* **14**:e1007023.

Fol M, Włodarczyk M, Druszczyńska M (2020) **Host epigenetics in intracellular pathogen**

infections *Int J Mol Sci* **21**:4573.

Fu M, Liang S, Wu J, Hua Y, Chen H, Zhang Z, Liu J, Li X, Zhang B, Zhao W, Wan C (2021)

An *Escherichia coli* effector protein EspF may induce host DNA damage via

interaction with SMC1 *Front Microbiol* **12**:682064.

Gómez Borrego J, Torrent Burgas M (2024) **Structural assembly of the bacterial essential**

interactome *Elife* **13**:e94919.

Guarnaccia AD, Rose KL, Wang J, Zhao B, Popay TM, Wang CE, Guerrazzi K, Hill S,

Woodley CM, Hansen TJ (2021) **Impact of WIN site inhibitor on the WDR5 interactome**

Cell Rep 34:108636.

Hammond DE, Kumar JD, Raymond L, Simpson DM, Beynon RJ, Dockray GJ, Varro A

(2018) **Stable isotope dynamic labeling of secretomes (SIDLS) identifies authentic**

secretory proteins released by cancer and stromal cells *Mol Cell Proteomics* **17**:1837-

1849.

Hanford HE, Von Dwingelo J, Abu Kwaik Y (2021) **Bacterial nucleomodulins: a**

coevolutionary adaptation to the eukaryotic command center *PLoS Pathog*

17:e1009184.

Hsu PL, Li H, Lau HT, Leonen C, Dhall A, Ong SE, Chatterjee C, Zheng N (2018) **Crystal**

structure of the COMPASS H3K4 methyltransferase catalytic module *Cell* **174**:1106-

1116.e9.

Hyun K, Jeon J, Park K, Kim J (2017) **Writing, erasing and reading histone lysine**

methylations *Exp Mol Med* **49**:e324.

Jose L, Ramachandran R, Bhagavat R, Gomez RL, Chandran A, Raghunandanan S,

Omkumar RV, Chandra N, Mundayoor S, Kumar RA (2016) **Hypothetical protein Rv3423.1**

of *Mycobacterium tuberculosis* is a histone acetyltransferase *FEBS J* **283**:265-281.

Khan AA, Khan Z (2021) **Bacterial nucleomodulins and cancer: an unresolved enigma**

Transl Oncol **14**:100922.

Lebreton A, Job V, Ragon M, Le Monnier A, Dessen A, Cossart P, Bierne H (2014)

Structural basis for the inhibition of the chromatin repressor BAHD1 by the bacterial

nucleomodulin LntA *mBio* **5**:e00775-13.

Penn BH, Netter Z, Johnson JR, Von Dollen J, Jang GM, Johnson T, Ohol YM, Maher C, Bell

SL, Geiger K (2018) **An Mtb-human protein-protein interaction map identifies a switch**

between host antiviral and antibacterial responses *Mol Cell* **71**:637-648.e5.

Pisu D, Huang L, Grenier JK, Russell DG (2020) **Dual RNA-Seq of Mtb-Infected**

Macrophages In Vivo Reveals Ontologically Distinct Host-Pathogen Interactions *Cell*

Rep **30**:335-350.e4.

Pourpre R, Lakisic G, Desgranges E, Cossart P, Pagliuso A, Bierne H (2022) **A bacterial**

virulence factor interacts with the splicing factor RBM5 and stimulates formation of

nuclear RBM5 granules *Sci Rep* **12**:21961.

Prokop A, Gouin E, Villiers V, Nahori MA, Vincentelli R, Duval M, Cossart P, Dussurget O

(2017) **OrfX, a nucleomodulin required for *Listeria monocytogenes* virulence** *mBio*

8:e01550-17.

Qiang L, Zhang Y, Lei Z, Lu Z, Tan S, Ge P, Chai Q, Zhao M, Zhang X, Li B (2023) **A**

mycobacterial effector promotes ferroptosis-dependent pathogenicity and

dissemination *Nat Commun* **14**:1430.

Radoshevich L, Cossart P (2018) *Listeria monocytogenes***: towards a complete picture**

of its physiology and pathogenesis *Nat Rev Microbiol* **16**:32-46.

Rennoll-Bankert KE, Garcia-Garcia JC, Sinclair SH, Dumler JS (2015) **Chromatin-bound**

bacterial effector ankyrin A recruits histone deacetylase 1 and modifies host gene

expression *Cell Microbiol* **17**:1640-1652.

Rolando M, Sanulli S, Rusniok C, Gomez-Valero L, Bertholet C, Sahr T, Margueron R,

Buchrieser C (2013) *Legionella pneumophila* **effector RomA uniquely modifies host**

chromatin to repress gene expression and promote intracellular bacterial replication

Cell Host Microbe **13**:395-405.

Rolando M, Wah Chung IY, Xu C, Gomez-Valero L, England P, Cygler M, Buchrieser C

(2023) **The SET and ankyrin domains of the secreted** *Legionella pneumophila* **histone**

methyltransferase work together to modify host chromatin *mBio* **14**:e0165523.

Schator D, Mondino S, Berthelet J, Di Silvestre C, Ben Assaya M, Rusniok C, Rodrigues-

Lima F, Wehenkel A, Buchrieser C, Rolando M (2023) **Legionella para-effectors target**

chromatin and promote bacterial replication *Nat Commun* **14**:2154.

Sharma G, Upadhyay S, Srilalitha M, Nandicoori VK, Khosla S (2015) **The interaction of**

mycobacterial protein Rv2966c with host chromatin is mediated through non-CpG

methylation and histone H3/H4 binding *Nucleic Acids Res* **43**:3922-3937.

Singh PR, Dadireddy V, Udupa S, Kalladi SM, Shee S, Khosla S, Rajmani RS, Singh A,

Ramakumar S, Nagaraja V (2023) **The *Mycobacterium tuberculosis* methyltransferase**

Rv2067c manipulates host epigenetic programming to promote its own survival *Nat*

Commun **14**:8497.

Sun H, Kamanova J, Lara-Tejero M, Galán JE (2016) **A Family of Salmonella Type III**

Secretion Effector Proteins Selectively Targets the NF-κB Signaling Pathway to

Preserve Host Homeostasis *PLoS Pathog* **12**:e1005484.

Wang H, Fan Z, Shliaha PV, Miele M, Hendrickson RC, Jiang X, Helin K (2023) **H3K4me3**

regulates RNA polymerase II promoter-proximal pause-release *Nature* **615**:339-348.

Wang H, Helin K (2025) **Roles of H3K4 methylation in biology and disease** *Trends Cell*

Biol **35**:115-128.

Wang J, Ge P, Qiang L, Tian F, Zhao D, Chai Q, Zhu M, Zhou R, Meng G, Iwakura Y (2017)

The mycobacterial phosphatase PtpA regulates the expression of host genes and

promotes cell proliferation *Nat Commun* **8**:244.

Xue H, Yao T, Cao M, Zhu G, Li Y, Yuan G, Chen Y, Lei M, Huang J (2019) **Structural basis**

of nucleosome recognition and modification by MLL methyltransferases *Nature*

573:445-449.

Yang D, He X, Li S, Liu J, Stabenow J, Zalduondo L, White S, Kong Y (2019) **Rv1075c of**

***Mycobacterium tuberculosis* is a GDSL-Like esterase and is important for intracellular**

survival *J Infect Dis* **220**:677-686.

Yaseen I, Kaur P, Nandicoori VK, Khosla S (2015) **Mycobacteria modulate host epigenetic machinery by Rv1988 methylation of a non-tail arginine of histone H3** *Nat Commun* **6**:8922.

Yu H, Lesch BJ (2024) **Functional roles of H3K4 methylation in transcriptional regulation** *Mol Cell Biol* **44**:505-515.

Yu L, Fang F, Dai X, Xu H, Qi X, Fang M, Xu Y (2017) **MKL1 defines the H3K4Me3 landscape for NF-κB dependent inflammatory response** *Sci Rep* **7**:191.

Zhao J, Liu Y, Lin F, Wang W, Yang S, Ge Y, Chen PR (2019) **Bioorthogonal engineering of bacterial effectors for spatial-temporal modulation of cell signaling** *ACS Cent Sci* **5**:145-152.

Figures

Figure legends

Figure 1. Identification of conserved nucleomodulins in mycobacteria through

functional screening.

(**A**) Schematic representation of the bioinformatic pipeline for identifying conserved

nucleomodulins in Mycobacterium species. Genomic sequences from *M. tuberculosis*

H37Rv, *M. tuberculosis* H37Ra, *M. avium*, *M. marinum*, and *M. bovis* BCG were analyzed.

Signal peptides were predicted using SignalP 5.0 (D-score ≥ 0.5), non-classical secretion

signals were identified using SecretomeP 2.0 (NN score ≥ 0.9), and nuclear localization

signals (NLSs) were predicted using cNLS Mapper (score ≥ 2.0). (**B**) Classification of

conserved cellular nucleomodulins. Secreted proteins were categorized based on the

presence or absence of predicted NLS motifs. (**C**) Subcellular localization of EGFP-tagged

candidate nucleomodulins. (left) Confocal images of Rv0846c, MgdE, Rv2577, and Rv1988

fused to EGFP (green); nuclei are stained with DAPI (blue). (right) Quantification of nuclear

EGFP intensity. Rv1988 and pEGFP were used as positive and negative controls,

respectively. Scale bar, 10 μm. Data are presented as mean ± SD (n = 25 cells/group).

Figure 2. Hypothetical protein MgdE enters the host nucleus via dual NLS.

(**A**) Domain architecture of MgdE. Schematic representation of MgdE with the annotated

functional domains, including a Tat signal peptide (1–37 aa, twin-arginine translocation

motif), a GDSL-like lipase/acylhydrolase catalytic domain (74–258 aa), and two nuclear

localization signals (NLS1: 108–111 aa; NLS2: 300–305 aa). (**B**) Phylogenetic and structural

conservation of MgdE. (left) Neighbor-joining phylogenetic tree of MgdE homologs across

Mycobacterium species (1,000 bootstrap replicates; values ≥50% shown). (right) Clustal

Omega sequence alignment highlighting conserved residues (≥90% identity, red). Species

abbreviations: *M. tuberculosis* (Mtu), *M. bovis* BCG (Mbb), *M. tuberculosis* H37Ra (Mra), *M.*

africanum (Maf), *M. tuberculosis* CDC1551 (Mtc), *M. tuberculosis* GM041182 (Tbm), *M.*

shinjukuense (Msh), *M. marinum* (Mma), *M. lupini* (Mli), *M. avium* (Mav), *M. manitobense*

(Mman), *M. intracellulare* (Min). (**C**) Expression and nuclear localization of MgdE. (upper)

qRT-PCR analysis of mgdE mRNA in HEK293T cells over 48 h post-transfection. (lower)

Western blot analysis of nuclear fractions showing time-dependent nuclear accumulation of

MgdE-EGFP. Histone H3 and β-actin served as nuclear and cytoplasmic markers,

respectively. (**D**) Subcellular localization of EGFP-tagged wild-type and NLS-deleted MgdE.

(left) Schematic representation of EGFP-tagged constructs, including wild-type MgdE and its

mutants (MgdE$^{\Delta NLS1}$, MgdE$^{\Delta NLS2}$, and MgdE$^{\Delta NLS1-2}$). (right) Confocal microscopy images of

HEK293T cells transfected with the indicated constructs for 36 h. EGFP fluorescence

(green) and nuclear staining with DAPI (blue) were visualized using an FLUOVIEW software

(v5.0). Scale bar, 10 μm. Images were acquired with a ×100 oil immersion objective (NA =

1.4). (**E**) Nuclear EGFP intensity of wild-type and mutant constructs in (**D**). Data are shown

as mean ± SD (n = 15 cells). (**F**) Western blot analysis of nuclear and cytoplasmic fractions

from HEK293T cells transfected with wild-type and mutant MgdE-EGFP confirmed their

nuclear localization. MgdE-EGFP was detected using an anti-GFP antibody, and histone H3

and β-actin served as nuclear and cytoplasmic markers, respectively. All experiments were

performed in three biologically independent replicates. Statistical significance was assessed

by two-tailed unpaired Student's t-tests, with $*P < 0.05$ considered statistically significant.

Figure 3. Nuclear localization of MgdE facilitates mycobacterial intracellular survival

in macrophages.

(**A**-**B**) Intracellular survival of *M. bovis* BCG strains. THP-1 human macrophages (A) and

RAW264.7 murine macrophages (B) were infected (MOI = 10) with wild-type BCG (WT),

MgdE deletion mutant (ΔMgdE), complemented strain (Comp-MgdE), or NLS-deficient

complement (Comp-MgdE$^{\Delta NLS1-2}$). Bacterial survival was assessed by CFU enumeration at 2,

24, 48, and 72 h post-infection. (**C-H**) Cytokine expression in the infected THP-1 cells. qRT-

PCR analysis of cytokine mRNA levels in THP-1 cells infected with WT or ΔMgdE strains for

4–24 h. Inflammatory cytokine expression in infected THP-1 cells. qRT-PCR analysis of

intracellular cytokine mRNA levels in THP-1 cells infected with WT or ΔMgdE strains for 4–

12 h. Target genes include *il1b* (C), *il6* (D), *il10* (E), *csf1* (F), *csf2* (G), and *csf3* (H). Data are

presented as mean ±SD of three biologically independent experiments, analyzed using two-

tailed unpaired Student's *t*-tests (*$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$).

Figure 4. MgdE directly interacts with ASH2L and WDR5, core components of the host

COMPASS complex.

(**A**) Yeast cells were co-transformed with bait (pGBKT7) and prey (pGADT7) plasmids

expressing wild-type MgdE and human COMPASS components (ASH2L, WDR5, RbBP5,

and DPY30). Growth was monitored on non-selective (-Leu/-Trp, left) and selective (-Leu/-

Trp/-Ade/-His + 200 ng/µL aureobasidin A, right) media. Controls: CK− (pGBKT7-*lam* +

pGADT7-T, negative); CK+ (pGBKT7-*p53* + pGADT7-T, positive). (**B**-**C**) Cells were co-

transfected with Flag-MgdE and HA-tagged ASH2L, WDR5, or RbBP5 (1:1 molar ratio). At

36 h post-transfection, the lysates were immunoprecipitated using (**B**) anti-HA or (**C**) anti-

Flag antibodies, followed by immunoblotting with anti-HA, anti-Flag, and anti-GAPDH

(loading control). The input lanes represent 5% of the total lysate. Data are representative of

three independent experiments.

A

B

D

C

Figure 5. The conserved residues D^{224} and H^{247} mediate the binding ability of MgdE to

WDR5.

(**A**) Y2H assay identifying interactions between MgdE mutants and COMPASS complex

subunits. Yeast cells were co-transformed with bait (pGBKT7) and prey (pGADT7) plasmids

expressing wild-type or mutant MgdE and human COMPASS subunits (ASH2L, WDR5,

RbBP5, and DPY30). Growth was assessed on non-selective (-Leu/-Trp, left) and selective

(-Leu/-Trp/-Ade/-His + 200 ng/μL aureobasidin A, right) media. Controls: CK− (pGBKT7-*lam*

+ pGADT7-T, negative) and CK+ (pGBKT7-*p53* + pGADT7-T, positive). (**B**) Co-IP analysis of

MgdE mutants with WDR5. HEK293T cells were co-transfected with Flag-tagged MgdE

mutants and HA-tagged WDR5 (1:1 molar ratio). Complexes were immunoprecipitated using

anti-HA antibody and protein A/G beads, followed by immunoblotting with anti-Flag and anti-

HA antibodies. (**C**) Nuclear distribution of wild-type and mutants MgdE. Confocal microscopy

of HEK293T cells expressing wild-type or D^{224}A/H^{247}A MgdE-EGFP at 12 and 24 h post-

transfection (hpt). Nuclear foci were visualized by EGFP (green) and DAPI (blue) staining.

Scale bar, 10 μm. Images were acquired with a ×100 oil immersion objective (NA = 1.4). (**D**)

Immunoblot analysis of H3K4me3 levels. HEK293T cells expressing wild-type or D^{224}A/H^{247}A

mutant MgdE were analyzed for changes in H3K4me3 levels over 0–24 h post-transfection.

Histone H3 was used as a loading control. The data represent two independent biological

replicates.

Figure 6. MgdE suppresses COMPASS complex-mediated inflammatory responses by

inhibiting H3K4 methylation.

(**A**) Volcano plot of DEGs. DEGs between MgdE-deleted strain (Ko*mgdE*) and wild-type

BCG (WT) were visualized in a volcano plot. Genes with |log$_2$-fold change| ≥ 1 and P < 0.05

were considered significant. The *x*-axis represents log$_2$fold change, and the *y*-axis shows -

log$_{10}$(P-value). (**B**) GO enrichment analysis of DEGs. GO analysis revealed the significant

enrichment of immune and inflammatory processes in Ko*mgdE*-infected macrophages

compared to that in WT-infected cells, including positive regulation of response to external

stimulus (GO:0032103), response to molecule of bacterial origin (GO:0002237), response to

virus (GO:0009615), response to lipopolysaccharide (GO:0032496), leukocyte migration

(GO:0050900), viral process (GO:0016032), regulation of response to biotic stimulus

(GO:0002831), T cell activation (GO:0042110), and mononuclear cell differentiation

(GO:1903131). (**C**) KEGG pathway enrichment analysis of DEGs. Chord diagram of KEGG

pathway enrichment analysis showing signaling pathways that are significantly enriched in

Ko*mgdE*, including cytokine-cytokine receptor interaction and JAK-STAT signaling as the top

enriched pathways. (**D**) Heatmap of the inflammatory gene expression. Heatmap depicting

log$_2$-foldchange levels of inflammatory genes involved in the JAK-STAT and cytokine

signaling pathways. Upregulated and downregulated genes in Ko*mgdE* are shown in green

and red, respectively. The data were *Z*-score normalized. (**E-G**) qRT-PCR analysis of

cytokine mRNA levels in THP-1 cells infected with WT, ΔMgdE, MgdE-complemented

(Comp-MgdE), and NLS-deleted complement (Comp-MgdE$^{\Delta NLS1-2}$) at 24 h post-infection.

Cytokines analyzed include *il6* (**E**), *il10* (**F**), and *il12b* (**G**). Data are presented as mean ± SD

Figure 7. Nuclear localization of MgdE is essential for mycobacterial survival in mice.

(**A**) Bacterial burden in the lungs of the infected mice. C57BL/6 mice (n = 6/group) maintained under SPF conditions were intratracheally infected with 1.0×10^7 colony-forming units (CFU) of *M. bovis* BCG strains, including wild-type (WT), MgdE-deleted (ΔMgdE), MgdE-complemented (Comp-MgdE), and NLS-deleted complement (Comp-MgdE$^{\Delta NLS1-2}$). Lung bacterial loads were quantified using CFU assays at 0, 14, 28, and 56 d post-infection. (**B**) H&E-stained lung sections from infected mice (as in **A**) revealed granulomatous inflammation. Scale bars: 200 μm. (**C-D**) Pro-inflammatory cytokine expression in mice spleen. qRT-PCR analysis of cytokine mRNA levels of *il6* (**C**) and *il1β* (**D**) in spleen tissues from infected mice (n = 6/group) at 2 and 28 d post-infection. (**E**) Mechanistic model showing how mycobacterial nucleomodulin MgdE hijacks the COMPASS complex to suppress H3K4me3 and promote immune evasion. Upon *M. bovis* BCG infection, the nucleomodulin MgdE is delivered into the host nucleus via its nuclear localization signal (NLS) and directly binds to the COMPASS complex subunits, ASH2L or WDR5. This interaction disrupts H3K4 trimethylation (H3K4me3) deposition, leading to the epigenetic suppression of pro-inflammatory cytokine transcription (e.g., *il6*), thereby facilitating the intracellular survival of the pathogen. Data are presented as mean ± SD from six biologically independent experiments. Statistical significance was determined by a two-tailed unpaired Student's *t*-test (*P < 0.05, **P < 0.01, and ***P < 0.001).

Supplementary figures

Supplementary figure 1. Comparative analysis of classical and non-classical secreted

proteins in mycobacterial species.

(**A**) Venn diagram showing the distribution of predicted classical secreted proteins in four

Mycobacterium species: *M. tuberculosis* H37Rv (Mtu) and *M. tuberculosis* H37Ra (Mra), *M.*

bovis BCG (Mbb), *M. marinum* (Mmar), and *M. avium* (Mav). Proteins were predicted using

SignalP 5.0, with a signal peptide score (D-score) ≥ 0.5. A total of 125 proteins were

conserved across all species. (**B**) Venn diagram showing the distribution of predicted non-

classical secreted proteins across the same Mycobacterium species, predicted using

SecretomeP 2.0 with a neural network (NN) score ≥ 0.9. Ten proteins were conserved

across all species. Bar graphs (left: classical, right: non-classical) summarize the total

number of predicted secreted proteins per species. **(C)** Detection of secreted MgdE in

culture supernatants. C-terminally Flag-tagged Ag85B (positive control) and MgdE were

expressed in *M. bovis* BCG strains. Immunoblot analysis of bacterial lysates and culture

supernatants showing the expression and secretion of Flag-tagged Ag85B and MgdE.

Supplementary Figure 2. Subnuclear localization of MgdE-EGFP.

Confocal microscopy was used to assess the nuclear distribution of MgdE-EGFP at various

time points post-transfection. Nuclei were stained with DAPI (blue); MgdE-EGFP is shown in

green. Scale bar: 10 µm.

Supplementary Figure 3. Deletion of the nuclear localization signal of MgdE does not

affect the growth of *M. bovis* BCG strains.

(**A-B**) Construction and validation of the MgdE-deleted strain of *M. bovis* BCG. (**A**)

Schematic diagram of the homologous recombination strategy used to delete *mgdE* from the

M. bovis BCG genome. (**B**) Wild-type and mutant strains were used as templates to amplify

the *mgdE* gene (600 bp upstream–600 bp downstream) by PCR. Lanes 1 and 3: wild-type

genomic DNA; lanes 2 and 4: ΔmgdE genomic DNA. (**C**) Growth curve analysis of *M. bovis*

BCG strains. Growth of BCG strains, including wild-type BCG (WT), MgdE-deleted (ΔMgdE),

MgdE-complemented (Comp-MgdE), and NLS-deleted complemented (Comp-MgdE$^{\Delta NLS1-2}$)

strains, was measured in 7H9 medium.

A

Function	COMPASS complexes	Predicted binding confidence (pLDDT)
The catalytic subunit	SET1A/B	-
Required for H3K4me3	ASH2L	0.47
Required for assembly	RBBP5	0.30
Required for assembly	WDR5	0.77
	DPY30	0.62
Components of the Set1 complexes	CXXC1	0.51
	WDR82	-
Components of the Set1 complexes	HCF1	0.37

B

MgdE-ASH2L	MgdE-DPY30	MgdE-RBBP5	MgdE-WDR5

Supplementary Figure 4. MgdE interacts with COMPASS complex subunits.

(**A**) Predicted binding affinities between MgdE and COMPASS core subunits. The predicted

local distance difference test (pLDDT) scores calculated using AlphaFold 2.2.0 for the

interactions between MgdE and the COMPASS subunits were as follows: ASH2L (pLDDT =

0.47), RbBP5 (pLDDT = 0.30), WDR5 (pLDDT = 0.77), and DPY30 (pLDDT = 0.62).

Confidence levels are categorized as follows: High confidence: pLDDT ≥ 0.7 (strong

predicted binding); Medium confidence: 0.5 ≤ pLDDT < 0.7 (moderate binding); Low

confidence: pLDDT < 0.5 (weak predicted binding). (**B**) Structural modeling of MgdE-

COMPASS interactions. AlphaFold generated models of the simulated binding interfaces are

shown, with MgdE highlighted in red and COMPASS subunits (ASH2L, WDR5, RbBP5, and

DPY30) shown in gray.

Supplementary Figure 5. Conserved residue mutations do not affect the binding of

MgdE to ASH2L.

Co-IP analysis of MgdE mutants with ASH2L. HEK293T cells were co-transfected with Flag-

tagged MgdE mutants and HA-tagged ASH2L (1:1 molar ratio). Co-IP was performed using

anti-HA antibody-bound protein A/G beads, and immunoprecipitated complexes were

analyzed by immunoblotting with anti-Flag and anti-HA antibodies.

Supplementary Figure 6. MgdE suppresses cellular inflammatory responses during *M.*

bovis* BCG infection.

(**A**) KEGG pathway enrichment analysis. KEGG pathway analysis identified significantly

enriched pathways in THP-1 cells infected with the Ko*mgdE* strain compared to those

infected with wild-type BCG (WT). DEGs were predominantly associated with immune and

signaling pathways, including thyroid hormone synthesis (hsa04918), AMPK signaling

pathway (hsa04152), PPAR signaling pathway (hsa03320), apelin signaling pathway

(hsa04371), cytokine-cytokine receptor interaction (hsa04060), JAK-STAT signaling pathway

(hsa04630), and hematopoietic cell lineage (hsa04640). (**B**) Elevated inflammatory gene

expression in Ko*mgdE*-infected cells. Key upregulated inflammatory genes in Ko*mgdE*- *vs*.

WT-infected cells are highlighted. (**C**) Functional enrichment analysis of upregulated genes.

Analysis using STRING and Cytoscape revealed regulation of inflammatory responses as a

top enriched biological process (*P* < 0.05).

Supplementary Figure 7. MgdE modulates inflammatory responses and bacterial

colonization in the spleens of infected mice.

Bacterial colonization in the spleen. CFU assays were performed on splenic homogenates

from mice infected as described in **Figure 7A**. Bacterial loads (CFU/g tissue) were quantified

at 2, 14, 21, 28, and 56 d post-infection. Data are presented as mean ± SD (n = 3).

Statistical significance was determined using a two-tailed unpaired Student's *t*-test (*P* < 0.05,

**$P < 0.01$, and ***$P < 0.001$).

950