Targeting Semaphorin 7a signaling in preclinical models of estrogen receptor-positive breast cancer

Rachel N Steinmetz[1,2], Veronica Wessells[1], Heather Fairchild[1], Traci R Lyons[1,2,3]

[1]Division of Medical Oncology, Department of Medicine, School of Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.
[2]Young Women's Breast Cancer Translational Program, University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA.
[3]University of Colorado Cancer Center, Aurora, CO, 80045, USA.

Running title: Targeting SEMA7A signaling in preclinical models of ER+ breast cancer

Corresponding author:
Name: Traci R. Lyons
Address: U. Colorado Anschutz, 12801 E 17th Ave, L18-8103 MS 8117, Aurora, CO 80045
Email: traci.lyons@cuanschutz.edu

Conflict of interest disclosures:
Rachel N Steinmetz – no conflict of interest to disclose.
Veronica Wessells – no conflict of interest to disclose.
Heather Fairchild – no conflict of interest to disclose.
Traci R Lyons – Cofounder and Chief Scientific Officer of Pearl Scientific.

TRANSLATIONAL RELEVANCE

Our studies propose inhibition of a novel target in estrogen receptor-positive (ER+) breast cancer (BC) with an anti-Semaphorin 7a (SEMA7A) monoclonal antibody (SmAbH1). We show efficacy of SmAbH1 as a single agent and in combination with endocrine therapy in preclinical models. We also show that targeting of SEMA7A signaling, including the use of well-known and novel PI3K inhibitors, could be an effective treatment strategy for patients with SEMA7A+ BC, particularly in combination with standard of care. Our mechanistic studies detail the cellular signaling pathway activated by SEMA7A and support the further investigation of these novel drug combinations for clinical use. Further, SEMA7A is a biomarker for poor prognosis and decreased patient survival, posing the need for a clinical diagnostic test for SEMA7A levels in BC patients, which we are currently developing.

ABSTRACT

Estrogen receptor-positive (ER+) breast cancer (BC) comprises over 70% of breast cancers and is the leading cause of BC-related deaths in women worldwide. Despite available therapies targeting ER in BC, recurrence occurs in many patients due to therapeutic resistance. Semaphorin 7a (SEMA7A) is a biomarker associated with poor prognosis and endocrine therapy resistance for BC patients. Survival analyses of ER+ BC patients on endocrine therapy confirm early recurrence in patients with SEMA7A+ tumors. Thus, we aim to establish novel treatment strategies to improve outcomes for patients with ER+ SEMA7A+ BC. In this paper we investigate the mechanisms by which SEMA7A promotes resistance to endocrine therapy and its potential as a therapeutic target for ER+ BC. Our studies suggest that SEMA7A binds to integrins β1 and β4 and activates AKT mediated pro-survival signaling via its RGD domain. Using syngeneic ER+ models, tumors in were treated with PI3K inhibitors (20mg/kg alpelisib; 10mg/kg GCT-007), alone or in combination with tamoxifen (0.5 mg/100uL peanut oil), which resulted in decreased tumor growth. Tumors were also treated with a combination of an anti-SEMA7A antibody (SmAbH1) (100-250mg/kg) and fulvestrant (83mg/kg), compared to single agents. Our results demonstrate that direct inhibition of SEMA7A via SmAbH1 significantly reduces tumor growth of SEMA7A+ tumors, and the combination with fulvestrant may be even more effective. Our studies suggest that patients with ER+SEMA7A+ tumors should be candidates for PI3K-targeted therapies or anti-SEMA7A-based therapy.

INTRODUCTION

The most prevalent type of breast cancer (BC) is estrogen receptor-positive (ER+), accounting for 70% of all BC diagnoses (1,2). The standard treatment for women with ER+ BC typically involves endocrine therapy, such as the selective ER degrader, fulvestrant, the selective ER modulator, tamoxifen, and aromatase inhibitors such as exemestane and letrozole (2,3). While endocrine therapy is effective in many patients, 22-52% of patients with ER+ BC will experience disease recurrence, depending on lymph node involvement (2,3). Thus, there is an urgent need to improve treatments for patients with ER+ BC. We and others have identified that Semaphorin 7a (SEMA7A), a neuro-endocrine signaling protein (4), is dysregulated in various diseases and cancer types, resulting in poor prognosis (5-13). While numerous studies have investigated the role of SEMA7A in triple-negative breast cancer (TNBC), which lack expression of ER (ER-), progesterone receptor (PR-), and amplification of HER2 (HER2-) (14-19), we have shown that ER can regulate expression of SEMA7A. As such, and have previously analyzed the importance of SEMA7A expression in ER+ BC to show that SEMA7A is expressed in ER+ BC where expression confers resistance to ER targeting therapies including fulvestrant and tamoxifen (2). Our previous studies have also demonstrated mechanisms of therapy resistance activated in ER+ SEMA7A+ tumors that drive reduced sensitivity to the endocrine therapy fulvestrant, including downregulation of ER, activation of pro-survival signaling through Bcl-2, and regulation of markers of drug-resistant stem cells (CD44+/CD24-) (2). We have also shown that long-term estrogen deprivation, such as that induced by endocrine therapies, results in upregulation of SEMA7A and suggest pro-survival signaling as a potential mechanism of resistance(2). In support of this, other studies have shown that pro-survival signaling, such as that mediated by phosphoinositide linked kinase-3 (PI3K), AKT, and other protein kinases, leads to mechanisms of resistance to endocrine therapies (20,21); the contribution of PI3K/AKT to SEMA7A induced phenotypes is still unknown. Therefore, understanding how SEMA7A+ ER+ BCs upregulate pro-survival signaling, and whether these mechanisms could be targetable in SEMA7A+ tumors, will provide insight into treatment strategies for this subset of patients.

In previous results, we observed that *SEMA7A* mRNA expression is elevated in 67% of luminal

breast cancer cell lines, compared to 50% and 36% in basal A and basal B, respectively (2) and that

increased expression results in poor outcomes in all types of breast cancer(2,16). In its immune-related

and other roles, SEMA7A is known to bind to two receptors: Plexin C1 and integrin β1 (22), and in

models of TNBC we have shown that the pro-tumor roles for SEMA7A are dependent on integrin β1

and integrin α6 (14,16). However, in the context of ER+ BC, whether SEMA7A binds integrin receptors

remains unknown. Known downstream targets of integrins are those involved in cell adhesion and cell

survival, including the PI3K/AKT pathway. In support of a role for SEMA7A- and integrin-mediated

activation of the PI3K pathway, studies in TNBC show activation of this pro-survival pathway (15). We

have also shown that ER+SEMA7A+ tumors are sensitive to a non-selective PI3K inhibitor (LY294002)

that is not utilized clinically. Clinical trials have also shown efficacy of alpelisib or alpelisib plus

fulvestrant in PIK3CA-mutated ER+ breast cancer (23,24). Therefore, analysis of clinically relevant

PI3K targeted therapies could provide insight into treatment options for patients with SEMA7A+ BC

(2,25).

Constitutively active PI3K signaling can lead to worse patient prognosis and multidrug

resistance, including fulvestrant resistance (20,26,27). We therefore sought to investigate the hypothesis

that PI3K inhibitors could be effective for ER+SEMA7A+ BC. In our results presented herein, we show

that breast cancer progression is increased in women with ER+SEMA7A+BC, resulting in poor

prognosis and highlighting the need to identify additional treatment options. Mechanistically, we show

for the first time that SEMA7A binds to integrin β1 and β4, in ER+ breast cancer cells and activates

PI3K signaling as well as other phospho-signaling pathways related to cell survival in an integrin-

dependent manner. Integrins can activate AKT directly through Integrin-linked kinase (ILK)

phosphorylation of AKT at serine 473 (28) and we have previously published that inhibitors of PI3K,

ILK, and/or integrins can reduce SEMA7A-mediated activation of pAKT (29); however, integrin and

ILK inhibitors are not clinically available, so we explored the effect of PI3K inhibitors in ER+

SEMA7A+ tumors. We reveal a partial reversal of endocrine therapy resistance in ER+SEMA7A+

models of breast cancer by PI3K inhibition. Further, we tested a novel anti-SEMA7A monoclonal

antibody for specifically reducing tumor growth in mice harboring SEMA7A+ tumors, both as a single

agent and combined with fulvestrant. Collectively, our results support targeting of SEMA7A, both

directly and indirectly through PI3K inhibition, as novel therapeutic strategies to improve patient

outcomes in ER+SEMA7A+BC.

MATERIALS & METHODS

<u>**Cell Culture:**</u>

MCF7 cells were obtained from K. Horwitz (University of Colorado Anschutz Medical Campus). ZR75-

1 cells were obtained from the University of Colorado Cancer Center Cell Technologies Shared

Resource (CTSR). TC11 cells were obtained from L. Schuler (University of Wisconsin School of

Veterinary Medicine). SSM2 cells were obtained from A. Borowsky (University of California Davis).

Cell lines were validated by fingerprinting analysis at the University of Colorado Anschutz/Barbara

Davis Center Molecular Biology Core. Cells were routinely tested for mycoplasma contamination

throughout the studies described herein with Lonza MycLonza MycoAlert PLUS (Walkersville, MD);

last tested 3-24-2025. Cells were cultured at 37°C and 5% CO2. MCF7 were grown in MEM with 5%

heat-inactivated fetal bovine serum (FBS; Peak Serum PS-FB3, Colorado USA) and 1%

penicillin/streptomycin (P/S; ThermoFisher Scientific). TC11 and ZR75-1 were grown in RMPI 1640

with 10% heat-inactivated FBS and 1% P/S. SSM2 were grown in DMEM/F12 and supplemented with

10% FBS, 1% L-glutamine, 1% P/S. 50 μM 2-mercaptoethanol, 0.3 μM hydrocortisone, 5 μg/ml insulin

and 10 ng/ml transferrin. Experiments were performed with cells below passage 30. For exogenous

SEMA7A treatments, recombinant SEMA7A (rSEMA7A) was purified from SEMA7A-Fc expressing

MDA-MB-231 cells in collaboration with the University of Colorado Cancer Center CTSR (14,30).

<u>**Transductions:**</u>

MCF7-SEMA7A Overexpressing (OE) and MCF7 empty vector (EV) control cell populations were generated by lentiviral transductions with pLX304-Blast-V5 vectors; CMV-driven, V5-tagged full-length Semaphorin 7a cDNA (666 aa) or empty vector, respectively, and selected for using blasticidin at 5 µg/mL (Dana Farber Cancer Institute & CU Anschutz Functional Genomics Core). TC11-control and TC11-SEMA7A knockdown cell lines were generated via lentiviral transduction with pLKO-shNonTargeting-puro, and pLKO-shSEMA7A-038 (KD#1) and pLKO-shSEMA7A-040 (KD#2), respectively, and selected for using puromycin at 2 µg/mL. SEMA7A expression was validated via western blot analysis for all transduced cell lines.

Cell Viability Assays:

Cells were plated in a 96-well (1,000 cells/well), 24-well (100,000 cells/well), 12-well (200,000 cells/well) or 6-well (500,000 cells/well) plate in triplicate and allowed to adhere at 37°C for 24 hours. After 24 hours, the cells were treated with vehicle or drug as follows: Fulvestrant and 4-hydroxytamoxifen were purchased from Millipore Sigma (Cat#129453-61-8 and Cat#68392-35-8, St. Louis, MO). For *in vitro* assays, fulvestrant and 4-hydroxytamoxifen treatments were at a final concentration of 8 nM and 5 µM in DMSO, respectively. Alpelisib (BYL719) was purchased from SelleckChem (Cat# S2814) and used at 0.5-1 µM in DMSO, as determined by our dose response and the literature (31). GCT-007 was obtained via collaboration with Global Cancer Technology (San Diego, CA) and used at 0.1-0.5 µM in DMSO as determined by our dose response. Anti-Semaphorin 7a monoclonal antibody; clone H1 (SmAbH1) was obtained from the University of Colorado Cancer Center CTSR. SmAbH1 or Mouse IgG1 (Bio X Cell; Cat# BP0083) was used at 1 µM in PBS as determined by our dose response and the literature (29,32,33). Percent confluence was determined via formalin fixation and 0.1% crystal violet staining and analyzed via ImageJ analysis. Data are normalized to control condition for each treatment. Experiments were performed in biological and technical triplicates and statistics were measured via repeated-measures two-way ANOVA with multiple comparisons.

Drug Titration Studies:

5,000 SEMA7A overexpressing or empty vector control MCF7 cells, or 10,000 TC11 shCtrl or

shSEMA7A cells, were seeded per well in a 96-well plate and allowed to adhere overnight. Vehicle or

increasing concentrations of drug were added. Growth was assessed using 0.1% crystal violet staining

after 24 hours of treatment. To calculate IC50 values, cell viability data for a range of concentrations

(minimum 6-points data curve) for each drug was plotted as a dose-response curve in GraphPad Prism

and analyzed using nonlinear regression three-parameters curve fit model of log(inhibitor) vs. response.

Experiments were performed in triplicate and two-tailed, unpaired t-test was performed on each IC50

value in triplicate experiments.

Co-immunoprecipitation:

Pierce Classic Magnetic IP/Co-IP Kit (ThermoFisher; Cat# 88804) and User Guide/CoIP Protocol

(Pierce Biotechnologies) was used to perform all CoIP experiments. Cells were cultured normally to

optimal confluence in T75 flasks. For RGDS experiments, RGDS peptide (Tocris Bioscience; Cat#3498,

50uM in milliQ H_2O) or control (milliQ H_2O) was added to culture media for 1 hour prior to making cell

lysates. Lysates were made using IP lysis/wash buffer and incubated on ice for 15 min followed by

centrifugation. Protein concentration of lysates was quantified using Qubit Protein Broad Range Assay

Kit (ThermoFisher; Cat# A50668). Per sample, 1.0 mg of lysate was pre-incubated with 30 uL IP

antibody: mouse IgG or SEMA7A (Santa Cruz; Sc-374432) and rocked overnight at 4°C. Per sample,

25μl of Pierce Protein A/G magnetic beads were washed and incubated with the antibody-lysate solution

and rocked for 1 hour at room temperature. Using a magnet, supernatant was removed, and the beads

were washed with lysis buffer three times. Beads were then eluted with Pierce Elution Buffer, and 10uL

(pH) Neutralization Buffer was added per sample. Non-reducing Lane Marker Sample Buffer (5X,

Pierce) was used as loading buffer for immunoblots following CoIP elution.

Immunoblotting:

Protein extracts were prepared from cell lines by lysing cells in RIPA with EDTA (Thermo Fisher; Cat#

J61529.AP) containing PhosSTOP phosphatase inhibitor (Roche; Cat# 4906837001), and complete

protease inhibitor (Roche; Cat# 11697498001). Protein concentration of lysates was determined using

Qubit Broad Range Protein Assay Kit (ThermoFisher; Cat# A50668). 30μg of total protein in Laemmli

protein sample buffer was heated for 10 minutes. Lysates were run on SDS-Page gels and transferred

onto methanol-activated PVDF membranes using the Trans-Blot Turbo Transfer System (Bio-Rad).

Membranes were blocked with 5% bovine serum albumin and probed with the following primary

antibodies according to manufacturer recommendation: SEMA7A (Santa Cruz; Cat#Sc-374432), pAKT

S473 (Cell Signaling; Cat# 4060), total AKT (Cell Signaling; Cat#2920), Integrin β4 (Cell Signaling;

Cat#14803), Integrin β1 (Cell Signaling; Cat#34971), ERα (AbCam; Cat#16660), α-tubulin (Cell

Signaling; Cat +#9099S) and GAPDH (Biolegend; Cat# 607902). Membranes were washed with 1X

Tris-buffered saline with Tween 20, incubated with a goat anti-rabbit HRP secondary antibody (Abcam;

Cat# 6721), and developed with enhanced luminescence detection system (Thermo Fisher; Cat# 32106)

on the Licor Odyssey. Densitometric O.D. values of three biological replicates were obtained using

ImageJ(34) and normalized to the loading control GAPDH or α-tubulin, and normalized to the first

sample.

Animal Experiments:

Power calculations based on pilot studies were utilized to determine the number of animals per group.

Tumors were measured using digital calipers every 2–3 days with the observer blinded to treatment

group, and tumor volume was calculated as (L × W × W × 0.5). Tumors were harvested at designated

study endpoints (Maximum tumor volume = $1000mm^3$). Mammary glands with intact tumor, axial

lymph nodes and lungs were harvested, formalin fixed, and paraffin embedded for

immunohistochemical analysis. All animal studies were performed under the approval of the UC

Anschutz Medical Campus Institutional Animal Care and Use Committee. Beeswax estrogen (E2) and

sham pellets were generated and administered based on previous literature (35). For the study that

combined fulvestrant and rSmAbH1-P, tumors were harvested and detached from the mammary gland;

photographs were taken next to ruler and scaled to 1cm scale bar.

MCF7 Model: MCF7 is a well-known human ER+ cell line used to study breast cancer, thus it was an

appropriate model to initiate our studies. 2,500,000 SEMA7A overexpressing or control MCF7 cells

were injected into the #4 mammary fat pads of 6–8-week-old female NCG mice (Cat# CRL572, Charles

River, Wilmington, MA). For drug treatment studies, fulvestrant dosing began when average tumor

volume reached ~100mm^3. At day 37, animals were randomized to treatment into size-matched groups

to receive either vehicle or PI3K inhibitor (alpelisib or GCT-007) daily.

TC11 Model: TC11 is a syngeneic ER+ murine model which we chose to use because it provides the

context of an intact immune system, and there are limited other ER+ mouse models. TC11 also

expresses endogenous SEMA7A, making it a relevant model for our research. 200,000 TC11 cells (wild

type, shCtrl or shSEMA7A #2) were injected into the bilateral #4 mammary fat pads of 6–8-week-old

female FVB/N mice (FVB/NCrl, Cat# CRL207, Charles River, Wilmington, MA). Mice were

randomized to treatment into size-matched groups at average tumor volume ~50-100mm^3. For the

estrogen/sham pilot study, mice were randomized to groups receiving E2 or sham pellets two days prior

to tumor injections.

SSM2 Model: SSM2 is another syngeneic ER+ murine model which we chose to use because it provides

the context of an intact immune system, and there are limited other ER+ mouse models. SSM2 also

expresses endogenous SEMA7A, making it a relevant model for our research. 1,000,000 SSM2 tumor

cells were injected into the #4 mammary fat pads of 6–8-week-old female 129 Sv-E mice (129 Sv-E,

Cat# CRL287, Charles River, Wilmington, MA). Mice were randomized to treatment into size-matched

groups at average tumor volume ~50mm^3.

Treatments: Fulvestrant was administered as follows: 2.5 mg in 90% sesame oil and 10% ethanol,

injected subcutaneously every 5 days. This dose is equivalent to approximately 83 mg/kg per mouse.

Using the FDA Dosing Guidance for Industry: Conversion of Animal Doses to Human Equivalent

Doses, equivalent to 6.7mg/kg in humans (2). Tamoxifen was given using the following dosing regimen:

0.5 mg/100uL peanut oil per mouse, via intraperitoneal (IP) injection every 3rd day (36). Alpelisib was

dissolved in 1% Carboxymethyl cellulose solution + 0.5% Tween 80 and administered via oral gavage at

20 mg/kg daily. GCT-007 was dissolved in 55% 1M NaCl, 30% PEG-400, 7.5% Solutol-HS + 7.5% N-

Methyl-2-pyrrolidone and administered via oral gavage at 10 mg/kg daily. SmAbH1/rSmAbH1 or

Mouse IgG1 (Bio X Cell; Cat# BP0083) was given in sterile PBS at 100-250 µg per 100µL PBS every

other day via IP injection (32,33).

Immunohistochemistry:

Tissue collected from the *in vivo* study endpoints as described above, was formalin-fixed and paraffin-

embedded. Tissue was stained with Hematoxylin and Eosin. Four-µm sections were pretreated with 10X

Dako Target Retrieval System. Slides were prepared in a Dako Autostainer with primary antibody

(Integrin β4: Cell Signaling Cat#14803, SEMA7A AbCam Cat # ab255602, 9EG7/CD29: BD

Pharmingen rat-anti-mouse CD29 clone 9EG7 Cat# 550531), followed by EnVision+ HRP Rabbit

(Agilent Technologies cat# K4003) antibody for 30 mins. For 9EG7 staining, rat-on-mouse HRP

polymer detection kit (Biocare cat# RT517H) was used for secondary. The tissue was incubated with

chromogen 3,3'diaminobenzidine (Vector Labs catalog #SK-4105) for 10 minutes, and then the tissue

was counter-stained with hematoxylin for 1 minute (Vector Labs catalog #H-3401-500). Stained images

were acquired using an Olympus Fluorescent Microscope IX83. Quantification was performed using

Cell Sense Analysis.

ELISA:

ELISA Buffer Kit (Thermo Cat #CNB0011) was used to perform an indirect ELISA where the plate was

coated with SEMA7A protein and probed with serial dilution concentrations of rSmAbH1-P to

determine binding specificity. A 96-well plate was pre-coated with 5ug/well of purified SEMA7A

(50ug/mL) protein in Coating Buffer A and incubated at 4 degrees overnight. Plate was washed with 1X

wash buffer and then probed with primary antibody (rSmAbH1-P) at serial dilution concentrations, from

20 to 0 ug/mL, and incubated at 4 degrees overnight. Plate was washed with 1X wash buffer three times

and then probed with secondary antibody (Goat anti-mouse IgG-HRP; Abcam Cat#ab97040) for 30

minutes at room temperature. Plate was washed three times with 1X wash buffer and absorbance at

425nm was read on plate reader.

Data Mining Analysis:

Kaplan Meir Plotter(37) was used to assess overall survival (OS), distant metastasis-free survival

(DMFS), and post-progression survival (PPS) data for BC patients in the ER+ dataset, and patients in

the ER+ dataset who received endocrine therapy (tamoxifen or fulvestrant). SEMA7A was queried and

stratified into high and low-expression groups using the auto-select best cutoff level function. The

generated data, plots, hazard ratio (HR), and LogRank P value were downloaded from KMplotter.com.

TCGA Human Data Analysis: We analyzed The Cancer Genome Atlas (TCGA) Breast Cancer study for

the mRNA expression of Semaphorin 7a (*SEMA7A),* integrin β1 *(ITGB1), and* integrin β4 *(ITGB4)*

collected from patient samples processed by RNAseq – IlluminaHiSeq. Gene expression analysis was

performed using the Xena Functional Genomics Explorer(38). The log2(norm_count+1) value for

mRNA expression was used to graph correlations between SEMA7A and the other genes in the TCGA

Breast Cancer dataset. Each data was analyzed using Pearson's correlation (rho) and p-values are

indicated on each plot.

BioSpa Assay:

219 500 MCF7 cells per well were plated in 96 well plates and incubated overnight to allow for adherence.

220 Cells were then treated with 1μM IgG1 or 1μM SmAbH1 and immediately placed in a BioSpa Live Cell

221 Imager (Agilent) for 4 days, with images of each well being taken every 4 h, at ×10 magnification. The

222 cell count of each well was measured via brightfield count in the BioSpa Software.

Reverse Phase Protein Array (RPPA):

For cell lysates (RPPA-1), MCF7 Empty Vector control or SEMA7A Overexpressing cells were

cultured, each in quadruplicate, in a 10cm dish for 72 h. After washing cells with PBS, lysates were

generated using RPPA Lysis Buffer (Baylor College of Medicine Antibody-Based Proteomics Core) and

frozen at -20°C. For tumor RPPA (RPPA-2), tumors were harvested and flash-frozen in liquid nitrogen

at necropsy and stored at -80°C. Lysates/tumors were shipped on dry ice to the Baylor College of

Medicine (BCM) Antibody-Based Proteomics Core (Houston, TX). Subsequent RPPA was performed

by the core facility as outlined here (https://www.bcm.edu/academic-centers/dan-l-duncan-

comprehensive-cancer-center/research/cancer-center-shared-resources/antibody-based-

proteomics/protocols-workflow) (39). Normalized fold change for each antibody was calculated by

(each signal value/AVG value of controls) and is depicted as log2(normalized fold change. Significance

was determined by performing an unpaired two-tailed t-test between the fold change values for each

antibody. Each sample was run in technical triplicates and n=4 samples per group. Samples with a

Coefficient of Variation >25% for technical triplicates were excluded from further analysis. Pathway

analysis was done using Gene Ontology Pathway Analysis.

Protein Ribbon Structures:

The protein structure of SEMA7A was analyzed in PyMOL with an academic research license. Ribbon

structure (purple) was created and modified to highlight the RGD motif (orange) and the antibody-

binding site (green) in PyMOL.

Synergy Scores:

Bliss Synergy scores for two-drug combination treatments were calculated using Synergyfinder.org;

scores $\leq$ -10 are considered antagonistic, between -10 and 10 are considered additive, and a score $\geq$ 10 is

considered synergistic (40). Further details on the methods used can be found at

https://synergyfinder.fimm.fi/synergy/synfin_docs/.

Novel drug treatments:

SmAbH1 (hybridoma): Mice were injected with SEMA7A protein and KLH peptide to enhance

immune response. B cells from mouse spleens were fused with myeloma cells to create a hybridoma;

positive hybridoma clones were selected and monoclonal antibodies were harvested. SmAbH1 Antibody

binds to amino acids 381-392 of SEMA7A (green; SFigure 1G) (29).

rSmAbH1 (recombinant): The ExpiCHO-S expression system (Gibco) was used for the production of

recombinant antibodies according to the manufacturer's protocol. After the 12 days of expression, the

culture media were harvested by centrifugation, filtered through a 0.2-micron filter, and underwent

affinity chromatography using MabSelect PrismA resin and the AKTA pure FPLC instrument for

purification of recombinant monoclonal antibody.

GCT-007: GCT-007 is a pan PI3K inhibitor and was provided to us for these studies by Global Cancer

Technology. (16776 Bernardo Center Drive #203, San Diego, CA 92128).

Statistical Analysis:

All quantitative and statistical analysis was performed using GraphPad Prism (Version 10.3.1), and the

methods of tests are labeled in the figure legends. The statistical tests used in this paper include unpaired

two-tailed t-test, Pearson's correlation, and ordinary one-way ANOVA with Tukey's multiple

comparisons test. All p-values were measured under two-tailed conditions. Significance was determined

by a p-value less than 0.05. $*p<0.05$, $**p<0.01$, $***p<0.001$ $****p<0.0001$.

Data Availability:

The data generated in this study are available upon request from the corresponding author.

RESULTS

SEMA7A is expressed in ER+ breast cancer and associated with decreased 5-year patient survival.

To test our hypothesis that SEMA7A predicts decreased response to therapy, and therefore poor

prognosis, we used Kaplan-Meier Plotter(37) to examine 5-year survival for ER+ BC patients with high

versus low tumor *SEMA7A* mRNA expression (Figure 1A-C). The probability of overall survival (OS)

was significantly decreased for patients with high *SEMA7A* expression (Figure 1A), as well as distant

metastasis-free survival (DMFS) (Figure 1B) and post-progression survival (PPS) (Figure 1C). Further,

relapse-free survival (RFS) (Figure 1D) and DMFS (Figure 1E) were decreased in ER+ BC patients with

high *SEMA7A* expression who received endocrine therapy.

SEMA7A activates AKT signaling and binds to Integrins β1 and β4 via RGD in ER+ tumor cells.

To investigate differential protein and phospho-protein expression in ER+ MCF7 breast cancer

cells transduced with an empty vector (EV) or SEMA7A overexpression (OE) vector, we utilized a

reverse phase protein array (RPPA-1) (41). We confirmed SEMA7A overexpression via western blot

prior to submission of lysates for RPPA (SFigure 1A). Using the data obtained from RPPA-1 we

performed Gene Ontology (GO) Analysis to identify upregulated pathways in the SEMA7A OE lysates

compared to EV; we observed that signaling related to cell survival, including AKT, mTOR, and

integrin signaling, was upregulated in SEMA7A OE cells (Figure 2A). Specifically, the RPPA data

suggest that overexpression of SEMA7A leads to increased integrins β1 and β4 as well as phospho-AKT

(pAKT) at serine 473 (S473) (Figure 2B). The RPPA also show that integrin-linked kinase, which can

activate AKT through S473 phosphorylation (42), the anti-apoptotic protein Bcl-2, and downstream

targets of AKT, such as p-P70S6K (S371), were all increased in SEMA7A OE lysates (Figure 2B).

Phospho-mTOR(S2448), which is also a downstream target of AKT, was also increased, but not with

statistical significance (p=0.0564)(Figure 2B). To validate that SEMA7A activates AKT signaling

through pAKT (S473) in ER+ breast cancer cells, we validated pAKT (S473) and total AKT using

immunoblot analysis on MCF7 EV and SEMA7A OE cells, or WT MCF7 treated with purified

SEMA7A (rSEMA7A); the pAKT to AKT ratio was significantly higher in SEMA7A OE cells

compared to EV and rSEMA7A-treated cells compared to controls (SFigure 1B-C). Expression of

ITGB1 and *ITGB4* mRNA also weakly correlated with *SEMA7A* expression in the TCGA Breast Cancer

Dataset (n=1247) (SFigure 1D-E). To investigate integrin β1 and β4 as binding partners for SEMA7A in

ER+ breast cancer cells, we performed co-immunoprecipitation (CoIP) using a SEMA7A specific

antibody to bind the protein and its binding partner(s). After CoIP, western blotting revealed binding of

both integrin β1 and β4 to SEMA7A, which was higher in SEMA7A OE cells compared to controls

(Figure 2C, SFigure 1G: CoIP input and unbound controls). To investigate whether SEMA7A binds to

integrins via its RGD (Arg-Gly-Asp) motif (SFigure 1F), which is known to facilitate integrin binding

(43), we conducted CoIP using MCF7 SEMA7A OE lysates treated with PBS or RGDS (Arg-Gly-Asp-

Ser) peptide, which inhibits RGD-integrin binding. Our results indicate that binding of SEMA7A to

integrin β1 and β4 was significantly reduced by the RGDS peptide (Figure 2D, SFigure 1H: CoIP input

and unbound contorls). These results were validated in an additional human ER+ breast cancer cell line,

ZR75-1 (SFigure 1I).

SEMA7A-mediated activation of AKT is dependent on RGD binding in ER+ tumor cells.

Based on our results showing increased pAKT (S473) in SEMA7A OE MCF7, we validated our

results in a syngeneic ER+ cell line. We obtained TC11 and SSM2 cells, which are ER+ mouse

mammary tumor cell lines (44,45)(generous gifts from Linda Schuler and Alexander Borowsky), and

validated ERα expression compared to MCF7 (SFigure 2A). We also compared SEMA7A expression in

the two cell lines to reveal that TC11 cells express slightly higher endogenous SEMA7A (SFigure 2B).

To determine whether SEMA7A was necessary to drive pAKT in the TC11 cell line, we compared cell

lines stably transduced with shCtrl and shSEMA7A lentiviral vectors (shSEMA7A KD#1 and KD#2).

We confirmed SEMA7A expression was decreased in both KD vectors, with a larger decrease in KD#2

(SFigure 2C). As expected, knockdown of SEMA7A decreased pAKT levels, and the ratio of pAKT to

AKT was decreased in a SEMA7A-dependent manner (Figure 3A). Additionally, we show that

recombinant SEMA7A (rSEMA7A) or conditioned media from SEMA7A OE cells rescued the decrease

in pAKT observed in the KD cells (Figure 3B). To further define whether AKT activation by SEMA7A

occurs via SEMA7A-integrin binding, we treated TC11 cells with rSEMA7A, RGDS peptide or both,

and measured pAKT/AKT expression (Figure 3C). Although the RGDS peptide alone inhibited pAKT

levels, our results also show that rSEMA7A increased pAKT, and this effect was inhibited by the RGDS

peptide (Figure 3C). We observed similar results with the RGDS in two additional ER+ cell lines,

MCF7 and ZR751 (SFigure 2D-E).

SEMA7A+ tumor growth is inhibited by combination treatment with endocrine therapy and PI3K
inhibitors.

Since there are no clinically available integrin or ILK inhibitors, we investigated other

therapeutic strategies to inhibit SEMA7A-mediated signaling. Increased pAKT levels in SEMA7A-

expressing cells suggest a potential increase in PI3K signaling activity (46). Thus, we hypothesized that

SEMA7A+ ER+ BC cells would exhibit increased sensitivity to treatment with PI3K inhibitors (PI3Ki).

To test this hypothesis, we determined the 50% inhibitory concentration (IC50) for alpelisib (p110α

inhibitor) and GCT-007 (pan-PI3K inhibitor in development for treatment in glioblastoma and breast

cancer(47)) in MCF7 EV and SEMA7A OE cells (SFigure 3A-B). The IC50 of alpelisib and GCT-007

in SEMA7A OE cells was lower, suggesting increased susceptibility to PI3Ki in SEMA7A-expressing

tumor cells. We also examined sensitivity to PI3Ki in TC11 shSEMA7A #2 KD cells compared to

controls (SFigure 3C-D). In support of our hypothesis, cell viability assays revealed that inhibition by

both alpelisib and GCT-007 is greater in SEMA7A OE cells, compared to controls (SFigure 3E). We

also confirmed that pAKT expression is reduced by both PI3Ki treatments in MCF7 and TC11 (SFigure

3F-G) cells.

Since PI3K inhibition is utilized clinically in conjunction with endocrine therapy for ER+ breast

cancers, we aimed to evaluate whether a combination of endocrine therapy and PI3K inhibition (PI3Ki)

would be more effective than either treatment alone in SEMA7A-expressing tumors. Based on previous

literature (2) and our IC50 curves (SFigure 3A-D; SFigure 4A-B), we treated ER+ TC11 cells with

fulvestrant, alpelisib, or the combination and measured cell viability to reveal that fulvestrant alone had

minimal effect, while single-agent alpelisib and GCT-007 resulted in a moderate reduction in cell

viability, and the combination treatments were the most effective (Figure 4A-B). Similar results were

observed with the combination of fulvestrant plus alpelisib or GCT-007 in another ER+ cell line, ZR75-

1 (SFigure 4C-D). Notably, TC11 tumor cells are reported to exhibit resistance to tamoxifen (44).

Therefore, we also tested the combination of tamoxifen and each PI3Ki in TC11 cells; our findings

reveal that tamoxifen alone has minimal effect, whereas the combination significantly reduced cell

viability compared to control, to a greater extent than either single agent (Figure 4C-D). Following *in*

vitro synergy assays, Bliss synergy scores were calculated for the treatment combinations of fulvestrant

(Figure 4E-F) and tamoxifen (Figure 4G-H) with PI3K inhibitors. Synergy scores between 0 and 10

indicate an additive effect, while scores above 10 indicate synergy between two treatments, and we

observed synergy with fulvestrant/alpelisib (Score = 11.3), tamoxifen/alpelisib (Score = 14.61) and

tamoxifen/GCT-007 (Score = 16.33), and an additive effect with fulvestrant/GCT-007 (Score = 3.52)

(40). To evaluate the combination of endocrine therapy and PI3K inhibition *in vivo*, MCF7 SEMA7A

OE tumor-bearing NCG mice were treated with fulvestrant alone or in combination with alpelisib or

GCT-007, resulting in a greater reduction of tumor growth with the combination treatments (SFigure

4E-F). We monitored mouse weight and moribound criteria and observed no evidence of toxicities with

either PI3K inhibitor based on these criteria (SFigure 4G-H). In data from the manufacturer, GCT-007

exhibits efficacy in immunocompetent mice using a model of glioblastoma (unpublished observations).

Therefore, we tested combination treatments in an ER+ (SFigure 4I), estrogen-responsive (44)(SFigure

4J) immune-competent model. TC11 tumor-bearing FVB/N mice were treated with tamoxifen, GCT-

007, or the combination; the combination was most effective at inhibiting tumor growth compared to

either single agent or controls (Figure 4I). Consistent with the previous study, mouse no evidence of

toxicity was observed (SFigure 4K).

Inhibition of SEMA7A using a monoclonal antibody (SmAbH1) reduces tumor growth and AKT

signaling in ER+ BC.

We next tested our hypothesis that resistance to endocrine therapy, mediated by SEMA7A (2),

could be reversed by direct targeting of SEMA7A using our novel anti-SEMA7A monoclonal antibody

(SmAbH1) that we have previously shown to have efficacy in ER-negative models of breast

cancer(32,33). Our *in vitro* BioSpa live imaging assay results show that cell viability is reduced by

SmAbH1 over time in MCF7 SEMA7A OE cells, but not in EV controls (Figure 5A). The IC50 for

SmAbH1 is also lower in SEMA7A OE compared to controls (SFigure 5A), suggesting SmAbH1 is

most effective in SEMA7A-expressing cells. Previous studies from our lab have confirmed binding

specificity of SmAbH1 to SEMA7A protein via ELISA (29). In contrast, TC11 SEMA7A shSEMA7A

KD#2 showed less response to SmAbH1 compared to TC11 shCtrl cells *in vitro* (Figure 5B). Our

previous studies suggest that SmAbH1 is not efficacious in immune-compromised hosts (32). Thus, we

evaluated SmAbH1 in the TC11 model *in vivo* (SFigure 5B-D). We observed inhibition of tumor growth

in SmAbH1-treated tumors only (SFigure 5B-C). We observed no evidence of toxicity based on mouse

weight and moribound criteria (SFigure 5D). We validated these findings in the SSM2/129Sv-E model,

an additional ER+ immune-competent model (48) with endogenous SEMA7A expression (SFigure 2B,

SFigure 5E). As further evidence of specificity of SmAbH1, we observed that SmAbH1 inhibited tumor

growth compared to the IgG control (Figure 5C) and this effect was specific to SEMA7A-expressing

shCtrl tumors only (Figure 5C). Similarly, tumor volume calculated at study endpoint and tumor weight

were significantly reduced only in the TC11 shCtrl tumors treated with SmAbH1 (Figure 5D-E). To

confirm a SEMA7A/integrin-mediated mechanism *in vivo*, we performed IHC staining on control and

SmAbH1-treated TC11 tumors (Figure 5F). To look at pAKT levels, we performed RPPA analysis

(RPPA-2) of lysates made from IgG-treated and SmAbH1-treated TC11 and SSM2 tumors to reveal

decreased expression of pAKT (S473) in SmAbH1-treated tumors (Figure 5G, SFigure 5F). Further,

quantification of tumor IHC staining showed that activated integrin β1 (via 9EG7 staining), integrin β4,

and SEMA7A protein expression were reduced in the SmAbH1-treated group (Figure 5H-J, SFigure 5G:

secondary-only IHC controls, SFigure 5H-J: n=5 per group).

Combination treatment with fulvestrant and SmAbH1 reduces SEMA7A+ ER+ tumor growth.

To test the hypothesis that the combination of SmAbH1 and fulvestrant may benefit patients, we

tested cell viability in TC11 and MCF7 cell lines. TC11 shCtrl cell viability was inhibited by SmAbH1

alone, and even more so by combination of fulvestrant and SmAbH1, whereas TC11 shSEMA7A KD#2

viability was not affected (Figure 6A). Further, treatment of MCF7 EV and SEMA7A OE cells with

fulvestrant, SmAbH1, or the combination showed that SEMA7A OE cells respond positively to

SmAbH1 as a single agent and even more robustly to the combination of fulvestrant and SmAbH1,

while control cells do not (SFigure 6A). Additionally, EV cells exhibit a moderate response to

fulvestrant alone, while SEMA7A OE cells do not (SFigure 6A). These results were validated in SSM2

and ZR75-1 cells (SFigure 6B-C). In contrast, TC11 shSEMA7A cells displayed increased sensitivity to

fulvestrant alone *in vitro* (Figure 6A), suggesting that loss of SEMA7A may re-sensitize TC11 tumor

cells to fulvestrant. We also calculated synergy scores for fulvestrant in combination with SmAbH1 in

TC11 cells (Figure 6B). SmAbH1 with fulvestrant resulted in a synergistic score (Bliss score>10), while

SmAbH1 with tamoxifen was only slightly additive (Bliss score<10) (SFigure 6D), thus we moved

forward with fulvestrant. To test our hypothesis *in vivo,* TC11 shCtrl and shSEMA7A KD#2 tumor cells

were implanted into FVB/N mice and groups with Vehicle, IgG1, fulvestrant, SmAbH1, or the

combination. Tumor growth of shCtrl tumors was reduced by SmAbH1 and the combination therapy,

but not by the IgG or vehicle controls or fulvestrant alone (Figure 6C). Consistent with our hypothesis,

growth of TC11 shSEMA7A tumors was not affected by SmAbH1 alone but was reduced by fulvestrant

and the combination therapy (Figure 6D). Previous results were obtained with a hybridoma derived

SmAbH1. Thus, we also tested a recombinant version of anti-SEMA7A antibody, rSmAbH1, for range

of reactivity and specificity via IC50 and ELISA assay (SFigure 6E-F). Synergy results revealed that

rSmAbH1-P and fulvestrant are synergistic *in vitro,* similar to what was observed with SmAbH1 and

fulvestrant (SFigure 6G). We then repeated the TC11 *in vivo* experiment with rSmAbH1-P and

fulvestrant; images of harvested tumors show visibly smaller tumors in the rSmAbH1-P-treated and

combination-treated groups compared to controls or fulvestrant alone (Figure 6E). Mouse weights were

monitored (SFigure 6H), and a 6-chemistry toxicity panel was run on mouse plasma and showed no

differences between rSmAbH1-P-treated treatment and control-treated mice (SFigure 6I-N).

DISCUSSION

Collectively, our results reveal mechanisms of action and therapeutic vulnerabilities of SEMA7A

expressing ER+BC. Previous studies implicated integrin β1 as a receptor for SEMA7A on cancer cells

and other cell types, our findings suggest that SEMA7A binds to both integrin β1 and β4 in ER+ breast

cancer cells (Figure 2). We also provide evidence that the RGD motif within SEMA7A mediates binding

to β-integrins. Furthermore, our findings with the RGDS blocking peptide suggest that SEMA7A

enhances phosphorylation and activation of AKT specifically through integrins, which were previously

shown to undergo conformational changes when bound by an RGD motif leading to intracellular

activation of signaling pathways such as PI3K/AKT, ILK, Fak, and MAPK/ERK (49). Our results are

consistent with previously results showing that integrin β1 activation of PI3K and MAPK signaling can

facilitate endocrine therapy resistance in breast cancer (50); thus blocking this SEMA7A mediated

pathway is a potential way to overcome endocrine therapy resistance in patients with ER+BC.

Limitations of the current study include that the synthetic RGDS peptide may have also caused non-

specific integrin inhibition beyond that attributed to SEMA7A. Further, while the RGD motif in

SEMA7A is known, the protein folding structure of SEMA7A when bound versus unbound to integrins

remains unknown. Future studies could further investigate this protein interaction through mutations and

further structural analyses.

Our findings provide premise for treating patients with SEMA7A+ breast cancer with more

targeted therapies, including those that target of PI3K and/or directly target SEMA7A. Given our

previous studies showing poor relapse-free survival rates in both ER+ and ER- breast cancer patients (2),

clinical trials informed by our data stand to impact thousands of patients annually. We observed a

correlation between SEMA7A overexpression and increased pAKT (S473) and downstream targets of

AKT, suggesting activation of downstream survival signaling by AKT. Since ILK was moderately

increased in SEMA7A OE cells and ILK is known to phosphorylate AKT at S473, we suspect

involvement of ILK downstream of SEMA7A/integrin signaling. Because there are no clinically

available integrin or ILK inhibitors, we suggest that clinically approved PI3K inhibitors be tested to

inhibit the SEMA7A-AKT pathway. Clinical use of alpelisib is currently approved for patients with an

E545K mutation in *PIK3CA* that causes constitutive activation of the PI3K p110α subunit (24,51).

However, we hypothesize that even in the absence of the *PIK3CA* E545K mutation, high SEMA7A

expression can cause hyperactivation of PI3K including p110α, thus SEMA7A-expressing tumors may

be sensitive to alpelisib (52). GCT-007, a pan-PI3K inhibitor that has shown preclinical efficacy in

glioblastoma, is not approved yet for clinical use, but our results suggest it may be efficacious in breast

cancer patients as well. Limitations of the clinical applicability of our study include the known toxicities

observed in some patients including hyperglycemia, rash, and diarrhea (53), and while we saw no overt

toxicities in our mouse studies, the potential toxicities of GCT-007 in humans is unknown. Additionally,

while we did not explore direct targeting of AKT future studies could determine that this is another

viable option for patients with ER+SEMA7A+BC. Pan-AKT inhibitors such as capivasertib and

ipatasertib, mTOR inhibitors such as everolimus, and other PI3K inhibitors such as a p110δ inhibitors

(idelalisib) could be explored in our SEMA7A+ER+BC models to optimize available treatment

combinations. Moreover, we have published that the Bcl-2 inhibitor venetoclax can slow tumor growth

in pre-clinical models, and our RPPA-1 results confirm increased Bcl-2 expression with SEMA7A

expression; thus, combination of SmAbH1 and venetoclax should also be explored.

Our results suggest that direct inhibition of SEMA7A via our monoclonal antibody, SmAbH1, is

a promising therapeutic strategy that should be further investigated with additional pre-clinical studies to

enable progression to first in human clinical trials. While the combination of fulvestrant and SmAbH1

did reach statistical significance compared to SmAbH1 alone, it is important to note that SmAbH1 did

not appear to interfere with fulvestrant, which is utilized clinically for ER+ breast cancers and at study

end the tumors were visibly smaller in the combination-treated group. These results suggest that patients

will obtain benefit from the addition of SmAbH1 to their current standard of care (continued endocrine

therapy) in the recurrent/metastatic setting and that after additional testing SmAbH1 could become a

novel front line therapy for ER+BC patients that express SEMA7A. Aside from reduced tumor growth,

we also saw that SmAbH1 decreased pAKT (S473) suggesting that SmAbH1 is functionally inhibiting

the pathway activated by SEMA7A/integrin interaction. We also performed additional in vivo

SmAb/fulvestrant studies with a newly generated recombinant antibody, rSmAbH1-P, which is

necessary for humanization and future IND-enabling studies. A limitation of our study is the use of

mouse cell lines for the *in vivo* models instead of human cell lines. We deliberately chose the TC11 and

SSM2 models to test SmAbH1 because they are immunocompetent and we have recently published that

SmAbH1 activates anti-tumor T cell responses by re-activating anti-tumor T cell responses (29).

We conclude that inhibition of SEMA7A mediated signaling in breast cancer, both direct and

indirect, are promising clinical treatment avenues that should be further investigated particularly in

patients with recurrent disease despite endocrine therapies. Furthermore, our previous results suggest

that SEMA7A positive tumors will not respond to CDK4/6 inhibition, which is FDA approved for ER+

breast cancer patients. Additionally, our results suggest that patients with ER+ BC may benefit from a diagnostic test that quantifies SEMA7A expression as we have published a threshold of SEMA7A expression by immunostaining fixed tissues that can predict for relapse in a small patient dataset(17). If such a diagnostic existed, patients with SEMA7A+ tumors could receive a SEMA7A-targeted therapy like patients with HER2+ disease that are treated with trastuzumab. Future directions could also include the development and optimization of a minimally invasive SEMA7A-detecting diagnostic (blood) test, which we have initiated. However, further investigation is needed to determine whether SEMA7A levels in tumors correlate with levels in patient serum samples. Cumulatively, our current and previous studies suggest a unique and druggable aspect in the biology of SEMA7A-expressing breast cancers that could transform treatment options for ER+BC.

ACKNOWLEDGEMENTS

- Technical support
  - Lyons Lab: Lyndsey Crump, Lauren Cozzens, Kelsey Kines, Alan Elder, Alex Becks (University of Colorado Anschutz)
  - Sarah Tarullo (Plymouth State University)
  - Porter lab: Weston Porter, Garhett Wyatt (Texas A&M)
- Provided Reagents:
  - Global Cancer Technology
  - Novartis
- Clinical mentor:
  - Virginia Borges (CU Anschutz, UC Health)
- Thesis committee members: Carol Sartorius, Benjamin Bitler, Katherine Fantauzzo, John Tentler, Jennifer Richer, Daniel Sherbenou
- Experimental advice:
  - Michael Oliphant

-    NIH/NCATS Colorado CTSA Grant Number TL1 TR002533 to RS

-    NIC R01CA282900-01 and R01HD108335-01A1 to TL

-    SPARK/REACH Program, NIH: U01HL152405 and OEIDIT/AIA: MA 2021-2028 to TL

-    ABN Ohana Gates Grubstake Award to TL

-    DOM Outstanding Early Career Scholars Program to TL

-    UCCC and DOM ASPIRE to TL

-    Men for the Cure PO1 pilot grant to TL

-    University of Colorado Cancer Center Support Grant #P30CA046934 to RS

o   Anh Le, Steve Anderson and the University of Colorado Anschutz CTSR (Cell
Technologies Shared Resource)

o   University of Colorado Anschutz CPSR (Comparative Pathology Shared Resource)

-    Cancer Prevention & Research Institute of Texas Core Facility Support Awards (PR210227, PI:
Dean Edwards)

o   Baylor College of Medicine Antibody-Based Proteomics Core

▪   Dean Edwards

▪   Shixia Huang

REFERENCES

1.   ACS. 2024 American Cancer Society: Cancer Facts and Figures 2024. In American Cancer
Society <https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/2024-
cancer-facts-figures.html>.

2.   Crump LS, Wyatt GL, Rutherford TR, Richer JK, Porter WW, Lyons TR. Hormonal Regulation
of Semaphorin 7a in ER[+] Breast Cancer Drives Therapeutic Resistance. Cancer
Research **2021**;81:187-98

3.   Pan H, Gray R, Braybrooke J, Davies C, Taylor C, McGale P*, et al.* 20-Year Risks of Breast-
Cancer Recurrence after Stopping Endocrine Therapy at 5 Years. N Engl J Med **2017**;377:1836-
46

4.   Pasterkamp RJ, Peschon JJ, Spriggs MK, Kolodkin AL. Semaphorin 7A promotes axon
outgrowth through integrins and MAPKs. Nature **2003**;424:398-405

5. Bao ZZ, Jin Z. Sema3D and Sema7A have distinct expression patterns in chick embryonic
development. Dev Dyn **2006**;235:2282-9

6. Kohler D, Granja T, Volz J, Koeppen M, Langer HF, Hansmann G*, et al.* Red blood cell-derived
semaphorin 7A promotes thrombo-inflammation in myocardial ischemia-reperfusion injury
through platelet GPIb. Nat Commun **2020**;11:1315

7. Manini I, Ruaro ME, Sgarra R, Bartolini A, Caponnetto F, Ius T*, et al.* Semaphorin-7A on
Exosomes: A Promigratory Signal in the Glioma Microenvironment. Cancers (Basel) **2019**;11

8. Kinehara Y, Nagatomo I, Koyama S, Ito D, Nojima S, Kurebayashi R*, et al.* Semaphorin 7A
promotes EGFR-TKI resistance in EGFR mutant lung adenocarcinoma cells. JCI Insight **2018**;3

9. Xie J, Wang H. Semaphorin 7A as a potential immune regulator and promising therapeutic target
in rheumatoid arthritis. Arthritis Res Ther **2017**;19:10

10. Vaidya B, Patel R, Muth A, Gupta V. Exploitation of Novel Molecular Targets to Treat
Idiopathic Pulmonary Fibrosis: A Drug Discovery Perspective. Curr Med Chem **2017**;24:2439-
58

11. Garcia-Areas R, Libreros S, Simoes M, Castro-Silva C, Gazaniga N, Amat S*, et al.* Suppression
of tumor-derived Semaphorin 7A and genetic ablation of host-derived Semaphorin 7A impairs
tumor progression in a murine model of advanced breast carcinoma. Int J Oncol **2017**;51:1395-
404

12. Gutierrez-Franco A, Eixarch H, Costa C, Gil V, Castillo M, Calvo-Barreiro L*, et al.* Semaphorin
7A as a Potential Therapeutic Target for Multiple Sclerosis. Mol Neurobiol **2016**

13. Roth JM, Kohler D, Schneider M, Granja TF, Rosenberger P. Semaphorin 7A Aggravates
Pulmonary Inflammation during Lung Injury. PLoS One **2016**;11:e0146930

14. Tarullo SE, Hill RC, Hansen KC, Behbod F, Borges VF, Nelson AC*, et al.* Postpartum breast
cancer progression is driven by semaphorin 7a-mediated invasion and survival. Oncogene
2020;39:2772-85

15. Rutherford TR, Elder AM, Lyons TR. Anoikis resistance in mammary epithelial cells is
mediated by semaphorin 7a. Cell Death Dis **2021**;12:872

16. Black SA, Nelson AC, Gurule NJ, Futscher BW, Lyons TR. Semaphorin 7a exerts pleiotropic
effects to promote breast tumor progression. Oncogene **2016**;35:5170-8

17. Borges VF, Hu J, Young C, Maggard J, Parris HJ, Gao D*, et al.* Semaphorin 7a is a biomarker
for recurrence in postpartum breast cancer. npj Breast Cancer **2020**;6:56

18. Elder AM, Tamburini BAJ, Crump LS, Black SA, Wessells VM, Schedin PJ*, et al.* Semaphorin
7A Promotes Macrophage-Mediated Lymphatic Remodeling during Postpartum Mammary
Gland Involution and in Breast Cancer. Cancer Res **2018**;78:6473-85

19. Kines KaF, Heather and Elder, Alan and Cozzens, Lauren and Strugar, Zachary and Porter,
Weston and Borges, Virginia and Lyons, Traci, . Aging-Induced Semaphorin 7a Promotes
Tgfβ1-Mediated Cell Plasticity and Breast Tumor Metastases. **2024**

20.   Burris HA, 3rd. Overcoming acquired resistance to anticancer therapy: focus on the
PI3K/AKT/mTOR pathway. Cancer Chemother Pharmacol **2013**;71:829-42

21.   Truong TH, Benner EA, Hagen KM, Temiz NA, Kerkvliet CP, Wang Y*, et al.* PELP1/SRC-3-
dependent regulation of metabolic PFKFB kinases drives therapy resistant ER+ breast cancer.
Oncogene **2021**;40:4384-97

22.   Suzuki K, Kumanogoh A, Kikutani H. Semaphorins and their receptors in immune cell
interactions. Nat Immunol **2008**;9:17-23

23.   Andre F, Ciruelos EM, Juric D, Loibl S, Campone M, Mayer IA*, et al.* Alpelisib plus fulvestrant
for PIK3CA-mutated, hormone receptor-positive, human epidermal growth factor receptor-2-
negative advanced breast cancer: final overall survival results from SOLAR-1. Ann Oncol
2021;32:208-17

24.   André F, Ciruelos E, Rubovszky G, Campone M, Loibl S, Rugo HS*, et al.* Alpelisib for
PIK3CA-mutated, hormone receptor–positive advanced breast cancer. New England Journal of
Medicine **2019**;380:1929-40

25.   Duarte A, Silveira GG, Soave DF, Costa JPO, Silva AR. The Role of the LY294002 - A Non-
Selective Inhibitor of Phosphatidylinositol 3-Kinase (PI3K) Pathway- in Cell Survival and
Proliferation in Cell Line SCC-25. Asian Pac J Cancer Prev **2019**;20:3377-83

26.   Yang SX, Polley E, Lipkowitz S. New insights on PI3K/AKT pathway alterations and clinical
outcomes in breast cancer. Cancer Treat Rev **2016**;45:87-96

27.   Bostner J, Karlsson E, Pandiyan MJ, Westman H, Skoog L, Fornander T*, et al.* Activation of
Akt, mTOR, and the estrogen receptor as a signature to predict tamoxifen treatment benefit.
Breast Cancer Res Treat **2013**;137:397-406

28.   Chen H, Mingming C, Pengcheng G, Xiangzhong Z, Ganggang L, Liting W*, et al.* GDC-0941
activates integrin linked kinase (ILK) expression to cause resistance to GDC-0941 in breast
cancer by the tumor necrosis factor (TNF)-α signaling pathway. Bioengineered **2022**;13:10944-
55

29.   Elder AM, Fairchild HR, Kines KT, Cozzens LM, Becks AR, Slansky JE*, et al.* Semaphorin7A
and PD-L1 cooperatively drive immunosuppression during mammary involution and breast
cancer. Cell Rep **2025**;44:115676

30.   Czopik AK, Bynoe MS, Palm N, Raine CS, Medzhitov R. Semaphorin 7A Is a Negative
Regulator of T Cell Responses. Immunity **2006**;24:591-600

31.   Gómez Tejeda Zañudo J, Mao P, Alcon C, Kowalski K, Johnson GN, Xu G*, et al.* Cell Line-
Specific Network Models of ER(+) Breast Cancer Identify Potential PI3Kα Inhibitor Resistance
Mechanisms and Drug Combinations. Cancer Res **2021**;81:4603-17

32.   Kines KaF, Heather and Elder, Alan and Cozzens, Lauren and Strugar, Zachary and Porter,
Weston and Borges, Virginia and Lyons, Traci, . Aging-Induced Semaphorin 7a Promotes
Tgfβ1-Mediated Cell Plasticity and Breast Tumor Metastases. Cell Press Sneak Peak **2024**

33.   Elder AM, Fairchild HR, Kines KT, Cozzens LM, Becks AR, Lyons TR. Semaphorin7A and
PD-L1 cooperatively drive immunosuppression during mammary involution and breast cancer.
bioRxiv **2025**:2025.01.03.631229

34.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nat
Methods **2012**;9:671-5

35.   DeRose YS, Gligorich KM, Wang G, Georgelas A, Bowman P, Courdy SJ*, et al.* Patient-derived
models of human breast cancer: protocols for in vitro and in vivo applications in tumor biology
and translational medicine. Curr Protoc Pharmacol **2013**;Chapter 14:Unit14 23

36.   Massarweh S, Osborne CK, Creighton CJ, Qin L, Tsimelzon A, Huang S*, et al.* Tamoxifen
resistance in breast tumors is driven by growth factor receptor signaling with repression of
classic estrogen receptor genomic function. Cancer Res **2008**;68:826-33

37.   Gyorffy B. Survival analysis across the entire transcriptome identifies biomarkers with the
highest prognostic power in breast cancer. Comput Struct Biotechnol J **2021**;19:4101-9

38.   Goldman MJ, Craft B, Hastie M, Repecka K, McDade F, Kamath A*, et al.* Visualizing and
interpreting cancer genomics data via the Xena platform. Nat Biotechnol **2020**;38:675-8

39.   Coarfa C, Grimm SL, Rajapakshe K, Perera D, Lu HY, Wang X*, et al.* Reverse-Phase Protein
Array: Technology, Application, Data Processing, and Integration. J Biomol Tech **2021**

40.   Zheng S, Wang W, Aldahdooh J, Malyutina A, Shadbahr T, Tanoli Z*, et al.* SynergyFinder Plus:
Toward Better Interpretation and Annotation of Drug Combination Screening Datasets.
Genomics Proteomics Bioinformatics **2022**;20:587-96

41.   Davis JB, Andes S, Espina V. Reverse Phase Protein Arrays. Methods Mol Biol **2021**;2237:103-
22

42.   Xu W, Yang Z, Lu N. A new role for the PI3K/Akt signaling pathway in the epithelial-
mesenchymal transition. Cell Adh Migr **2015**;9:317-24

43.   Ludwig BS, Kessler H, Kossatz S, Reuning U. RGD-Binding Integrins Revisited: How Recently
Discovered Functions and Novel Synthetic Ligands (Re-)Shape an Ever-Evolving Field. Cancers
(Basel) **2021**;13

44.   Jallow F, O'Leary KA, Rugowski DE, Guerrero JF, Ponik SM, Schuler LA. Dynamic
interactions between the extracellular matrix and estrogen activity in progression of ER+ breast
cancer. Oncogene **2019**;38:6913-25

45.   Mori H, Chen JQ, Cardiff RD, Pénzváltó Z, Hubbard NE, Schuetter L*, et al.* Pathobiology of the
129:Stat1 (-/-) mouse model of human age-related ER-positive breast cancer with an immune
infiltrate-excluded phenotype. Breast Cancer Res **2017**;19:102

46.   Mayer IA, Arteaga CL. The PI3K/AKT Pathway as a Target for Cancer Treatment. Annu Rev
Med **2016**;67:11-28

47.   Technology GC. 2024 Global Cancer Technology. <https://globalcancertechnology.com/>.

48.   Mori H, Chen JQ, Cardiff RD, Pénzváltó Z, Hubbard NE, Schuetter L, *et al.* Pathobiology of the
651   129:Stat1−/− mouse model of human age-related ER-positive breast cancer with an immune
652   infiltrate-excluded phenotype. Breast Cancer Research **2017**;19:102

653   49.   Nieberler M, Reuning U, Reichart F, Notni J, Wester HJ, Schwaiger M, *et al.* Exploring the Role
654   of RGD-Recognizing Integrins in Cancer. Cancers (Basel) **2017**;9

655   50.   Pontiggia O, Sampayo R, Raffo D, Motter A, Xu R, Bissell MJ, *et al.* The tumor
656   microenvironment modulates tamoxifen resistance in breast cancer: a role for soluble stromal
657   factors and fibronectin through β1 integrin. Breast Cancer Res Treat **2012**;133:459-71

658   51.   Chang DY, Ma WL, Lu YS. Role of Alpelisib in the Treatment of PIK3CA-Mutated Breast
659   Cancer: Patient Selection and Clinical Perspectives. Ther Clin Risk Manag **2021**;17:193-207

660   52.   Armaghani AJ, Han HS. Alpelisib in the Treatment of Breast Cancer: A Short Review on the
661   Emerging Clinical Data. Breast Cancer (Dove Med Press) **2020**;12:251-8

662   53.   Wilhoit T, Patrick JM, May MB. Alpelisib: A Novel Therapy for Patients With PIK3CA-
663   Mutated Metastatic Breast Cancer. J Adv Pract Oncol **2020**;11:768-75

Figure 1. SEMA7A is expressed in ER+ breast cancer and is associated with decreased 5-year patient survival.

Survival analysis was done via Kaplan-Meier Plotter, which was assessed for *SEMA7A* mRNA expression levels in breast cancer patients in the ER+ subtype dataset (n=3499). Data are shown as KM plots; curves are stratified by *SEMA7A* high (red) and low (black) expression. Hazard ratios (HR) and log-rank p-values are included.

A) Overall survival; ER+ subtype (OS; n=221).

B) Distant metastasis-free survival; ER+ subtype (DMFS; n=248).

C) Post-progression survival; ER+ subtype (PPS; n=34).

Further analysis was done within the ER+ subtype to include only patients who had received endocrine therapy (n=1831).

D) Relapse-free survival; ER+ and received endocrine therapy (RFS; n=385).

E) Distant metastasis-free survival: ER+ and received endocrine therapy (DMFS; n=205).

Figure 2. SEMA7A activates AKT signaling and binds to Integrins β1 and β4 via RGD in ER+ tumor cells.

A) Gene Ontology (GO) Pathway Analysis of RPPA results, showing top pathways upregulated in MCF7 SEMA7A OE cell lysates, as fold change compared to EV, p<0.05.

B) Log2(fold change) of integrin and AKT-related proteins in MCF7 SEMA7A OE versus EV lysates from RPPA-1 data. Multiple two-tailed unpaired t-tests.

C) Immunoblot for Integrin β4, Integrin β1 and SEMA7A in elutions following CoIP of MCF7 EV and SEMA7A OE lysates for IgG or SEMA7A. ii. Quantification of IP shown as fold change compared to EV. Two-tailed unpaired t-test.

D) Immunoblot for Integrin β4, Integrin β1 and SEMA7A in elutions following CoIP of MCF7 lysates for IgG or SEMA7A, after treatment with PBS (-) or 50 μM RGDS (+). Quantification of IP shown as fold change compared to control. Two-tailed unpaired t-test.

Error bars are mean +/- SD. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 3. SEMA7A-mediated activation of AKT is dependent on RGD binding in ER+ tumor cells.

A) Representative immunoblot for pAKT (S473), AKT and GAPDH in TC11 shCtrl and shSEMA7A (KD#1 and KD#2) cell lines. Bar graph: ratio of pAKT to total AKT. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

B) Representative immunoblot for pAKT (S473), AKT and GAPDH in TC11 shCtrl and shSEMA7A KD#2 cell lines after treatment for 24 hours with 20 µg/mL purified rSEMA7A, or conditioned media from SEMA7A OE cells (S7OE CM). Bar graph: ratio of pAKT to total AKT. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

C) Representative immunoblot for pAKT (S473), AKT and GAPDH in TC11 WT cell lysates after treatment for 1 hour with 20 µg/mL rSEMA7A, 50 µM RGDS peptide, or both. Bar graph: ratio of pAKT to total AKT. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

Error bars are mean +/- SD. *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Figure 4. SEMA7A+ tumor growth is inhibited by combination treatment with endocrine therapy and PI3K inhibitors.

A) Percent cell viability quantified via crystal violet staining after 48-hour treatment of TC11 cells with DMSO, 8nM fulvestrant, 1μM alpelisib, or the combination, normalized to control. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

B) Percent cell viability quantified via crystal violet staining after 48-hour treatment of TC11 cells with DMSO, 8nM fulvestrant, 0.3μM GCT-007, or the combination, normalized to control. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

C) Percent cell viability quantified via crystal violet staining after 48-hour treatment of TC11 cells with DMSO, 5 μM 4-hydroxytamoxifen, 1μM alpelisib, or the combination, normalized to control. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

D) Percent cell viability quantified via crystal violet staining after 48-hour treatment of TC11 cells with DMSO, 5 μM 4-hydroxytamoxifen, 0.3 μM GCT-007, or the combination, normalized to control. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

E) Bliss synergy score for fulvestrant and alpelisib in TC11 cells.

F) Bliss synergy score for fulvestrant and GCT-007 in TC11 cells.

G) Bliss synergy score for (4-hydroxy)tamoxifen and alpelisib in TC11 cells.

H) Bliss synergy score for (4-hydroxy)tamoxifen and GCT-007 in TC11 cells.

I) Average TC11 tumor volume over time in FVB/N mice, in groups treated with Vehicle, GCT-007, tamoxifen, or the combination. GCT = GCT-007, TAM = tamoxifen. Bar graph: tumor volume at the study endpoint (Day 13). Ordinary one-way ANOVA with Tukey's multiple comparisons test.

Error bars are mean +/- SD (A-D), or mean +/- SEM (I). *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Figure 5. Direct inhibition of SEMA7A using a monoclonal antibody (SmAb H1) reduces tumor growth and AKT signaling in ER+ BC.

A) Brightfield cell count measured by BioSpa Live Imaging over 96 hours of MCF7 EV and SEMA7A OE cells treated with mouse IgG1 or 1 μM SmAb H1. Two-tailed unpaired t-test at time = 96 hours.

B) Representative crystal violet staining of TC11 shCtrl and shSEMA7A KD#2 cells after treatment for 48 hours with mouse IgG1 or 1 μM SmAb H1. Bar graph: quantification of all replicates. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

C) Average tumor volume in TC11 tumor-bearing mice over time, with groups receiving IgG1 or SmAb H1 (n=5 per group). Two-tailed unpaired t-test at day 4, 6, and 8 between (shCtrl) IgG and (shCtrl) SmAb H1 groups..

D) Tumor volume calculated using necropsy tumor photos in TC11 shCtrl and shSEMA7A-tumor bearing mice over time with groups receiving IgG1 or SmAb H1 (n=10 per group). Ordinary one-way ANOVA with Tukey's multiple comparisons test.

E) Weight of the tumor (and attached mammary gland) at study endpoint (n=10/group). Ordinary one-way ANOVA with Tukey's multiple comparisons test.

F) Immunohistochemistry staining of ex vivo TC11 tumors from IgG-treated and SmAb H1-treated groups. Scale bars = 50μm. Hematoxylin and eosin (H&E) staining (i-ii), active integrin B1 (9EG7) staining; brown (iii-iv), integrin B4 staining; brown (v-vi), and SEMA7A staining; brown (vii-viii).

G) Log2(fold change) of pAKT (S473) expression in SmAb H1-treated versus IgG-treated TC11 tumors from RPPA-2 data. Two-tailed unpaired t-test.

H) Quantification of 9EG7 (active ITGB1) IHC staining from all tumors (n=5 per group); Percent positive out of tumor area. Two-tailed unpaired t-test.

I) Quantification of ITGB4 IHC staining from all tumors (n=5 per group); Percent positive out of tumor area. Two-tailed unpaired t-test.

J) Quantification of SEMA7A IHC staining from all tumors (n=5 per group); percent positive out of tumor area. Two-tailed unpaired t-test.

Error bars are mean +/- SD (A-B), or mean +/- SEM (C-E, G-I). *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Figure 6. Combination treatment with fulvestrant and SmAb H1 reduces SEMA7A+ ER+ tumor growth.

A) Cell viability quantified via crystal violet staining after 48-hour treatment of TC11 shCtrl and TC11 shSEMA7A KD#2 cells with DMSO, 8 nM fulvestrant, 1μM SmAb H1, or the combination, normalized to control. Ordinary one-way ANOVA with Tukey's multiple comparisons test.

B) Bliss synergy score for fulvestrant and SmAb H1 in TC11 cells.

C) Average tumor volume in TC11 shCtrl tumor-bearing FVB/N mice over time with groups receiving Vehicle F, IgG1, fulvestrant, SmAb H1 or the combination (n=5 per group). Bar graph: tumor volume at study endpoint (Day 7). Ordinary one-way ANOVA with Tukey's multiple comparisons test.

D) Average tumor volume in TC11 shSEMA7A KD#2 tumor-bearing FVB/N mice over time with groups receiving Vehicle F, IgG1, fulvestrant, SmAb H1 or the combination (n=5 per group). Bar graph: tumor volume at study endpoint (Day 7). Ordinary one-way ANOVA with Tukey's multiple comparisons test.

E) Representative photographs of harvested TC11 tumors treated with Vehicle F, IgG1, fulvestrant, rSmAb H1 or the combination (n=5 per group). Scale bar = 1cm.

Error bars are mean +/- SD (A), or mean +/- SEM (C-D). $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$.