Impact of Group Size and Habitat Disturbance on Parasitic Infection in Free-

ranging Proboscis Monkeys

[1,a]Muhammad Nur Fitri-Suhaimi, [2,a*]Liesbeth Frias, [3b]Elke Zimmermann, [4]Primus Lambut,

[5]Joseph Tangah, [6]Henry Bernard, [7]Vijay Subbiah Kumar, [1,6,8**]Ikki Matsuda

[1]*Wildlife Research Center of Kyoto University, 2-24 Tanaka-Sekiden-cho, Sakyo, Kyoto 606-*

8203, Japan

[2]*Department of Infectious Diseases and Public Health, Jockey Club College of Veterinary*

Medicine and Life Sciences, City University of Hong Kong, Hong Kong SAR

[3]*Institute of Zoology, University of Veterinary Medicine Hannover, Hannover, Lower Saxony,*

Germany,

[4]*Sabah Wildlife Department, Wisma Muis, Kota Kinabalu, Sabah 88100, Malaysia*

[5]*Sabah Forestry Department, Forest Research Centre, Sandakan, Sabah, Malaysia*

[6]*Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Jalan UMS,*

Kota Kinabalu, Sabah, 88400, Malaysia

[7]*Biotechnology Research Institute, Universiti Malaysia Sabah, Jalan UMS, Kota Kinabalu,*

Sabah 88400, Malaysia

[8]*Chubu Institute for Advanced Studies, Chubu University, Kasugai, Aichi, Japan*

[a]*Co-first Author*

[b]*Elke Zimmermann passed away on 25th July 2019*

[*]*Correspondence to: Liesbeth Frias, Department of Infectious Diseases and Public Health,*

Jockey Club College of Veterinary Medicine and Life Sciences, City University of Hong

Kong, To Yuen Building, 31 To Yuen Street, Kowloon Tong, Kowloon, Hong Kong SAR

[**]*Correspondence to: Ikki Matsuda, Wildlife Research Center of Kyoto University, 2-24*

Tanaka-Sekiden-cho, Sakyo, Kyoto 606-8203, Japan

E-mail addresses: matsuda.ikki.5a@kyoto-u.ac.jp (IM); liesbeth.frias@gmail.com (LF)

Abstract

Group-living primates experience the benefits and costs associated with sociality, including an

elevated risk of parasite transmission. However, the relative influence of group type (i.e., social

structure), group size, and habitat disturbance on parasitic infection remains unclear,

particularly in Southeast Asian primates. In this study, the abundance of intestinal parasites in

proboscis monkeys (*Nasalis larvatus*) inhabiting a riverine forest along the Menanggul River,

Sabah, Malaysian Borneo, was investigated. Fecal samples (n = 160) were collected from one-

male-multifemale and all-male groups in areas with varying levels of anthropogenic

disturbance, with efforts made to ensure that each sample originated from a different individual.

In addition, the effects of group type, group size, and sampling location on parasite abundance

were evaluated using fecal egg counts and Bayesian models. Three dominant parasite species

groups (*Trichuris* sp., *Strongyloides fuelleborni*, and *Oesophagostomum aculeatum*) with an

overall infection prevalence of 81.25% were identified. Results showed that group type did not

significantly affect parasite abundance. However, group size showed a positive correlation with

the abundance of *Trichuris* sp. and a negative correlation with *S. fuelleborni* and *O. aculeatum*.

In addition, our models revealed that the infection load of *Trichuris* sp. decreased with

increasing distance from the river mouth, which was used as a proxy for a disturbance gradient,

whereas *O. aculeatum* exhibited higher infection load at greater distances, indicating lower

prevalence in more disturbed downstream areas. Thus, parasite abundance in proboscis

monkeys may be shaped by social and environmental factors, with taxa-specific responses

likely reflecting differences in environmental persistence and transmission ecology.

Keywords: Host-parasite interactions, Anthropogenic impact, Primates, Social organization,

Zoonotic risk, Wildlife disease ecology

Introduction

Highly social animals such as primates exhibit complex group dynamics that influence their health, behavior, and interactions with pathogens (Kappeler et al., 2015). Although group-living promotes social support and cooperative behaviors that can mitigate some health risks (Silk, 2007), it also imposes costs, including an increased potential for pathogen transmission and heightened susceptibility to disease caused by social stress (Altizer et al., 2003). Thus, the relationship between sociality and disease risk is complicated. Factors such as group structure, contact patterns, and social hierarchies all play key roles in determining infection dynamics (Nunn et al., 2015). Notably, this relationship is bidirectional in primates. While social factors such as group size and composition influence parasite transmission, parasite infection can also feed back to shape host social behavior. Consequently, infection prevalence and intensity can affect social interaction and group dynamics (Vitone et al., 2004; Nunn, 2012; Rushmore et al., 2017).

Apart from social dynamics, anthropogenic environmental changes, such as deforestation and provisioning, can also influence parasite transmission by altering host movement patterns, population densities, and contact rates (Gillespie and Chapman, 2006; Becker et al., 2018). These ecological modifications can create conditions that either facilitate or hinder parasite spread, depending on the nature and extent of human-induced disturbances. Therefore, understanding the mechanism by which environmental, social, and ecological factors contribute to parasitic infections in social animals is crucial for developing effective conservation and management strategies (Chapman et al., 2005; Gillespie and Chapman, 2006).

On the contrary, research on parasitic infection and sociality in wild primates has yielded inconsistent findings, while the effects of anthropogenic environmental changes on parasitic infections in primates have been relatively consistent (Gillespie and Chapman, 2006). The relationship between parasite load and group size is complex, as it depends on various factors, such as host susceptibility and environmental conditions (e.g. Chapman et al., 2009;

Klaus et al., 2018). Although larger groups might increase the likelihood of infection because

of higher social interactions, group mobility and seasonal changes may reduce the risk of

parasite transmission (e.g., increased precipitation during the rainy season washing away feces

and parasites), thereby complicating the host–parasite dynamics (Nguyen et al., 2015;

Radespiel et al., 2015; Knauf and Jones-Engel, 2020).

Although knowledge of parasitism in wild primates is expanding, it remains limited,

particularly in Southeast Asia (Hopkins and Nunn, 2007; Cooper and Nunn, 2013). In Borneo,

which is a highly biodiverse island in Southeast Asia, numerous endemic species are

experiencing population declines (Wilcove et al., 2013), and the large arboreal proboscis

monkey (*Nasalis larvatus*) inhabiting mangroves, peat swamps, and riverine forests is no

exception to this trend (Lhota et al., 2019; Bernard et al., 2025). Several studies have provided

insights into primate–parasite interactions in Sabah, including the proboscis monkey. This

primate has unique parasite associations (Hasegawa et al., 2020) and infection dynamics that

reflect its specialized ecological adaptations (Klaus et al., 2017) and interactions with

sympatric primates (Frias et al., 2019; Frias et al., 2021). Previous studies have reported a high

prevalence of helminth eggs in fecal samples, identifying five major parasite groups, with co-

infections being common (Salgado-Lynn, 2010; Klaus et al., 2017; Frias et al., 2021). In

addition, parasite species richness has been reported to be significantly higher in semi-wild

populations than in wild ones. This increase likely results from the high levels of human

disturbance, which can alter the mechanism by which parasites are transmitted. By contrast,

group type and group size have been reported not to significantly affect parasite richness (Klaus

et al., 2018). Despite the increasing knowledge of parasitic infections in proboscis monkeys,

the fine-scale variation in infection patterns –across individuals or groups inhabiting distinct

microhabitats and experiencing different levels of anthropogenic disturbance– remains unclear.

In this study, the abundance of intestinal parasites in proboscis monkeys was

investigated, with particular emphasis on the effects of group type, group size, and habitat

disturbance. Fecal samples collected over a 1-year period were analyzed to identify the

parasites present and to examine the mechanism by which group size and social structure (i.e.,

group type: one-male-multifemale or all-male groups) influence parasite loads in a forest

subjected to varying degrees of anthropogenic disturbance. Considering that proboscis

monkeys exhibit seasonal variation in foraging behavior, ranging patterns, and intergroup

spacing at our study site (Matsuda et al., 2009a; Matsuda et al., 2009b; Matsuda et al., 2010a),

a year-round sampling approach may provide a comprehensive assessment of parasite

dynamics. Thus, extended sampling could reveal a positive correlation between group size and

parasitic infection, which is contrary to previous findings (Klaus et al., 2018). In addition, the

study forest along the tributary river exhibits varying degrees of anthropogenic disturbance

between the upstream and downstream areas. In particular, downstream areas are more strongly

influenced by tourism-related boat traffic, human activity, and proximity to oil palm plantations,

whereas upstream areas retain relatively higher plant diversity (Jose et al., 2024). The impact

of habitat disturbance on parasite abundance can be assessed by comparing the fecal samples

collected from individuals across these areas. Given that research in our study area has

documented alterations in the gut microbiota of proboscis monkeys living in disturbed versus

undisturbed areas (Jose et al., 2024), we hypothesize that parasite abundance would also differ

between these environments because of human-induced disturbances.

Methods

Study site and subjects

This study was conducted in a riverine forest along the Menanggul River, a tributary of the

Kinabatangan River, in Sabah, Malaysian Borneo (118°30′ E, 5°30′ N). The study site has a

mean daily temperature range of 24°C–30°C and an annual average precipitation of 2474 mm

(Matsuda et al., 2019). The river exhibits daily water-level fluctuations of approximately 1 m,

while seasonal inundation results in an average increase exceeding 3 m (Matsuda et al., 2010b).

The study area spans 4 km along the Menanggul River and supports a minimum population of

200 proboscis monkeys, forming a multilevel society composed of one-male-multifemale

groups and all-male groups, which frequently assemble in the riverside (Matsuda et al., 2024).

The riparian zone primarily consists of secondary forest in the southern portion, while much of

the northern portion has been converted to oil palm plantations, except for a narrow, protected

riparian strip. Accordingly, in this tributary, forests extending upstream of the river mouth

exhibit a higher plant diversity (Jose et al., 2024). The proboscis monkeys in the study area are

well habituated to human presence, as the Menanggul River has been a major tourist attraction

for over a decade and serves as a frequent route for boats and observers.

Sample collection

Between June 2015 and April 2016, boat surveys were conducted in the late afternoon on

approximately 2–3 days per week to locate proboscis monkeys, as they typically return to

riverside trees to sleep (Matsuda et al., 2011), and to record group composition and the GPS

coordinates of sleeping locations. The following morning, before the monkeys awakened, these

sleeping locations were revisited. Given that proboscis monkeys often defecate just before

moving into the forest, the area around their sleeping sites was carefully searched for fresh

feces after the monkeys had departed their sleeping trees. Multiple one-male-multifemale

groups often remained close to one another along the river, making it challenging to determine

to which group the feces found on the ground belonged. Therefore, we inferred the group

associated with the feces based on the location of the feces and the group observed in the

previous day's survey. Only fecal samples presumed to originate from adult individuals were

included, based on fecal size, which allows for differentiation between adults and immature

individuals despite the challenges in distinguishing feces from neighboring groups (Matsuda

et al., 2014). Consequently, between June 2015 and April 2016, 160 fecal samples were

collected. Individual identity was subsequently confirmed through genetic analyses, which

verified that the samples originated from genetically distinct individuals (Matsuda et al., 2024).

At least nine groups were identified in the study area, although the number of groups these

fecal samples were collected from remains unclear, as group identification was not complete

at the time the samples were collected (Matsuda et al., 2024). The fecal samples were preserved

in tubes with 70% ethanol.

Parasite detection and identification

A modified protocol was used to concentrate parasite eggs via formalin-ethyl acetate

sedimentation, and fecal samples were analyzed using the sequential sedimentation–flotation

method (Frias et al., 2021). The final pellet was resuspended in saline, and 1 mL of aliquot was

transferred into a vial and placed on a magnetic stirrer to ensure thorough homogenization

during analysis. In estimating the number of eggs per gram of feces (EPG) in each sample, an

aliquot was collected from the homogeneous suspension and placed in a McMaster counting

chamber for examination at 100× magnification. EPG was calculated on the basis of the

average of five replicate counts of parasite eggs observed within the chamber's grid, using the

known weight of fecal sediment in a 0.15 mL volume. After quantifying the eggs, each sample

was centrifuged again; the supernatant was discarded, and the concentrated pellet was

resuspended in Sheather's solution with a specific gravity of 1.27. Two slides were examined

to reduce the likelihood of overlooking less abundant helminth eggs. Parasite identification was

performed using standard identification keys (Modrý et al., 2018).

Parasite prevalence and abundance

The prevalence of a given parasite was calculated as the percentage of hosts infected by that

parasite. Parasite abundance was estimated for each sample using the EPG rounded to the

nearest integer as a proxy measure, following the terminology of Bush et al. (1997).

Factors influencing parasite abundance

A Bayesian model was developed using the "brms" package (Bürkner, 2018) to evaluate the

mechanism by which group type (one-male-multifemale or all-male groups), group size, and

fecal sampling locations affect parasite abundance. Sampling locations were represented by

distance from the river mouth to upstream locations, which served as a proxy for the

disturbance gradient along the river system. In our analysis, parasite abundance was used as a

dependent variable and modelled using a Poisson distribution with a log link. Group type was

incorporated as a group-level effect, whereas group size and the distance of the sampling

location to the river mouth were included as continuous variables. All models were controlled

for host group identity. Distinct groups were not consistently identifiable throughout the study

period. Therefore, we assigned samples to groups based on their spatial and temporal proximity

along the river. This approach was guided by observations: the mean distance between

consecutive sleeping sites for each group was approximately 350 m in the study area (Matsuda

et al., 2024). In addition, changes in food availability over time can influence spatial patterns

along the river (Matsuda et al., 2010a). Based on these criteria, two samples collected within

350 meters of each other and during the same month were classified as originating from the

same group. Groups with fewer than three observations were excluded from the analysis.

Weakly informative priors were assigned to population-level effects, following a

Student's t-distribution. The posterior distributions of the model parameters were estimated

using Markov chain Monte Carlo methods. For this purpose, four chains were constructed, each

consisting of 2000 steps, with 1000-step warm-up periods. Consequently, a total of 4000 steps

were retained to estimate the posterior distributions. The independent continuous variables

were scaled to aid model fitting and coefficient interpretation. Chain convergence was assessed

visually, and the potential scale reduction factor (Rhat) values were examined to ensure they

were less than or equal to 1 for all parameters (Bürkner, 2018). This modeling process was

repeated for the three parasites identified within the host population.

Results

Parasite prevalence and abundance

Parasites were detected in 130 of the 160 analyzed samples, yielding an overall parasite prevalence of 81.25%. The eggs of *Trichuris* sp. (78.12%, n = 125), *Strongyloides fuelleborni* (10.62%, n = 17), *Oesophagostomum aculeatum* (5%, n = 8), *Enterobius* (*Colobenterobius*) *serratus* (1.25%, n = 2), and *Spirurida* sp. (0.63%, n = 1) were identified. The mean abundance of the top three prevalent parasites was recorded at 168.2 EPG (range: 0–984), 9.41 EPG (range: 0–650), and 2.83 EPG (range: 0–109), respectively.

Factors influencing parasite abundance

Of the 160 fecal samples examined, 124 were included in the Bayesian analyses after excluding samples from groups with fewer than three observations, as these could not be reliably incorporated into the group-level models. The type of host group did not have a statistically significant effect on parasite abundance across any of the parasite groups (Fig. 1A). However, host group size emerged as an important predictor of parasite abundance, with varying effects across parasite groups. The abundance of *Trichuris* sp. was positively correlated with group size, whereas that of *S. fuelleborni* and *O. aculeatum* was negatively correlated with group size (Fig. 1B). In addition, our models indicated that the distance from the river mouth to upstream locations, used as a proxy for a disturbance gradient, markedly influenced the parasite abundance for *Trichuris* sp. and *O. aculeatum*. In particular, as the distance from the river increased, the abundance of *Trichuris* sp. decreased, whereas *O. aculeatum* exhibited an opposite trend, with higher counts linked to greater distances from the river mouth (Fig. 1C). A summary of the results from the models is provided in Table 1.

Discussion

In this study, the relationship among social structure (i.e., group type: one-male-multifemale

and all-male groups), the degree of anthropogenic habitat disturbances within a 4-km stretch

along the river, and parasite load in free-ranging proboscis monkeys was examined. A high

prevalence of intestinal parasites was observed (81.25%), which is consistent with the previous

prevalences recorded for proboscis monkeys along the Kinabatangan River in Sabah [i.e.,

96.6% in Salgado-Lynn (2010), 92.3% in Klaus et al. (2017), and 80.2% in Frias et al. (2021)]

and in Bako National Park in Sarawak [i.e., 95.4% in Adrus et al. (2019)]. The predominant

parasite groups in this study included *Trichuris* sp., *Strongyloides fuelleborni*, and

Oesophagostomum aculeatum, which have also been identified as dominant in other fecal

analyses of proboscis monkeys, in wild and captive settings (Table 2), reinforcing a consistent

pattern of parasitic infections across different populations. Nonetheless, given the substantial

differences in the diets of proboscis monkeys between riverine and mangrove forests (Matsuda

et al., 2019), habitat-linked variation in food use may also influence parasite exposure. Because

habitat type determines not only the food resources available to proboscis monkeys but also

forest height and frequency of ground use or sediment contact during foraging (Feilen and

Marshall, 2014, 2020; Atmoko et al., 2021), these ecological differences may affect

transmission dynamics of environmentally transmitted parasites. Thus, whether similar trends

in parasite prevalence are observed among proboscis monkeys across different habitat types on

the island of Borneo should be examined in the future.

However, no significant association was found between parasite abundance and group

type (i.e., one-male-multifemale and all-male groups) in proboscis monkeys in this study,

consistent with previous findings (Klaus et al., 2018), suggesting that sex differences may have

a limited influence on parasitic infection. This interpretation is further supported by a study

conducted at the same site that examined gut microbial composition in fecal samples and found

no significant sex differences (Jose et al., 2024). Given that gut microbiota, such as intestinal

parasites, would be transmitted through direct or indirect social contact and shared use of space

and food resources, the absence of sex-specific patterns in both types of organisms indicates

that behavioral and spatial overlap between males and females minimizes sex-related

divergence in microbial and parasite exposure.

Contrary to the findings of Klaus et al. (2018), our study revealed that group size might

be a potential factor influencing parasite abundance for a particular parasite species, namely,

Trichuris sp. This finding supports the hypothesis that larger groups may exhibit higher

infection rates, probably because of increased contact rates and overlapping space use among

individuals (Snaith et al., 2008; Rifkin et al., 2012; Kappeler et al., 2015). Although the exact

mechanism remains unclear, a comparable trend was observed in the gut microbial diversity in

the same population. A larger group size was associated with greater alpha diversity (Jose et

al., 2024), suggesting that higher host density may increase exposure to intestinal parasites and

microbes. This association may reflect the increased direct interactions among individuals and

result from enhanced environmental contamination in frequently reused areas. Given that all

the parasites identified in this study could be transmitted via fecal-oral routes, the accumulation

of feces in shared sleeping or foraging sites could be a key factor driving infection. Limited

territoriality and habitual reuse of sleeping trees by proboscis monkeys –often shared across

one-male-multifemale and all-male groups in multilevel societies– likely facilitates such

transmission (Matsuda et al., 2024). Furthermore, the differences in habitat structure and spatial

dynamics among the study sites may explain the contrasting results reported by Klaus et al.

(2018). They focused on groups inhabiting the main channel of the Kinabatangan River, where

the river's width (>200 m) limits cross-group movement and shared space use. By contrast, our

study site along narrow tributaries (15–20 m wide) may allow more frequent crossings and

higher spatial overlap among groups (Matsuda et al., 2008; Matsuda et al., 2010a), thereby

promoting indirect transmission through contaminated environments.

On the contrary, *S. fuelleborni* and *O. aculeatum* were negatively correlated with group

size because of their differing ecological tolerances. *S. fuelleborni*, which is a parasitic

nematode, may be more influenced by habitat disturbance or seasonal changes, with larger

groups potentially reducing parasite exposure through increased host mobility and spatial

separation from contaminated areas (Gillespie and Chapman, 2006; Radespiel et al., 2015).

Similarly, *O. aculeatum* may require more stable environmental conditions for successful

transmission, as the development and survival of free-living stages in *Oesophagostomum* spp.

are influenced by factors such as temperature and related habitat conditions (Sengupta et al.,

2016; Senanayake et al., 2025). The relatively undisturbed and pristine upstream habitats may

therefore provide better conditions for transmission compared with downstream areas, whereas

larger groups in disturbed downstream areas could hinder transmission (Kappeler et al., 2015).

More broadly, these contrasting patterns may reflect a trade-off between contamination

density and spatial avoidance. Larger groups may increase fecal contamination at frequently

reused sites, favoring parasites with environmentally resistant stages such as *Trichuris*.

Conversely, increased host mobility and shorter residence time at specific patches may reduce

exposure to parasites with more fragile free-living stages, such as *S. fuelleborni* and *O.*

aculeatum. These findings indicate that host social dynamics and environmental factors must

be considered, as species-specific responses shape infection patterns (Rifkin et al., 2012;

Nguyen et al., 2015).

Our findings revealed a more complex relationship between parasite abundance and

habitat disturbance than initially anticipated. Contrary to expectations that anthropogenic

habitat disturbance would universally elevate parasite loads (Gillespie and Chapman, 2006),

we observed that the impact of disturbance varied across different parasite taxa. In particular,

the abundance of *Trichuris* sp. decreased with increasing distance from the river mouth,

indicating that higher parasite loads occur in downstream areas with more intense human

activity and tourism. Conversely, *O. aculeatum* displayed the opposite trend, with greater

abundance in less disturbed upstream forests. These contrasting patterns indicate that different

parasite species may respond to habitat disturbance through distinct ecological mechanisms.

However, we acknowledge that distance from the river mouth serves as a proxy that likely

integrates multiple environmental gradients beyond anthropogenic disturbance alone. These

may include natural variation in soil composition, flooding regimes, vegetation structure, and

microclimatic stability, all of which could influence parasite spatial responses observed

between *Trichuris* sp. and *O. aculeatum.*

This interpretation would be consistent with differences in environmental persistence

between parasite taxa, as *Trichuris* eggs are relatively resistant, whereas *O. aculeatum* may

depend more strongly on stable microenvironmental conditions for successful development and

transmission. Considering the elevated *Trichuris* sp. abundance in downstream areas, more

disturbed habitats may not be attributable to the increased host density or spatial overlap, as

proboscis monkeys in these areas do not necessarily aggregate at higher densities compared

with the upstream areas. Aggregation likely occurs in upstream areas, where the river is

narrower, providing a safer environment for proboscis monkeys because they can escape

terrestrial predators, such as clouded leopards, by jumping to the opposite bank (Matsuda et al.,

2010a). Given its greater safety from predators, this environment likely facilitates aggregation

in upstream areas, making them a more favorable location for proboscis monkey groups to

congregate. This interpretation is consistent with the species' highly resilient egg morphology,

which supports environmental persistence even under degraded conditions (Gillespie and

Chapman, 2006; Snaith et al., 2008). Morphological robustness may enable *Trichuris* sp. to

maintain transmission success across a range of habitat types, including more disturbed

downstream areas. By contrast,   successful transmission of *O. aculeatum* requires

embryonation, hatching, and survival of infective larval stages in the environment, processes

that are influenced by environmental conditions such as temperature and moisture (Rose and

Small, 1980; Radespiel et al., 2015; Sengupta et al., 2016; Senanayake et al., 2025).

Accordingly, the relatively pristine conditions in upstream sites likely provide a favorable

microenvironment for their development.

Interestingly, these spatial patterns in parasite abundance are contrary to prior findings

from the same population, which identified clear site-based differences in gut microbial

composition, with lower microbial diversity in the downstream areas (Jose et al., 2024). Gut

microbiota and intestinal parasites share common transmission pathways, including social

contact and environmental exposure. However, their ecological sensitivities appear to diverge.

In general, gut microbiota are more responsive to dietary shifts and subtle ecological changes

(e.g., Amato et al., 2013; Hayakawa et al., 2018; Lee et al., 2019; Barelli et al., 2020). By

contrast, parasite abundance in our study is influenced by a combination of host social

dynamics (e.g., group size and space use) and species-specific environmental tolerances. This

distinction emphasizes the importance of adopting a taxon-specific approach when interpreting

the host–environment–parasite relationship. Furthermore, it indicates that parasites and

microbiota may differentially influence the ecological and health dynamics of primates (Rifkin

et al., 2012; Nguyen et al., 2015).

These contrasting patterns between gut microbiota diversity and *Trichuris* abundance

may reflect differences in the ecological processes governing their assembly. Gut microbiota

are strongly shaped by dietary and environmental inputs, which can be altered by habitat

disturbance (Amato et al., 2013; Hayakawa et al., 2018). In contrast, helminth infections

depend more directly on encounters with infective stages in the environment and patterns of

fecal contamination (Nunn et al., 2011). Parasites with environmentally resistant eggs, such as

Trichuris, may therefore persist and accumulate even in disturbed habitats where microbial

exposure diversity declines. Disturbed environments may thus simultaneously reduce

opportunities for microbial acquisition while maintaining or even increasing exposure to

environmentally persistent helminths.

Understanding the interplay among social organization, habitat characteristics, and

parasitic infection in wild primates deepens our ecological insights. It informs broader

questions in evolutionary biology and conservation. The findings of this study underscore the

complex, context-dependent nature of host–parasite dynamics. They highlight how intrinsic factors such as group size and behavior can mediate exposure risks, even in the face of external environmental change. Given the escalating habitat disturbance, a nuanced understanding is crucial for predicting disease risks in wildlife populations and for designing conservation strategies that account for social and ecological variability. Integrative approaches that consider behavioral ecology alongside environmental gradients may also be essential for safeguarding the health and persistence of socially complex yet increasingly vulnerable species such as the proboscis monkey.

Finally, the helminths detected in this study are of potential zoonotic relevance, especially in ecotourism landscapes such as the Menanggul River, where frequent contact occurs among humans, livestock, and wild primates. The genus *Trichuris* may include species closely related to *T. trichiura*, a well-known human pathogen that can infect non-human primates (Rivero et al., 2020). Likewise, *S. fuelleborni* and *O. aculeatum* have been reported in both human and non-human hosts in Southeast Asia (Yalcindag et al., 2021; Chan et al., 2023). However, because parasite identification in the present study was based solely on egg morphology, neither host specificity nor actual zoonotic transmission risk can be confirmed at this stage. Molecular characterisation will therefore be required to determine whether the detected parasites belong to human-infective lineages. Nevertheless, the presence of these helminths in areas accessible to humans highlights the need for further investigation of their transmission dynamics and potential public health relevance under increasing anthropogenic pressure (Gillespie and Chapman, 2006).

Acknowledgements

We express our sincere thanks to the Sabah Biodiversity Centre and the Sabah Wildlife Department for granting permission to carry out this research. We are also deeply indebted to the Sabah Forestry Department for facilitating the use of its field facilities. We particularly

391   thank the research assistants, Asnih Binti Etin and Jasrudy Bin Mandu, for their support in the

392   field. Finally, we are grateful to the reviewers for their fruitful comments that improved this

393   manuscript.

Author contribution

MNF: conceptualization, writing—original draft, methodology; LF: conceptualization,

writing—review and editing, methodology, visualization, formal analysis, investigation; EZ:

conceptualization, methodology; PL: resources, writing—review and editing; JT: resources,

writing—review and editing; HB: resources, writing—review and editing; VK: resources,

writing—review and editing; IM: conceptualization, data curation, funding acquisition,

investigation, methodology, project administration, resources, supervision, validation,

visualization, writing—original draft. All authors (except deceased EZ) gave final approval for

publication and agreed to be held accountable for the work performed therein.

Funding

406   This study was partially funded by the Japan Society for the Promotion of Science KAKENHI

407   (nos. 24H00774 to I.M.; no. 23K27254 to G.H.) and Core-to-Core Program, Asia-Africa

408   Science Platforms (JPJSCCB20250006 to IM).

Data availability

411   All data generated or analysed during this study are included in this published article.

Declarations

Ethics approval

The animal study was reviewed and approved by the Sabah Biodiversity Centre and the Sabah

Wildlife Department [JKM/MBS.1000-2/2 JLD.3 (120)].

Conflict of interest

The authors declare no competing interests.

Generative AI and AI-assisted technologies in the writing process

During the preparation of this manuscript, the authors used ChatGPT-5.2 to refine the clarity

and logical flow of the text. The authors carefully reviewed, corrected, and approved all content

generated, and take full responsibility for the final published version.

References

Adrus, M., Zainuddin, R., Ahmad Khairi, N.H., Ahamad, M., Abdullah, M.T., 2019. Helminth parasites occurrence in wild proboscis monkeys (*Nasalis larvatus*), endemic primates to Borneo Island. J Med Primatol 48, 357-363. doi: 10.1111/jmp.12437

Altizer, S., Nunn, C.L., Thrall, P.H., Gittleman, J.L., Antonovics, J., Cunningham, A.A., Dobson, A.P., Ezenwa, V., Jones, K.E., Pedersen, A.B., Poss, M., Pulliam, J.R.C., 2003. Social Organization and Parasite Risk in Mammals: Integrating Theory and Empirical Studies. Annual Review of Ecology, Evolution, and Systematics 34, 517-547. doi: 10.1146/annurev.ecolsys.34.030102.151725

Amato, K.R., Yeoman, C.J., Kent, A., Righini, N., Carbonero, F., Estrada, A., Gaskins, H.R., Stumpf, R.M., Yildirim, S., Torralba, M., Gillis, M., Wilson, B.A., Nelson, K.E., White, B.A., Leigh, S.R., 2013. Habitat degradation impacts black howler monkey (*Alouatta pigra*) gastrointestinal microbiomes. ISME J. 7, 1344-1353. doi: 10.1038/ismej.2013.16

Atmoko, T., Mardiastuti, A., Bismark, M., Prasetyo, L.B., Iskandar, E., 2021. Land cover and Proboscis monkey habitats in Berau Delta, East Kalimantan. IOP Conference Series: Earth and Environmental Science 739. doi: 10.1088/1755-1315/739/1/012062

Barelli, C., Albanese, D., Stumpf, R.M., Asangba, A., Donati, C., Rovero, F., Hauffe, H.C., 2020. The gut microbiota communities of wild arboreal and ground-feeding tropical primates are affected differently by habitat disturbance. mSystems 5. doi: 10.1128/mSystems.00061-20

Becker, D.J., Streicker, D.G., Altizer, S., 2018. Using host species traits to understand the consequences of resource provisioning for host-parasite interactions. J Anim Ecol 87, 511-525. doi: 10.1111/1365-2656.12765

Bernard, H., Mohammad-Shom, S., Kulanthavelu, M., Sha, J.C.M., Malim, T.P., Abram, N.K., Matsuda, I., 2025. Monitoring the population and distribution of the proboscis monkey (*Nasalis larvatus*) in the Klias Peninsula, Sabah, Borneo, Malaysia: insights from an 18-year study. Primates 66, 277-294. doi: 10.1007/s10329-025-01183-7

Bürkner, P.-C., 2018. Advanced Bayesian Multilevel Modeling with the R Package brms. The R Journal 10, 395-411. doi:

Bush, A.O., Lafferty, K.D., Lotz, J.M., Shostak, A.W., 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. The Journal of Parasitology 83. doi: 10.2307/3284227

Chan, A.H.E., Kusolsuk, T., Watthanakulpanich, D., Pakdee, W., Doanh, P.N., Yasin, A.M., Dekumyoy, P.,
Thaenkham, U., 2023. Prevalence of *Strongyloides* in Southeast Asia: a systematic review and meta-
analysis with implications for public health and sustainable control strategies. Infect Dis Poverty 12, 83.
doi: 10.1186/s40249-023-01138-4

Chapman, A.C., Rothman, J.M., Hodder, S.A.M., 2009. Can parasite infections be a selective force
influencing primate group size? A test with red colobus, in: Huffman, M.A., Chapman, C.A. (Eds.),
Primate Parasite Ecology: The Dynamics and Study of Host-Parasite Relationships. Cambridge
University Press, Cambridge, pp. 423-440.

Chapman, C.A., Gillespie, T.R., Goldberg, T.L., 2005. Primates and the Ecology of their Infectious Diseases:
How will Anthropogenic Change Affect Host-Parasite Interactions? Evolutionary Anthropology: Issues,
News, and Reviews 14, 134-144. doi: 10.1002/evan.20068

Cooper, N., Nunn, C.L., 2013. Identifying future zoonotic disease threats: Where are the gaps in our
understanding of primate infectious diseases? Evol Med Public Health 2013, 27-36. doi:
10.1093/emph/eot001

Feilen, K.L., Marshall, A.J., 2014. Sleeping site selection by proboscis monkeys (*Nasalis larvatus*) in West
Kalimantan, Indonesia. Am J Primatol 76, 1127-1139. doi: 10.1002/ajp.22298

Feilen, K.L., Marshall, A.J., 2020. Responses to spatial and temporal variation in food availability on the
feeding ecology of proboscis monkeys (*Nasalis larvatus*) in West Kalimantan, Indonesia. Folia Primatol
(Basel), 1-18. doi: 10.1159/000504362

Frias, L., Hasegawa, H., Chua, T.H., Sipangkui, S., Stark, D.J., Salgado-Lynn, M., Goossens, B., Keuk, K.,
Okamoto, M., MacIntosh, A.J.J., 2021. Parasite community structure in sympatric Bornean primates.
International Journal for Parasitology. doi: 10.1016/j.ijpara.2021.03.003

Frias, L., Stark, D.J., Salgado Lynn, M., Nathan, S., Goossens, B., Okamoto, M., MacIntosh, A.J.J., 2019.
Molecular characterization of nodule worm in a community of Bornean primates. Ecology and
Evolution. doi: 10.1002/ece3.5022

Gillespie, T.R., Chapman, C.A., 2006. Prediction of parasite infection dynamics in primate metapopulations
based on attributes of forest fragmentation. Conserv Biol 20, 441-448. doi: 10.1111/j.1523-
1739.2006.00290.x

Hasegawa, H., Frias, L., Peter, S., Hasan, N.H., Stark, D.J., Lynn, M.S., Sipangkui, S., Goossens, B.,
Matsuura, K., Okamoto, M., Macintosh, A.J.J., 2020. First description of male worms of *Enterobius*

483 (*Colobenterobius*) *serratus* (Nematoda: Oxyuridae), the pinworm parasite of proboscis monkeys.

484 Zootaxa 4722, 287-294. doi: 10.11646/zootaxa.4722.3.6

485 Hayakawa, T., Nathan, S., Stark, D.J., Saldivar, D.A.R., Sipangkui, R., Goossens, B., Tuuga, A., Clauss, M.,

486 Sawada, A., Fukuda, S., Imai, H., Matsuda, I., 2018. First report of foregut microbial community in

487 proboscis monkeys: are diverse forests a reservoir for diverse microbiomes? Environmental

488 Microbiology Reports 10, 655-662. doi: 10.1111/1758-2229.12677

489 Hopkins, M.E., Nunn, C.L., 2007. A global gap analysis of infectious agents in wild primates. Diversity and

490 Distributions 13, 561-572. doi: 10.1111/j.1472-4642.2007.00364.x

491 Jose, L., Lee, W., Hanya, G., Tuuga, A., Goossens, B., Tangah, J., Matsuda, I., Kumar, V.S., 2024. Gut

492 microbial community in proboscis monkeys (*Nasalis larvatus*): implications for effects of geographical

493 and social factors. Royal Society Open Science 11. doi: 10.1098/rsos.231756

494 Kappeler, P.M., Cremer, S., Nunn, C.L., 2015. Sociality and health: impacts of sociality on disease

495 susceptibility and transmission in animal and human societies. Philos Trans R Soc Lond B Biol Sci 370.

496 doi: 10.1098/rstb.2014.0116

497 Klaus, A., Strube, C., Roper, K.M., Radespiel, U., Schaarschmidt, F., Nathan, S., Goossens, B.,

498 Zimmermann, E., 2018. Fecal parasite risk in the endangered proboscis monkey is higher in an

499 anthropogenically managed forest environment compared to a riparian rain forest in Sabah, Borneo.

500 PLoS One 13, e0195584. doi: 10.1371/journal.pone.0195584

501 Klaus, A., Zimmermann, E., Röper, K.M., Radespiel, U., Nathan, S., Goossens, B., Strube, C., 2017. Co-

502 infection patterns of intestinal parasites in arboreal primates (proboscis monkeys, *Nasalis larvatus*) in

503 Borneo. International Journal for Parasitology: Parasites and Wildlife. doi:

504 10.1016/j.ijppaw.2017.09.005

505 Knauf, S., Jones-Engel, L., 2020. Neglected Diseases in Monkeys: From the Monkey-Human Interface to

506 One Health. Springer Cham, Switzerland AG.

507 Lee, W., Hayakawa, T., Kiyono, M., Yamabata, N., Hanya, G., 2019. Gut microbiota composition of

508 Japanese macaques associates with extent of human encroachment. Am J Primatol 81, e23072. doi:

509 10.1002/ajp.23072

510 Lhota, S., Sha, J.C.M., Bernard, H., Matsuda, I., 2019. Proboscis monkey conservation: beyond the science,

511 in: Behie, A.M., Teichroeb, J.A., Malone, N.M. (Eds.), Primate Research and Conservation in the

512 Anthropocene. Cambridge University Press, Cambridge pp. 182-196.

Matsuda, I., Kubo, T., Tuuga, A., Higashi, S., 2010a. A Bayesian analysis of the temporal change of local

density of proboscis monkeys: implications for environmental effects on a multilevel society. American

Journal of Physical Anthropology 142, 235-245. doi: 10.1002/ajpa.21218

Matsuda, I., Murai, T., Grueter, C.C., Tuuga, A., Goossens, B., Bernard, H., Yahya, N.K., Orozco-terWengel,

P., Salgado-Lynn, M., 2024. The multilevel society of proboscis monkeys with a possible patrilineal

basis. Behavioral Ecology and Sociobiology 78. doi: 10.1007/s00265-023-03419-2

Matsuda, I., Nakabayashi, M., Otani, Y., Yap, S.W., Tuuga, A., Wong, A., Bernard, H., Wich, S.A., Kubo,

T., 2019. Comparison of plant diversity and phenology of riverine and mangrove forests with those of

the dryland forest in Sabah, Borneo, Malaysia, in: Nowak, K., Barnett, A.A., Matsuda, I. (Eds.), Primates

in Flooded Habitats: Ecology and Conservation. Cambridge University Press, Cambridge, pp. 15-28.

Matsuda, I., Tuuga, A., Akiyama, Y., Higashi, S., 2008. Selection of river crossing location and sleeping site

by proboscis monkeys (*Nasalis larvatus*) in Sabah, Malaysia. American Journal of Primatology 70,

1097-1101. doi: 10.1002/ajp.20604

Matsuda, I., Tuuga, A., Bernard, H., 2011. Riverine refuging by proboscis monkeys (*Nasalis larvatus*) and

sympatric primates: Implications for adaptive benefits of the riverine habitat. Mammalian Biology 76,

165-171. doi: 10.1016/j.mambio.2010.03.005

Matsuda, I., Tuuga, A., Hashimoto, C., Bernard, H., Yamagiwa, J., Fritz, J., Tsubokawa, K., Yayota, M.,

Murai, T., Iwata, Y., Clauss, M., 2014. Faecal particle size in free-ranging primates supports a

'rumination' strategy in the proboscis monkey (*Nasalis larvatus*). Oecologia 174, 1127-1137. doi:

10.1007/s00442-013-2863-9

Matsuda, I., Tuuga, A., Higashi, S., 2009a. The feeding ecology and activity budget of proboscis monkeys.

American Journal of Primatology 71, 478-492. doi: 10.1002/ajp.20677

Matsuda, I., Tuuga, A., Higashi, S., 2009b. Ranging behavior of proboscis monkeys in a riverine forest with

special reference to ranging in inland forest. International Journal of Primatology 30, 313-325. doi:

10.1007/s10764-009-9344-3

Matsuda, I., Tuuga, A., Higashi, S., 2010b. Effects of water level on sleeping-site selection and inter-group

association in proboscis monkeys: why do they sleep alone inland on flooded days? Ecological Research

25, 475-482. doi: 10.1007/s11284-009-0677-3

Modrý, D., Pafčö, B., Petrzělková, K.J., Hasegawa, H., 2018. Parasites of Apes: an Atlas of Coproscopic

Diagnostics. Chimaira, Frankfurt.

Nguyen, N., Fashing, P.J., Boyd, D.A., Barry, T.S., Burke, R.J., Goodale, C.B., Jones, S.C., Kerby, J.T.,
Kellogg, B.S., Lee, L.M., Miller, C.M., Nurmi, N.O., Ramsay, M.S., Reynolds, J.D., Stewart, K.M.,
Turner, T.J., Venkataraman, V.V., Knauf, Y., Roos, C., Knauf, S., 2015. Fitness impacts of tapeworm
parasitism on wild gelada monkeys at Guassa, Ethiopia. Am J Primatol 77, 579-594. doi:
10.1002/ajp.22379

Nunn, C.L., 2012. Primate disease ecology in comparative and theoretical perspective. Am J Primatol 74,
497-509. doi: 10.1002/ajp.21986

Nunn, C.L., Jordan, F., McCabe, C.M., Verdolin, J.L., Fewell, J.H., 2015. Infectious disease and group size:
more than just a numbers game. Philos Trans R Soc Lond B Biol Sci 370. doi: 10.1098/rstb.2014.0111

Nunn, C.L., Thrall, P.H., Leendertz, F.H., Boesch, C., 2011. The spread of fecally transmitted parasites in
socially-structured populations. PLoS One 6, e21677. doi: 10.1371/journal.pone.0021677

Radespiel, U., Schaber, K., Kessler, S.E., Schaarschmidt, F., Strube, C., 2015. Variations in the excretion
patterns of helminth eggs in two sympatric mouse lemur species (*Microcebus murinus* and *M.*
ravelobensis) in northwestern Madagascar. Parasitol Res 114, 941-954. doi: 10.1007/s00436-014-4259-
0

Rahman, F.G., Nurliani, A., Rezeki, A., Rusmiati, R., 2023. Identification of intestine parasite worms eggs
in feces proboscis monkey (*Nasalis larvatus*). Proceeding of International Conference on Biology
Education, Natural Science, and Technology 1, 541-549. doi:

Ramadhani, M., Nurhidayah, N., Kurniawati, D.A., Purmadi, M., Rafiq, F.A., Kusumaningtyas, E.,
Endrawati, D., Hidayatik, N., Prastiya, R.A., Mariya, S.S., Mufasirin, Hastutiek, P., 2024.
Gastrointestinal helminths of captive proboscis monkey (*Nasalis larvatus*) in Surabaya zoo, Indonesia.
J Med Primatol 53, e12719. doi: 10.1111/jmp.12719

Rifkin, J.L., Nunn, C.L., Garamszegi, L.Z., 2012. Do animals living in larger groups experience greater
parasitism? A meta-analysis. Am Nat 180, 70-82. doi: 10.1086/666081

Rivero, J., Cutillas, C., Callejon, R., 2020. *Trichuris trichiura* (Linnaeus, 1771) From Human and Non-
human Primates: Morphology, Biometry, Host Specificity, Molecular Characterization, and Phylogeny.
Front Vet Sci 7, 626120. doi: 10.3389/fvets.2020.626120

Rose, J.H., Small, A.J., 1980. Observations on the development and survival of the free-living stages of
Oesophagostomum dentatum both in their natural environments out-of-doors and under controlled
conditions in the laboratory. Parasitology 81, 507-517. doi: 10.1017/s0031182000061898

Rushmore, J., Bisanzio, D., Gillespie, T.R., 2017. Making New Connections: Insights from Primate-Parasite

Networks. Trends Parasitol 33, 547-560. doi: 10.1016/j.pt.2017.01.013

Salgado-Lynn, M., 2010. Primate viability in a fragmented landscape: genetic diversity and parasite burden

o f long-tailed macaques and proboscis monkeys in the Lower Kinabatangan Floodplain, Sabah,

Malaysia. Cardiff University, Cardiff.

Senanayake, N.S., Boyle, L., O'Driscoll, K., Menant, O., Butler, F., 2025. Effects of season, age and parasite

management practices on gastro - intestinal parasites in pigs kept outdoors in Ireland. Ir Vet J 78, 12.

doi: 10.1186/s13620-025-00297-0

Sengupta, M.E., Thapa, S., Thamsborg, S.M., Mejer, H., 2016. Effect of vacuum packing and temperature

on survival and hatching of strongyle eggs in faecal samples. Vet Parasitol 217, 21-24. doi:

10.1016/j.vetpar.2015.12.014

Silk, J.B., 2007. The adaptive value of sociality in mammalian groups. Philos Trans R Soc Lond B Biol Sci

362, 539-559. doi: 10.1098/rstb.2006.1994

Snaith, T.V., Chapman, C.A., Rothman, J.M., Wasserman, M.D., 2008. Bigger groups have fewer parasites

and similar cortisol levels: a multi-group analysis in red colobus monkeys. Am J Primatol 70, 1072-

1080. doi: 10.1002/ajp.20601

Vitone, N.D., Altizer, S., Nunn, C.L., 2004. Body size, diet and sociality influence the species richness of

parasitic worms in anthropoid primates. Evolutionary Ecology Research 6, 183-199. doi:

Wilcove, D.S., Giam, X., Edwards, D.P., Fisher, B., Koh, L.P., 2013. Navjot's nightmare revisited: logging,

agriculture, and biodiversity in Southeast Asia. Trends Ecol Evol 28, 531-540. doi:

10.1016/j.tree.2013.04.005

Yalcindag, E., Stuart, P., Hasegawa, H., Streit, A., Dolezalova, J., Morrogh-Bernard, H., Cheyne, S.M.,

Nurcahyo, W., Foitova, I., 2021. Genetic characterization of nodular worm infections in Asian Apes.

Sci Rep 11, 7226. doi: 10.1038/s41598-021-86518-2

Table and figure legends

Table 1. Summary of the fitted models.

Table 2. Comparison between the parasite species detected and parasite prevalence (%) in fecal

samples of proboscis monkeys in each study.

Fig. 1. Posterior distributions of the differences in parasite abundance for *Trichuris* sp. (top),

S. fuelleborni (middle), and *O. aculeatum* (bottom) in relation to host group type (A). The green

shaded areas represent the posterior distributions, with the black dot indicating the median

estimate and the black horizontal line shown in boldface representing the 99% credible interval.

The vertical dashed line at zero represents no difference in infection rates among the groups.

The predicted probability of abundance for the same parasite groups in relation to host group

size (B) and sampling location represented by the distance from the river mouth (used as a

proxy for disturbance gradient) (C). Credible intervals are set at 89%.

Table 1. Summary of fitted models

	Parameter	Est.	Est. error	95% CI (Lower)	95% CI (Upper)	Rhat	Bulk ESS	Tail ESS
	Random Effects							
	sd (Intercept)	0.92	0.22	0.6	1.44	1	1600	2336
	Fixed Effects							
Trichuris sp.	Intercept	4.32	0.51	3.29	5.32	1	3515	4008
	group type	0.81	0.55	-0.27	1.93	1	3151	3511
	group size (scaled)	0.16	0.01	0.14	0.19	1	6937	4494
	distance to river (scaled)	-0.7	0.14	-0.97	-0.44	1	2949	3362
	Random Effects							
	sd (Intercept)	5.74	1.67	3.37	9.71	1	2691	3424
	Fixed Effects							
S. fuelleborni	Intercept	-1.68	3.25	-8.21	4.77	1	3101	3670
	group type	-1.09	3.7	-8.69	5.88	1	2916	3602
	group size (scaled)	-1.62	0.12	-1.86	-1.38	1	9301	5682
	distance to river (scaled)	1.03	0.71	-0.35	2.44	1	5057	4881
	Random Effects							
	sd (Intercept)	10.54	3.96	5	20.28	1	2116	3425
	Fixed Effects							
O. aculeatum	Intercept	-5.12	6.63	-18.87	7.67	1	3150	3920
	group type	-0.47	7.58	-16.08	14.32	1	2693	3620
	group size (scaled)	-1.48	0.19	-1.87	-1.11	1	7942	5445
	distance to river (scaled)	3.46	1.4	0.71	6.27	1	3953	4708

614 **Table 2.** Comparison of the parasite species detected and parasite prevalence (%) in fecal samples of proboscis monkeys in each study

	Bekantan Rescue Center Sahabat Bekantan Foundation	Surabaya Zoo	Labuk Bay Proboscis Monkey Sanctuary	Bako National Park	LKWS[1] (Lots 1-7)	LKWS (Lot 6)	LKWS (Lots 1-4)	LKWS (Lot 4)
	(Rahman et al., 2023)	(Ramadhani et al., 2024)	(Klaus et al., 2017; Klaus et al., 2018)	(Adrus et al., 2019)	(Frias et al., 2021)	(Klaus et al., 2017; Klaus et al., 2018)	(Salgado-Lynn, 2010)	(this study)
Forest type	N/A	N/A	Mangrove	Mangrove	Riverine forest	Riverine forest	Riverine forest	Riverine forest
Living-condition	Captive	Captive	Semi-free ranging	Free-ranging	Free-ranging	Free-ranging	Free-ranging	Free-ranging
No. of individuals	3	< 27	< 195	Unknown	Unknown	< ca. 200	Unknown	160
No. of samples	60	20	90	65	91	652	146	160
Cestodes								
Diphyllobotrium sp.	–	–	–	1.5	–	–	–	–
Dipylidium-like morphs	–	–	–	–	–	–	9.6	–
Hymenolepis nana	–	5.0	–	–	–	–	–	–
Hymenolepis sp.	–	–	–	3.1	–	–	–	–
Taenia sp.	–	–	–	–	–	–	28.8	–
Nematodes								
Anatrichosoma spp.	–	–	2.8	–	–	1.4	2.1	–
Ascaris lumbricoides	–	–	2.2	–	–	8.6	–	–

	Bekantan Rescue Center Sahabat Bekantan Foundation (Rahman et al., 2023)	Surabaya Zoo (Ramadhani et al., 2024)	Labuk Bay Proboscis Monkey Sanctuary (Klaus et al., 2017; Klaus et al., 2018)	Bako National Park (Adrus et al., 2019)	LKWS[1] (Lots 1-7) (Frias et al., 2021)	LKWS (Lot 6) (Klaus et al., 2017; Klaus et al., 2018)	LKWS (Lots 1-4) (Salgado-Lynn, 2010)	LKWS (Lot 4) (this study)
Ascaris sp.	–	5.0	–	30.8	–	–	67.1	–
Enterobius (Colobenterobius) serratus	–	–	–	–	1.25	–	–	1.25
Enterobius spp.	–	–	–	–	–	5.5	–	–
Oesophagostomum aculeatum	–	–	–	–	–	–	–	5.0
Oesophagostomum sp.	5.0	–	31.4	33.8	–	22.8	–	–
oxyurid	–	–	–	–	–	–	4.1	–
Physaloptera sp.	–	–	–	3.1	–	–	–	–
Spirurida sp.	–	–	–	–	2.1	–	–	0.63
strongylid	–	–	4.6	–	–	–	45.2	–
Strongylida	–	–	43.3	–	29.6	3.1	–	–
Strongyloides fuelleborni	–	–	–	–	–	–	–	10.6
Strongyloides stercoralis	–	–	–	4.6	–	–	–	–
Strongyloides spp.	–	5.0	65.6	21.5	20.8	32.7	5.5	–

	Bekantan Rescue Center Sahabat Bekantan Foundation (Rahman et al., 2023)	Surabaya Zoo (Ramadhani et al., 2024)	Labuk Bay Proboscis Monkey Sanctuary (Klaus et al., 2017; Klaus et al., 2018)	Bako National Park (Adrus et al., 2019)	LKWS[1] (Lots 1-7) (Frias et al., 2021)	LKWS (Lot 6) (Klaus et al., 2017; Klaus et al., 2018)	LKWS (Lots 1-4) (Salgado-Lynn, 2010)	LKWS (Lot 4) (this study)
Trichostrongylus spp.	10.0	–	36.0	9.2	–	48.5	–	–
Trichuris trichiura	15.0	–	–	44.6	–	–	–	–
Trichuris spp.	–	85.0	21.8–47.1	35.4	80.2	82.1	91.8	78.1
Trematodes								
Fasciola sp.	–	–	–	1.5	–	–	–	–
Schistosoma sp.	–	–	–	1.5	–	–	–	–

[1]Lower Kinabatangan Wildlife Sanctuary

Fig. 1. Posterior distributions of the differences in parasite abundance for *Trichuris* sp. (top), *S. fuelleborni* (middle), and *O. aculeatum* (bottom)
in relation to host group type (A). The green shaded areas represent the posterior distributions, with the black dot indicating the median estimate
and the black bold horizontal line representing the 99% credible interval. The vertical dashed line at zero represents no difference in infection rates
between groups. Predicted probability of abundance for *the same parasite groups* in relation to host group size (B) and sampling location
represented by the distance from the river mouth (used as a proxy for disturbance gradient) (C). Credible intervals are set at 89%.

623
624