The RNA splicing factor PRPF8 is required for left-right organiser cilia function and

determination of cardiac left-right asymmetry via regulation of *Arl13b* splicing

Fangfei Jiang[1]†, Michael Boylan[1]†, Dale W. Maxwell[1,2]†, Wasay Mohiuddin Shaikh

Qureshi[1], Charlie F. Rowlands[1], Gennadiy Tenin[1,9], Karen Mitchell[1], Louise A. Stephen[1],

Elton J. R. Vasconcelos[3], Dapeng Wang[3], Tong Chen[4], Junzhe Zha[5], Jingshu Liu[1], Nouf

Althali[1,6], Dragos V. Leordean[1], Meurig T. Gallagher[7], Basudha Basu[8], Katarzyna

Szymanska[8], Advait Veeraghanta[2], Bernard Keavney[9-11], Martin J. Humphries[5], Jamie

Ellingford[1,12], David Smith[7], Colin A. Johnson[8], Raymond T. O'Keefe[1], Sudipto Roy[2,13],

Kathryn E. Hentges[1]*

Author Affiliations:

1. Division of Evolution, Infection and Genomics, BHF Manchester Centre of Research
Excellence, School of Biological Sciences, Faculty of Biology, Medicine, and Health,
Manchester Academic Health Science Centre, University of Manchester, Manchester, M13
9PT, UK
17

18   2. Institute of Molecular and Cell Biology (IMCB), Agency for Science, Technology and
19   Research (A*STAR), Proteos, 61 Biopolis Drive, Singapore 138673, Singapore
20

21   3. LeedsOmics, University of Leeds, Leeds, LS2 9JT, UK
22

23   4. Division of Molecular and Cellular Function, School of Biological Sciences, Faculty of
24   Biology, Medicine, and Health, Manchester Academic Health Science Centre, University of
25   Manchester, Manchester, M13 9PT, UK
26

5. Division of Cell Matrix Biology and Regenerative Medicine, Manchester Cell-Matrix

Centre, School of Biological Sciences, Faculty of Biology, Medicine, and Health, Manchester

Academic Health Science Centre, University of Manchester, Manchester, M13 9PT, UK

6. Biology Department, Science College, King Khalid University, Abha, Saudi Arabia.

7. Centre for Systems Modelling and Quantitative Biomedicine, University of Birmingham,

Edgbaston, Birmingham, UK and Centre for Human Reproductive Science, University of

Birmingham, Birmingham, UK

8. Division of Molecular Medicine, Leeds Institute of Medical Research, University of Leeds,

Leeds, LS9 7TF, UK

9. Division of Cardiovascular Sciences, BHF Manchester Centre of Research Excellence,

School of Medical Sciences, Faculty of Biology, Medicine, and Health, The University of

Manchester, Manchester, M13 9PL, UK

10. Manchester Heart Centre, Manchester University NHS Foundation Trust, Manchester

Academic Health Science Centre, UK

11. NIHR Manchester Biomedical Research Centre, UK

112. Genomics England, London, E14 4AB, UK

13. Department of Paediatrics, National University of Singapore, 1E Kent Ridge Road,

Singapore 119228, Singapore

† = equal contribution
*= author for correspondence, Tel +44 161 275 1490, email:

Kathryn.hentges@manchester.ac.uk

Short title: PRPF8 mutation alters cilia function and organ laterality

Present addresses are listed in supplemental table 6.

Abstract

Cilia function in the left-right organizer (LRO) is critical for determining internal organ

asymmetry in vertebrates. To further understand the genetics of left-right asymmetry, we

isolated a mouse mutant with laterality defects, *l11Jus27,* from a random mutagenesis screen.

l11Jus27 mutants carry a missense mutation in the pre-mRNA processing factor, *Prpf8*.

cephalophŏnus (cph) mutant zebrafish, carrying a protein truncating mutation in *prpf8*,

phenocopy the laterality defects of *l11Jus27* mutants. *Prpf8* mutant mouse and fish embryos

have increased expression of an alternative transcript encoding the cilium-associated protein,

ARL13B, that lacks exon 9. In zebrafish, over-expression of the *arl13b* transcript lacking exon

9 perturbed cilium formation and caused laterality defects. The shorter ARL13B protein

isoform lacked interactions with intraflagellar transport proteins. Our data suggest that PRPF8

plays a prominent role in LRO cilia by through the regulation of alternative splicing of

ARL13B, thus uncovering a new mechanism for cilia-linked developmental defects.

Key words:

PRPF8, cilia; left-right axis, splicing, IFT, ARL13B, mutagenesis screen, ciliopathy, retinitis
pigmentosa, embryonic development

Introduction

Cilia are specialised hair-like organelles required for many cellular functions (1). Motile cilia produce a rhythmic waving or beating motion, whereas non-motile cilia act as sensory antennae for cells (1, 2). Diseases caused by cilia dysfunction are collectively known as ciliopathies. Mutations in the wide range of genes required for cilia function cause ciliopathies, with most phenotypes displaying both genetic heterogeneity and variable expressivity (3). In humans, mutations in several spliceosomal pre-mRNA processing factor proteins cause autosomal dominant Retinitis Pigmentosa (RP) (4-6), a ciliopathy causing progressive retinal atrophy and loss of sight with age. RP-causing mutations in Pre-mRNA Processing Factor 8 (PRPF8) produce multiple splicing defects (5, 7-11). However, PRPF8 may have a more direct role in cilia functionality. PRPF8 localises to the basal body of primary cilia, and knocking down PRPF8 in human and mouse cells (12, 13), zebrafish embryos (13) and *C. elegans* (14), inhibits ciliogenesis.

In vertebrate embryogenesis, specialised motile cilia in the LRO play a critical role in left-right axis specification (2, 15). In mice, laterality is established at the ventral node (2), the mammalian LRO. The node is a teardrop shaped pit at the posterior end of the notochord with motile cilia; these cilia generate a leftward flow of fluid within the node cavity thought to be detected by immotile cilia on adjacent perinodal 'crown' cells (16-20). In response, expression of the genes *Cerl2* and *Nodal*, becomes restricted preferentially on the right and left side of the node, respectively (21-23). Nodal in the left lateral plate mesoderm (LPM) induces expression of the Nodal inhibitors *Lefty1* and *Lefty2* (24, 25) at the midline and left LPM, respectively (26). Nodal in the LPM also induces expression of the transcription factor *Pitx2* (27). *Pitx2* is considered to be the effector of Nodal signalling, and is required for the correct positioning and

morphogenesis of organs including the heart, lungs, gut and blood vessels (28-30), but is

dispensable for heart looping directionality (reviewed in (31)) .

A recessive lethal mouse mutant with laterality defects was isolated from a random chemical

mutagenesis screen: the *l11Jus27* mouse (32, 33). The causative mutation in this mouse is an

asparagine (N) to serine (S) missense mutation at residue 1531 in *Prpf8*, the mouse orthologue

of human *PRPF8*. *Prpf8$^{N1531S/N1531S}$* embryos (hereafter referred to as *Prpf8^{N1531S}* mutants or

Prpf8^{N1531S} hom) have impaired LRO cilia formation and motility. Zebrafish *cph* mutants

(carrying a Prpf8 truncation mutation) phenocopy laterality defects seen in *Prpf8^{N1531S}* mutant

mouse embryos. Moreover, modelling the analogous *Prpf8^{N1531S}* mutation in yeast resulted in

splicing defects. Bulk RNA-seq analysis of *Prpf8^{N1531S}* mutant mouse embryos showed

increased skipping of exon 9 in the transcript of the Arf-like GTPase *Arl13b;* the ARL13B

protein is specifically localised to the ciliary membrane (34, 35) where it functions in regulating

cilia length, and thus impacts upon beat pattern and flow (36-38), as well as ciliary trafficking

via interactions with intraflagellar transport (IFT) proteins (39, 40). Importantly, over-

expression of the shorter a*rl13b* transcript (lacking exon 9) in wild type zebrafish re-capitulated

many aspects of the *Prpf8^{N1531S}* mutant phenotype. Additionally, the ARL13B protein lacking

the region encoded by exon 9 showed reduced binding to IFT partners. Combined, these results

suggest that altering *Arl13b* alternative transcript ratios may be the mechanistic basis for the

laterality defects observed in *Prpf8* mutants. The data presented here reveal specific functions

for PRPF8 in cilia function and embryonic patterning, which may also provide new insights

into the role of PRPF8 in disease pathology in patients with RP.

Results

Phenotypic analysis of *l11Jus27* mouse mutants

The *l11Jus27* mouse line was isolated from a random chemical mutagenesis screen, which

employed a balancer chromosome to facilitate mapping of mutations (32). *l11Jus27*

heterozygotes are viable and fertile. At E8.25, *l11Jus27* homozygous embryos appear grossly

normal (Supplemental Fig. 1a-b) but are developmentally delayed (Supplemental Fig. 1c). At

E9.5 the *l11Jus27* homozygotes have an open neural tube, distended heart tube, and a high

incidence of reversed cardiac looping (Fig. 1a). At E10.5 less severely affected mutants have

milder cardiac defects and developmental delay (Supplemental Fig. 1d-f). More severely

delayed embryos have also failed to undergo chorioallantoic fusion (Supplemental Fig. 1f).

l11Jus27 homozygous embryos have impaired yolk sac vascularisation at E9.5 (Supplemental

Fig. 1g-h) and frequently do not have patent umbilical vessels (Supplemental Fig 1i-j),

probably leading to the failure in chorioallantoic fusion observed in mutant embryos at E10.5

(Supplemental Fig 1j; penetrance reported in Supplemental Table 1). The majority of

homozygous mutant embryos died by E11.5, and none were isolated after E12.5.

The *l11Jus27* line contains a *Prpf8^{N1531S}* mutation

The *l11Jus27* mutant mouse line was isolated from a balancer chromosome mutagenesis

screen; therefore the causative mutation was known to reside in a 24cM region of mouse

chromosome 11 (32). Meiotic mapping further refined the *l11Jus27* candidate interval

(Supplemental Fig. 1k) (41). Genome sequencing of an E10.5 homozygous mutant embryo

revealed a novel point mutation of an A to G transition within the genetic candidate interval at

position chr11: 75,391,978 (mm39), which produces a N1531S missense mutation in *Prpf8*.

This sequence alteration was confirmed in an additional 3 homozygous mutant embryos and

was not present in wild type C57BL/6 or 129S5 animals (Fig. 1b). Thus, we concluded the

l11Jus27 line carries a PRPF8 N1531S missense mutation, and *l11Jus27* homozygous embryos

will now be referred to as *Prpf8* hom. We confirmed by quantitative RT-PCR that the N1531S

mutation does not affect *Prpf8* transcript levels (Fig. 1c). PRPF8 is highly conserved (42), and

residue N1531 is conserved between multiple eukaryotes (Fig. 1d). The N1531S amino acid

substitution is in a linker region near the endonuclease-like domain (Fig. 1e; (43)) and is

predicted to be damaging by PolyPhen and SIFT (44, 45) in both mouse and human proteins.

At E10.5, mutants appeared to display a reduction in PRPF8 protein levels, but this was not

significantly different (Fig. 1f-g; one-way ANOVA p=0.4212)(12). To test if the PRPF8

mutation altered protein stability, we expressed an N-terminal FLAG epitope-tagged PRPF8

construct in mammalian cells (Fig. 1h) and performed a cycloheximide assay to determine the

PRPF8 degradation rate. We detected a reduced level of epitope tagged PRPF8^{N1531S} protein

compared to wild type protein at each time point in the assay, but the overall degradation rate

of the N1531S mutant was similar to the wild type construct (Wilcoxon test p=0.125; Fig. 1i-

j), and thus did not indicate significant evidence of increased protein degradation.

172   Nodal cilia are present but dysmotile in *Prpf8* ^{N1531S} mutants

Heart looping defects are associated with ciliopathies (46, 47). Scanning electron microscopy

(SEM) revealed that control and *Prpf8^{N1531S}* mutant embryos possessed abundant nodal cilia at

the 2-3 somite stage (Fig. 2a-d). Control embryos had stereotypical pit shaped nodes (Fig. 2a,

n=7), whereas *Prpf8^{N1531S}* mutant embryonic nodes appeared unusually flat (Fig. 2b, *n*=6/8),

suggesting that node morphogenesis is disrupted in *Prpf8^{N1531S}* mutant embryos. While there

was a trend towards fewer cilia in *Prpf8^{N1531S}* mutant embryos, this trend was not statistically

significant (Fig. 2e; unpaired t-test of late head fold (LHF) stage data, p=0.1934; for 2-3 somite

data p=0.1494). Transmission electron microscopy (TEM) analysis of nodal cilia did not reveal

any overt differences between homozygous *Prpf8^{N1531S}* mutant embryos and controls

(Supplemental Fig 2). Because mutations in the spliceosome have been found to affect the rate

of cell division (48), we performed immunofluorescence staining for phospho-histone H3 to

identify cells in mitosis. There was no difference in the percentage of dividing cells between

genotypes (Fig. 2f; unpaired t-test p=0.7918). Using acetylated tubulin as a marker of cilia (Fig.

2g-h), we found that node cilia length in homozygous $Prpf8^{N1531S}$ mutant embryos was

significantly shorter than in heterozygous control embryos (Fig. 2i, unpaired t-test p=0.0408).

LRO fluid flow generation is related to cilia length (38). Therefore, we evaluated whether flow

was established correctly in $Prpf8^{N1531S}$ mutant embryo nodes. We placed microbeads in the

node of embryos mounted in a chamber slide to detect the direction of nodal flow. While almost

all heterozygotes established flow towards the left of the node (n=5/6; Fig. 2j; Supplemental

Movie File 1), almost all $Prpf8^{N1531S}$ mutants failed to establish any directional fluid flow

(n=5/6; Fig. 2k); instead the microbeads showed random Brownian motion (Supplemental

Movie file 2). We next investigated whether cilia motility was impaired in mutant embryos.

Directly visualising cilia movement using high-speed videomicroscopy revealed that wild type

embryos had strongly beating cilia (n=5 Supplemental Movie File 3), as did $Prpf8^{N1531S/+}$

embryos (n=4/4). However, $Prpf8^{N1531S}$ homozygotes had very few motile cilia (n=3;

Supplemental Movie File 4). We quantified the number of motile cilia (Fig. 2l) and found

significant differences between the three genotypes (p=0.0058, Welch's ANOVA). Posthoc

testing revealed that the number of motile cilia in $Prpf8^{N1531S}$ homozygotes was significantly

lower compared to $Prpf8^{+/+}$ (p=0.0222) and $Prpf8^{N1531S/+}$ embryos (p=0.0487), but not between

$Prpf8^{N1531S/+}$ and $Prpf8^{+/+}$ embryos.

PRPF8 has been reported to localise to the base of cilia (12-14). To determine if the $Prpf8^{N1531S}$

missense mutation affected protein localisation, we used immunofluorescence to analyse node

cilia. We confirmed that PRPF8 protein can be detected at the base of some cilia in

heterozygous (Supplemental Fig. 3 a-c) and $Prpf8^{N1531S}$ homozygous mutant embryos within

the node (Supplemental Fig. 3 g-i). A transverse projection of the Z-stack of confocal images

confirmed that the node in heterozygous embryos formed a pit like structure (Supplemental 3d-

f) but in homozygotes the node was flatter (Supplemental 3 j-l). These findings suggest that

$Prpf8^{N1531S}$ homozygous mutants have defects in node structure, cilia motility, and nodal fluid

flow that are consistent with downstream defects in L-R axis formation.

Laterality gene expression analysis reveals pathway defects in $Prpf8^{N1531S}$ mutants

We next determined whether the L-R axis pathway was disturbed by the defects in the node.

We performed whole-mount *in situ* hybridisation for several genes that are important for L-R

axis formation and found that $Prpf8^{N1531S}$ mutant embryos had highly aberrant laterality gene

expression patterns. *Shh*, needed for L-R axis establishment (49), was expressed in the midline

and node of both control and $Prpf8^{N1531S}$ mutant embryos (Fig. 3a-b), though half of $Prpf8^{N1531S}$

mutants displayed fainter and discontinuous midline *Shh* expression (*n*=3/6) *Cerl2* (Fig. 3c-e)

and *Nodal* (Fig. 3f-h) expression was highly variable, with normal, reversed, symmetric and

absent expression patterns observed for both genes in the mutants. The LPM of $Prpf8^{N1531S}$

mutants usually lacked *Nodal* expression (*n*=19/21), and when *Nodal* was expressed, the

expression domain did not include the anterior LPM. Accordingly, both *Lefty1* and *Lefty2* were

not expressed in $Prpf8^{N153S1}$ mutant embryos (*n*=10/10) (Fig 3i-j). *Pitx2* expression in the LPM

and subsequent secondary heart field (50), which is induced by Nodal activity, was variable,

with normal (n=13/38), reversed (n=8/38), bilateral (n=8/38) and absent (n=9/38) expression

in $Prpf8^{N153S1}$ mutant embryos (Fig 3k-p). *Pitx2* is normally expressed on the same side as the

developing ventricle. We quantified both *Pitx2* expression sidedness and heart looping

direction in mutant embryos (Fig 3q). In embryos where the sidedness of *Pitx2* expression was

unambiguous, 29% (n=6/21) expressed *Pitx2* on the opposite side of the embryo from the

inflow tract of the heart (Fig 3r). These data led us to hypothesise that the *Prpf8^{N1531S}* mutants

with reversed cardiac looping exhibited heterotaxy rather than *situs inversus*. To test this

hypothesis, we examined the expression of *Barx1* in the developing stomach (51), which is

found left of the embryonic midline. Both control and mutant embryos, including those with

reversed cardiac looping (n=6), maintained *Barx1* expression left of the midline, (Fig 3s-t),

supporting the hypothesis that *Prpf8^{N1531S}* mutant embryos model heterotaxy, not *situs*

inversus. The early lethality of *Prpf8^{N1531S}* mutant embryos precluded examining the laterality

of other organ systems.

Phenocopy of *Prpf8 ^{N1531S}* laterality defects in *prpf8* mutant zebrafish

To evaluate if laterality defects present in the *Prpf8^{N1531S}* mouse showed evolutionary

conservation, we examined the ENU mutant zebrafish strain, *cph* (52). This mutant line

carries an E396* substitution in Prpf8, predicting a severely truncated protein (52). Whereas

wild type (Fig. 4a) and heterozygous embryos are phenotypically normal, *cph* homozygous

mutant embryos displayed a curled down body axis and delayed heart looping (Fig. 4b). *cph*

homozygotes displayed randomised cardiac looping: 27% of *cph* mutants had correctly

looped, 42% unlooped, and 31% reversed looped hearts, compared to 97%, 1% and 2%,

respectively, for wild type fish (Fig. 4c).

Laterality gene expression was altered in *cph* homozygous mutant zebrafish in a manner

resembling *Prpf8^{N1531S}* mutant mouse embryos. Homozygous *cph* mutant embryos had absent

or mis-localised expression of *lefty2* (*lft2*) (Fig. 4d-f) and *southpaw* (*spaw*), a zebrafish *Nodal*

homologue (Fig. 4g-i). Additionally, *dand5,* the zebrafish *Cerl2* homologue, which is

normally expressed more prominently on the right side of the embryo (Fig. 4j), displayed

altered expression patterns in *cph* embryos (Fig. 4k-l). The range of abnormal expression

patterns amongst mutant embryos for these key laterality genes is reminiscent of expression

defects found in *Prpf8^{N1531S}* mouse mutants. Overall, expression defects and heart looping

abnormalities found in *cph* zebrafish suggest that the role of PRPF8 in laterality

establishment and heart looping is conserved amongst vertebrates.

We next evaluated cilia formation and function in Kupffer's vesicle (KV), the zebrafish LRO.

Within KVs of wild type and *cph* zebrafish, motile cilia were present (Fig 4m), and no

significant reduction in overall number of cilia was found in mutants compared to wild type

controls (Fig 4n). However, there was a significant reduction in cilia length in *cph* mutants

relative to wild type controls (Fig 4o; unpaired t-test. P<0.0001), similar to our findings in

Prpf8^{N1531S} mutant mouse embryos (Fig 2i). High speed live video imaging of KVs of wild

type and *cph* homozygous mutants revealed that *cph* embryos exhibited a wide variation in

individual cilia motility patterns. Some cilia were motile, similar to wild type (Supplemental

Movie File 5), but others displayed abnormal motility, and immotile cilia were also observed

(Supplemental Movie File 6). Tracking movement of endogenous particles detectable in KV

fluid revealed a net anticlockwise movement in control KVs (Fig 4p, Supplemental Movie

File 7), but no such net movement was observed in *cph* mutant KVs (Fig 4q, Supplemental

Movie File 8), indicating a disruption in flow.

Effect of the *Prpf8^{N1531S}* mutation on mRNA splicing

Since PRPF8 is a component of the U5 snRNP that forms the spliceosomal catalytic centre, we

investigated whether the *Prpf8^{N1531S}* substitution affected splicing. Yeast cells containing a

N1603S missense substitution in yeast Prp8 (homologue of PRPF8), analogous to the mouse

Prpf8^{N1531S} allele, grew similarly to wild type yeast, indicating that the Prp8^{N1603S} mutation is

viable (Fig. 5a). We used the *ACT1-CUP1* splicing reporter (53) in Prp8^{N1603S} mutants to

determine which step of splicing was affected. Primer extension analysis of the *ACT1-CUP1*

splicing reporter revealed that with a branch site mutation (A259C), increased lariat

intermediate was detected in Prp8^{N1603S} cells compared to Prp8^{WT} (Fig. 5b), indicating the first

step of splicing had occurred. However, with a 3' splice site mutation (A302U), reduced levels

of mature mRNA were present in Prp8^{N1603S} cells (Fig. 5b), indicative of a defect in the second

step of splicing. This assay demonstrated that the Prp8^{N1603S} mutant is a first step allele of Prp8,

as it promoted the first step of splicing at the expense of the second step. We then examined

growth of Prp8^{N1603S} vs Prp8^{WT} yeast colonies with *ACT1-CUP1* reporters with either 3' splice

site, 5' splice site or branch point site mutations (Fig. 5c) on media with increasing

concentrations of copper. Copper tolerance is correlated with the amount of correctly spliced

mature *CUP1* transcript from the *ACT1-CUP1* construct and therefore the efficiency of splicing

(53). Prp8^{N1603S} cells could not tolerate copper as well as Prp8^{WT} cells, regardless of where the

mutation was in the *ACT1-CUP1* reporter construct. These data suggest that the Prpf8^{N1531S}

allele does not suppress any splice site mutations and is likely to be a hypomorphic mutation.

RNA-seq analysis to identify splicing defects in *Prpf8^{N1531S}* mutant mouse embryos

Mutations in PRPF8 are reported to cause mis-splicing in yeast, zebrafish and humans (7, 42,

52, 54). The yeast data we present suggest that the *Prpf8^{N1531S}* mutation could cause mis-

splicing in mouse embryos. We therefore performed bulk RNA-seq analysis on individual

whole embryos at E10.5: wild type (n=3), E10.5 *Prpf8^{N1531S}* mutants with correct heart looping

(n=3), and E10.5 *Prpf8^{N1531S}* mutants with reversed heart looping (n=3). Analysis of differential

exon skipping events between the two strains using the rMATS software package (55)

identified 498 statistically significant mid-penetrance exon skipping events (average difference

in exon inclusion between strains ≥ 20%; FDR-corrected p-value<0.05), across 431 unique

genes (Supplemental Table 2). Gene ontology (GO) enrichment analysis revealed the most

significant enrichment (24/51 genes, 2.43-fold) to be for genes associated with the biological

process "plasma membrane bounded cell projection morphogenesis", consistent with a model

of ciliary dysfunction (Supplemental Table 3).

To further narrow the field of key candidate genes that could underlie the laterality defects

detected in *Prpf8*[N1531S] mutant mice, individual GO terms were attached to differentially spliced

genes identified by rMATS. Of the 431 unique genes harbouring exon skipping events, three -

Arl13b, *Rfx3* and *Rpgrip1l* - were upstream of or directly involved in "determination of

left/right symmetry", although only *Arl13b* was associated with "heart looping" and "left/right

axis specification". Because *Arl13b* is involved in both general L-R axis determination and

cardiac looping, and has known roles in the function of motile and non-motile cilia, the *Arl13b*

exon skipping event was prioritised for additional investigation.

In *Prpf8*[N1531S] homozygous mutants, there is splicing between exons 8 and 10 of the *Arl13b*

gene in approximately 51% of reads mapping to the exon 8 splice donor site, whereas in wild

type embryos the corresponding event is detected in approximately 30% of reads (Fig. 6a).

This exon skipping event deletes the 69bp in-frame exon 9 from the transcript. An analysis of

the *Arl13b* transcript revealed that the omission of exon 9 is predicted to remove a 23 amino

acid segment of the Proline Rich Region from the C-terminal end of ARL13B (Fig. 6b). We

have termed this *Arl13b* transcript, lacking exon 9, the "short isoform", and the transcript

including exon 9 the "long isoform". The short isoform is not annotated in the fish or mouse

UCSC genome browsers, although a single mouse EST clone (DT914850) does have the same

sequence as the *Arl13b* short isoform we have identified. To our knowledge the function of this isoform has not been studied previously. Using a single pair of PCR primers located in exons 8 and 10, we amplified both short and long *Arl13b* isoforms by RT-PCR in wildtype mouse and zebrafish embryos (Fig. 6c). There was an increased ratio of the *Arl13b* short isoform transcript to long transcript in both *Prpf8^{N1531S}* homozygous mouse and *cph* homozygous mutant fish embryos (Fig. 6c). Sequencing of these RT-PCR products revealed the precise deletion of exon 9 from the short isoform product (Fig. 6d). We also detected the *Arl13b* short isoform in multiple adult mouse tissues, most notably in the testes (Fig. 6e-f).

The ARL13B short isoform localises to cilia

To determine if the shift in *ARL13B* isoform usage could affect phenotype, we generated *ARL13B* short or long isoform human transcript sequences with N-terminal GFP fusions. These were transfected into hTERT-RPE1 cells for comparison with endogenous ARL13B. Endogenous ARL13B co-localised with acetylated tubulin at the cilium in hTERT-RPE1 cells (Fig. 7a left). We detected protein produced from the GFP-tagged ARL13B short isoform co-localised with endogenous acetylated tubulin at the cilium in hTERT-RPE1 cells (Fig. 7a centre), demonstrating the short isoform produces a stable protein that is localised correctly. Similar localisation results were seen for the GFP tagged ARL13B long isoform (Fig. 7a right). The short isoform cilia localisation is consistent with the ARL13B cilia localisation signal being within amino acids 347-363 (56), which are included in the short isoform transcript.

Over-expression of the *arl13b* short isoform in zebrafish embryos mimics the *cph* mutant phenotype

We next sought to determine if increased expression of the zebrafish *arl13b* short isoform

causes cilia defects. We injected wild type zebrafish embryos with mRNA encoding a C-

terminal Arl13b-GFP fusion protein of either the short or long isoform. At 12 hours post-

injection, embryos exhibiting GFP were selected for further experiments. Half of the selected

embryos were fixed immediately for immunofluorescence to analyse Arl13b protein

localisation and KV cilia length, while the remaining embryos were monitored until 48 hours

post-injection to investigate phenotypes indicative of cilia dysfunction: cardiac looping and

pericardial oedema. In injected zebrafish embryos both Arl13b long and short isoform GFP

fusion proteins localised to KV cilia, showing co-localisation with acetylated tubulin (Fig 7b).

Measuring KV cilia length for each injection group, and plotting the average cilia length per

KV, revealed no significant change in KV cilia length in wild type fish injected with 80ng/ul

of *arl13b* long isoform GFP-tagged mRNA as compared to uninjected control fish (Fig 7c;

One-way ANOVA test, p=0.9901). However, cilia within the KV of fish injected with the

arl13b short isoform were significantly shorter than those in the uninjected control group (Fig

7c; One-way ANOVA test, p=0.0002), similar to findings for *cph* KV cilia length. The number

of cilia per KV was not significantly different between any of the groups (Fig 7d).

We next analysed cardiac looping at 48hpf in embryos injected with *arl13b* long and short

isoforms compared to uninjected controls. Immunofluorescence microscopy for the cardiac

marker MF20 revealed varied heart looping directionality. The uninjected control group heart

tube was correctly looped (Fig 7e top left); embryos from *arl13b* short isoform mRNA

injection showed a variety of cardiac laterality patterns including correct looping (34%),

delayed development and incomplete looping (66%) and reversed looping (2%) (Fig 7e; heart

looping direction listed on top left of image). No significant changes were seen in *arl13b* long

isoform injected embryos (Fig 7f; Fisher's exact test, p= 0.4949), whereas a significant

proportion of *arl13b* short isoform injected embryos displayed cardiac looping abnormalities

(Fig 7f; Fisher's exact test p<0.0001). Bright-field images of wild type uninjected fish

captured at 48hpf showed the expected cardiac morphology (Fig 7g top), whereas 56% of

arl13b short isoform injected embryos displayed pericardial oedema (Fig 7g bottom). The

percentage of the *arl13b* short isoform injected embryos showing pericardial oedema at 48hpf

was significantly increased compared to uninjected controls (Fig 7h; Fisher's exact test,

p<0.0001).

Over-expression of *arl13b* long isoform rescued phenotypic defects of *prpf8* loss in

zebrafish embryos

To determine if disruption of *arl13b* isoform usage is, at least in part, the mechanism

underlying the cardiac laterality defects in *cph* mutant fish, we utilised a *prpf8* splice

blocking morpholino (sbMO) targeting exon 7 (Supplemental Fig. 3), which provokes cardiac

laterality defects in injected zebrafish (57). KV cilia were identified by acetylated tubulin

immunofluorescence (Fig.8a). Compared to WT uninjected control embryos, the sbMO

injected embryo had significantly decreased KV cilia length (Fig. 8b; Unpaired t-test,

P<0.0001), but no significant difference was detected in cilia number per KV (Fig. 8c;

Unpaired t-test; p = 0.4868). Immunofluorescence microscopy at 48hpf showed cardiac

looping defects in embryos injected with *prpf8* sbMO (Fig. 8d), while co-injection of *prpf8*

sbMO with *arl13b* long isoform mRNA reduced the frequency of cardiac looping defects.

Co-injection of *prpf8* sbMO with *arl13b* short isoform mRNA did not rescue cardiac

development. Injection of a control MO showed no cardiac defects (Fig. 8d). Quantification

of cardiac defects revealed significant differences between the sbMO and control embryos

(Fig. 8e, Fisher's exact test p <0.0001), but not between embryos co-injected with *prpf8*

sbMO and *arl13b* long isoform mRNA when compared to control embryos (Fig. 8e; Fisher's

exact test, p >0.05). Additionally, a high incidence of severe pericardial oedema and tail

curvature defects was seen in embryos injected solely with *prpf8* sbMO, but embryos co-

injected with *prpf8* sbMO and *arl13b* long isoform mRNA exhibited a significant reduction

in the percentage of these defects (Fig. 8f-g; Fisher's exact test p <0.0001). The 72hpf

survival rate for *prpf8* sbMO embryos co-injected with *arl13b* long isoform mRNA was

significantly increased compared to *prpf8* sbMO injection alone (Fig. 8h; Fisher's exact test p

<0.0001). Morphological analysis of wild type uninjected control embryos or embryos co-

injected with *prpf8* sbMO and *arl13b* long isoform mRNA did not reveal obvious anatomical

defects (Fig. 8i) at 72hpf, whereas *prpf8* sbMO injected embryos exhibited pericardial

oedema (black arrowhead), head morphological defects (blue arrowhead), eye defects (green

arrowhead) and tail defects (orange arrowhead), as did embryos co-injected with *prpf8* sbMO

and *arl13b* short isoform mRNA.

The ARL13B short isoform displays reduced binding to IFT-B1 complex proteins

The short isoform of ARL13B lacks part of the proline rich region near the C-terminus of the

protein. The proline rich region binds IFT46 and IFT56 (58), components of the IFT-B1

complex involved in anterograde ciliary trafficking (59). We therefore sought to examine

differences in protein interactions between the long and short isoforms of ARL13B. Using a

GFP-trap system (60), we expressed either the GFP tag alone, or human long or short

ARL13B-GFP fusion protein isoforms in RPE1 cells, followed by GFP affinity purification-

mass spectrometry of three biological replicates. Network analysis of biological processes

that are enriched in proteins associated with the long isoform compared to the short isoform

revealed a network associated with cilia function (Fig. 8j; Supplemental Fig. 5). The short

isoform of ARL13B is depleted for binding to IFT22, IFT25, IFT46, IFT52, IFT56, IFT70a,

IFT74, and IFT81, which are all proteins found within the IFT-B1 complex (Fig. 8k;

Supplemental Fig. 5). Overall, these data suggest that the correct balance between ARL13B

isoforms is critical for maintaining cilia structure and function during development.

Discussion

We report a novel, evolutionarily conserved role for *Prpf8* in LRO cilia function and determination of organ laterality in vertebrate embryos. We identified a shift in *Arl13b* isoform usage as a potential mechanism explaining the laterality defects observed in *Prpf8* mutant mouse and zebrafish embryos. Consistent with this finding, *Arl13b* mouse mutants (61, 62) display cardiac looping defects, highly reminiscent of *Prpf8^{N1531S}* embryos described here (Fig 1), supporting the hypothesis that altered *Arl13b* isoform usage can perturb left-right axis formation and embryonic development. We did not observe changes in PRPF8 protein localisation to cilia (Supplemental Fig 2), suggesting that any as yet undefined ciliary, non-splicing roles of PRPF8 (12) may be intact in the *Prpf8^{N1531S}* homozygous mouse mutant.

To our knowledge, other characterised mouse strains with mutations in *Prpf8* do not resemble the *Prpf8^{N153S}* embryonic lethal laterality defect phenotype that we have detected (63-65). Previously characterised mutants were generated to model human RP, containing variants in the globular Jab1/MPN domain of PRPF8 (10). The N1531S mutation lies within the linker domain of PRPF8, near the endonuclease-like domain (43), and thus may have a different effect on PRPF8 function than mutations in the Jab1/MPN domain. Although many missense variants throughout yeast Prp8 have been characterised (42), additional study is needed in mammalian models to reveal the complexity of transcriptome and phenotypic defects that may arise from variants in differing protein domains. However, an iPSC model of a heterozygous *PRPF8* c.6926 A > C (p.H2309P) RP-associated mutation caused impaired alternative splicing and enhanced cryptic splicing predominantly in ciliary and retinal-specific transcripts (11), which is consistent with findings from the mouse N1531S variant. Although we have shown several aspects of the *Prpf8^{N1531S}* and *cph* mutant phenotypes are recapitulated by over-expression of the *arl13b* short isoform mRNA, we cannot exclude

contributions from other genes to the mutant phenotypes. For example, we detected splicing

alterations in the transcription factor *Rfx3* in *Prpf8^{N1531S}* mutants. *Rfx3* knockout embryos have

laterality and node cilia elongation defects (66), but their overall morphology, frequency of

situs inversus and incompletely penetrant embryonic lethality (66) are dissimilar to the

phenotypes of *Prpf8^{N1531S}* and *cph* mutants.

We demonstrated that increased usage of a previously uncharacterised transcript isoform of

arl13b, lacking exon 9, disrupted ciliogenesis and provoked laterality defects when over-

expressed in wild type zebrafish. We detect the short isoform in all tissues tested, in varying

ratios compared to the long isoform. Additional research is needed to reveal the roles of the

two different ARL13B isoforms in different tissues and cell types. In humans, pathological

variants in many ciliary genes, including *ARL13B*, cause Joubert syndrome (6, 67, 68), a well-

recognized ciliopathy (69). Laterality defects have been reported in some Joubert syndrome

patients, although individuals with laterality defects did not have variants in *ARL13B* (70).

Interactions between ARL13B and the IFT complex govern the speed of IFT entry into cilia,

and consequently, ciliary length (36, 71). Mutations in *Arl13b* and some IFT components in

zebrafish and mouse have been noted to result in short cilia (62, 72-74), suggesting that a

similar mechanism may play a role in the reduction in cilia length in zebrafish KV cilia in

embryos over-expressing the *arl13B* short isoform mRNA. Additionally, missense variants in

IFT52 cause cilia elongation defects in IMCD3 cells (75). Although the short isoform of the

ARL13B protein identified in *Prpf8^{N1531S}* and *cph* mutants localises to cilia, we demonstrated

that its interactions with IFT-B components, including IFT52, are reduced. Thus, the shorter

cilia detected in mutant LROs are consistent with a loss of IFT function.

Cilia motility is driven by dynein function. We found motility defects in the LRO cilia in

both $Prpf8^{N1531S}$ mutant mouse and cph fish embryos, yet the mouse node cilia appear to have

intact dynein inner and outer arms. Defective cilia motility with preservation of dynein

protein cilia localisation has previously been reported for $Dnaaf1^{m4Bei}$ and $Lrrc48^{m6Bei}$ mouse

mutants (76). Therefore, although the mechanism for cilia motility defects in $Prpf8$ mutants is

not fully understood, there is a precedent for the disruption of cilia motility without dynein

arrangement defects. Additionally, the shift in ARL13B isoform usage results in reduced

binding between ARL13B and IFT-B complex proteins, including IFT74, IFT22 and IFT81.

IFT74 has physical interactions with IFT22 and IFT81, and a binary interaction with the inner

dynein arm protein DNALI1 (77) proposed to be required for cilia motility (78). Patients with

hypomorphic IFT74 variants show cilia motility defects, although not laterality defects (79).

A mouse IFT74 mouse null mutant shows mid-gestation lethality and cardiac oedema, but

laterality defects were not reported for these mutants (80). To our knowledge, it is not clear if

these proteins are localised to the LRO cilia during development, but this is an area for future

investigation. There is high conservation of protein content between cilia of different types

and different localisations (81), which notably includes ARL13B and the IFT proteins found

to be affected in PRPF8 mutants. Together, our findings suggest a mechanistic link between

altered PRPF8 protein function in the spliceosome, aberrant splicing of $Arl13b$ transcript and

abnormalities in IFT and cilia function, culminating in conserved laterality defects in mutant

embryos.

Materials and Methods

l11Jus27 mouse model

The $Prpf8^{N1531S}$ strain is derived from the *l11Jus27* strain (32) and was backcrossed for at least 2 generations to the 129S5/SvEvBrd line carrying the chromosomal inversion Inv(11)8Brd$^{Trp-Wnt3}$, and then maintained through sibling intercrosses for at least 6 generations. Mouse colonies were maintained at the Biological Service Facility at the University of Manchester, UK, according to Home Office requirements and with local ethical approval (project licence number PP3720525). Mice were euthanised using a Schedule 1 method following UK Home Office regulations. Genotyping was performed as described (41).

Mouse embryo morphological analysis

Embryos were isolated from timed pregnant female mice by dissection in PBS. Morphology was analysed using a Leica dissecting microscope. Whole embryos were incubated in 1 ng/mL propidium iodide staining solution for 5 min, followed by 3 X 5 minute washes in PBS, before imaging with a DAPI filter to identify heart tube laterality.

Genome sequencing

Genomic DNA was extracted from an E10.5 *l11Jus27* homozygous embryo using ISOLATE II Genomic DNA Kit (Bioline). 1 μg of DNA was sequenced at The Manchester Centre for Genomic Medicine (MCGM), Saint Mary's Hospital, Manchester, UK, on the Illumina HiSeq platform. Reads were then aligned to the mouse MGSCv37 (mm9) genome assembly. Variants in the candidate region on chr 11 were analysed. Known variants in 129 and B6 mouse strains

were excluded, as well as variants present in fewer than 50% of reads. The remaining variants

were annotated with Ensembl Variant Effect Predictor (VEP). A single novel exonic mutation

was present in 100% of mapped reads: an A to G transition at position Chr11: 75,391,978

(mm39). Sanger sequencing was performed to verify the sequence change in three additional

individual embryos (University of Manchester Genomic Technologies Core Facility).

Mutation Analysis

Conservation of protein sequence across different species was evaluated using multiple

sequence alignments in Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/). Mutation

pathogenicity was predicted using SIFT (45) and PolyPhen2 (44). The Prpf8 protein

Alphafold structure was downloaded from AlphFold Protein Structure Database:

https://alphafold.ebi.ac.uk/entry/Q6P2Q9; and opened as entry AF-Q99PV0-F1-v4 in

PyMOL software (Version 3.1.1).

Quantitative PCR

Prpf8 expression was quantified using a *Prpf8* TaqMan probe (Applied Biosystems Cat:

4331182) on a DNA Engine Opticon 2 continuous fluorescence detector (BioRad). The raw

data were processed on Opticon monitor 3 (BioRad) software and subsequently analysed

using Excel (Microsoft). Samples were normalised against expression levels of *Gapdh*

(primers listed in Supplemental Table 4).

Scanning Electron Microscopy

Embryos at were dissected in PBS and fixed in 2.5% glutaraldehyde, 4% paraformaldehyde in

0.1M HEPES (pH7.4) at 4°C from overnight up to 1 month. Embryos were then washed 5x 5

minutes in dH_2O, then incubated for 1 hour in OsO_4 in 0.1M HEPES (pH7.4). They were then

washed 5x 5 minutes in dH$_2$O and taken through an ethanol dehydration series before critical

point drying. Embryos were mounted ventral side up on stubs using graphite sticky tape and

conductive epoxy. Samples were imaged at high vacuum at 10KV on a Quanta FEG 250

electron microscope (FEI).

Transmission Electron Microscopy

Embryos were fixed and processed as for SEM. Embedding, sectioning and imaging was

performed as previously described (82). Samples were imaged on a Tecnai 12 Biotwin

transmission electron microscope.

Nodal cilia videomicroscopy

E8.25 embryos were dissected individually in warmed DMEM modified with HEPES without

glutamine and sodium pyruvate (Sigma D6171), supplemented with 10% heat inactivated FBS

(Gibco 10500064). Embryos at the 1-3 somite stage were selected for imaging and were

transferred to a homemade chamber slide with the ventral node facing the microscope

objective. To visualise the nodal flow, the media was replaced with a 1:100 dilution of latex

microbeads (Sigma L1398) in media. A coverslip was then placed over the chamber and the

slides were incubated at 37°C for 10 minutes before imaging on a Leica DMRB at 40X using

a Leica MC170HD camera. Five-minute videos were taken using VGA2USB (Epiphan Video)

with an MPEG4 decompressor at maximum quality. Video analysis was performed using the

Manual Tracker plugin from Fiji (83). Movies were converted to .tiff stacks of 150 frames from

the first minute of filming and the paths of 5 beads were tracked.

For visualising nodal cilia movement directly, embryos were imaged on a Nikon TE2000 PFS

microscope using a 100x/1.49 Apo TIRF objective. Videos lasting 20 seconds were collected

on a FASTCAM SA3 (Photron) at 125 frames per seconds and analysed using Fiji. Imaging

data were enhanced for visual analysis by performing 11-frame running-average background

subtraction followed by inversion and thresholding, using custom routines coded in MATLAB

R2015a with Image Processing toolbox (Mathworks, Natick, MA). Motile cilia were counted

manually by inspection of 125 frame (1 s) sequences taken at each focal plane.

Mouse and zebrafish embryo whole mount *in situ* hybridisation

Whole mount *in situ* hybridisation was performed according to established protocols in mouse

(84) and zebrafish (85). Control and mutant embryos were pooled and hybridised with

digoxigenin labelled probes and incubated with alkaline phosphatase conjugated anti-

digoxigenin antibodies. Hybridised embryos were then developed with NBT-BCIP until

desired staining was achieved. *In situ* probes were synthesised using either plasmid or PCR

templates. *In situ* probe primer sequences for PCR templates are listed in Supplemental Table

4.

Mouse embryo immunofluorescence staining

Antibodies used for immunostaining are found in Supplemental Table 5. Embryos previously

stored at -20C in methanol were rehydrated through a methanol series, washed 3x 20 minutes

in PBS and blocked for 90 minutes at room temperature in 5% horse serum and 0.5% Triton

X-100 in PBS. Control and mutant embryos were pooled and then incubated overnight at 4°C

in 5% horse serum, 0.5% Triton X-100 in PBS with primary antibodies. Embryos were then

washed in 3x20 minutes in PBS and then incubated with secondary antibodies in PBS with 1%

horse serum for 2 hours at room temperature. Embryos were washed 3x10 minutes in PBS and

then incubated with Cy5 labelled streptavidin in PBS for 30 minutes. Embryos were washed

3x20 minutes in PBS with 1% horse serum and then stained with DAPI in PBS, before a final

10 minute wash in ice-cold PBS. Embryos were then individually placed in dishes of PBS and

the anterior portion of the embryo was removed. The posterior portion, including the node,

was transferred to a homemade chamber slide, excess PBS removed, and embryos mounted in

Prolong Gold Antifade (Life Sciences, P36934). The anterior portion was used to re-genotype

the samples. The embryo was arranged ventral side up and a coverslip was placed on top, with

gentle pressure applied to remove bubbles and excess mounting medium. Slides were cured

for at least 2 days at room temperature.

Embryos were imaged on a Leica TCS SP5 AOBS upright confocal microscope using a

63x/1.40 HCX PL Apo objective and a 2.0x digital zoom using sequential imaging in a

2048x2048 format.

Mouse Embryo Western Blot

E10.5 mouse embryos were lysed in RIPA buffer supplemented with protease inhibitors. As

described earlier, protein lysates were separated by SDS-PAGE and transferred to PVDF

membranes (86). Western blotting was performed using antibodies against PRPF8 and GAPDH

(loading control). Signal detection was carried out using LI-COR Odyssey system and the

image was quantified using the LI-COR Image Studio software.

PRPF8 FLAG-tagged plasmid generation and transfection

IMAGE clone 5587081 containing human PRPF8 was used for generating PRPF8 constructs.

Mouse and human PRPF8 proteins differ by only 3 of the total 2335 amino acids. We

generated the N1531S variant by site directed mutagenesis (Q5 Site Directed Mutagenesis

kit; NEB), and corrected a A759G missense sequence change present in the IMAGE clone.

Primer sequences are listed in Supplemental Table 4. To correct the A759G sequence

change, primers creating a new restriction site for Eco47III were utilised. Rather than

creating the exact *l11Jus27* mutation (aat to agt), we used an alternate codon sequence which

generated the same amino acid mutation (aat to agc), but created a BseYI restriction site

allowing screening for colonies containing the correct mutation prior to sequence verification.

Oligonucleotides containing a FLAG tag sequence, a start codon, a Kozak sequence and a

Cla1 restriction site were cloned into the corrected wild type and the N1531S mutant PRPF8

plasmids at an AgeI restriction site.

PRPF8 protein stability analysis

HEK293T cells were cultured as described previously (86). The cells were seeded at a

density of 0.1×10^6/well in 6 well plates for transfection. The next day the cells were

transfected with 2μg of *PRPF8* WT-Flag or N1531S-Flag plasmid using Fugene HD reagent

(Promega). 48hours later the transfection media was removed, and cells were treated with

100μM cycloheximide in cell culture media. The cells were lysed at different time points (0,

4, 8, 16 hours) in RIPA buffer. The immunoblotting was performed using anti Flag and anti

β-Actin (ACTB) antibodies as described previously (86). Signal detection was carried out

using LI-COR Odyssey system. Image was quantified using the LI-COR Image Studio

software.

Zebrafish *cph* mutants

Fish were maintained at the Institute of Molecular and Cell Biology, A*STAR Zebrafish

Facility and the University of Manchester Biological Services Facility (IMCB protocol

number: 221702 and Manchester project licence PP8367714) at a temperature of 28.5°C and a

standard light/dark cycle of 14 hours and 10 hours respectively. Pair mating was used to obtain

embryos. Embryos were incubated at 28.5°C in egg water containing methylene blue until the

desired stage.

Genotyping *cph* mutants

cph mutant zebrafish were genotyped by PCR amplification of exon 8 and 9 of the *prpf8* gene

(52), followed by purification using QIAquick PCR and Gel Purification Kit (Cat. No 28104),

and a restriction digest using AccI enzyme performed as per manufacturer's instructions.

Samples were ran on a 2% agarose gel. WT fish produced one band, heterozygotes produced 3

bands and *cph* homozygotes produced 2 bands (Supplemental Figure 6).

Zebrafish whole mount immunofluorescence

Embryos were transferred into a glass cavity dish and rehydrated from methanol by washing

in consecutively lower methanol concentrations (75%, 50%, 25%) in PBS. Embryos were

subsequently washed in PBS for 3x 5 minutes. PBS was removed and embryos treated with -

20°C acetone and placed at -20°C for 7 minutes (this step was not used for embryos fixed with

Dent's fix). Embryos were washed for 3x 5 minutes in room temperature PBS. Embryos were

incubated for 1 hour at room temperature in blocking buffer (2% sheep serum in PBDT) on an

orbital shaker at 70 rpm. Blocking buffer was removed and primary antibody diluted in PBDT

added (Supplemental Table 5). Embryos were placed on an orbital shaker at 70rpm overnight

at 4°C. The following morning primary antibody was removed and embryos washed in room

temperature PBDT 3x 30 minutes on an orbital shaker at 70rpm. PBDT was removed and

secondary antibodies diluted in fresh PBDT were added. Embryos were placed on an orbital

shaker at 70rpm for 5 hours at room temperature. During the last 30 minutes DAPI was added

(1:2000). Secondary antibodies were removed and embryos washed 3x 30 minutes with room

temperature PBDT. PBDT was removed and embryos were submerged in 70% glycerol for

storage at 4°C. Primary antibodies for immunostaining found in Supplemental Table 5.

Measurement of Node and KV LRO Cilia

To detect nodal cilia, mouse 1-4 somite embryos were stained for Ac-tubulin and DAPI and

subsequently mounted and imaged as described earlier. Zebrafish 14hpf embryos were stained

with Ac-tubulin, Gamma-Tubulin and DAPI as per zebrafish whole mount

immunofluorescence protocol. KV was dissected out and mounted on a slide. Mounted KVs

were imaged on an Olympus Fluoview 1000 confocal microscope and Leica TCS SP8 AOBS

inverted confocal microscope at x100 zoom and a z-stack images were taken for the entire

vesicle. Mouse and fish images were processed using ImageJ with Simple Neurite Tracer

utilised to measure each cilium from the base to the tip in 3 dimensions. After imaging, embryos

were retrieved from the slides, washed with PBS and used for genotyping.

Zebrafish KV Motile Cilia Live Imaging

Embryos from *cph* heterozygous fish incross were collected and placed in egg water at 28.5°C

to develop until ~14hpf. A mould designed to hold embryos was made using 2% agarose in

egg water and left to set. Once embryos had developed to the correct stage, embryos with well-

developed KVs were dechorionated and placed into the mould. Embryos were covered with

egg water and placed on a Zeiss Imager M2 live imaging microscope. All embryos were imaged

on a 100x water dipping lens at ~150-200fps using Metamorph software. Embryos were then

placed into separate wells of a 24 well plate coated with 2% agarose in egg water to stop the

embryos sticking to the plastic. Recordings of each embryo were saved by well number.

Embryos were left to develop until the mutant phenotype became apparent and recordings were

correlated with the phenotype. Recordings were analysed using ImageJ and slowed to 15fps.

RNA sequencing analysis

Total RNA was isolated from wild type and *Prpf8^{N1531S}* mutant embryos at E10.5 using Trizol

according to manufacturer's instructions (Ambion). RNA was prepared from 3 embryos of

each genotype. RNA quality was evaluated using TapeStation.

Paired-end RNA sequencing of E10.5 mouse embryo samples was performed on the Illumina

HiSeq 3000 platform in the University of Leeds Next Generation Sequencing Facility. The

FASTQ files were deposited into the Sequence Read Archive (SRA) under the accession

number PRJNA690736. For RNA-Seq data, the adapter sequences were removed using

Cutadapt (1.9.1) (87) , and the filtering and trimming of sequences were done using

PRINSEQ (0.20.4) (88). The clean reads were aligned to the UCSC mouse genome (mm10)

(89) using STAR (2.5.1b) (90). The processing of BAM files was done using SAMtools

(1.3.1) (91) and only uniquely mapped alignments were kept for further analysis. Read counts

for protein-coding genes were generated through the featureCounts function provided by

Rsubread package (1.22.0) (92) and RPKM values were calculated using rpkm function of

edgeR package (3.14.0) (93). For those genes with RPKM >= 1 in at least one sample, a

value of 1 was added to all RPKM values before the log2 transformation. Principal

component analysis was carried out on the quantile normalised values and the first two

principal components were plotted. We observed three expression outliers based on poor

clustering: one from the wild type, one from the mutant correct looped, and one from the

mutant reversed loop strain, which were excluded from downstream splicing analysis.

Differential expression between groups of samples were conducted using DESeq2 package

(1.12.0) (94) based on the raw counts for genes. Differential alternative splicing events were

identified using rMATS (4.0.2) (55).

Differential splicing analysis

Splicing events in RNA-seq data from the three wild-type day 10.5 embryos were compared against those in the three $Prpf8^{N1531S}$ mutant embryos using rMATS v4.0.2. Results were filtered to retain only splicing events exhibiting a difference in exon level inclusion (IncLevelDifference $\geq$ 0.2 or $\leq$ -0.2), and an FDR < 0.05.

Gene ontology (GO) analysis

GO enrichment was performed using the PANTHER GO overrepresentation test (http://geneontology.org/). Genes harbouring exon skipping events with a difference in inclusion of $\geq$ 20% between the wild-type and $Prpf8^{N1531S}$ mutant embryos were evaluated against all *Mus musculus* genes in the GO Ontology Database. GO terms exhibiting a positive enrichment were retained. Genes present in the rMATS output were annotated with their respective GO annotations, retaining those annotations labelled as "biological_process" and with relationships to their genes of "involved_in" and "enables".

***Arl13b* isoform RT-PCR quantification**

qPCR could not be utilised for isoform quantification as no primers or probe could be generated specifically for the "short" isoform of *Arl13b*. To quantify relative isoform levels, *Arl13b* RT-PCR was performed with a species-specific single primer set (Supplemental Table 4) that amplifies both the short and long isoforms. Gel electrophoresis was carried out to separate isoforms, with gels imaged on a Li-Cor Odessey and optical intensity quantified using Image Studio software with background gel intensity set to 0. The ratio of optical intensity of the "short" isoform to the "long" isoform was then calculated for each sample, and fold change

751 calculated for the mutants as compared to wild type. Data are shown for at least three individual

752 embryos of each genotype.

Generation of Prp8^{N1603S} yeast and yeast splicing analysis

The N1603S mutation was introduced into pRS413-PRP8 by site directed mutagenesis (95).

The pRS413-PRP8, pRS413-PRP8^{N1603S} or pRS413 plasmid was transformed into a haploid

yeast strain where both the *CUP1* and *PRP8* genes were deleted and *PRP8* function was

complemented with pRS416-PRP8. Growth on 5-fluoro-orotic acid (5-FOA) containing plates

was used to select against and remove the complementing pRS416-PRP8 plasmid to leave the

pRS413 plasmids as the sole source of *PRP8* and growth properties observed at 16°C, 30°C

and 37°C. The resulting haploid strains with pRS413-PRP8 or pRS413-PRP8^{N1603S} were

transformed with different *ACT1-CUP1* splicing reporters (53) and growth assayed on plates

with increasing concentrations of $CuSo_4$. Primer extension was carried out (53) to monitor the

two steps of splicing from the *ACT1-CUP1* splicing reporters.

ARL13B mammalian expression constructs

767 The long and short isoforms of human *ARL13B* were synthesised by GenScript and inserted as

768 a N-terminal EGFP fusion protein into a mammalian expression plasmid with a CMV

769 promoter. Sequence was confirmed by whole plasmid sequencing (Genewiz).

hTERT-RPE1 cell transfections and imaging

Cell Culture

773 hTERT-RPE1 cells were cultured in DMEM/F12 media containing 10% (v/v) foetal bovine

774 serum (FBS) and 2mM L-glutamine. Cells were split 1 in 10 using trypsin-EDTA when

775 reaching 80-90% confluence. Cells were incubated at 37°C, 5% CO_2.

Transfection

hTERT-RPE1 cells were transfected with *GFP*-tagged *ARL13B* long and short isoform plasmid using Lipofectamine^TM 3000 Transfection Reagent (Invitrogen), following the manufacturer's instructions. The transfection medium was replaced with 0.2% FBS starvation medium after 24h and cells were incubated a further 24h-48h to induce cilia before use.

Immunofluorescence

hTERT-RPE1 cells on coverslips were fixed in 4% paraformaldehyde for 10 minutes at RT. Cells were then permeabilised with 0.3% Triton X-100 in PBS for 30 minutes at RT and blocked in SuperBlock blocking buffer (Thermo Fisher Scientific) for 1h at RT. Cells were incubated in the primary antibody (mouse anti-acetylated tubulin, Supplemental Table 5) diluted in 'Solution 1 for primary antibody' from SignalBoost Immunoreaction Enhancer Kit (Merck) overnight at 4°C before washing three times in 0.1% PBS-Tween20 and incubation in the secondary antibody (Alexa Fluor donkey anti-mouse 594 IgG (H+L) AffiniPure, Jackson ImmunoResearch) diluted in 'Solution 2 for secondary antibody' from SignalBoost Immunoreaction Enhancer Kit for 1h at RT. Cells were then washed three times in 0.1% PBS-Tween20 followed by incubation in DAPI for 5min. Coverslips were air dried before mounting with Prolong Gold antifade reagent (Thermo Fisher Scientific). Cells were imaged on a Leica TCS SP8 AOBS inverted confocal microscope. Images were processed with Fiji.

Zebrafish *arl13b* mRNA injections

Zebrafish *arl13b* long isoform was amplified by RT-PCR from zebrafish cDNA and cloned into PCS2+ xlt Vector using Hifi DNA assembly kit (New England Biolabs). Inverse PCR was used to generate the *arl13b* short isoform lacking exon 9. Plasmid inserts were sequenced (Genewiz). RNA was transcribed from a linearised plasmid using the mMessage Machine kit (Invitrogen). RNA was precipitated and stored at -80°C. RNA was diluted in DEPC-H_2O to the

desired concentration, and 0.5nl of 80ng/µl *arl13b* long or short isoform mRNAs was

microinjected into zebrafish embryos at the single cell stage. Injected embryos were collected

and placed in fresh egg water and left to develop at 28.5°C to the desired developmental stages.

Embryos with green florescence were selected at 12hpf (using a LEICA M165 FC

stereomicroscope) for use in further analysis.

***prpf8* splice blocking morpholino knockdown and rescue**

Splicing blocking morpholinos targeting exon 7 of zebrafish *prpf8* (57) were synthesized by

GeneTools (Supplemental Table 4). A 300nmol *prpf8* sbMO stock solution was made in sterile

water to a final concentration of 1mM, and zebrafish embryos were injected with 0.5 nl of a

1:2 dilution of sbMO stock at the single-cell stage. For the rescue of *prpf8* morphants, after the

injection of morpholinos, 0.5nl of *arl13b-GFP* long isoform or *arl13b-GFP* short isoform

mRNA (200ng/ul) was injected into each morpholino injected embryo. Wild type embryos

were injected with 0.5nl of a 1:2 dilution of Gene Tools standard MO as a control

(Supplemental Table 4). Embryos were cultured in fish embryo medium (0.3g/L NaCl) until

12hpf. Embryos exhibiting green fluorescence by microscopy (LEICA M165 FC) were

selected for further study. Embryos were incubated at 28.5°C until the desired developmental

stages to observe the phenotype using brightfield microscopy (LEICA M80).

ARL13B Affinity purification-mass spectrometry

Cloning and Vector Preparation

For GFP-trap affinity purification-mass spectrometry (AP-MS) analysis, the *ARL13B*_long

and *ARL13B*_short isoforms were cloned into the CMV-EGFP-RGCC-SV40 poly(A)-

NeoR/KanR vector (gifted by Prof. Ben Goult, University of Liverpool) by replacing the

RGCC gene sequence with the respective *ARL13B* isoform sequences. Cloning was

performed using SacI-HF and XmaI restriction enzymes (New England Biolabs, NEB, USA)

and HiFi DNA Assembly, according to the manufacturer's instructions. The control vector,

expressing GFP alone, was generated by excising the RGCC sequence using MfeI-HF and

BspEI restriction enzymes (NEB) and inserting a short filler sequence using HiFi DNA

Assembly.

Transfection

$1x10^6$ hTERT-RPE1 cells were seeded in 10cm plates one day before transfection.

Transfection was performed using Lipofectamine 3000 transfection reagent (Invitrogen,

L3000015) according to the manufacturer's protocol. For each plate, 8µg of plasmid DNA

was used. After 10 hours of transfection, the transfection medium was replaced with fresh

complete medium to allow the cells to recover. At 24 hours post-transfection, the medium

was changed to low-serum medium (0.2% FBS) for 24 hours to induce ciliogenesis before

harvesting.

Cell Harvesting and Peptide Sample Preparation

hTERT-RPE1 cells expressing GFP-vector, GFP-ARL13B short isoform and GFP-ARL13B

long isoform were harvested in RIPA buffer (Thermo Scientific, 89900, USA) supplemented

with EDTA-free protease inhibitor cocktail (Roche, REF: 04693159001), PhosSTOP

phosphatase inhibitor (Roche, REF: 04906837001), DNase (50 U/ml, Thermo Scientific, Cat:

EN0521), and 2.5 mM MgCl2 (NEB, B0510A). Cell lysates were clarified by centrifugation.

GFP pull-down, trypsin digestion, and peptide preparation for AP-MS analysis were

performed according to the protocol provided with the ChromoTek iST Myc-Trap® Kit

(Proteintech, Cat no: ytak-iST).

Mass Spectrometry Analysis

Samples were analyzed using a Thermo Exploris 480 mass spectrometer equipped with a

FAIMS Pro interface and coupled to a U3000 nanoUPLC system. Each sample underwent a

1-hour LC-MS run, performed at the Biological Mass Spectrometry Facility, University of

Manchester.

Data Analysis

Mass spectrometry data were processed using Fragpipe (v22.0), incorporating MSFragger

(v4.1), IonQuant (v1.10.27), and DiaTracer (v1.1.5) within a label-free quantification (LFQ)

and match-between-runs (MBR) workflow. MaxLFQ intensity values were exported from

Fragpipe and analyzed using the R Shiny application "Manchester Proteome Profiler"

(developed by Dr. Stuart Cain, unpublished). Proteins were identified as significantly

enriched based on a fold-change threshold of >4 and an adjusted p-value cutoff of <0.05.

Statistical analysis

Statistical analyses and the generation of graphs was performed on GraphPad Prism 10 for

Mac, GraphPad Software, La Jolla, California, USA.

Data Availability

RNAseq data can be accessed in the Sequence Reads Archive. Mass spectrometry data is

available in the PRIDE database. Supplemental Movie Files are located on Zenodo with doi:

10.5281/zenodo.15482494.

Acknowledgements

We thank Monica Justice for the *l11Jus27* mouse line and Graham Lieschke for the *cph* fish

line. We thank Dominic Norris for advice on EM imaging of cilia, and for the *Nodal*, *Cerl2*,

Lefty1 and *Lefty2* mouse *in situ* hybridisation probe plasmids. We thank Megan Davey for

advice on immunostaining of cilia. We thank Adam Hurlstone, Martin Lowe, Shane Herbert

and Karel Dorey for advice on zebrafish experiments and access to equipment. Some

computational data analysis work was undertaken on ARC4, part of the High-Performance

Computing facility at the University of Leeds, UK. The authors wish to thank the staff in the

University of Manchester FBMH Electron Microscopy Core Facility, Biological Mass

Spectrometry Facility (RRID: SCR_020987), and Peter March in the BioImaging Core Facility

for their assistance, and the Wellcome Trust for funding to support core facility equipment. We

thank Stuart Cain for assistance with mass spectrometry data analysis and Xiaofan Xiong for

uploading data to PRIDE. We thank the Biological Services Facility at the University of

Manchester for mouse and fish husbandry. We thank the IMCB zebrafish facility for

maintenance of zebrafish strains.

Funding Statement

This work was supported by BHF grants PG/06/144/21898, PG/10/87/28624 and

PG/18/28/33632 to KEH, Fight for Sight grant 5163/5164 to KEH, and funding from the

Agency for Science, Technology and Research (A*STAR) of Singapore to S.R. KEH and BK

acknowledge support from a British Heart Foundation Research Excellence Award

(RE/24/130017). BK was supported by a BHF personal chair (CH/13/2/30154). DWM was

supported by a University of Manchester-A*STAR Research Attachment Program (ARAP)

Ph.D. fellowship. This work was supported in part by the Wellcome Trust [105610/Z/14/Z].

MB, LS and KM were supported by BBSRC DTP studentships awarded to the University of

Manchester. MTG and DJS acknowledge EPSRC award EP/N021096/1.

Author Contributions

FJ, MB, DM, WMSQ, CFR, GT, JZ, JL, KM, LAS, NA, DVL, EJRV, DW, MTG, BB, KS,

AV, DS, KEH performed research. FJ, MB, DM, KEH wrote original manuscript draft. BK,

MJH, JE, DS, CAJ, ROK, SR and KEH provided supervision and obtained funding. All

authors read and edited the manuscript draft and approved submission.

Competing Interests

No competing interests

Materials and Correspondence - KEH

References

1. Mirvis M, Stearns T, James Nelson W. Cilium structure, assembly, and disassembly regulated by the cytoskeleton. The Biochemical journal. 2018;475(14):2329-53.

2. Shinohara K, Hamada H. Cilia in Left-Right Symmetry Breaking. Cold Spring Harb Perspect Biol. 2017;9(10).

3. Waters AM, Beales PL. Ciliopathies: an expanding disease spectrum. Pediatr Nephrol. 2011;26(7):1039-56.

4. Ferrari S, Di Iorio E, Barbaro V, Ponzin D, Sorrentino FS, Parmeggiani F. Retinitis pigmentosa: genes and disease mechanisms. Curr Genomics. 2011;12(4):238-49.

5. Wheway G, Lord J, Baralle D. Splicing in the pathogenesis, diagnosis and treatment of ciliopathies. Biochim Biophys Acta Gene Regul Mech. 2019;1862(11-12):194433.

6. Maxwell DW, O'Keefe RT, Roy S, Hentges KE. The role of splicing factors in retinitis pigmentosa: links to cilia. Biochem Soc Trans. 2021;49(3):1221-31.

7. Tanackovic G, Ransijn A, Thibault P, Abou Elela S, Klinck R, Berson EL, et al. PRPF mutations are associated with generalized defects in spliceosome formation and pre-mRNA splicing in patients with retinitis pigmentosa. Hum Mol Genet. 2011;20(11):2116-30.

8. Korir PK, Roberts L, Ramesar R, Seoighe C. A mutation in a splicing factor that causes retinitis pigmentosa has a transcriptome-wide effect on mRNA splicing. BMC Res Notes. 2014;7:401.

9. Arzalluz-Luque A, Cabrera JL, Skottman H, Benguria A, Bolinches-Amoros A, Cuenca N, et al. Mutant PRPF8 Causes Widespread Splicing Changes in Spliceosome Components in Retinitis Pigmentosa Patient iPSC-Derived RPE Cells. Front Neurosci. 2021;15:636969.

10. Malinova A, Cvackova Z, Mateju D, Horejsi Z, Abeza C, Vandermoere F, et al. Assembly of the U5 snRNP component PRPF8 is controlled by the HSP90/R2TP chaperones. J Cell Biol. 2017;216(6):1579-96.

11. Atkinson R, Georgiou M, Yang C, Szymanska K, Lahat A, Vasconcelos EJR, et al. PRPF8-mediated dysregulation of hBrr2 helicase disrupts human spliceosome kinetics and 5-splice-site selection causing tissue-specific defects. Nat Commun. 2024;15(1):3138.

12. Wheway G, Schmidts M, Mans DA, Szymanska K, Nguyen TM, Racher H, et al. An siRNA-based functional genomics screen for the identification of regulators of ciliogenesis and ciliopathy genes. Nat Cell Biol. 2015;17(8):1074-87.

13. Kim JH, Ki SM, Joung JG, Scott E, Heynen-Genel S, Aza-Blanc P, et al. Genome-wide screen identifies novel machineries required for both ciliogenesis and cell cycle arrest upon serum starvation. Biochim Biophys Acta. 2016;1863(6 Pt A):1307-18.

14. Xu K, Ou G. Cilia regeneration requires an RNA splicing factor from the ciliary base. Cell Regen. 2022;11(1):29.

15. Okada Y, Takeda S, Tanaka Y, Belmonte JI, Hirokawa N. Mechanism of nodal flow: a conserved symmetry breaking event in left-right axis determination. Cell. 2005;121(4):633-44.

16. Sulik K, Dehart DB, Iangaki T, Carson JL, Vrablic T, Gesteland K, Schoenwolf GC. Morphogenesis of the murine node and notochordal plate. Dev Dyn. 1994;201(3):260-78.

17. Nonaka S, Tanaka Y, Okada Y, Takeda S, Harada A, Kanai Y, et al. Randomization of left-right asymmetry due to loss of nodal cilia generating leftward flow of extraembryonic fluid in mice lacking KIF3B motor protein. Cell. 1998;95(6):829-37.

18. Okada Y, Nonaka S, Tanaka Y, Saijoh Y, Hamada H, Hirokawa N. Abnormal nodal flow precedes situs inversus in iv and inv mice. Mol Cell. 1999;4(4):459-68.

19. McGrath J, Somlo S, Makova S, Tian X, Brueckner M. Two populations of node monocilia initiate left-right asymmetry in the mouse. Cell. 2003;114(1):61-73.

20.   Katoh TA, Omori T, Mizuno K, Sai X, Minegishi K, Ikawa Y, et al. Immotile cilia
mechanically sense the direction of fluid flow for left-right determination. Science.
2023;379(6627):66-71.
21.   Marques S, Borges AC, Silva AC, Freitas S, Cordenonsi M, Belo JA. The activity of
the Nodal antagonist Cerl-2 in the mouse node is required for correct L/R body axis. Genes
Dev. 2004;18(19):2342-7.
22.   Collignon J, Varlet I, Robertson EJ. Relationship between asymmetric nodal
expression and the direction of embryonic turning. Nature. 1996;381(6578):155-8.
23.   Nakamura T, Saito D, Kawasumi A, Shinohara K, Asai Y, Takaoka K, et al. Fluid
flow and interlinked feedback loops establish left-right asymmetric decay of Cerl2 mRNA.
Nat Commun. 2012;3:1322.
24.   Saijoh Y, Adachi H, Sakuma R, Yeo CY, Yashiro K, Watanabe M, et al. Left-right
asymmetric expression of lefty2 and nodal is induced by a signaling pathway that includes
the transcription factor FAST2. Mol Cell. 2000;5(1):35-47.
25.   Norris DP, Brennan J, Bikoff EK, Robertson EJ. The Foxh1-dependent autoregulatory
enhancer controls the level of Nodal signals in the mouse embryo. Development.
2002;129(14):3455-68.
26.   Yamamoto M, Mine N, Mochida K, Sakai Y, Saijoh Y, Meno C, Hamada H. Nodal
signaling induces the midline barrier by activating Nodal expression in the lateral plate.
Development. 2003;130(9):1795-804.
27.   Shiratori H, Sakuma R, Watanabe M, Hashiguchi H, Mochida K, Sakai Y, et al. Two-
step regulation of left-right asymmetric expression of Pitx2: initiation by nodal signaling and
maintenance by Nkx2. Mol Cell. 2001;7(1):137-49.
28.   Kitamura K, Miura H, Miyagawa-Tomita S, Yanazawa M, Katoh-Fukui Y, Suzuki R,
et al. Mouse Pitx2 deficiency leads to anomalies of the ventral body wall, heart, extra- and
periocular mesoderm and right pulmonary isomerism. Development. 1999;126(24):5749-58.
29.   Semina EV, Reiter R, Leysens NJ, Alward WL, Small KW, Datson NA, et al. Cloning
and characterization of a novel bicoid-related homeobox transcription factor gene, RIEG,
involved in Rieger syndrome. Nat Genet. 1996;14(4):392-9.
30.   Liu C, Liu W, Lu MF, Brown NA, Martin JF. Regulation of left-right asymmetry by
thresholds of Pitx2c activity. Development. 2001;128(11):2039-48.
31.   Franco D, Sedmera D, Lozano-Velasco E. Multiple Roles of Pitx2 in Cardiac
Development and Disease. J Cardiovasc Dev Dis. 2017;4(4).
32.   Kile BT, Hentges KE, Clark AT, Nakamura H, Salinger AP, Liu B, et al. Functional
genetic analysis of mouse chromosome 11. Nature. 2003;425(6953):81-6.
33.   Hentges KE, Nakamura H, Furuta Y, Yu Y, Thompson DM, O'Brien W, et al. Novel
lethal mouse mutants produced in balancer chromosome screens. Gene Expr Patterns.
2006;6(6):653-65.
34.   Cevik S, Hori Y, Kaplan OI, Kida K, Toivenon T, Foley-Fisher C, et al. Joubert
syndrome Arl13b functions at ciliary membranes and stabilizes protein transport in
Caenorhabditis elegans. J Cell Biol. 2010;188(6):953-69.
35.   Song P, Perkins BD. Developmental expression of the zebrafish Arf-like small
GTPase paralogs arl13a and arl13b. Gene Expr Patterns. 2018;29:82-7.
36.   Lu H, Toh MT, Narasimhan V, Thamilselvam SK, Choksi SP, Roy S. A function for
the Joubert syndrome protein Arl13b in ciliary membrane extension and ciliary length
regulation. Dev Biol. 2015;397(2):225-36.
37.   Gigante ED, Taylor MR, Ivanova AA, Kahn RA, Caspary T. ARL13B regulates
Sonic hedgehog signaling from outside primary cilia. Elife. 2020;9.
38.   Pintado P, Sampaio P, Tavares B, Montenegro-Johnson TD, Smith DJ, Lopes SS.
Dynamics of cilia length in left-right development. R Soc Open Sci. 2017;4(3):161102.

39. Cevik S, Sanders AA, Van Wijk E, Boldt K, Clarke L, van Reeuwijk J, et al. Active
transport and diffusion barriers restrict Joubert Syndrome-associated ARL13B/ARL-13 to an
Inv-like ciliary membrane subdomain. PLoS Genet. 2013;9(12):e1003977.
40. Fu W, Wang L, Kim S, Li J, Dynlacht BD. Role for the IFT-A Complex in Selective
Transport to the Primary Cilium. Cell Rep. 2016;17(6):1505-17.
41. Ridge LA, Mitchell K, Al-Anbaki A, Shaikh Qureshi WM, Stephen LA, Tenin G, et
al. Non-muscle myosin IIB (Myh10) is required for epicardial function and coronary vessel
formation during mammalian development. PLoS Genet. 2017;13(10):e1007068.
42. Grainger RJ, Beggs JD. Prp8 protein: at the heart of the spliceosome. RNA.
2005;11(5):533-57.
43. Galej WP, Oubridge C, Newman AJ, Nagai K. Crystal structure of Prp8 reveals active
site cavity of the spliceosome. Nature. 2013;493(7434):638-43.
44. Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense
mutations using PolyPhen-2. Curr Protoc Hum Genet. 2013;Chapter 7:Unit7 20.
45. Ng PC, Henikoff S. SIFT: Predicting amino acid changes that affect protein function.
Nucleic Acids Res. 2003;31(13):3812-4.
46. Kennedy MP, Omran H, Leigh MW, Dell S, Morgan L, Molina PL, et al. Congenital
heart disease and other heterotaxic defects in a large cohort of patients with primary ciliary
dyskinesia. Circulation. 2007;115(22):2814-21.
47. Rachel RA, Li T, Swaroop A. Photoreceptor sensory cilia and ciliopathies: focus on
CEP290, RPGR and their interacting proteins. Cilia. 2012;1(1):22.
48. Valcarcel J, Malumbres M. Splicing together sister chromatids. EMBO J.
2014;33(22):2601-3.
49. Tsiairis CD, McMahon AP. An Hh-dependent pathway in lateral plate mesoderm
enables the generation of left/right asymmetry.
50. Nowotschin S, Liao J, Gage PJ, Epstein JA, Campione M, Morrow BE. Tbx1 affects
asymmetric cardiac morphogenesis by regulating Pitx2 in the secondary heart field.
51. Tissier-Seta JP, Mucchielli ML, Mark M, Mattei MG, Goridis C, Brunet JF. Barx1, a
new mouse homeodomain transcription factor expressed in cranio-facial ectomesenchyme
and the stomach. Mech Dev. 1995;51(1):3-15.
52. Keightley MC, Crowhurst MO, Layton JE, Beilharz T, Markmiller S, Varma S, et al.
In vivo mutation of pre-mRNA processing factor 8 (Prpf8) affects transcript splicing, cell
survival and myeloid differentiation. FEBS Lett. 2013;587(14):2150-7.
53. Lesser CF, Guthrie C. Mutational analysis of pre-mRNA splicing in Saccharomyces
cerevisiae using a sensitive new reporter gene, CUP1. Genetics. 1993;133(4):851-63.
54. Kurtovic-Kozaric A, Przychodzen B, Singh J, Konarska MM, Clemente MJ, Otrock
ZK, et al. PRPF8 defects cause missplicing in myeloid malignancies. Leukemia.
2015;29(1):126-36.
55. Shen S, Park JW, Lu ZX, Lin L, Henry MD, Wu YN, et al. rMATS: robust and
flexible detection of differential alternative splicing from replicate RNA-Seq data. Proc Natl
Acad Sci U S A. 2014;111(51):E5593-601.
56. Mahajan D, Madugula V, Lu L. Rab8 and TNPO1 are ciliary transport adaptors for
GTPase Arl13b by interacting with its RVEP motif containing ciliary targeting sequence. J
Biol Chem. 2023;299(5):104604.
57. Boylan M. A Novel Point Mutation in Prpf8 Causes Defects in Left-Right Axis
Establishment in the Mouse. United Kingdom: University of Manchester; 2015.
58. Nozaki S, Katoh Y, Terada M, Michisaka S, Funabashi T, Takahashi S, et al.
Regulation of ciliary retrograde protein trafficking by the Joubert syndrome proteins
ARL13B and INPP5E. J Cell Sci. 2017;130(3):563-76.

59.   Lacey SE, Foster HE, Pigino G. The molecular structure of IFT-A and IFT-B in
anterograde intraflagellar transport trains. Nat Struct Mol Biol. 2023;30(5):584-93.
60.   Zhang S, Williamson NA, Bogoyevitch MA. Complementary proteomics strategies
capture an ataxin-1 interactome in Neuro-2a cells. Sci Data. 2018;5:180262.
61.   Garcia-Garcia MJ, Eggenschwiler JT, Caspary T, Alcorn HL, Wyler MR, Huangfu D,
et al. Analysis of mouse embryonic patterning and morphogenesis by forward genetics. Proc
Natl Acad Sci U S A. 2005;102(17):5913-9.
62.   Caspary T, Larkins CE, Anderson KV. The graded response to Sonic Hedgehog
depends on cilia architecture. Dev Cell. 2007;12(5):767-78.
63.   Graziotto JJ, Farkas MH, Bujakowska K, Deramaudt BM, Zhang Q, Nandrot EF, et
al. Three gene-targeted mouse models of RNA splicing factor RP show late-onset RPE and
retinal degeneration. Invest Ophthalmol Vis Sci. 2011;52(1):190-8.
64.   Krausova M, Kreplova M, Banik P, Cvackova Z, Kubovciak J, Modrak M, et al.
Retinitis pigmentosa-associated mutations in mouse Prpf8 cause misexpression of circRNAs
and degeneration of cerebellar granule cells. Life Sci Alliance. 2023;6(6).
65.   Farkas MH, Lew DS, Sousa ME, Bujakowska K, Chatagnon J, Bhattacharya SS, et al.
Mutations in pre-mRNA processing factors 3, 8, and 31 cause dysfunction of the retinal
pigment epithelium.
66.   Bonnafe E, Touka M, AitLounis A, Baas D, Barras E, Ucla C, et al. The transcription
factor RFX3 directs nodal cilium development and left-right asymmetry specification.
67.   Cantagrel V, Silhavy JL, Bielas SL, Swistun D, Marsh SE, Bertrand JY, et al.
Mutations in the cilia gene ARL13B lead to the classical form of Joubert syndrome. Am J
Hum Genet. 2008;83(2):170-9.
68.   Thomas S, Cantagrel V, Mariani L, Serre V, Lee JE, Elkhartoufi N, et al.
Identification of a novel ARL13B variant in a Joubert syndrome-affected patient with retinal
impairment and obesity. Eur J Hum Genet. 2015;23(5):621-7.
69.   Spahiu L, Behluli E, Grajcevci-Uka V, Liehr T, Temaj G. Joubert syndrome:
Molecular basis and treatment. J Mother Child. 2022;26(1):118-23.
70.   Gana S, Serpieri V, Valente EM. Genotype-phenotype correlates in Joubert
syndrome: A review. Am J Med Genet C Semin Med Genet. 2022;190(1):72-88.
71.   Engel BD, Ludington WB, Marshall WF. Intraflagellar transport particle size scales
inversely with flagellar length: revisiting the balance-point length control model. J Cell Biol.
2009;187(1):81-9.
72.   Pazour GJ, Dickert BL, Vucica Y, Seeley ES, Rosenbaum JL, Witman GB, Cole DG.
Chlamydomonas IFT88 and its mouse homologue, polycystic kidney disease gene tg737, are
required for assembly of cilia and flagella. J Cell Biol. 2000;151(3):709-18.
73.   Krock BL, Perkins BD. The intraflagellar transport protein IFT57 is required for cilia
maintenance and regulates IFT-particle-kinesin-II dissociation in vertebrate photoreceptors. J
Cell Sci. 2008;121(11):1907-15.
74.   Hou Y, Qin H, Follit JA, Pazour GJ, Rosenbaum JL, Witman GB. Functional analysis
of an individual IFT protein: IFT46 is required for transport of outer dynein arms into
flagella. J Cell Biol. 2007;176(5):653-65.
75.   Dupont MA, Humbert C, Huber C, Siour Q, Guerrera IC, Jung V, et al. Human IFT52
mutations uncover a novel role for the protein in microtubule dynamics and centrosome
cohesion. Hum Mol Genet. 2019;28(16):2720-37.
76.   Ha S, Lindsay AM, Timms AE, Beier DR. Mutations in Dnaaf1 and Lrrc48 Cause
Hydrocephalus, Laterality Defects, and Sinusitis in Mice. G3 (Bethesda). 2016;6(8):2479-87.
77.   McCafferty CL, Papoulas O, Lee C, Bui KH, Taylor DW, Marcotte EM, Wallingford
JB. An amino acid-resolution interactome for motile cilia identifies the structure and function
of ciliopathy protein complexes. Dev Cell. 2025;60(6):965-78 e3.

78.   Rashid S, Breckle R, Hupe M, Geisler S, Doerwald N, Neesen J. The murine Dnali1
gene encodes a flagellar protein that interacts with the cytoplasmic dynein heavy chain 1.
Mol Reprod Dev. 2006;73(6):784-94.
79.   Fassad MR, Rumman N, Junger K, Patel MP, Thompson J, Goggin P, et al. Defective
airway intraflagellar transport underlies a combined motile and primary ciliopathy syndrome
caused by IFT74 mutations. Hum Mol Genet. 2023;32(21):3090-104.
80.   Bakey Z, Cabrera OA, Hoefele J, Antony D, Wu K, Stuck MW, et al. IFT74 variants
cause skeletal ciliopathy and motile cilia defects in mice and humans. PLoS Genet.
2023;19(6):e1010796.
81.   Wheway G, Parry DA, Johnson CA. The role of primary cilia in the development and
disease of the retina. Organogenesis. 2014;10(1):69-85.
82.   Canty EG, Lu Y, Meadows RS, Shaw MK, Holmes DF, Kadler KE. Coalignment of
plasma membrane channels and protrusions (fibripositors) specifies the parallelism of tendon.
J Cell Biol. 2004;165(4):553-63.
83.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.
Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.
84.   Henrique D, Adam J, Myat A, Chitnis A, Lewis J, Ish-Horowicz D. Expression of a
Delta homologue in prospective neurons in the chick. Nature. 1995;375(6534):787-90.
85.   Thisse B, Pflumio, S., Fürthauer, M., Loppin, B., Heyer, V., Degrave, A., Woehl, R.,
Lux, A., Steffan, T., Charbonnier, X.Q. and Thisse, C. Expression of the zebrafish genome
during embryogenesis (NIH R01 RR15402). [ZFIN Direct Data Submission]. In press 2001.
86.   Wood KA, Rowlands CF, Qureshi WMS, Thomas HB, Buczek WA, Briggs TA, et al.
Disease modeling of core pre-mRNA splicing factor haploinsufficiency. Hum Mol Genet.
2019;28(22):3704-23.
87.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. 2011. 2011;17(1):3.
88.   Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets.
Bioinformatics. 2011;27(6):863-4.
89.   Frankish A, Diekhans M, Ferreira AM, Johnson R, Jungreis I, Loveland J, et al.
GENCODE reference annotation for the human and mouse genomes. Nucleic Acids Res.
2019;47(D1):D766-D73.
90.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
91.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
92.   Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and
better for alignment and quantification of RNA sequencing reads. Nucleic Acids Res.
2019;47(8):e47.
93.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2010;26(1):139-40.
94.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.
95.   Kunkel TA, Roberts JD, Zakour RA. Rapid and efficient site-specific mutagenesis
without phenotypic selection. Methods Enzymol. 1987;154:367-82.
1154

1155

Figures and Legends

Fig 1. Genetic analysis of *l11Jus27* mice. A) Propidium iodide stained E9.5 *l11Jus27*

heterozygote showing correct leftward looping of heart (heart tube outlined in white; lv= left

ventricle), *l11Jus27* homozygote showing reversed looping of heart (heart tube outlined in

white; lv = morphological left ventricle), *l11Jus27* homozygote showing open neural tube (blue

arrow) and distended heart tube (orange arrow). B) Sanger sequencing of individual mouse

embryos confirming that the N1531S substitution is present in heterozygous embryos and

homozygous embryos carrying the *l11Jus27* phenotype, but not in wild type mice from either

a 129S5 (129) or C57BL/6 (B6) background. C) qPCR quantification of *Prpf8* expression

levels in *Prpf8^{N1531S}* heterozygous and homozygous mutant whole embryos at E10.5. Results

from the average of three biological and three technical replicates are presented. Expression

levels were normalised to *Gapdh* expression. D) Clustal analysis of PRPF8 protein orthologues

in mouse, zebrafish, human and *Saccharomyces cerevisiae* showing that residue 1531 is

conserved across multiple eukaryotes. Identical residues are highlighted in yellow. E) Protein

structure of PRPF8 with amino acid replacement at residue 1531 shown in red. F) Western blot

confirms PRPF8 protein is present in *Prpf8^{N1531S}* heterozygotes and homozygous mutant whole

embryos with either correct heart looping (CL) or reversed heart looping (RL) at E10.5. G)

Quantification of PRPF8 protein levels from Western blots. Each sample was normalised to

GAPDH protein expression levels. H) Western blot analysis of mammalian cells transfected

with epitope tagged PRPF8 constructs at specified time points after cycloheximide treatment.

I) Quantification of PRPF8 expression levels normalised to ACTB for wild type and

PRPF8^{N1531S} alleles following cycloheximide treatment. J) PRPF8 WT and N1531S protein

half-life percentage decrease as compared to protein levels present at 0h. Numbers report

percentage of original protein remaining at each time point. Differences in degradation rate are

non-significant (Wilcoxon test p=0.125).

1182

1183

Fig 2. Analysis of *Prpf8^{N1531S}* heterozygous and homozygous node cilia. A-D) SEM images

of 2-3 somite stage *Prpf8^{N1531S}* heterozygous (A, C) and homozygous (B, D) mouse embryos.

A) The pit shape of the node (yellow boxes) is obvious in *Prpf8^{N1531S}* heterozygotes.

Prpf8^{N1531S} mutant embryos (B) have a flat node. Scale bar = 100μm. A', B') Higher

magnification images of boxed region in A and B. The node (yellow outline) is ciliated in both

genotypes. Scale bar = 10um. C-D) Higher magnification images of the node at 2-3 somite

stage. No gross morphological difference in cilia was observed between genotypes. Scale bar

= 5μm. E) Heterozygotes and homozygotes have equivalent numbers of nodal cilia at the LHF

and 2-3 somite stage (unpaired t-test). F) Cell proliferation was unaltered between genotypes

(unpaired t-test). G-H) Mouse node cilia labelled with acetylated tubulin in heterozygous

control (G) and $Prpf8^{N1531S}$ mutant embryos (H). Nuclei are stained with DAPI. I) Measurement

of node cilia length reveals that $Prpf8^{N1531S}$ mutant cilia are significantly shorter than

heterozygous controls (unpaired t-test p=0.0408). J-K) Tracking the travel over time of five

microbeads in the node revealed directional movement in $Prpf8^{N1531S}$ heterozygotes (J) but only

Brownian motion in $Prpf8^{N1531S}$ homozygotes (K). Direction of movement is indicated with an

arrow. A= anterior, L = left. L) Quantification of motile nodal cilia in $Prpf8^{+/+}$, $Prpf8^{N1531S}$

heterozygous and $Prpf8^{N1531S}$ homozygous embryos at the 2-3 somite stage revealed significant

differences between the three genotypes (p=0.0058, Welch's ANOVA). Post-hoc testing with

Dunnett's T3 test showed that the number of motile cilia in $Prpf8^{N1531S}$ homozygotes was

significantly lower compared to $Prpf8^{+/+}$ (p=0.0222) and $Prpf8^{N1531S/+}$ embryos (p=0.0487),

but not between $Prpf8^{N1531S/+}$ and $Prpf8^{+/+}$ embryos (p=0.9332).

1206

1207

1208

1209

1210

Fig 3. Wholemount *in situ* hybridisation expression analysis of laterality genes in

Prpf8^{N1531S} **heterozygous and homozygous mutant embryos.** A) *Prpf8^{N1531S}* heterozygote

showing *Shh* midline and node expression (black arrow). B) *Prpf8^{N1531S}* homozygote showing

intact *Shh* expression at the node (black arrow), but discontinuous midline staining (white

arrows). C) E8.5 *Prpf8^{N1531S}* heterozygote with upregulation of *Cerl2* on the right vs the left of

the node. D) *Prpf8^{N1531S}* homozygote with reversed *Cerl2* expression. E) Graphical

representation of *Cerl2* expression patterns seen in *Prpf8^{N1531S}* homozygotes revealing a high

degree of variability. F) E8.5 *Prpf8^{N1531S}* heterozygote showing *Nodal* expression pattern with

upregulation left of the node and left LPM. G) *Prpf8^{N1531S}* homozygote showing reduced and

reversed perinodal *Nodal* expression and no expression in the LPM. H) Graphical

representation of *Nodal* perinodal expression in *Prpf8^{N1531S}* homozygotes revealing variable

expression patterns. I) E8.5 *Prpf8^{N1531S}* heterozygote showing *Lefty1, 2* expression along the

midline and left LPM, detected with a probe complimentary to both mRNAs; no expression

was detected in *Prpf8^{N1531S}* homozygotes (J). K-P) *Pitx2* expression in the LPM (black

arrowheads) at E8.75 (K,L), E9.0 (M,N) and E9.5 (O,P) with $Prpf8^{N1531S}$ heterozygotes

showing left-sided *Pitx2* expression (K,M,O) and $Prpf8^{N1531S}$ homozygotes showing

randomised Pitx2 expression (L,N,P). Q) Graphical representation of data comparing sidedness

of LPM *Pitx2* expression with heart looping direction at E9.5 revealing a disconnect between

expression sidedness (*Pitx2*) and heart looping (heart). C= correct, R= reversed, B= bilateral,

A= absent. R) E9.5 expression of *Pitx2* in $Prpf8^{N1531S}$ homozygote showing discordance

between heart looping and *Pitx2* expression sidedness (arrow). S,T) E10.5 expression of *Barx1*

labelling the developing stomach in $Prpf8^{N1531S}$ heterozygote (S) and in $Prpf8^{N1531S}$

homozygote (T). Dashed line indicates the midline, and the heart is outlined in white. L= left,

R= right. Scale bars = 0.5mm.

1235

1236

Fig 4. Developmental laterality and cilia defects in *cph* zebrafish. A) Wild type zebrafish at

48hpf with straight body axis (left panel). MF20 staining to identify heart tube is shown in

green (right two panels). Red arrow shows looping direction of heart. B) *cph* mutant zebrafish

with curled down body axis (black arrow) and pericardial oedema (blue arrow) indicative of

cilia defects. MF20 staining shows delayed heart tube looping (green; right two panels). C) *cph*

mutant fish show a significance increase in heart looping defects compared to wild type

controls. D) Expression of the laterality gene *lefty2* in wild type zebrafish at 22hpf (white

arrowheads indicate location of gene expression in panels E-L). E) Gene expression patterns

of the laterality gene *lefty2* are altered in *cph* mutant fish, showing a variety of expression

patterns. F) Statistical summary of *lefty2* expression pattern defects in *cph* fish. G) Expression

of the laterality gene *southpaw* in wild type zebrafish at 18hpf. H) Gene expression patterns of

the laterality gene *southpaw* are altered in *cph* mutant fish, showing a variety of expression

patterns. I) Statistical summary of *southpaw* expression pattern defects in *cph* fish. J)

Expression of the laterality gene *dand5* in wild type zebrafish at 8hpf. K) Gene expression

patterns of the laterality gene *dand5* are altered in *cph* mutant fish, showing a variety of

expression patterns. L) Statistical summary of *dand5* expression pattern defects in *cph* fish. M)

Acetylated tubulin (green) and gamma tubulin (blue) staining in wild type fish KV (left) and

cph fish KV (right). DAPI staining for nuclei is shown in red (Scale bar = 10µ). N)

Quantification of average cilia number per KV between wild type control and *cph* fish. O)

Quantification of average cilia length within individual KVs demonstrates *cph* KV cilia are

significantly shorter than wild type KV cilia (n=14 and n=18 KVs respectively). Data are

represented as mean ± SD. Significance was evaluated using unpaired t-test. P<0.0001 P) Still

images from videos showing particle movement in wild type fish KV (representative of n=5,

Scale bar = 10µm, Time lapsed = ~3s). Q) Still images from movies showing particle

movement in *cph* fish KV (representative of n=5, Scale bar = 10µm, Time lapsed = ~3s). Green

and red arrows indicate the start and end position of the bead, respectively. L left, R right.

1263

1264

Fig 5. Investigation of Prp8 N1603S substitution phenotype in yeast. A) Growth of haploid

yeast strains with genomic *PRP8* deletion and complementing pRS413 plasmids with either no

PRP8, PRP8 wild-type (PRP8*) or PRP8 N1603S. Left sector: Strains transformed with empty

pRS413 fail to grow, confirming genomic *PRP8* deletion is non-viable; middle and right

sectors: yeast strains are viable when transformed with pRS413 containing either wild type or

mutated (N1603S) *PRP8*. Plates were grown at 30°C. B) Primer extension analysis in Prp8 and

Prp8[N1603S] strains containing *ACT1-CUP1* reporters. Top panel displays lariat accumulation in

the *ACT1-CUP1* A259C branch site reporter in the Prp8[N1603S] strain. Bottom panel displays

depletion of mature mRNA product in the *ACT1-CUP1* A302U 3'SS reporter in the Prp8[N1603S]

strain. These results are consistent with the N1603S mutation representing a first step allele of

Prp8. C) Comparison of splicing efficiencies using *ACT1-CUP1* splicing reporters. More

efficient splicing of a reporter correlates with increased copper resistance and therefore growth

at higher concentrations of copper sulphate (CuSO4). Haploid yeast strains with *CUP1* and

PRP8 deletions with either wild type or N1603S Prp8 complementing plasmids were

transformed with *ACT1-CUP1* splicing reporters containing mutations at the 5'SS (G5A), BPS

(C256A and A259C) and the 3'SS (A302G and A302U) and grown on plates with increasing

concentrations of CuSO4; in each case Prp8 N1603S was less efficient at splicing the reporter

compared to wild type Prp8. Graphs indicate the highest concentration of copper where all

three replicates survived.

Fig 6. *Arl13b* transcript alterations present in *Prpf8* mutant mice and fish. A) Sashimi plot

illustrating relative read coverage of splice junctions between exons 8 and 10 of the murine

Arl13b transcript in three WT mice (top, green) and three *Prpf8^N1351S^* homozygous mouse

mutants (bottom, red). Numbers represent the mean read coverage across each junction for the

three replicates of each genotype. *Prpf8^{N1351S}* homozygous mutants exhibit a marked increase

in skipping of exon 9, producing greater levels of the *Arl13b* short isoform. B) Schematic

representation of ARL13B predicted protein products found in wild type embryos and

Prpf8^{N1531S} mouse mutants. The region of the transcript encoding amino acids 380-403 from

the wild type sequence is removed from the short isoform. C) RT-PCR using a single primer

pair displaying long and short isoforms of *Arl13b* in mouse (left) and zebrafish (right) embryos.

Quantification of the relative proportions of the *Arl13b* short and long isoforms plotted as fold

change relative to wild type are shown below the agarose gels. D) Sequencing chromatograph

of a*rl13b* RT-PCR product from wild type zebrafish (top) and *cph* embryos (bottom). The

region corresponding to exon 9 is highlighted in the long isoform sequence. The junction

between exon 8 and 10 is highlighted in the short isoform sequence. E) RT-PCR using a single

primer pair displaying *Arl13b* long and short isoform distribution in wild type and *Prpf8^{N1531S}*

heterozygous adult mouse organs. F) Quantification of the ratio of relative expression levels of

Arl13b long and short isoform in wild type and *Prpf8^{N1531S}* heterozygous adult mouse organs.

1304

1305

Fig 7. Analysis of *ARL13B* short and long isoform over-expression in human cells and

wild type zebrafish embryos. A) Co-localization of human acetylated tubulin (green) and

endogenous ARL13B (red; left panels); co-localization of transfected GFP-tagged human ARL13B short isoform (green) with acetylated tubulin (red; centre panels); co-localisation of transfected GFP-tagged human ARL13B long isoform (green) with acetylated tubulin (red; right panels) in hTERT-RPE1 cells. Nuclei are stained with DAPI (blue). Scale bar 5μm. B) Immunofluorescence images of KV cilia in three groups: wild type uninjected embryos (top), embryos injected with *arl13b-gfp* short isoform mRNA (middle), and embryos injected with *arl13b-gfp* long isoform mRNA (bottom). Cilia are stained with acetylated tubulin (labelling axonemes), and GFP to label the injected Arl13b protein isoforms. Nuclei are stained with DAPI. C) Quantification of average cilia length per KV in wild type uninjected control embryos (n=9); *arl13b* long isoform injected embryos (n=6), and *arl13b* short isoform injected embryos (n=6). No significant difference was detected between the *arl13b* long isoform injected embryos and the uninjected wildtype embryos (one-way ANOVA, $p > 0.05$). Cilia length in *arl13b* short isoform injected embryos was significantly decreased compared to uninjected wildtype embryos (one-way ANOVA, $p < 0.05$). Scale bar = 20μm. D) Quantification of cilia number per KV in control and injected embryos. No significant differences were detected between groups (one-way ANOVA $p > 0.05$). E) Heart directionality at 48hpf detected by MF20 immunofluorescence (red arrows). Some *arl13b* short isoform injected injected embryos show delayed cardiac development and have not achieved fully looped hearts at 48hpf, but resemble the jogging asymmetry typically present at 24hpf. Injection condition is shown above the images, observed heart directionality is shown on the images. F) A significant difference in heart defects frequency was presented among the three group (Chi-square test, $p < 0.0001$). Pairwise comparisons then were performed between groups. Embryos injected with the *arl13b* short isoform mRNA (n=50) displayed a significantly increased percentage of heart directionality defects compared with the *arl13b* long isoform mRNA injected (n=50) or wild type uninjected group (n=50)

(Fisher's exact test, p < 0.0001), while there is no significant difference of heart defect

frequency observed between the *arl13b* long isoform mRNA injected group and the wildtype

uninjected group (Fisher's exact test, p = 0.4949) . G) Brightfield image of wild type

uninjected embryo (bottom) at 48hpf lacking oedema in pericardium (orange arrowhead).

arl13b short isoform mRNA injected embryo (bottom) at 48hpf shows pericardial oedema

(orange arrowhead). H) Wild type embryos (n=50) do not display pericardial oedema at

48hpf, nor do embryos injected with the *arl13b* long isoform (n=50). Embryos injected with

the *arl13b* short isoform mRNA (n=50) showed a significant increase in pericardial oedema

compared with the uninjected control group (Fisher's exact test, p < 0.0001).

1343

1344

Fig 8. Analysis of *Prpf8* sbMO knockdown phenotypes and rescue by *arl13B* long

isoform mRNA co-injection. A) Immunofluorescence of KV cilia stained using an antibody

against acetylated tubulin labelling axonemes (green) in WT and *prpf8* sbMO knockdown

embryos. Nuclei are stained with DAPI (blue). Scale bar is 10μm. B) Quantification of cilia

length per KV in WT (n=10) and *prpf8* splice blocking morpholino (sbMO) knockdown

embryos (n=8). All cilia detected with acetylated tubulin staining were measured, and the

average length per KV is plotted. *prpf8* sbMO knockdown embryos show a significant

decrease in cilia length (Unpaired t-test, $p < 0.0001$). C) Quantification of cilia number per

KV showed no significant difference between groups (Unpaired t-test; $p = 0.4868$). D)

Cardiac looping morphogenesis detected by MF20 immunofluorescence. sbMO embryos and

arl13b short isoform mRNA rescue embryos showed heart looping defects. The co-injection

of *arl13b* long isoform mRNA with sbMO partially rescued heart looping defects. Control

MO injected embryos did not display heart directionality defects. E-H) Analysis of

phenotypes in wildtype uninjected embryos (n=94), *prpf8* sbMO knockdown embryos

(n=104), sbMO co-injection with *arl13b* short isoform mRNA (n=44), sbMO co-injection

with *arl13b* long isoform mRNA (n=79), and control MO injected embryos (n=84).

Statistical analysis was first performed using a Chi-square test to assess morphological

defects among five groups. Where significant difference was observed, pairwise comparisons

were conducted between two groups by Fisher's test. E) The sbMO knockdown embryos with

arl13b long isoform mRNA injection displayed a significantly decreased percentage of heart

directionality defects compared to the sbMO knockdown group (Fisher's exact test, $p <$

0.0001). F) The percentage of pericardial oedema at 48hpf is significantly decreased in the

arl13b long isoform mRNA injection rescue group compared to the sbMO knockdown group

(Fisher's exact test, $p < 0.0001$). G) The percentage of tail defects is significantly decreased

in the *arl13b* long isoform overexpression rescue group compared to the sbMO knockdown

group (Fisher's exact test, p < 0.0001). H) Embryo survival at 72 hpf is significantly

increased in the long isoform co-injection group compared to the sbMO knockdown group

(Fisher's exact test, p < 0.0001). I) Brightfield images of WT embryo, *prpf8* sbMO, *arl13b*

long isoform co-injected embryo, *arl13b* short isoform co-injected embryo and control MO

injected embryo. Black arrowheads indicate pericardial oedema, blue arrowheads indicate

head morphological defects, green arrowheads indicate eye defects, and red arrowheads

indicate tail defects seen in *prpf8* sbMO injected embryos. J) Network analysis of enriched

biological processes at GO Level 5 for proteins binding to the long isoform of ARL13B.

Many processes related to cilia function are present. K) Volcano plot of mass spectrometry

data showing proteins that are enriched for binding the long isoform of ARL13B compared to

the short isoform, based on a fold-change cutoff of 4 and an adjusted p-value cutoff of 0.05.

The X-axis represents the log2 fold change, and the Y-axis represents the log2 adjusted p-

value. Note that many members of the IFT-B complex are significantly enriched in this

comparison.

1385

1386