Testing for a wildlife reservoir of divergent SARS-CoV-2 in white-tailed deer

Lauren Crawshaw[1]*, Jonathon D. Kotwa[2], Simon P. Jeeves[3], Courtney Loomer[4], Antonia

Dibernardo[4], Ashley Stewart[1], Sasha L. Newar[1], Emily Chien[2], Winfield Yim[2], Peter

Kruczkiewicz[5], Oksana Vernygora[5], Oliver Lung[5,6], Albrecht I. Schulte-Hostedde[7], Larissa

Nituch[1], Finlay Maguire[2,8,9], Bradley Pickering[5,10], Claire M. Jardine[3,11], Heather Coatsworth[4],

Samira Mubareka[2,12,*†], and Jeff Bowman[1,*†]

[1] Wildlife Research and Monitoring Section, Ontario Ministry of Natural Resources, Peterborough, Ontario, Canada

[2] Sunnybrook Research Institute, University of Toronto, Toronto, Ontario, Canada

[3] Department of Pathobiology, University of Guelph, Guelph, Ontario, Canada

[4] National Microbiology Laboratory, Public Health Agency of Canada, Winnipeg, Manitoba, Canada

[5] National Centre for Foreign Animal Disease, Canadian Food Inspection Agency, Winnipeg, Manitoba, Canada

[6] Department of Biological Sciences, University of Manitoba, Winnipeg, Manitoba, Canada

[7] School of Natural Sciences, Laurentian University, Sudbury, Ontario, Canada

[8] Faculty of Computer Science, Dalhousie University, Halifax, Nova Scotia, Canada

[9] Department of Community Health and Epidemiology, Dalhousie University, Halifax, Nova Scotia, Canada

[10] Department of Medical Microbiology and Infectious Diseases, University of Manitoba, Winnipeg, Manitoba, Canada

[11] Canadian Wildlife Health Cooperative, Guelph, Ontario, Canada

[12] Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario, Canada

* Corresponding authors

lauren.crawshaw@ontario.ca

samira.mubareka@sunnybrook.ca

jeff.bowman@ontario.ca

[†]These authors jointly supervised this work.

Abstract

The 2021 discovery of a divergent lineage (B.1.641) of severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) in white-tailed deer (*Odocoileus virginianus*) from Ontario raised

concerns that deer were a potential reservoir. To assess whether white-tailed deer continued to

be infected with B.1.641 and to test for spillover into other species, we established a

surveillance program in Ontario by sampling wildlife via existing monitoring programs and

through active surveillance of captive and wild animals. Between 2022 to 2024, we tested 2,839

animals, identifying one active SARS-CoV-2 infection (a likely spillover of a recombinant

XBB.2.3.11.3 lineage), but no cases of B.1.641. Overall, 93 animals (6.8%) tested positive for

SARS-CoV-2 antibodies, including 89 white-tailed deer, two Virginia opossums (*Didelphis*

virginiana), one American mink (*Neogale vison)*, and one river otter (*Lontra canadensis*). In

Southwestern Ontario, where B.1.641 was originally detected, 15.2% of deer samples were

seropositive. Generalized Linear Models demonstrated that seropositive deer were more likely

to be found in areas with a higher fall deer harvest and human population density, and closer to

previous B.1.641 cases. Our data suggest that deer-associated B.1.641 may have caused a

relatively localized epizootic without forming a stable reservoir. This study underscores the

importance of One Health-focused surveillance.

Keywords: SARS-CoV-2; severe acute respiratory syndrome coronavirus 2; COVID-19;

spillover; zoonoses; zoonotic diseases; One Health; white-tailed deer; reservoir; genomics;

sequencing; serology; PCR

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is responsible for coronavirus

disease 2019 (COVID-19) and an estimated 7 million human deaths worldwide (World Health

Organization 2024). SARS-CoV-2 continues to be a major public health concern post-pandemic,

and the effectiveness of human-centred interventions, such as vaccination, risk being

undermined by a variant persisting outside of human populations. When a pathogen is able to

infect multiple host species, it is more likely to be an emerging human pathogen (Singh et al.

2023). Understanding which species are susceptible to SARS-CoV-2 can shed light on how the

virus is able to spill into humans (Palmer et al. 2021). SARS-CoV-2 has a broad host range,

infecting species from almost all orders of mammal (Abdel-Moneim and Abdelwhab 2020; Zhou

et al. 2020; Kuchipudi et al. 2023; Z.H. Tan et al. 2024). The propensity of SARS-CoV-2 to infect

numerous species can be partially explained by the virus' ability to bind to the angiotensin-

converting enzyme 2 (ACE2) host cell receptor (F. Wu et al. 2020), which is conserved across

many mammalian species (Damas et al. 2020; Zhou et al. 2020).

Given the potential for SARS-CoV-2 spillback from humans to animal populations, there

remains a pressing need to understand the risk of the virus infecting novel species (C.C.S. Tan

et al. 2024). Transmission of SARS-CoV-2 from humans into other mammalian species

occurred early in the pandemic. The first human-to-animal transmission events of SARS-CoV-2

occurred in mink farms of the Netherlands and Denmark in 2020, likely transmitted from farm

workers (Oreshkova et al. 2020; Oude Munnink et al. 2021; Wolters et al. 2022). Since then,

there have been many reports of infected animal populations, from both domesticated and wild

species. For example, in a study of wildlife in Virginia, USA, 8 of 23 native mammalian species

tested positive or seropositive. SARS-CoV-2 infection was strongly correlated with urbanization

and prevalence was high in synanthropic species such as Virginia opossums (*Didelphis*

virginiana) and raccoons (*Procyon lotor*) (Goldberg et al. 2024). Infection of 11 cattle in

Germany coincided with a large wave of the Delta variant of concern in humans in fall to early

winter 2021 (Wernike et al. 2022). In Nigeria, SARS-CoV-2 antibodies were detected in feral

dogs, rabbits, and pangolins (Agusi et al. 2024), in the context of an estimated 80%

seroprevalence among humans (Kolawole et al. 2022).

The broad host range of SARS-CoV-2 and frequent spillback from humans to wildlife has led to

concern regarding the establishment of a potential reservoir. We follow the suggestion of

Haydon et al. (2002) and define a reservoir as having two characteristics: an ability to maintain

infection with a zoonotic pathogen, and the presence of an interface with humans. Reservoirs

may undermine public health efforts and make strain-specific vaccination more challenging by

providing a continual source of virus capable of transmitting back into human populations

(Haydon et al. 2002), gaining genomic diversity through recombination with other variants (Wei

et al. 2021) or adaptive immune-evasive mutations. Reservoirs have the potential to spread a

pathogen back to vulnerable human populations even after the pathogen has been reduced or

eradicated (Haydon et al. 2002). Such is the case with Middle East respiratory syndrome

(MERS-CoV), a betacoronavirus that first emerged in 2012 in Jordan and Saudi Arabia. While

MERS outbreaks in human populations are typically self limiting (Pustake et al. 2022), the

pathogen is widely enzootic in dromedary camels (Mackay and Arden 2015), with camel-to-

camel transmission exclusively perpetuating the epidemic and evolution of the virus (Dudas et

al. 2018).

With the concern about SARS-CoV-2 spillback into wildlife, many jurisdictions in North America

initiated surveillance early in the pandemic. White-tailed deer (*Odocoileus virginianus,* hereafter

deer) were flagged as an important species for surveillance, given their abundance, highly

social and transient behaviour (Peterson et al. 2017), frequent contact with humans (Pratt et al.

2025), and high affinity for ACE2 virus binding (Damas et al. 2020). Challenge studies found

that deer fawns are not only highly susceptible to SARS-CoV-2, but they could transmit the

infection to other fawns through droplets or aerosol (Palmer et al. 2021; Martins et al. 2022),

and the virus can be transmitted vertically from doe to fetus (Cool et al. 2022). Surveillance

studies found infected deer in wild free-ranging herds in 23 US states (Feng et al. 2023), with

dozens of spillover events from humans and evidence of sustained and continual deer-to-deer

transmission (Hale et al. 2022; Kuchipudi et al. 2022; Marques et al. 2022; Pickering et al. 2022;

Caserta et al. 2023; Feng et al. 2023; McBride et al. 2023). SARS-CoV-2 antibodies were

detected in almost half the deer sampled (40%) across four US states in 2021, with the highest

prevalence in Michigan (67%) (Chandler et al. 2021). Reports have shown the circulation of

nearly extinct variants of concern like Alpha or Gamma (or both) persisting in deer many months

after their last detection in humans in the US population (Marques et al. 2022; Caserta et al.

2023; Feng et al. 2023; McBride et al. 2023).

Recognizing the potential for transmission of SARS-CoV-2 from humans to deer, we initiated

surveillance of free-ranging deer harvested by hunters in southern Ontario, Canada. Surveys in

autumn 2021 produced evidence of sustained deer-to-deer transmission within 80 km of

Michigan, and a divergent lineage of SARS-CoV-2 infecting deer, Phylogenetic Assignment of

Named Global Outbreak (PANGO) lineage B.1.641. This lineage contains 76 nucleotide

mutations relative to ancestral SARS-CoV-2 (Wuhan Hu.1) and 49 mutations compared to its

closest ancestor in Global Initiative on Sharing All Influenza Data (GISAID) as of March 2022,

making this one of the most divergent SARS-CoV-2 lineages sequenced at the time of

discovery. B.1.641 shares a common ancestor with human- and mink-derived sequences from

Michigan sampled one year prior (autumn 2020). The B.1.641 lineage also stands out for being

the first confirmed case of deer-to-human transmission documented up to that point, making it

the first confirmed wildlife-to-human spillover event of SARS-CoV-2. The associated human

case (ON-PHL-21-44225), which shared the majority (80/90) of mutations with the deer

samples, was detected in autumn 2021 in a person with known contact with deer a week prior to

symptom onset (Pickering et al. 2022).

Viruses that evolve in the same species show similar mutational spectra due to selective

pressure from the host environment (Shan et al. 2020; Wei et al. 2021; Rudar et al. 2024).

Analysis of mutational spectra in B.1.641, especially the prevalence of C-to-U base pair

changes, suggest that this virus evolved for some time in deer between 2020 and 2021.

Consequently, we infer that this lineage is deer-adapted, and has rapidly accumulated

mutations, suggesting the virus was able to propagate freely in a highly susceptible host

population (Pickering et al. 2022). Given the possibility of sustained infection of deer with

B.1.641, we were interested in the potential establishment of a deer reservoir of SARS-CoV-2,

and the role that a deer host might play in altering the evolutionary trajectory of this lineage.

Mutations in the spike protein account for nine of the 76 mutations of B.1.641 (Pickering et al.

2022). Spike protein mutations can improve affinity for ACE2 and expand the host range

(Oreshkova et al. 2020; Wei et al. 2021; McBride et al. 2023; Wang et al. 2023; Goldberg et al.

2024). Among US states with infected deer, the virus has shown repeated adaptation to the

deer host through substitutions in the spike protein and other proteins (Feng et al. 2023). The

potential for deer-adapted mutations to increase risk to humans remains a concern.

In response to the potential establishment of a reservoir of B.1.641 in deer in Ontario, we

initiated a surveillance program to sample a wide array of captive and wild animal species

across the same general area where the B.1.641 lineage had been discovered. We carried out

active surveillance in southern Ontario across a suite of mammal species, with a specific focus

on deer given the potential for reservoir establishment in this species. We examined the role of

deer as a host and reservoir, including whether B.1.641 has been sustained in deer or

transmitted into other species. In anticipation of finding more infected deer, we planned to

sequence viral genomes, aiming to characterize virus mutational spectra and build our

understanding of the host response. We hypothesized that sustained infection of deer with

B.1.641 would lead to viral genomes in deer closely related to B.1.641, rather than infection with

lineages more commonly circulating in the human population at the time of the study. Likewise,

spillover of B.1.641 from deer to other animal species would lead to the discovery of lineages

closely related to B.1.641.

Materials & methods

Sample collection

Between November 2019 and June 2024, we collected mid-turbinate, nasopharyngeal, or oral

swabs and blood samples from live and deceased animals across southern Ontario. Samples

were sourced through live-trapping, roadkill collection, existing ecological monitoring projects,

and from captive facilities (i.e. wildlife rehabilitation facilities, private zoos). Sampling priority

was given to species known to be susceptible to SARS-CoV-2, including cervids (Hale et al.

2022; Kuchipudi et al. 2022), mink (*Neogale vison)* (Oude Munnink et al. 2021), deer mice

(*Peromyscus maniculatus*), and white-footed mice (*Peromyscus leucopus*) (Bosco-Lauth et al.

2021; Fagre et al. 2021; Griffin et al. 2021) as well as species suspected to be susceptible (e.g.

canids, Virginia opossums, mustelids). We also opportunistically sampled mammalian species

as they were available, including those with unknown or limited evidence of susceptibility (e.g.

beaver, *Castor canadensis*). Two sizes of sterile swabs (Hardy Diagnostics™ Flex Minitip

Nasopharyngeal Flock Swab , Puritan™ PurFlock™ Ultra Flocked Swabs) (Thermo Fisher

Scientific Inc; https://www.fishersci.ca) were used to collect oral or nasal swabs, with the minitip

swabs used to collect oral swabs from animals the size of *Peromyscus*.

Wildlife rehabilitators and zoos

Mid-turbinate or oral swabs were collected from captive animals at 11 registered wildlife

rehabilitators and five regional zoos. The type of swab collected depended on the species, size,

and ability to humanely access the nasal passage. To limit the demand on the facility and

number of samples, only one or two animals were tested from litters of juveniles, and neonates

were excluded. While most of the samples were collected by the visiting research personnel, six

facilities were also provided with sampling kits and a protocol to collect additional samples.

Pre-existing surveillance and ecology programs

Most samples were collected through pre-existing wildlife projects, with the largest source being

deceased deer tested through Ontario's Chronic Wasting Disease (CWD) surveillance program,

as described by Pickering et al. (2022). CWD testing is carried out each November and

December in a subset of administrative areas within Ontario, known as Wildlife Management

Units (WMUs). Each year a handful of WMUs are selected for CWD surveillance based on a

provincial CWD risk model (Ontario Ministry of Natural Resources 2019). Every deer is assigned

to one 10 x 10 km square location where it was harvested, which corresponds to a vector grid

layer sourced from the Ontario Breeding Bird Atlas (Birds Canada 2024). Since 2021, SARS-

CoV-2 molecular detection and serology have been incorporated into the province's seasonal

CWD project through the collection of nasopharyngeal swabs and blood samples. Archival

retropharyngeal lymph nodes (RPLN) and blood collected in 2019 and 2020 were also

retrospectively tested.

Opportunistic sampling for SARS-CoV-2 took place during ongoing ecological studies run by the

Ontario Ministry of Natural Resources (MNR) and Trent University. During the MNR's rabies

vaccine serosurveillance program, SARS-CoV-2 swabs and blood were collected from target

species (i.e., raccoons and striped skunks, *Mephitis mephitis*) in the fall of 2020 and 2021, and

from any mink or fox bycatch during their trap-vaccinate-release (TVR) project in the summer

and fall of 2022, 2023, and 2024. Between 2022 and 2024, other projects involved in collecting

SARS-CoV-2 samples included projects placing GPS transmitters on eastern wild turkeys

(Meleagris gallopavo silvestris), deer, moose (*Alces alces)*, the Algonquin wolf (*Canis lupus*

lycaon), and eastern coyote (*Canis latrans*), as well as wild boar removed from the landscape

(*Sus scrofa*). The majority of *Peromyscus* samples came from trap lines in Algonquin Provincial

Park as part of the Algonquin Small Mammal Project (Fryxell et al. 1998).

Roadkill and carcasses

From 2022 to early 2024, swabs and blood were collected from any fresh target species found

dead on roadsides during live-trapping work (i.e., deer, skunk, red fox, mink). This also included

any deer roadkill picked up and reported by participating municipal patrol lots. When possible,

blood was collected directly from the heart muscle. Four deceased deer were sampled for

SARS-CoV-2 after strange behaviour prior to death, two of which tested positive for epizootic

hemorrhagic disease. Additionally, local trappers provided carcasses of mink and other

furbearers (beaver (*C. canadensis*) , fisher (*Pekania pennanti*), river otter (*Lontra canadensis*))

between 2021 and 2023 where tissue samples, swabs, and blood were collected for SARS-

CoV-2 testing.

Live-trapping study area and trapping risk model

We carried out active surveillance through live-trapping of small mammals and mesocarnivores.

Given the geographic expanse of southern Ontario, the challenge in accessing wildlife hosts,

and the low prevalence of SARS-CoV-2 up to May 2021 (Greenhorn et al. 2022), we built an

ecological model to predict where humans, wildlife, and the virus were more likely to be in

contact (Kuchipudi et al. 2023), which narrowed the geographic scope for active surveillance.

Our model included: (1) proximity to the 17 SARS-CoV-2 B.1.641 PCR-positive deer from 2021

(Pickering et al. 2022); (2) approximate distance to mink farms and (3) deer farms; (4) distance

to the state of Michigan (U.S. Census Bureau 2018); and (5) an independently derived estimate

of terrestrial landscape connectivity referred to as "current density" (Pither et al. 2023).

Spatial analyses were carried out using ArcGIS Pro 3.0.3 (Esri 2022). Euclidean distance was

calculated for the three vector-based point layers (i.e., deer and mink farms, deer PCR cases) to

the same resolution and coordinate system. Pixel values of every raster (100 m resolution) were

then reclassified to a scale from 10 to 1, with higher scores representing higher potential risk of

SARS-CoV-2 transmission (e.g., the closer to a deer or mink farm, the higher the score). The

spatial grid vector layer (Birds Canada 2024) used for CWD surveillance was used as the

sampling unit, and mean zonal statistics were calculated for each grid cell for all five variables.

Each grid cell was then averaged to obtain a mean risk score, with proximity to the 17 PCR-

positive deer given double the weight as the other variables, given our priority to sample near

previous B.1.641 positive cases:

$$Risk\ Score\ = \frac{Dist_{Mink\ Farm}\ +\ Dist_{Deer\ Farm}\ +\ Dist_{Michigan}\ +\ Current\ Density\ +\ 2 \cdot Dist_{PCR-Positive\ Deer}}{5}$$

The results from the risk model identified an area with the highest risk score (mean score >8),

covering 17,000 km^2, which included the metropolitan area of the City of London, Ontario. This

risk area was prioritized for sample collection via live-trapping. The trapping zone included a

Komoka Provincial Park and six provincial green space management areas, called

Conservation Authorities, which intersected the Thames River waterways.

Live-trapping surveys

Live trapping was carried out across a 5,194 km² region of southern Ontario over three years

(June - October 2022 and 2023, one week in February 2024). Many trapping sites from 2022

were revisited in 2023. During the trapping period, small mammals (i.e., rodents, shrews) and

mesocarnivores (i.e., raccoons, skunks, opossum, mink) were live-trapped on private land and

government managed greenspace, with permission obtained through informed consent by

landowners, parks, and conservation authorities. Various sizes of Tomahawk live trap (sizes

102, 103, 105, and 106) (Tomahawk Live Trap; https://www.livetrap.com) were baited with

sardines and scent lures. Small mammals and shrews were trapped using Sherman live traps

(25.4 cm x 7.6 cm x 7.6 cm) (H.B. Sherman Traps Inc; https://shermantraps.com) baited with a

mix of sunflower seeds, peanuts, dried insects, and seafood. Each day, 40-60 Tomahawk traps

and 30-40 Sherman traps were set in riparian or forest habitat for four consecutive nights per

week. Traps were placed in sheltered areas for protection from the elements and secured with

cables to prevent rolling. Traps were checked each morning and reset for the next day, with

Sherman traps closed during the day and reset in the evenings. If evening temperatures fell

below -20°C, traps would remain closed overnight. Adequate thermal conditions for trapping in

the fall and winter were maintained by making sure all traps were sheltered from precipitation

and wind by covering traps with plastic and supplying each trap with either cotton bedding or

straw. For animals sampled on the Algonquin Small Mammal Project, Sherman and Longworth

(Longworth Scientific Instrument Co; https://www.penlon.com) live traps were baited with water-

soaked sunflower seeds set on established transects between May and August of each

sampling year.

Ethics approval

Animal care and handling was approved for each year of the study under the MNR Wildlife Animal Care Committee ((WACC Protocols #23-358 and #24-491), the Trent University Animal Care Committee (#28123), and the Laurentian University Animal Care Committee (#6011106), Ontario, Canada. Live-trapping and sampling of animals was conducted in accordance with applicable laws and regulations.

Animal handling

To obtain samples, animals were carefully restrained in the trap or transferred into a semi-soft funnel or large plastic bag, allowing for gentle immobilization of the animal during oral swabbing and marking. Trap locations were changed every week, while the Algonquin Small Mammal Project used the same established trap lines each year. Every capture was marked with livestock paint or chalk, unless a trapping site was used for more than one consecutive week, in which case an ear tag was applied instead (sizes 1005-1 for *Peromyscus*, skunk and mink; and 1005-3 for raccoons) (National Band & Tag Company; https://www.nationalband.com). Trappers wore medical grade face masks and nitrile gloves during processing to prevent the possible spread of SARS-CoV-2, and tools (pliers, clippers) were disinfected between animals.

Sample collection

Swabs were gently inserted into the mouth or nasal mid-turbinates and rotated for three to five seconds. Ungulates were the only animals where it was possible to humanely swab the nasal cavity, and nasopharyngeal samples were only collected from deceased deer. A small subset of deer, mink and *Peromyscus* had an additional exterior rectal swab taken, given potential of detecting the virus in *Peromyscus* (Griffin et al. 2021), deer (Martins et al. 2022) and ferrets (*Mustela furo*) (Gortázar et al. 2021) through this route. Where possible, live-trapped

mesocarnivores had a small blood sample (0.05-0.2 ml) collected on a piece of Advantec

Nobuto filter paper from the tail artery or a nail clipping (Advantec MFS, Inc;

https://www.advantecmfs.com). Additional tissues collected over the project included a small

section of the right and left RPLNs from CWD-tested deer (2019-2020), cloacal swabs from wild

turkeys, fecal samples from live-trapped deer, and lung and intestinal tissues from harvested

mink.

Swab tips were stored in sterile 2 ml tubes containing 1 ml of universal transport medium (UTM;

Sunnybrook Research Institute). Organs, RPLNs, and fecal material were stored in sterile dry

tubes. All samples taken in the field were kept on ice packs until they could be transferred to

storage at -80 °C. Blood samples collected on filter paper were allowed to air dry then placed in

coin envelopes before eventually being frozen at -80 °C.

298 **Diagnostic testing**

299 *RNA extraction and PCR testing*

The majority of RNA extractions were undertaken by staff at the Sunnybrook Research Institute

(Toronto, Ontario, Canada) and performed using 140 µL of swab sample spiked with Armored

RNA enterovirus (Asuragen; https://www.asuragen.com) using the Nuclisens EasyMag using

Generic Protocol 2.0.1 (bioMérieux Canada Inc., St-Laurent, QC, Canada) according to

manufacturer's instructions. Tissue samples were thawed, weighed, minced with a scalpel, and

homogenized in 600 µL of lysis buffer using the Next Advance Bullet Blender (Next Advance,

Troy, NY, USA) and a 5 mm stainless steel bead at 5 m/s for 3 minutes. RNA from 30 mg tissue

samples was extracted via the Nuclisens EasyMag using Specific Protocol B 2.0.1; RNA was

eluted in 50 µL. RNA extractions for *Peromyscus* swab samples in UTM were performed at the

Department of Pathobiology, University of Guelph (Guelph, Ontario, Canada). RNA was

extracted from 140 µl of swab UTM using the QIAamp Viral RNA Mini Kit (QIAGEN;

https://ww.qiagen.com) according to the manufacturer's protocol. Extractions were spiked with 1

µl Armored RNA Enterovirus (Asuragen; https://www.asuragen.com). Final elution volume of

extracted RNA was 60 µl in AVE buffer. All RT-PCR was carried out by Sunnybrook Hospital

Research Institute (SRI).

A subset of certain species (i.e. mink, fox, skunk), or animals exhibiting abnormal behaviour

were screened for highly pathogenic avian influenza virus (HPAIV) by the Animal Health Lab

(University of Guelph, Ontario). Only HPAIV-negative samples were submitted to SRI. RT-PCR

was used for SARS-CoV-2 RNA detection as previously described by Pickering et al. (2022);

two gene targets were used for SARS-CoV-2 RNA detection: the 5' untranslated region (UTR)

and the envelope (E) gene (Corman et al. 2020; LeBlanc et al. 2020). All samples were run in

duplicate and samples with cycle thresholds (Ct) < 40 for both SARS-CoV-2 targets and

armored RNA enterovirus in at least one replicate were considered positive. A subset of

Peromyscus spp. (N=282) samples were tested for SARS-CoV-2 RNA with a nested

pancoronavirus RT-PCR assay as previously described (Kotwa et al. 2025).

SARS-CoV-2 antibody testing

Blood samples were shipped to the National Microbiology Laboratory in Winnipeg, Manitoba for

SARS-CoV-2 antibody testing. Once received, the strips were sorted and evaluated for degree

of saturation. The saturated portion of each strip was cut into 4-5 pieces using sterile scissors

and dropped into a 2 ml sample tube and eluted in 1X phosphate-buffered saline (PBS) at 4°C

overnight. Fully saturated (100%) strips were eluted in 0.360 ml of PBS, 75% saturated strips in

0.270 ml of PBS and 50% saturated strips in 0.180 ml of PBS. These elution volumes were

considered to be equivalent to a single 10-fold dilution of serum. The eluates were tested with

the Genscript SARS-CoV-2 Surrogate Virus Neutralization Test (Catalog L00847-A) (Genscript;

https://www.genscript.com) following the manufacturer's instructions. A percent inhibition

greater than or equal to 30% was considered positive for the presence of SARS-CoV-2

neutralizing antibodies as per the kit protocol.

Whole-genome sequencing

Whole-genome sequencing on RT-PCR positive samples were conducted at SRI and was

performed as follows. LunaScript RT Supermix 5X kit (New England Biolabs, NEB;

https://www.neb.ca) was used to convert RNA samples to cDNA for amplification with the

SARS-CoV-2 ARTIC protocol. ARTIC v4.1 10μM primer pools were used in conjunction with Q5

High-Fidelity 2X Master Mix (NEB) for the PCR reaction. Cycling conditions were as follows:

98°C for 30s, 35 cycles of 98°C for 15s, and 63°C for 5min. Post amplicon generation, the PCR

products of the primer pools were combined followed by a 1X bead clean-up using AMPure XP

beads (Beckman Coulter; https://www.beckmancoulter.com). Amplicon concentrations were

determined using the Qubit dsDNA quantitation, High Sensitivity Kit (Thermo Fisher;

https://www.thermofisher.com) on a Qubit 4 Fluorometer. Sequencing libraries were generated

using the Illumina DNA prep kit as per manufacturer's instructions. Libraries were loaded on a

MiniSeq (Illumina; https://www.illumina.com) using the MiniSeq Mid Output Kit to perform

paired-end sequencing (2x149bp, 300 cycles). Original material from positive samples was sent

to the Canadian Food Inspection Agency (CFIA) for confirmatory RT-PCR testing.

Genomic analysis

Reference-mapped assembly and genomic analysis of paired-end Illumina reads was

conducted with the nf-core/viralrecon Nextflow workflow (v2.6.0) (Di Tommaso et al. 2017; Patel

et al. 2023). Briefly, nf-core/viralrecon ran the following steps: sequencing read quality control

with FASTQC (v0.11.9) (Andrews 2010); read quality filtering and adapter trimming with fastp

(v0.23.2) (Chen et al. 2018); read alignment to Wuhan-Hu-1 SARS-CoV-2 reference

(MN908947.3) (L. Wu et al. 2020) with Bowtie2 (v2.4.4) (Langmead and Salzberg 2012); NEB

VarSkip primer trimming from read alignments with iVar (v1.4) (Grubaugh et al. 2019); read

alignment coverage and statistics calculation with Mosdepth (v0.3.3) (Pedersen and Quinlan

2018) and Samtools (v1.16.1) (Li et al. 2009); variant calling and consensus sequence

generation with Bcftools (v1.16) (Danecek et al. 2021); variant effect prediction and annotation

with SnpEff (v5.0) (Cingolani, Platts, et al. 2012) and SnpSift (v4.3.1t) (Cingolani, Patel, et al.

2012); PANGO lineage (Rambaut et al. 2020) assignment with Pangolin (v4.3) (O'Toole et al.

2021) using UShER (v0.6.2) (Turakhia et al. 2021) Scorpio (v0.3.17) and Constellations

(v.0.1.2) (Colquhoun et al. 2023). The results from nf-core/viralrecon were summarized into an

XLSX report with xlavir (v1.0.2) (https://github.com/CFIA-NCFAD/xlavir). Nextclade (v3.12.0)

analysis was performed on the nf-core/viralrecon assembled consensus sequence of the PCR-

positive deer (wcov1242165) using the SARS-CoV-2 "nextstrain/sars-cov-2/wuhan-hu-1/orfs"

Nextclade dataset (last updated 2025-04-01 08:20:12 (UTC)) (Aksamentov et al. 2021).

Phylogenetic Analysis

Phylogenetic placement of the SARS-CoV-2 isolated from one positive deer was inferred

relative to other publicly available sequences from white-tailed and mule deer (N=811; GISAID

accessed 2025-03-19) as well as the 96 closest sequence matches recovered by UShER

(dataset of 16,884,360 genomes from GISAID, GenBank, COG-UK, and CNCB (2025-03-26)).

The Wuhan-Hu-1 (GenBank accession MN908947.3) sequence was included as an outgroup.

The multiple sequence alignment was performed using Nextclade v.3.12.0 and maximum

likelihood phylogenetic inference analysis was performed with IQ-TREE v.2.2.3 (Minh et al.,

2020). The best-fit nucleotide substitution model (GTR+F+R6) was determined by IQ-TREE

ModelFinder (Kalyaanamoorthy et al. 2017) under the Bayesian Information Criterion (BIC).

Node support was estimated using the ultrafast bootstrap and SH-aLRT with 1000 replicates for

each test (Hoang et al. 2017). Tree visualization was performed using FigTree

(http://tree.bio.ed.ac.uk/software/figtree) and Inkscape v.1.0.2 (https://inkscape.org).

Regression modeling

Understanding the spatial patterns of infection on the landscape can shed light on the current

state of SARS-CoV-2 in southern Ontario wildlife and identify risk factors for transmission. We

ran regression analysis on all serological data collected since 2022 (after the discovery of the

B.1.641 lineage) from deer south of latitude 47.3°N. In addition, we also explored biological

characteristics of the animals to see if there was any association with past infection. All

analyses were carried out in R (version 4.3.2) and RStudio (2024.04.02, Build 764) (R Core

Team 2024), and mapping and graphs in R and ArcGIS (Esri 2022).

Zonal statistics

Landscape variables, some of which were also used in the risk model, included: (1) proximity to

the 17 B.1.641 PCR-positive deer from 2021; (2) Euclidean distance to the state of Michigan

(km); (3) proportion of each grid cell classified as deer/non-deer habitat (Kennedy-Slaney et al,

unpublished report, 2018) derived by reclassifying the Ontario Land Cover Compilation Data

(Ontario Ministry of Natural Resources 2014); (4) annual fall deer harvest in Ontario by WMU

(Ontario Ministry of Natural Resources 2024); (5) current density (Pither et al. 2023); (6)

average cases of human SARS-CoV-2 in the fall of each year (2020-2023) (Ontario Agency for

Health Protection and Promotion (Public Health Ontario) 2024); and (7) human population

(Statistics Canada 2023a) (Table 1). Due to the limited data on deer abundance in Ontario,

annual deer hunting harvest was used as a proxy for deer abundance. The human population

variable was created by joining the 2021 census data (Statistics Canada 2023a) to a spatial

layer of the smallest census subdivisions, known as dissemination areas (Statistics Canada

2023b). Prior to running zonal statistics, the three polygon layers (human SARS-CoV-2 cases,

deer harvest, and deer habitat) were converted to raster grids at a 100 m resolution. Each deer

sample fell within one of the 10 x 10 km grid cells used in the risk model (Birds Canada 2024),

which were used as the main sampling units to extract a zonal statistic (a weighted mean) for

each of the landscape variables. In cases where the grid cells overlapped with large lakes (>5

km²), the grid cell layer was first clipped using a high resolution waterbody layer (Ontario

Ministry of Natural Resources 2020).

Sampling unit selection

Prior to running the models, a second sampling unit (30 x 30 km) was created to explore

possible bias of having a single 10x10 grid cell represent the presence of a highly mobile deer

species with a wide potential home range, with some yearling males reported to range over

6000 hectares (Lesage et al. 2000). This was particularly important where the true deer

locations occur near the edges of the 10 x 10 km grid cell. This second sampling unit was

created by buffering each grid cell by 10 km. Zonal statistics were repeated with these two

buffer sampling units for variables unrelated to distance and compared to the grid-based

sampling unit for correlation and relationship with the dependent variable.

Zonal mean of the larger sampling unit were highly correlated with the same variable run with

the grid (>0.75), indicating high similarities with the buffer and grid cell zonal statistics. Due to

this similarity, plus less multicollinearity between the original 10 x 10 grid variables of interest,

the original 10 x 10 km grid sampling unit was selected moving forward with the models.

Modeling

We used Generalized Linear Mixed Models (GLMMs) through R package *lme4* version 1.1-37

(Bates et al. 2015) to identify landscape features associated with a serological history of

infection in wild deer. The 10 x 10 km grid cell was used as the random effect to account for

non-independence between samples within these larger groups. Candidate models were

selected by first identifying highly correlated variables and retaining the variable that were

perceived as biologically significant or with greater univariate significance to the dependent

variable. Multicollinearity was checked by calculating the variance inflation factor (VIF) function

(R package *car* version 3.1-3) (Fox and Weisberg 2019) and comparing correlation plots (R

package *PerformanceAnalytics* version 2.0.8) (Peterson and Carl 2024). For example, high

correlation was observed between distance to PCR deer positives, distance to Michigan, and

deer habitat. Distance to deer PCR positives, being the primary variable of interest, was

retained in the model over distance to Michigan and deer habitat. The final full model included

five variables: (1) Distance to 17 PCR positive deer (2021); (2) Deer harvest in the fall; (3)

average COVID-19 cases reported in the fall by public health unit by year; (4) average human

population by year; and (5) year of sample (Table 1). Some variables were first rescaled by 10

or 1000 to allow the model to converge.

Table 1. Potential variables for use in the regression models. Some variables were removed
due to multicollinearity.
446

Class	Variable	Description/Biological relevance
Binary outcome	SARS-CoV-2 exposure*	Serology result (PVE+/NVE-) of each deer since January 2022. The dependent variable of interest.
Raster	Distance to 17 PCR+ deer*	Euclidean distance to 17 PCR+ deer of B.1.641 lineage sampled in 2021 (Pickering et al. 2022)
Vector to Raster	Annual fall deer harvest*	Average count of deer harvested each fall by WMU (2022-2023) (Ontario Ministry of Natural Resources 2024). Models were tested with each year, and an averaged variable containing all years.
Vector to Raster	Human population*	Mean human population derived from 2021 Census data (Statistics Canada 2023a) per dissemination area (Statistics Canada 2023b).
Vector to Raster	Human Density	Human population per square kilometre, derived from 2021 Census data (Statistics Canada 2023a) per dissemination area (Statistics Canada 2023b).
Vector to Raster	Human COVID-19 cases*	Daily change in human cases of COVID-19 reported to public health units each fall (September - December 2022-2023) (Ontario Agency for Health Protection and Promotion (Public Health Ontario) 2024). Models were tested with each year, and a variable containing all years.
Factor	Year*	Year sample was collected (2022-2024)
Raster	Distance to Michigan	Euclidean distance to the state of Michigan (U.S. Census Bureau 2018)
Raster	Deer habitat	Binary reclassification of the Ontario Land Cover Compilation Data (Ontario Ministry of Natural Resources 2014) as deer habitat or non-habitat (Kennedy-Slaney et al. 2018, unpublished)
Raster	Current density*	Ecological connectivity layer derived from costs from anthropogenic and natural barriers, with higher values equating lower cost for terrestrial animals to travel across the landscape (Pither et al. 2023)

*Variables used in final full model

For the GLMM models, samples with missing data for key variables were removed (i.e. samples

in 2024 where data was not yet available for COVID-19 cases and fall deer harvest). The one

deer that tested PCR positive but seronegative was retained in the serology data as a

"negative", as this wouldn't bias the results towards rejecting the null hypothesis that there was

no association between seropositivity and the landscape. The model was split into training and

testing data (20/80%; N=195/785). Rather than using an automated stepwise approach, which

is not recommended in GLMMs, model selection was driven by comparing the AIC of multiple

models, a measure of goodness of fit that penalizes overfitting. Using the *drop1* function of

package *lme4* (Bates et al. 2015), five new models were created from the full model using the

training data, each with just one variable was dropped. The model with the lowest AIC value

was retained and compared to the previous model using an ANOVA test. If the model reduced

the AIC and the ANOVA test was non-significant, this model was retained and the *drop1*

process repeated until the most parsimonious model remained. All figures were generated with

R package *ggplot2* (Wickham 2016), and all code and non-restricted datasets will be made

available on OSF upon publication.

Results

SARS-CoV-2 surveillance in southern Ontario

Between January 2022 and June 2024, 2,839 animals were tested for current or past infection

of SARS-CoV-2 across southern Ontario using serological or molecular analyses. Only animals

with location data were retained in the sample. The study area covered 148,460 km², with

SARS-CoV-2 infection detected in wildlife across 40,960 km² (28% of the study area). Among

the 15 projects that sourced samples between 2022 and 2024 (Table 2), 39.0% (N=1107) were

deer harvested through the CWD program run by MNR, followed by animals intentionally live-

trapped for SARS-CoV-2 or sampled through the MNR Rabies Program TVR project (N=482;

17%), animals sampled at wildlife rehabilitators (N=327; 11.5%), and *Peromyscus* sampled at

Algonquin Provincial Park (N =211; 7.4%). Overall, 76% of the animals sampled at wildlife

rehabilitators were orphaned juveniles. Surveillance of roadkill accounted for 5.2% of samples

collected (N=148). Between 2021 and 2023, 35 mink and 57 other furbearer carcasses (beaver,

fisher, river otter) were provided by local trappers for testing (N=92; 3.2%). The remaining

samples were collected on assorted ecology projects run by MNR or Trent University, or

ungulates submitted through the Canadian Wildlife Health Cooperative (CWHC).

Table 2. Breakdown of samples by project 2022-2024

Project	2022	2023	2024	Total	% All samples
Chronic Wasting Disease surveillance	626	481		1107	39.0%
Wildlife in rehabilitation	156	160	11	327	11.5%
SARS-CoV-2 surveillance live-trapping (2023)		288		288	10.1%
Algonquin Park Small Mammal Project	75	136		211	7.4%
SARS-CoV-2 surveillance live-trapping (2022)	156			156	5.5%
Roadkill, hunted or euthanized (not CWD)	47	85	16	148	5.2%
White-tailed deer live-trapping	96	44		140	4.9%
Rabies dRIT laboratory testing	133			133	4.7%
Zoo animals	87	23	8	118	4.2%
Mustelid and furbearer carcass samples	86	6		92	3.2%
Ungulates tested by CWHC	23	27	7	57	2.0%
Rabies TVR project	5	23		28	1.0%
Eastern coyote and Algonquin wolf live-trapping	13		1	14	0.5%
Highly Pathogenic Avian Influenza outbreak response live-trapping (February 2024)			10	10	0.4%
Wild turkey live-trapping	10			10	0.4%
Total	**1513**	**1273**	**53**	**2839**	100%

Overall, 95.7% of the samples since 2022 were native wildlife species (Table 3 and 4). White-

tailed deer were the most common species sampled (N = 1446; 50.9%). Among all cervids, the

majority were sourced through the CWD Program (1107; 72.2%), followed by live-trapping

(N=140; 9.1%), roadkill or hunted (N=84; 5.5%), zoos (N=76; 5.0%), wildlife rehabilitators

(N=69; 4.5%), and carcasses submitted to CWHC (N=57; 3.7%). Other frequently sampled

species were deer mice (N=211; 7.4%), raccoons (N=192; 6.8%), and striped skunks (N=155;

5.5). Sixty-eight mink were sampled (2.4%) between 2022 and 2024. Most animals were male

(50.1%, 40.5% female, 9.4% unknown) and adult (69.5%) (26.7% juveniles, 3.8% unknown).

Table 3. Summary of the serological tests by species and species class. In 2021, two deer were
both seropositive and PCR-positive.

	Species	2019	2020	2021	2022	2023	2024	Combined Sample 2019-2024)	Combined Sample (2022-2024)
				Total Samples (# Seropositive) / **%**					
NATIVE	White-tailed deer (*Odocoileus virginianus*)	50 (0)	160 (1) *1%*	89 (6) *7%*	463 (51) *11%*	517 (36) *7%*	11 (2) *18%*	1290 (96) *7%*	991 (89) *9%*
	Raccoon (*P. lotor*)		140 (0)	135 (0)	10 (0)	77 (0)	3 (0)	365 (0)	90 (0)
	Virginia opossum (*D. virginiana*)				33 (1) *3%*	65 (0)	7 (1) *14%*	105 (2) *2%*	105 (2) *2%*
	American mink (*N. vison*)			52 (0)	37 (0)	12 (0)	3 (1) *33%*	104 (1) *1%*	52 (1) *2%*
	Striped skunk (*M. mephitis*)		36 (0)	19 (0)	18 (0)	14 (0)		87 (0)	32 (0)
	North American beaver (*C. canadensis*)				39 (0)			39 (0)	39 (0)
	Eastern coyote (*Canis latrans*) or Algonquin wolf (*C. lupus lycaon*)				13 (0)	1 (0)	1 (0)	15 (0)	15 (0)
	North American river otter (*L. canadensis*)				11 (1) *9%*			11 (1) *9%*	11 (1) *9%*
	Wild turkey (*M. gallopavo*)				10 (0)			10 (0)	10 (0)
	Fisher (*P. pennanti*)				6 (0)	1 (0)		7 (0)	7 (0)
	Red fox (*V. vulpes*)				1 (0)	2 (0)		3 (0)	3 (0)
	White-footed mouse (*P. leucopus*)					2 (0)		2 (0)	2 (0)
	Long-tailed weasel (*Mustela frenata*)					1 (0)		1 (0)	1 (0)
	American red squirrel (*Tamiasciurus hudsonicus*)					1 (0)		1 (0)	1 (0)
EXOTIC	Barbary sheep (*Ammotragus lervia*)						5 (0)	5 (0)	5 (0)
	Eurasian wild boar (*S. scrofa*)						2 (0)	2 (0)	2 (0)
	Porcupine caribou (*Rangifer tarandus granti*)						2 (0)	2 (0)	2 (0)
	Linnaeus's two-toed sloth (*Choloepus didactylus*)						1 (0)	1 (0)	1 (0)
	Total	50 (0)	336 (1) *0%*	295 (6) *2%*	641 (53) *8%*	693 (36) *5%*	35 (4) *11%*	2050 (100) *5%*	1369 (93) *7%*

492   **Table 4.** Summary of PCR tests by species and species class. In 2021, two deer were both
493   seropositive and PCR-positive. The 17 positive deer in 2021 were those previously reported by
494   Pickering et al. (2022).

	Species	2019	2020	2021	2022	2023	2024	Combined Sample 2019-2024)	Combined Sample (2022-2024)
				Total Samples (#PCR Positive) / %					
NATIVE	White-tailed deer (*O. virginianus*)		238 (0)	406 (17) *4%*	774 (0)	613 (1) *0%*	29 (0)	2060 (18) *1%*	1416 (1) *0%*
	Raccoon (*P. lotor*)	169 (0)	189 (0)		102 (0)	87 (0)	3 (0)	550 (0)	192 (0)
	Striped skunk (*M. mephitis*)	51 (0)	138 (0)		115 (0)	38 (0)		342 (0)	153 (0)
	Deer mouse (*P. maniculatus*)				75 (0)	136 (0)		211 (0)	211 (0)
	Virginia opossum (*D. virginiana*)				58 (0)	83 (0)	7 (0)	148 (0)	148 (0)
	White-footed mouse (*P. leucopus*)				55 (0)	87 (0)		142 (0)	142 (0)
	American mink (*N. vison*)			53 (0)	43 (0)	22 (0)	3 (0)	121 (0)	68 (0)
	Eastern chipmunk (*Tamias striatus*)				19 (0)	48 (0)		67 (0)	67 (0)
	Big brown bat (*Eptesicus fuscus*)				10 (0)	46 (0)		56 (0)	56 (0)
	Red fox (*V. vulpes*)				28 (0)	21 (0)		49 (0)	49 (0)
	North American beaver (*C. canadensis*)				43 (0)			43 (0)	43 (0)
	Groundhog (*Marmota monax*)				14 (0)			14 (0)	14 (0)
	Meadow vole (*Microtus pennsylvanicus*)				3 (0)	10 (0)		13 (0)	13 (0)
	Eastern coyote (*C. latrans*) or Algonquin wolf (*C. lupus lycaon*)				9 (0)	3 (0)		12 (0)	12 (0)
	North American porcupine (*Erethizon dorsatum*)				7 (0)	5 (0)		12 (0)	12 (0)
	North American river otter (*L. canadensis*)				11 (0)			11 (0)	11 (0)
	Wild turkey (*M. gallopavo*)				10 (0)			10 (0)	10 (0)
	Fisher (*P. pennanti*)				7 (0)	2 (0)		9 (0)	9 (0)
	Moose (*Alces alces*)				7 (0)	2 (0)		9 (0)	9 (0)
	Meadow jumping mouse (*Zapus hudsonius*)				1 (0)	7 (0)		8 (0)	8 (0)
	Northern short-tailed shrew (*Blarina brevicauda*)				7 (0)			7 (0)	7 (0)
	Short-tailed weasel (*Mustela erminea*)				2 (0)	8 (0)		10 (0)	10 (0)
	American red squirrel (*T. hudsonicus*)				1 (0)	2 (0)		3 (0)	3 (0)
	Eastern grey squirrel (*Sciurus carolinensis*)				2 (0)			2 (0)	2 (0)
	House mouse (*Mus musculus*)				2 (0)			2 (0)	2 (0)
	Norway rat (*Rattus norvegicus*)				1 (0)	1 (0)		2 (0)	2 (0)
	American black bear (*Ursus americanus*)				1 (0)			1 (0)	1 (0)
	Elk (*Cervus canadensis*)				1 (0)			1 (0)	1 (0)
	Hoary bat (*Lasiurus cinereus*)				1 (0)			1 (0)	1 (0)
	Little brown bat (*Myotis lucifugus*)					1 (0)		1 (0)	1 (0)
	Long-tailed weasel (*N. frenata*)					1 (0)		1 (0)	1 (0)

Southern flying squirrel (*Glaucomys Volans*)			1 (0)			1 (0)	1 (0)
Eurasian fallow deer (*Dama dama*)			48 (0)	23 (0)		71 (0)	71 (0)
Plains bison (*Bison bison bison*)			8 (0)			8 (0)	8 (0)
Barbary sheep (*A. lervia*)					5 (0)	5 (0)	5 (0)
Domestic goat (*Capra aegagrus hircus*)			4 (0)			4 (0)	4 (0)
Domestic sheep (*Ovis aries*)			4 (0)			4 (0)	4 (0)
Red deer (*Cervus elaphus*)			2 (0)	2 (0)		4 (0)	4 (0)
Donkey (*Equus africanus*)			3 (0)			3 (0)	3 (0)
Pygmy goat (*C. aegagrus hircus*)			3 (0)			3 (0)	3 (0)
Capybara (*Hydrochoerus hydrochaeris*)			2 (0)			2 (0)	2 (0)
Eurasian wild boar (*S. scrofa*)					2 (0)	2 (0)	2 (0)
Kunekune pig (*Sus scrofa domesticus*)			2 (0)			2 (0)	2 (0)
Llama (*Lama glama*)			2 (0)			2 (0)	2 (0)
American miniature horse (*Equus caballus*)			2 (0)			2 (0)	2 (0)
Peccary (*Tayassu* genus)			2 (0)			2 (0)	2 (0)
Porcupine caribou (*R. tarandus granti*)					2 (0)	2 (0)	2 (0)
Alpaca (*Vicugna pacos*)			1 (0)			1 (0)	1 (0)
Bactrian camel (*Camelus bactrianus*)			1 (0)			1 (0)	1 (0)
Boer goat (*C. aegagrus hircus*)			1 (0)			1 (0)	1 (0)
Domestic pig (*S. scrofa domesticus*)				1 (0)		1 (0)	1 (0)
Linnaeus's two-toed sloth (*C. didactylus*)					1 (0)	1 (0)	1 (0)
Mink (farmed) (*N. vison*)				1 (0)		1 (0)	1 (0)
Red deer/elk hybrid	1 (0)					1 (0)	0 (0)
Total	459 (0)	786 (17) **2%**	1495 (0)	1250 (1) **0%**	52 (0)	4042 (18) **0%**	2797 (1) **0%**

(Left margin vertical category label: **EXOTIC**)

Serology and PCR results

Between January 2022 and June 2024, 93 animals tested positive for SARS-CoV-2 antibodies (Table 3), and viral RNA was detected through PCR testing in one deer in central Ontario in November 2023 (Table 4). The PCR-positive deer was seronegative and located over 235 km away from nearest PCR-positive deer case from 2021 (Figure 1). The majority of seropositive animals were detected in 2022 (N= 53; 57.0%), followed by 2023 (N= 36; 38.7%) and 2024 (N=4 ; 4.3%). Among the 93 seropositive animals, 89 were deer, and the remaining included new species detections for the region: one river otter (March 2022), two Virginia opossum

(October 2022, February 2024), and an American mink (February 2024) (Figure 2). Only one

animal sampled during live-trapping tested seropositive, a Virginia opossum sampled in

February 2024 near Windsor, Ontario, close to the Michigan/Ontario border. The two

serologically positive cases in February 2024 (mink, opossum) were separated by 480 km.

The PCR-positive deer was detected during annual CWD surveillance. The CWD project

accounted for 79.6% (N=74) of the serologically positive animal cases, followed by roadkill or

juvenile deer collected during CWD sampling (12.9%; N=12), live-deer trapping and collaring

(N=5; 5.3%), collected through furbearer trapping (1.1%; N=1), and during Avian influenza

sampling (1.1%; N=1). No animals from rabies surveillance, *Peromyscus* live-trapping, canid

live-trapping, ungulates submitted by CWHC, or any captive facilities were PCR positive or

seropositive.

Figure 1. Distribution of SARS-CoV-2 positive deer (89 seropositive and one PCR-positive deer; ID= wcov1242165) relative to the sampling effort among deer only within each 10 x 10 km grid cell. Provincial and state borders were sourced by Statistics Canada (Statistics Canada 2023c) and the U.S. Census Bureau (U.S. Census Bureau 2018).

554   **Figure 2.** Distribution of four SARS-CoV-2 serological positives mammals, excluding deer,
555   relative to the sampling effort of these species within each 10 x 10 km grid cell. Provincial and
556   state borders were sourced by Statistics Canada (Statistics Canada 2023c) and the U.S.
557   Census Bureau (U.S. Census Bureau 2018).

Sequence analysis

A high-quality genome sequence was assembled for the one PCR-positive deer (sample

wcov1242165) following sequencing and bioinformatics analysis with 99.6% genome coverage

at 10X or greater depth of coverage and 23,235X mean coverage depth. Both UShER and

Nextclade assigned wcov1242165 to the GS.3 PANGO lineage, alias of recombinant lineage

XBB.2.3.11.3, first detected in the USA in May 2023 according to GISAID (Elbe and Buckland-

Merrett 2017). Compared to Wuhan-Hu-1, the wcov1242165 genome had 33 synonymous

nucleotide substitutions and 75 amino acid substitutions, as well as 4 inframe deletions affecting

ORF1a, S and N gene amino acid sequences (ORF1a:del3675/3677, S:del25/27, S:del144,

N:del31/33) commonly found in lineage GS.3 sequences according to outbreak.info

(Gangavarapu et al., 2023). The wcov1242165 genome had several amino acid mutations not

typically found in GS.3 sequences (ORF1a:T403I, ORF1a:P3371S, ORF1a:L3588F,

ORF1a:A4092V, ORF1b:E1601D, S:A27S, S:G72E, ORF9b:P10S) according to outbreak.info.

Of these mutations, ORF1ab:A4092V, S:G72E and 6 nucleotide mutations (T3232C, C12540T,

C19113T, G21777A, C29659T, C29719T) were unique to wcov1242165 compared to nearest

neighbour sequences from UShER phylogenetic placement analysis against 16,884,360

genomes from GISAID, GenBank, COG-UK and CNCB (as of 2025-03-26) and the

sarscov2phylo 13-11-20 tree with newer sequences added by UShER.

Phylogenetic analysis recovered the new wcov1242165 sequence from deer deeply nested

within a clade of SARS-CoV-2 sequences from humans (Figure 3, A-B). The most closely

related viral sequences were from humans sampled in Ohio and Texas, USA, and Ontario,

Canada, in 2023. Phylogenetic analysis revealed that the new wcov1242165 genome from deer

was distinct from other publicly available deer-derived SARS-CoV-2 sequences, which were not

recovered in the same cluster of human-derived SARS-CoV-2 as the new sequence from deer.

Figure 3. Maximum likelihood phylogeny of SARS-CoV-2 genomes from white-tailed deer. (A) Phylogeny showing all publicly available SARS-CoV-2 genomes from deer (N=811) and (B) 96 closest matches to the WTD-0037-1242165 identified by UShER and a close-up of a subtree showing placement of the new viral sequence from deer deeply nested within a group of viral sequences from human hosts. Tips of the phylogeny are coloured according to the host type (human or deer). The new viral sequence from deer is labeled in red font.

Temporal and spatial patterns in SARS-CoV-2 detections

The majority (90.3%) of seropositive animals since 2022 were found in the region of

Southwestern Ontario (Figure 1 and 2). The seroprevalence among deer tested for antibodies

was high (15.2%) in this region. Among Virginia opossums, the only other positive animal in the

southwest, the prevalence was 2.0%. Overall, 10.5% (N=84) of animals tested for antibodies in

Southwestern Ontario were seropositive.

Prior to 2022, 7 deer (2.3% of deer sampled for antibodies) across the whole study area were

seropositive, compared to 89 deer during or after 2022 (9.0%) (Tables 3). Between 2019 and

2021, only deer and raccoons were tested, so we don't have a time comparison for all species.

Overall, prior to 2022, 1.0% of all animals tested for antibodies were seropositive, compared to

6.8% since 2022.

We compared the prevalence of SARS-CoV-2 PCR and seropositive cases from 2019-2024

against reported human COVID-19 cases reported each fall (September - December) in the

same public health units where the samples were collected (Ontario Agency for Health

Protection and Promotion (Public Health Ontario) 2024). No human COVID-19 data were

available for fall of 2024, as sample collection ended in June 2024. The proportion of

seropositive tests in this study increased from 0.3% in 2020 to 11.4% 2024, with a drop in 2023

(5.2%). Although seroprevalence was highest in 2024, only 35 serological tests were carried out

this year. Just by observation, the peak activity in human cases occurred in the fall in 2021,

coinciding with the highest yearly proportion of the samples that tested PCR positive.

Differences in seropositivity by sex and age were compared across the whole serological

dataset (2022-2024; N =2,839) among deer tested for antibodies using Fisher's exact test. Sex

and age were not significantly associated with seropositivity (p>0.05).

Regression modeling

Model selection

The most parsimonious model (lowest AIC through stepwise selection) explaining seropositivity of deer included three variables retained in the model, which were all significant with presence of SARS-CoV-2 on the landscape (distance to PCR positive deer, fall deer harvest, and human population) (Table 5). Distance to the previous 17 PCR positive deer was the most significant variable ($p=0.0008$), with every 10 km distance away from a previous positive PCR case, or the equivalent of one grid 10 x 10 km cell, the odds of an animal being seropositive decreased by 7.6% (Odds Ratio (OR): 0.92; 95% CI: 0.88 – 0.97).

Table 5. GLMM model coefficients of final model

	Estimate	Std. Error	Z value	P Value
Intercept	-3.6508	0.7240	-5.0422	<0.001
Distance to 17 PCR+ deer (2021)	-0.0790	0.0235	-3.3598	<0.001
Annual fall deer harvest (per 1000 animals)	0.4890	0.2034	2.4045	0.0162
Human population (per 1000 people)	1.1641	0.4172	2.7906	0.0053

The annual fall deer harvest was positively correlated with seropositivity, where deer sampled in grid cells with higher average hunting numbers were more likely to be seropositive. For every 1000 deer harvested in a WMU, the odds of one animal being seropositive increased by 63% (OR: 1.63; 95% CI: 1.1 – 2.5). Confidence intervals were calculated using the R *Stats* package (R Core Team, 2023). Population was found to be significant in the multivariate model, but not in univariate analysis. Population was not highly correlated with other variables, so this result was not due to multicollinearity. The results indicate that for every 1000 people, the odds a deer tested seropositive increased three times (OR: 3.2; 95% CI: 1.4 – 7.8).

Model diagnostics

Diagnostics were carried out of the mixed model using R package *PerformanceAnalytics* and *DHARMa*, which was created for assessing GLMMs by simulating standardized residuals (Hartig 2024). Dharma plots of residual vs predicted values showed no significant problems and no evidence of significant outliers, overdispersion or underdispersion. In GLMMs, the models assume a linear relationship between continuous predictor variables and the logit of the outcome. We ran scatterplots and found a linear trend with deer positives and human population, but some violation for deer harvest, which is not surprising with ecological models.

Model performance and accuracy

Model performance of the final model was assessed by making predictions with the test data and determining the model's accuracy with R package *lme4* (Bates et al. 2015). The *Predict* function uses population level fixed effects, where the corresponding random effect components are set to zero.

To test the ability of this model to correctly classify samples as positive or negative, we identified an optimal threshold of 0.1, which maximizes sensitivity and specificity. This threshold was used to create an ROC curve using R package *pROC* (Robin et al. 2011), where the area under the curve (AUC) represents the probability that a model will correctly distinguish between true positives and true Negatives, where a perfect score of 1 indicates perfect separability. The model had an AUC value of 0.71, which is considered acceptable discrimination (Mandrekar 2010). The model has low sensitivity (60%), but moderate specificity (88%), indicating lower ability to detect true positives (i.e., a higher false negative rate).

Discussion

We initiated an active surveillance program following the discovery in 2021 of a divergent

lineage of SARS-CoV-2 in deer. Analyses of B.1.641 suggested that the lineage may have

evolved in deer for up to one year, accumulating 76 mutations compared to ancestral SARS-

CoV-2 and appearing to have originated in Michigan mink farms (Pickering et al. 2022). Despite

sampling 1,446 Ontario deer during our active surveillance program since 2022, we have not

detected additional B.1.641 infections, or infections with related B.1 viruses. The only SARS-

CoV-2 infection we have observed in deer from Ontario after 2021 is a single XBB infection in

central Ontario. Consequently, our data do not support the conclusion that deer in Ontario are a

reservoir of B.1.641 in particular, or SARS-CoV-2 in general.

We define a reservoir as having an ability to maintain infection with a zoonotic pathogen, and

the presence of an interface for transmission to humans (Haydon et al. 2002). While it is clear

that there is ample opportunity for transmission of SARS-CoV-2 from deer to humans, such as

through hunting and backyard feeding of deer (Pratt et al. 2025), the lack of any viruses related

to B.1.641 in our study suggests that the virus has not been maintained in the deer population

following the relatively localised year-long outbreak previously observed (Pickering et al. 2022).

Instead, subsequent infections (e.g., the XBB infection we observed) are likely a result of

additional spillovers from humans. A similar pattern of multiple spillover events has been

observed (Kuchipudi et al. 2022; Caserta et al. 2023), including detection of Omicron variants at

a time where it accounted for 90% of human cases (Vandegrift et al. 2022), although there is

also evidence of sustained transmission within deer (Caserta et al. 2023; Feng et al. 2023).

However, we consider that evidence is accumulating which indicates that deer populations do

not sustain SARS-CoV-2 infections indefinitely.

Across Ontario, we estimated that SARS-CoV-2 seropositivity among deer was about 9.0%, but

in Southwestern Ontario, the estimated seropositivity was 15.2% (Figure 1 and 2). It is clear that

there was some transmission of B.1.641 among deer, both because the lineage appeared to

evolve for about a year prior to its detection in 2021 and because the best predictor in our

regression model for a seropositive deer was the distance to a previous B.1.641 case. This

variable in combination with variables describing human population density and deer harvest

density allowed us to estimate that for every 10 km distance away from a previous B.1.641

case, the odds of a deer being seropositive decreased by 7.6%. The regression model paints a

picture of B.1.641 spreading among deer for some period of time in areas with high deer and

human population densities. The infections eventually ceased to spread however, and this lack

of sustained transmission may be due to patterns in contact between animals or the adaptive

immune response of deer. The decrease in SARS-CoV-2 in deer and the parallel increase in

seropositivity raises the question of whether previous infection provides protective immunity,

which may explain why prevalence of infection decreased from 2021 to 2024. Although titres will

drop with time, neutralizing antibodies have been found to persist in captive white-tailed deer for

at least 13 months (Hamer et al. 2022).There is much that we do not yet know about how the

deer immune system responds to SARS-CoV-2 infections (Kotwa et al. 2023) or how

anthropogenic and natural landscape barriers that influence deer dispersal consequently affect

disease transmission (Peterson et al. 2017).

In addition to the seropositive deer detected in proximity to known B.1.641 infections, we also

detected a small number of other seropositive species (two opossums, a mink, and an otter).

With the exception of one opossum, none of the positive species occurred in close proximity to

known B.1.641 cases, and as such, we have no evidence to suggest B.1.641 was spilling from

deer into other wildlife species. In general, the seroprevalence was low in non-deer species,

which may reflect independent spillover events to these wildlife species without onward and

sustained intra-species transmission. Low estimates of seropositivity might also be due to a lack

of seroconversion, or waning antibodies. For example, an opossum resampled after a month

had a low percent of neutralizing antibodies (51.1%; Goldberg et al., 2024). While the Surrogate

Virus Neutralization Test (sVNT) used to detect neutralization antibodies has been highly

sensitive (95-100%) to Covid-19 in humans (Tan et al. 2020), this may vary based on species,

method of testing and field conditions, and even mutations since this test was first established

(Hewitt et al. 2025). For example, there is some evidence that Nobuto strips may underestimate

seroprevalence for pre-Omicron variants (Hewitt et al. 2025). Another factor that cannot be

accounted for is the potential cross-reactions with other endemic coronaviruses that could be

affecting the specificity of the assay. Little is known about endemic coronaviruses circulating in

these species. As for low prevalence of virus detected through PCR testing, it is possible that

active infection was missed due to the means of swabbing. In experimentally inoculated skunks,

viral titres were greater in nasal samples compared to oral samples (Bosco-Lauth et al. 2021),

the latter of which were primarily used in our study, although this exposure to high titred

innoculum does not reflect natural infection pressure.

Human SARS-CoV-2 cases peaked in Ontario during 2021, and not surprisingly this was also

the peak of SARS-CoV-2 infections in Ontario's wildlife. Superficially, this appears to support

the idea that high infection burden in the human population was associated with spillover to

wildlife (e.g., Vandegrift et al., 2022). All of the documented positive cases of SARS-CoV-2 in

wild animals during 2021 were deer with B.1.641. However, as of April 2021, Delta became the

dominating circulating variant in the human population (Ontario Agency for Health Protection

and Promotion (Public Health Ontario) 2022), and we have not documented any Delta infections

among wildlife in Ontario. The B.1.641 infections appear to have been evolving in deer for up to

a year (Pickering et al. 2022). We did observe a peak in seropositivity among wildlife samples in

2022, a year after the peak in B.1.641 cases, mostly, but not exclusively arising from deer

samples. We cannot disentangle whether the seropositive cases represent spillovers from the

human population, or sylvatic transmission among wildlife species.

As has been observed in other North American jurisdictions since Omicron became the

dominant SARS-CoV-2 clade (Feng et al. 2023; McBride et al. 2023), we found recent evidence

of a deer with an Omicron infection (Figure 3). This does not appear associated with the

previous B.1.641 infections, but is instead likely a recent spillover from the human population.

The infected deer was sampled in central Ontario in a region distant from any other documented

cases of deer with SARS-CoV-2 and with an abundance of cottages, providing opportunities for

human-to-deer transmission. The lack of additional cases in deer is likely associated with the

reduced prevalence of infection in the human population after 2021. It is also possible, however,

that deer are less susceptible to infection with the Omicron lineages of SARS-CoV-2 than with

older lineages. One possible explanation for such an effect is that APOBEC family of proteins,

which lead to C>U mutations through RNA editing, are prominent features of deer immune

response to infections with pre-Omicron lineages (Pickering et al. 2022; Kotwa et al. 2023).

APOBECs have also been shown however, to promote viral replication and propagation (Kim et

al. 2022). Therefore, if APOBEC-mediated RNA editing is less involved in deer immune

response to Omicron, this might reduce prevalence of infection among deer. There remains

considerable uncertainty about the differential immune response of deer to lineages of SARS-

CoV-2.

Filling the knowledge gaps in viral genomic surveillance and the interface between human and

animal disease can help us better prepare for emerging diseases and can improve our

understanding of host jumping (Mubareka et al. 2023; C.C.S. Tan et al. 2024). When disease

surveillance systems exist across animal and human health sectors however, they are often

carried out by separate agencies or industries with different regulations and reporting

requirements, making communication and data sharing difficult. To address these barriers, we

applied a One Health approach to our surveillance program, with the aim to reduce the existing

gaps separating huma and animal disease knowledge. We established a multidisciplinary team,

with cross-sectoral expertise in animal, human, and environmental health (Lee et al. 2024). This

follows Kuchipudi et al.'s call for a robust One Health-centered SARS-CoV-2 surveillance of

cross-species transmission in non-human species (Kuchipudi et al. 2023).

We recognize that our study has potential limitations. For example, on live wild animals, we

principally used oral swabs for virus detection, and this may not have been the optimal virus

target in some species. Our serological assay was not validated for use across all species in the

study. We also need to establish a better understanding of the immune response of deer which

might have led to the eventual decline of B.1.641, and perhaps to less prevalent Omicron

infections. With limited data on serological and viral infection, sex and age-mediated differences

could be missed, especially when 7% of the animals were of unknown sex, which may have

limited the influence on transmission and behavior.

Wildlife reservoirs of SARS-CoV-2 can alter virus ecology, leading to divergent lineages with

potential health consequences for the human population. Effects of the B.1.641 lineage on

humans are not yet well understood, and consequently there has been considerable interest in

assessing the possibility that deer were becoming a reservoir of this virus. We established a

multi-disciplinary team to undertake a One Health-centred surveillance program to evaluate the

risks of reservoir establishment. To date, evidence suggests that B.1641 has not continued to

circulate among deer, and therefore, we have no evidence that deer are a reservoir of SARS-

CoV-2 in Ontario.

Acknowledgements

We want to thank those who made sampling wildlife possible, including students and volunteers

assisting on the SARS-CoV-2 project (Laurelie, Brennan, Alexis, Sean, Thomas, Greg, Sarah,

Maissie, Shelby, Sarah, Rebekah, Janet), MNR researchers who took on additional sampling

work to increase our study area, as well as wildlife rehabilitators (Another Chance, Aspen Valley

Wildlife Sanctuary, Erie Wildlife Rescue, Fur-Ever Wild Rehabilitation, Hobbitstee Wildlife

Refuge, Itty Bitty Critters Wildlife Rehabilitation, Procyon Wildlife, Salthaven Wildlife

Rehabilitation & Education Centre, Shades of Hope Wildlife Refuge, Woodlands Wildlife

Sanctuary, Wings Rehabilitation Centre, with special thanks to Carol, Colleen, Kelly, and Dave),

zoos (Greenview Park & Zoo, Marineland, Waterford Deer Park, West Perth Animal Park,

Riverview Park & Zoo), Conservation Authorities (Ausable Bayfield CA, Catfish Creek CA, Kettle

Creek CA, Lower Thames Valley CA, Upper Thames River CA, St Clair Region CA), provincial

and national parks (Algonquin, Komoka, Pinery, Point Pelee, Short Hills, Rondeau), the CWHC

(B. Stevens and L. Shirose), and municipalities (City of London, City of St. Thomas, Town of St.

Mary's, and Oxford County). Those who contributed harvested deer samples, we thank

members of the Haudenosaunee Confederacy and Haudenosaunee Wildlife and Habitat

Authority, Kettle and Stony Point First Nation, licenced Ontario deer hunters, and Appin BBQ

and Catering. We also thank Norfolk Field Naturalists and the Niagara SPCA and Humane

Society.

We recognize that our study area in present-day Ontario is located on lands of traditional

multiple Indigenous nations, and that accessing wildlife on these lands has been possible in part

by the preservation of greenspaces across Ontario and through stewardships with these

Indigenous communities.

Author contributions

Conceptualization: L.C., J.D.K., O.L., F.M., B.,P., C.J., S.M., J.B.
Data Curation: L.C., J.D.K., S.P.J., C.L., E.C., W.Y., P.K., O.V., O.L., A.I.S., C.J., H.C., S.M., J.B.
Formal Analysis: L.C., J.D.K., S.P.J., C.L., A.D., P.K., O.V., H.C., S.M., J.B.
Funding Acquisition: O.L., A.I.S., B.,P., C.J., S.M., J.B.
Investigation: L.C., J.D.K., S.P.J., C.L., A.D., A.S., S.L.N., E.C., W.Y., H.C., J.B.
Methodology: L.C., J.D.K., C.L., A.D., P.K., O.V., O.L., S.M., J.B.
Project Administration: J.D.K., C.J., H.C., S.M., J.B.
Resources: J.D.K., O.L., C.J., H.C., S.M., J.B.
Software: L.C., J.D.K., P.K., O.V.
Supervision: L.C., J.D.K., O.L., C.J., H.C., S.M., J.B.
Validation: J.D.K., C.L., A.D.
Visualization: L.C., P.K., O.V.
Writing – Original Draft Preparation: L.C., J.D.K., S.P.J., W.Y., O.V., O.L., H.C., J.B.,
Writing – Review & Editing: L.C., J.D.K., S.P.J., C.L., A.D., A.S., S.L.N., E.C., W.Y., P.K., O.V.,
O.L., A.I.S., F.M., B.,P., C.J., H.C., S.M., J.B.

Declaration of interests

842 The authors declare no competing interests.

References

Abdel-Moneim AS, and Abdelwhab EM. 2020. Evidence for SARS-CoV-2 infection of animal

hosts. Pathogens, **9**(7): 529. https://doi.org/10.3390/pathogens9070529.

Agusi ER, Schön J, Allendorf V, Eze EA, Asala O, Shittu I, et al. 2024. SARS-CoV and SARS-

CoV -2 cross-reactive antibodies in domestic animals and wildlife in Nigeria suggest circulation

of sarbecoviruses. One Health, **18**: 100709. https://doi.org/10.1016/j.onehlt.2024.100709.

Aksamentov I, Roemer C, Hodcroft EB, and Neher RA. 2021. Nextclade: clade assignment,

mutation calling and quality control for viral genomes. Journal of Open Source Software, **6**(67):

3773. https://doi.org/10.21105/joss.03773.

Andrews S. 2010. FastQC: A quality control tool for high throughput sequence data.

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/. Available from

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Bates D, Mächler M, Bolker B, and Walker S. 2015. Fitting linear mixed-effects models using

lme4. Journal of Statistical Software, **67**: 1–48. https://doi.org/10.18637/jss.v067.i01.

Birds Canada. 2024. Ontario Breeding Bird Atlas: Atlas 10x10km grid shapefile [dataset]. Port

Rowan, ON. https://www.birdsontario.org.

Bosco-Lauth AM, Root JJ, Porter SM, Walker AE, Guilbert L, Hawvermale D, et al. 2021.

Peridomestic mammal susceptibility to severe acute respiratory syndrome coronavirus 2

infection. Emerging Infectious Diseases, **27**(8): 2073–2080.

https://doi.org/10.3201/eid2708.210180.

Caserta LC, Martins M, Butt SL, Hollingshead NA, Covaleda LM, Ahmed S, et al. 2023. White-

tailed deer (*Odocoileus virginianus*) may serve as a wildlife reservoir for nearly extinct SARS-

CoV-2 variants of concern. Proceedings of the National Academy of Sciences, **120**(6):

e2215067120. https://doi.org/10.1073/pnas.2215067120.

Chandler JC, Bevins SN, Ellis JW, Linder TJ, Tell RM, Jenkins-Moore M, et al. 2021. SARS-

CoV-2 exposure in wild white-tailed deer (*Odocoileus virginianus*). Proceedings of the National

Academy of Sciences, **118**(47): e2114828118. https://doi.org/10.1073/pnas.2114828118.

Chen S, Zhou Y, Chen Y, and Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics, **34**(17): i884–i890. https://doi.org/10.1093/bioinformatics/bty560.

Cingolani P, Patel VM, Coon M, Nguyen T, Land SJ, Ruden DM, et al. 2012. Using *Drosophila*

melanogaster as a model for genotoxic chemical mutational studies with a new program,

SnpSift. Frontiers in Genetics, **3**(35). https://doi.org/10.3389/fgene.2012.00035.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. 2012. A program for

annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. Fly, **6**(2): 80–

92. https://doi.org/10.4161/fly.19695.

Colquhoun R, Jackson B, O'Toole Á, and Rambaut A. 2023. SCORPIO: A utility for defining and

classifying mutation constellations of virus genomes. Bioinformatics, **39**(10): btad575.

https://doi.org/10.1093/bioinformatics/btad575.

Cool K, Gaudreault NN, Morozov I, Trujillo JD, Meekins DA, McDowell C, et al. 2022. Infection

and transmission of ancestral SARS-CoV-2 and its alpha variant in pregnant white-tailed deer.

Emerging Microbes & Infections, **11**(1): 95–112.

https://doi.org/10.1080/22221751.2021.2012528.

Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. 2020. Detection of

2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. Eurosurveillance, **25**(3): 2000045.

https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045.

Damas J, Hughes GM, Keough KC, Painter CA, Persky NS, Corbo M, et al. 2020. Broad host

range of SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in vertebrates.

Proceedings of the National Academy of Sciences, **117**(36): 22311–22322.

https://doi.org/10.1073/pnas.2010146117.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. 2021. Twelve years of

SAMtools and BCFtools. GigaScience, **10**(2). https://doi.org/10.1093/gigascience/giab008.

Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, and Notredame C. 2017.

Nextflow enables reproducible computational workflows. Nature Biotechnology, **35**(4): 316–319.

https://doi.org/10.1038/nbt.3820.

Dudas G, Carvalho LM, Rambaut A, and Bedford T. 2018. MERS-CoV spillover at the camel-

human interface. eLife, **7**: e31257. https://doi.org/10.7554/eLife.31257.

Elbe S, and Buckland-Merrett G. 2017. Data, disease and diplomacy: GISAID's innovative

contribution to global health. Global Challenges, **1**(1): 33–46. https://doi.org/10.1002/gch2.1018.

Esri. 2022. ArcGIS Pro (Version 3.0.3). Redlands, CA. https://www.esri.com/en-

us/arcgis/products/arcgis-pro/overview.

Fagre A, Lewis J, Eckley M, Zhan S, Rocha SM, Sexton NR, et al. 2021. SARS-CoV-2 infection,

neuropathogenesis and transmission among deer mice: Implications for spillback to New World

rodents. PLoS pathogens, **17**(5): e1009585. https://doi.org/10.1371/journal.ppat.1009585.

Feng A, Bevins S, Chandler J, DeLiberto TJ, Ghai R, Lantz K, et al. 2023. Transmission of

SARS-CoV-2 in free-ranging white-tailed deer in the United States. Nature Communications,

14(4078). https://doi.org/10.1038/s41467-023-39782-x.

Fox J, and Weisberg S. 2019. An R Companion to Applied Regression (3rd ed.). Sage,

Thousand Oaks, CA. https://www.john-fox.ca/Companion.

Fryxell JM, Falls JB, Falls EA, and Brooks RJ. 1998. Long-term dynamics of small-mammal

populations in Ontario. Ecology, **79**(1): 213–225. https://doi.org/10.2307/176876.

Gangavarapu K, Latif AA, Mullen JL, Alkuzweny M, Hufbauer E, Tsueng G, et al. 2023.

Outbreak.info genomic reports: Scalable and dynamic surveillance of SARS-CoV-2 variants and

mutations. Nature Methods, **20**(4): 512–522. https://doi.org/10.1038/s41592-023-01769-3.

Goldberg AR, Langwig KE, Brown KL, Marano JM, Rai P, King KM, et al. 2024. Widespread

exposure to SARS-CoV-2 in wildlife communities. Nature Communications, **15**(6210).

https://doi.org/10.1038/s41467-024-49891-w.

Gortázar C, Barroso-Arévalo S, Ferreras-Colino E, Isla J, de la Fuente G, Rivera B, et al. 2021.

Natural SARS-CoV-2 infection in kept ferrets, Spain. Emerging Infectious Diseases, **27**(7):

1994–1996. https://doi.org/10.3201/eid2707.210096.

Greenhorn JE, Kotwa JD, Bowman J, Bruce L, Buchanan T, Buck PA, et al. 2022. SARS-CoV-2

wildlife surveillance in Ontario and Québec. Canada Communicable Disease Report, **48**(6):

243–251. https://doi.org/10.14745/ccdr.v48i06a02.

Griffin BD, Chan M, Tailor N, Mendoza EJ, Leung A, Warner BM, et al. 2021. SARS-CoV-2

infection and transmission in the North American deer mouse. Nature Communications,

12(3612). https://doi.org/10.1038/s41467-021-23848-9.

Grubaugh ND, Gangavarapu K, Quick J, Matteson NL, De Jesus JG, Main BJ, et al. 2019. An

amplicon-based sequencing framework for accurately measuring intrahost virus diversity using

PrimalSeq and iVar. Genome Biology, **20**(8). https://doi.org/10.1186/s13059-018-1618-7.

Hale VL, Dennis PM, McBride DS, Nolting JM, Madden C, Huey D, et al. 2022. SARS-CoV-2

infection in free-ranging white-tailed deer. Nature, **602**: 481–486.

https://doi.org/10.1038/s41586-021-04353-x.

Hamer SA, Nunez C, Roundy CM, Tang W, Thomas L, Richison J, et al. 2022. Persistence of

SARS-CoV-2 neutralizing antibodies longer than 13 months in naturally infected, captive white-

tailed deer (*Odocoileus virginianus*), Texas. Emerging Microbes & Infections, **11**(1): 2112–2115.

https://doi.org/10.1080/22221751.2022.2112913.

Hartig F. 2024. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models [online]. https://CRAN.R-project.org/package=DHARMa.

Haydon DT, Cleaveland S, Taylor LH, and Laurenson MK. 2002. Identifying reservoirs of

infection: A conceptual and practical challenge. Emerging Infectious Diseases, **8**(12): 1468–

1473. https://doi.org/10.3201/eid0812.010317.

Hewitt J, Wilson-Henjum G, Chandler JC, Phillips AT, Diel DG, Walter WD, et al. 2025.

Evaluation of SARS-CoV-2 antibody detection methods for wild Cervidae. Preventive Veterinary

Medicine, **241**: 106522. https://doi.org/10.1016/j.prevetmed.2025.106522.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, and Vinh LS. 2017. UFBoot2: Improving

the ultrafast bootstrap approximation. Molecular Biology and Evolution, **35**(2): 518–522.

https://doi.org/10.1093/molbev/msx281.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, and Jermiin LS. 2017. ModelFinder:

fast model selection for accurate phylogenetic estimates. Nature Methods, **14**(6): 587–589.

https://doi.org/10.1038/nmeth.4285.

Kim K, Calabrese P, Wang S, Qin C, Rao Y, Feng P, et al. 2022. The roles of APOBEC-

mediated RNA editing in SARS-CoV-2 mutations, replication and fitness. Scientific Reports,

12(14972). https://doi.org/10.1038/s41598-022-19067-x.

Kolawole OM, Tomori O, Agbonlahor D, Ekanem E, Bakare R, Abdulsalam N, et al. 2022. SARS

CoV-2 seroprevalence in selected states of high and low disease burden in Nigeria. JAMA

network open, **5**(10): e2236053. https://doi.org/10.1001/jamanetworkopen.2022.36053.

Kotwa JD, Bhuinya A, Yim W, Gallo G, Singh P, Liu Q, et al. 2025. High host specificity of

alphacoronaviruses in Nearctic, insectivorous bats. npj Viruses, **3**(38).

https://doi.org/10.1038/s44298-025-00115-y.

Kotwa JD, Lobb B, Massé A, Gagnier M, Aftanas P, Banerjee A, et al. 2023. Genomic and

transcriptomic characterization of delta SARS-CoV-2 infection in free-ranging white-tailed deer

(*Odocoileus virginianus*). iScience, **26**(11): 108319. https://doi.org/10.1016/j.isci.2023.108319.

Kuchipudi SV, Surendran-Nair M, Ruden RM, Yon M, Nissly RH, Vandegrift KJ, et al. 2022.

Multiple spillovers from humans and onward transmission of SARS-CoV-2 in white-tailed deer.

Proceedings of the National Academy of Sciences, **119**(6): e2121644119.

https://doi.org/10.1073/pnas.2121644119.

Kuchipudi SV, Tan C, van Dorp L, Lichtveld M, Pickering B, Bowman J, et al. 2023. Coordinated

surveillance is essential to monitor and mitigate the evolutionary impacts of SARS-CoV-2

spillover and circulation in animal hosts. Nature Ecology & Evolution, **7**: 956–959.

https://doi.org/10.1038/s41559-023-02082-0.

Langmead B, and Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nature

Methods, **9**(4): 357–359. https://doi.org/10.1038/nmeth.1923.

LeBlanc JJ, Gubbay JB, Li Y, Needle R, Arneson SR, Marcino D, et al. 2020. Real-time PCR-

based SARS-CoV-2 detection in Canadian laboratories. Journal of Clinical Virology, **128**:

104433. https://doi.org/10.1016/j.jcv.2020.104433.

Lee Y, Jeeves S, Crawshaw L, Kotwa J, Pickering B, Mubareka S, et al. 2024. The Wildlife

Emerging Pathogens Initiative: Wild EPI and One Health. iScience, **27**(7): 110317.

https://doi.org/10.1016/j.isci.2024.110317.

Lesage L, Crête M, Huot J, Dumont A, and Ouellet J-P. 2000. Seasonal home range size and

philopatry in two northern white-tailed deer populations. Canadian Journal of Zoology, **78**(11):

1930–1940. https://doi.org/10.1139/z00-117.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. 2009. The Sequence

Alignment/Map format and SAMtools. Bioinformatics, **25**(16): 2078–2079.

https://doi.org/10.1093/bioinformatics/btp352.

Mackay IM, and Arden KE. 2015. Middle East respiratory syndrome: An emerging coronavirus

infection tracked by the crowd. Virus Research, **202**: 60–88.

https://doi.org/10.1016/j.virusres.2015.01.021.

Mandrekar JN. 2010. Receiver operating characteristic curve in diagnostic test assessment.

Journal of Thoracic Oncology, **5**(9): 1315–1316.

https://doi.org/10.1097/JTO.0b013e3181ec173d.

Marques AD, Sherrill-Mix S, Everett JK, Adhikari H, Reddy S, Ellis JC, et al. 2022. Multiple

introductions of SARS-CoV-2 Alpha and Delta variants into white-tailed deer in Pennsylvania.

mBio, **13**(5): e0210122. https://doi.org/10.1128/mbio.02101-22.

Martins M, Boggiatto PM, Buckley A, Cassmann ED, Falkenberg S, Caserta LC, et al. 2022.

From Deer-to-Deer: SARS-CoV-2 is efficiently transmitted and presents broad tissue tropism

and replication sites in white-tailed deer. PLOS Pathogens, **18**(3): e1010197.

https://doi.org/10.1371/journal.ppat.1010197.

McBride DS, Garushyants SK, Franks J, Magee AF, Overend SH, Huey D, et al. 2023.

Accelerated evolution of SARS-CoV-2 in free-ranging white-tailed deer. Nature

Communications, **14**(5105). https://doi.org/10.1038/s41467-023-40706-y.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. 2020.

IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era.

Molecular Biology and Evolution, **37**(5): 1530–1534. https://doi.org/10.1093/molbev/msaa015.

Mubareka S, Amuasi J, Banerjee A, Carabin H, Copper Jack J, Jardine C, et al. 2023.

Strengthening a One Health approach to emerging zoonoses. FACETS, **8**: 1–64.

https://doi.org/10.1139/facets-2021-0190.

Ontario Agency for Health Protection and Promotion (Public Health Ontario). 2022. Confirmed

SARS-CoV-2 variant of concern cases from November 29, 2020 to November 12, 2021.

Queen's Printer for Ontario, Toronto, ON.

Ontario Agency for Health Protection and Promotion (Public Health Ontario). 2024. Ontario

Respiratory Virus Tool [dataset]. King's Printer for Ontario, Toronto, ON.

https://www.publichealthontario.ca/en/Data-and-Analysis/Infectious-Disease/Respiratory-Virus-

Tool.

Ontario Ministry of Natural Resources. 2014. Ontario Land Cover Compilation Data (Version

2.0) [dataset]. Queen's Printer for Ontario, Peterborough, ON.

https://geohub.lio.gov.on.ca/documents/7aa998fdf100434da27a41f1c637382c.

Ontario Ministry of Natural Resources. 2019. Chronic Wasting Disease prevention and response

plan. King's Printer for Ontario, Peterborough, ON.

Ontario Ministry of Natural Resources. 2020. Ontario Hydro Network: Waterbody (Version 1.2)

[dataset]. Queens Printer for Ontario, Peterborough, ON.

https://geohub.lio.gov.on.ca/datasets/mnrf::ontario-hydro-network-ohn-waterbody.

Ontario Ministry of Natural Resources. 2024. White-tailed deer hunting activity and harvest -

Ontario Data Catalogue [dataset]. King's Printer for Ontario, Peterborough, ON.

https://data.ontario.ca/dataset/white-tailed-deer-hunting-activity-and-harvest.

Oreshkova N, Molenaar RJ, Vreman S, Harders F, Oude Munnink BB, Hakze-van der Honing

RW, et al. 2020. SARS-CoV-2 infection in farmed minks, the Netherlands, April and May 2020.

Eurosurveillance, **25**(23): pii=200100. https://doi.org/10.2807/1560-

7917.ES.2020.25.23.2001005.

O'Toole Á, Scher E, Underwood A, Jackson B, Hill V, McCrone JT, et al. 2021. Assignment of

epidemiological lineages in an emerging pandemic using the pangolin tool. Virus Evolution, **7**(2):

veab064. https://doi.org/10.1093/ve/veab064.

Oude Munnink BB, Sikkema RS, Nieuwenhuijse DF, Molenaar RJ, Munger E, Molenkamp R, et

al. 2021. Transmission of SARS-CoV-2 on mink farms between humans and mink and back to

humans. Science, **371**: 172–177. https://doi.org/10.1126/science.abe5901.

Palmer MV, Martins M, Falkenberg S, Buckley A, Caserta LC, Mitchell PK, et al. 2021.

Susceptibility of white-tailed deer (*Odocoileus virginianus*) to SARS-CoV-2. Journal of Virology,

95(11): e00083-21. https://doi.org/10.1128/JVI.00083-21.

Patel H, Monzón S, Varona S, Espinosa-Carrasco J, Garcia MU, Nf-Core Bot, et al. 2023. nf-

core/viralrecon: nf-core/viralrecon (Version 2.6.0) - Rhodium Raccoon [online].

https://zenodo.org/records/7764938.

Pedersen BS, and Quinlan AR. 2018. Mosdepth: Quick coverage calculation for genomes and

exomes. Bioinformatics, **34**(5): 867–868. https://doi.org/10.1093/bioinformatics/btx699.

Peterson BE, Storm DJ, Norton AS, and Van Deelen TR. 2017. Landscape influence on

dispersal of yearling male white-tailed deer. The Journal of Wildlife Management, **81**(8): 1449–

1456. https://doi.org/10.1002/jwmg.21318.

Peterson BG, and Carl P. 2024. PerformanceAnalytics: Econometric tools for performance and

risk analysis. R package (Version 2.0.8-1) [online].

https://github.com/braverock/performanceanalytics.

Pickering B, Lung O, Maguire F, Kruczkiewicz P, Kotwa JD, Buchanan T, et al. 2022. Divergent

SARS-CoV-2 variant emerges in white-tailed deer with deer-to-human transmission. Nature

Microbiology, **7**: 2011–2024. https://doi.org/10.1038/s41564-022-01268-9.

Pither R, O'Brien P, Brennan A, Hirsh-Pearson K, and Bowman J. 2023. Predicting areas

important for ecological connectivity throughout Canada. PLoS One, **18**(2): e0281980.

https://doi.org/10.1371/journal.pone.0281980.

Pratt A, Prezioso T, Mateus-Pinilla N, Pepin KM, and Smith R. 2025. Interactions between

humans and white-tailed deer in Illinois: A cross-sectional survey. EcoHealth, **22**: 147–160.

https://doi.org/10.1007/s10393-024-01694-7.

Pustake M, Tambolkar I, Giri P, and Gandhi C. 2022. SARS, MERS and CoVID-19: An overview

and comparison of clinical, laboratory and radiological features. Journal of Family Medicine and

Primary Care, **11**(1): 10–17. https://doi.org/10.4103/jfmpc.jfmpc_839_21.

R Core Team. 2024. R: A Language and Environment for Statistical Computing. Vienna,

Austria. https://www.R-project.org.

Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, et al. 2020. A dynamic

nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. Nature

Microbiology, **5**: 1403–1407. https://doi.org/10.1038/s41564-020-0770-5.

Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J-C, et al. 2011. pROC: An open-

source package for R and S+ to analyze and compare ROC curves. BMC bioinformatics, **12**: 77.

https://doi.org/10.1186/1471-2105-12-77.

Rudar J, Kruczkiewicz P, Vernygora O, Golding GB, Hajibabaei M, and Lung O. 2024.

Sequence signatures within the genome of SARS-CoV-2 can be used to predict host source.

Microbiology Spectrum, **12**(4): e03584-23. https://doi.org/10.1128/spectrum.03584-23.

Shan C, Yao Y-F, Yang X-L, Zhou Y-W, Gao G, Peng Y, et al. 2020. Infection with novel

coronavirus (SARS-CoV-2) causes pneumonia in Rhesus macaques. Cell Research, **30**: 670–

677. https://doi.org/10.1038/s41422-020-0364-z.

Singh BB, Ward MP, and Dhand NK. 2023. Host characteristics and their influence on zoonosis,

disease emergence and multi-host pathogenicity. One Health, **17**: 100596.

https://doi.org/10.1016/j.onehlt.2023.100596.

Statistics Canada. 2023a. Census Profile. 2021 Census of Population [dataset]. Statistics

Canada Catalogue no. 98-316-X2021001. Ottawa, ON. https://www12.statcan.gc.ca/census-

recensement/2021/dp-pd/prof/index.cfm?Lang=E.

Statistics Canada. 2023b. Dissemination Area Boundary File, 2021 Census [dataset]. Statistics

Canada Catalogue no. 92-196-X. Ottawa, ON.

https://www150.statcan.gc.ca/n1/en/catalogue/92-169-X.

Statistics Canada. 2023c. Provincial Boundary Files, 2021 Census [dataset]. Statistics Canada

Catalogue no. 92-160-X, Ottawa, ON. https://www12.statcan.gc.ca/census-

recensement/2021/geo/sip-pis/boundary-limites/index2021-eng.cfm?year=21.

Tan CCS, Van Dorp L, and Balloux F. 2024. The evolutionary drivers and correlates of viral host

jumps. Nature Ecology & Evolution, **8**: 960–971. https://doi.org/10.1038/s41559-024-02353-4.

Tan CW, Chia WN, Qin X, Liu P, Chen MI-C, Tiu C, et al. 2020. A SARS-CoV-2 surrogate virus

neutralization test based on antibody-mediated blockage of ACE2-spike protein-protein

interaction. Nature Biotechnology, **38**: 1073–1078. https://doi.org/10.1038/s41587-020-0631-z.

Tan ZH, Yong KY, and Shu J-J. 2024. Predicting potential SARS-CoV-2 spillover and spillback

in animals. Journal of Microbiology, Immunology, and Infection = Wei Mian Yu Gan Ran Za Zhi,

57(2): 225–237. https://doi.org/10.1016/j.jmii.2024.01.002.

Turakhia Y, Thornlow B, Hinrichs AS, De Maio N, Gozashti L, Lanfear R, et al. 2021. Ultrafast

Sample placement on Existing tRees (UShER) enables real-time phylogenetics for the SARS-

CoV-2 pandemic. Nature Genetics, **53**: 809–816. https://doi.org/10.1038/s41588-021-00862-7.

U.S. Census Bureau. 2018. Cartographic Boundary Files - 2018 TIGER/Line State Shapefile

[dataset]. U.S. Department of Commerce. Washington, D.C.

https://www.census.gov/geographies/mapping-files/time-series/geo/carto-boundary-file.html.

Vandegrift KJ, Yon M, Surendran Nair M, Gontu A, Ramasamy S, Amirthalingam S, et al. 2022.

SARS-CoV-2 Omicron (B.1.1.529) infection of wild white-tailed deer in New York City. Viruses,

14(12): 2770. https://doi.org/10.3390/v14122770.

Wang Q, Ye S-B, Zhou Z-J, Li J-Y, Lv J-Z, Hu B, et al. 2023. Key mutations on spike protein

altering ACE2 receptor utilization and potentially expanding host range of emerging SARS-CoV-

2 variants. Journal of Medical Virology, **95**(1): e28116. https://doi.org/10.1002/jmv.28116.

Wei C, Shan K-J, Wang W, Zhang S, Huan Q, and Qian W. 2021. Evidence for a mouse origin

of the SARS-CoV-2 Omicron variant. Journal of Genetics and Genomics, **48**(12): 1111–1121.

https://doi.org/10.1016/j.jgg.2021.12.003.

Wernike K, Böttcher J, Amelung S, Albrecht K, Gärtner T, Donat K, et al. 2022. Antibodies

against SARS-CoV-2 suggestive of single events of spillover to cattle, Germany. Emerging

Infectious Diseases, **28**(9): 1916–1918. https://doi.org/10.3201/eid2809.220125.

Wickham H. 2016. Elegant Graphics for Data Analysis. Springer-Verlag. Houston, TX.

https://ggplot2.tidyverse.org.

Wolters WJ, de Rooij MMT, Molenaar RJ, de Rond J, Vernooij JCM, Meijer PA, et al. 2022.

Manifestation of SARS-CoV-2 infections in mink related to host-, virus- and farm-associated

factors, the Netherlands 2020. Viruses, **14**(8): 1754. https://doi.org/10.3390/v14081754.

World Health Organization. 2024. WHO Coronavirus (COVID-19) dashboard: Number of

COVID-19 deaths reported to WHO (cumulative total) [dataset]. Datadot. Geneva, Switzerland.

https://data.who.int/dashboards/covid19/deaths.

Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, et al. 2020. A new coronavirus associated

with human respiratory disease in China. Nature, **579**: 265–269. https://doi.org/10.1038/s41586-

020-2008-3.

Wu L, Chen Q, Liu K, Wang J, Han P, Zhang Y, et al. 2020. Broad host range of SARS-CoV-2

and the molecular basis for SARS-CoV-2 binding to cat ACE2. Cell Discovery, **6**: 68.

https://doi.org/10.1038/s41421-020-00210-9.

Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. 2020. A pneumonia outbreak

associated with a new coronavirus of probable bat origin. Nature, **579**: 270–273.

https://doi.org/10.1038/s41586-020-2012-7.

1132