Intracellular formate is a critical mediator of the antimicrobial fate of *Salmonella* Typhimurium Against meropenem and ciprofloxacin

Debapriya Mukherjee[1], Sakshi Vasant Wagle[2], Pallab Ghosh[1#], Abhishek Biswas[1#], Dipshikha Chakravortty[1,3*]

[1]Department of Microbiology and Cell Biology, Division of Biological Sciences, Indian Institute of Science, Bangalore, India.

[2]Department of Biological Sciences, Indian Institute of Science Education and Research, Bhopal, India

[3]School of Biology, Indian Institute of Science Education and Research, Thiruvananthapuram, 695551.

#Equal contribution

*Corresponding author: dipa@iisc.ac.in

Tel: 0091 80 2293 2842

Fax: 0091 80 2360 2697

Abstract

The emergence of multidrug-resistant *Salmonella enterica* serovar Typhimurium (STM) poses a critical threat to global health. While classical antimicrobial resistance (AMR) mechanisms are well-characterized, alternative metabolites and metabolic pathways also contribute to resistance. This study highlights the role of intracellular formate, maintained by pyruvate-formate lyase (PflB), in modulating susceptibility of STM to meropenem and ciprofloxacin. Loss of *pflB* disrupts intracellular pH (ipH), impairing efflux pump activity, increasing reactive oxygen species (ROS), and causing membrane depolarization. These effects enhance antibiotic susceptibility but are reversed by extracellular formate supplementation. Mechanistically, disrupted ipH activates the RpoE–*csrB* axis, leading to downregulation of efflux pump genes *acrB* and *tolC*. Additionally, the BarA/SirA two-component system can also modulate the pathway. Our findings uncover a non-canonical role of formate in regulating efflux-mediated AMR, emphasizing its potential as a metabolic target in combating resistant STM strains.

Keywords: PflB, Antimicrobial response (AMR), Efflux pumps, CsrA/*csrB* axis

Introduction

The widespread misuse and overuse of antibiotics over recent decades have substantially contributed to the accelerated emergence of bacterial antimicrobial resistance (AMR) [1, 2]. Among resistant pathogens, multidrug-resistant (MDR) *Salmonella* has emerged as a significant global health concern due to its association with elevated morbidity and mortality rates. In 2019, AMR-related *Salmonella* infections were estimated to have caused approximately 1,50,000 deaths worldwide, with 23,700 cases directly attributable to MDR strains [3]. *Salmonella enterica* subspecies *enterica* serovar Typhimurium (STM) is one of the most prevalent serotypes implicated in MDR infections and plays a critical role in the declining efficacy of commonly used antibiotics, posing a substantial threat to public health systems globally [4-7].

Salmonella employs a range of mechanisms to evade antibiotic action, including enzymatic degradation of antibiotics, efflux pump-mediated drug expulsion, structural modification of antimicrobial agents, and alteration or protection of drug targets[8, 9]. While these resistance strategies have been well characterized in recent studies, the role of bacterial metabolism in modulating antimicrobial resistance (AMR) in *Salmonella* remains underexplored. In our previous work, we elucidated the role of the *pflB* gene—encoding pyruvate-formate lyase (PflB)—in STM pathogenesis. PflB contributes to the maintenance of the intracellular pool of formate, a short-chain fatty acid that plays a critical role in regulating intracellular pH (ipH). In the *pflB* deletion mutant (STM Δ*pflB*), dysregulation of ipH results in membrane perturbation, activation of the extra cytoplasmic sigma factor RpoE, increased expression of the small RNA *csrB*, and subsequent repression of flagellar gene expression[10]. Further, in a serendipitous finding in our laboratory, we observed that deletion of *pflB* also sensitizes STM to the antibiotics meropenem and ciprofloxacin. This unexpected finding prompted a detailed investigation into the role of intracellular formate homeostasis in AMR regulation.

In this study, we investigated the role of both the intracellular formate pool and extracellular

formate supplementation in modulating antibiotic resistance in STM and evaluated their

potential as novel therapeutic targets. Our findings indicate that membrane damage resulting

from dysregulated intracellular pH (ipH) in STM influences the expression of the efflux pump

acrB through activation of the sigma factor RpoE. Additionally, extracellular formate

supplementation leads to downregulation of *acrB* expression via the BarA/SirA-regulated

CsrA/*csrB* regulatory pathway. To the best of our knowledge, this is the first report

demonstrating the impact of intracellular formate levels on antibiotic susceptibility in STM,

and how intracellular and extracellular formate pools distinctly modulate its antimicrobial

response.

Materials and Methods

Bacterial strains and culture conditions

The wild-type *Salmonella enterica* subsp. *enterica* serovar Typhimurium strain 14028S used

in this study was kindly provided by Professor Michael Hensel (Max von Pettenkofer Institute

for Hygiene and Medical Microbiology, Germany). All bacterial strains were preserved as

glycerol stocks at $-80\,°C$ and revived on Luria-Bertani (LB) agar plates (HiMedia)

supplemented with the appropriate antibiotics when necessary (50 µg/ml kanamycin, 50 µg/ml

ampicillin, or 25 µg/ml chloramphenicol). Single colonies were inoculated into LB broth

supplemented with the appropriate antibiotics and incubated overnight at $37\,°C$ with shaking

at 170 rpm. The overnight cultures were then diluted 1:100 in fresh LB medium and grown

under the same conditions for 6 hours to obtain bacteria in the logarithmic growth phase. The

optical density at 600 nm (OD_{600}) was subsequently adjusted to 0.1 in Mueller-Hinton Broth

(MHB), corresponding to approximately 10^6 colony-forming units (CFU), for use in

downstream experiments. A detailed list of bacterial strains and plasmids used in this study is

provided in Table 1.

Table 1

Bacterial strains/plasmids	Characteristics	Source
Salmonella enterica subspecies *enterica* serovar Typhimurium strain 14028S	Wild type	Gift from Prof. M. Hensel
STM Δ*pflB*	Kan^R	Laboratory stock
STM Δ*pflB*/pQE60: *pflB*	Kan^R, Amp^R	Laboratory stock
STM Δ*fliC*	Kan^R	Laboratory stock
STM Δ*rpoE*	Chl^R	Laboratory stock
STM Δ*pflB*Δ*rpoE*	Kan^R, Chl^R	Laboratory stock
STM Δ*sirA*	Chl^R	Laboratory stock
STM Δ*pflB*Δ*sirA*	Kan^R, Chl^R	Laboratory stock
pQE60	Low copy number plasmid, Amp^R	Laboratory stock

Determination of MIC of meropenem, ciprofloxacin, ceftazidime

Logarithmic phase bacterial cultures, adjusted to an OD$_{600}$ of 0.1, were inoculated into 96-well

microtiter plates containing Mueller-Hinton Broth (MHB) supplemented with serial dilutions

of antibiotics. The concentrations used were as follows: meropenem (8 to 0.0156 mg/L)

(Sigma), ciprofloxacin (0.5 to 0.0009 mg/L) (Sigma), and ceftazidime (128 to 0.125 mg/L).

Plates were incubated at 37 °C with shaking at 170 rpm for 14-16 hours. Bacterial growth was

assessed by measuring OD$_{600}$ using a Tecan microplate reader.

Estimation of bacterial viability using resazurin assay

Following antibiotic exposure and incubation for 14-16 hours, 20 µL of resazurin solution (prepared from a 0.2 mg/mL stock) was added to each well and incubated for 2 hours in the dark. Thereafter, fluorescence intensity was measured using a Tecan microplate reader with excitation at 540 nm and emission at 590 nm. Cell viability was determined based on fluorescence and expressed as a percentage of untreated controls.

Flow cytometry mediated quantification of live/dead population, intracellular reactive oxygen species (ROS), membrane depolarization, and ipH

After 14–16 hours of antibiotic treatment, bacterial cells were stained with a panel of fluorescent dyes: Propidium Iodide (PI, 1 µg/mL) for live/dead discrimination, dichlorodihydrofluorescein diacetate (DCF$_2$DA, 10 µM) to assess intracellular reactive oxygen species (ROS), Bis-(1,3-Dibutylbarbituric Acid) Trimethine Oxonol (DiBAC$_4$, 1 µg/mL) for measuring membrane depolarization, and 2′,7′-Bis-(2-Carboxyethyl)-5-(and-6)-Carboxyfluorescein Acetoxymethyl Ester (BCECF-AM, 20 µM) for intracellular pH (ipH) comparison. Stained samples were incubated in the dark for 15 minutes and subsequently analyzed using the BD FACSVerse flow cytometer (BD Biosciences, USA). Data including dot plots, density plots, median fluorescence intensity (MFI), and the percentage of positively stained cells were acquired and processed using BD FACSVerse software.

RNA isolation and RT-qPCR

Logarithmic-phase bacterial cultures were harvested by centrifugation at 6,000 rpm for 10 minutes at 4 °C, and the resulting cell pellets were resuspended in TRIzol reagent. Samples were stored at −80 °C until further processing. Total RNA was extracted using the chloroform–isopropanol method, and RNA pellets were washed with 70% ethanol and resuspended in 30 µL

of DEPC-treated Milli-Q water. RNA concentration was measured using a NanoDrop spectrophotometer (Thermo Fisher Scientific), and RNA integrity was verified by electrophoresis on a 2% agarose gel. For removal of genomic DNA, 2 µg of total RNA was treated with TURBO DNase (Thermo Fisher Scientific) at 37 °C for 30 minutes, followed by inactivation with 5 mM Na₂EDTA at 65 °C for 10 minutes. First-strand cDNA synthesis was performed using the PrimeScript RT reagent kit (Takara, Cat. No. RR037A) according to the manufacturer's protocol. Quantitative real-time PCR (qRT-PCR) was carried out using the TB Green RT-qPCR kit (Takara) on a QuantStudio 5 Real-Time PCR System. Gene expression was analysed using intragenic primers and normalized to *16S rRNA* levels. Primer sequences are provided in Table 2.

Table 2

Primer name	Sequence (5'-3')
16S *rRNA* RT FP	GTGAGGTAACGGCTCACCAA
16S *rRNA* RT RP	TAACCGCAACACCTTCCTCC
acrB RT FP	CTCTATCACCATCGTATCGG
acrB RT RP	AGTAATCCGTGACCTCATCC
tolC RT FP	TAACGAAGCGCGTAGTCCTT
tolC RT FP	AGCACATCAATAGCGTTCA
ompF RT FP	TGAATCCTATACCGATATGG
ompF RT RP	CCTGCTGATCGTTAGTACGC
emrB RT FP	TGGCGTATCCAGCAGCCTGA
emrB RT RP	GATGCCAATCACCAGTAGCG

Efflux pump activity by Nile Red Assay

Efflux pump activity was quantified following the method outlined by Lyu *et al*[11]. In brief,

bacterial cultures in the logarithmic growth phase were adjusted to an optical density of 0.1 at

600 nm (OD_{600}). A volume of 100 µl from each normalized culture was dispensed into wells of

a 96-well microplate and incubated with Nile Red dye at a final concentration of 48 µg/ml. The

plate was incubated in the dark under shaking conditions for 3 hours. Fluorescence

measurements were then obtained using a TECAN microplate reader, with excitation and

emission wavelengths set at 549 nm and 628 nm, respectively. Fluorescence intensity values

were normalized against absorbance readings at 600 nm.

Atomic Force Microscopy

After 14–16 hours of antibiotic exposure, bacterial cells were drop-cast onto sterile glass

coverslips. Once air-dried, the cells were fixed by treating them with 3.5% paraformaldehyde

for 10 minutes. Residual salts were removed by washing the samples three times with double-

autoclaved MilliQ water. The coverslips were then dried in a vacuum desiccator prior to

imaging with the NX10 Atomic Force Microscope (AFM). Image processing and

measurements of cell length and height were performed using XEI software.

Animal experiments

All animal experiments were conducted with prior approval from the Institutional Animal

Ethics Clearance Committee (IAEC) at the Indian Institute of Science, Bangalore. The institute

is registered under CPCSEA with registration number 48/1999/CPCSEA. All procedures

strictly adhered to the guidelines set by the Committee for the Purpose of Control and

Supervision of Experiments on Animals (CPCSEA). The ethical approval number for this study

is CAF/Ethics/853/2021.

A modified protocol based on the method described by Roy Chowdhury *et al.* was employed

to evaluate the organ burden following infection and antibiotic treatment with the wild-type

(WT) and mutant bacterial strains[12]. Briefly, 10^6 cells from overnight cultures were orally

gavaged into 6–8-week-old C57BL/6 mice on day 0 (0 dpi). Mice received 1 mg/kg of the

respective antibiotics on days 2 and 4 post-infection. On day 5 post-infection, the animals were

ethically euthanized, and the intestine, liver, and spleen were collected. These organs were

homogenized, and serial dilutions of the homogenates were plated on *Salmonella-Shigella* (SS)

agar. Bacterial load was quantified and expressed as $\log_{10}$ CFU per gram of tissue.

To compare mortality rates between STM WT and STM Δ*pflB* under antibiotic exposure,

groups of mice (n=5) were orally administered 10^6 cells from stationary phase cultures of each

strain. Starting from day 2 post-infection, 1 mg/kg of the respective antibiotics was

administered every alternate day till the mice survived. Survival was monitored daily and

represented as percent survival. A 125 mM solution of sodium formate was supplemented in

the drinking water provided to the animals in both experimental setups.

Estimation of Formate in mouse serum by Gas Chromatography-Mass Spectrometry

To quantify formate levels in mouse serum following supplementation with 125 mM sodium

formate in drinking water, a protocol adapted from Hughes *et al* was employed[13]. Male

C57BL/6 mice (6–8 weeks old) were provided with 125 mM sodium formate in their drinking

water. At five days post-infection, blood was collected via retro-orbital puncture. After clotting,

serum was separated by centrifugation at $10,000 \times$ g for 20 minutes and subsequently used for

formate quantification.

Sodium formate (Sigma) served as an external standard. Both standards and biological samples

were derivatized prior to mass spectrometry analysis. To protonate the formate, 2 M

hydrochloric acid was added to each sample in a 1:1 ratio. Liquid-liquid extraction was

performed to isolate formic acid into ethyl acetate (Sigma-Aldrich). Residual water was

removed by passing the ethyl acetate extract through an anhydrous sodium sulfate column

(Sigma-Aldrich). The extract was then incubated at 80 °C for 1 hour with the derivatizing

reagent   N,O-Bis(trimethylsilyl)trifluoroacetamide   (Sigma-Aldrich)   containing   1%

chlorotrimethylsilane (Fluka). Derivatized samples were transferred to autosampler vials for

analysis using Gas Chromatography-Mass Spectrometry (GC-MS; Agilent Technologies,

7890A GC, 5975C MS).

Injection parameters were as follows: injection temperature of 200 °C, injection split ratio of

1:40, and injection volume of 1 μL. The oven temperature program began at 40 °C (held for 4

minutes), increased to 120 °C at a rate of 5 °C/min, and then to 200 °C at 25 °C/min, with a

final hold of 3 minutes. Helium was used as the carrier gas at a constant flow rate of

1.2 mL/min. Chromatographic separation was performed using a 30 m × 0.25 mm × 0.40 μm

DB-624 column (Agilent). The interface temperature was maintained at 220 °C, and the

electron impact ion source operated at 230 °C with an ionization energy of 70 eV and 1235 EM

volts.

For qualitative analysis, selected ion monitoring in single quadrupole mode was employed with

a scan at m/z 103. The retention time for formate and its deuterated analog was approximately

5.9 minutes. The target and reference (qualifier) ions for formate were m/z = 103 and m/z =

75, respectively. Sample concentrations were determined by interpolating from a standard

calibration curve generated using known concentrations.

Determination of the fraction of antibiotic persister

The determination of antibiotic persister was performed by modifying a protocol described by Drescher et al[14]. Logarithmic phase cultures of STM WT and STM $\Delta pflB$ strains were treated with 15X and 25X the minimum inhibitory concentration (MIC) of meropenem (2 and ciprofloxacin. To determine the persister population, aliquots were collected at specified timepoints post-treatment and plated for colony-forming unit (CFU) enumeration. Similarly, 24-hour-old biofilm cultures were exposed to the same antibiotic concentrations, and persister cells were quantified via plating. The resulting CFU counts were normalized against the pre-inoculum values to calculate fold viability relative to the 0-hour timepoint. Data were expressed as $\log_{10}$(fold decrease in CFU relative to 0 h).

Statistical Analysis

Each experiment was independently repeated between 1 and 12 times, as specified in the figure legends. Statistical analyses were conducted using an unpaired Student's t-test, or two-way ANOVA, as indicated in the respective legends. Data from mouse experiments were analysed using the Mann–Whitney U test. A p-value of less than 0.05 was considered statistically significant. Results are presented as Mean ± SD or Mean ± SEM, as noted in the figure legends. All data analysis and graphical representations were performed using GraphPad Prism version 8.4.2.

Results

Deletion of *pflB* enhances susceptibility of STM to meropenem and ciprofloxacin, whereas supplementation of extracellular formate also heightens susceptibility of STM to these antibiotics

To evaluate the susceptibility of STM Δ*pflB* to meropenem and ciprofloxacin, both the wild-

type and mutant strains were exposed to varying concentrations of the antibiotics. The STM

Δ*pflB* mutant showed significantly increased sensitivity to meropenem, particularly at

concentrations of 1 mg/L and 0.5 mg/L **(Figure 1A)**. While the partial restoration of bacterial

growth upon sodium formate supplementation was not statistically significant, it suggests that

intracellular formate deficiency may contribute to the heightened sensitivity to meropenem.

Interestingly, when formate was added to wild-type cells—despite their intact intracellular

formate levels—a significant reduction in survivability was observed under antibiotic stress at

the same concentrations. Comparable results were obtained with ciprofloxacin, where

statistically significant effects were noted at 0.03125, 0.015625, 0.0078, and 0.0039 mg/L

(Figure 1C). These findings were further supported by resazurin assay data **(Figure 1B, D)**.

The enhanced susceptibility of STM ΔpflB and its recovery in the presence of sodium formate

was also true for ceftazidime, a third generation cephalosporin[15]**(Figure S1)**.

To investigate how deletion of *pflB* affects STM's antibiotic susceptibility, we treated

logarithmic-phase cultures overnight with sub-lethal doses of meropenem (0.5 mg/L) and

ciprofloxacin (0.015625 mg/L)and analyzed Propidium Iodide (PI) uptake via flow cytometry.

STM Δ*pflB* showed a higher PI-positive population than wild-type (WT), which was

significantly reduced with sodium formate supplementation. In contrast, formate

supplementation increased PI-positive cells in WT upon antibiotic exposure **(Figure 2A, B)**.

Given that bactericidal antibiotics induce reactive oxygen species (ROS) irrespective of their

mechanisms, we assessed intracellular ROS using DCF_2DA staining[16-18]. Consistent with PI

staining results, STM Δ*pflB* exhibited elevated DCF_2DA-positive populations under antibiotic

stress, which was mitigated by formate. Similarly, formate supplementation increased ROS

levels in WT cells exposed to antibiotics **(Figure S2 A, B)**. These findings suggest that loss of

pflB, and thus intracellular formate, increases STM's susceptibility to meropenem and

ciprofloxacin, while exogenous formate to wild-type cells also enhances antibiotic sensitivity.

Deletion of *pflB* was associated with a transcriptional downregulation of antibiotic efflux

pumps, which was also common for formate supplemented in wild-type cells

Previous studies have demonstrated that cytoplasmic acidification is a key factor contributing

to antibiotic resistance in *Escherichia coli*[19]. Similarly, work by Panta *et al*. has shown that

maintenance of optimal cytoplasmic pH is critical for colistin resistance in various bacterial

species, including *S. enterica*[20]. To investigate whether intracellular pH dynamics play a role

in antibiotic susceptibility in our system, we exposed bacterial strains to sub-lethal

concentrations of meropenem and ciprofloxacin and measured intracellular pH using BCECF-

AM, a cell-permeable, dual-excitation ratiometric dye[21] **(Figure 2C)**. A reduced 488/405

fluorescence ratio in STM WT cells following antibiotic exposure indicated cytoplasmic

acidification under antibiotic stress. Consistent with previous observations, STM Δ*pflB*

exhibited a higher baseline 488/405 ratio in the absence of stress, which remained unchanged

upon antibiotic treatment, suggesting a resistance to cytoplasmic acidification.

Supplementation with sodium formate partially restored the 488/405 ratio in STM Δ*pflB*,

indicating a modulatory effect of formate on intracellular pH in the presence of antibiotics.

While formate supplementation alone did not significantly alter the intracellular pH of STM

WT, an increase in the 488/405 ratio was observed upon antibiotic exposure, further supporting

the role of formate in buffering pH under stress conditions.

Our previous work demonstrated increased membrane depolarization in STM Δ*pflB* compared

to WT cells[10]. Given that antimicrobial peptides exert bactericidal effects via membrane

depolarization we hypothesized that the enhanced antibiotic susceptibility of STM Δ*pflB* may

be attributable, at least in part, to its depolarized membrane state[22-26]. Notably, membrane

depolarization was further exacerbated upon exposure to sub-lethal concentrations of

meropenem and ciprofloxacin, while sodium formate partially alleviated this effect in STM

$\Delta pflB$. In contrast, formate supplementation had no impact on membrane potential in WT cells

under unstressed conditions, although increased depolarization was observed upon antibiotic

exposure **(Figure S2 C, D)**.

Efflux-mediated resistance represents a major mechanism of antibiotic failure in Gram-

negative bacteria. *Salmonella* harbors multiple efflux systems, among which the TolC-

dependent resistance-nodulation-division (RND) family plays a central role in expelling a

broad range of clinically relevant antibiotics[27-30]. As proton motive force (PMF) and pH

homeostasis are critical for the functioning of these pumps, and membrane depolarization can

disrupt PMF, we hypothesized that the reduced antibiotic efflux in STM $\Delta pflB$ may stem from

impaired PMF[11]. To test this, we employed a nile red efflux assay. STM $\Delta pflB$ retained

significantly higher levels of nile red fluorescence, indicating decreased efflux activity.

Formate supplementation reduced nile red retention, suggesting partial restoration of efflux

capacity. Similarly, formate-treated STM WT cells exhibited increased nile red fluorescence

upon antibiotic exposure, consistent with reduced efflux function **(Figure 2D)**.

To determine whether reduced efflux activity was due solely to membrane depolarization or

also involved transcriptional regulation, we measured mRNA levels of *acrB* and *tolC*, two key

components of the RND efflux system in *Salmonella*[31]. Deletion of *pflB* resulted in

transcriptional downregulation of both genes, with partial recovery upon formate

supplementation. Interestingly, formate supplementation in STM WT also led to reduced *acrB*

and *tolC* expression **(Figure 2 E, F)**. These findings suggest that while membrane

depolarization may directly impair efflux pump activity, additional transcriptional regulatory

mechanisms may also contribute to reduced efflux and heightened antibiotic susceptibility.

Outer membrane proteins (Omps) have been linked to antibiotic susceptibility in *Salmonella*,

with reduced expression of the large porin OmpF commonly associated with resistance[21, 32, 33].

Interestingly, we observed downregulation of *ompF* in STM Δ*pflB* **(Figure 2G)**, despite its

increased antibiotic sensitivity. This suggests that the heightened susceptibility in STM Δ*pflB*

occurs independently of Omps.

Formate supplementation in drinking water and intrabacterial formate deficiency both

enhanced bacterial clearance by antibiotics *in-vivo*

To investigate whether our *in-vitro* findings held true in an *in-vivo* setting, we administered 10^6

CFU of STM WT and STM Δ*pflB* orally to C57BL/6 mice. Alongside this, the mice received

125 mM sodium formate in their drinking water **(Figure 3A)**. Previous studies have estimated

that mice require approximately 40 mmol of formate daily. Given that C57BL/6 mice typically

consume around 30 ml of water per day, a 1 M concentration would theoretically be necessary

to meet this demand. However, prior research has indicated that providing 500 mM sodium

formate in drinking water can lead to weight loss in mice. Therefore, we chose a safer

concentration of 125 mM, which corresponds to about 10% of the estimated daily requirement,

to stay within physiological limits[34]. In our pilot study, this supplementation level was

sufficient to significantly elevate serum formate levels by 5 days post-infection (dpi) **(Figure**

S3).

Our results showed that combining formate supplementation with meropenem treatment led to

a greater reduction in bacterial load compared to meropenem alone, across the intestine, liver,

and spleen of C57BL/6 mice. Consistent with earlier findings, the STM Δ*pflB* mutant exhibited

a lower organ burden[10]; however, formate supplementation did not restore its bacterial counts.

Meropenem further suppressed bacterial levels, and, contrary to our *in-vitro* findings, formate

in the drinking water did not rescue the reduced bacterial numbers. Notably, combining formate

with meropenem did not lead to a reduction in organ colonization for STM Δ*pflB* as seen in the

STM WT, suggesting that STM Δ*pflB* may utilize extracellular formate to support survival

under antibiotic pressure **(Figure 3B)**. Similar patterns were observed with ciprofloxacin

treatment in STM WT and STM Δ*pflB*-infected mice **(Figure 3C)**. These findings were further

corroborated with ceftazidime exposure, where comparable effects were observed in mice

infected with STM WT and STM Δ*pflB* under formate-supplemented conditions **(Figure 3D)**.

Deletion of *rpoE* from STM Δ*pflB* reversed the enhanced antibiotic susceptibility,

associated with a recovery in the transcript levels of *acrB* and *tolC*

In our previous work, we demonstrated that deletion of *pflB* disrupts intracellular pH

homeostasis, leading to membrane depolarization. This perturbation activated the

extracytoplasmic sigma factor RpoE, which in turn upregulated transcription of the small RNA

csrB. The *csrB* RNA antagonizes the activity of the global regulator CsrA by direct binding,

thereby impairing processes such as flagellar biosynthesis[10]. Notably, CsrA has been shown to

stabilize the *acrAB* transcript by interacting with its 5′ untranslated region (UTR)[35]. Given the

increased membrane depolarization observed in STM Δ*pflB* upon antibiotic exposure, we

hypothesized that RpoE may regulate *acrB* expression indirectly through modulation of the

CsrA–*csrB* regulatory circuit.

To test this, we exposed the STM Δ*pflB*Δ*rpoE* strain to sublethal concentrations of meropenem

and ciprofloxacin. Compared to STM Δ*pflB*, the double knockout exhibited a significantly

reduced proportion of propidium iodide (PI)-positive cells **(Figure 4A, B)**. This was

accompanied by a decrease in intracellular reactive oxygen species (ROS), as assessed using

DCF$_2$DA staining **(Figure S4A, B)**.

Interestingly, deletion of *rpoE* in the STM Δ*pflB* background did not restore membrane

potential under basal conditions. However, upon antibiotic challenge, we observed a significant

reduction in DiBAC$_4$-positive cells, indicative of partial recovery in membrane polarization

(Figure S4C, D). The improved viability of STM Δ*pflB*Δ*rpoE* relative to STM Δ*pflB* under

antibiotic stress was further supported by atomic force microscopy, which revealed increased

cellular length and height following exposure to both meropenem and ciprofloxacin **(Figure**

S5A, B, C).

These phenotypic changes correlated with the restoration of *acrB* and *tolC* transcript levels in

STM Δ*pflB*Δ*rpoE*, suggesting enhanced efflux activity **(Figure 4C, D)**. This was further

confirmed by increased nile red efflux, indicative of improved pump function **(Figure 4E)**.

Given the role of EmrB in fluoroquinolone resistance, we also examined its expression **(Figure**

4F)[36]. As previous studies have not identified a direct regulatory relationship between CsrA

and *emrB*, we measured *emrB* transcript levels and observed no significant change in STM

Δ*pflB*Δ*rpoE* relative to STM Δ*pflB*. These findings suggest that the observed phenotypic rescue

is primarily mediated through the CsrA–*csrB* axis. Deletion of *rpoE* in STM Δ*pflB* led to

further downregulation of *ompF*, indicating that while *ompF* suppression does not explain the

increased antibiotic sensitivity of STM Δ*pflB*, its further downregulation upon *rpoE* deletion

may contribute to the restored resistance observed in STM Δ*pflB*Δ*rpoE* **(Figure 4G)**.

Deletion of *sirA* abolishes the antibiotic susceptibility of STM Δ*pflB*, along with reduction

in ROS levels and membrane depolarization

To further confirm that small RNA *csrB* regulates antibiotic susceptibility, we examined an

additional upstream regulator of *csrB*. The BarA/SirA two-component system, which is also

known to activate the expression of *csrB* and *csrC* small RNAs, has previously been shown to

enhance the expression of HilD, a key regulator of genes within the *Salmonella* Pathogenicity

Island-1[37-39]. We deleted *sirA* in the STM Δ*pflB* background and used flow cytometry to

quantify propidium iodide-positive cells, revealing a marked reduction in dead cells in the STM

$\Delta pflB\Delta sirA$ strain compared to STM $\Delta pflB$ **(Figure 5A, B)**. Interestingly, this decrease in cell

death did not correspond with a reduction in membrane depolarization, indicating that the

reduced antibiotic sensitivity in STM $\Delta pflB\Delta sirA$ was independent of changes in membrane

potential **(Figure S6A, B)**. A drop in DCF_2DA-positive cells further supported the observation

of diminished antibiotic susceptibility following *sirA* deletion in STM $\Delta pflB$. Notably, STM

$\Delta sirA$ alone showed an increase in DCF_2DA-positive cells after exposure to meropenem and

ciprofloxacin **(Figure S6 C, D)**. This prompted an investigation into efflux pump expression.

We observed that *sirA* deletion in STM $\Delta pflB$ led to elevated expression of *acrB*, *tolC*, *emrB*,

and *ompF* **(Figure 5C, D, E, F)**. A nile red assay confirmed restored efflux pump activity in

STM $\Delta pflB\Delta sirA$ **(Figure 5G)**. Interestingly, deletion of *sirA* alone resulted in decreased *acrB*

transcript levels, which may explain the increased DCF_2DA-positive population upon

treatment with meropenem and ciprofloxacin in STM $\Delta sirA$ **(Figure S6 C, D)**. However, no

overall change in efflux activity was detected in STM $\Delta sirA$, possibly due to compensatory

roles by other efflux pumps **(Figure 5G)**.. These findings suggest that targeting upstream

regulators of *csrB* can effectively reverse the antibiotic susceptibility observed in STM $\Delta pflB$.

Deletion of *pflB* impaired the long-term survival of STM under high antibiotic

concentrations, a similar effect observed when formate was supplemented to the wild type

Bacterial persistence in the presence of antibiotics, distinct from antibiotic resistance, remains

a persistent challenge in effectively eliminating pathogens in clinical settings[40]. Since our data

demonstrated that *pflB* deletion increased the antibiotic susceptibility of STM, we aimed to

investigate its impact on long-term bacterial survival under high antibiotic stress. Log-phase

bacterial cultures, with or without formate supplementation, were exposed to 15× and 25× MIC

concentrations of meropenem and ciprofloxacin. Exposure to 15× MIC of meropenem led to a

rapid decline in bacterial countsin planktonic culture, with STM $\Delta pflB$ exhibiting the fastest

reduction, reaching statistical significance at 24 hours post-treatment **(Figure 6A)**. Formate

supplementation restored survival levels of STM Δ*pflB* to those of the wild type, while

supplementation in STM WT also caused a significant reduction in CFUs. Interestingly, by 48

and 72 hours, all strains—regardless of formate treatment—showed similar fold decreases in

bacterial counts. This effect was specific to planktonic cultures treated with meropenem and

was not observed in biofilm forms or in ciprofloxacin-treated planktonic and biofilm cultures

(Figure 6B, C, D).

Given that the increased susceptibility in earlier experiments was linked to elevated

intracellular pH, we examined whether this condition persisted in long-term meropenem-

treated cells **(Figure 6E)**. CFUs from 72-hour survivors were stained with BCECF-AM to

assess cytosolic acidification. Under untreated conditions, STM Δ*pflB* showed a significantly

higher 488/405 fluorescence ratio, indicating elevated intracellular pH. However, after

meropenem treatment, all strains—regardless of formate treatment—exhibited similar high

levels of cytosolic acidification. These results suggest that intracellular pH plays a key role in

bacterial persistence and survival under prolonged, high-dose antibiotic exposure.

Discussion

Salmonella has become a significant food-borne pathogen exhibiting widespread drug

resistance, with the rise of multidrug-resistant strains becoming increasingly concerning[41-43].

The growing resistance to existing treatments has complicated clinical management and

highlights the urgent need for new antimicrobials or repurposing the existing ones[44, 45].

Historically, antimicrobial resistance (AMR) research has largely focused on molecular

mechanisms[46, 47]. These include how *Salmonella* and other bacteria resist antibiotics through

mechanisms such as modifying antibiotic targets, enzymatic degradation of drugs, and changes

in membrane permeability[48-51].

More recently, attention has shifted towards the role of bacterial metabolism and metabolic by-

products in influencing AMR. For instance, studies have shown that ceftriaxone-resistant

Salmonella Derby displays altered metabolic activity, particularly in glutathione biosynthesis,

with notable decreases in oxidized glutathione and citrulline levels. Supplementing these

metabolites externally was found to restore ceftriaxone susceptibility in multidrug-resistant

Salmonella Derby[52].

Several studies have also highlighted the significance of pH in influencing antimicrobial

resistance in *Salmonella* and other bacterial species[11, 19, 20, 53]. This underlines the importance

of investigating metabolic pathways that are crucial for pH regulation. The enzyme pyruvate-

formate lyase (PflB), which catalyzes the conversion of pyruvate into formate and acetyl-CoA,

plays a key role in maintaining intracellular pH (ipH)[10]. Consistent with this, our research

showed that *Salmonella* Typhimurium lacking *pflB* was more vulnerable to ciprofloxacin and

meropenem. Interestingly, this increased susceptibility was partially alleviated by

supplementing formate externally.

While it is well known that pH affects the proton motive force (PMF), thereby influencing the

activity of efflux pumps, our data indicate that pH also regulates efflux pump expression

through transcriptional downregulation of *acrB* and *tolC*[11]. In earlier work, we observed that

ipH disruption-induced membrane damage activated the RpoE sigma factor, leading to

increased expression of the small RNA *csrB*. Deleting *rpoE* reversed the non-flagellated

phenotype seen in STM Δ*pflB*[10]. In the current study, we further found that *rpoE* deletion also

reversed the increased antibiotic sensitivity of STM Δ*pflB*. A similar effect was seen with the

deletion of *sirA*, another upstream regulator of *csrB*. These findings suggest that *pflB* regulates

both *Salmonella* invasion and antibiotic resistance through a shared regulatory pathway.

Other studies in *E. coli* have demonstrated that adenosine can suppress the stringent response,

enhance PMF, promote respiration, and increase antibiotic-induced killing of tolerant and

stationary phase persister cells—a phenomenon described as "persister awakening"[54]. Since

persistence poses a major hurdle in treating infections, we investigated the long-term survival

of STM Δ*pflB* under high-dose antibiotic exposure. In planktonic cultures, STM Δ*pflB*

exhibited a faster killing rate than the wild-type, a phenotype reversed by external formate.

Moreover, formate supplementation to STM WT enhanced bacterial killing. Cytoplasmic pH

analysis using BCECF-AM staining after 72 hours revealed that surviving bacteria—regardless

of formate exposure—shared similar pH levels when treated with meropenem. This implies

that cytoplasmic pH plays a key role in bacterial survival under antibiotic stress, and that

surviving bacteria may carry mutations allowing pH regulation, although characterizing these

mutations was beyond the scope of this study and warrants future investigation.

Overall, the effectiveness of antibiotics is closely tied to the metabolic state of bacteria, with

certain pathways being linked to specific resistance profiles. Mapping these pathways and

targeting them with selective inhibitors that shift the metabolic state of resistant bacteria toward

that of susceptible ones could offer a promising strategy to combat antibiotic resistance.

Acknowledgements

We thank the Departmental Real-Time PCR, Divisional Mass Spectrometry, Divisional Flow

Cytometry Facility, Divisional Atomic Force Microscopy, and the Central Animal Facility at

IISc for supporting experimental work. Dr. Sunitha assisted in formate quantification via GC-

MS. Flow cytometry data acquisition was supported by Mr. Munish and Mrs. Navya. Atomic

Force Microscopy was supported by Mrs. Monisha.

Author Contributions

DM and DC conceived the study and designed the experiments. DM conducted all experiments, data acquisition, analysis, figure preparation, and wrote the original draft. SVW, PG, and AB helped DM with experiments. DM and DC contributed to manuscript editing. DC supervised the project and secured funding. All authors reviewed and approved the final manuscript.

Declaration of Interest

The authors declare no conflicts of interest.

Funding Information

We gratefully acknowledge financial support from the Department of Biotechnology (DBT) and the Department of Science and Technology (DST), Ministry of Science and Technology. DC thanks DAE for the SRC Outstanding Investigator Award, ASTRA Chair Professorship, and TATA Innovation Fellowship. Additional support came from the DBT-IISc Partnership Program. Infrastructure support from ICMR (Molecular Medicine Center), DST (FIST), and UGC-CAS is also acknowledged. DM is supported by IISc fellowship. PG is supported by DBT-JRF fellowship. AB acknowledges UGC fellowship. The funders had no involvement in study design, data collection, analysis, or publication decisions.

Data Availability

Data supporting this study are available from the authors upon request.

Figure legends

Figure 1

Figure 1: Deletion of *pflB* increases STM's sensitivity to meropenem and ciprofloxacin,

while extracellular formate supplementation similarly enhances antibiotic susceptibility.

Meropenem and ciprofloxacin susceptibility of STM WT, STM Δ*pflB*, and STM Δ*pflB*/pQE60:

pflB, with or without formate treatment, was assessed using the microfold dilution method (A,

C) and the resazurin assay (B, D).

(Unpaired two-tailed Student's t-test for column graphs, Two-way Anova for grouped data,

Mann-Whitney U-test for animal experiment data (*** p < 0.0001, *** p < 0.001, ** p<0.01,

* p<0.05))

Figure 2

Figure 2: Increased antibiotic sensitivity following *pflB* deletion was linked to

transcriptional downregulation of efflux pumps, a pattern also observed in formate-

treated wild-type cells.

Dot plots (SSC-A vs PI-A) and bar graphs (A, B) illustrate increased bacterial killing with

meropenem and ciprofloxacin following *pflB* deletion and formate supplementation in STM

WT. BCECF dye analysis shows disrupted cytosolic pH in STM Δ*pflB* under antibiotic stress,

which is restored by exogenous formate (C). Nile red assay quantifies efflux pump activity

(D), while transcript levels of *acrB*, *tolC*, and *ompF* are measured in STM WT and STM

Δ*pflB*, with or without formate treatment (E, F, G).

(Unpaired two-tailed Student's t-test for column graphs, Two-way Anova for grouped data,

Mann-Whitney U-test for animal experiment data (*** p < 0.0001, *** p < 0.001, ** p<0.01,

* p<0.05))

Figure 3

Figure 3: Both formate supplementation via drinking water and internal formate

deficiency enhanced antibiotic-mediated bacterial clearance *in vivo*.

Schematic outlining the experimental design (A). Formate supplementation improved STM

WT clearance from the intestine, liver, and spleen following treatment with meropenem (B),

ciprofloxacin (C), and ceftazidime (D). STM Δ*pflB* showed increased *in vivo* killing upon

antibiotic treatment, while exogenous formate helped maintain its bacterial load in mice (B, C,

D).

(Unpaired two-tailed Student's t-test for column graphs, Two-way Anova for grouped data,

Mann-Whitney U-test for animal experiment data (*** p < 0.0001, *** p < 0.001, ** p<0.01,

* p<0.05))

Figure 4

Figure 4: Deletion of *rpoE* in STM Δ*pflB* reversed its increased antibiotic sensitivity, accompanied by restored *acrB* and *tolC* transcript levels.

Dot plots (SSC-A vs PI-A) and bar graphs (A, B) show reduced antibiotic-mediated killing in STM Δ*pflB* upon *rpoE* deletion. Transcript levels of key efflux pump genes *acrB* and *tolC* were compared between STM Δ*pflB*Δ*rpoE* and STM Δ*pflB* (C, D). Efflux activity was assessed using the Nile red assay (E), and expression of *emrB* and *ompF* was also evaluated (F, G). (Unpaired two-tailed Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (*** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$))

Figure 5

Figure 5: Deleting *sirA* in STM Δ*pflB* similarly reversed its heightened antibiotic sensitivity and restored *acrB* and *tolC* expression levels.

Dot plots (SSC-A vs PI-A) and bar graphs (A, B) show reduced antibiotic-mediated killing in STM Δ*pflB* upon *sirA* deletion. Transcript levels of efflux pump genes *acrB, tolC*, *emrB,* and *ompF* were compared between STM Δ*pflB*Δ*sirA* and STM Δ*pflB* (C, D, E, F). Efflux activity was assessed using the Nile red assay (G).

(Unpaired two-tailed Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (*** p < 0.0001, *** p < 0.001, ** p<0.01, * p<0.05))

Figure 6

Figure 6: Deletion of *pflB* reduced STM's long-term survival under high antibiotic concentrations, an effect also seen with formate supplementation in the wild type.

Line graphs show $\log_{10}$ fold reduction in CFU relative to 0h for planktonic (A, B) and biofilm (C, D) cultures of STM WT and STM $\Delta pflB$, with or without formate supplementation, following exposure to high doses of meropenem and ciprofloxacin at 4, 8, 24, 48, and 72 hours. BCECF staining was performed on 72h colonies, with or without meropenem treatment (E).

(Unpaired two-tailed Student's t-test for column graphs, Two-way Anova for grouped data, Mann-Whitney U-test for animal experiment data (*** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$))

Figure S1

Figure S1: Deletion of *pflB* increases STM's susceptibility to the third-generation

cephalosporin ceftazidime, and similar sensitivity is observed with extracellular formate

supplementation.

Ceftazidime susceptibility in STM WT, STM Δ*pflB*, and STM Δ*pflB*/pQE60:*pflB*, with or

without formate, was evaluated using the microfold dilution method.

(Unpaired two-tailed Student's t-test (column graphs), two-way ANOVA (grouped data), and

Mann–Whitney U-test (animal data).

(***p < 0.0001, **p < 0.001, **p < 0.01, *p < 0.05))

Figure S2

N=3, n=3

Figure S2: Enhanced antibiotic sensitivity due to *pflB* deletion correlates with increased

DCF2DA and DiBAC4 staining, indicating elevated ROS levels and membrane

depolarization—effects also seen in formate-treated STM WT.

Dot plots (SSC-A vs DCF$_2$DA/DiBAC-A) (A, C) and corresponding bar graphs (B, D) show

higher intracellular ROS and membrane depolarization following meropenem and

ciprofloxacin treatment in STM ΔpflB and formate-supplemented STM WT.

(Unpaired two-tailed Student's t-test (column graphs), two-way ANOVA (grouped data), and

Mann–Whitney U-test (animal data).

(***$p < 0.0001$, **$p < 0.001$, **$p < 0.01$, *$p < 0.05$))

Figure S3

Figure S3: Formate supplementation increased formate levels in serum of treated mice

Bar graph showing formate levels in mice serum after exposure to formate through drinking

water

(Unpaired two-tailed Student's t-test (column graphs), two-way ANOVA (grouped data), and

Mann–Whitney U-test (animal data).

(***$p < 0.0001$, **$p < 0.001$, **$p < 0.01$, *$p < 0.05$))

Figure S4

N=3, n=4

N=3, n=4

Figure S4: Recovered antibiotic sensitivity due to *rpoE* deletion in STM Δ*pflB* correlates with reduced intracellular ROS levels but it did not have any effect on membrane depolarization

Dot plots (SSC-A vs DIBAC-A/ DCF$_2$DA) (A, C) and corresponding bar graphs (B, D) compares intracellular ROS and membrane depolarization following meropenem and ciprofloxacin treatment in STM Δ*pflB* and STM Δ*pflB*Δ*rpoE*.

(Unpaired two-tailed Student's t-test (column graphs), two-way ANOVA (grouped data), and Mann–Whitney U-test (animal data).

(***p < 0.0001, **p < 0.001, **p < 0.01, *p < 0.05))

Figure S5

Figure S5: The recovery in antibiotic sensitivity upon *rpoE* deletion in STM Δ*pflB* was

also evident from morphological changes under Atomic Force Microscopy

AFM-assissted visualization of morphology of STM WT, STM Δ*pflB*, STM Δ*rpoE*, STM

Δ*pflB*Δ*rpoE* upon expsure to sublethal concentrations of meropenem and ciprofloxacin (A) and

quantification of bacterial length (B) and height (C) from the same.

(Unpaired two-tailed Student's t-test (column graphs), two-way ANOVA (grouped data), and

Mann–Whitney U-test (animal data).

(***$p < 0.0001$, **$p < 0.001$, **$p < 0.01$, *$p < 0.05$))

Figure S6

Figure S6: *sirA* deletion mediated reduction of antibiotic susceptibility in STM Δ*pflB* was associated with reduced intracellular ROS levels, but without any effect on membrane depolarization

Dot plots (SSC-A vs DIBAC-A/ DCF₂DA) (A, C) and corresponding bar graphs (B, D) shows a comparative analysis between intracellular ROS and membrane depolarization following meropenem and ciprofloxacin treatment in STM Δ*pflB* and STM Δ*pflB*Δ*sirA*.

References:

1.   Zhu S, Yang B, Jia Y et al. Comprehensive analysis of disinfectants on the horizontal transfer of antibiotic resistance genes. *J Hazard Mater* 2023; **453**: 131428.

2.   Brauner A, Fridman O, Gefen O et al. Distinguishing between resistance, tolerance and
persistence to antibiotic treatment. *Nat Rev Microbiol* 2016; **14**: 320-30.
3.   Antimicrobial Resistance C. Global burden of bacterial antimicrobial resistance in 2019:
a systematic analysis. *Lancet* 2022; **399**: 629-55.
4.   Azmatullah A, Qamar FN, Thaver D et al. Systematic review of the global epidemiology,
clinical and laboratory profile of enteric fever. *J Glob Health* 2015; **5**: 020407.
5.   Kim TH, Hwang HJ, Kim JH. Development of a Novel, Rapid Multiplex Polymerase Chain
Reaction Assay for the Detection and Differentiation of Salmonella enterica Serovars Enteritidis
and Typhimurium Using Ultra-Fast Convection Polymerase Chain Reaction. *Foodborne Pathog*
Dis 2017; **14**: 580-6.
6.   Zhang H, Yao S, Sheng R et al. A cascade amplification strategy for ultrasensitive
Salmonella typhimurium detection based on DNA walker coupling with CRISPR-Cas12a. *J Colloid*
Interface Sci 2022; **625**: 257-63.
7.   Wu S, Ji J, Sheng L et al. Lysine and valine weaken antibiotic resistance in Salmonella
Typhimurium induced by disinfectant stress. *J Hazard Mater* 2024; **480**: 135858.
8.   Punchihewage-Don AJ, Ranaweera PN, Parveen S. Defense mechanisms of Salmonella
against antibiotics: a review. *Front Antibiot* 2024; **3**: 1448796.
9.   Chowdhury AR, Mukherjee D, Chatterjee R et al. Defying the odds: Determinants of the
antimicrobial response of Salmonella Typhi and their interplay. *Mol Microbiol* 2024; **121**: 213-29.
10.   Mukherjee D, Chakravortty D. A Tale of Twins: The Distinct Roles of Intracellular and
Extracellular Formate in Regulating Flagellar and Pathogenicity Island-1 Genes in
Salmonella Typhimurium. *bioRxiv* 2024: 2024.01.30.577997.
11.   Lyu Z, Yang A, Villanueva P et al. Heterogeneous Flagellar Expression in Single Salmonella
Cells Promotes Diversity in Antibiotic Tolerance. *mBio* 2021; **12**: e0237421.
12.   Chowdhury AR, Mukherjee D, Singh AK et al. Loss of outer membrane protein A (OmpA)
impairs the survival of Salmonella Typhimurium by inducing membrane damage in the presence
of ceftazidime and meropenem. *J Antimicrob Chemother* 2022; **77**: 3376-89.
13.   Hughes ER, Winter MG, Duerkop BA et al. Microbial Respiration and Formate Oxidation as
Metabolic Signatures of Inflammation-Associated Dysbiosis. *Cell Host Microbe* 2017; **21**: 208-19.
14.   Drescher SPM, Gallo SW, Ferreira PMA et al. Salmonella enterica persister cells form
unstable small colony variants after in vitro exposure to ciprofloxacin. *Sci Rep* 2019; **9**: 7232.
15.   Papich MG. Saunders handbook of veterinary drugs. *North Carolina* 2016; **12**: 162-71.
16.   Dwyer DJ, Kohanski MA, Hayete B et al. Gyrase inhibitors induce an oxidative damage
cellular death pathway in Escherichia coli. *Mol Syst Biol* 2007; **3**: 91.
17.   Kohanski MA, Dwyer DJ, Hayete B et al. A common mechanism of cellular death induced
by bactericidal antibiotics. *Cell* 2007; **130**: 797-810.
18.   Dwyer DJ, Belenky PA, Yang JH et al. Antibiotics induce redox-related physiological
alterations as part of their lethality. *Proc Natl Acad Sci U S A* 2014; **111**: E2100-9.
19.   Reyes-Fernandez EZ, Schuldiner S. Acidification of Cytoplasm in Escherichia coli
Provides a Strategy to Cope with Stress and Facilitates Development of Antibiotic Resistance. *Sci*
Rep 2020; **10**: 9954.
20.   Panta PR, Doerrler WT. A link between pH homeostasis and colistin resistance in bacteria.
Sci Rep 2021; **11**: 13230.
21.   Roy Chowdhury A, Sah S, Varshney U et al. Salmonella Typhimurium outer membrane
protein A (OmpA) renders protection from nitrosative stress of macrophages by maintaining the
stability of bacterial outer membrane. *PLoS Pathog* 2022; **18**: e1010708.
22.   Epand RF, Pollard JE, Wright JO et al. Depolarization, bacterial membrane composition,
and the antimicrobial action of ceragenins. *Antimicrob Agents Chemother* 2010; **54**: 3708-13.
23.   Te Winkel JD, Gray DA, Seistrup KH et al. Analysis of Antimicrobial-Triggered Membrane
Depolarization Using Voltage Sensitive Dyes. *Front Cell Dev Biol* 2016; **4**: 29.

24. Han FF, Gao YH, Luan C et al. Comparing bacterial membrane interactions and
antimicrobial activity of porcine lactoferricin-derived peptides. *J Dairy Sci* 2013; **96**: 3471-87.
25. Soren O, Brinch KS, Patel D et al. Antimicrobial Peptide Novicidin Synergizes with
Rifampin, Ceftriaxone, and Ceftazidime against Antibiotic-Resistant Enterobacteriaceae In Vitro.
Antimicrob Agents Chemother 2015; **59**: 6233-40.
26. Cheng M, Huang JX, Ramu S et al. Ramoplanin at bactericidal concentrations induces
bacterial membrane depolarization in Staphylococcus aureus. *Antimicrob Agents Chemother*
2014; **58**: 6819-27.
27. Masi M, Refregiers M, Pos KM et al. Mechanisms of envelope permeability and antibiotic
influx and efflux in Gram-negative bacteria. *Nat Microbiol* 2017; **2**: 17001.
28. Du D, Wang-Kan X, Neuberger A et al. Multidrug efflux pumps: structure, function and
regulation. *Nat Rev Microbiol* 2018; **16**: 523-39.
29. Ramaswamy VK, Vargiu AV, Malloci G et al. Molecular Rationale behind the Differential
Substrate Specificity of Bacterial RND Multi-Drug Transporters. *Sci Rep* 2017; **7**: 8075.
30. Rosenberg EY, Ma D, Nikaido H. AcrD of Escherichia coli is an aminoglycoside efflux
pump. *J Bacteriol* 2000; **182**: 1754-6.
31. Webber MA, Bailey AM, Blair JM et al. The global consequence of disruption of the AcrAB-
TolC efflux pump in Salmonella enterica includes reduced expression of SPI-1 and other
attributes required to infect the host. *J Bacteriol* 2009; **191**: 4276-85.
32. Ziervogel BK, Roux B. The binding of antibiotics in OmpF porin. *Structure* 2013; **21**: 76-87.
33. Choi KM, Kim MH, Cai H et al. Salicylic Acid Reduces OmpF Expression, Rendering
Salmonella enterica Serovar Typhimurium More Resistant to Cephalosporin Antibiotics.
Chonnam Med J 2018; **54**: 17-23.
34. Dhayade S, Pietzke M, Wiesheu R et al. Impact of Formate Supplementation on Body
Weight and Plasma Amino Acids. *Nutrients* 2020; **12**.
35. Ricci V, Attah V, Overton T et al. CsrA maximizes expression of the AcrAB multidrug
resistance transporter. *Nucleic Acids Res* 2017; **45**: 12798-807.
36. Gu Y, Huang L, Wu C et al. The Evolution of Fluoroquinolone Resistance in Salmonella
under Exposure to Sub-Inhibitory Concentration of Enrofloxacin. *Int J Mol Sci* 2021; **22**.
37. Salvail H, Groisman EA. The phosphorelay BarA/SirA activates the non-cognate regulator
RcsB in Salmonella enterica. *PLoS Genet* 2020; **16**: e1008722.
38. Nava-Galeana J, Nunez C, Bustamante VH. Proteomic analysis reveals the global effect
of the BarA/SirA-Csr regulatory cascade in Salmonella Typhimurium grown in conditions that
favor the expression of invasion genes. *J Proteomics* 2023; **286**: 104960.
39. Gudapaty S, Suzuki K, Wang X et al. Regulatory interactions of Csr components: the RNA
binding protein CsrA activates csrB transcription in Escherichia coli. *J Bacteriol* 2001; **183**: 6017-
27.
40. Huemer M, Mairpady Shambat S, Brugger SD et al. Antibiotic resistance and persistence-
Implications for human health and treatment perspectives. *EMBO Rep* 2020; **21**: e51034.
41. Algarni S, Ricke SC, Foley SL et al. The Dynamics of the Antimicrobial Resistance
Mobilome of Salmonella enterica and Related Enteric Bacteria. *Front Microbiol* 2022; **13**: 859854.
42. Dyson ZA, Klemm EJ, Palmer S et al. Antibiotic Resistance and Typhoid. *Clin Infect Dis*
2019; **68**: S165-S70.
43. Wu S, Yang Y, Wang T et al. Effects of acid, alkaline, cold, and heat environmental stresses
on the antibiotic resistance of the Salmonella enterica serovar Typhimurium. *Food Research*
International 2021; **144**: 110359.
44. Kuang X, Hao H, Dai M et al. Serotypes and antimicrobial susceptibility of Salmonella spp.
isolated from farm animals in China. *Front Microbiol* 2015; **6**: 602.
45. Kebede A, Kemal J, Alemayehu H et al. Isolation, Identification, and Antibiotic
Susceptibility Testing of Salmonella from Slaughtered Bovines and Ovines in Addis Ababa
Abattoir Enterprise, Ethiopia: A Cross-Sectional Study. *Int J Bacteriol* 2016; **2016**: 3714785.

46.   Christou A, Aguera A, Bayona JM et al. The potential implications of reclaimed wastewater
reuse for irrigation on the agricultural environment: The knowns and unknowns of the fate of
antibiotics and antibiotic resistant bacteria and resistance genes - A review. *Water Res* 2017; **123**:
448-67.
47.   Sunuwar J, Azad RK. A machine learning framework to predict antibiotic resistance traits
and yet unknown genes underlying resistance to specific antibiotics in bacterial strains. *Brief*
Bioinform 2021; **22**.
48.   Devanga Ragupathi NK, Muthuirulandi Sethuvel DP, Shankar BA et al. Draft genome
sequence of bla(TEM-1)-mediated cephalosporin-resistant Salmonella enterica serovar Typhi
from bloodstream infection. *J Glob Antimicrob Resist* 2016; **7**: 11-2.
49.   Greninger AL, Chatterjee SS, Chan LC et al. Whole-Genome Sequencing of Methicillin-
Resistant Staphylococcus aureus Resistant to Fifth-Generation Cephalosporins Reveals
Potential Non-mecA Mechanisms of Resistance. *PLoS One* 2016; **11**: e0149541.
50.   Palzkill T. Structural and Mechanistic Basis for Extended-Spectrum Drug-Resistance
Mutations in Altering the Specificity of TEM, CTX-M, and KPC beta-lactamases. *Front Mol Biosci*
2018; **5**: 16.
51.   Sanders CC. beta-Lactamase stability and in vitro activity of oral cephalosporins against
strains possessing well-characterized mechanisms of resistance. *Antimicrob Agents Chemother*
1989; **33**: 1313-7.
52.   Ji J, Wu S, Sheng L et al. Metabolic reprogramming of the glutathione biosynthesis
modulates the resistance of Salmonella Derby to ceftriaxone. *iScience* 2023; **26**: 107263.
53.   Wu S, Ji J, Carole NVD et al. Combined metabolomics and transcriptomics analysis
reveals the mechanism of antibiotic resistance of Salmonella enterica serovar Typhimurium after
acidic stress. *Food Microbiol* 2023; **115**: 104328.
54.   Kitzenberg DA, Lee JS, Mills KB et al. Adenosine Awakens Metabolism to Enhance Growth-
Independent Killing of Tolerant and Persister Bacteria across Multiple Classes of Antibiotics.
mBio 2022; **13**: e0048022.

774