Cell loss disrupts mechanical homeostasis to drive retinal pigment epithelium ageing-like phenotype *in vitro*

Teodora Piskova[1], Aleksandra N. Kozyrina[2], Giedrė Astrauskaitė[2], Mohamed Elsafi Mabrouk[3], Sebastian Schepl[1], Stacy Lok Sze Yam[1], Ragul Ravithas[1], Wolfgang Wagner[3], Massimo Vassalli[2] and Jacopo Di Russo[1,4]

[1]Institute of Molecular and Cellular Anatomy, RWTH Aachen University, Germany
[2] Centre for the Cellular Microenvironment, James Watt School of Engineering, University of Glasgow, Glasgow, UK
[3]Institute of Stem Cell Biology, University Hospital of RWTH Aachen, Aachen, Germany
[4]DWI-Leibniz-Institute for Interactive Materials, Aachen, Germany

Keywords: Postmitotic epithelium, Ageing, Mechanical homeostasis, RPE, Phagocytosis

Correspondence:

Jacopo Di Russo

jdirusso@ukaachen.de

Abstract

Tissue homeostasis relies on mechanical feedback loops balanced by cell loss and proliferation. However, maintaining this balance becomes particularly challenging in postmitotic tissues, where alternative mechanisms replace compensatory proliferation. In the postmitotic retinal pigment epithelium (RPE), these mechanisms include significant structural adaptations over time. Yet, how these adaptations relate to epithelial mechanical homeostasis and age-associated functional decline remains poorly understood. To establish the relationship between structural changes, mechanical homeostasis and function, we developed an *in vitro* reductionistic model mimicking age-related reduction in RPE cell density. Inducing large-scale apoptosis in postmitotic stem cell-derived RPE monolayers recapitulates structural hallmarks of aged tissue, such as reduced cell height, shortened microvilli and cytoskeletal reorganisation. This new structure acquires a new mechanical equilibrium, evidenced by tissue stiffening and enhanced junctional contractility. Functionally, the monolayers display impaired vision-supporting phagocytosis of photoreceptor outer segments. Mechanistically, modulation of actin nucleators, Arp2/3 and formins, demonstrates that apicolateral monolayer deformation is critical for phagocytosis and may be compromised in aged RPE. Our findings suggest that a shift in mechanical homeostasis due to cell loss is a major driver of age-related RPE functional decline. Importantly, we show that structural remodelling in ageing alone can compromise tissue function, independent of other stressors.

1 INTRODUCTION

The structure and form of a biological entity determine its function, as described by the form-function paradigm [1,2]. This paradigm applies across various length scales, from the molecular [3,4] and sub-cellular levels [5,6] to the tissue and organ levels, where proper architecture and cellular connectivity are essential for functionality [7–9]. In recent decades, alongside form and organisation, increasing recognition as key parameters influencing function and homeostasis has also been given to the mechanical properties of tissues [10–12]. These properties are determined by the cellular organisation and environment [13–15], and their regulation within a preferred range engages mechanosensitive feedback loops that sustain physiological function [16–18], a concept referred to as mechanical homeostasis [19–21]. Despite its importance, mechanical homeostasis remains poorly understood in postmitotic tissues such as the central nervous system and the retina, which must adapt structurally to naturally occurring cell death to maintain the tissue's integrity and functional state [22–25].

An emblematic example of this adaptation is the age-related changes occurring in the retinal pigment epithelium (RPE) [26]. In the retina, the RPE is located between the light-detecting photoreceptor cells and the choroid capillaries and provides vital metabolic support to photoreceptor cells [27]. In ageing, this support is challenged by the steady rate of apoptosis in RPE [28–30], which needs to be actively compensated by the remaining cells, leading to major structural changes of the tissue. These include monolayer thinning, with the average RPE cell height reported to decrease, while cell area increases [29,31,32]. Furthermore, the RPE actin cytoskeleton remodels, leading to local fibres thickening or thinning with the formation of transcellular stress fibres [33]. Finally, ageing has been shown to be associated with changes in RPE apical microvilli, which become shorter and sparser [31,34]. In temporal correlation to these structural changes, the functional ability of the RPE to support photoreceptor cells changes towards a reduced ability to phagocyte and process photoreceptor outer segments (POS) fragments, with important implications for retinal degenerative pathologies [35,36].

While it has been proposed that the age-related changes in RPE organisation stem from a time-associated reduction in cell density, direct evidence supporting this hypothesis remains elusive. Moreover, whether such alterations drive a shift in tissue mechanical homeostasis that implies functional decline remains a critical unresolved question. Notably, we have recently shown that the strength of both cell-substrate and cell-cell adhesion forces in RPE cells directly relates to their ability to phagocytose photoreceptor outer segments (POS), with higher actomyosin contractility leading to reduced phagocytic activity [21]. Finally, because rodent models and human post-mortem tissue studies cannot reveal the precise effects of reduced cell density on monolayer structure from other ageing-related processes, such as oxidative stress or extracellular matrix (ECM) remodelling, it is necessary to establish a reductionist, human-relevant *in vitro* model.

To address this knowledge and technical gap, we present an optimised *in vitro* model
that employs large-scale apoptosis on demand to promote a reduction in cell density
in postmitotic human induced pluripotent stem cell-derived RPE (iRPE). This density-
reduced iRPE (dr-iRPE) displays structural changes such as enlarged and thinner
cells with shortened apical microvilli and cytoskeleton reorganisation, alongside
increased tissue stiffness, enhanced junctional contractility and a remodelled apical
cortex. The mechanical changes coincide with impaired POS phagocytosis, reflected
by reduced apicolateral deformation, decreased number of internalised particles and
increased fragment size. Pharmacological modulation of actin nucleation allowed us
to partially rescue function, suggesting that functional alteration may result from a
shift in force balances within the RPE that prioritises lateral structural reinforcement
instead of apical plasticity needed for POS phagocytosis.

RESULTS

Apoptosis-generated cell density reduction leads to structural remodelling in
RPE *in vitro* and phenocopies structural features of the ageing RPE

Inspired by the natural reduction in cell density in RPE *in vivo* [28,30], we aimed to
replicate cell density reduction *in vitro* by inducing large-scale apoptosis and to study
the adaptation of the epithelium to the cell loss. To model a native RPE monolayer,
iRPE was employed as a recognised and reliable model for the native tissue [37–40].
Importantly, to investigate cellular force balance, iRPE were cultured on 4 kPa soft
polyacrylamide hydrogels with basement membrane-like coating, thereby
recapitulating the physiological mechanical and biochemical cues of the native
substrate as previously characterized [21] (Sup. Fig. 1A). In these conditions, iRPE
assumes a characteristic honeycomb-like organisation, apicobasal polarisation (Sup.
Fig. 1B), and arrest of proliferation (0,01 to 0,41% within 24 h) (Fig. 1A and Sup.
Fig.1C). To facilitate on-demand apoptosis, on the 10th day of culture we transduced
the cells with an adeno-associated virus (AAV) carrying a construct designed to
promote the expression of FKBP-caspase 8 (Fig. 1B, Sup. Fig. 1D). FKBP-caspase 8
is pro-apoptotic fusion protein that can be activated on demand through AP20187-
induced dimerization, initiating the cellular suicide program [41]. Therefore, after
reaching a stable expression, on the 15th day, we incubated the monolayer with the
compound AP20187, inducing monolayer-scale apoptosis that led to the extrusion of
2% of the cells within the observation time (Fig. 1B-C, Sup. Fig. 1E and Video 1). By
the 20th day (96 hours after apoptosis completion), the iRPE monolayer showed a
reduction of cell density of approximately 8% (Fig. 1D), which was not compensated
by increased proliferation. In fact, we quantified significantly lower proliferation rates
compared to control monolayers (Fig. 1A). As a result, the obtained monolayer was
referred to as density-reduced iRPE or dr-iRPE.

Morphologically, dr-iRPE presents cells with a larger area on average (Fig. 1E, Sup.
Fig. 1F and G) without experiencing any fluidification or unjamming, as evident by

cellular area distribution and shape factor (Sup. Fig. 1H and I), recapitulating
morphological changes observed in human RPE ageing [29]. Furthermore, we noticed
that one-tenth of cells within dr-iRPE presented transcellular stress fibres (Fig. 1F),
which were also reported to occur frequently in human RPE ageing [33]. Next, we
quantified the cell height in fixed and mounted samples (Fig. 1G and H), showing that
dr-iRPE is significantly thinner than the controls (9.1 vs 11.6 μm), in agreement with
observations from the ageing RPE [31,42]. Lastly, motivated by the documented
changes of microvilli size and number in the ageing RPE *in vivo* [31,34], we evaluated
ezrin as a polarity marker and microvilli organiser [43–45]. Qualitative analyses of ezrin
staining revealed a distinct morphology of the cellular apices between dr-iRPE and
control (Fig. 1I), which can also be quantitatively captured by the reduced protein
level detected in the cells by western blot (Fig. 1J). Moreover, the examination of the
microvilli length by confocal microscopy showed approximately one-third shorter
microvilli in the dr-iRPE compared to the control (Fig. 1K). Based on the studies
performed on Ezrin$^{-/-}$ mice [43], these observed differences strongly suggest that in the
dr-iRPE conditions, the microvilli are less elaborated and more disorganised. Finally,
the quantification of the active form of ezrin (phosphorylated ezrin), which connects
the actin cortex to the membrane [46], revealed a significant decrease in the dr-iRPE
(Fig. 1L).

To exclude the influence of the viral delivery on the observed structural changes, we
included a matched viral control, which promoted the expression of GFP (GFP-iRPE)
(Sup. Fig. 2A). Compared to this control, the dr-iRPE was still showing a reduced cell
density and cell height (Sup. Fig. 2B, C and D), as well as significant differences in
both organisation and length of ezrin-positive microvilli (Sup. Fig. 2E). Next, we
addressed the observation that despite most of the cells within the monolayer
expressed the reporter after transduction, only a few of the cells ultimately underwent
apoptosis during the AP20187 treatment (Video 1). To exclude any possible effects
of the leftover recombinant protein in the remaining cells, we included an additional
'mixed' control in the analyses, consisting of an iRPE monolayer made by a mixed
population of transduced (10%) and naïve cells. This condition allowed us to target
the caspase activation only on a subpopulation of cells (Sup. Fig. 2F and G).
Interestingly, with this approach, apoptosis still caused a significant cell density and
height reduction, confirming that these features are independent of the possible
presence of inactive recombinant protein in the cells (Sup. Fig. 2H and I). Lastly, we
asked whether the changes were induced by secondary factors released in the
medium after the large-scale apoptosis induction [47]. Hence, we transferred a medium
pre-conditioned by the dr-iRPE sample during and after apoptosis onto control
monolayers (Sup. Fig. 2J). Nonetheless, the quantification still showed no effect of
the "conditioned medium" on cell morphology, density, or height (Sup. Fig. 2K, L and
M).

Altogether, we present an *in vitro* model based on cell density reduction via on-
demand apoptosis, and we demonstrate that the RPE monolayer responds to large-
scale apoptosis with structural adaptation. This includes the formation of transcellular
actin fibres, reduced cell height, larger cell area and shorter and remodelled

microvilli. Interestingly, these cellular responses appear to result from intrinsic tissue
stretching and preservation of cell volume rather than from cellular hypertrophy.
Finally, our model recapitulates key morphological features of the aged RPE *in vivo*,
suggesting a causal relationship between cell loss and the structural changes
observed in ageing.

Figure 1: Structural adaptation of iRPE to cell density reduction. A: Quantification of proliferating cells within
the monolayers on day 20 (n=4). **B:** Experimental timeline highlights key steps of the experiment, including cell
seeding on hydrogels (day 1), AAV-mediated transduction with FKBP-casp8-encoding plasmid (day 10 to 11),
apoptosis induction by the addition of the dimerising drug AP20187 (day 15 to 16), and day of read-out (day 20).
C: Quantification of extruded cells during the apoptosis induction (n=3). **D:** Quantification of cell density within the
monolayers on day 20 (n=5). **E:** Representative optical sections of iRPE and dr-iRPE stained for F-actin, which
reveals cell organisation and sizes. **F:** Zoomed-in images from the iRPE and dr-iRPE with arrows indicating the
local presence of transcellular stress fibres in the dr-iRPE condition (down). Quantification of cells with
transcellular fibres (right, n=4). **G:** Orthogonal cross-sections of F-Actin micrographs. **H:** Quantification of cell
height (n=4). **I:** Monolayer apical signal of ezrin highlighting microvilli organisation. **J:** Immunoblots showing ezrin
and GAPDH as a housekeeping protein (left) and quantification of total ezrin protein from immunoblots,

normalised to GAPDH (right). Paired values from the same experiment are connected with lines (n=4). **K:**
Representative orthogonal views of iRPE monolayers stained for ezrin showing the apical microvilli (left) and the
quantification of their average length (right, n=4). **L:** Immunoblots showing phosphorylated ezrin and GAPDH as a
housekeeping protein (left) and their quantification (right). Paired values from the same experiment are connected
with lines (n=4). Datapoints represent average values per experiment (± SEM in case of bar graphs). Statistical
significance in A, C, D, F, H and K was tested using a paired t-test, where *: p<0.05, **: p<0.01. Statistical
significance in J and L was tested using a ratio paired t-test, where *: p<0.05.

Cell density reduction compromises RPE phagocytotic activity

One of RPE's main functions *in vivo* is the daily engulfment and internalisation of
oxidised fragments of photoreceptor outer segments (POS), often referred to in the
literature as POS phagocytosis [48] or, due to the active nature of nibbling involved, as
trogocytosis [49]. Along the lines of the form-function paradigm, we asked whether the
structural adaptation observed in response to cell density reduction would alter
iRPE's ability to internalise POS. Thus, following an established assay [40,50], we
incubated control and dr-iRPE with fluorescently labelled porcine POS fragments in
the presence of bridging molecules Protein-S and MFG-E8 for 4 h (Fig. 2A). During
this time, some of the fragments bound to the apical surface of the iRPE and some
got internalised by the cells (Fig. 2A and B). After 3D segmentation, we used the
apical actin signal to differentiate between bound and internalised fragments (Sup.
Fig. 3A). Quantification revealed that the dr-iRPE exhibited significantly fewer
internalised particles, both within the field of view or when normalised by cell number
(Fig. 2C, Sup. Fig. 3B). Next, motivated by a reported trend of larger POS
phagosomes in the RPE of aged animals [51], we further quantified the volume of the
internalised photoreceptor fragments. Surprisingly, we found the average volume of
internalised fragments to be larger in dr-iRPE monolayers (Fig. 2B and D). Regarding
the bound particles, neither their number nor their average volume was found to be
significantly different (Sup. Fig. 3C). Similar differences in internalised particle
number and volume were observed in the mixed cell population subjected to cell
density reduction, with fewer internalised particles and a larger average particle
volume (Sup. Fig. 3E, G). Finally, the medium transfer from the apoptotic-induced
samples showed no significant impact on either the internalised particle count or on
the fragment volume (Sup. Fig. 3F and H).

Altogether, these data demonstrate that structurally remodelled dr-iRPE exhibit
deficient internalisation of POS fragments, internalising fewer and larger fragments.
By phenocopying the phagocytic deficits observed in ageing RPE, this model
suggests that reduced cell density alone can drive functional decline in the remaining
cells.

Figure 2: Density reduced iRPE shows altered internalisation of POS fragments. A: Schematics of POS fragment internalisation assay. Isolated and labelled porcine POS fragments are fed to iRPE together with bridging molecules at timepoint 0 and analysed four hours later. To distinguish between bound and internalised fragments, we define a cut-off height. **B:** Representative orthogonal cross-sections and single optical slices of iRPE and dr-iRPE at t = 4 h. Zoom-in of the POS signal from the inlet showcases the size of the fragments. **C:** Quantification of fragments internalised within the time of the assay. Values are shown as times thousand per field of view (n=6). **D:** Quantification of the average volume of internalised fragments. Data points represent average values per experiment (n=6). Statistical significance in C and D was tested using a ratio paired t-test, where **: $p < 0.01$. **E:** Representative orthogonal cross-sections of F-actin labelled monolayers in a standard culture condition vs during the POS internalisation assay. Images are not paired. Schematic illustration and arrows highlight the distinct cross-sectional morphology of cells, with control cell bodies bulging out above the level of cell-cell junctions. **F:** Representative orthogonal cross-sections of Myosin IIB labelled monolayers and FITC-labelled POS fragments. Myosin IIB is enriched in proximity to the POS (arrows). **G:** Representative apical view of iRPE cells in control and density reduced condition, stained for Myosin IIB during internalisation of POS-FITC. In contrast to many small phagocytic cups localised apically and laterally in iRPE (arrows), dr-iRPE shows fewer apical-only cup-like structures with more localised myosin signal (arrowheads). **H:** Representative 3D segmentation of apical F-actin surface of monolayers during internalisation assay (left) and quantification of apical actin area within the same field of view (right). Datapoints represent average values per experiment (n=3). Statistical significance was tested using a ratio paired t-test, where *: $p < 0.05$. **I:** Representative apical and orthogonal cross sections of F-actin-labelled monolayers during internalisation. Arrows indicate the presence of protruding actin-rich structures in dr-iRPE.

Cell density reduction leads to altered actomyosin plasticity and mechanical properties

POS phagocytosis relies on the active remodelling of the apical actin network, which plays a crucial role in fragment internalisation and phagosome size [40,49,50]. Actin dynamics are essential throughout the process, guiding key steps from initial binding and ensheathment to the final scission of the phagosome [48,52]. Thus, we examined the morphology of the apical actin during internalisation, on a few-cell scale in our 3D segmentation and on a larger monolayer scale. In the side view, we noticed the apical actin of control iRPE monolayers experienced a substantial deformation during the internalisation, such as bulging and enlargement of the protruding cell portion above the cell junctions and clear formation of myosin IIB-positive cups (Fig. 2E-H). In contrast, dr-iRPE did not experience much evident deformation, and phagocytotic cups were less resolved (Fig. 2F and G). Next, we noticed that cells in the dr-iRPE displayed a different apical actin topography compared to the control, having an overall smoother surface, as quantified by area per field of view (Fig. 2H). Finally, on the monolayer scale, we noticed that the dr-iRPE frequently presented large actomyosin-rich apical structures (Fig. 2G and I, Sup. Fig. 4A and B), which were absent in iRPE samples.

These observations suggest that iRPE cells in the density-reduced condition exhibit altered ability in remodelling their actomyosin cytoskeleton when internalising POS fragments. Consequently, we wondered whether the differences stemmed from distinct patterns of actin-cytoskeleton-related signalling. Hence, we examined the myocardin-related transcription factor A (MRTFA), which is known to regulate actin-associated signalling [53]. We observed a short-term translocation of MRTFA into the nucleus at day 17, shortly after the large-scale cell loss, which went back to levels comparable to the control at day 20, or the day of the read-out (Fig. 3A and B). This demonstrates that the cell density reduction leads to a transient transcriptional adaptation, possibly conveying the observed structural remodelling of dr-iRPE. We then examined the mRNA expression of actin-associated proteins after bulk RNA sequencing (Fig. 3C, Sup. Fig. 5A). Examining more precisely the expression of these proteins showed an array of significantly differentially expressed actin-related genes in the dr-iRPE. Upregulated genes in dr-iRPE contained genes related to actin branching, polymerisation, depolymerisation, and crosslinking (*ARPC2, PFN1, FHOD3, INF2, CFL2, ACTN1, FSCN1, FLNA*). Downregulated genes included ezrin and the formins *DIAPH1, DIAPH2, DAAM1* and *DAAM2* (Fig. 3D and E). Next, to further assess the parallel between our model and ageing RPE, we compared global gene expression changes in iRPE and dr-iRPE with those reported in ageing RPE from two independent studies [54,55] (Sup. Fig. 5 B-D). Considering the significant differences from our reductionist approach and *in vivo* human data, as well as the differences between the results of the studies, it is worth noting that our model may mimic further aspects of RPE ageing besides the actomyosin remodelling discussed in this work.

Altogether, these insights point to a distinct signature of actin-associated proteins in
dr-iRPE that implies altered actomyosin network organisation, consistent with the
observed changes in subapical F-actin topography. Such reorganisation may prevent
the formation of the prominent apical deformations or bulges required for efficient
phagocytosis, instead promoting the emergence of localised actin-rich structures.

Figure 3: A different fingerprint of actin-associated proteins in density reduced iRPE. A: Quantification of
nuclear to cytoplasmic MRTFA fluorescence intensities and representative micrographs of MRTFA staining one
day after apoptosis induction (day 17). **B:** Quantification of MRTFA fluorescence intensities and representative
micrographs of MRTFA staining on the day of readout (day 20). Datapoints in A and B represent average values
per experiment ± SEM originating from n=3 independent experiments. Statistical significance was tested using a
ratio paired t-test, where *: p<0.05. **C:** Hierarchically clustered heatmap of RNA transcript counts of actin-related
proteins from n=4 independent experiments of iRPE vs dr-iRPE samples. **D:** Graphs showing the Log2Fold
change in RNA transcript expression of branched (left) and linear (right) actin nucleator proteins for dr-iRPE vs.
iRPE. **E:** Graphs showing the Log2Fold change in RNA transcript expression of actin assembly/disassembly
regulators (left), actin membrane linkers (middle) and crosslinkers and bundlers (right) for dr-iRPE vs. iRPE.
Differentially expressed genes in D and E are marked with *: $p_{adj}<0.1$, **: $p_{adj} <0.01$, ***: $p_{adj} <0.001$, and ****: p_{adj}
<0.0001 as a result of the use of negative binomial generalised linear models and Wald test.

As we previously showed that RPE mechanical equilibrium influences the epithelium
phagocytotic activity [50], we sought to determine whether the observed actin network
remodelling would alter the mechanical properties of dr-iRPE. To characterise the
viscoelastic properties of the monolayer as a multicellular system, we conducted
nanoindentation using a spherical tip with a diameter of 20 μm, allowing to probe a
contact area of approximately 10 μm in diameter (Fig. 4A, Sup. Fig. 6A, Video 2).
This approach allowed us to measure the Effective Young's modulus of the
uppermost 2.5 μm of the monolayer, which approximately corresponds to the apical-
most 9% of the cell height (average height of live iRPE is 27.8 ± 0.6 μm). The

Effective Young's modulus of dr-iRPE was roughly one-third higher (694 Pa vs 513
Pa), pointing to a stiffer tissue (Fig. 4B). Next, using the same nanoindentation setup,
we quantified both storage and loss moduli at different probing frequencies to
characterise the monolayer viscoelasticity [56]. With this approach, we found a
significant increase in the storage modulus of the dr-iRPE while the loss modulus
remained unchanged (Fig. 4C, Sup. Fig. 6B). Similar differences were observed
when dr-iRPE was compared to the virus control or when the condition with mixed
cell population was analysed without or with cell extrusion (Sup. Fig 6D and E). In the
latter case, the differences in stiffness and storage modulus were smaller and only
apparent at the lower probed frequencies (1 and 2 Hz). Lastly, medium transfer
controls seemed to experience softening rather than stiffening, excluding an effect of
apoptotic debris on the observed stiffening in dr-iRPE (Sup. Fig. 6F). Collectively,
these measurements revealed a reinforced cellular interconnectivity that results in a
less deformable monolayer, i.e. stiffer, as well as more elastic, suggesting an
actomyosin-based reinforcement [57].

As mechanical properties and contractility are often linked [58,59], we examined the
levels of phosphorylated myosin light chain 2 (pMLC2) to obtain a general idea about
actomyosin contractility [60]. Immunoblotting revealed general increased levels of
pMLC2 in dr-iRPE (Sup. Fig. 6C), pointing to overall higher actomyosin contractility,
especially localised at the cell-cell junctions as shown by immunofluorescent staining
(Fig. 4D and D^I). This observation was further supported by enrichment of vinculin at
the cell-cell and at the tricellular junctions (Fig. 4E and E^I), pointing to higher
contractile force at cell-cell junctions [61].

Figure 4: Density reduced iRPE presents a new biomechanical state. A: Image illustrating the probing window (10 x 2.5 µm) during nanoindentation on an orthogonal cross-section of F-Actin-labelled live iRPE. **B:** Average Effective Young's Modulus measured by nanoindentation scans in iRPE and dr-iRPE (n=4). **C:** Average storage modulus at different oscillation frequencies (n=4). **D:** Representative images of pMLC2-labelled iRPE monolayers with grey values represented in the colour bar. **D^I:** Quantification of junctional pMLC2 signal (n=4). **E:** Representative images of vinculin-labelled iRPE monolayers with annotated regions enlarged. **E^I:** Quantification of junctional fluorescence signal of vinculin normalised to cytoplasmic fluorescence signal (n=3). **F:** Image illustrating the geometry of low-resolution Brillouin z-scans on an orthogonal cross-section of F-Actin-labelled live iRPE. **G:** Representative line profiles of the z scans showing the Brillouin shift values starting from the hydrogel, across the iRPE monolayers and reaching into the cell medium. Averages of z-scans are represented with thicker lines. Highlighted in light lilac and salmon are the approximate basal and apical cell regions, respectively. **H:**

Nanoindentation enabled us to characterise the biomechanical properties of the RPE
from the apical side. Our indentation setup was designed to probe multicellular length
scale deformation, where the indentation of the 20 μm tip at the selected depth is
counteracted by intercellular junctions [62] (Fig. 4A, Sup. Fig. 6A, Video 2). To
characterise subcellular mechanical properties of the RPE monolayer non-invasively,
i.e., independently from intercellular adhesion, we applied Brillouin micro-
spectroscopy. Here, the detection of photons, inelastically scattered by thermally-
induced acoustic phonons, enables the probing of viscoelastic properties within the
sample at microscale resolution [63]. Using Brillouin microscopy, we first performed
low-resolution z-scans through iRPE and dr-iRPE monolayers, including the
surrounding hydrogel and medium (Fig. 4F). These line profiles revealed the
presence of a peak in the Brillouin shift, corresponding to the cellular basal side,
which gradually decreased towards the apical cell region (Fig. 4G). The quantification
of fitted peak values of these line profiles revealed significantly lower Brillouin shift in
dr-iRPE compared to iRPE (Fig. 4H). Moreover, based on the peaks shown in Figure
4G and confirmed by live-cell brightfield imaging, we identified the apical and basal z-
levels for conducting 2D high resolution raster scans (Fig. 4I). The 2D raster scans
(50 x 50 μm) revealed mechanical compartmentalisation into areas with different
Brillouin shifts, which may correspond to cellular and intercellular regions (Fig. 4J).
Notably, this compartmentalisation was much more evident at the basal sides. The
quantification of the mean Brillouin shifts revealed a difference between the
conditions, with dr-iRPE cells showing lower average values at the apical (6.30 vs
6.36 GHz) and basal (6.35 vs 6.37 GHz) sides of the cells (Fig. 4K). Interestingly,
when comparing apical and basal shifts within the same conditions, dr-iRPE cells
showed significantly different apical shifts compared to basal, which was not the case
in iRPE (Fig. 4L). As the Brillouin shift reflects cytoskeletal organisation and water
content[63–65], this supports the idea that the apical sides of the cells undergo the most
pronounced reorganisation in dr-iRPE. This is consistent with microvilli and apical
actin remodelling observed in dr-iRPE, i.e., transcellular stress fibres and F-actin
surface topography.

As the intercellular junctions showed an increased pMLC2, thereby an increased
contractility and stiffness, we sought to isolate the shifts of "cell edges" from those of
"cell bodies". As this distinction was not clear in the apical scans, we performed the
analysis on the basal sides, where the distribution of shift values revealed two distinct
modes (Sup. Fig. 7A and B). Quantification based on segmented Brillouin shift plots
shows that, regardless of the experimental condition, the interstitial regions of the
monolayer ("cell edges") consistently exhibit lower shifts than the cell cores ("cell
bodies"). This also confirms a trend toward reduced core shifts in dr-iRPE

independently of the interstitium (Sup. Fig. 7C and D). Notably, the shift quantified at
the interstitium was unexpectedly lower than the cell bodies, likely reflecting the
absence of well-defined intercellular junctions at the basal side, and instead, the
presence of loosely interdigitated basal membranes, as seen in F-actin staining (Sup.
Fig.7E).

Altogether, biomechanical characterisation at the two different length scales showed
that, in addition to the structural and functional adaptation, the reduction of cell
density causes RPE to adopt a different mechanical state. This mechanical state is
associated with increased contractility at the apical intercellular junctions, which
confers a stiffer and more elastic tissue, as evident by the nanoindentation. Brillouin
shift analyses at the subcellular level also confirmed distinct mechanical states
between the two conditions, most pronounced at the apical surface. Taken together,
this apical prevalence of the mechanical changes may result from the intimate link
between the epithelial junctional complex and the organisation of the apical domains,
which in the case of the RPE may create a mechanical feedback loop between tissue
integrity and function.

Modulating actin remodelling differentially impacts POS phagocytosis

Motivated by our observations on cortical actin topography, mechanics, and the
differential expression of actin-related regulators, we aimed to directly investigate the
cause-effect relationship between POS internalisation and actin remodelling. To this
end, we targeted actin nucleation, a key process influencing cortical thickness and
membrane tension [66]. For the experimental design, we considered the upregulation
of Arp2/3 subunit mRNA expression and the downregulation of several formins in dr-
iRPE (Fig. 3D), along with the distinct actin structures observed during internalisation.
Dr-iRPE exhibited a larger, flatter actin surface topography (Fig. 2H), suggesting an
abundance of heavily branched actin, similar to that found in lamellipodia. In contrast,
control samples displayed more numerous, smaller, and protruding actin structures
(Fig. 2H), resembling filopodia, which are characterised by linear actin cores and
highly dependent on formin activity [67]. Based on these findings, we inhibited Arp2/3
activity using the compound CK-666[68] and inhibited formin activity using the
compound SMIFH2 [69].

Formin inhibition in iRPE did not affect the cell height but significantly stiffened the
monolayer on a tissue level (Sup. Fig. 8A, B). In dr-iRPE, the inhibition did not
produce a change in cell height, but led to a softer monolayer (Sup. Fig. 8C, D).
When tested for the POS fragments internalisation, the quantification showed a
highly variable effect on the number of internalised particles, which nevertheless
showed an overall significant reduction in the number of both internalised and bound
POS fragments (Fig. 5A and B; Sup. Fig. 8E and F). In contrast, no difference was
detected in SMIFH2-treated dr-iRPE (Sup. Fig. 8G). Next, we quantified the
fragments' size, which was significantly affected by SMIFH2 only in iRPE, resulting in
larger fragments on average (Fig. 5A and B, Sup. Fig. 8H). Since SMIFH2 affected
internalisation of POS only in iRPE, we further analysed 3D-segmented actin and

monolayer deformation during internalisation. SMIFH2-treated iRPE monolayers
displayed a smoother apical actin topography; however, this was not captured by the
quantification of the total apical actin area (Fig. 5C; Sup. Fig. 8I). Furthermore,
SMIFH2-treated iRPE monolayers also bulged out upwards similarly to untreated
controls, but showed denser F-actin signal at the junction, suggesting a lateral
reinforcement comparable to dr-iRPE (Fig. 5D). In addition, the bulging cells showed
a more irregular apical morphology (Fig. 5D).

Although SMIFH2 is widely accepted as a reliable formin inhibitor, it has been
reported to have a concentration-dependent off-target effect on Myosin II activity [70].
To evaluate whether the effects observed with 20 µM SMIFH2 were not due to partial
Myosin II inhibition, we conducted SMIFH2 treatments across a range of
concentrations (5–100 µM). We then assessed iRPE and dr-iRPE monolayers for
changes in stiffness and phagocytic activity (Sup. Fig. 8J, L, N, P, R and T). As a
positive control for Myosin II inhibition, we used blebbistatin (Sup. Fig. 8K, M, O, Q, S
and U). In iRPE, stiffness quantification confirmed a significant monolayer stiffening
with 20 µM SMIFH2, whereas blebbistatin treatment induced an expected softening
as cellular contractility is reduced (Sup. Fig. 8J and K). Differently, dr-iRPE treated
with SMIFH2 displayed an overall softening effect, most prominent at 100 µM,
consistent with the trend observed following blebbistatin treatment (Sup. Fig. 8L and
M). Finally, analyses of the POS internalisation process revealed that both SMIFH2
at all concentrations and blebbistatin reduced the number of internalised POS
particles in iRPE. Still, only the low concentrations of SMIFH2 (5 and 20 µM) led to
increased average phagosome size, while blebbistatin did not affect fragment size
(Sup. Fig. 8N-Q).

In dr-iRPE, SMIFH2 significantly reduced the number of internalised particles only at
5 µM with no effect at higher concentrations, whereas blebbistatin reduced it (Sup.
Fig. 8R, S). Average phagosome size was clearly reduced by SMIFH2 treatment in
dr-iRPE, in opposite direction to iRPE (Sup. Fig. 8T), while blebbistatin did not affect
the phagosome size (Sup. Fig. 8 U). Overall, while we cannot fully exclude the
possibility of minor off-target inhibition of Myosin II by 20 µM SMIFH2, we are
confident that Myosin II inhibition does not drive our results. This confidence stems
from the clear differences in trends for iRPE and dr-iRPE regarding both stiffness and
internalised POS number and volume, as well as the contrasting effects seen with
blebbistatin treatments.

The inhibition of Arp2/3 using CK-666 in iRPE did not change the cell height or
stiffness, whereas in dr-iRPE, the treatment led to a slightly increased cell height
(Sup. Fig. 9A and B), but no significant changes in monolayer stiffness (Sup. Fig. 9C
and D). During phagocytosis, no effect was detected when iRPE were treated, but in
dr-iRPE, the treatment led to an increased number of internalised particles (Fig. 5E
and F; Sup. Fig. 9E and G), while the average internalised particle size and the
number of bound particles remained comparable (Fig. 5E, G, Sup. Fig. 9F and H). As
the phagocytic process was affected only in dr-iRPE, we proceeded to analyse the
morphology of the 3D-segmented cortical actin during internalisation. The analyses

showed the occasional presence of thinner and longer ruffles (Fig. 5G), which
resulted in a significant increase in the quantified apical actin area (Sup. Fig. 9I).
Furthermore, orthogonally, the treated dr-iRPE showed some degree of 'bulging'
during phagocytosis, which was more pronounced than in the untreated dr-iRPE (Fig.
5H).

Overall, these results demonstrate that modulating actin cytoskeleton remodelling by
interfering with actin nucleation effectively altered cortical actin responsiveness
during internalisation. These manipulations were sufficient to replicate or rescue
differences in photoreceptor internalisation observed in dr-iRPE. Particularly, Arp2/3
activity appears to be related to cell 'bulging', modulating apical actin area and cell
height, which impacts the number of internalised particles, while formins activity
modulates tissue-level stiffness, affecting both the number and size of the
internalised particles.

These insights strongly indicate that dr-iRPE cells establish a new mechanical
equilibrium, in which the actin cytoskeleton acts as a critical link between tissue
structure, mechanics, and the execution of phagocytosis. The altered organisation of
the actomyosin network enhances contractility at cell-cell junctions, and
simultaneously reduces the capacity for apical remodelling required for the formation
of phagosomes. We therefore conclude that, in response to reduced cell density, the
RPE monolayer reroutes actin-related processes toward reinforcing tissue integrity.
This shift diminishes the plasticity of the actin-myosin network, which is essential for
efficient POS phagocytosis (Fig. 6).

Figure 5: Actin modulation via nucleator inhibitors affects internalisation of photoreceptor outer segments. A: Number of internalised fragments (left) and their average volume (right) for iRPE vs iRPE treated with 20 µM SMIFH2 (n=8). **B:** Representative orthogonal cross-sections and single optical slices of iRPE and SMIFH2-treated iRPE at t = 4 h. Zoom-in of the POS signal from the inlet showcases the size of the fragments. **C:** Representative 3D segmented apical F-actin surface of monolayers during POS internalisation assay for iRPE vs SMIFH2-treated iRPE. **D:** Representative orthogonal cross-sections of F-actin labelled monolayers during POS internalisation assay of untreated and SMIFH2-treated iRPE. Arrows indicate the junctional remodelling to induce cell bulging, arrowheads indicate the junctional reinforcement, and stars highlight apical irregularities. **E:** Number of internalised fragments (left) and their average volume (right) for dr-iRPE vs dr-iRPE treated with 500 nM CK-666 (n=4). **F:** Representative orthogonal cross-sections and single optical slices of dr-iRPE and CK666-treated dr-iRPE at t = 4 h. Zoom-in of the POS signal from the inlet showcases the size of the fragments **G:** 3D segmented apical F-actin surface of monolayers during POS internalisation assay for untreated vs CK666-treated dr-iRPE. **H:** Representative orthogonal cross-sections of F-actin labelled monolayers during photoreceptor fragment internalisation assay of untreated and CK666-treated dr-iRPE. Arrows indicate the junctional remodelling to

477 induce cell bulging, while arrowheads indicate the junctional reinforcement. Datapoints represent average values
478 per experiment. Statistical significance was tested using a ratio paired t-test, where *: p<0.05.

DISCUSSION

In this work, we present an *in vitro* model that recapitulates structural and functional
features of the aged RPE. The model is inspired by the natural loss of cells through
apoptosis within the native RPE during ageing. Comparing dr-iRPE with iRPE, we
demonstrate significant changes in biomechanical features that evidence a shift in
RPE mechanical equilibrium related to the structural remodelling to accommodate the
reduced cell density. Most importantly, this shift has implications for the function of
POS internalisation, potentially leading to impaired tissue performance. We identify
remodelling of the actin cytoskeleton as a key factor mediating both the structural
tissue adaptation to cell loss and the observed functional differences, suggesting that
tissue preservation may be prioritised at the expense of functional trade-offs in
ageing RPE. Ultimately, this structural adaptation disrupts the actomyosin
machinery's ability to effectively remodel, thereby compromising efficient phagosome
formation and POS phagocytosis (Fig. 6).

Figure 6: Cell density reduction leads to a shift in RPE mechanical homeostasis. The density reduction in
iRPE leads to structural remodelling and biomechanical alterations that recapitulate the tissue architecture of the
aged native RPE, suggesting a shift in mechanical homeostasis. These changes correlate with a reduced number
of internalised POS fragments, which are larger in size, highlighting functional decline. Actin cytoskeleton
remodelling mediates structural reinforcement and functional alteration, suggesting a trade-off where tissue
structural preservation compromises the apico-lateral plasticity essential for phagosome formation and POS
phagocytosis.

Ageing is a multifactorial phenomenon that impairs the capacity for dynamic
homeostatic adjustments [71]. Only recently has the scientific community
acknowledged altered mechanical properties as a hallmark of ageing [72], raising
interest in mechanobiological studies in ageing. With the aim of studying the effect of
reduced cell density on the structural and functional adaptation of RPE, our model
highlights the importance of tissue organisation and mechanical properties in ageing.

By using epigenetically 'young' [73] cells derived from human induced pluripotent stem
cells (hiPSCs), which have undergone rejuvenation-associated reprogramming, we
reduced the complexity down to examining how tissue architecture alone impacts
mechanical homeostasis and functional capacity, putting aside more complex
aspects of biochemical signalling such as oxidative stress. In this system, we were
able to recapitulate structural aspects identical to reported remodelling of the aged
RPE, such as cell height reduction [31], shortening of the apical microvilli [74] and
presence of actin stress fibers [33] by only reducing cell density. This remodelling is
possible due to the nearly postmitotic nature of iRPE cells, which cannot use
compensatory proliferation as a homeostatic mechanism, but are believed to rely on
compensatory cellular hypertrophy [75,76]. We noted that cellular 'hypertrophy' in our
model is not merely cell volume enlargement, but 'tissue stretching from within'. We
observed cell area enlargement paired with cell height reduction, pointing to
individual cell volume conservation, which may enrich the current notion of
compensatory hypertrophy in postmitotic epithelia [76]. Individual cell area
enlargement could also explain the observed microvilli shortening, as microvilli
provide membrane reservoirs capable of unfolding to facilitate cell shape changes or
to buffer mechanical stress [77,78]. In light of the cell density changes in RPE
pathologies such as proliferative vitreoretinopathy and age-related macular
degeneration[79,80], future studies should explore the relationship between the degree
of cell loss and the structural, functional or cell-cycle-related adaptations.

The changes induced by cell density reduction are also paired with a shift in the
range of mechanical equilibrium. This shift is evidenced by multicellular length-scale
stiffening and increased elasticity, with softening at the apical side at the subcellular
length scale. Tissue stiffening, in particular increased elasticity, has been reported as
a response of epithelial cells to stretch [77,81], at least in the short term. In our system,
the mechanical changes are observed in the long term, four days after the apoptosis-
mediated deformation. While these data reveal how RPE cells *in vitro* compensate for
large-scale cell loss, the situation *in vivo* is undoubtedly more complex to decipher.
Multiple factors influence native tissue during ageing, making a direct comparison
between our reductionistic model and physiological RPE ageing premature.
Nevertheless, alterations in actin cytoskeletal organisation have been reported in
aged RPE [33], supporting the notion that changes in mechanical homeostasis,
including tissue stiffening, also occur *in vivo*. Such adaptation may arise from
reduced cell density and the resulting need to reinforce tissue integrity. This is
particularly evident in the formation of transcellular actin stress fibres, observed both
in our model and in aged native tissue [33]. In fact, the occurrence of transcellular actin
stress has been reported as a sign of junctional remodelling [82] and limiting cell size or
elongation during morphogenesis [83]. Moreover, the strengthening of cell-cell
junctions is a known mechanism to support cells in resisting forces [84]. Interestingly,
the non-invasive probing technique of Brillouin microscopy revealed a mismatch
between tissue reinforcement and apical cortex remodelling. Although the results of
indentation measurements cannot be directly compared to light-based measurements
[63,64,85], the observed lower Brillouin shift may be explained by significant changes in

the apical cortex composition and in the apical microvilli, which are rich in actin and
tubulin filaments. These filaments enhance the Brillouin shift due to their structural
stiffness compared to the underlying cell [65]. A more elaborate apical cortex may also
possess a different level of hydration, which has recently been shown to be a crucial
factor affecting Brillouin, but not indentation-based measurements[64]. Furthermore,
differential remodelling of the apical and lateral cortical actin networks is likely, as is
known to happen during morphogenesis [86]. Notably, the quantified reduction of
phosphorylated ezrin indicates decreased cortical tension in dr-iRPE, consistent with
both our Brillouin measurements and existing literature [87]. Finally, we were surprised
to find that many structural and mechanical adaptations are associated mainly with
the apicolateral domains. In fact, a direct interplay between the structural and
mechanical remodelling of cell–cell and cell–ECM adhesions has been widely
reported [88–93]. Except for epithelial buckling during morphogenesis [90] or pathology [94],
this homeostatic balance is generally linear: reinforcement and increased tension at
intercellular junctions are typically accompanied by enhanced cell–ECM adhesion
and increased traction forces [89]. Nevertheless, in our case, no apparent
reinforcement has been detected at the basal side of the monolayer. This may be
attributed to our study conditions, where iRPE cells are cultured on polyacrylamide
hydrogels, which provide strictly controlled physical and biochemical cues with a
limited possibility for cell-mediated remodelling. Future studies need to elucidate the
contribution of ECM cues and cell-mediated ECM remodelling as part of the puzzle of
age-related structural remodelling of the outer retina.

Our work suggests that age-associated structural phenotype goes hand in hand with
actomyosin organisation and remodelling capacity, which affects the RPE function of
POS fragments internalisation. Phenotypically, we observe an overall higher
contractility and shifts in the expression of nucleators and other actin-related proteins.
The apical domain of RPE cells requires high remodelling capacity due to its
essentiality for the internalisation function [49]. In this study, we observed an
apicobasal cell remodelling with bulging and lateral relocalisation of cell-cell junctions
during the POS internalisation assay, which has not been reported before. Extensive
studies on phagocytosis in macrophages have demonstrated that efficient phagocytic
cup formation requires coordinated mechanical remodelling of the plasma membrane
and cortical actin [95–97]. In contrast to isolated cells, RPE cells cannot remodel their
apical membrane without simultaneously remodelling the interaction with their
neighbours, due to their tight epithelial organisation. Therefore, unlike single-cell
phagocytosis, phagocytosis in the RPE can be envisioned as a more collective
phenomenon, in which neighbouring cells coordinate lateral remodelling to support
apical bulging and phagosome formation. The observed phenomena may arise from
the synergy of conditions in our culture system, where we observe mature cells of
nearly three weeks cultured on soft substrates, as opposed to the majority of other
studies, which are being conducted on stiff cell culture plastics or glass substrates. In
our model of cell density reduction, we observed a lack of apical bulging and
prominent actin-rich protrusions during internalisation. The absence of bulging may
stem from the difficulty in disassembling junction-associated structures, which are

more reinforced and stable, or from their connection to highly crosslinked or bundled
actin structures. A logical explanation is that increased contractility at the cell-cell
junction, necessary to maintain tissue integrity, may redirect actin dynamics from the
apical side. This could limit the cytoskeletal and membrane flexibility required for
efficient apical processes, including phagosome formation and the development of
sufficient tension to engulf POS fragments. The observed formation of large actin-rich
protrusions exclusively in dr-iRPE is reminiscent of "arms" structures described in the
early embryo [98] and may reflect an altered remodelling capacity, or plasticity, of the
membrane-cortex complex. The formation of these protrusions is driven by Arp2/3
activity [98], which is highly versatile both in maintaining junctional integrity [99],
regulating cortical thickness and tension [66], as well as playing a major role in the
internalisation process [48,52]. Its activity may be diverted towards a new housekeeping
maintenance of cortical and junctional actin structure, as opposed to on-demand
availability for internalisation. The role of formins in RPE function remains unexplored
to date, but our data suggest that they both influence the efficiency of POS
phagocytosis and phagocytic cup size when inhibited. This effect may occur through
alterations in membrane curvature or surface ruffling, impacting the contact area for
internalisation. Notably, inhibition of Myosin II with blebbistatin did not affect the size
of internalised POS fragments, but only their number, thereby excluding a role for
contractility in regulating phagosome size.

Our findings demonstrate that structural adaptation to cell loss reprogrammes actin
organisation, stabilising filaments and redirecting actin towards lateral junctions. This
reinforcement limits cytoskeletal responsiveness at the apical surface, constraining
the dynamic remodelling required for POS phagocytosis. Thus, the monolayer adopts
a new mechanical homeostasis that prioritises structural integrity over functional
plasticity. We propose that this stability–flexibility trade-off is a hallmark of postmitotic
tissues, offering protection at the expense of specialised functions. This
mechanobiological shift provides insight into ageing-related decline in tissues such as
the retina and brain [24,25,100–103], and may extend to proliferative epithelia. Future work
should determine how density and architecture-driven changes in mechanics
contribute to progressive functional loss across tissues.

METHODS

Preparation of cell hydrogel substrates
The surface of 20 mm glass-bottom dishes (Cellvis, D35-20-0-N) or 6-well glass
bottom plates (Cellvis, P06-1.5H-N) was activated to allow hydrogel attachment by
etching with 0.1 M NaOH (Merck, 106498), followed by reaction with 4% (3-
Aminopropyl)triethoxysilane (Sigma Aldrich, 440140) in isopropanol (VWR, 20842)
and final incubation with 1% Glutaraldehyde (Merck, 354400) in double distilled water
(ddH2O) (B. Braun, 0082479E).

Polyacrylamide hydrogels of 4 kPa stiffness were prepared from an 5%/ 0.1%
Acrylamide/Bis-Acrylamide (Bio-Rad 1610140, Bio-Rad 1610142) mixture in ddH2O

by radical polymerisation upon addition of the initiators ammonium persulfate (Sigma
Aldrich, 248614) at 0.5% (vol/vol) and N'-Tetramethylethylenediamine (Sigma
Aldrich, 411019) at 0.05% (vol/vol), as previously described [104].

Hydrogels were trimmed to size of 4 mm before functionalisation with Laminin-511
(Biolamina, LN511) and Laminin-332 (Biolamina, LN332) at a concentration of
15 µg/ml respectively. Briefly, a functionalisation solution was prepared from ddH2O,
0.5 M HEPES (Carl Roth, 9105) pH 6, 100% ethanol (Sigma Aldrich, 32205) and
0.2% (vol/vol) Bis-Acrylamide. The solution was degassed for 20 min and was
completed by adding 0.2% Di(trimethylopropane)tetraacrylate (Sigma Aldrich,
408360), 3% (wt/vol) Hydroxy-4'-(2-hydroxyethoxy)-2-methylpropiophenone (Sigma,
41089 and 0.03% (wt/vol) Acrylic acid N-hydroxysuccinimide ester (Sigma, A8060) as
previously described [105]. Hydrogels were dehydrated with 70% ethanol for 5 min,
incubated under UV for 10 min with the functionalisation solution and washed with
25 mM HEPES and PBS for a total of 20 min before adding the protein solution. The
gels were incubated with the protein solution overnight at 4ºC.

Culture of human induced pluripotent stem-cell derived RPE
HiPSCs-derived RPE cells (Fujifilm Cellular Dynamics, R1102) were seeded in 24-
wellplates coated with 2.5 µg/ml vitronectin XF (StemCell Technologies, 07180).
Cells were cultured in MEM-alpha with nucleosides (Thermo Fisher, 12571063),
supplemented with 1x Glutamax (Thermo Fisher, 35050061), 5% (vol/vol) Knockout
Serum replacement (Gibco, 10828028), 1% (vol/vol) N2-Supplement (Gibco,
17502048), 55 nM Hydrocortisone (Sigma Aldrich, H6009), 250 µg/ml Taurine
(Sigma Aldrich, T8691), 14 pg/ml Triiodo-L-thyronine (Sigma Aldrich, T5516) and
25 µg/ml gentamycin (Gibco, 15750060) following the manufacturer's instructions.
The medium was exchanged every other day. For seeding on hydrogels, cells were
detached from the 24-wellplate by 30 min incubation with 5 mM PBS/EDTA (Sigma
Aldrich, E6511), followed by 5 min incubation with TrypLE Express (Gibco,126050).
Cells were seeded on the UV-sterilized functionalised hydrogels at 5000-6000
cells/mm^2. The medium of cells on hydrogels was exchanged halfway twice a week.

On-demand apoptosis for generation of density-reduced iRPE (dr-iRPE)
Apoptosis was induced by expression and on-demand dimerisation of recombinant
FKBP-caspase8 (FKBP-casp8) fusion protein [41]. The FKBP-casp8 containing
plasmid was delivered packaged in AAV5 (VectorBuilder, VB230127-1043eaq).
Briefly, on day 10 after seeding on the hydrogel, cells were transduced with the virus
at 7×10^6 infectious units per cell for 24 h. On day 15, 120 h after the transduction,
FKBP-casp8 was dimerised by addition of 100 nM AP20187 in medium (Sigma
Aldrich, SML2838) for 24 h. Paired control monolayers were also subjected to the
AP20187 treatment. The read-out was performed on day 20, or 96 h after apoptosis
treatment finalisation.

Virus, mixed and medium-transfer controls

To exclude effects of the AAV as a delivery system, we used an AAV5-packaged construct that promotes the expression of green fluorescent protein (GFP) (VectorBuilder, VB10000-9287ffw) as a control. Cells were transduced and treated the same way as described for FKBP-casp8 expressing cells, including same timing of viral transduction, AP20187 treatment and read-out.

To obtain a mixed population of transfected and naïve cells, two days prior to seeding on hydrogels, we transduced hiPSCs-derived RPE in a 24-well plate at 7×10^6 infectious units per cell for 24 h with the FKBP-casp8 vector (VectorBuilder, VB230127-1043eaq). The virus solution was replaced with fresh medium after 24 h. After another 24 h, we seeded a 10%/90% mixture of the transduced hiPSCs-derived RPE cells and naïve hiPSCs-derived RPE on the hydrogels as described above. On day 15, corresponding to the same apoptosis timing of dr-iRPE generation, apoptosis was induced by addition of 100 nM AP20187 (Sigma Aldrich, SML2838) for 24 h. The read-out was performed at 96 h after apoptosis treatment finalisation.

To exclude possible effects of the release of apoptotic debris during the apoptosis, 16 h after AP20187 addition, the medium from dr-iRPE samples was transferred onto control conditions. The medium was then replaced after apoptosis treatment finalisation. 24 h after apoptosis finalisation, the medium from dr-iRPE samples was again transferred to control samples and the samples were maintained in this conditioned medium until read-out.

EdU incorporation-based proliferation assay

Proliferation was assessed by evaluating the incorporation of the base 5-Ethinyl-2'-Desoxyuridin (EdU) with the Click-iT EdU kit (Thermo Fisher, C10337). Cells were incubated with 1:1 mixture of 20 µg/ml EdU and cell culture supernatant for 24 h. The subsequent EdU development reaction was performed following the manufacturer's instructions. Monolayers were imaged completely as overviews of stitched tiles at inverted epi-fluorescent microscope Zeiss Axio Observer 7. Positive cells were identified by using the particle analysis tool in Fiji and their number was normalised to the total number of cells, as detected by DNA labelling by DAPI (1 µg/ml, Invitrogen D1306).

Photoreceptor fragment phagocytosis assay

Photoreceptor outer segment fragments were isolated from porcine eyes obtained from a local slaughterhouse and labelled with FITC (Thermo Fisher, F1906) as previously described [106]. For the assay, the vial was thawed and washed with medium at 2300 g for 5 min. The pellet was resuspended in complete hiPSCs-derived RPE medium together with the bridging molecules human Protein-S (Coachrom, pp012A) at 2 µg/ml and human MGF-8 (R&D Systems, 2767-MF-050) at 2.4 µg/ml. An approximate ratio of ten photoreceptor outer segment fragments per RPE cell seeded was used. Cells were incubated for 4 h, washed four times with DPBS with Ca^{2+} and Mg^{2+} (Gibco, 14040117) and fixed with 2% paraformaldehyde

(Merck, 818715) for 8 min at room temperature (RT). Samples were then
permeabilised with 0.3% Triton-X-100 (Sigma Aldrich, T8787) for 2 min and labelled
for actin with Phalloidin AF568 (Molecular probes, A12380). Three z-stacks per
monolayer were recorded at Zeiss LSM710 Confocal Microscope with ZEN Black
software with 40x water immersion objective (ZEISS, 420862-9970-000) in Airyscan
super resolution mode. The subsequent analysis was performed in Imaris v10.2.0
(Oxford Instruments).

Nanoindentation and dynamic mechanical analysis

Nanoindentation was performed on a Chiaro Nanoindentor (Optics 11 Life) mounted
on an inverted epifluorescence microscope Zeiss Axio Observer 7 with environmental
control. Measurements were performed on living cells in displacement control mode
using a spherical probe of 20 µm diameter and 0.025 N/m stiffness. Three matrix
scans of 16 indentations were recorded per monolayer, indenting 6 µm into the cells.
Dynamic mechanical measurements were performed at 1, 2, 4 and 10 Hz to allow for
viscoelastic characterisation. For the analysis, indentation portions between the
contact point and 2.5 µm were fitted using the Hertz model in the Optics 11 Analysis
Software, allowing a range for the contact point fit of 30% of the maximum load.
Effective Young's modulus values of fit R^2 better than 0.95 were used for further
statistics. Loss and storage moduli were determined for the different frequencies (1
Hz, 2 Hz, 4 Hz and 10 Hz) using the software default methods.

Immunofluorescent labelling and imaging

Monolayers on hydrogels were fixed with 2% paraformaldehyde (Merck, 818715) for
8 min at RT and permeabilised with 0.3% Triton-X-100 (Sigma Aldrich, T8787) for 2
min at RT. Samples were then blocked with 1% BSA (Serva, 11930) in PBS for 30
min at RT, followed by incubation with primary antibodies (Table 1) in blocking
solution overnight at 4ºC. After washing, secondary antibodies (Table 1) diluted in
PBS were added for 2 h at RT. Samples were washed and mounted in Mowiol
mounting medium (Carl Roth, 9002-89-5) before imaging. Unless started otherwise,
samples were imaged at Zeiss LSM710 Confocal Microscope with 40x water
immersion objective (ZEISS, 420862-9970-000) in Airyscan super resolution mode as
z-stacks of 0.5 µm optical slicing. Uniform imaging settings were used between
samples within the same experiment. Images were then inspected and further
processed for different analyses using Fiji [107].

Bulk RNA sequencing

RNA isolation for RNA sequencing was performed using the RNeasy Micro RNA
isolation kit (Qiagen, 74004) with DNase treatment following the manufacturers
protocol. Bulk RNA sequencing was performed by the Life & Brain NGS Core Facility
(Bonn, Germany) after QuantSeq 3'-mRNA Library Prep/ Lexogen enrichment at
NovaSeq 6000 with a depth of 20M reads. The raw data was processed using nf-
core/rnaseq v3.12.0 (https://doi.org/10.5281/zenodo.1400710) of the nf-core

collection of workflows [108]. The pipeline was executed with Nextflow v23.04.1 [109]. The
complete data set is available on the Gene Expression Omnibus repository (GEO
accession number GSE297557). Differential expression analysis was performed
using DESeq2 package [110] in R, applying an alpha of 0.05. Genes were considered
differentially expressed at an adjusted p-value lower than 0.1. Heatmaps were
generated from the DESeq data set after z-normalisation using the package
pheatmap [111] and a custom-curated gene list. Functional enrichment of terms related
to differentially expressed genes was performed using the g:Profiler online tool
[112] using the terms GO molecular function, GO cellular component and GO biological
process as well as the pathways KEGG, Reactome and WikiPathways. For
comparison, publicly available data on the ageing human RPE was obtained from
Butler et al.[55] and Cai et al. [54].

Immunoblotting

Samples were harvested using RIPA buffer [113] with phosphatase inhibitors
PhosSTOP™ (Roche, 4906845001) and were subjected to DNA digestion for 15 min
by Pierce Universal Nuclease for cell lysis (Thermo Fischer, 88700) prior to addition
of Lämmli buffer [114]. Samples were then heated to 95 ºC for 7 min and ca. 10 µg
were loaded onto a 10 to 14% (depending on target protein molecular weight)
polyacrylamide gel [115] for SDS-PAGE together with a ProSieve Quadcolor Protein
Marker (Lonza, 00193837). Proteins were blotted onto Immobilion-P PFDV
membrane (Thermo Fischer 88518) and labelled with primary and secondary
antibodies for different targets (Table 1). Bands were detected at the Fusion Solo
(Vilber Lourmat) after detection with the Pierce™ ECL Western blotting substrate
(Thermo Fischer, 32106). The signal was quantified in Fiji [107] by using equal-sized
region of interest (ROI) for all bands and measuring the mean grey value. The
measurements for the proteins of interest were normalised to the housekeeping
protein GAPDH.

Modulation of actin nucleator and myosin II activity

Actin nucleation by Arp2/3 was modulated by using CK-666 (Tocris Bioscience,
#3950) at 500 nM in medium. After 1 h incubation, the monolayers were
characterised by nanoindentation. For phagocytosis studies, the monolayers were
incubated for 30 min with 500 nM CK-666, followed by incubation with the
photoreceptor fragments and the bridging molecules in CK-666 containing medium
for 4 h. For modulating formin activity, 20 µM SMIFH2 (Tocris Bioscience, #4401)
was used instead keeping the same timing and procedure. DMSO at a paired
concentration was used as carrier control. For the incubation with different SMIFH2
concentrations, a series of dilutions was prepared starting from a 100 µM solution,
resulting in a dilution series of 100, 40, 20 and 5 µM concentrations. For
nanoindentation and POS functional studies, the same timing and procedure were
used as described above with DMSO as a carrier control. Myosin II inhibition was
performed using 5 µM blebbistatin (Abcam, ab120425) for 1 h before nanoindentation
and for 30 min before POS addition in blebbistatin-containing medium (see above).

Analysis of extrusion events

Extrusion events were quantified from live cell imaging of apoptosis induction on SiR-Actin (Spirochrome, SC001) labelled cells. The last 15 hours of apoptosis induction were used for the quantification which relies on extrusion events being visible as very bright spots. The video was aligned to correct for stage shifts, smoothed using a Gaussian filter and overlayed in a maximum intensity z-projection in Fiji. The resulting image was thresholded to generate a binary mask and extruded cells were counted using the particle analysis tool in Fiji.

Analysis of cell density

Cell density was quantified from snaps of nuclear staining obtained on epi-fluorescence microscope at 20x magnification. Three to four snaps were used per condition per experiment. Nuclear signal was smoothed using a Gaussian filter, then detected by thresholding. The binary mask was subjected to watersheding and the number of nuclei was counted by using the particle analysis tool in Fiji. Density was calculated by dividing the obtained nuclei count to the area of the field of view. Data originates from at least three independent experiments.

Analysis of cell height

Cell height was determined from super resolution z-stacks of F-actin staining in cross-sectional view. Cells were imaged fixed after mounting in Mowiol, which leads to lower absolute height compared to the height detectable in live-cell imaging. Ten cells per stack were measured with the line tool in three stacks per condition from three independent experiments.

Analysis of cell outline segmentation and shape factor calculation

Cellular outlines from F-actin labelling were segmented using cellpose [116]. The generated regions of interest (ROI) were imported in Fiji and the cell area and perimeter were measured to calculate the dimensionless shape factor [117,118].

Analysis of stress fibres

Cells with stress fibres were manually counted using the multipoint tool in Fiji and normalised to the cell number as calculated after segmentation of cells via cellpose.

Analysis of microvilli length

Microvilli length was characterised from cross-sectional views of ezrin labelling in Fiji. Briefly, at least three cross-sections from at least three stacks per condition from three independent experiments were blurred with a Gaussian blur and a threshold was applied to capture the structure. The resulting masks were selected via the magic wand tool and an eclipse fit was performed. The minor axis was used as an approximate measure of the microvilli length.

Photoreceptor fragment phagocytosis analysis

The photoreceptor fragment phagocytosis was characterised using the Software
Imaris v10.2.0 (Oxford Instruments). Briefly, the actin labelling was used to create a
surface capturing only the apical actin portion of the monolayer. The average z-
position of the surface was determined as a cuff-off height to distinguish internal vs
external particles. Separately, photoreceptor fragments were segmented as a surface
and the cut-off height was used to distinguish between internalised and external
fragments. The count and volume of the fragments of both populations was recorded
and used for further analysis.

Quantification of apical actin area

The apical actin area of 300 x 300 pixel field of view was segmented from z-stacks of
phalloidin staining during photoreceptor internalisation as a surface in Imaris v10.2.0
(Oxford Instruments). For the region of interest, only a set number of slices including
the same upper portion of the cell monolayer were used. The covering area, as
internally determined by the software, was then noted.

Quantification of vinculin junctional to cytoplasmic ratio

Junctional signal of vinculin was quantified as a ratio of junctional to cytoplasmic
mean grey value in single optical sections. For this, paired F-actin signal was used to
segment cell outlines via cellpose as described above. The generated ROIs were
imported in Fiji and a custom macro was used to 'slim-down' the ROIs, invert the
mask, watershed the signal and generate ROIs that encompass the junctional
regions around cells. The generated junctional ROIs were used to measure grey
value as an average per image. The 'slimmed-down' cell ROIs were used to measure
the cytoplasmic signal. The average values of junctional and cytoplasmic signals per
image were used to generate the junctional to cytoplasmic vinculin measurements.

Quantification of junctional pMLC2

Paired samples from the same experiment were stained with the same antibody
solution and imaged with the same imaging settings. In single optical sections at the
junctional level, pMLC2 signal was thresholded in Fiji making sure the junctional
signal is captured. The mean grey value was measured only within the thresholded
region. At least three images were measured per condition per experiment.

Quantification of MRTFA

MRTFA was quantified as nuclear to cytoplasmic ratio of the mean grey value. For
this, paired cell nuclei signal (DAPI staining) was segmented via cellpose to generate
nuclear ROIs. Nuclear MRTFA was measured as average of the mean grey value of
all nuclei ROIs within the image. Cytoplasmic MRTFA was measured after setting
nuclear ROIs pixels as zero and thresholding the rest of the image. Then, the mean
grey value of the thresholded, non-zero region was measured.

Brillouin microscopy

Brillouin imaging was performed using a commercial setup based on an inverted Nikon Ti-2 Eclipse microscope using a vertically polarised, continuous-wave laser (Cobolt Flamenco, λ = 660 nm, ~ 100 mW) beam with a temperature-tuned air-spaced etalon (LightMachinery) attenuated by 0.6 optical density (OD) filter, resulting in approximately 9 mW incident power at the sample. Light was coupled into a confocal circulator (LightMachinery, HF-13311) and directed to the microscope through a polarising beam splitter and a quarter wave plate (QWP) yielding circularly polarised light. Measurements were performed on a motorised translation stage (Prior Scientific, H117E2NN) using an air objective (Nikon S Plan Fluor 20X NA 0.45), achieving an estimated 0.73 μm lateral and 6.52 μm axial resolution. Brightfield and fluorescence imaging were performed sequentially using a Teledyne sCMOS camera and epi-fluorescence illumination (Nikon D-LEDi).

Backscattered Rayleigh and Brillouin signals were collected with the same objective and routed back through the confocal circulator. The Brillouin signal was acquired using a LightMachinery HyperFine spectrometer (HF-8999-PK-660), equipped with a "pump killer" (PK) module with an air-spaced, pressure-tuned etalon to suppress the elastically scattered Rayleigh light and laser line reflection (~ 60 dB attenuation). Frequency-shifted Brillouin Stokes and anti-Stokes components were directed into the spectrometer, comprised of a virtually imaged phase array (VIPA) and a reflective echelle diffraction grating for spectral dispersion. The cross-dispersed orders were focused onto the CMOS camera sensor (Hamamatsu ORCA-Fusion, C14440-20UP) simultaneously acquiring the full spectrum. The typical acquisition time of a single spectrum is 500 ms, with a spectral resolution < 1 pm.

Spectral data acquisition and raster scanning was facilitated with LightMachinery LabView-based software. The x-y pixel coordinates on the CMOS sensor are converted to wavelength (or frequency) and a Lorentzian fit is applied over each Brillouin peak as described by equation (1).

$$I(\omega) = \frac{I_0}{\pi} \frac{\Gamma \omega_B^2}{(\omega^2 - \omega_B^2)^2 + \Gamma^2 \omega_B^2} \#(1)$$

The final Brillouin shift ω_B and linewidth Γ values for a given spectrum were computed by averaging the corresponding values from the Stokes and anti-Stokes peaks forming a single voxel in the raster scan. Z-scans were acquired as a 3x3 matrix in a step size of 20 μm and z-step of 2 μm, z-range of 80 μm, while apical and basal scans were acquired over the region of 50 x 50 μm with a lateral step size of 1.5 μm. The z-position of apical and basal Brillouin scans within the monolayer was selected based on the appearance of the cells in brightfield and SiR-actin fluorescence images. The scans were limited to 15 minutes to obtain an optimal raster scan resolution while maintaining the cell viability. It should be noted that although the average laser power at sample is within the range used in biological

samples, by comparing brightfield images taken before and after the Brillouin scan,
some deformation of the sample was observed. This could have occurred due to the
heat of the laser affecting the gel or the cells as the RPE cells' pigmentation
contributes to the linear laser light absorption.

Brillouin shift data analysis

For segmentation, the Brillouin shift maps from basal layers were linearly scaled and
converted into 8-bit grayscale images, proportionally reflecting the original Brillouin
shift values. The grayscale images were segmented using cellpose to identify the
individual structures within the images. Following segmentation, two populations
were defined: "core", corresponding to segmented regions, and "interstitium",
corresponding to areas between them. These modified images were then imported
into a custom-written MATLAB script, where appropriate binary masks were created
and applied to the original Brillouin shift maps to separate the "core" and the
"interstitium". For each raster scan, the mean Brillouin shift and the standard
deviation was calculated separately for the two regions. The 3 x 3 raster z-scans
resulted in 9 representative z-axis line profiles per region, capturing mechanical
transition across the sample. Each profile typically displayed a stiffness change from
the hydrogel to the monolayer and then from the monolayer to the surrounding cell
media. For each sample, 5 of these scans were obtained.

The Brillouin Shift data was imported into a custom-written MATLAB script to identify
the local peaks or, where absent, plateaus in the Brillouin shift profiles
(https://github.com/GiedreAstra/Example_E3_zscan_RPE.git). Each line profile was
smoothed with a Savitzky-Golay filter, the peak was detected using MATLAB's
findpeaks function, with a minimum width imposed to prevent false peak
identification. A gradient of the filtered z-line was taken. To refine peak localisation, a
Gaussian fit was applied across a range of filtered Brillouin shift values. The fitting
range was bound by the z-positions at which the local maximum and the minimum of
the derivative of the z-line was within an appropriate range from the previously
identified peak. The fitted peak value and position were retrieved. In the case of no
peak detection, the code identifies the onset of a drop in the plateau by considering
the change in the derivative over a moving average window. The corresponding
plateau value was then recorded. The performance of the script was checked for
individual z-profiles to exclude outliers.

STATISTICAL ANALYSIS

Statistical analysis was performed using GraphPad Prism on paired averaged values
originating from at least three independent biological experiments. Datapoints from at
least three technical replicates or from matrix measurements within the samples,
epithelial monolayers, were averaged into representative values from a biological
replicate. Assuming normal distribution, the data was tested with a paired, two-sided
test depending on experimental design with statistical tests and number of biological
replicates for each graph given in the captions of the figures.

DECLARATION OF AI USE

The preparation of this manuscript has been aided by AI-assisted technologies to improve its readability and language.

ACKNOWLEDGMENT

We would like to acknowledge all the members of the REMeD group in the past years for their contributions to the discussion of the work. Moreover, we appreciated Adam Breitscheidel's technical support in graphic design, Gloria's excellent help in imaging and insightful feedback from Rudolf Leube. We would like to acknowledge the support of Chris Davis from the Centre for Virus Research of the University of Glasgow for providing a lab space in Garscube campus to enable AAV work needed for the sample preparation for Brillouin microscopy. This work was supported by the Two-Photon Imaging Facility of the Interdisciplinary Center for Clinical Research (IZKF) within the Faculty of Medicine at RWTH Aachen University.

FUNDINGS

This work in Di Russo's laboratory was supported by: A grant from the Interdisciplinary Centre for Clinical Research within the Faculty of Medicine at the RWTH Aachen University and German Research Foundation (DFG; RU 2366/3-1). Furthermore, J.D.R. and W.W. are supported by the DFG within 363055819/GRK2415.

The collaboration with Vassalli's lab for Brillouin microscopy was supported by: RS MacDonald Facility Access funding grant by the Scottish Universities Life Sciences Alliance (SULSA). A travel grant by the German Academic Exchange Service (DAAD) program number 57745342 and a travel grant by Boehringer Ingelheim Fonds.

T.P. and A.N.K. are supported by Add-on Fellowships of the Joachim Herz Foundation.

AUTHORS CONTRIBUTION

JDR conceived and supervised the project. TP designed and performed all the experiments. GA, ANK and TP performed Brillouin measurements. GA and ANK performed Brillouin analysis. MM performed initial transcriptomics data analysis and advised on the transcriptomic exploration. RR supported the immunoblotting experiments. SS supported with confocal imaging. LY supported the establishment of viral transduction. WW and MV supported the work with expertise and advice. JDR and TP wrote the manuscript, and all authors discussed and commented on it.

COMPETING INTERESTS

Authors declare that they have no competing interests.

988 DATA AND MATERIALS AVAILABILITY

989 All data are available in the main text or the supplementary materials. Furthermore,
990 the complete data set of RNA bulk sequencing is available on the Gene Expression
991 Omnibus repository (GEO accession number GSE297557).

Supplementary figures

Suppl. figure 1: The iRPE and dr-iRPE model and cell morphometric characterisation. A: Schematics of iRPE cell culture system displaying the monolayer obtained on 4 kPa hydrogels functionalised with basement membrane mimicking coating. **B:** Cross-section of iRPE after 20 days of culture labelled for F-actin (red) and ezrin (green) showing characteristic apicobasal polarisation. **C:** Representative micrograph of proliferation assay showing iRPE cells in brightfield and proliferating cell nuclei labelled by EdU-AF488 (green). **D:** Map of plasmid used for the expression of pro-apoptotic FKBP-casp8 protein and delivered to iRPE cells by AAV5. **E:** Representative micrographs of SiR-actin labelled iRPE (white) expressing mApple as fluorescent reporter (red) at the timepoint of AP20187 addition and 10 h afterwards. Extrusion events of apoptotic cells at the latter point are indicated with yellow arrows. **F:** Quantification of average cell area (n=4). **G:** Relative frequency distribution of cell areas (from 3263 (iRPE) and 2821 (dr-iRPE) cells from n=4). **H:** Standard deviation of cell areas within an experiment (n=4). **I:** Average cell shape factor (n=4). Datapoints represent average values ± SEM per experiment. Statistical significance was tested using a paired t-test, where **: $p < 0.01$.

1008

**Suppl. figure 2: Controls for influence of virus, recombinant FKBP-casp8 protein expression and debris
released into medium. A:** Vector map of plasmid used as viral control, which promotes the expression of GFP.
B: Representative micrographs of F-actin in GFP-iRPE compared to dr-iRPE (top) and cross sections (down). **C:**
Quantification of average cell density for GFP-iRPE (n=5). **D:** Quantification of cell height for GFP-iRPE (n=4). **E:**
Representative micrographs showing ezrin as maximum intensity projection in top (left) and side views (middle)
and the quantification of ezrin-positive microvilli length (right, n=4). **F:** Experimental schematics of mixed
population control. The micrographs show representative monolayers of mApple-positive transduced cells within
the naïve cells at day 4 and at day 20 (the readout day). **G:** Representative micrographs of F-actin labelled mixed
population iRPE without apoptosis induction (noAP) and after induction of apoptosis (AP). **H:** Quantification of
average cell density for the mixed population control (n=3). **I:** Quantification of cell height for the mixed population
control (n=3). **J:** Experimental schematics of medium-transfer control **K:** Representative micrographs of F-actin
labelled control monolayers vs. monolayers subjected to medium transfer. **L:** Quantification of average cell
density for the medium transfer control (n=3). **M:** Quantification of cell height for the medium transfer control
(n=3). Datapoints represent average values per experiment ± SEM. Statistical significance was tested using a
paired t-test, where *: p<0.05, **: p<0.01.

1024

1025

Suppl. figure 3: POS internalisation assay. A: Workflow for 3D segmentation, cut-off height determination and
classification into 'bound' vs 'internalised' particles. **B:** Quantification of number of internalised POS particles
normalised to cell number (n=6). **C:** Quantification of bound POS particles number in the field of view (n=6). **D:**
Quantification of average bound POS particle volume (n=6). **E:** Representative micrographs of iRPE at t = 4 h of
POS internalisation assay as orthogonal cross-sections (up) or as single optical slices in the middle of cells (lower
two panels) for the "mixed" control. **F:** Representative micrographs of iRPE at t = 4 h of POS internalisation assay
as orthogonal cross-sections (up) or as single optical slices in the middle of cells (lower two panels) for the
medium-transfer control (MedTr). **G:** Quantification of POS fragments internalised within the time of the assay per

field of view (left) and quantification of the average volume of internalised fragments (right) for the "mixed"
population control (n=3). **H:** Quantification of POS fragments internalised within the time of the assay per field of
view (left) and quantification of the average volume of internalised fragments (right) for the medium-transfer
control (n=3). Datapoints represent average values per experiment. Statistical significance was tested using a
paired t-test, where *: p<0.05, **: p<0.01.

1039

1040 **Suppl. figure 4: Differential apical actomyosin remodelling between iRPE and dr-iRPE during POS**
1041 **internalisation assay. A:** Representative micrographs of the apical side of monolayers during internalisation
1042 showing F-actin and photoreceptors. Arrows point to the presence of protruding actin-rich structures in dr-iRPE.
1043 **B:** Representative orthogonal cross sections of iRPE and dr-iRPE monolayers during internalisation showing F-
1044 actin and FITC-labelled POS. Arrows point to actin-rich structures.

B

Gene symbol	Protein name	Direction change
Overlapping DEGs with reported age-related changes in humans		
Cai H. et al. 2012 (DOI:10.3389/fnagi.2012.00008)		
DKK1	dickkopf WNT signaling pathway inhibitor 1(DKK1)	up
LAMA3	laminin subunit alpha 3(LAMA3)	up
TYMS	thymidylate synthetase(TYMS)	up
PLEKHG4	pleckstrin homology and RhoGEF domain containing G4(PLEKHG4)	down
SCN2B	sodium voltage-gated channel beta subunit 2(SCN2B)	down
Butler J. M. et al. 2021 (DOI: 10.1111/jcmm.16569)		
AKAP6	A-kinase anchoring protein 6(AKAP6)	up
BMP8B	bone morphogenetic protein 8b(BMP8B)	up
CCDC51	coiled-coil domain containing 51(CCDC51)	up
CTSV	cathepsin V(CTSV)	up
DUSP6	dual specificity phosphatase 6(DUSP6)	up
KL	klotho(KL)	up
SLC6A20	solute carrier family 6 member 20(SLC6A20)	up
TNFRSF10C	TNF receptor superfamily member 10c(TNFRSF10C)	up
TRPV4	transient receptor potential cation channel subfamily V member 4(TRPV4)	up
CPAMD8	C3 and PZP like alpha-2-macroglobulin domain containing 8(CPAMD8)	down
CSRP2	cysteine and glycine rich protein 2(CSRP2)	down
DCT	dopachrome tautomerase(DCT)	down
GAS1	growth arrest specific 1(GAS1)	down
TMEM150C	transmembrane protein 150C (TMEM150C)	down

1045

Suppl. figure 5: Density reduced iRPE shows a differential gene expression from iRPE, which is partly overlapping with ageing patients. A: Base mean vs Log2Fold change for significantly differentially expressed actin-associated genes in dr-iRPE vs iRPE (n=4). **B:** Differentially expressed genes (DEGs) overlapping with the results from studies on ageing human RPE from Cai H. et al. [54] and Butler J. M. et al. [55]. Gene names are labelled in blue or red for downregulated or upregulated, respectively. **C:** Volcano plot of upregulated DEGs between dr-iRPE and iRPE. **D:** Volcano plot of downregulated DEGs between dr-iRPE and iRPE. Cut-off lines were drawn at log2fold=1, corresponding to a fold change of two, and at log10P = -2, corresponding to a p of 0.001 and padj of 0.1, which we set as cut-off significance value for DEGs. Genes are assigned to groups based on their log2fold and log10fold values as highlighted in the legend.

1055

Suppl. figure 6: Biomechanical characterisation of iRPE conditions. A: Experimental setup for performing nanoindentation. The micrograph shows a top view of SiR-actin labelled cell monolayer and cantilever with tip. **B:** Average loss modulus of iRPE and dr-iRPE (n=4). **C:** Immunoblot of pMLC2 and GAPDH as a housekeeping protein and quantification of total pMLC2 protein from immunoblots, normalised to GAPDH (n=4). **D:** Average Effective Young's modulus, storage and loss moduli for virus control (n=4). **E:** Average Effective Young's modulus, storage and loss moduli for treated (noAP) and untreated (AP) mixed population iRPE (n=4). **F:** Average Effective Young's modulus, storage and loss moduli for iRPE and conditioned medium treated iRPE (MedTr-iRPE) (n=4). Datapoints represent average values ± SEM per experiment. Statistical significance was tested using a paired t-test, where *: p<0.05 and ***: p<0.001.

Suppl. figure 7: Distribution of Brillouin shift values in apical and basal layers of iRPE and density reduced iRPE and segmentation of Brillouin shift values from basal maps. A: Representative histograms of iRPE and dr-iRPE basal and apical Brillouin shift values acquired from 50 x 50 μm regions of interest and the respective plots. **B:** Representative plots of segmented Brillouin shift maps from basal scans with two population modes defined by Cellpose segmentation (see Methods). **C:** Representative histogram of the plots shown in B illustrating the identified populations. **D:** Quantification of Brillouin shift population modes named as "Interstitium" and "Core" in basal iRPE and dr-iRPE monolayers (n=3). **E:** Representative F-Actin micrographs of basal iRPE and dr-iRPE monolayers with annotated regions enlarged. Datapoints in D represent average values per experiment ± SEM. Statistical significance was tested using a one-way ANOVA, where *: p<0.05 and **: p<0.01.

Suppl. figure 8: Linear nucleator inhibition with SMIFH2 during photoreceptor phagocytosis. A:
Quantification of cell height for iRPE vs SMIFH2-treated iRPE (n=3). **B:** Average Effective Young's Modulus,
derived from 2.5 µm nanoindentation with a 20 µm spherical tip for iRPE vs SMIFH2-treated iRPE monolayers
(n=8). **C:** Quantification of cell height for dr-iRPE vs SMIFH2-treated dr-iRPE (n=4). **D:** Average Effective Young's
Modulus for dr-iRPE vs SMIFH2-treated dr-iRPE monolayers (n=8). **E:** Quantification of the number of
internalised photoreceptor particles normalised to cell number for iRPE vs SMIFH2-treated iRPE (n=4). **F:**
Quantification of the number of bound particles in the field of view for iRPE vs SMIFH2-treated iRPE (n=4). **G:**
Quantification of the number of internalised photoreceptor particles for dr-iRPE vs SMIFH2-treated dr-iRPE (n=8).
H: Quantification of the average volume of internalised photoreceptor particles for dr-iRPE vs SMIFH2-treated dr-
iRPE (n=8). **I:** Quantification of apical actin area per field of view during internalisation in iRPE vs SMIFH2-treated
iRPE (n=4). **J:** Average Effective Young's Modulus of iRPE monolayers after 1 h treatment with SMIFH2
concentrations of 5, 20, 40 or 100 µM (n=4). **K:** Average Effective Young's Modulus of iRPE monolayers after 1 h
treatment with 5 µM blebbistatin (n=4). **L:** Average Effective Young's Modulus of dr-iRPE monolayers after 1 h
treatment with SMIFH2 concentrations of 5, 20, 40 or 100 µM (n=4). **M:** Average Effective Young's Modulus of dr-
iRPE monolayers after 1 h treatment with 5 µM blebbistatin (n=4). **N:** Number of internalized POS particles of
iRPE monolayers after treatment with SMIFH2 concentrations of 5, 20, 40 or 100 µM (n=4). **O:** Number of
internalized POS particles of iRPE monolayers after treatment with 5 µM blebbistatin (n=4). **P:** Average volume of

internalized particles of iRPE monolayers after treatment with SMIFH2 concentrations of 5, 20, 40 or 100 µM (n=4). **Q:** Average volume of internalized particles of iRPE monolayers after treatment with 5 µM blebbistatin (n=4). **R:** Number of internalized POS particles of dr-iRPE monolayers after treatment with SMIFH2 concentrations of 5, 20, 40 or 100 µM (n=4). **S:** Number of internalized POS particles of dr-iRPE monolayers after treatment with 5 µM blebbistatin (n=4). **T:** Average volume of internalized particles of dr-iRPE monolayers after treatment with SMIFH2 concentrations of 5, 20, 40 or 100 µM (n=4). **U:** Average volume of internalized particles of dr-iRPE monolayers after treatment with 5 µM blebbistatin. Datapoints represent average values (± SEM in case of bar graphs) per experiment. Statistical significance in A, B, C, D, I, K and M was tested using a paired t-test, where *: $p<0.05$ and **: $p<0.01$. Statistical significance in E, F, G, H, O, Q, S and U was tested using a ratio paired t-test, where **: $p<0.01$. Statistical significance in J, L, N, P, R and T was tested using a One-way ANOVA with multiple comparisons, where *: $p<0.05$ and **: $p<0.01$.

1107

Suppl. figure 9: Branched nucleator inhibition with CK666 during photoreceptor phagocytosis. A: Quantification of cell height for dr-iRPE vs CK666-treated dr-iRPE (n=4). **B:** Average Effective Young's Modulus, derived from 2.5 µm nanoindentation with a 20 µm spherical tip for dr-iRPE vs CK666-treated dr-iRPE (n=3). **C:** Quantification of cell height for iRPE vs CK666-treated iRPE (n=4). **D:** Average Effective Young's Modulus, derived from 2.5 µm nanoindentation with a 20 µm spherical tip for iRPE vs CK666-treated iRPE (n=4). **E:** Quantification of the number of internalised photoreceptor particles normalised to cell number for dr-iRPE vs CK666-treated dr-iRPE monolayers (n=4). **F:** Quantification of the number of bound particles in the field of view for dr-iRPE vs CK666-treated dr-iRPE (n=4). **G:** Number of internalized POS particles of iRPE vs CK666-treated iRPE (n=4). **H:** Average volume of internalized particles of iRPE vs CK666-treated iRPE (n=4). **I:** Quantification of apical actin area per field of view of monolayers during internalisation in dr-iRPE vs CK666-treated dr-iRPE (n=3). Datapoints represent average values (± SEM in case of bar graphs) per experiment. Statistical significance in A - D and I was tested using a paired t-test, where *: p<0.05 and **: p<0.01. Statistical significance in E - H and I was tested using a ratio paired t-test, where *: p<0.05.

1121

1122

1123

1124

1125 **Supplementary information**

1126

1127 **Table 1: Antibodies and dyes used for immunofluorescence (IF) or immunoblotting (WB).**

Name	Source	Concentration/Dilution
Primary antibodies		
Anti phospho-myosin light chain 2 (Ser19)	Cell Signalling, 3671	1:50 (IF); 1:1000 (WB)
Anti vinculin	Sigma, V4139	15 µg/ml (IF)
Anti ezrin	abcam, ab4069	10 µg/ml (IF); 1 µg/ml (WB)
Anti phospho ezrin (Thr567)	Invitrogen, PA5-37763	5 µg/ml (IF); 1 µg/ml (WB)
Anti non muscle myosin heavy chain IIB	BioLegend, 909901	4 µg/ml (IF)
Secondary antibodies		
AF488 anti mouse IgG (H+L)	Invitrogen, A11001	8 µg/ml (IF)
AF594 anti mouse IgG (H+L)	Invitrogen, A11005	8 µg/ml (IF)
AF594 anti rabbit IgG (H+L)	Invitrogen, A11012	8 µg/ml (IF)
AF647 anti rabbit IgG (H+L)	Jackson/Dianova, 111-605-144	7,5 µg/ml (IF)
AF 488 anti rabbit IgG (H+L)	Jackson/Dianova, 711-546-152	8 µg/ml (IF)
HRP anti rabbit IgG	Abcam, ab99696	1:5000 (WB)
HRP anti mouse IgG (H+L)	Jackson/Dianova, 115-035-003	1:5000 (WB)
Fluorescent dyes		
Phalloidin-iFluor 488	Abcam, ab176753	1:200 (IF)
Phalloidin-iFluor 647	Abcam, ab176759	1:200 (IF)
Phalloidin AF568	Molecular probes, A12380	1:200 (IF)

1128

Videos Legends

Video 1: On-demand activation of FKBP/casp8 by AP20187 leads to monolayer-
scale extrusion events. Live-iRPE monolayer time-lapse expressing FKBP/casp8
(mApple-positive red cells) and labelled with actin dye SiR-Actin (white). Upon the
addition of the AP20187 compound, cells are extruded due to apoptosis induction.
Extrusion events include neighbouring cells adopting a characteristic rosette-like
morphology, while the dying cell is extruded out of the monolayer, forming a bright
blob.

1137

Video 2: Live-cell time-lapse of iRPE monolayer during nanoindentation. The SiR-Actin labelled iRPE monolayer is probed by nanoindentation in a matrix scan (left). A zoom-in view (right) reveals the junctional deformation occurring during the measurement.

References

1. Thomson, J. A. On Growth and Form. **100**, 21–22 (1917).

2. Ball, P. *Shapes: Nature's Patterns.* (Oxford University Press, USA, 2009).

3. Bepler, T. & Berger, B. Learning the protein language: Evolution, structure, and function. *Cell Syst.* **12**, 654-669.e3 (2021).

4. Gligorijević, V. *et al.* Structure-based protein function prediction using graph convolutional networks. *Nat. Commun.* **12**, 3168 (2020).

5. Yang, J., Griffin, A., Qiang, Z. & Ren, J. Organelle-targeted therapies: a comprehensive review on system design for enabling precision oncology. *Signal Transduct. Target. Ther.* **7**, 379 (2022).

6. Glancy, B., Kim, Y., Katti, P. & Willingham, T. B. The Functional Impact of Mitochondrial Structure Across Subcellular Scales. *Front. Physiol.* **11**, 541040 (2020).

7. Morgan, E. F. & Gerstenfeld, L. C. Marcus and Feldman's Osteoporosis. 15–35 (2021) doi:10.1016/b978-0-12-813073-5.00002-2.

8. Baum, G. L. *et al.* Development of structure–function coupling in human brain networks during youth. *Proc. Natl. Acad. Sci.* **117**, 771–778 (2020).

9. Simpson, C. L., Patel, D. M. & Green, K. J. Deconstructing the skin: cytoarchitectural determinants of epidermal morphogenesis. *Nat. Rev. Mol. Cell Biol.* **12**, 565–580 (2011).

10. Anlaş, A. A. & Nelson, C. M. Tissue mechanics regulates form, function, and dysfunction. *Curr. Opin. Cell Biol.* **54**, 98–105 (2018).

11. Ayad, N. M. E., Kaushik, S. & Weaver, V. M. Tissue mechanics, an important regulator of development and disease. *Philosophical Transactions Royal Soc B* **374**, 20180215 (2019).

12. Guillot, C. & Lecuit, T. Mechanics of Epithelial Tissue Homeostasis and Morphogenesis. *Science* **340**, 1185–1189 (2013).

13. Biggs, L. C., Kim, C. S., Miroshnikova, Y. A. & Wickström, S. A. Mechanical Forces in the Skin: Roles in Tissue Architecture, Stability, and Function. *J. Investig. Dermatol.* **140**, 284–290 (2020).

14. Guimarães, C. F., Gasperini, L., Marques, A. P. & Reis, R. L. The stiffness of
living tissues and its implications for tissue engineering. *Nat. Rev. Mater.* **5**, 351–
370 (2020).

15. Gómez-Gálvez, P., Anbari, S., Escudero, L. M. & Buceta, J. Mechanics and self-
organization in tissue development. *Semin. Cell Dev. Biol.* **120**, 147–159 (2021).

16. Hannezo, E. & Heisenberg, C.-P. Mechanochemical Feedback Loops in
Development and Disease. *Cell* **178**, 12–25 (2019).

17. Kaliman, S. *et al.* Mechanical Regulation of Epithelial Tissue Homeostasis. *Phys*
Rev X **11**, 031029 (2021).

18. DuFort, C. C., Paszek, M. J. & Weaver, V. M. Balancing forces: architectural
control of mechanotransduction. *Nat Rev Mol Cell Bio* **12**, 308–319 (2011).

19. Ishikawa, J. *et al.* Mechanical homeostasis of liver sinusoid is involved in the
initiation and termination of liver regeneration. *Commun. Biol.* **4**, 409 (2021).

20. Eichinger, J. F. *et al.* Mechanical homeostasis in tissue equivalents: a review.
Biomech. Model. Mechanobiol. **20**, 833–850 (2021).

21. Kozyrina, A. N. *et al.* Laminin-defined mechanical status modulates retinal
pigment epithelium phagocytosis. *EMBO Rep.* 1–27 (2025) doi:10.1038/s44319-
025-00475-9.

22. Tamori, Y. & Deng, W.-M. Tissue Repair through Cell Competition and
Compensatory Cellular Hypertrophy in Postmitotic Epithelia. *Dev. Cell* **25**, 350–
363 (2013).

23. Campello, L. *et al.* Aging of the Retina: Molecular and Metabolic Turbulences and
Potential Interventions. *Annu Rev Vis Sc* **7**, 1–32 (2021).

24. Terry, R. D., DeTeresa, R. & Hansen, L. A. Neocortical cell counts in normal
human adult aging. *Ann. Neurol.* **21**, 530–539 (1987).

25. Freeman, S. H. *et al.* Preservation of Neuronal Number Despite Age-Related
Cortical Brain Atrophy in Elderly Subjects Without Alzheimer Disease. *J.*
Neuropathol. Exp. Neurol. **67**, 1205–1212 (2008).

26. Piskova, T., Kozyrina, A. N. & Di Russo, J. Mechanobiological implications of
age-related remodelling in the outer retina. *Biomater. Adv.* **147**, 213343 (2023).

27. Sparrrow, J. R., Hicks, D. & Hamel, C. P. The Retinal Pigment Epithelium in
Health and Disease. *Curr. Mol. Med.* **10**, 802–823 (2010).

28. Panda-Jonas, S., JONAS, J. B. & JAKOBCZYK-ZMIJA, M. Retinal Pigment
Epithelial Cell Count, Distribution, and Correlations in Normal Human Eyes. *Am. J.*
Ophthalmol. **121**, 181–189 (1996).

29. Bhatia, S. K. *et al.* Analysis of RPE morphometry in human eyes. *Mol Vis* **22**,
898–916 (2016).

30. Priore, L. V. D., Kuo, Y.-H. & Tezel, T. H. Age-related changes in human RPE cell
density and apoptosis proportion in situ. *Invest Ophth Vis Sci* **43**, 3312–8 (2002).

31. Katz, M. L. & Robison, W. G. Age-related changes in the retinal pigment
epithelium of pigmented rats. *Exp Eye Res* **38**, 137–151 (1984).

32. Ortolan, D. *et al.* Single-cell–resolution map of human retinal pigment epithelium
helps discover subpopulations with differential disease sensitivity. *Proc. Natl.*
Acad. Sci. **119**, e2117553119 (2022).

33. Tarau, I.-S., Berlin, A., Curcio, C. A. & Ach, T. The Cytoskeleton of the Retinal
Pigment Epithelium: from Normal Aging to Age-Related Macular Degeneration. *Int*
J Mol Sci **20**, 3578 (2019).

34. Lai, Y. L. & Rana, M. W. A Study of Photoreceptor–Retinal Pigment Epithelium
Complex: Age-Related Changes in Monkeys. *P Soc Exp Biol Med* **181**, 371–381
(1986).

35. Inana, G. *et al.* RPE phagocytic function declines in age-related macular
degeneration and is rescued by human umbilical tissue derived cells. *J Transl Med*
16, 63 (2018).

36. Ma, J. Y. W. *et al.* Aging induces cell loss and a decline in phagosome
processing in the mouse retinal pigment epithelium. *Neurobiol. Aging* **128**, 1–16
(2023).

37. Brandl, C. *et al.* In-Depth Characterisation of Retinal Pigment Epithelium (RPE)
Cells Derived from Human Induced Pluripotent Stem Cells (hiPSC).
NeuroMolecular Med. **16**, 551–564 (2014).

38. Miyagishima, K. J. *et al.* In Pursuit of Authenticity: Induced Pluripotent Stem
Cell-Derived Retinal Pigment Epithelium for Clinical Applications. *STEM CELLS*
Transl. Med. **5**, 1562–1574 (2016).

39. Song, M. J. *et al.* Bioprinted 3D outer retina barrier uncovers RPE-dependent
choroidal phenotype in advanced macular degeneration. *Nat. Methods* **20**, 149–
161 (2023).

40. Müller, C., Charniga, C., Temple, S. & Finnemann, S. C. Quantified F-Actin
Morphology Is Predictive of Phagocytic Capacity of Stem Cell-Derived Retinal
Pigment Epithelium. *Stem Cell Rep* **10**, 1075–1087 (2018).

41. Oberst, A. *et al.* Inducible Dimerization and Inducible Cleavage Reveal a
Requirement for Both Processes in Caspase-8 Activation*. *J. Biol. Chem.* **285**,
16632–16642 (2010).

42. Ko, F. *et al.* Associations with Retinal Pigment Epithelium Thickness Measures in
a Large Cohort Results from the UK Biobank. *Ophthalmology* **124**, 105–117
(2017).

43. Bonilha, V. L., Rayborn, M. E., Saotome, I., McClatchey, A. I. & Hollyfield, J. G.
Microvilli defects in retinas of ezrin knockout mice. *Exp. Eye Res.* **82**, 720–729
(2006).

44. Kivelä, T., Jääskeläinen, J., Vaheri, A. & Carpén, O. Ezrin, a membrane-
organizing protein, as a polarization marker of the retinal pigment epithelium in
vertebrates. *Cell Tissue Res.* **301**, 217–223 (2000).

45. Bonilha, V. L., Finnemann, S. C. & Rodriguez-Boulan, E. Ezrin Promotes
Morphogenesis of Apical Microvilli and Basal Infoldings in Retinal Pigment
Epithelium. *J. Cell Biol.* **147**, 1533–1548 (1999).

46. Fievet, B. T. *et al.* Phosphoinositide binding and phosphorylation act sequentially
in the activation mechanism of ezrin. *J. Cell Biol.* **164**, 653–659 (2004).

47. Tixeira, R. *et al.* Defining the morphologic features and products of cell
disassembly during apoptosis. *Apoptosis* **22**, 475–477 (2017).

48. Zihni, C. Phagocytosis by the retinal pigment epithelium: New insights into
polarized cell mechanics. *BioEssays* **47**, e2300197 (2025).

49. Umapathy, A. *et al.* Spatiotemporal Live-Cell Analysis of Photoreceptor Outer
Segment Membrane Ingestion by the Retinal Pigment Epithelium Reveals Actin-
Regulated Scission. *J. Neurosci.* **43**, 2653–2664 (2023).

50. Kozyrina, A. N. *et al.* Laminin-defined mechanical status modulates retinal
pigment epithelium phagocytosis. *EMBO Rep.* 1–27 (2025) doi:10.1038/s44319-
025-00475-9.

51. Ma, J. Y. W. *et al.* Aging induces cell loss and a decline in phagosome
processing in the mouse retinal pigment epithelium. *Neurobiol. Aging* **128**, 1–16
(2023).

52. Kwon, W. & Freeman, S. A. Phagocytosis by the Retinal Pigment Epithelium:
Recognition, Resolution, Recycling. *Front Immunol* **11**, 604205 (2020).

53. Olson, E. N. & Nordheim, A. Linking actin dynamics and gene transcription to
drive cellular motile functions. *Nat. Rev. Mol. Cell Biol.* **11**, 353–365 (2010).

54. Cai, H., Fields, M. A., Hoshino, R. & Priore, L. V. D. Effects of Aging and
Anatomic Location on Gene Expression in Human Retina. *Front. Aging Neurosci.*
4, 8 (2012).

55. Butler, J. M., Supharattanasitthi, W., Yang, Y. C. & Paraoan, L. RNA‐seq analysis
of ageing human retinal pigment epithelium: Unexpected up‐regulation of visual
cycle gene transcription. *J. Cell. Mol. Med.* **25**, 5572–5585 (2021).

56. Chaudhuri, O. Viscoelastic hydrogels for 3D cell culture. *Biomater. Sci.* **5**, 1480–
1490 (2017).

57. Khalilgharibi, N. *et al.* Stress relaxation in epithelial monolayers is controlled by
the actomyosin cortex. *Nat. Phys.* **15**, 839–847 (2019).

58. Nagayama, M., Haga, H., Takahashi, M., Saitoh, T. & Kawabata, K. Contribution
of cellular contractility to spatial and temporal variations in cellular stiffness. *Exp.*
Cell Res. **300**, 396–405 (2004).

59. Wang, N. *et al.* Cell prestress. I. Stiffness and prestress are closely associated in
adherent contractile cells. *Am. J. Physiol.-Cell Physiol.* **282**, C606–C616 (2002).

60. Watanabe, T., Hosoya, H. & Yonemura, S. Regulation of Myosin II Dynamics by
Phosphorylation and Dephosphorylation of Its Light Chain in Epithelial Cells. *Mol.*
Biol. Cell **18**, 605–616 (2007).

61. Leerberg, J. M. & Yap, A. S. Vinculin, cadherin mechanotransduction and
homeostasis of cell–cell junctions. *Protoplasma* **250**, 817–829 (2013).

62. Brückner, B. R. & Janshoff, A. Importance of integrity of cell-cell junctions for the
mechanics of confluent MDCK II cells. *Sci. Rep.* **8**, 14117 (2018).

63. Prevedel, R., Diz-Muñoz, A., Ruocco, G. & Antonacci, G. Brillouin microscopy: an
emerging tool for mechanobiology. *Nat Methods* **16**, 969–977 (2019).

64. Passeri, A. A. *et al.* Beyond Water Content: Unraveling Stiffness in Hydrated
Materials by a Correlative Brillouin–Raman Approach. *ACS Photonics* **12**, 3794–
3802 (2025).

65. Geisler, F. *et al.* Intestinal intermediate filament polypeptides in C. elegans:
Common and isotype-specific contributions to intestinal ultrastructure and function.
Sci. Rep. **10**, 3142 (2020).

66. Chugh, P. *et al.* Actin cortex architecture regulates cell surface tension. *Nat. cell*
Biol. **19**, 689–697 (2017).

67. Lee, S. Y., Fujioka, T., Osuga, M., Nishimura, T. & Suetsugu, S. Plasma
Membrane Shaping. 245–263 (2023) doi:10.1016/b978-0-323-89911-6.00019-4.

68. Hetrick, B., Han, M. S., Helgeson, L. A. & Nolen, B. J. Small Molecules CK-666
and CK-869 Inhibit Actin-Related Protein 2/3 Complex by Blocking an Activating
Conformational Change. *Chem. Biol.* **20**, 701–712 (2013).

69. Rizvi, S. A. *et al.* Identification and Characterization of a Small Molecule Inhibitor
of Formin-Mediated Actin Assembly. *Chem. Biol.* **16**, 1158–1168 (2009).

70. Nishimura, Y. *et al.* The Formin Inhibitor, SMIFH2, Inhibits Members of the
Myosin Superfamily. *J. Cell Sci.* **134**, jcs.253708 (2021).

71. Pomatto, L. C. D. & Davies, K. J. A. The role of declining adaptive homeostasis in
ageing. *J. Physiol.* **595**, 7275–7309 (2017).

72. Schmauck-Medina, T. *et al.* New hallmarks of ageing: a 2022 Copenhagen
ageing meeting summary. *Aging* **14**, 6829–6839 (2022).

73. Cornacchia, D. & Studer, L. Back and forth in time: Directing age in iPSC-derived
lineages. *Brain Res.* **1656**, 14–26 (2017).

74. Bonilha, V. L. Age and disease-related structural changes in the retinal pigment
epithelium. *Clin Ophthalmol Auckl N Z* **2**, 413–424 (2008).

75. Chen, M. *et al.* Retinal pigment epithelial cell multinucleation in the aging eye – a
mechanism to repair damage and maintain homoeostasis. *Aging Cell* **15**, 436–445
(2016).

76. Tamori, Y. & Deng, W.-M. Compensatory cellular hypertrophy: the other strategy
for tissue homeostasis. *Trends Cell Biol* **24**, 230–237 (2014).

77. Bodenschatz, J. F. E. *et al.* Epithelial cells sacrifice excess area to preserve
fluidity in response to external mechanical stress. *Commun. Biol.* **5**, 855 (2022).

78. Figard, L., Wang, M., Zheng, L., Golding, I. & Sokac, A. M. Membrane Supply
and Demand Regulates F-Actin in a Cell Surface Reservoir. *Dev. Cell* **37**, 267–278
(2016).

79. Liao, M. *et al.* Multi-omics profiling of retinal pigment epithelium reveals
enhancer-driven activation of RANK-NFATc1 signaling in traumatic proliferative
vitreoretinopathy. *Nat. Commun.* **15**, 7324 (2024).

80. Zhang, Q. *et al.* Comparison of histologic findings in age-related macular
degeneration with RPE flatmount images. *Mol. Vis.* **25**, 70–78 (2019).

81. Trepat, X. *et al.* Viscoelasticity of human alveolar epithelial cells subjected to
stretch. *Am. J. Physiol.-Lung Cell. Mol. Physiol.* **287**, L1025–L1034 (2004).

82. Morris, T., Sue, E., Geniesse, C., Brieher, W. M. & Tang, V. W. Synaptopodin
stress fiber and contractomere at the epithelial junction. *J. Cell Biol.* **221**,
e202011162 (2022).

83. López-Gay, J. M. *et al.* Apical stress fibers enable a scaling between cell
mechanical response and area in epithelial tissue. *Science* **370**, eabb2169 (2020).

84. Bays, J. L., Campbell, H. K., Heidema, C., Sebbagh, M. & DeMali, K. A. Linking
E-cadherin mechanotransduction to cell metabolism through force-mediated
activation of AMPK. *Nat. Cell Biol.* **19**, 724–731 (2017).

85. Gutmann, M. *et al.* Beyond comparison: Brillouin microscopy and AFM-based
indentation reveal divergent insights into the mechanical profile of the murine
retina. *J. Phys.: Photonics* **6**, 035020 (2024).

86. Zhang, P., Medwig-Kinney, T. N. & Goldstein, B. Architecture of the cortical
actomyosin network driving apical constriction in C. elegans. *J. Cell Biol.* **222**,
e202302102 (2023).

87. Brückner, B. R., Pietuch, A., Nehls, S., Rother, J. & Janshoff, A. Ezrin is a Major
Regulator of Membrane Tension in Epithelial Cells. *Sci. Rep.* **5**, 14700 (2015).

88. Goodwin, K. *et al.* Cell–cell and cell–extracellular matrix adhesions cooperate to
organize actomyosin networks and maintain force transmission during dorsal
closure. *Mol. Biol. Cell* **28**, 1301–1310 (2017).

89. Maruthamuthu, V., Sabass, B., Schwarz, U. S. & Gardel, M. L. Cell-ECM traction
force modulates endogenous tension at cell–cell contacts. *Proc National Acad Sci*
108, 4708–4713 (2011).

90. Valencia-Expósito, A., Khalilgharibi, N., Millán, A. M.-A., Mao, Y. & Martín-
Bermudo, M. D. Local weakening of cell-extracellular matrix adhesion triggers
basal epithelial tissue folding. *EMBO J.* **44**, 2002–2024 (2025).

91. Zuidema, A., Wang, W. & Sonnenberg, A. Crosstalk between Cell Adhesion
Complexes in Regulation of Mechanotransduction. *BioEssays* **42**, 2000119 (2020).

92. Mui, K. L., Chen, C. S. & Assoian, R. K. The mechanical regulation of integrin–
cadherin crosstalk organizes cells, signaling and forces. *J Cell Sci* **129**, 1093–
1100 (2016).

93. Weber, G. F., Bjerke, M. A. & DeSimone, D. W. Integrins and cadherins join
forces to form adhesive networks. *J Cell Sci* **124**, 1183–93 (2011).

94. Rooij, J. de, Kerstens, A., Danuser, G., Schwartz, M. A. & Waterman-Storer, C.
M. Integrin-dependent actomyosin contraction regulates epithelial cell scattering.
J. Cell Biol. **171**, 153–164 (2005).

95. Herant, M., Heinrich, V. & Dembo, M. Mechanics of neutrophil phagocytosis:
behavior of the cortical tension. *J. Cell Sci.* **118**, 1789–1797 (2005).

96. Herant, M., Heinrich, V. & Dembo, M. Mechanics of neutrophil phagocytosis:
experiments and quantitative models. *J. Cell Sci.* **119**, 1903–1913 (2006).

97. Vorselen, D., Labitigan, R. L. D. & Theriot, J. A. A mechanical perspective on
phagocytic cup formation. *Curr Opin Cell Biol* **66**, 112–122 (2020).

98. Hoijman, E. *et al.* Cooperative epithelial phagocytosis enables error correction in
the early embryo. *Nature* **590**, 618–623 (2021).

99. Li, J. X. H., Tang, V. W. & Brieher, W. M. Actin protrusions push at apical
junctions to maintain E-cadherin adhesion. *Proc. Natl. Acad. Sci.* **117**, 432–438
(2020).

100. Gao, H. & Hollyfield, J. G. Aging of the human retina. Differential loss of neurons
and retinal pigment epithelial cells. *Invest Ophth Vis Sci* **33**, 1–17 (1992).

101. Curcio, C. A., Millican, C. L., Allen, K. A. & Kalina, R. E. Aging of the human
photoreceptor mosaic: evidence for selective vulnerability of rods in central retina.
Investig. Ophthalmol. Vis. Sci. **34**, 3278–96 (1993).

102. Curcio, C. A. & Drucker, D. N. Retinal ganglion cells in Alzheimer's disease and
aging. *Ann. Neurol.* **33**, 248–257 (1993).

103. Eliasieh, K., Liets, L. C. & Chalupa, L. M. Cellular Reorganization in the Human
Retina during Normal Aging. *Investig. Opthalmology Vis. Sci.* **48**, 2824 (2007).

104. Tse, J. R. & Engler, A. J. Preparation of Hydrogel Substrates with Tunable
Mechanical Properties. *Curr. Protoc. Cell Biol.* **47**, 10.16.1-10.16.16 (2010).

105. Przybyla, L., Lakins, J. N., Sunyer, R., Trepat, X. & Weaver, V. M. Monitoring
developmental force distributions in reconstituted embryonic epithelia. *Methods*
94, 101–113 (2016).

106. Parinot, C., Rieu, Q., Chatagnon, J., Finnemann, S. C. & Nandrot, E. F. Large-
Scale Purification of Porcine or Bovine Photoreceptor Outer Segments for
Phagocytosis Assays on Retinal Pigment Epithelial Cells. *J. Vis. Exp.* (2014)
doi:10.3791/52100.

107. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.
Nat. Methods **9**, 676–682 (2012).

108. Ewels, P. A. *et al.* The nf-core framework for community-curated bioinformatics
pipelines. *Nat. Biotechnol.* **38**, 276–278 (2020).

109. Tommaso, P. D. *et al.* Nextflow enables reproducible computational workflows.
Nat. Biotechnol. **35**, 316–319 (2017).

110. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

111. Kolde, R. *Pheatmap: Pretty Heatmaps*. (2018).

112. Raudvere, U. *et al.* g:Profiler: a web server for functional enrichment analysis
and conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, W191–W198
(2019).

113. Janes, K. A. An analysis of critical factors for quantitative immunoblotting. *Sci.*
Signal. **8**, rs2 (2015).

114. LAEMMLI, U. K. Cleavage of Structural Proteins during the Assembly of the
Head of Bacteriophage T4. *Nature* **227**, 680–685 (1970).

115. Brunelle, J. L. & Green, R. One-dimensional SDS-polyacrylamide gel
electrophoresis (1D SDS-PAGE). *Methods Enzym.* **541**, 151–9 (2014).

116. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist
algorithm for cellular segmentation. *Nat. Methods* **18**, 100–106 (2021).

117. Bi, D., Lopez, J. H., Schwarz, J. M. & Manning, M. L. A density-independent
rigidity transition in biological tissues. *Nat. Phys.* **11**, 1074–1079 (2015).

118. Park, J.-A. *et al.* Unjamming and cell shape in the asthmatic airway epithelium.
Nat. Mater. **14**, 1040–1048 (2015).

1429

1430

a

POS fragments

RPE

t = 0

Bound

$h_{cut-off}$

Internalized

t = 4 h

b

iRPE   dr-iRPE

Apical

Basal

Side

Internalized

20 μm

5 μm

F-Actin   POS

c

Internalized fragments (x 10³)

**

iRPE   dr-iRPE

d

Avg. fragments volume (μm³)

**

iRPE   dr-iRPE

e

iRPE   dr-iRPE

Side

10 μm

Standard culture

iRPE   dr-iRPE

F-Actin

Side

10 μm

During POS internalization

f

iRPE   dr-iRPE

Side

Side

Myosin IIB   POS-FITC

5 μm

g

iRPE   dr-iRPE

Top

Myosin IIB   POS-FITC

5 μm

h

iRPE   dr-iRPE

F-Actin

5 μm

Top

Apical actin area (x10³ μm²)

*

iRPE   dr-iRPE

i

iRPE   dr-iRPE

Top

20 μm

Side

F-Actin

10 μm

iRPE

dr-iRPE

Mechanical Homeostasis

Apico-lateral plasticity

iRPE

dr-iRPE

Structural reinforcement

POS fragment   Actomyosin network