Optimization of Bio-Orthogonal Non-Canonical Amino acid Tagging (BONCAT) for effective low-disruption labelling of Arabidopsis proteins *in vivo*

Nicholas Hassan[1], Shelly Braun[2], Mohana Talasila[1], Richard P. Fahlman[2] and R. Glen Uhrig[1,2#]

[1] Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada
[2] Department of Biochemistry, University of Alberta, Edmonton, Alberta, Canada

[#] Senior and corresponding author:

Dr. R. Glen Uhrig
Department of Biological Sciences,
University of Alberta,
Edmonton, Alberta, Canada, T6G 2E9
Email: ruhrig@ualberta.ca
Phone: +1 780 492 3088
Fax : +1 780 492-9234ruhrig@ualberta.ca

Running Title: Developing Bio-Orthogonal Non-Canonical Amino acid Tagging (BONCAT) for *in-vivo* labelling of newly synthesized plant proteins in plants

Key Words: BONCAT, Click Chemistry, Mass Spectrometry, Quantitative Proteomics

ABSTRACT

Plants require robust responses in protein synthesis to adapt to variable environmental

conditions. Measurement of newly synthesized proteins has been successfully facilitated with

Bio-Orthogonal Non-Canonical Amino acid Tagging (BONCAT) across a multitude of organisms.

Here, we use non-canonical amino acids (NCAAs) L-azidohomoalanine (AHA) or L-

homopropargylglycine (HPG) incorporation in place of methionine residues into the actively

translating *Arabidopsis* proteome, allowing for that subset of proteins to be enriched for mass

spectrometry quantification. Although this technique has seen occasional use in plants,

optimization of the protocol to maximize functionality while minimizing organismal stress has not

yet been established. Here, we provide evidence for successful implementation through the

liquid immersion of seedlings in AHA or HPG-containing media that functions with significantly

lower concentrations than the literature standard. Our approach splits acute exposure and

incorporation phase of labelling to mitigate potential negative impacts of prolonged NCAA

exposure without compromising effective enrichment capacity, and demonstrate that this results

in an unperturbed growth phenotype for AHA-treated seedlings. Finally, we demonstrate the

capacity of this modified approach to enrich newly synthesized proteins from the whole

proteome under standard stress conditions. These improvements allow for a broader use of

BONCAT technologies in molecular plant research, affording a deeper understanding of the

newly synthesized proteome without negatively impacting plant health.

INTRODUCTION

Balancing protein synthesis and degradation defines proteostasis in all cells, which in turn defines the function of all organisms. Rapid adjustment of protein levels and proteome plasticity is essential for survival in a changing environment, especially for plants, which are carefully tuned to their environmental changes. Defining newly synthesized proteins, or the nascent proteome, is crucial to understanding plant cell regulation and adaptation to environmental stressors. However, conventional strategies for labelling the newly synthesized proteome, such as radiolabelling, lack depth of analysis (Tivendale et al., 2021), while characterization of the nascent proteome by enrichment-based mass spectrometry (MS) approaches can better quantify changes in lower abundant proteins. To fill this niche, Bio-Orthogonal Non-Canonical Amino acid Tagging (BONCAT), has been developed to label nascent proteins with a methionine analog, allowing for enrichment of newly synthesized proteins (Dieterich et al., 2006; Parker and Pratt, 2020).

BONCAT utilizes a combination of azide and alkyne groups to form irreversible conjugations via a copper-catalyzed azide-alkyne cycloaddition (CuAAC; reviewed in Chen et al., 2023). To employ this in a biological setting, unnatural methionine analogs L-azidohomoalanine (AHA, azide) or L-homopropargylglycine (HPG, alkyne) are supplied to live organisms. These non-canonical amino acids (NCAAs) are incorporated into proteins during synthesis by the relatively promiscuous methionine tRNA (Figure 1A; Bagert et al., 2014). Labelled proteins are then extracted and "clicked" to the opposing azide or alkyne conjugated to biotin, effectively biotinylating newly synthesized proteins to facilitate enrichment via streptavidin coupled matrix for subsequent quantification via liquid chromatography (LC)-MS/MS. While BONCAT was initially deployed in mammalian cells, it has since found use in a wide variety of organisms, such as live mice, bacteria, and archaea (Dieterich et al., 2006; Kern and Ferreira-Cerca, 2022; Lindivat et al., 2021; Saleh et al., 2019). However, unlike other organisms, BONCAT implementation in plants has not been fully realized.

In the model plant *Arabidopsis thaliana* (Arabidopsis), BONCAT was first implemented in 2017. Here, AHA and HPG were applied to Arabidopsis seedlings through the flooding of agar plates with 1 mM AHA (Glenn et al., 2017). This yielded thousands of enriched proteins after only a few hours of exposure. However, subsequent endeavours found that Arabidopsis reacts metabolically to NCAAs similar to excess methionine, stunting plant growth and/or impacting plant health during prolonged exposure (Tivendale et al., 2021). Since then, only a handful of studies have utilized BONCAT in plants, with protocol updates generally focused on seedling exposure time to high concentrations of NCAAs (Li et al., 2023; Linster et al., 2022; Lyu et al.,

2023; Xie et al., 2023; Yu et al., 2020). Together, these studies indicate that optimization of

BONCAT in plants remains incomplete, as a wider variety of experimental parameters need to

be tested to strike a balance between effective NCAA labelling concentrations and minimal

disturbance plant health. Additionally, protocol simplification efforts remain unexplored,

including: a) using the same biotin-conjugated clicked extract for both anti-biotin blots and

enrichment for LC-MS/MS analysis to minimize variability and b) exploring the exclusivity of the

azide-alkyne cycloaddition, including eliminating the need for repeated precipitation or sample

processing prior to remove metabolites prior to "clicking" (Ma et al., 2018).

Here, we present an adapted protocol for BONCAT labelling of Arabidopsis seedlings in

liquid MS medium that allows for more reliable seedling exposure to, and removal of, AHA and

HPG (Figure 1B). Specifically, we demonstrate that the titration of AHA and HPG as well as

optimization of labelling time an effective approach to balance labelling while minimizing

exposure to NCAAs. Lastly, we perform extensive seedling viability assays to ensure minimal

disruption of plant growth using the modified protocol, and evaluate the newly synthesized

proteome using the modified protocol under high and low acute salt and mannitol stress via LC-

MS/MS.

RESULTS AND DISCUSSION

AHA and HPG label Arabidopsis seedlings at 50 µM or less using short application

pulses

To determine concentrations of AHA and HPG suitable for application pulses, 5 d-old

seedlings were acclimated to liquid medium overnight before being exposed to 0, 10, 25, 50,

100, or 250 µM AHA or HPG for 30 min (see Materials and Methods). Washed seedlings were

then allowed to incorporate NCAAs into the nascent proteome for an additional 3 h before whole

plants were flash frozen and extracted. Copper-catalyzed click chemistry was then performed *in*

vitro to conjugate NCAA tags to biotin, and total proteome biotinylation was detected via

immunoblot. Anti-biotin immunoblotting against untagged controls was selected as the preferred

method for monitoring qualitative labelling efficiency, as whole proteome analysis may not detect

AHA tags that have a mass shift similar to salt adducts of methionine (Liu et al., 2022). Here,

AHA was found to have the best above-background labelling at all tested concentrations (Figure

2A). Based on these results, a concentration of 50 µM AHA or HPG was selected as an optimal

concentration due to clearly visible labelling with minimal required reagent given the known

metabolic stress of NCAAs (Tivendale et al., 2021).

We then performed time-course labelling to select ideal labelling periods. Labelling periods
were divided into acute exposure time (NCAA present in medium) and incorporation time (NCAA
removed from medium), with NCAA proteome incorporation assessed independently. This
revealed that increased exposure time increases labelling over the course of 24 h at both 50 µM
AHA and HPG, with above-background labelling visible within the first 15 min (Figure 2B).
Similarly, increased incorporation time with a set initial exposure of 30 min appears to increase
labelling for up to 6 h after removal of tags using an initial 50 µM of either NCAA, suggesting
residual NCAA likely remains present in the plant's tissues after wash and removal from the
medium, with that the remaining reagent continuing to be incorporated for several hours (Figure
2C). From this data, a 30 min exposure followed by 3 h of NCAA incorporation were selected as
optimal labelling times for 50 µM of either NCAA, with a shorter acute time to mitigate potential
negative effects of exposure on the seedlings. Interestingly, concentrations of AHA as low as 10
µM also showed increases in labelling across longer exposure times, but labelling stagnated
with prolonged incorporation (Figure 2D). Therefore, a NCAA concentration of 10 µM is likely too
low for excess NCAA to accumulate in plant tissue prior to its removal, but may be more suitable
for long-term exposure labelling, such as overnight incubations for pulse-chase assay.

Short exposure times prevent problems associated with prolonged exposure to AHA
Viability assays of NCAA treated Arabidopsis seedlings was performed by assessing fold-
change in root length as a proxy for plant growth, in order to measure the potential negative
effects of AHA and HPG. Here, 5 d-old Arabidopsis seedlings were transplanted onto 0.5x MS
agar plates containing AHA or HPG and allowed to grow for multiple days. All tested
concentrations from 10 to 250 µM AHA and HPG decreased plant root growth across a 3-day
exposure, with HPG being significantly worse than AHA, even at comparably low concentrations
(Figure 3A, Supplemental Figure 1A). However, with only 24 h of exposure, 10 µM AHA did not
show a significant reduction in root growth. As this concentration shows above-background
labelling within a few hours but does not continue to incorporate the NCAA after being washed
(Figure 2D), 10 µM AHA may find an appropriate use in pulse-chase assays, where its capacity
for 24-h acute exposure labelling could be used to label plants overnight for degradation
experiments the following day.
Differing approaches for *in vivo* AHA and HPG viability treatments have been deployed
previously. For example, use of 1 mM AHA or HPG to acutely flood Arabidopsis seedlings
growing on MS-agar plates revealed a negative impact on whole seedling growth and cell
culture viability, in addition to perturbed metabolism (Tivendale et al., 2021). However, other

eukaryotes demonstrate a more mixed or neutral response to NCAA exposure. For example, 2
mM AHA and 1 mM HPG have no negative viability effects on monkey and mouse-derived cells,
while HeLa cells exhibit perturbation responses to AHA that seem to be mitigated by the
presence of methionine in the growth medium (Bagert et al., 2014; Su Hui Teo et al., 2016;
Wisse et al., 2017). On the molecular level, mouse-derived cell lines exhibit moderate stress
signals (e.g. increased HEAT SHOCK PROTEIN 70 and 90 abundance) when treated with 50
µM AHA, but no significant increase in global protein ubiquitination and/or apoptosis / cell death
associated with a reduction in viability (Morey et al., 2021).

In contrast, Tivendale et al. (2021) found AHA to be more toxic than HPG, especially in plant
cell cultures. Bacteria have shown similar concentration-dependent sensitivity to NCAAs, but in
a manner more consistent with this study. AHA appears to be tolerable to log-phase *E. coli* at
higher concentrations than HPG, where HPG reduced bacterial growth at concentrations above
2.8 µM while AHA was tolerated up to the 200 µM range for prolonged exposure (Landor et al.,
2023). This effect differs slightly between prototrophs and auxotrophs, where bacteria unable to
synthesize essential compounds appear to be more negatively affected by HPG (Jecmen et al.,
2023). Both AHA and HPG mildly affect bacterial metabolism as well, but are not degraded as
the expected products of methionine (Steward et al., 2020).

It has been proposed that negative long-term effects of NCAAs on organism or cell growth
may be irrelevant if incorporation into nascent proteins and immediate toxicity does not occur
during the length of the experiment (Landor et al., 2023). To date, viability assays in plants have
examined organism or cellular responses over prolonged exposure times, which is not always
representative of the experimental conditions. Therefore, we repeated viability assays with a
more representative exposure (short acute exposure in liquid 0.5x MS), followed by seedling
washing and return to 0.5x MS - agar plates. All tested concentrations of AHA (up to 250 µM) did
not reduce plant growth compared to controls after a 30-min AHA exposure (Figure 3B-C,
Supplemental Figure 1B), suggesting that the long-term exposure growth defects are not a
result of immediate toxicity of the NCAA. In contrast, 50 µM and 100 µM concentrations of HPG
showed a slight reduction in growth compared to control under the same conditions (Figure 3C).
By limiting exposure to 30 min and allowing incorporation to continue for < 6 h, viability
concerns can be avoided without compromising in labelling efficiency, particularly in the use
case of AHA.

AHA-labelled proteins under salt and osmotic stress

To benchmark the quantitative labelling efficiency of the optimized protocol under different
experimental conditions, we applied two abiotic stress conditions during AHA labelling. The
experiment was performed with either high stress (150 mM salt or 300 mM mannitol) or low
stress (50 mM salt or 100 mM mannitol) conditions, as the effects of applied stressors may be
intensified by liquid submersion. Here, we find 549 proteins enriched in at least one condition
under high stress conditions, 797 under low stress conditions, with a greater percentage of
overlapping proteins between groups under lower stress versus high stress (55.1% vs. 35.1%;
Figure 4A). Hundreds of these proteins represent Class I hits, which we define as below
detection in the MS2 spectra of negative control samples (Figure 4B). Further principal
component analyses found AHA-treated samples cluster away from untreated (bead binding
control) samples, but distance between clusters of stressed and unstressed samples vary with
abiotic stress intensity (Figure 5A and B).

Additionally, using positive AHA-labeled control samples, we can disentangle stress specific
responses. Network analysis of dynamically changing salt and mannitol responsive proteins,
resolved multiple protein networks related to abiotic stimulus response (Figure 4C). Gene
ontology (GO) enrichment analysis further validated enrichment of abiotic stress response
biological processes under the stress conditions, including wax biosynthesis and water-related
regulation (Figure 4D, Supplemental Figure 2, Supplemental Figure 3). Consistent with prior
findings, responses to salt and mannitol at the protein level appear to differ from each other
(Figure 5C and D, Gallo et al., 2023).

SUMMARY

In this study, we established a simple and usable workflow for the implementation of
BONCAT in plants. Although 50 µM AHA was selected as the optimal labelling concentration,
the protocol is flexible and modifiable based on need, with modified protocols able to be tracked
for efficiency at multiple stages. Our BONCAT implementation showed an ability clearly define
stress and unstressed conditions in relatively short periods that mitigates potential negative
plant responses. This unveils outstanding new opportunities for applications where enrichment
of a subsect of the proteome is required for sufficient depth of analysis. Lastly, the potential for
NCAA-tag incorporation at concentrations substantially below the literature standard has
positive new implications for the use of BONCAT in plants moving forward.

MATERIALS AND METHODS
Plant growth conditions

Arabidopsis thaliana seeds (Col-0) were liquid sterilized by immersion in 70% (v/v) ethanol
for 2 min followed by 30% (v/v) bleach for 10 min. Sterilized seeds were vernalized in H_2O in the
dark at 4^oC for 3 d. Germination plates were prepared using half-strength MS basal salts
(Caisson Laboratories Inc. Murashige & Skoog MSP01), pH adjusted to 5.8 with aqueous KOH,
with 0.08% (w/v) agar (Caisson laboratories Inc. Phytoblend PTP01).

For click chemistry, vernalized seeds were stratified on plates and allowed to grow with
plates positioned vertically in 24 h light at 22^oC for 5 days. Approximately 100 mg of 5 d-old
seedlings were immersed in autoclaved half-strength MS media in 24-well plates and placed
under constant light for 16 h to acclimatize. MS media was removed and replaced with 50 µM
filter sterilized L-azidohomoalanine or L-homopropargylglycine (Vector Labs, CCT-1066 and
CCT-1076) from 0.1 M stocks dissolved in H_2O and diluted in 0.5x MS media. Seedlings were
then incubated for 30 min of exposure time in constant light before being washed once in 0.5x
MS and incubated in constant light in 0.5x MS for 3 hours of "incorporation" time. Labelled
seedlings were removed from media, rinsed in 0.5x MS, and quickly touch dried before being
flash frozen in liquid N_2 and stored at -80^oC until subsequent steps.

To perform root length assays, seeds were sown individually on 0.5x MS agar plates
positioned vertically in dark boxes with seeds at the light interface to emulate roots growing in
the dark and shoots growing in the light. Seedling growth conditions consisted of a 12 h
photoperiod, 22^oC and 100 µmol/m^{2}s light. After 5 d for the long-term reagent exposure,
seedlings were transplanted along the same light/dark interface to 0.5x MS plates containing 0,
10, 25, 50, or 250 µM AHA or HPG. For the short-term exposure, seedlings were placed into
liquid 0.5x MS overnight to equilibrate, then immersed for 30 min in 0.5x liquid MS containing 0,
10, 25, 50, or 250 µM AHA, before being washed in 0.5x MS and re-plated on to vertical plates.
Root length was recorded for 7 d using digital calipers, and fold-change in individual seedling
root length relative to day 0 (transplanting day) was plotted. Recording of root length stopped
after day 3 for the long-term exposure due to statistically significant differences between all
concentrations relative to controls.

Optimization of plant labelling

To determine optimal concentrations of label, variable concentrations of 10, 25, 50, 100, or
250 µM of AHA or HPG were applied as above. Optimal concentrations for labelling were
chosen based on a combination of intensity of labelled protein present on streptavidin
immunoblots and minimal observed growth retardation on 0.5x MS agar containing the same
concentration of label. For time-course experimentation, acute exposure times were varied

between 15 min, 30 min, or 1, 2, 4, or 22 h. Incorporation times were varied between 30 minutes

or 1, 2, 3, 6, or 24 h. Optimal incubation times were chosen based on minimal required times to

achieve effective labelling at the desired concentration as visualized via immunoblot, and

verified to cause insignificant or no growth inhibition to the seedlings based on viability studies.

Protein extraction and quantification

Flash frozen tissue was ground using Geno/Grinder (SPEX SamplePrep) for 30 s at 1200

rpm with 3mm glass beads and extracted in either 50mM HEPES (pH 7.5) with 1% (w/v) SDS or

50 mM Tris (pH 8.0) with 1% (w/v) SDS, at a ratio of 1:2 w/v ground tissue to extraction buffer.

Following optimization, HEPES was selected as the extraction buffer for all samples due to

better click labelling observed. Samples were heated at $95^{o}C$ for 10 min and 1000 rpm on

Eppendorf ThermoMixer F2.0, cooled, and centrifuged at 16,000 xg for 10 min at room

temperature to remove tissue particulates. Protein amounts were quantified using Bradford

assay with samples diluted to 0.2% (w/v) SDS against BSA controls.

Click chemistry reaction

A total of 250 μg of protein from each sample was diluted in extraction buffer with 1% (w/v) or

no SDS to a volume of 346 μL ensuring a final concentration of 0.5% (w/v) SDS to maintain

protein solubility. To each reaction, the following was added: 8 μL of 50 mM tris(2-

carboxyethyl)phosphine (TCEP)-HCl (Sigma Cat. No: 51805-45-9; diluted from 1 M stock in H_2O

same day of use), 30 μL of 1.7 mM tris[(1-benzyl-1H-1,2,3-triazol-4-yl)methyl]amine (TBTA)

(Vector Labs Cat. No: CCT-1061; in 20% (v/v) dimethyl sulfoxide (DMSO) and 80% (v/v) tert-

butanol, stored at room temperature as in Ma et al., 2018), 8 μL of CuSO4 (Sigma Cat. No:

451657-10G; in H_2O, stored at $-20^{o}C$), and 8 μL of 5mM biotin-PEG3-azide (Vector Labs Cat.

No: CCT-AZ104; for HPG samples, in DMSO) or biotin-PEG4-alkyne (Vector Labs Cat. No:

CCT-TA105; for AHA samples, in DMSO). Click reactions were allowed to proceed for 1 h at

room temperature, shaking at 700 rpm and protected from light. Reactions were then quenched

via chelation with 20 μL of 0.5 M ethylenediaminetetraacetic acid (EDTA) (pH 8.0, stored at

$4^{o}C$). Labelling validation was determined via streptavidin immunoblot using 6 μg of protein run

on 10% SDS-PAGE gels transferred to 0.45 μM polyvinylidene fluoride (PVDF) membrane.

Blots were blocked in 5% fat-free milk in Tris-buffered saline (TBS) for 1 h, at which point HRP-

conjugated streptavidin (Thermo Scientific Cat. No. N100) diluted 1:2000 in Tris-buffered saline

with 0.1% Tween-20 (TBST) was added for an additional hour. Membranes were washed in

TBST and visualized using Clarity Western ECL Substrate (Bio-Rad Cat No. 170-5060).
Colloidal Coomassie Blue-stained gels were used as loading controls.

Enrichment of clicked proteins

Excess biotin and small molecules were removed from clicked samples using Cytiva PD
Spintrap G-25 columns (Cat. No: 28918004). Samples were bound to Cytiva streptavidin
Sepharose beads (Cytiva Cat. No. 17511301); all depicted experiments proceeded using non-
magnetic Sepharose beads. All beads were initially washed 3 times in 50 mM HEPES, pH 7.5,
for 1 min while rotating at room temperature. For all steps, supernatant was removed by
centrifugation for 1 min at 600 xg. Up to 200 µg of flow-through protein from the Spintrap
columns was loaded onto equilibrated streptavidin beads and allowed to bind while rotating for
16 hours at 4^{o}C. The supernatant was removed and retained at -20^{o}C to assess capture
efficiency.

Capture efficiency was checked before digestion via streptavidin-detected dot blot
comparison of clicked protein, Spintrap column flow-through, and supernatant after overnight
bead binding. After confirmation of a signal reduction of more than 70% in the supernatant after
binding, beads were washed sequentially with the following solutions at 1mL each: 50 mM
HEPES pH 7.5 (2x washes), 1 M potassium chloride, ice-cold 100 mM sodium carbonate (2x
washes), 2 M urea in 50 mM Tris-HCl pH 8.0 (2x wash), and 50 mM triethylammonium
bicarbonate (TEAB; Sigma Cat. No: 15715-58-9) in HPLC-grade water (3x washes). All washes
were completed at room temperature with rotation for 3 min.

Peptides were generated through on bead digestion using trypsin diluted 1:100 in 50 mM
TEAB overnight at 37^{o}C; the supernatant was removed before the beads were washed with 50
mM TEAB and the supernatant pooled with previously extracted peptides. The extract was
reduced with 10 mM DTT at 95^{o}C for 5 min while shaking at 450 rpm, alkylated with 30 mM
iodoacetamide (IA) at room temperature for 30 min, and the reaction quenched with 10 mM DTT
for 10 minutes. Samples were then acidified to 0.5% (v/v) TFA to a pH of less than 2 and
centrifuged at 16,000 xg for 1 min before the supernatant was removed and dried down.
Peptides were re-suspended in 3% (v/v) acetonitrile / 0.1% (v/v) formic acid for desalting via
ZipTip (Sigma Cat. No. ZTC18S960) before being dried and stored at -80^{o}C. Desalted peptides
were the re-suspended in 3% (v/v) acetonitrile / 0.1% (v/v) formic acid for mass spectrometry
analysis.

LC MS/MS and Data Analysis

Peptides were analyzed using a FAIMSpro mounted Fusion Lumos Orbitrap mass spectrometer (Thermo Fisher Scientific) in data independent acquisition (DIA) mode. Peptides were injected using Easy-nLC 1200 system (LC140; Thermo Fisher Scientific) and an Acclaim PepMap 100 C18 trap column (Cat# 164750; Thermo Fisher Scientific) followed by a 50 cm Easy-Spray PepMap C18 analytical column (ES903; Thermo Fisher Scientific) warmed to 50 °C. Peptides were eluted at 0.3 µl/min using a segmented solvent B gradient of 0.1% (v/v) formic acid in 80% (v/v) acetonitrile from 4 to 41% solvent B (0–60min). The FAIMSpro was used with a fixed gas flow of 3.5 L/min with a 3-CV setting of -30, -50, -70. A positive ion spray voltage of 2.3 kV was used with an ion transfer tube temperature of 300 °C and an RF lens setting of 40%. Direct DIA acquisition was performed as previously described (Mehta et al., 2022). Full scan MS[1] spectra (350–1400 m/z) were acquired with a resolution of 120,000 at 200 m/z with a normalized automatic gain control (AGC) target of 100% and IT set to automatic. Fragment spectra were acquired at resolution of 30,000 across 28 38.5 m/z windows overlapping by 1 m/z using a dynamic maximum injection time and an AGC target value of 2000%, with a minimum number of desired points across each peak set to 6. Higher-energy collisional dissociation fragmented was performed using a fixed 27% fragmentation energy. Downstream data analysis of dDIA acquisitions was performed using Spectronaut ver. 19 (Biognosys AG) with default settings as previously described (Mehta et al., 2022), with all searches made against a custom-made decoy (reversed) version of the Arabidopsis protein database from Araport 11 (27,533 protein encoding genes; ver. 2022-09-14). Briefly, search parameters included the following: trypsin digest permitting two missed cleavages, fixed modifications (carbamidomethyl (C)), variable modifications (oxidation (M)) and a peptide spectrum match, peptide and protein false discovery threshold of 0.01.

Bioinformatic Data Analysis

Significantly changing proteins (Log$_2$FC of MS2 spectra > 0.58 over negative control) were pooled with Class I protein hits (absent in MS2 of negative controls) to form the foreground with a background of all identified proteins (Supplemental Table 1). Gene Ontology (GO) enrichments were performed using the Ontologizer (Bauer et al., 2008; http://ontologizer.de/, Supplemental Table 2) using a Benjamini-Hochberg parent-child union with an analysis cutoff of p > 0.01, and visualized in R 4.5.0. Proteins below differential abundance in both positive and negative controls was used for network analysis of stress-exclusive proteins via the StringApp plugin for Cytoscape v. 3.10.3, with an edge score cut-off of 0.5(Doncheva et al., 2019,

321 https://cytoscape.org/). STRING Enrichment using *StringApp* was used to assign gene
322 ontologies. PCA analysis was performed using *SRplot* (Tang et al., 2023), with other
323 visualizations produced using a combination of *ggplot2*, Biorender (https://www.biorender.com/),
324 GraphPad Prism v. 8.4.3 (https://www.graphpad.com/), and Affinity Designer v. 1.10.5
325 (https://affinity.serif.com/en-us/designer/).
326

327 **ACKNOWLEDGEMENTS**
328 The authors would like to thank the Canadian Foundation for Innovation (CFI) and Natural
329 Sciences and Engineering Council of Canada (NSERC) for funding this work. We would also
330 like to thank Jack Moore and the Alberta Proteomics and Mass Spectrometry Facility (APM) for
331 mass spectrometry assistance and upkeep.
332

AUTHOR CONTRIBUTIONS
Nicholas Hassan: Investigation, Conceptualization, Methodology, Data curation, Formal
analysis, Visualization, Writing—original draft. *Shelly Braun*: Writing—review & editing,
Conceptualization. *Mohana Talasila*: Formal analysis, Data curation, Visualization. *Richard
Fahlman*: Supervision, Writing—review & editing, Material Resources *R. Glen Uhrig*:
Supervision, Conceptualization, Writing—review & editing, Funding acquisition.
339

340 **DATA AVAILABILITY**
341 Raw data have been deposited to the ProteomeExchange Consortium
342 (http://proteomecentral.proteomexchange.org) *via* the PRoteomics IDEntification Database
343 (PRIDE; https://www.ebi.ac.uk/pride/) partner repository with the data set identifiers PXD064210
344

REFERENCES
Bagert, J.D., Xie, Y.J., Sweredoski, M.J., Qi, Y., Hess, S., Schuman, E.M., Tirrell, D.A., 2014.
Quantitative, time-resolved proteomic analysis by combining bioorthogonal noncanonical
amino acid tagging and pulsed stable isotope labeling by amino acids in cell culture.
Molecular and Cellular Proteomics 13, 1352–1358.
https://doi.org/10.1074/mcp.M113.031914

Chen, M.M., Kopittke, P.M., Zhao, F.J., Wang, P., 2023. Applications and opportunities of click
chemistry in plant science. Trends Plant Sci 29, 167–178.
https://doi.org/10.1016/j.tplants.2023.07.003

Dieterich, D.C., Link, A.J., Graumann, J., Tirrell, D.A., Schuman, E.M., Sharpless, K.B., 2006.
Selective identification of newly synthesized proteins in mammalian cells using

bioorthogonal noncanonical amino acid tagging (BONCAT). National Academy of Sciences
103, 9482–9487.

Doncheva, N.T., Morris, J.H., Gorodkin, J., Jensen, L.J., 2019. Cytoscape StringApp: Network
Analysis and Visualization of Proteomics Data. J Proteome Res 18, 623–632.
https://doi.org/10.1021/acs.jproteome.8b00702

Gallo, M.C.R., Li, Q., Talasila, M., Uhrig, R.G., 2023. Quantitative Time-Course Analysis of
Osmotic and Salt Stress in Arabidopsis thaliana Using Short Gradient Multi-CV FAIMSpro
BoxCar DIA. Molecular and Cellular Proteomics 22.
https://doi.org/10.1016/j.mcpro.2023.100638

Glenn, W.S., Stone, S.E., Ho, S.H., Sweredoski, M.J., Moradian, A., Hess, S., Bailey-Serres, J.,
Tirrell, D.A., 2017. Bioorthogonal noncanonical amino acid tagging (BONCAT) enables
time-resolved analysis of protein synthesis in native plant tissue. Plant Physiol 173, 1543–
1553. https://doi.org/10.1104/pp.16.01762

Jecmen, T., Tuzhilkin, R., Sulc, M., 2023. Photo-Methionine, Azidohomoalanine and
Homopropargylglycine Are Incorporated into Newly Synthesized Proteins at Different Rates
and Differentially Affect the Growth and Protein Expression Levels of Auxotrophic and
Prototrophic E. coli in Minimal Medium. Int J Mol Sci 24.
https://doi.org/10.3390/ijms241411779

Kern, M., Ferreira-Cerca, S., 2022. Differential Translation Activity Analysis Using Bioorthogonal
Noncanonical Amino Acid Tagging (BONCAT) in Archaea. Ribosome Biogenesis: Methods
and Protocols 229–246. https://doi.org/10.1007/978-1-0716-2501-9_14

Landor, L.A.I., Bratbak, G., Larsen, A., Tjendra, J., Våge, S., 2023. Differential toxicity of
bioorthogonal non-canonical amino acids (BONCAT) in Escherichia coli. J Microbiol
Methods 206. https://doi.org/10.1016/j.mimet.2023.106679

Li, X., Chai, X., Lyu, H.N., Fu, C., Tang, H., Shi, Q., Wang, J., Xu, C., 2023. Chemical
proteomics reveals an ISR-like response elicited by salicylic acid in Arabidopsis. New
Phytologist. https://doi.org/10.1111/nph.18646

Lindivat, M., Bratbak, G., Larsen, A., Hess-Erga, O.K., Hoell, I.A., 2021. Flow Cytometric
Analysis of Bacterial Protein Synthesis: Monitoring Vitality After Water Treatment. Front
Microbiol 12. https://doi.org/10.3389/fmicb.2021.772651

Linster, E., Forero Ruiz, F.L., Miklankova, P., Ruppert, T., Mueller, J., Armbruster, L., Gong, X.,
Serino, G., Mann, M., Hell, R., Wirtz, M., 2022. Cotranslational N-degron masking by
acetylation promotes proteome stability in plants. Nat Commun 13.
https://doi.org/10.1038/s41467-022-28414-5

Liu, C., Wong, N., Watanabe, E., Hou, W., Biral, L., Decastro, J., Mehdipour, M., Aran, K.,
Conboy, M.J., Conboy, I.M., 2022. Mechanisms and Minimization of False Discovery of
Metabolic Bioorthogonal Noncanonical Amino Acid Proteomics. Rejuvenation Res 25, 95–
109. https://doi.org/10.1089/rej.2022.0019

Lyu, H.N., Fu, C., Chai, X., Gong, Z., Zhang, J., Wang, Jiaqi, Wang, Jigang, Dai, L., Xu, C.,
2023. Systematic thermal analysis of the Arabidopsis proteome: Thermal tolerance,

organization, and evolution. Cell Syst 14, 883-894.e4.
https://doi.org/10.1016/j.cels.2023.08.003

Ma, Y., McClatchy, D.B., Barkallah, S., Wood, W.W., Yates, J.R., 2018. Quantitative analysis of
newly synthesized proteins. Nat Protoc 13, 1744–1762. https://doi.org/10.1038/s41596-
018-0012-y

Mehta, D., Scandola, S., Uhrig, R.G., 2022. BoxCar and Library-Free Data-Independent
Acquisition Substantially Improve the Depth, Range, and Completeness of Label-Free
Quantitative Proteomics. Anal Chem 94, 793–802.
https://doi.org/10.1021/acs.analchem.1c03338

Morey, T.M., Esmaeili, M.A., Duennwald, M.L., Rylett, R.J., 2021. SPAAC Pulse-Chase: A Novel
Click Chemistry-Based Method to Determine the Half-Life of Cellular Proteins. Front Cell
Dev Biol 9. https://doi.org/10.3389/fcell.2021.722560

Parker, C.G., Pratt, M.R., 2020. Click Chemistry in Proteomic Investigations. Cell 180, 605–632.
https://doi.org/10.1016/j.cell.2020.01.025

Saleh, A.M., Jacobson, K.R., Kinzer-Ursem, T.L., Calve, S., 2019. Dynamics of Non-Canonical
Amino Acid-Labeled Intra- and Extracellular Proteins in the Developing Mouse. Cell Mol
Bioeng 12, 495–509. https://doi.org/10.1007/s12195-019-00592-1

Steward, K.F., Eilers, B., Tripet, B., Fuchs, A., Dorle, M., Rawle, R., Soriano, B.,
Balasubramanian, N., Copié, V., Bothner, B., Hatzenpichler, R., 2020. Metabolic
Implications of Using BioOrthogonal Non-Canonical Amino Acid Tagging (BONCAT) for
Tracking Protein Synthesis. Front Microbiol 11. https://doi.org/10.3389/fmicb.2020.00197

Su Hui Teo, C., Serwa, R.A., O'Hare, P., 2016. Spatial and Temporal Resolution of Global
Protein Synthesis during HSV Infection Using Bioorthogonal Precursors and Click
Chemistry. PLoS Pathog 12. https://doi.org/10.1371/journal.ppat.1005927

Tang, D., Chen, M., Huang, X., Zhang, Guicheng, Zeng, L., Zhang, Guangsen, Wu, S., Wang,
Y., 2023. SRplot: A free online platform for data visualization and graphing. PLoS One 18.
https://doi.org/10.1371/journal.pone.0294236

Tivendale, N.D., Fenske, R., Duncan, O., Millar, A.H., 2021. In vivo homopropargylglycine
incorporation enables sampling, isolation and characterization of nascent proteins from
Arabidopsis thaliana. The Plant Journal 107, 1260–1276.
https://doi.org/https://doi.org/10.1111/tpj.15376

Wisse, L.E., Penning, R., Zaal, E.A., van Berkel, C.G.M., ter Braak, T.J., Polder, E., Kenney,
J.W., Proud, C.G., Berkers, C.R., Altelaar, M.A.F., Speijer, D., van der Knaap, M.S., Abbink,
T.E.M., 2017. Proteomic and metabolomic analyses of vanishing white matter mouse
astrocytes reveal deregulation of ER functions. Front Cell Neurosci 11.
https://doi.org/10.3389/fncel.2017.00411

Xie, Z., Zhao, S., Li, Ying, Deng, Y., Shi, Y., Chen, X., Li, Yue, Li, H., Chen, C., Wang, X., Liu, E.,
Tu, Y., Shi, P., Tong, J., Gutierrez-Beltran, E., Li, J., Bozhkov, P. V., Qian, W., Zhou, M.,
Wang, W., 2023. Phenolic acid-induced phase separation and translation inhibition mediate
plant interspecific competition. Nat Plants 9, 1481–1499. https://doi.org/10.1038/s41477-
023-01499-6

Yu, H., Kong, X., Huang, H., Wu, W., Park, J., Yun, D.J., Lee, B. ha, Shi, H., Zhu, J.K., 2020.
STCH4/REIL2 Confers Cold Stress Tolerance in Arabidopsis by Promoting rRNA
Processing and CBF Protein Translation. Cell Rep 30, 229-242.e5.
https://doi.org/10.1016/j.celrep.2019.12.012

 FIGURES

Figure 1. Visualization of BONCAT technology. A) Incorporation mechanism of AHA and
HPG into the nascent proteome. B) Workflow for click labelling. Whole seedlings are
acclimatized to liquid 0.5X MS media for 24 hours before treatment with AHA or HPG. Acute
exposure time in the modified protocol is limited to 30 minutes, followed by 3 hours of
incorporation time. Protein is extracted from seedlings for click chemistry in vitro, followed by
visualization via anti-biotin blot to determine qualitative labelling before streptavidin enrichment
for LC-MS/MS.

Figure 2. Streptavidin-detected anti-biotin blots of biotin-clicked Arabidopsis whole
lysate samples. Beneath each blot is a cropped segment of a Coomassie Blue gel loading
control. Each experiment was performed in triplicate, but single replicates are shown for
simplicity. A) Titration of AHA and HPG concentrations with a 30-minute acute exposure and 3-
hour incorporation time. B) 50µM AHA and HPG tested with exposure times varying from 15
minutes to 22 hours, all followed by a 3-hour incorporation. C) 50µM AHA and HPG tested with
incorporation times varying from 30 minutes to 24 hours, each preceded with a 30-minute
exposure time. D) Low concentration AHA (10µM) test of varying incorporation times with a 30-
minute exposure (left) or varying exposure times with a 3-hour incorporation (right).

Figure 3. Viability assays of Arabidopsis seedlings exposed to methionine analogs.
Average fold change in root length of seedlings with standard deviation error bars is plotted, and
t-tests between experimental and control groups were performed with p-value significance <
0.05 shown. A) n = 72 5 d (days)-old seedlings per condition were transplanted to agar plates
containing AHA or HPG for 1-3 d to simulate extensive exposure times. High concentrations
after 3 days (left), low concentrations after 3 days (middle) and low concentrations after 24
hours (right) are shown. B) n = 54 seedlings per condition were acclimatized to liquid media for
24 h, then exposed for 30 min in AHA- or HPG-containing liquid media before being returned to
solid medium and grown for 7 d. C) Snapshot of days 5 and 7 for all conditions from
summarized data in 2B to show significant changes.

Figure 4. Summary mass spectrometry data for 50µM AHA-labelled 3-hour salt or
mannitol stressed Arabidopsis seedlings. A) Venn diagrams of valid (>1.5X FC or below
detection in negative control) protein groups between positive control (AHA treated, unstressed)
and stress conditions in either high stress (150mM salt and 300mM mannitol, left, n = 549) or
low stress (50mM salt, 100mM mannitol, right, n = 797) conditions. B) Exclusively Class I hits

from 1A that are absent in negative controls (n = 147 and n = 300. C) STRING network of valid

protein hits enriched in the low stress salt and/or mannitol conditions that were below detection

or below threshold in both positive and negative controls. Only nodes with >2 edges were kept.

Increasing line thickness is edge score ranging from 0.5-1. D) Gene ontology biological

processes (Benjamini-Hochberg parent-child union) for high stress (left, p < 0.0025) and low

stress (right, p < 0.005) conditions. All proteins enriched in one condition over negative controls

were included. Highlighted in red are terms directly related to stress response.

Figure 5. Proteomics data representation of salt and mannitol stressed AHA-treated

seedlings. A) PCA plot of high stress (150mM salt, orange, and 300mM mannitol, purple) 50µM

AHA-treated seedlings, as well as control treated (blue) and control untreated (grey) samples.

Elipses are 68% confidence. B) Same as A but with low stress samples (50mM salt, 100mM

mannitol). C) Euclidian distance-clustered heatmap of differentially expressed proteins that were

significantly enriched (>0.58 Log_2 fold change) in all 3 AHA-treated conditions over untreated

control in the high stress conditions, without Class I hits that were absent in the negative control

(n = 117). D) Same as C, but for only terms enriched in all 3 groups in low stress conditions (n =

219).

SUPPLEMENTAL FIGURES

Supplemental Figure 1. Representative images of seedling samples during AHA and HPG

viability assays.

Supplemental Figure 2. Enlarged gene ontology of high stress set AHA-treated seedlings.

Supplemental Figure 3. Enlarged gene ontology of low stress set AHA-treated seedlings.

SUPPLEMENTAL TABLES

Supplemental Table 1. Enriched AHA-treated low and high salt and mannitol stress

proteomics data.

Supplemental Table 2. Ontologizer GO output data for AHA-treated low and high salt and

mannitol stressed proteins.

Supplemental Table 3. Differentially expressed proteins above significance threshold
divided by group between positive control, salt stressed, and mannitol stressed
seedlings in both high and low stress.

Figure 1. Visualization of BONCAT technology. A) Incorporation mechanism of AHA and HPG into the nascent proteome. B) Workflow for click labelling. Whole seedlings are acclimatized to liquid 0.5X MS media for 24 hours before treatment with AHA or HPG. Acute exposure time in the modified protocol is limited to 30 minutes, followed by 3 hours of incorporation time. Protein is extracted from seedlings for click chemistry in vitro, followed by visualization via anti-biotin blot to determine qualitative labelling before streptavidin enrichment for LC-MS/MS.

Figure 2. Streptavidin-detected anti-biotin blots of biotin-clicked Arabidopsis whole lysate samples. Beneath each blot is a cropped segment of a Coomassie Blue gel loading control. Each experiment was performed in triplicate, but single replicates are shown for simplicity. A) Titration of AHA and HPG concentrations with a 30-minute acute exposure and 3-hour incorporation time. B) 50µM AHA and HPG tested with exposure times varying from 15 minutes to 22 hours, all followed by a 3-hour incorporation. C) 50µM AHA and HPG tested with incorporation times varying from 30 minutes to 24 hours, each preceded with a 30-minute exposure time. D) Low concentration AHA (10µM) test of varying incorporation times with a 30-minute exposure (left) or varying exposure times with a 3-hour incorporation (right).

Figure 3. Viability assays of Arabidopsis seedlings exposed to methionine analogs. Average fold change in root length of seedlings with standard deviation error bars is plotted, and t-tests between experimental and control groups were performed with p-value significance < 0.05 shown. A) n = 72 5 d (days)-old seedlings per condition were transplanted to agar plates containing AHA or HPG for 1-3 d to simulate extensive exposure times. High concentrations after 3 days (left), low concentrations after 3 days (middle) and low concentrations after 24 hours (right) are shown. B) n = 54 seedlings per condition were acclimatized to liquid media for 24 h, then exposed for 30 min in AHA- or HPG-containing liquid media before being returned to solid medium and grown for 7 d. C) Snapshot of days 5 and 7 for all conditions from summarized data in 2B to show significant changes.

Figure 4. Summary mass spectrometry data for 50µM AHA-labelled 3-hour salt or mannitol stressed Arabidopsis seedlings. A) Venn diagrams of valid (>1.5X FC or below detection in negative control) protein groups between positive control (AHA treated, unstressed) and stress conditions in either high stress (150mM salt and 300mM mannitol, left, n = 549) or low stress (50mM salt, 100mM mannitol, right, n = 797) conditions. B) Exclusively Class I hits from 1A that are absent in negative controls (n = 147 and n = 300. C) STRING network of valid protein hits enriched in the low stress salt and/or mannitol conditions that were below detection or below threshold in both positive and negative controls. Only nodes with >2 edges were kept. Increasing line thickness is edge score ranging from 0.5-1. D) Gene ontology biological processes (Benjamini-Hochberg parent-child union) for high stress (left, p < 0.0025) and low stress (right, p < 0.005) conditions. All proteins enriched in one condition over negative controls were included. Highlighted in red are terms directly related to stress response.

Figure 5. Proteomics data representation of salt and mannitol stressed AHA-treated seedlings. A) PCA plot of high stress (150mM salt, orange, and 300mM mannitol, purple) 50μM AHA-treated seedlings, as well as control treated (blue) and control untreated (grey) samples. Elipses are 68% confidence. B) Same as A but with low stress samples (50mM salt, 100mM mannitol). C) Euclidian distance-clustered heatmap of differentially expressed proteins that were significantly enriched (>0.58 Log_2 fold change) in all 3 AHA-treated conditions over untreated control in the high stress conditions, without Class I hits that were absent in the negative control (n = 117). D) Same as C, but for only terms enriched in all 3 groups in low stress conditions (n = 219).