Optimization of Bio-Orthogonal Non-Canonical Amino acid Tagging

(BONCAT) for effective low-disruption labelling of Arabidopsis

proteins *in vivo*

Nicholas Hassan[1], Shelly Braun[2], Mohana Talasila[1], Richard P. Fahlman[2] and R. Glen

Uhrig[1,2#]

[1] Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

[2] Department of Biochemistry, University of Alberta, Edmonton, Alberta, Canada

[#] Senior and corresponding author:

Dr. R. Glen Uhrig

Department of Biological Sciences,

University of Alberta,

Edmonton, Alberta, Canada, T6G 2E9

Email: ruhrig@ualberta.ca

Phone: +1 780 492 3088

Fax : +1 780 492-9234ruhrig@ualberta.ca

28 **Running Title:** Developing Bio-Orthogonal Non-Canonical Amino acid Tagging

29 (BONCAT) for *in-vivo* labelling of newly synthesized plant proteins in plants

32 **Key Words:** BONCAT, Click Chemistry, Mass Spectrometry, Quantitative Proteomics

ABSTRACT

Plants, as sessile organisms, require robust responses in protein synthesis to adapt to variable environmental conditions. Measurement of newly synthesized proteins has been successfully facilitated with Bio-Orthogonal Non-Canonical Amino acid Tagging (BONCAT) across a multitude of organisms. Here, we use non-canonical amino acids (NCAAs) L-azidohomoalanine (AHA) or L-homopropargylglycine (HPG) incorporation in place of methionine residues into the actively translating *Arabidopsis* proteome, allowing for that subset of proteins to be enriched for mass spectrometry quantification. Although this technique has seen occasional use in plants, optimization of the protocol to maximize functionality while minimizing organismal stress has not yet been established. Here, we provide evidence for successful implementation through the liquid immersion of seedlings in AHA or HPG-containing media that functions with significantly lower concentrations than the literature standard. Our approach splits acute exposure and incorporation phase of labelling to mitigate potential negative impacts of prolonged NCAA exposure without compromising effective enrichment capacity, and demonstrate that this results in an unperturbed growth phenotype for AHA-treated seedlings. Finally, we demonstrate the capacity of this modified approach to enrich newly synthesized proteins from the whole proteome under standard stress conditions. These improvements allow for a broader use of BONCAT technologies in molecular plant research, affording a deeper understanding of the newly synthesized proteome without negatively impacting plant health.

INTRODUCTION

The balance between protein synthesis and degradation defines proteostasis in all cells, which in turn defines the functions and capacities of all organisms. Rapid adjustment of protein levels and plasticity of the proteome is essential for survival in a changing environment, especially for plants, which are carefully tuned to their immediate environment. Defining newly synthesized proteins, or the nascent proteome, is crucial to understanding regulation of the plant cell and adaptation to environmental stressors, and can provide a quantitative measurements of the effectors produced within a cell. However, conventional strategies for labelling the newly synthesized proteome for detection via mass spectrometry (MS), such as radiolabelling, often lack depth of analysis due to abundant, older peptides obscuring detection of lower-abundance proteins of interest (Tivendale et al., 2021). Isolation of the nascent proteome would allow for a closer view of proteins lower in abundance, increasing sensitivity of analysis. To fill this niche, Bio-Orthogonal Non-Canonical Amino acid Tagging (BONCAT), has been developed to label nascent proteins with a methionine analog to allow for enrichment of newly synthesized proteins (Dieterich et al., 2006; Parker & Pratt, 2020).

Azide and alkyne groups can be irreversibly conjugated to each other via a copper-catalyzed azide-alkyne cycloaddition (CuAAC), which utilizes the efficient, biologically non-reactive click chemistry mechanism (reviewed in Chen et al., 2023). To employ this in a biological setting, unnatural methionine analogs L-azidohomoalanine (AHA, azide) or L-homopropargylglycine (HPG, alkyne) are supplied to live organisms. These non-canonical amino acids (NCAAs) are incorporated into proteins during synthesis by the relatively promiscuous methionine tRNA in a moderately methionine-depleted environment (Figure 1A), but complete methionine depletion in eukaryotes is generally not recommended, as perturbations to the proteome can be mitigated in a methionine-containing environment by limiting NCAA incorporation beyond levels required for enrichment (Bagert et al., 2014). After application, proteins are then extracted and "clicked" to the opposing azide or alkyne conjugated to biotin, effectively biotinylating newly synthesized proteins and facilitating enrichment via streptavidin resin for subsequent quantification via liquid chromatography (LC)-MS/MS. While BONCAT was initially developed for use in mammalian cells, it has since found use in a variety of organisms, such as live mice, bacteria, and archaea (Dieterich et al., 2006; Kern & Ferreira-Cerca, 2022; Lindivat et al., 2021; Saleh et al., 2019). However, unlike other organisms, BONCAT implementation in plants has not been fully realized.

In the model plant *Arabidopsis thaliana* (Arabidopsis), BONCAT method was first implemented in 2017. Here, AHA and HPG were applied to Arabidopsis seedlings through the flooding of agar plates with 1 mM AHA (Glenn et al., 2017). This yielded thousands of enriched

proteins after only a few hours of exposure time. However, subsequent endeavours found that
Arabidopsis reacts metabolically to these compounds in a similar way to excess methionine,
stunting growth and/or impacting plant health during prolonged exposure (Tivendale et al.,
2021). Since then, only a handful of studies have utilized BONCAT in plants, and changes to the
protocol have generally been limited to exposure time of the seedlings to the tags, with few
studies endeavouring alternate implementations using less than 1 mM of the NCAAs (Li et al.,
2023; Linster et al., 2022; Lyu et al., 2023; Xie et al., 2023; Yu et al., 2020). Together, these
studies indicate that optimization of BONCAT in plants remains incomplete, as a wider variety of
times and concentrations need to be tested to strike a balance between effective labelling
concentrations and, ideally, minimal disturbance to the health of the seedling plants. Additionally,
protocol modifications for simplicity remain unexplored, including: a) using the same biotin-
conjugated clicked extract for both anti-biotin blots and enrichment for mass spectrometry to
minimize variability instead of clicking to a separate fluorophore for blot detection and b)
exploring the exclusivity of the azide-alkyne cycloaddition, such as eliminating the need for
repeated precipitation or processing prior to the click reaction to remove metabolites (Ma et al.,
2018).

Here, we present an adapted protocol for BONCAT labelling of Arabidopsis seedlings in
liquid 0.5x MS medium that allows for more reliable seedling exposure to, and removal of, AHA
and HPG (Figure 1B). Specifically, we demonstrate this through experimentation involving the
titration of AHA and HPG as well as optimization of labelling time and approach to balance
effective labelling with minimized exposure these methionine analogs. Lastly, we perform
extensive seedling viability assays to ensure minimal disruption of plant growth with the
modified protocol, and evaluate the newly synthesized proteome using the modified protocol
under high and low salt and mannitol stress via LC-MS/MS.

RESULTS AND DISCUSSION

AHA and HPG label Arabidopsis seedlings at 50 µM or less using short application
pulses

To determine concentrations of AHA and HPG suitable for application pulses, 5 d-old
seedlings were acclimated to liquid medium overnight before being exposed to 0, 10, 25, 50,
100, or 250 µM AHA or HPG for 30 min (see Materials and Methods). Washed seedlings were
then allowed to incorporate any remaining tag into the nascent proteome for an additional 3 h
before whole plants were flash frozen and extracted. Copper-catalyzed click chemistry was then
performed *in vitro* to conjugate NCAA tags to biotin, and total proteome biotinylation was

detected via immunoblot. Anti-biotin immunoblotting against untagged controls was selected as
the preferred method for monitoring qualitative labelling efficiency, as whole proteome analysis
may not detect AHA tags that have a mass shift similar to salt adducts of methionine (Liu et al.,
2022). Although HPG appeared to have lower background labelling than AHA, the signal from
AHA was observed to be higher, as above-background labelling is visible at all tested
concentrations of AHA and above 25 µM for HPG (Figure 2A). Correspondingly, a concentration
of 50 µM AHA or HPG was selected as an optimal concentration due to clearly visible labelling
with minimal required reagent, primarily to minimize metabolic stress on the seedlings.

We then performed time-course labelling to select ideal labelling periods. Labelling periods
were divided into acute exposure time (NCAA present in medium) and incorporation time (NCAA
removed from medium), with NCAA proteome incorporation assessed independently. This
revealed that an increase in exposure time led to an increase in labelling over the course of 24
hours for both 50 µM AHA and HPG, but above-background labelling was visible via immunoblot
within the first 15 minutes (Figure 2B). Similarly, increased incorporation time with a set initial
exposure of 30 min appears to increase labelling for up to 6 h after removal of tags using an
initial 50 µM of either NCAA, suggesting residual NCAA may remain present in the plant's
tissues after wash and removal from the medium and that the remaining reagent continues to be
incorporated for several hours (Figure 2C). From this data, a 30 min exposure followed by 3 h of
NCAA incorporation were selected as optimal labelling times for 50 µM of either NCAA, with a
shorter acute time to mitigate potential negative effects of exposure on the seedlings.
Interestingly, low concentrations of AHA (10 µM) also showed increases in labelling across
longer exposure times, but labelling stagnated with prolonged incorporation (Figure 2D).
Therefore, this concentration is likely too low for excess NCAA to accumulate in plant tissue
prior to its removal, but may be more suitable for long-term exposure labelling, such as
overnight incubations for pulse-chase assay.

Short exposure times prevent problems associated with prolonged exposure to AHA
Viability measurements of NCAA treated Arabidopsis seedlings was performed using fold-
change in root length as a proxy for plant health, in order to measure the potential negative
effects of AHA and HPG. Here, 5 d-old Arabidopsis seedlings were transplanted onto 0.5x MS
agar plates containing AHA or HPG and allowed to grow for multiple days. All tested
concentrations from 10 to 250 µM AHA and HPG decreased plant root growth across a 3-day
exposure, with HPG being significantly worse than AHA, even at comparably low concentrations
(Figure 3A). However, with only 24 h of exposure, 10 µM AHA did not show a significant

reduction in root growth. As this concentration shows above-background labelling within a few
hours but does not continue to incorporate the NCAA after being washed (Figure 2D), 10 µM
AHA may find an appropriate use in pulse-chase assays, where its capacity for 24-h acute
exposure labelling could be used to label plants overnight for degradation experiments the
following day.

Differing approaches have been proposed for *in vivo* AHA and HPG viability treatments in the
past. Use of a flooding approach to supply 1 mM AHA or HPG to Arabidopsis seedlings, a
similar step to acute exposure in this experiment, has previously been shown to negatively
impact whole seedling growth and cell culture viability, in addition to perturbing metabolism over
a multi-day duration, which may be related to the strict metabolic balance of methionine in
plants (Tivendale et al., 2021). However, other eukaryotes demonstrate a more mixed or neutral
response to exposure. For example, 2 mM AHA and 1 mM HPG have been shown to have no
negative viability effects on monkey and mouse-derived cells, and experimentation on HeLa
cells has shown that perturbation of the proteome in response to AHA can be mitigated as long
as methionine remains present in the medium (Bagert et al., 2014; Su Hui Teo et al., 2016;
Wisse et al., 2017). On the molecular level, Morey et al (2021) found mouse-derived cell lines
exhibit moderate stress signals in the form of higher levels of heat shock proteins HSP70 and
HSP90 when treated with 50 µM AHA, but no significant instability or increase in global protein
ubiquitination during AHA treatment was observed, as well as no apoptosis or cell death that
would be expected with a significant reduction in viability (Morey et al., 2021).

In contrast, Tivendale et al. (2021) found that AHA was more toxic than HPG, especially in
plant cell cultures. Interestingly, bacterial studies have shown similar results to this study, with
AHA being tolerable to log-phase *E. coli* at higher concentrations than HPG, where HPG
reduced bacterial growth at concentrations above 2.8 µM while AHA was tolerated up to the 200
µM range (Landor et al., 2023). This effect seems to differ slightly between prototrophs and
auxotrophs, where bacteria unable to synthesize essential compounds appear to be more
negatively affected by HPG (Jecmen et al., 2023). Both AHA and HPG have been shown to
mildly affect bacterial metabolism as well, but are not degraded as the expected products of or
processed as methionine (Steward et al., 2020). Contrast in viability between these organisms
and plants in different studies is likely related to difference in methods used, as well as
metabolic effects.

It has been proposed that negative long-term effects of NCAAs on organism or cell growth
may be irrelevant if incorporation into nascent proteins and immediate toxicity does not occur
during the length of the experiment (Landor et al., 2023). All prior viability assays looked at

156 organism or cellular responses over a prolonged exposure time, which is not representative of

157 the experimental conditions. To accomplish a more experimentally valid condition, we repeated

158 viability assays with a short acute exposure in liquid 0.5x MS, followed by seedling washing and

159 return to agar plates for 7 d to simulate prolonged incorporation time. All tested concentrations

160 of AHA (up to 250 µM) did not exhibit reduced growth compared to control after implementation

161 of a 30-min exposure and this protocol (Figure 3B-C), suggesting that the long-term exposure

162 growth defects are not a result of immediate toxicity of the NCAA. In contrast, 50 µM and 100

163 µM concentrations of HPG showed a slight reduction in growth compared to control under the

164 same conditions (Figure 3C). By limiting exposure to 30 min and allowing incorporation to

165 continue for < 6 h, viability concerns of the plants can be circumvented without compromising in

166 labelling efficiency, particularly in the use case of AHA.

168 ***AHA-labelled proteins under salt and osmotic stress***

169 To benchmark quantitative labelling efficiency of the optimized protocol in different conditions,

170 we applied either salt or osmotic stress during an AHA labelling period with unlabelled protein as

171 a bead binding control and unstressed AHA-labelled seedlings as a positive control. The

172 experiment was performed with either high stress (150 mM salt or 300 mM mannitol) or low

173 stress (50 mM salt or 100 mM mannitol) conditions, as the effects of applied stressors may be

174 intensified in liquid submersion. Here, we find more than 400 proteins enriched in at least one

175 condition under high stress conditions, and more than 600 under low stress conditions, with a

176 greater number of overlapping proteins between groups in the lower stress condition (Figure

177 4A). Many of these proteins are Class I hits, which we define as below detection in the negative

178 control (Figure 4B). Additionally, using the positive control, we can dissect stress specific

179 responses. Network analysis of the hits exclusive to salt and mannitol stress, even in the lower

180 stress conditions, resolves multiple protein networks related to abiotic stimulus response (Figure

181 4C). Gene ontology enrichment analysis also resolved numerous abiotic stress response

182 biological processes in the stress conditions than versus the control, including biological

183 processes such as wax biosynthesis and water-related regulation (Figure 4D, highlighted).

185 **SUMMARY**

186 In this study, we established a workflow for the use of AHA and HPG to label and enrich the

187 newly synthesized proteome of Arabidopsis in less than 4 h without reducing seedling viability.

188 Although 50 µM AHA was selected as the optimal labelling concentration, the protocol is flexible

189 and modifiable based on need, and modified protocols can be tracked for efficiency at multiple

stages via blotting from the same clicked sample that is used for enrichment. Enriched samples

show clear definition between stress and unstressed conditions in relatively short periods. As a

technique that allows for isolation of the newly synthesized proteome, establishing a BONCAT

workflow in Arabidopsis that mitigates negative plant responses unveils outstanding

opportunities for applications where enrichment of a subsect of the proteome is required for

sufficient depth of analysis. Lastly, the potential for NCAA tag incorporation at concentrations

substantially below literature standard has positive implications for the use of this technique in

plants moving forward.

MATERIALS AND METHODS

Plant growth conditions

Arabidopsis thaliana seeds (Col-0) were liquid sterilized by immersion in 70% (v/v) ethanol

for 2 min followed by 30% (v/v) bleach for 10 min. Sterilized seeds were vernalized in H_2O in the

dark at $4^{o}C$ for 3 d. Germination plates were prepared using half-strength Murashige & Skoog

(MS) basal salts, pH adjusted to 5.8 with aqueous KOH, with 0.08% (w/v) agar.

For click chemistry, vernalized seeds were stratified on plates and allowed to grow with

plates positioned vertically in 24-h light at $22^{o}C$ for 5 days. Approximately 100 mg of 5 d-old

seedlings were immersed in autoclaved half-strength MS media in 24-well plates and placed

under constant light for 16 hours to acclimatize. MS media was removed and replaced with

50uM filter sterilized L-azidohomoalanine or L-homopropargylglycine (Vector Labs, CCT-1066

and CCT-1076) from 0.1M stocks dissolved in H_2O and diluted in 0.5x MS media. Seedlings

were then incubated for 30 minutes of exposure time in constant light before being washed once

in 0.5x MS and incubated in constant light in 0.5x MS for 3 hours of "incorporation" time.

Labelled seedlings were removed from media, rinsed in 0.5x MS, and quickly touch dried before

being flash frozen in liquid N_2 and stored at $-80^{o}C$ until subsequent steps.

For root length assay of reagent toxicity, seeds were sown individually on 0.5x MS agar

plates positioned vertically in dark boxes with seeds at the light interface to emulate roots

growing in the dark and shoots growing in the light. Seedling growth conditions consisted of a

12 h photoperiod, $22^{o}C$ and 100 $\mu mol/m^2s$ light. After 5 d for the long-term reagent exposure,

seedlings were transplanted along the same light/dark interface to 0.5x MS plates containing 0,

10, 25, 50, or 250 µM AHA or HPG. For the short-term exposure, seedlings were placed into

liquid 0.5x MS overnight to equilibrate, then immersed for 30 minutes in 0.5x liquid MS

containing 0, 10, 25, 50, or 250 µM AHA, before being washed in 0.5x MS and re-plated on to

vertical plates. Root length was recorded for 7 d using digital calipers, and fold change in

individual seedling root length relative to day 0 (transplanting day) was plotted. Recording of
root length stopped after day 3 for the long-term exposure due to statistically significant
differences between all concentrations relative to controls.
227

Optimization of plant labelling

To determine optimal concentrations of label, variable concentrations of 10, 25, 50, 100, or
250 µM of AHA or HPG were applied as above. Optimal concentrations for labelling were
chosen based on a combination of intensity of labelled protein present on streptavidin
immunoblots and minimal observed growth retardation on 0.5X MS agar containing the same
concentration of label. For time-course experimentation, acute exposure times were varied
between 15 min, 30 min, or 1, 2, 4, or 22 h. Incorporation times were varied between 30 minutes
or 1, 2, 3, 6, or 24 h. Optimal incubation times were chosen based on minimal required times to
achieve effective labelling at the desired concentration as visualized via immunoblot, and
verified to cause insignificant or no growth inhibition to the seedlings based on viability studies.
238

Protein extraction and quantification

Flash frozen tissue was ground using Geno/Grinder (SPEX SamplePrep) for 30 s at 1200
rpm with 3mm glass beads and extracted in either 50mM HEPES (pH 7.5) with 1% (w/v) SDS or
50 mM Tris (pH 8.0) with 1% (w/v) SDS, at a ratio of 1:2 w/v ground tissue to extraction buffer.
Following optimization, HEPES was selected as the extraction buffer for all samples due to
better click labelling observed. Samples were heated at $95^{o}C$ for 10 min and 1000 rpm on
Eppendorf ThermoMixer F2.0, cooled, and centrifuged at 16,000 xg for 10 min at room
temperature to remove tissue particulates. Protein amounts were quantified using Bradford
assay with samples diluted to 0.2% (w/v) SDS against BSA controls.
248

Click chemistry reaction

A total of 250 µg of protein from each sample was diluted in extraction buffer with 1% (w/v) or
no SDS to a volume of 346 µL ensuring that a final concentration (in 400uL total) of 0.5% (w/v)
SDS to maintain protein solubility in solution. To each reaction, the following was added: 8uL of
50 mM tris(2-carboxyethyl)phosphine (TCEP)-HCl (Sigma; diluted from 1 M stock in H_2O same
day of use), 30 µL of 1.7 mM tris[(1-benzyl-1H-1,2,3-triazol-4-yl)methyl]amine (TBTA) (Vector
Labs CCT-1061; in 20% (v/v) dimethyl sulfoxide (DMSO) and 80% (v/v) tert-butanol, stored at
room temperature as in Ma et al., 2018), 8 µL of CuSO4 (Sigma; in H_2O, stored at $-20^{o}C$), and 8
µL of 5mM biotin-PEG3-azide (Vector Labs CCT-AZ104; for HPG samples, in DMSO) or biotin-

PEG4-alkyne (Vector Labs CCT-TA105; for AHA samples, in DMSO). Click reactions were
allowed to proceed for 1 h at room temperature, shaking at 700 rpm and protected from light.
Reactions were then quenched via chelation with 20 µL of 0.5 M ethylenediaminetetraacetic
acid (EDTA) (pH 8.0, stored at 4^{o}C).

Validation of click chemistry labelling was determined via streptavidin immunoblot. Here, 6 µg
of protein was run on 10% SDS-PAGE gels then transferred to 0.45 µM polyvinylidene fluoride
(PVDF) membrane semi-dry over 45 min at 15 V. Blots were blocked in 5% fat-free milk in Tris-
buffered saline (TBS) for 1 h, at which point HRP-conjugated streptavidin diluted 1:2000 in Tris-
buffered saline with 0.1% Tween-20 (TBST) was added for an additional hour. Membranes were
washed in TBST and visualized using Clarity Western ECL Substrate. Colloidal Coomassie
Blue-stained gels were used as loading controls.

Enrichment of clicked proteins

Excess biotin was removed from clicked samples using Cytiva PD Spintrap G-25 columns.
Bead binding comparison was done using Invitrogen streptavidin-C1 magnetic beads and Cytiva
streptavidin Sepharose beads; all depicted experiments proceeded using non-magnetic
Sepharose beads. All beads were initially washed 3 times in 50mM HEPES, pH 7.5, for 1 min
while rotating at room temperature. For all steps, supernatant was removed from magnetic
beads by adhering beads to a magnetic rack, and from other types of beads by centrifugation
for 1 minute at 600 xg. Up to 200 µg of flow-through protein from the Spintrap columns was
loaded onto equilibrated streptavidin beads and allowed to bind while rotating for 16 hours at
4^{o}C. The supernatant was removed and retained at -20^{o}C to assess capture efficiency.

Capture efficiency was checked before digestion via streptavidin-detected dot blot
comparison of clicked protein, Spintrap column flow-through, and supernatant after overnight
bead binding. After confirmation of a signal reduction of more than 70% in the supernatant after
binding, beads were washed sequentially with the following solutions at 1mL each: 50 mM
HEPES pH 7.5 (2x washes), 1 M potassium chloride, ice-cold 100 mM sodium carbonate (2x
washes), 2 M urea in 50 mM Tris-HCl pH 8.0 (2x wash), and 50 mM triethylammonium
bicarbonate (TEAB) in HPLC-grade water (3x washes). All washes were completed at room
temperature with rotation for 3 min.

Peptides were generated through on bead digestion using trypsin diluted 1:100 in 50 mM
TEAB overnight at 37^{o}C; the supernatant was removed before the beads were washed with 50
mM TEAB and the supernatant pooled with previously extracted peptides. The extract was
reduced with 10 mM DTT at 95^{o}C for 5 min while shaking at 450 rpm, alkylated with 30 mM

iodoacetamide (IA) at room temperature for 30 min, and the reaction quenched with 10 mM DTT

for 10 minutes. Samples were then acidified to 0.5% (v/v) TFA to a pH of less than 2 and

centrifuged at 16,000 xg for 1 min before the supernatant was removed and dried down.

Peptides were re-suspended in 3% (v/v) acetonitrile / 0.1% (v/v) formic acid for desalting via

ZipTip (Sigma cat. no. ZTC18S960) before being dried and stored at -80°C. Desalted peptides

were the re-suspended in 3% (v/v) acetonitrile / 0.1% (v/v) formic acid for mass spectrometry

analysis.

LC MS/MS and Data Analysis

Peptides were analyzed on a FAIMSpro mounted Fusion Lumos Orbitrap mass

spectrometer (Thermo Fisher Scientific) in data independent acquisition (DIA) mode. Re

suspended peptides were injected using Easy-nLC 1200 system (LC140; Thermo Fisher

Scientific) and an Acclaim PepMap 100 C18 trap column (Cat# 164750; Thermo Fisher

Scientific) followed by a 50 cm Easy-Spray PepMap C18 analytical column (ES903; Thermo

Fisher Scientific) warmed to 50 °C. Peptides were eluted at 0.3 µl/min using a segmented

solvent B gradient of 0.1% (v/v) formic acid in 80% (v/v) acetonitrile (A998, Thermo Fisher

Scientific) from 4 to 41% solvent B (0–60min). The FAIMSpro was used with a fixed gas flow of

3.5 L/min with a 3-CV setting of -30, -50, -70. A positive ion spray voltage of 2.3 kV was used

with an ion transfer tube temperature of 300 °C and an RF lens setting of 40%. Direct DIA

acquisition was performed as previously described (Mehta et al., 2022). Full scan MS^1 spectra

(350–1400 *m/z*) were acquired with a resolution of 120,000 at 200 *m/z* with a normalized

automatic gain control (AGC) target of 100% and IT set to automatic. Fragment spectra were

acquired at resolution of 30,000 across 28 38.5 *m/z* windows overlapping by 1 *m/z* using a

dynamic maximum injection time and an AGC target value of 2000%, with a minimum number of

desired points across each peak set to 6. Higher-energy collisional dissociation fragmented was

performed using a fixed 27% fragmentation energy. Downstream data analysis of dDIA

acquisitions was performed using Spectronaut ver. 19 (Biognosys AG) with default settings as

previously described (Mehta et al., 2022), with all searches made against a custom-made decoy

(reversed) version of the Arabidopsis protein database from Araport 11 (27,533 protein encoding

genes; ver. 2022-09-14). Briefly, search parameters included the following: trypsin digest

permitting two missed cleavages, fixed modifications (carbamidomethyl (C)), variable

modifications (oxidation (M)) and a peptide spectrum match, peptide and protein false discovery

threshold of 0.01.

Bioinformatic Data Analysis

Gene Ontology (GO) enrichments were performed using the Ontologizer
(http://ontologizer.de/; Bauer et al., 2008) using a Benjamini-Hochberg parent-child union with
an analysis cutoff of p > 0.01, and visualized in R 4.5.0. Significantly changing proteins (Log2FC
of MS2 spectra > 0.58 over negative control) were pooled with Class I protein hits (absent in
MS2 of negative controls) to form the foreground with a background of all identified proteins
(Supplemental Table 1). A subset of the same dataset that was below threshold abundance in
both positive and negative controls was used for network analysis of stress-exclusive proteins
via the StringApp plugin for Cytoscape v. 3.10.3, with an edge score cutoff of 0.5 (Doncheva et
al., 2019, https://cytoscape.org/). The STRING Enrichment within StringApp was used to assign
outlines based on gene ontology. Non-network visualizations were produced using the *ggplot2*
R package, Biorender (https://www.biorender.com/), GraphPad Prism v. 8.4.3
(https://www.graphpad.com/), and Affinity Designer v. 1.10.5 (https://affinity.serif.com/en-
us/designer/).

ACKNOWLEDGEMENTS

The authors would like to thank the Canadian Foundation for Innovation (CFI) and Natural
Sciences and Engineering Council of Canada (NSERC) for funding this work. We would also
like to thank Jack Moore and the Alberta Proteomics and Mass Spectrometry Facility (APM) for
mass spectrometry assistance and upkeep.

AUTHOR CONTRIBUTIONS

Nicholas Hassan: Investigation, Conceptualization, Methodology, Data curation, Formal
analysis, Visualization, Writing—original draft. *Shelly Braun*: Writing—review & editing,
Conceptualization. *Mohana Talasila*: Formal analysis, Data curation, Visualization. *Richard
Fahlman*: Supervision, Writing—review & editing, Material Resources *R. Glen Uhrig*:
Supervision, Conceptualization, Writing—review & editing, Funding acquisition.

DATA AVAILABILITY

Raw data have been deposited to the ProteomeExchange Consortium
(http://proteomecentral.proteomexchange.org) *via* the PRoteomics IDEntification Database
(PRIDE; https://www.ebi.ac.uk/pride/) partner repository with the data set identifiers PXD064210

REFERENCES

Bagert, J. D., Xie, Y. J., Sweredoski, M. J., Qi, Y., Hess, S., Schuman, E. M., & Tirrell, D. A.
(2014). Quantitative, time-resolved proteomic analysis by combining bioorthogonal
noncanonical amino acid tagging and pulsed stable isotope labeling by amino acids in cell
culture. *Molecular and Cellular Proteomics*, *13*(5), 1352–1358.
https://doi.org/10.1074/mcp.M113.031914

Bauer, S., Grossmann, S., Vingron, M., & Robinson, P. N. (2008). Ontologizer 2.0--a
multifunctional tool for GO term enrichment analysis   and data exploration.
Bioinformatics (Oxford, England), *24*(14), 1650–1651.
https://doi.org/10.1093/bioinformatics/btn250

Chen, M. M., Kopittke, P. M., Zhao, F. J., & Wang, P. (2023). Applications and opportunities of
click chemistry in plant science. *Trends in Plant Science*, *29*(2), 167–178.
https://doi.org/10.1016/j.tplants.2023.07.003

Dieterich, D. C., Link, A. J., Graumann, J., Tirrell, D. A., Schuman, E. M., & Sharpless, K. B.
(2006). Selective identification of newly synthesized proteins in mammalian cells using
bioorthogonal noncanonical amino acid tagging (BONCAT). *National Academy of Sciences*,
103(25), 9482–9487. www.pnas.orgcgidoi10.1073pnas.0601637103

Doncheva, N. T., Morris, J. H., Gorodkin, J., & Jensen, L. J. (2019). Cytoscape StringApp:
Network Analysis and Visualization of Proteomics Data. *Journal of Proteome Research*,
18(2), 623–632. https://doi.org/10.1021/acs.jproteome.8b00702

Glenn, W. S., Stone, S. E., Ho, S. H., Sweredoski, M. J., Moradian, A., Hess, S., Bailey-Serres,
J., & Tirrell, D. A. (2017). Bioorthogonal noncanonical amino acid tagging (BONCAT)
enables time-resolved analysis of protein synthesis in native plant tissue. *Plant Physiology*,
173(3), 1543–1553. https://doi.org/10.1104/pp.16.01762

Jecmen, T., Tuzhilkin, R., & Sulc, M. (2023). Photo-Methionine, Azidohomoalanine and
Homopropargylglycine Are Incorporated into Newly Synthesized Proteins at Different Rates
and Differentially Affect the Growth and Protein Expression Levels of Auxotrophic and
Prototrophic E. coli in Minimal Medium. *International Journal of Molecular Sciences*,
24(14). https://doi.org/10.3390/ijms241411779

Kern, M., & Ferreira-Cerca, S. (2022). Differential Translation Activity Analysis Using
Bioorthogonal Noncanonical Amino Acid Tagging (BONCAT) in Archaea. *Ribosome
Biogenesis: Methods and Protocols*, 229–246. https://doi.org/10.1007/978-1-0716-2501-
9_14

Landor, L. A. I., Bratbak, G., Larsen, A., Tjendra, J., & Våge, S. (2023). Differential toxicity of
bioorthogonal non-canonical amino acids (BONCAT) in Escherichia coli. *Journal of
Microbiological Methods*, *206*. https://doi.org/10.1016/j.mimet.2023.106679

Li, X., Chai, X., Lyu, H. N., Fu, C., Tang, H., Shi, Q., Wang, J., & Xu, C. (2023). Chemical
proteomics reveals an ISR-like response elicited by salicylic acid in Arabidopsis. In *New
Phytologist* (Vol. 237, Issue 5, pp. 1486–1489). John Wiley and Sons Inc.
https://doi.org/10.1111/nph.18646

Lindivat, M., Bratbak, G., Larsen, A., Hess-Erga, O. K., & Hoell, I. A. (2021). Flow Cytometric
Analysis of Bacterial Protein Synthesis: Monitoring Vitality After Water Treatment. *Frontiers*
in Microbiology, *12*. https://doi.org/10.3389/fmicb.2021.772651

Linster, E., Forero Ruiz, F. L., Miklankova, P., Ruppert, T., Mueller, J., Armbruster, L., Gong, X.,
Serino, G., Mann, M., Hell, R., & Wirtz, M. (2022). Cotranslational N-degron masking by
acetylation promotes proteome stability in plants. *Nature Communications*, *13*(1).
https://doi.org/10.1038/s41467-022-28414-5

Liu, C., Wong, N., Watanabe, E., Hou, W., Biral, L., Decastro, J., Mehdipour, M., Aran, K.,
Conboy, M. J., & Conboy, I. M. (2022). Mechanisms and Minimization of False Discovery of
Metabolic Bioorthogonal Noncanonical Amino Acid Proteomics. *Rejuvenation Research*,
25(2), 95–109. https://doi.org/10.1089/rej.2022.0019

Lyu, H. N., Fu, C., Chai, X., Gong, Z., Zhang, J., Wang, J., Wang, J., Dai, L., & Xu, C. (2023).
Systematic thermal analysis of the Arabidopsis proteome: Thermal tolerance, organization,
and evolution. *Cell Systems*, *14*(10), 883-894.e4. https://doi.org/10.1016/j.cels.2023.08.003

Ma, Y., McClatchy, D. B., Barkallah, S., Wood, W. W., & Yates, J. R. (2018). Quantitative
analysis of newly synthesized proteins. *Nature Protocols*, *13*(8), 1744–1762.
https://doi.org/10.1038/s41596-018-0012-y

Mehta, D., Scandola, S., & Uhrig, R. G. (2022). BoxCar and Library-Free Data-Independent
Acquisition Substantially Improve the Depth, Range, and Completeness of Label-Free
Quantitative Proteomics. *Analytical Chemistry*, *94*(2), 793–802.
https://doi.org/10.1021/acs.analchem.1c03338

Morey, T. M., Esmaeili, M. A., Duennwald, M. L., & Rylett, R. J. (2021). SPAAC Pulse-Chase: A
Novel Click Chemistry-Based Method to Determine the Half-Life of Cellular Proteins.
Frontiers in Cell and Developmental Biology, *9*. https://doi.org/10.3389/fcell.2021.722560

Parker, C. G., & Pratt, M. R. (2020). Click Chemistry in Proteomic Investigations. *Cell*, *180*(4),
605–632. https://doi.org/10.1016/j.cell.2020.01.025

Saleh, A. M., Jacobson, K. R., Kinzer-Ursem, T. L., & Calve, S. (2019). Dynamics of Non-
Canonical Amino Acid-Labeled Intra- and Extracellular Proteins in the Developing Mouse.
Cellular and Molecular Bioengineering, *12*(5), 495–509. https://doi.org/10.1007/s12195-
019-00592-1

Steward, K. F., Eilers, B., Tripet, B., Fuchs, A., Dorle, M., Rawle, R., Soriano, B.,
Balasubramanian, N., Copié, V., Bothner, B., & Hatzenpichler, R. (2020). Metabolic
Implications of Using BioOrthogonal Non-Canonical Amino Acid Tagging (BONCAT) for
Tracking Protein Synthesis. *Frontiers in Microbiology*, *11*.
https://doi.org/10.3389/fmicb.2020.00197

Su Hui Teo, C., Serwa, R. A., & O'Hare, P. (2016). Spatial and Temporal Resolution of Global
Protein Synthesis during HSV Infection Using Bioorthogonal Precursors and Click
Chemistry. *PLoS Pathogens*, *12*(10). https://doi.org/10.1371/journal.ppat.1005927

Tivendale, N. D., Fenske, R., Duncan, O., & Millar, A. H. (2021). In vivo homopropargylglycine
incorporation enables sampling, isolation and characterization of nascent proteins from

Arabidopsis thaliana. *The Plant Journal*, *107*(4), 1260–1276.
https://doi.org/https://doi.org/10.1111/tpj.15376

Wisse, L. E., Penning, R., Zaal, E. A., van Berkel, C. G. M., ter Braak, T. J., Polder, E., Kenney,
J. W., Proud, C. G., Berkers, C. R., Altelaar, M. A. F., Speijer, D., van der Knaap, M. S., &
Abbink, T. E. M. (2017). Proteomic and metabolomic analyses of vanishing white matter
mouse astrocytes reveal deregulation of ER functions. *Frontiers in Cellular Neuroscience*,
11. https://doi.org/10.3389/fncel.2017.00411

Xie, Z., Zhao, S., Li, Y., Deng, Y., Shi, Y., Chen, X., Li, Y., Li, H., Chen, C., Wang, X., Liu, E., Tu,
Y., Shi, P., Tong, J., Gutierrez-Beltran, E., Li, J., Bozhkov, P. V., Qian, W., Zhou, M., &
Wang, W. (2023). Phenolic acid-induced phase separation and translation inhibition
mediate plant interspecific competition. *Nature Plants*, *9*(9), 1481–1499.
https://doi.org/10.1038/s41477-023-01499-6

Yu, H., Kong, X., Huang, H., Wu, W., Park, J., Yun, D. J., Lee, B. ha, Shi, H., & Zhu, J. K.
(2020). STCH4/REIL2 Confers Cold Stress Tolerance in Arabidopsis by Promoting rRNA
Processing and CBF Protein Translation. *Cell Reports*, *30*(1), 229-242.e5.
https://doi.org/10.1016/j.celrep.2019.12.012

FIGURES

Figure 1. Visualization of BONCAT technology. A) Incorporation mechanism of AHA and
HPG into the nascent proteome.

Figure 2. Streptavidin-HRP anti-biotin blots of biotin-clicked Arabidopsis whole lysate
samples.

Figure 3. Viability assays of Arabidopsis seedlings exposed to methionine analogs.

Figure 4. Summary mass spectrometry data for 50µM AHA-labelled 3-hour salt or
mannitol stressed Arabidopsis seedlings.

SUPPLEMENTAL TABLES

Supplemental Table 1. Enriched AHA-treated low and high salt and mannitol stress
proteomics data.

Figure 1. Visualization of BONCAT technology. A) Incorporation mechanism of AHA and HPG into the nascent proteome. B) Workflow for click labelling. Whole seedlings are acclimatized to liquid 0.5X MS media for 24 hours before treatment with AHA or HPG. Acute exposure time in the modified protocol is limited to 30 minutes, followed by 3 hours of incorporation time. Protein is extracted from seedlings for click chemistry in vitro, followed by visualization via anti-biotin blot to determine qualitative labelling before streptavidin enrichment for LC-MS/MS.

Figure 2. Streptavidin-detected anti-biotin blots of biotin-clicked Arabidopsis whole lysate samples. Beneath each blot is a cropped segment of a Coomassie Blue gel loading control. Each experiment was performed in triplicate, but single replicates are shown for simplicity. A) Titration of AHA and HPG concentrations with a 30-minute acute exposure and 3-hour incorporation time. B) 50µM AHA and HPG tested with exposure times varying from 15 minutes to 22 hours, all followed by a 3-hour incorporation. C) 50µM AHA and HPG tested with incorporation times varying from 30 minutes to 24 hours, each preceded with a 30-minute exposure time. D) Low concentration AHA (10µM) test of varying incorporation times with a 30-minute exposure (left) or varying exposure times with a 3-hour incorporation (right).

Figure 3. Viability assays of Arabidopsis seedlings exposed to methionine analogs. Average fold change in root length of seedlings with standard deviation error bars is plotted, and t-tests between experimental and control groups were performed with p-value significance <0.05 shown. A) n = 72 5 d (days)-old seedlings per condition were transplanted to agar plates containing AHA or HPG for 1-3 d to simulate extensive exposure times. High concentrations after 3 days (left), low concentrations after 3 days (middle) and low concentrations after 24 hours (right) are shown. B) n = 54 seedlings per condition were acclimatized to liquid media for 24 h, then exposed for 30 min in AHA- or HPG-containing liquid media before being returned to solid medium and grown for 7 d. C) Snapshot of days 5 and 7 for all conditions from summarized data in 2B to show significant changes.

Figure 4. Summary mass spectrometry data for 50µM AHA-labelled 3-hour salt or mannitol stressed Arabidopsis seedlings. A) Venn diagrams of valid (>1.5X FC or below detection in negative control) protein groups between positive control (AHA treated, unstressed) and stress conditions in either high stress (150mM salt and 300mM mannitol, left) or low stress (50mM salt, 100mM mannitol, right) conditions. B) Exclusively Class I hits from 1A that are absent in negative controls. C) STRING network of valid protein hits enriched in the low stress salt and/or mannitol conditions that were below detection or below threshold in both positive and negative controls. Only nodes with >2 edges were kept. Increasing line thickness is edge score ranging from 0.5-1. D) Gene ontology biological processes (Benjamini-Hochberg parent-child union) for high stress (left, p > 0.0025) and low stress (right, p > 0.005) conditions. All proteins enriched in one condition over negative controls were included. Highlighted in red are terms directly related to stress response.