Title: Harnessing microbes for crop production: TS201 enhances maize yield and reduces pest damage

Authors: Man P. Huynh[1*], Khanh-Van Ho[2], Anne Z. Phillips[3], Amanda M Fox Ernwall[4], Dalton Ludwick[4], Deborah Finke[1], Zhentian Lei[2], Allison Jack[3], Bruce E. Hibbard[1,4,5], and Natalie W. Breakfield[3*]

Affiliations:

[1]Division of Plant Science & Technology, University of Missouri, Columbia, Missouri, 65211, USA

[2]Metabolomics Center, University of Missouri, Columbia, Missouri, 65211, USA

[3]NewLeaf Symbiotics, St. Louis, Missouri, 63132, USA

[4]Plant Genetics Research Unit, USDA-Agricultural Research Service, Columbia, Missouri, 65211, USA

[5]Biological Control of Insects Research Laboratory, USDA-Agricultural Research Service, Columbia, Missouri, 65203, USA

*Corresponding authors: manhuynh@missouri.edu, nbreakfield@newleafsym.com

Abstract: Microbial technologies are increasingly adopted to improve sustainable agriculture amid escalating economic, regulatory, and ecological pressures, yet few, if any, are supported by mechanistic understanding that translates to real-world performance. Here, we report TS201, a U.S. EPA–registered bioinsecticide, composed of *Methylorubrum extorquens*, that enhances maize yield and resilience under pest pressure. Across seven years (2016–2022) of field trials at 22 U.S. locations, TS201 increased yield and reduced lodging. Large-scale evaluations at 81 sites in eight U.S. states (2023–2024) confirmed its agronomic benefit. Mechanistically, TS201 induced biosynthesis of methyl anthranilate, a volatile insect-repellent compound, and triggered pest avoidance of treated roots. These findings reveal a novel plant-microbe-insect interaction and establish a systems biology framework for harnessing microbial plant resilience to advance crop production.

MAIN TEXT

Plant-associated microbiomes host diverse microbial taxa that often form symbiotic relationships with their host plants, shaping development and influencing resilience to environmental stressors (*1*). These interactions underpin a fast-growing class of agricultural biologicals enhancing crop productivity while reducing reliance on synthetic inputs (*2*). Despite their promise, most microbial products remain poorly characterized beyond proof-of-concept, and few, if any, are supported by mechanistic insights that translate to real-world agronomic outcomes.

This disconnection limits broader adoption to advance sustainable agriculture that feeds a growing
population in an era of intensifying regulatory, ecological, and economic pressures (*3*).

Among beneficial microbes, pink-pigmented facultative methylotrophs (PPFMs) in the
Methylobacterium and *Methylorubrum* genera are widespread in the phyllosphere and rhizosphere
(*4, 5*). These single-carbon metabolizers influence crop physiology by modulating nutrient uptake,
hormone balance, and immune signaling that can improve crop productivity (*6*). Here, we
investigate a bacterial solution TS201, composed of *Methylorubrum extorquens* that enhances
maize yield and mitigates pest damage evaluated under real-world field conditions.

44 TS201: field discovery and performance

We conducted several field trials across the U.S. Midwest to evaluate various naturally
occurring PPFMs as biological inoculants (biostimulants) to enhance maize performance. In one
such trial under high pressure of corn rootworms (CRW, *Diabrotica* spp.), the most serious pests
of maize (*7*), a subsequent wind event caused extensive lodging in control plots lacking PPFM
treatment. Remarkably, plots treated with selective PPFM strains remained upright. Lodging is
one of the most visible and economically consequential outcomes of CRW damage, as larval root
feeding weakens plant stability and, under environmental stressors such as wind or rain, causes
plants to lodge or topple, resulting in yield losses and harvest inefficiencies (*8, 9*). This observation
prompted multi-year field trials to identify the most effective strain capable of boosting yield and
mitigating CRW damage. Through this process, a PPFM species, *M. extorquens* strain NLS0042,
designated TS201, was selected and further investigated and developed into a commercial product.

From 2016 to 2022, we performed extensive small-plot field trials at 22 locations across
the U.S. Midwest, encompassing a range of low to high corn rootworm pressures, to evaluate the
impact of TS201 on maize yield. All trial sites included locations where CRW pressure exceeded
the economic injury threshold as indicated by nodal injury scores (NIS) of 0.25, a common
measure of CRW damage (*10, 11*), in untreated control plots. Across these sites over seven years,
TS201 significantly increased maize yield (220 kg/ha or 3.5 bu/A) and reduced CRW damage, as
shown by decreased NIS values, indicating enhanced root protection under pest pressure (Table 1,
table S1). Yield and pest suppression benefits were consistent across both conventional corn
hybrids and those expressing insecticidal traits targeting CRW, including Bt toxins and lethal
dsRNA constructs (table S2). TS201 also reduced lodging by 20% under moderate CRW pressure
(1.2 NIS, from 59% to 39%) and by 36% under high pressure (2.7 NIS, from 65% to 29%). Based
on its demonstrated biopesticidal efficacy, TS201 was registered in 2022 as a biological insecticide
(TS201™, EPA Reg. No. 95699-2, Terrasym®, NewLeaf Symbiotics, St. Louis, MO) with the
U.S. Environmental Protection Agency (EPA) under the Federal Insecticide, Fungicide, and
Rodenticide Act (FIFRA) (*12*).

Table 1. TS201 improves maize yield and root protection in small-plot field trials under corn rootworm
pressure. Field trials were conducted over seven growing seasons (2016-2022) at 22 locations across the
U.S. Midwest. Data represents phenotypic responses in TS201 treatment relative to untreated control.

Values summarize key agronomic and root protection traits observed in replicated small-plot trials. NIS: Nodal injury score.

Metric		TS201 treatment compared to control	P-value	Trialing scale
Performance	Yield	+220 kg/ha 11,706 kg/ha (control) vs. 11,926 kg/ha (TS201)	0.0108	22 locations over 7 years
Root damage	Nodal injury score (NIS)	-0.11 NIS 1.34 (control) vs. 1.23 (TS201)	0.0007	22 locations over 7 years

Following U.S. EPA registration, we conducted large-scale on-farm evaluations of TS201 during the 2023 and 2024 growing seasons across 81 geographically and agronomically diverse environments in eight U.S. states (Fig. 2a). TS201 was applied in combination with Terrasym® 450, a PPFM biostimulant containing *Methylobacterium gregans* and evaluated under commercial-scale conditions using grower standard practices (GSP), which included a range of maize hybrids, management strategies, and pesticide applications as determined by each grower. Across all locations, the combined treatment with GSP resulted in a significant yield increase of +351 kg/ha (5.6 bu/A; $P < 0.001$) and achieved a 72% win rate over GSP alone ($P < 0.001$; Fig. 2b). Importantly, these trials were conducted regardless of actual CRW pressure, enabling assessment of agronomic performance and economic return under real-world agronomic conditions. These results demonstrate that TS201 consistently enhanced yield across diverse genotypes, soil microbiomes, and climatic conditions. The integration of TS201 into crop production systems complements sustainable integrated pest management (IPM) frameworks, which emphasize productivity, resilience, nutrient use efficiency, and reduced ecological footprint (*13*).

Figure 1. Large-scale on-farm evaluation of TS201 + Terrasym 450 in maize during 2023 and 2024 growing seasons across eight U.S. states. a) Geographic distribution of 81 commercial evaluation sites within eight U.S. states: Colorado, Iowa, Illinois, Indiana, Maryland, Minnesota, Nebraska, and Wisconsin. Trials were conducted regardless of corn rootworm pressure to assess real-world agronomic performance. Map background adapted from Google Maps (Map data ©2025 Google) with markers indicating county-level trial locations. b) Piano chart showing yield improvement of TS201 + Terrasym 450 + grower standard practice (GSP) as compared to GSP alone across large-scale on-farm trials (ranging from 4.05 to 40.5 ha per site, n=81).

Mechanisms of action

Prompted by prior observations of reduced CRW feeding damage and plant lodging, we conducted a small-plot field trial in 2024 to evaluate the effects of TS201 on maize root system architecture when applied in combination with Terrasym 450. The trial experienced in low but economically relevant CRW pressure, with a mean NIS of 0.26, slightly above the economic threshold of 0.25 (*10, 11*). Application of TS201 + Terrasym 450 resulted in consistent numerical increases in total root volume, surface area, cumulative root length, and root tip number relative to the untreated control, independent of visible feeding damage (Fig. 2, tables S3 & S4). The most pronounced and statistically significant enhancements were observed in nodal positions 1 through 3, where total root length increased by 25%, primarily driven by a 39% increase in fine root length (≤0.5 mm diameter) (table S4). These findings suggest that TS201 may provide early protection by preserving finer, actively growing roots that are preferentially targeted by CRW larvae (*7, 14*).

Figure 2. Enhanced maize root development observed under TS201 + Terrasym 450 treatment compared to untreated control. Root systems were collected from a small-plot field trial conducted in 2024 (10 plants per plot per replicate, four replicates per treatment). Following excavation and pressure washing, roots were imaged through a structured phenotyping workflow: intact root crowns imaged from below (top panel), then nodal segments were dissected and imaged (bottom panel). Images shown are representative of each treatment.

To investigate the molecular mechanisms by which TS201 influences maize physiology under CRW pressure, we performed RNA-Seq analysis on root tissues collected from a field trial in Illinois. Maize was planted and treated with TS201 in April, and root samples were harvested in mid-July, coinciding with root digs conducted to determine CRW feeding damage. RNA-Seq was conducted on TS201-treated root samples from plots that exhibited both reduced NIS and increased grain yield relative to untreated control plots (table S5). Differential gene expression analysis revealed five genes significantly upregulated in TS201-treated roots compared to controls (table S6), several of which are annotated as being involved in stress response and defense pathways based on the B73 maize reference genome. An additional eight genes were significantly downregulated (table S7). Notably, two of the upregulated genes were associated with the anthranilate biosynthetic pathway, prompting the hypothesis that TS201 treatment may induce the production of methyl anthranilate, a volatile compound previously documented to repel western corn rootworm (WCRW, *Diabrotica virgifera virgifera*) (*15*).

To determine whether TS201 elicits volatile defenses against herbivory, we performed two-choice olfactometer assays using WCRW (*16*), the most invasive and agriculturally significant pest within CRW species (*7, 17*). Although no behavioral differences were detected at 3 days post-treatment, by day 10, TS201-treated roots exhibited robust repellency to WCRW larvae compared to controls (Figs. 3a, c–f). This repellent effect was independent of prior root feeding and required a lag phase of more than three days post-application, suggesting that TS201 triggers volatile-mediated defense mechanisms through induced systemic resistance (ISR) (*18*) rather than damage-induced responses via systemic acquired resistance (SAR) (*19*). Notably, prior herbivory reduced

larval avoidance (Figs. 3b, 3e), indicating that feeding may modulate volatile release or impose a
fitness cost in maize. ISR, mediated by jasmonic acid and ethylene pathways, primes rapid, energy-
efficient defense without constitutive gene activation (*20*). Emerging evidence also indicates that
ISR orchestrates systemic metabolic reprogramming, hormone crosstalk, and oxidative signaling
to enhance plant resilience (*21*). Although PPFMs are known to promote growth and pathogen
resistance via ISR (*22-24*), their role in mitigating insect herbivory has remained largely
unexplored. Our results demonstrate, for the first time, that a PPFM species can elicit a robust ISR
response in maize that deters an herbivorous pest.

Figure 3. Percentage of larvae approaching experimental treatments in two-choice bioassays. Control:
untreated seed without western corn rootworm (WCRW) infestation; TS201: TS201-treated seed without
WCRW infestation; WCRW: untreated seed with WCRW infestation; TS201 + WCRW: TS201-treated
seed with WCRW infestation. Bars with different letters are significantly different ($P < 0.05$). Mean ± SEM
(n=12 independent choice assays).

To identify volatiles associated with WCRW repellency, we analyzed VOCs from TS201-
treated and control plants using gas chromatography–quadrupole time-of-flight mass spectrometry
(GC-QTOF-MS). Of the 18 VOCs detected (tables S8 & S9), methyl anthranilate was significantly
induced by TS201 treatment (Fig. 4). Its relative abundance was markedly higher in TS201-treated
plants, while the remaining 17 VOCs showed only minor or non-significant changes (Fig. 4). VOC
levels were normalized to plant biomass, confirming that increased methyl anthranilate production
was not due to plant size. Notably, this induction occurred independently of WCRW herbivory,
with elevated emissions observed even in the absence of insect feeding. This emission pattern
corresponded with the insect choice assays conducted on the same set of plants, where significantly

fewer WCRW larvae were recovered from TS201-treated plants (fig. S1). Assays with purified
methyl anthranilate further validated its strong, dose-dependent repellency against WCRW, with
no significant repellency observed in untreated or solvent controls (fig. S2).

Figure 4. TS201 induces methyl anthranilate emission in maize. A list of all detected volatiles is available
at Supplementary Table 7. Mean ± SEM (n ≥ 4). Untransformed data were presented, whereas analyses
were performed with log10 transformed data. Signficant differences between the control and treated plants
are indicated by *($P < 0.05$), **($P<0.01$), and ***($P<0.001$).

Our results provide compelling evidence that TS201 activates the anthranilate biosynthetic
pathway, leading to the production of methyl anthranilate, a volatile compound previously
documented to repel WCRW (Fig. 5) (*15*). This mechanism represents a major advancement in
IPM strategies for CRW, a pest complex that has evolved resistance to nearly all existing control
strategies (*25-27*). Unlike conventional insecticides and Bt traits that impose direct lethal pressure
and drive rapid resistance evolution, TS201 primes plant defenses through ISR, triggering
endogenous pathways without direct toxicity, thereby minimizing selection pressure and
potentially preserving non-target organisms (*28-30*). While biological control agents including
Heterorhabditis bacteriophora and various biostimulants have shown variable efficacy and
inconsistent field performance (*31-34*), TS201 consistently enhances maize protection across
diverse agroecosystems, independent of pest pressure or environmental variability. These findings
establish TS201 as a scalable, ecologically rational addition to sustainable CRW management,
delivering strategic compatibility with current IPM frameworks and consistent efficacy across
diverse geographic regions.

Figure 5. Schematic of TS201-induced enhancement of maize defense against western corn rootworm.

Plant-associated microbiomes represent a promising avenue for advancing climate-resilient, pest-tolerant cropping systems. Microbe-induced priming may concurrently enhance yield and strengthen plant defense under real-world agronomic conditions. Bridging field performance with mechanistic insight could accelerate the development of microbial solutions that reengineer plant resilience through ecologically informed partnerships for sustainable agriculture.

REFERENCES AND NOTES

1. M. del Carmen Orozco-Mosqueda, M. del Carmen Rocha-Granados, B. R. Glick, G. Santoyo, Microbiome engineering to improve biocontrol and plant growth-promoting mechanisms. *Microbiol. Res.* **208**, 25-31 (2018).
2. C. Brito-Lopez, N. van der Wielen, M. Barbosa, R. Karlova, Plant growth–promoting microbes and microalgae-based biostimulants: sustainable strategy for agriculture and abiotic stress resilience. *Philos. Trans. B* **380**, 20240251 (2025).
3. A. Khoulati *et al.*, Harnessing biostimulants for sustainable agriculture: innovations, challenges, and future prospects. *Discov. Agric.* **3**, 56 (2025).
4. S. E. Lindow, M. T. Brandl, Microbiology of the phyllosphere. *Appl. Environ. Microbiol.* **69**, 1875-1883 (2003).
5. U. Kutschera, Plant-associated methylobacteria as co-evolved phytosymbionts: a hypothesis. *Plant Signal. Behav.* **2**, 74-78 (2007).
6. C. Zhang, M.-Y. Wang, N. Khan, L.-L. Tan, S. Yang, Potentials, utilization, and bioengineering of plant growth-promoting *Methylobacterium* for sustainable agriculture. *Sustainability* **13**, 3941 (2021).
7. H. C. Chiang, Bionomics of the northern and western corn rootworms. *Annu. Rev. Entomol.* **18**, 47-72 (1973).

8. B. P. Spike, J. J. Tollefson, Yield response of corn subjected to western corn rootworm (Coleoptera: Chrysomelidae) infestation and lodging. *J. Econ. Entomol.* **84**, 1585-1590 (1991).

9. E. Levine, H. Oloumi-Sadeghi, Management of *diabroticite* rootworms in corn. *Annu. Rev. Entomol.* **36**, 229-255 (1991).

10. J. D. Oleson, Y.-L. Park, T. M. Nowatzki, J. J. Tollefson, Node-injury scale to evaluate root injury by corn rootworms (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* **98**, 1-8 (2005).

11. B. Wright, Dig and rate roots to evaluate corn rootworm control. University of Nebraska-Lincoln Extension. https://cropwatch.unl.edu/dig-and-rate-roots-evaluate-corn-rootworm-control-0/ (2008).

12. EPA, Summary of the Federal Insecticide, Fungicide, and Rodenticide Act. Washington, D.C., United States: U. S. Environmental Protection Agency. www.epa.gov/laws-regulations/summary-federal-insecticide-fungicide-and-rodenticide-act (1996).

13. E. K. Mitter, M. Tosi, D. Obregón, K. E. Dunfield, J. J. Germida, Rethinking crop nutrition in times of modern microbiology: innovative biofertilizer technologies. *Front. Sustain. Food Syst.* **5**, 606815 (2021).

14. S. Strnad, M. Bergman, Distribution and orientation of western corn rootworm (Coleoptera: Chrysomelidae) larvae in corn roots. *Environ. Entomol.* **16**, 1193-1198 (1987).

15. E. J. Bernklau, B. E. Hibbard, A. P. Norton, L. B. Bjostad, Methyl anthranilate as a repellent for western corn rootworm larvae (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* **109**, 1683-1690 (2016).

16. T. F. Branson, E. E. Ortman, Feeding behavior of larvae of the western corn rootworm: normal larvae and larvae maxillectomized with laser radiation. *Ann. Entomol. Soc. Am.* **62**, 808-812 (1969).

17. A. Eben, Ecology and evolutionary history of *Diabrotica* beetles-overview and update. *Insects* **13**, 156 (2022).

18. C. M. Pieterse *et al.*, Induced systemic resistance by beneficial microbes. *Annu. Rev. Phytopathol.* **52**, 347-375 (2014).

19. W. E. Durrant, X. Dong, Systemic acquired resistance. *Annu. Rev. Phytopathol.* **42**, 185-209 (2004).

20. A. Martinez-Medina *et al.*, Recognizing plant defense priming. *Trends Plant Sci.* **21**, 818-822 (2016).

21. Y. Yu *et al.*, Induced systemic resistance for improving plant immunity by beneficial microbes. *Plants* **11**, 386 (2022).

22. M. Madhaiyan *et al.*, Plant growth–promoting *Methylobacterium* induces defense responses in groundnut (*Arachis hypogaea* L.) compared with rot pathogens. *Curr. Microbiol.* **53**, 270-276 (2006).

23. M. Madhaiyan *et al.*, Growth promotion and induction of systemic resistance in rice cultivar Co-47 (*Oryza sativa* L.) by *Methylobacterium* spp. *Bot. Bull. Acad. Sin.* **45**, (2004).

24. R. Poorniammal, S. Sundaram, K. Kumutha, *In vitro* biocontrol activity of *Methylobacterium extorquens* against fungal pathogens. *Int. J. Plant Prot.* **2**, 59-62 (2009).

25. A. J. Gassmann, Resistance to Bt maize by western corn rootworm: Effects of pest biology,
the pest–crop interaction and the agricultural landscape on resistance. *Insects* **12**, 136
(2021).
26. K. J. Paddock, C. A. Robert, M. Erb, B. E. Hibbard, Western corn rootworm, plant and
microbe interactions: a review and prospects for new management tools. *Insects* **12**, 171
(2021).
27. Z. Ye *et al.*, Too much of a good thing: Lessons from compromised rootworm Bt maize in
the US Corn Belt. *Science* **387**, 984-989 (2025).
28. J. Köhl, R. Kolnaar, W. J. Ravensberg, Mode of action of microbial biological control
agents against plant diseases: relevance beyond efficacy. *Front. Plant Sci.* **10**, 845 (2019).
29. G. Bonanomi, M. Lorito, F. Vinale, S. L. Woo, Organic amendments, beneficial microbes,
and soil microbiota: toward a unified framework for disease suppression. *Annu. Rev.*
Phytopathol. **56**, 1-20 (2018).
30. S. Timmusk, L. Behers, J. Muthoni, A. Muraya, A.-C. Aronsson, Perspectives and
challenges of microbial application for crop improvement. *Front. Plant Sci.* **8**, 49 (2017).
31. Š. Modic, P. Žigon, A. Kolmanič, S. Trdan, J. Razinger, Evaluation of the field efficacy of
Heterorhabditis bacteriophora Poinar (Rhabditida: Heterorhabditidae) and synthetic
insecticides for the control of western corn rootworm larvae. *Insects* **11**, 202 (2020).
32. S. Toepfer *et al.*, A critical evaluation of host ranges of parasitoids of the subtribe
Diabroticina (Coleoptera: Chrysomelidae: Galerucinae: Luperini) using field and
laboratory host records. *Biocontrol Sci. Technol.* **18**, 483-504 (2008).
33. S. I. Tarigan, J. Kiss, T. György, N. P. Y. Doan, S. Toepfer, Effects of microbial
biostimulants on maize and its pest, the western corn rootworm, *Diabrotica virgifera*
virgifera. *Agronomy* **14**, 2239 (2024).
34. G. Jaffuel *et al.*, Protecting maize from rootworm damage with the combined application
of arbuscular mycorrhizal fungi, *Pseudomonas* bacteria and entomopathogenic nematodes.
Scientific reports **9**, 3127 (2019).

ACKNOWLEDGMENTS

The authors thank the many NewLeaf employees both past and present who have worked
on this project over the years including Vanessa Pugliessa for making the summary figure, Dayna
Collett for running small plot trials 2018-2022, Roger Bowman for trial work in 2023-2024 and
Ashley Haddox for supporting root phenotyping in 2024. We would also like to thank our trialing
partners and contract researchers, especially Jay Pershing who consulted on this project for several
years providing his extensive expertise in corn rootworm management. Funding for this project
was provided by NewLeaf Symbiotics. This article reports the results of research only. Mention of
trade names or commercial products is solely for the purpose of providing specific information
and does not imply recommendation or endorsement by the USDA or the University of Missouri.
USDA is an equal opportunity provider and employer.

Author contributions:

Conceptualization: M.P.H., B.E.H., and N.W.B.

Methodology: M.P.H., B.E.H., A.J., and N.W.B.

Investigation: M.P.H., K-V.H., A.Z.P., A.F.E., D.C.L., A.J., and N.W.B.

Visualization: M.P.H., K-V.H., A.Z.P., and N.W.B.

Funding acquisition: M.P.H., D.F., B.E.H., and N.W.B.

Project administration: M.P.H., B.E.H., and N.W.B.

Supervision: M.P.H., B.E.H., and N.W.B.

Writing-original draft: M.P.H. and N.W.B.

Writing-review & editing: all authors.

Competing interests: Anne Phillips, Allison Jack and Natalie Breakfield are employees of
NewLeaf Symbiotics, St. Louis, MO, USA. All other authors declare no competing interests.

Data and materials availability: All raw transcriptomic sequencing data has been deposited
in the Sequence Read Archive database on NCBI with BioSample Accessions
SAMN48399254, SAMN48399255, SAMN48399256, SAMN48399257, SAMN48399258,
SAMN48399259 under BioProject accession number PRJNA1260573. All other pertinent data
are found in the figures and tables. Request for data and additional information should be
submitted to the corresponding author.

SUPPLEMENTARY MATERIALS:

Materials and Methods

Figs. S1 to S2

Tables S1 to S9

References (35–59)

Supplementary Materials for

Harnessing microbes for crop production: TS201 enhance maize yield and reduces pest damage

Man P. Huynh *et al.*

Corresponding authors: Man P. Huynh, manhuynh@missouri.edu; Natalie W. Breakfield, nbreakfield@newleafsym.com

The PDF file includes

MATERIALS AND METHODS

Biological materials

The EPA registered biopesticide product TS201, containing *Methylorubrum extorquens* strain NLS0042 (2.0%, 1×10^9 CFU/g) as an active ingredient, was provided by NewLeaf Symbiotics (St. Louis, MO). This bacterial strain was originally sourced from a soybean plant in Saint Louis County, MO. Maize hybrids (P1197, Pioneer, Johnston, IA) were purchased from local seed suppliers unless otherwise indicated.

Western corn rootworm (WCRW, *Diabrotica virgifera virgifera*) neonates were obtained from a non-diapausing colony maintained at the Plant Genetics Research Unit-USDA-ARS in Columbia, MO (*35, 36*). The eggs were obtained in Petri dishes consisting of eggs and soil and were incubated at 25 ± 1 °C in darkness in an incubator (Percival, Perry, IA). Neonates that hatched within 24 h were used for insect assays.

Field experiments

From 2016 to 2022, a series of 22 small-plot field trials were conducted across multiple locations in the U.S. Midwest with documented histories of elevated WCRW and northern corn rootworm (NCRW, *D. barberi*) damage, capturing variability in climate, soil type, and pest pressure. TS201 was evaluated as a standalone treatment to establish proof of concept under diverse field environments. During this pre-registration phase, trials were conducted under USDA-APHIS jurisdiction, with testing limited to a maximum of 4.05 ha (10 acres) per year in compliance with U.S. federal guidelines while U.S. Environmental Protection Agency (EPA) registration was in process.

The small-plot field experiments were designed as a randomized complete block design (RCBD) with four row plots 12 m (40 ft) in length and 4 to 6 replicates per location. TS201 was applied via multiple delivery methods to evaluate its efficacy and consistency under agronomically relevant deployment applications, including in furrow (309 g/ha, 1×10^9 CFU/g wettable powder), liquid seed treatment (1×10^6 CFU/seed application rate, 1×10^9 CFU/g wettable powder) or planter box application (0.5 oz/ unit seed or 14 g/ 8×10^4 maize seed, 1×10^9 CFU/g powder with seed lubricity agent). In each trial, the cooperating contract researcher selected the maize hybrid and agronomic management practices with base insecticide treatments specifically excluded, and these plots served as the untreated control. TS201 application was the sole treatment variable distinguishing the treated plots from the untreated control. In a subset of these trials, the same maize hybrids were evaluated with and without belowground corn rootworm (CRW) protection traits such as Bt traits, with TS201 application was the only difference from the untreated control. Trials were conducted under continuous maize cultivation (corn-on-corn rotations) with select sites incorporating squash trap crops to facilitate consistent CRW feeding pressure. Root damage was evaluated in mid-July to early August, corresponding with the observed emergence of adult CRW

beetles on maize plants, using the Iowa State University NIS system. To determine the efficacy of TS201 under CRW pressure, only field trials in which untreated control plots exhibited an NIS score greater than 0.25, a common economic damage threshold, were analyzed. Grain yield data were collected as bu/A and converted to kg/ha using the conversion of 1 bushel as 25.4 kg (*37*).

In 2020, an on-farm experiment was carried out to evaluate the impact of TS201 on maize lodging under CRW pressure. The trial included nine replicates in distinct locations in IA, IL, MN, and CO. At each site, 0.1 ha (¼-acre) strips were assigned to compare the untreated control with an in-furrow application of TS201 (309 g/ha, 1 x 10^9 CFU/g). TS201 was applied to a corn hybrid without CRW protection traits. Roots were sampled along two defined transects (37 m) in each treatment, 20 roots total per treatment, pressure washed and rated using the NIS system. Out of nine locations, two in IA exhibited larval root injury exceeding the economic threshold (NIS > 0.25) and were included in lodging analysis. Based on NIS values, trial sites were categorized into low (0.25–1), moderate (1–2), or high (2–3) levels of CRW pressure levels. Lodging was quantified as the percentage of lodged plants within two, 37 m predefined transects for each treatment.

Following EPA registration, field evaluations of TS201 shifted from small-plot studies to large-scale, on-farm evaluations under commercial production conditions during the 2023 and 2024 growing seasons. The large-scale field experiment included 81 replicates, each representing a maize field at different locations within eight U.S. states (Fig. 2a). Across 81 trials (≥ 4.05 ha each), conducted in collaboration with contract researchers and commercial growers, TS201 was applied in combination with Terrasym 450 as a planter box treatment (0.5 oz per unit of seed of each PPFM isolate at 1 x 10^9 CFU/g with seed lubricity agent) as an addition to the grower standard practice (GSP) and evaluated in comparison to GSP alone for CRW management. GSP reflected real-world production systems incorporating diverse IPM strategies, including combinations of belowground CRW protection traits (e.g., Bt, RNAi), in-furrow insecticides, crop rotation, and prior-season aerial applications targeting adult beetles. Trials were primarily conducted in corn-on-corn systems in regions with a reported history of CRW pressure. Experimental designs included full-field side-by-sides, split-planter setups, and replicated strips ranging from 4.05 ha to over 40.5 ha. Win rate was determined by dividing the number of trials with a positive yield over GSP by the total number of trials and converting to a percentage. Yield data were collected using precision monitors or weigh wagons and converted from bu/A to kg/ha as described above. Data were expressed as differences relative to the GSP control.

In 2024, building on prior observations of reduced CRW feeding damage and plant lodging, a small-plot field trial was conducted to evaluate the impact of TS201 on maize root system architecture when applied in combination with Terrasym 450. The trial was designed as a RCBD with four replicates. To determine baseline CRW pressure, roots from untreated control plots were sampled, pressure washed, and scored using the NIS system. Three weeks later, 10 root crowns per replicate were collected from both TS201 and Terrasym 450 treated and control plots to assess

root architecture under CRW pressure. Roots were imaged using a structured phenotyping
workflow: intact crowns imaged from below in the transverse plane and dissected nodal segments
imaged per individual crown to enable accurate imaging of total root length, root size classes, and
numbers of root tips. Root phenotyping images were analyzed using RhizoVision Explorer v2.0.3
(*38*). All images included a size marker for normalization and to convert pixel dimensions to metric
units (mm). Dissected roots were grouped by node: 1-3, 4, 5, and 6. No node 7 roots were present
in this study. Yield data were not collected, as the trial employed destructive sampling for detailed
root phenotyping.

Transcriptomic analysis

Root tissue samples from untreated and TS201-treated maize plants (one sample per plot;
six replicates per treatment) were collected from different RCBD plots in a field trial. Total RNA
was extracted using RNeasy Plant Mini Kits (74104, Qiagen, Germantown, MD), treated with
DNase using the RNase-Free DNase Set (Catalog #15200, Qiagen), and assessed for RNA quality
using an Agilent 2100 Bioanalyzer with RNA 6000 Nano Kits (5067-1511, Agilent Technologies,
Santa Clara, CA).

Root samples from TS201-treated plots that exhibited both increased yield and reduced
nodal injury scores (NIS) compared to control plots (three plots total) were used for gene
expression profiling using QuantSeq FWD 3′ mRNA-Seq using 75bp reads with V2 Illumina
chemistry (Lexogen, Vienna, Austria). QuantSeq is a tag-based sequencing method that captures
and sequences the 3′ end of polyadenylated transcripts, allowing accurate gene expression
quantification via read counting (*39*). The B73 v.4 maize reference genome was used for read
alignment. Differential gene expression analysis was performed using the BlueBee Genomics
platform which utilizes DESeq2; genes were considered differentially expressed if the $\log_2$ (fold
change) was greater than 1.5 and the adjusted *P*-value was less than 0.1 (*40*). All raw sequencing
data have been deposited in the NCBI Sequence Read Archive database with BioProject accession
number PRJNA1260573.

Insect choice assays

Untreated maize seeds were rinsed and incubated in distilled water overnight in a Percival
incubator maintained at $25 \pm 1°C$ to promote germination. After incubation, seeds were dried out
and treated with TS201 at a concentration of 1×10^6 CFU/seed, or with sterile distilled water (0.01
mL/seed, untreated seed). Subsequently, 3 kernels were planted per 50-mL conical tube and grown
in the Percival incubator for 10 days. Plants were watered with 2 mL of distilled water every other
day. The experiment included four treatment groups: untreated seed without WCRW infestation
(control), serving as the control, TS201-treated seed without WCRW infestation, untreated seed
with WCRW infestation, and TS201-treated seed with WCRW infestation. To allow for ISR
priming to occur, seeds were allowed to germinate and grow for 7 days prior to insect infestation.
Insect feeding was designed for 3-days to allow the potentiated ISR mediated-defense response to

occur, if required. On day 7, 8 WCRW neonates (<24 h old) were transferred to plants assigned to infestation treatments. On day 10, plants were removed from the soil using a fine paintbrush and used for insect choice assays.

Insect choice assays were conducted using a modified olfactometer system adapted from Branson and Ortman (*41*). The olfactometer consisted of three 50×15 mm Petri dishes connected in a linear arrangement via 12 mm segments of Teflon tubing (8 mm I.D. $\times$ 10 mm O.D.). The bioassays were designed as a completely randomized experiment. Ten WCRW neonates were introduced into the central arena, while treatment and control plants were randomly assigned to the terminal arenas at either end of the olfactometer. The experiment was conducted in complete darkness in the Percival incubator, and larval positions were recorded after 30 minutes to assess behavioral responses. The experiment was run in three cohorts at different times, with each cohort consisting of four replicates, for a total of 12 replicates. Each replicate used unexposed larvae and freshly prepared plant seedlings.

Larval behavioral response was quantified by calculating the proportion of larvae that responded (PLR), using the formula: PLR = Nr / Nt, where *Nr* is the number of larvae found in each zone (treatment side, control side, or infested center) at the end of the assay (30 min), and *Nt* is the total number of larvae introduced.

GC-MS analysis

Volatile organic compounds from control and TS201-treated maize plants were collected using a modified headspace sampling system based on the method described previously (*42*). Briefly, maize seeds were treated with TS201 or sterile water, and the treated seeds were planted without WCRW infestation in 50-mL conical tubes as previously described. At 10 days post-germination, seedlings were gently removed from the soil using a fine paintbrush. Maize seedlings were transferred into a 50-mL collection chamber (5 seedlings per chamber) fitted with Teflon tubing at both ends. One end of each chamber was connected to a flow meter (B07B6HQDDW, Wuhan, China), which was linked to a Whatman® carbon/HEPA filter capsule (WHA67041500, Sigma-Aldrich, St. Louis, MO) and a 6-arm air delivery pump. The opposite end was connected to a Tenax® TA thermal desorption tube (35/60 mesh, 0.25 in. O.D. $\times$ 3.5 in. length, Tenax® TA, 29747-U, Sigma-Aldrich) for VOC trapping. Air was drawn through the chamber at a flow rate of 2 mL/min for 90 minutes to collect volatiles. VOCs from blank chambers containing sterile water and purified methyl anthranilate (≥98% purity, W268224, Sigma-Aldrich) were used as negative and positive controls, respectively. Adsorbed compounds were eluted from the Tenax tubes with 2 mL of methanol (Optima™ LC/MS Grade, A456, Fisher Chemical, Waltham, MA) for GC-MS analysis. The volatile assay was conducted as a completely randomized experiment with more than 4 replicates. Each replicate consisted of freshly prepared, unexposed maize seedlings randomly assigned to individual chambers within the headspace sampling system.

162   Headspace volatiles were analyzed using an Agilent 7890B gas chromatograph (GC, Agilent Technologies, Santa Clara, CA) coupled to an Agilent 7250 quadrupole time-of-flight mass spectrometer (QTOF-MS). Sample injection was performed using a Gerstel multipurpose sampler. The GC-MS operating conditions were as follows: helium was used as the carrier gas at a constant flow rate of 1 mL/min; injection volume was 1 µL with a split ratio of 5:1; and the inlet temperature was maintained at 280 °C. The oven temperature program began at 50 °C (held for 2 minutes), ramped at 6 °C/min to 215 °C, then increased at 5 °C/min to 315 °C, where it was held for 12 minutes before cooling back to 50 °C. Separation was performed using an Agilent DB-5MS capillary column (60 m × 250 µm i.d. × 0.25 µm film thickness). The MS transfer line was maintained at 250 °C, and electron ionization (EI) was conducted at 70 eV. Mass spectra were acquired in full-scan mode over a mass range of 40-650 m/z.

173   Data processing, including deconvolution, alignment, and compound identification, was performed using MS-DIAL software, with spectral matching against an in-house mass spectral library. The peak areas of identified compounds were integrated using Agilent MassHunter Qualitative Analysis 10.0 and normalized to plant biomass (mg) to determine their relative concentrations.

Insect choice assays with methyl anthranilate

179   To determine the behavioral response of WCRW larvae to methyl anthranilate (MA), insect choice assays were conducted using the olfactometer system described above. In each assay, purified methyl anthranilate (≥98% purity; Sigma-Aldrich) was applied to one side of the olfactometer arena at concentrations of 0 (solvent control, methanol), 0.1, 1, or 10 µg with and without untreated maize seedlings, while the opposite side contained untreated maize seedlings. Ten neonate WCRW larvae (<24 h old) were introduced into the central arena of the olfactometer, and larval distribution was recorded after 30 minutes in complete darkness. Each treatment was replicated at least five times. The proportion of larvae responding was calculated as described above to determine the repellent effect of methyl anthranilate at each concentration.

Statistical analysis

189   The VOC data were analyzed using Student's unpaired t-test (PROC TTEST, SAS 9.4, SAS Institute, Cary, NC). Compounds were considered significantly induced when $P < 0.05$ and $\log_{10}$(fold change) > 1. Other data (i.e., small-plot field trial, root phenotyping, large-scale field trial, insect choice data) were analyzed using analysis of variance (ANOVA) followed by pairwise comparisons of least squares means (LSMEANs) or Fisher's least significant difference (LSD) for multiple comparisons. Analyses were conducted using a generalized linear mixed model in PROC GLIMMIX (SAS 9.4) or JMP v18.1.1 (JMP Statistical Discovery LLC, Cary, NC), with treatment specified as a fixed effect and year, location, block, or replication as random effects. For win rate analysis, trial-level outcomes were treated as binary responses (win = 1, loss = 0) and modeled using a binomial distribution. Normality and homogeneity of variance were evaluated using

199 univariate analysis in PROC UNIVARIATE (SAS 9.4). Datasets that did not meet these
200 assumptions were transformed prior to analysis: percentage variables were arcsine square-root
201 transformed, and numeric variables were square-root transformed, to meet the corresponding
202 distributional requirements.

204 **SUPPLEMENTARY FIGURES**

205
206 **Figure S1.** Percentage of larvae approaching experimental treatments in two-choice bioassays.
207 Bars with different letters are significantly different ($P < 0.05$). Mean ± SEM (n=17 independent
208 choice assays).

Figure S2. Percentage of larvae approaching experimental treatments in two-choice bioassays. Bars with different letters are significantly different ($P < 0.05$). Mean ± SEM (n=10 independent choice assays). MA: methyl anthranilate. Methyl anthranilate exhibited a strong, dose-dependent repellent effect against WCRW larvae, with few or no larvae found in its treatments. No significant difference was observed between untreated seeds and those with the solvent blank added.

 SUPPLEMENTARY TABLES

Table S1. Effects of TS201 on agronomic performance and corn rootworm damage relative to untreated control (UTC). Field trials were conducted over seven growing seasons (2016-2022) at 22 locations across the U.S. Midwest. Data represent results from 22 small plot field trials. *Least square mean values when only TS201 and control were used in the linear mixed effects model; [+] Least square mean values when TS201, control, and all other treatments (i.e., other examined PPFMs, insecticides) in the trials were used in the linear mixed effects model.

Treatment	Yield				Nodal injury score			
	Kg/ha	Std Error	Difference from control	P-value	Score	Std Error	Difference from control	P-value
Control	11,706	191.4			1.34	0.068		
TS201*	11,926	190.8	+220 kg/ha (3.5 bu/A)	0.0108	1.23	0.069	-0.11	0.0007
Control	11,593	302.6			1.40	0.147		
TS201[+]	11,813	301.9	+220 kg/ha (3.5 bu/A)	0.0760	1.29	0.146	-0.11	0.0217

223 **Table S2.** Effects of TS201 on agronomic performance and corn rootworm damage by corn
224 rootworm (CRW) belowground traits relative to untreated control. *Data represent results from 22
225 small plot field trials in 2016-2022; [+]Data represents results from 14 small plot trials over 2016-
226 2019 and 2021.

Maize hybrids	Treatment	Yield				Nodal injury score			
		Kg/ha	Std Error	Difference from control	P-value	Score	Std Error	Difference from control	P-value
Without belowground CRW traits*	Control	11,492	210.3	+258 kg/ha (4.1 bu/A)		1.51	0.073		
	TS201	11,750	209.7		0.0309	1.40	0.073	-0.11	0.0014
With belowground CRW traits[+]	Control	12,296	217.2	+188 kg/ha (3.0 bu/A)		0.97	0.092		
	TS201	12,484	215.9		0.1326	0.87	0.091	-0.10	0.0063

228 **Table S3.** Full root crowns: Root phenotypes associated with the application of TS201 and
229 Terrasym® 450 compared to an untreated control in a single location small plot trial with a
230 baseline NIS of 0.26 in untreated control plots.

Metric	Control	TS201 + Terrasym 450	Pct. Diff
Solidity (Network Area: Convex area)	0.4119	0.4123	+0.10
Network Area (total pixels)	29058.1	31002.2	+6.69
Convex Area	71584.2	78179.9	+9.21
Surface Area	233119.6	251782.5	+8.01
Volume	455752.9	510522.2	+12.02
Perimeter	16377.5	16975.0	+3.65

231 Units are in mm unless the number represents a ratio. All comparisons are not significant.

Table S4. Dissected roots: Root phenotypes associated with the application of TS201 and
Terrasym 450 compared to an untreated control in a single location small plot trial with a
baseline NIS of 0.26 in the UTC plots.

Node(s)	Metric	Control	TS201	Pct Diff fm ctrl	P-value
123	Number of Root Tips	1219.48	1601.53	*31.33**	0.0519
	Total Root Length	**11381.36**	**14251.61**	*25.22***	**0.0376**
	Branching Frequency per mm	0.24	0.24	3.51	0.1804
	Volume	15111.81	18751.49	*24.08**	0.0511
	Surface Area	**37838.75**	**46406.15**	*22.64***	**0.0418**
	Average Diameter	1.08	1.05	-2.91	0.3306
	Total root length: diameters 0.0-0.5mm	**2398.85**	**3332.01**	*38.90***	**0.0037**
	Total root length: diameters 0.5-2.0mm	7859.23	9575.07	*21.83**	0.0934
	Total root length: diameters 2.0-5.0mm	1096.16	1297.49	18.37	0.1630
	Total root length: diameters 5.0+mm	27.12	47.03	73.42	0.2058
4	Number of Root Tips	1930.93	2130.83	10.35	0.3826
	Total Root Length	18434.72	20139.92	9.25	0.4511
	Branching Frequency per mm	0.29	0.29	-0.27	0.9140
	Volume	33572.35	41919.52	24.86	0.1134
	Surface Area	66204.58	76888.42	16.14	0.2481
	Average Diameter	**1.16**	**1.23**	*5.47***	**0.0228**
	Total root length: diameters 0.0-0.5mm	4933.37	5078.32	2.94	0.6369
	Total root length: diameters 0.5-2.0mm	11034.75	11945.99	8.26	0.5899
	Total root length: diameters 2.0-5.0mm	2256.71	2804.16	*24.26**	0.0950
	Total root length: diameters 5.0+mm	209.90	311.45	48.38	0.1100
5	Number of Root Tips	3097.38	3489.55	12.66	0.2035
	Total Root Length	29739.06	32068.50	7.83	0.2887
	Branching Frequency per mm	0.30	0.31	0.22	0.6446
	Volume	**73582.43**	**78565.72**	*6.77***	**0.0137**
	Surface Area	120045.50	128068.52	6.68	0.1546
	Average Diameter	1.30	1.28	-1.36	0.6199
	Total root length: diameters 0.0-0.5mm	8199.80	8909.41	8.65	0.3791

	Total root length: diameters 0.5-2.0mm	16218.40	17455.97	7.63	0.3497
	Total root length: diameters 2.0-5.0mm	4583.48	4939.69	7.77	0.1555
	Total root length: diameters 5.0+mm	737.38	763.44	3.53	0.2112
6	Number of Root Tips	1539.60	1443.00	-6.27	0.8086
	Total Root Length	16261.98	14517.62	-10.73	0.6537
	Branching Frequency per mm	0.25	0.25	-1.55	0.6977
	Volume	52474.57	47981.96	-8.56	0.6999
	Surface Area	75456.17	68772.79	-8.86	0.7014
	Average Diameter	1.65	1.69	2.45	0.7914
	Total root length: diameters 0.0-0.5mm	3940.54	3348.43	-15.03	0.5194
	Total root length: diameters 0.5-2.0mm	8519.04	7600.73	-10.78	0.6673
	Total root length: diameters 2.0-5.0mm	3263.54	3097.09	-5.10	0.8145
	Total root length: diameters 5.0+mm	538.86	471.37	-12.53	

Units are in mm unless the number represents a ratio. Significant interactions are bold. * $P < 0.1$; **$P < 0.05$.

237 **Table S5.** Field phenotypes of roots used in the RNAseq experiment.

Treatment	Yield (kg/ha)	Std Error	Difference from control	P-value	Nodal injury score	Std Error	Difference from control	P-value
Control	10,474	1479			0.35	0.08		
TS201	13,851	306	+3,377	0.09	0.14	0.09	-0.21	0.15

239 **Table S6.** Up-regulated genes in TS201-treated maize roots relative to untreated control roots.
240 TF= transcription factor, ROS=reactive oxygen species, ET=ethylene.

v4 Gene Model ID	Gene Symbol	Full Name	Putative Function	log2 Fold change	Adjusted P-value
Zm00001d050455	pco096884	anthranilate N-benzoyltransferase protein 1	Anthranilate synthesis	1.63	0.0147
Zm00001d003395	ans1	anthranilate synthase homolog1	Anthranilate synthesis	1.55	0.0957
Zm00001d002799	hb66	Homeobox-transcription factor 66	TF in development	2.07	0.0995
Zm00001d049860	nactf75	NAC-transcription factor 75	TF in drought stress tolerance	1.73	0.0957
Zm00001d023617	lac23	laccase23	Enzyme in lignification/ defense	2.18	0.0957
Zm00001d024852	acco3	1-aminocyclopropane-1-carboxylate oxidase3	Enzyme in ET production	2.07	0.0408
Zm00001d002434	aox3	alternative oxidase3	Enzyme in ROS stress tolerance/ defense	1.85	0.0578
Zm00001d002052	NA	Probable isoaspartyl peptidase/ L-asparaginase 2	Nitrogen metabolism/ defense	2.48	<0.0001
Zm00001d011319	ZmABCG48	Probable ABCG Transporter 48	Transport	1.78	0.0408

242 **Table S7.** Down regulated differentially expressed genes in TS201-treated roots.

v4 Gene Model ID	Gene Symbol	Full Name	log2 Fold change	Adjusted P-value
Zm00001d015307	NA	Uncharacterized protein	-2.23	0.006
Zm00001d017454	NA	NA	-2.08	0.058
Zm00001d013259	zmm15	Zea mays MADS-box 15	-2.07	0.058
Zm00001d034045	mads4	MADS-transcription factor 4	-2.04	0.048
Zm00001d027763	IDP1425	plant-specific domain TIGR01615 family protein	-2.03	0.007
Zm00001d015092	NA	NA	-1.89	0.096
Zm00001d032457	NA	NA	-1.86	0.041
Zm00001d048137	NA	NA	-1.86	0.015
Zm00001d007045	pco137705	protein coding	-1.85	0.041
Zm00001d029636	cl945_1	protein coding	-1.82	0.015
Zm00001d035304	limtf10	LIM-transcription factor 10	-1.75	0.006
Zm00001d034621	pco116267	Protein trichome birefringence-like 32coding	-1.68	0.100
Zm00001d043477	cesa11	cellulose synthase 11	-1.65	0.096
Zm00001d009506	NA	NA	-1.58	0.100
Zm00001d026287	bnlg2190	UDP-N-acetylglucosamine diphosphorylase, bnlg2190	-1.52	0.096
Zm00001d043414	NA	NA	-1.52	0.058
Zm00001d000133	NA	NA	-1.45	0.096
Zm00001d019670	mmp187	kinesin-like protein KIN-4A	-1.44	0.096

244 **Table S8.** Volatile compounds annotated via GC-MS analysis.

Compound	Formula	Retention time	Quant Mass	log10 FoldChange	P-value
1-Octen-3-ol	$C_8H_{16}O$	9.415	57.03320782	1.13	0.3195
Decane	$C_{10}H_{22}$	9.885	57.06964660	0.96	0.4231
(Z)-3-Hexenyl acetate	$C_8H_{14}O_2$	10.011	67.05367851	1.03	0.8210
Tetradecane	$C_{14}H_{30}$	10.191	57.06973394	0.88	0.0040
Tetrahydro-β-citronellene	$C_{10}H_{20}$	10.449	57.06957392	0.99	0.8237
p-Cymene	$C_{10}H_{14}$	10.719	119.0841019	0.98	0.8453
R-(+)-Limonene	$C_{10}H_{16}$	10.844	67.05382538	1.00	0.9776
Dihydrocitronellol	$C_{10}H_{22}O$	11.924	69.06966019	0.88	0.0018
α-Terpinene	$C_{10}H_{16}$	12.358	93.06890615	1.02	0.8063
Undecane	$C_{11}H_{24}$	12.552	57.06959788	0.93	0.0437
Dodecane	$C_{12}H_{26}$	15.37	57.06970382	0.95	0.0750
Pentadecane	$C_{15}H_{32}$	17.444	71.08510437	0.95	0.0493
Tritriacontane	$C_{33}H_{68}$	18.705	57.06973076	1.01	0.8138
Methyl anthranilate	$C_8H_9NO_2$	19.502	92.02540970	2.93	<0.0001
Hexacosane	$C_{26}H_{54}$	24.173	71.08522415	1.17	0.1235
Dotriacontane	$C_{32}H_{66}$	24.417	71.08522288	1.16	0.0125
Octacosane	$C_{28}H_{58}$	28.066	57.06988525	1.08	0.2463
Hexedecane	$C_{16}H_{34}$	29.027	71.08506775	0.97	0.5955

246 **Table S9.** Bioactivities of volatile compounds detected from maize.

Compound	Pathway	Sub-pathway	Bioactivities	References
1-Octen-3-ol	Fatty acid metabolism	Oxylipin pathway	Induction of plant defense, insecticidal activity	(43); (44)
Decane	Fatty acid metabolism	Alkane biosynthesis	.	NA
(Z)-3-Hexenyl acetate	Fatty acid metabolism	Lipoxygenase pathway	Induction of plant defense, insect host and oviposition preference	(45); (46)
Tetradecane	Fatty acid metabolism	Alkane biosynthesis	Induction of plant defense, insect attractant	(47); (48)
Tetrahydro-β-citronellene	Terpenoid metabolism	Monoterpenoid biosynthesis	.	NA
p-Cymene	Terpenoid metabolism	Monoterpenoid biosynthesis	Insecticidal activity, insect repellent	(49); (50)
R-(+)-Limonene	Terpenoid metabolism	Monoterpenoid biosynthesis	Insect repellent	(51); (52)
Dihydrocitronellol	Monoterpene metabolism	Citronellol reduction	.	NA
α-Terpinene	Terpenoid metabolism	Monoterpenoid biosynthesis	Insecticidal activity	(53)
Undecane	Fatty acid metabolism	Alkane biosynthesis	Oviposition preference	(54)
Dodecane	Fatty acid metabolism	Alkane biosynthesis	Insect attractant, mating stimulant	(55); (56)
Pentadecane	Fatty acid metabolism	Alkane biosynthesis	Insect attractant	(57)
Tritriacontane	Fatty acid metabolism	Cuticular alkane biosynthesis	.	NA
Methyl anthranilate	Amino acid metabolism	Anthranilate biosynthesis (shikimate pathway)	Insect repellent	(58); (59)
Hexacosane	Fatty acid metabolism	Wax biosynthesis	.	NA
Dotriacontane	Fatty acid metabolism	Wax biosynthesis	.	NA
Octacosane	Fatty acid metabolism	Wax biosynthesis	.	NA
Hexedecane	Fatty acid metabolism	Alkane biosynthesis	.	NA

248 **REFERENCES**

249 35. M. P. Huynh, B. E. Hibbard, K.-V. Ho, K. S. Shelby, Toxicometabolomic profiling of
250 resistant and susceptible western corn rootworm larvae feeding on Bt maize seedlings.
251 *Scientific reports* **12**, 1-13 (2022).

36. L. N. Meihls *et al.*, Increased survival of western corn rootworm on transgenic corn within three generations of on-plant greenhouse selection. *Proc. Natl. Acad. Sci.* **105**, 19177-19182 (2008).

37. A. Johanns, Metric conversions. Iowa State University Extension and Outreach. https://www.extension.iastate.edu/AGDm/wholefarm/html/c6-80.html (accessed on May 22, 2025). (2024).

38. A. Seethepalli *et al.*, RhizoVision Explorer: open-source software for root image analysis and measurement standardization. *AoB Plants* **13**, plab056 (2021).

39. F. Ma *et al.*, A comparison between whole transcript and 3'RNA sequencing methods using Kapa and Lexogen library preparation methods. *BMC Genomics* **20**, 1-12 (2019).

40. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1-21 (2014).

41. T. F. Branson, E. E. Ortman, Feeding behavior of larvae of the western corn rootworm: normal larvae and larvae maxillectomized with laser radiation. *Ann. Entomol. Soc. Am.* **62**, 808-812 (1969).

42. S. Som, D. S. Willett, H. T. Alborn, Dynamics of belowground volatile diffusion and degradation. *Rhizosphere* **4**, 70-74 (2017).

43. H. Chen *et al.*, 1-Octen-3-ol, a self-stimulating oxylipin messenger, can prime and induce defense of marine alga. *BMC Plant Biol.* **19**, 1-16 (2019).

44. K. Cui *et al.*, Toxicological effects of the fungal volatile compound 1-octen-3-ol against the red flour beetle, *Tribolium castaneum* (Herbst). *Ecotoxicol. Environ. Saf.* **208**, 111597 (2021).

45. M. Ameye *et al.*, Priming of wheat with the green leaf volatile Z-3-hexenyl acetate enhances defense against *Fusarium graminearum* but boosts deoxynivalenol production. *Plant Physiol.* **167**, 1671-1684 (2015).

46. J. Wang *et al.*, A green leaf volatile,(Z)-3-hexenyl-acetate, mediates differential oviposition by *Spodoptera frugiperda* on maize and rice. *BMC Biol.* **21**, 140 (2023).

47. Y. Pan *et al.*, The herbivore-induced plant volatile tetradecane enhances plant resistance to *Holotrichia parallela* larvae in maize roots. *Pest Manag. Sci.* **78**, 550-560 (2022).

48. H. Yin, W. Li, M. Xu, D. Xu, P. Wan, The role of tetradecane in the identification of host plants by the mirid bugs *Apolygus lucorum* and *Adelphocoris suturalis* and potential application in pest management. *Front. Physiol.* **13**, 1061817 (2022).

49. Q.-H. Xie *et al.*, Bioactivities of thymol and p-cymene from the essential oil of *Adenosma buchneroides* against three stored-product insects. *Environ. Sci. Pollut. Res.* **30**, 110841-110850 (2023).

50. P. M. Bleeker *et al.*, The role of specific tomato volatiles in tomato-whitefly interaction. *Plant Physiol.* **151**, 925-935 (2009).

51. N. J. Conboy *et al.*, Volatile organic compounds as insect repellents and plant elicitors: an integrated pest management (IPM) strategy for glasshouse whitefly (*Trialeurodes vaporariorum*). *J. Chem. Ecol.* **46**, 1090-1104 (2020).

52. S. A. Belliard *et al.*, Identification of host plant volatile stimulants of *Anastrepha fraterculus* male courtship behavior. *Front. Ecol. Evol.* **10**, 943260 (2022).

53. S.-S. Cheng *et al.*, Variations in insecticidal activity and chemical compositions of leaf essential oils from *Cryptomeria japonica* at different ages. *Bioresour. Technol.* **100**, 465-470 (2009).

297   54.   C. Hajdu *et al.*, Volatile-mediated oviposition preference for healthy over root-infested
298   plants by the European corn borer. *Plant Cell Environ.* **47**, 2228-2239 (2024).

299   55.   J.-Y. Lv *et al.*, Active volatile components of the preferred hosts are potential attractants
300   to *Hyphantria cunea* adults. *Pestic. Biochem. Physiol.* **202**, 105910 (2024).

301   56.   H. Feng *et al.*, Plant volatile organic compounds attractive to *Lygus pratensis*. *Open Life*
302   *Sci.* **17**, 362-371 (2022).

303   57.   B. Bohman, A. M. Weinstein, R. Mozuraitis, G. R. Flematti, A.-K. Borg-Karlson,
304   Identification of (Z)-8-heptadecene and n-pentadecane as electrophysiologically active
305   compounds in *Ophrys insectifera* and its *Argogorytes* pollinator. *Int. J. Mol. Sci.* **21**, 620
306   (2020).

307   58.   E. J. Bernklau, B. E. Hibbard, A. P. Norton, L. B. Bjostad, Methyl anthranilate as a
308   repellent for western corn rootworm larvae (Coleoptera: Chrysomelidae). *J. Econ.*
309   *Entomol.* **109**, 1683-1690 (2016).

310   59.   A. Afify, B. Horlacher, J. Roller, C. G. Galizia, Different repellents for *Aedes aegypti*
311   against blood-feeding and oviposition. *PLoS One* **9**, e103765 (2014).