Long-term surveillance suggests multiple hybridization events by nuclear reassortment and
accelerated intercontinental spread of wheat yellow rust.

Mogens Støvring Hovmøller[1*#], Tine Thach[1*], Julian Rodriguez-Algaba[1], Jens Grønbech

Hansen[1], Marcel Meyer[2], David Hodson[3], Kumarse Nazari[4], Robert F. Park[5], Rita Tam[6],

Mareike Möller[1&6], Benjamin Schwessinger [6], John P. Rathjen [6], Paula Silva[7], Venancio Riella[7]

and Annemarie Fejer Justesen[1]

[1] Department of Agroecology, Global Rust Reference Center, Aarhus University, Slagelse,

Denmark; [2]Crop Science, Institute of Crop Science and Resource Conservation, University of

Bonn, Germany, [3]CIMMYT Nepal, Khumaltar, Lalipur, Nepal; [4] Turkey-ICARDA Regional Cereal

Rust Research Center, Izmir, Turkey; [5] Plant Breeding Institute, School of Life and Environmental

Science, The University of Sydney, Australia; [6] Research School of Biology, The Australian

National University, Canberra, Australia, [7] Instituto Nacional de Investigación Agropecuaria

(INIA), Colonia, Uruguay

*Shared 1st author

[#] corresponding author

Word count (plain text): 6186

Number of Figures: 4 (all in colour)

Number of Tables: 2

Supporting information

Number of Figures: 5

Number of Tables: 7

Summary

- Drivers of evolution of crop pathogens leading to novel diversity and long-distance

dispersal (LLD) may have strong implications on food security and sustainable plant

disease control at global scales. However, consolidated evidence is often lacking due to

absence of regular pathogen surveys beyond national capacities.

• Our study documents world-wide connectivity between populations of *Puccinia*

striiformis, causing yellow rust on cereals and grasses. Twelve cases of inter-

continental spread of wheat yellow rust, including seven cases with major impact on

disease epidemics in recipient areas, were detected from genotypic analyses of 3240

pathogen samples from 41 countries at six continents.

• In three cases, the most plausible mechanism for novel pathogen diversity was

somatic hybridization via nuclear reassortment between co-existing lineages on shared

host cultivars, spreading by LDD from Europe to South America and Australia,

respectively, a few years after first detection. Several high-impact incursions from

South Asia into East Africa were also observed, including a genotype with a dramatic

impact on wheat breeding programs of global relevance.

• The present study stresses an urgent need for coordinated crop pathogens monitoring

across borders. Only global efforts will enable prevention and control of pathogens

that represent major challenges for food security at regional and global scales.

Key words

Puccinia striiformis, hybridization, world-wide spread, trajectory simulations, global

surveillance, haplo-phased genome, SSR haplotype

Introduction

An increase in human travel and trade and the escalating speed of changing climate and

weather patterns have renewed attention on disease epidemiology in humans, animals and

crop plants, e.g., the US National Academies of Sciences (Olsen et al., 2011), the United

Nations Food and Agriculture Organisation (FAO, 2024). This is exemplified by human

vulnerability to pandemics as highlight by the coronavirus SARS-CoV-2 (Dillon et al., 2021;

Chavda et al., 2022). Long distance dispersal (LDD) is an important epidemiological feature of

many pathogens, including those affecting crops, because even rare LDD events can have a

major impact on crop production (Brown and Hovmøller, 2002; Carvajal-Yepes et al., 2019).

Recent reviews suggest that LDD events of cereal rust pathogens are increasing in frequency

at the intercontinental scale, possibly due to intensified crop production, climate change

resulting in milder winters in temperate regions, and increased travel and commerce (Park,

2015; Jin et al., 2020; Hovmøller et al., 2023). The negative impact of more frequent LDD

events may delay or prevent meeting the 70% increase in food and agricultural production by

2050 demanded by global population growth (van Dijk et al., 2021).

One major obstacle for the detection of LDD events in plant pathology is the lack of regular

and extensive pathogen population surveys beyond national capacities (Jeger and Pautasso,

2008; Ristaino et al., 2021), including common sampling protocols, analytic tools and data

sharing (Patpour et al., 2022). This makes it difficult to establish the exact geographical and

evolutionary origins of a new pathogen variant as it spreads into recipient areas. Population

genetic and genomic analyses are useful approaches to calculate distances and migration

rates within and among pathogen populations (Zhan et al., 2003; Ali et al., 2014a; Bebber et

al., 2014). At the same time, tracking of individual multi-locus genotypes (MLGs) represents a

powerful strategy for the detection of LDD events especially in clonal populations, which tend

to preserve haplotypes (Hovmøller et al., 2002).

The rust fungi (*Pucciniales*) are the most species rich and complex order of plant pathogens,

consisting of almost 8000 species. They are characterized by a true biotrophic lifestyle and

are individually adapted to their host plant at species and cultivar levels (Flor, 1971; Anikster,

1985; Aime et al., 2017). *Puccinia striiformis* f. sp. *tritici (Pst)*, causing yellow (or stripe) rust in

wheat, is dikaryotic, heteroecious and macrocyclic requiring two taxonomically unrelated

hosts to complete its life cycle (Jin et al., 2010). However, sexual reproduction on the

alternate host, *Berberis spp.,* is observed rarely in nature, even in western China (Wang et al.,

2016), despite multiple reports of signatures of sexual reproduction in populations in the

near-Himalayan region, which are considered the center of diversity for *Pst* (Duan et al.,

2010; Ali et al., 2014b; Thach et al., 2016; Huang et al., 2021). Somatic hybridization by fusion

of hyphae, leading to nuclear reassortment which may preserve the entire haplotype, has so

far only been reported under experimental conditions in *Pst* (Little and Manners, 1967; Little

and Manners, 1969; Lei et al., 2017), unlike the wheat leaf rust pathogen *P. triticina* (Park and

Wellings, 2012; Sperschneider et al., 2023), the wheat stem rust pathogen *P. graminis* f. sp.

tritici (Li et al., 2019), and *P. coronata* causing oat crown rust (Henningsen et al., 2024).

The use of molecular markers in recent time has contributed significantly to the

understanding of the global landscape of wheat rust fungi. In North America, yellow rust was

first discovered in 1915, although the founder population is not known (Line, 2002), however,

random samples from the 1970s, 1980s and 1990s suggested a common origin of the North

American and the European PstS0 population (Markell and Milus, 2008; Ali et al., 2014a;

Thach et al., 2016). The initial spread of *Pst* from Europe to Australia in 1978, also PstS0, is

well documented (Wellings and McIntosh, 1990; Hovmøller et al., 2008; Ding et al., 2021).

The worldwide spread of two warm temperature-adapted strains of *Pst* in 2000 and following

years (termed PstS1 and PstS2, respectively), collectively represent one of the largest and

most rapid spread of a crop pathogen up until that time (Hovmøller et al., 2008). Timewise,

these large-scale dispersal events of *Pst* coincided with the emergence of *Pgt* race "Ug99" in

East Africa that was virulent on more than 80% of wheat lines developed by The International

Maize and Wheat Improvement Center (CIMMYT) (McIntosh and Pretorius, 2011). In

response, the Borlaug Global Rust Initiative (BGRI) was formed and laid the foundation for

more extensive and regular wheat rust disease and pathogen surveys. These included Africa

and Asia, which were connected with national rust survey programs in Europe and elsewhere

by the establishment of the Global Rust Reference Center in 2008 (Hovmøller, 2011). This

allowed analyses of the spread and evolution of *Pst* and *Pgt* across borders and continents,

including the return of *Pgt* into Europe and connectivity to epidemics elsewhere (Patpour et

al., 2022), including Western Siberia 2015-2016 (Shamanin et al., 2016) and East Africa

(Olivera et al., 2015). In addition, independent surveys have reported large scale

epidemiological consequences of new incursions of *Pst* into South America in 2017 (Anibal

Carmona et al., 2019; Riella et al., 2024) and Australia in 2017-2018 (Ding et al., 2021).

The present study investigated drivers of evolution of the *P. striiformis* population at the

global scale based on genotypic analyses of 3180 samples collected at four continents (37

countries) between 2009 and 2023, complemented by genotype information of additional 60

samples representing long-term surveillance efforts in Europe, North America (Milus et al.,

2015a) and Australia (Ding et al., 2021). LDD events were investigated by the potential

sharing of genotypes in putative source and recipient areas, considering the time of first

detection as well as accumulation of genetic diversity in the considered populations. Likely

evolutionary origin of successful LDD migrants were explored in detail in specific cases,

including hypothetical hybridization events by nuclear reassortment, which has not been

demonstrated in *Pst* under non-experimental conditions. The plausibility of wind-dispersal

along hypothesized transmission routes was explored by conducting trajectory simulations

considering typical spore survival times and mean wind patterns. The prospects for

prevention and mitigation efforts to reduce the increasing impact of crop pathogen spread at

intercontinental scale are discussed.

Materials and Methods

Collection and recovery of yellow rust samples

Pathogen sampling reflected the coordinated efforts in wheat rust surveillance facilitated by

research programs within the Borlaug Global Rust Initiative, European Union research and

network initiatives as well as national programs in Europe and Australia and recent studies in

North America **(Fig. S1).**

A total of 3180 samples of yellow rust mainly from wheat (and occasionally other cereals and

grasses) were collected in 37 countries on four continents between 2009 and 2023 (Table S1).

The number of samples varied across years and locations, reflecting occurrence and sampling

efforts of the yellow rust disease. Two hundred thirteen samples from the Middle East and

West Asia were provided by The Regional Cereal Rust Research Center (RCRRC), ICARDA,

Turkey. An additional pre-existing 60 isolates served as references of previously identified *Pst*

groups, and *Pst* populations in USA (Milus et al. (2015b) and Australia (Ding et al. (2021).

Recovery of incoming samples and multiplication of urediniospores followed the procedure by

Thach et al. (2025), after which they were harvested, dried, and stored in liquid nitrogen (-

196°C) or freezer (-80°C) until further use. For analysis of diversity over time and space, samples

were grouped in two time periods (2009-2016) and (2017-2023) and seven geographical

regions, in addition to reference samples from North America, Australia and Europe.

DNA extraction and SSR genotyping

From 2009 to 2012, DNA was extracted from dried urediniospores (approx. 20 mg/sample) according to Thach et al. (2016). Since 2013, extraction was based on dried leaf segments with single rust lesions taken from multiplication plants or directly from incoming samples according to Thach et al. (2025). Genotyping was done using 19 simple sequence repeat (SSR) co-dominant markers and allele sizes were manually scored, allowing up to two missing markers per sample (Thach et al., 2025). Samples with allele sizes and/or combinations suggesting more than a single genotype were excluded from the dataset. Detection of novel or exotic genotypes in a region/country was confirmed by additional independent DNA extractions from original sample material and genotyping. Two hundred thirteen samples from the Middle East (2018-2020) collected by Turkey-ICARDA Regional Cereal Rust Research Center were SSR genotyped by EUROFINS, France. Scoring of allele sizes and data alignment to existing format were carried out at the Global Rust Reference Center (GRRC). Assignment of multilocus genotype identity (MLG #) was done for samples without missing data using poppr package version 2.9.3 implemented in the R environment (Kamvar et al., 2014; Kamvar et al., 2015); for samples with missing data at a maximum of two marker loci, new MLGs were considered in case scored allele sizes revealed additional variability, whereas missing observations were not taken into account for MLG identity. Duplicate MLGs derived from the same bulk sample were removed from the dataset.

Definition of genetic groups

Nineteen previously defined genetic groups within *Pst* (PstS0-PstS18), each consisting of closely related MLGs and associated races (Thach et al., 2025), were used to illustrate regional pathogen differences and changes between two time periods, 2009-2016 and 2017-2023, respectively. Historically, the definition and naming of a new group has been inspired by the potential epidemic impact across large areas Hovmøller et al. (2010), and a diversification from previous world-wide groups by a significant number of alleles (typically five SSR alleles or more) and/or group-specific diagnostic markers (Walter et al., 2016; Ali et al., 2017). An updated summary of named *Pst* groups is available at the GRRC website:

Phylogenetic analysis

The dataset for phylogenetic analysis was compiled by including one representative sample

from each MLG (clone corrected). An unrooted phylogenetic Neighbor Joining tree was

constructed to visualize global diversity within *Pst*, including the positions of major genetic

groups defined as described above. The calculations were based on calculated genetic

distances using poppr package version 2.9.3 implemented in the R environment, 1200

bootstraps and cutoff at 50 (Kamvar et al., 2014), where Bruvo's distance for stepwise

mutation for microsatellites was calculated according to Bruvo et al. (2004).

Population diversity and structure

To investigate the power of discrimination of applied markers, a genotype accumulation curve

was calculated in the poppr R package based on samples without missing data (n= 2588), and

random resampling 10,000 times without replacement (Kamvar et al., 2014) (Fig. S2).

F_{st} values representing divergence between the geographic populations in two time periods (I:

2009-2016 and II: 2017-2023) were calculated based on the full dataset (3180 samples) and a

clone-corrected dataset with a single MLG representative per population and time period (283

samples) according to (Peakall and Smouse, 2006; 2012).

Tests for somatic hybridization by nuclear reassortment

Nuclear-specific haplotypes of hypothetical parental and hybrid isolates were inferred based

on the position of SSR markers and allele sizes on a chromosomal map derived from

haplotype-phased genome data of a PstS0 isolate (Plant Breeding Institute accession number

415 collected in 1982) (Schwessinger et al., 2018; Tam et al., 2025). The position of SSR

markers were determined by blast analysis (-word_size 12 -evalue 1 -dust no -outfmt 6) of 20

primer sequence pairs against the PstS0 genome (Table S2) and visualized by a phenogram

according to Wolfgang et al. (2013), including the lengths of individual SSR alleles (Fig. S3).

The analyses took advantage of multiple hypothetical hybridization events involving shared

SSR haplotypes, as well as time and geo-referenced sampling supporting both temporal and

spatial dimensions of evolution within the considered geographical populations.

Whole genome confirmation of a somatic hybridization event

The hypothesized hybridization resulting in PstS10 inferred by SSR and PstS0 genome data

were validated by two newly generated reference assemblies of isolates representing PstS7

and PstS10, respectively. The fully phased nuclear-assigned reference genome for an

Australian sampled PstS10 isolate (Ding et al. 2021) was generated as described previously for

PstS0 (Tam et al., 2025) with the exception of using PacBio HiFi and Oxford Nanopore

Technology (ONT) R10.4 based long-reads and Hi-C as input for Verkko genome assembler

(Rautiainen et al., 2023). The haplotype-phased reference genome of PstS7 used ONT R10.4.1

simplex reads and Hi-C with hifiasm --ont (Cheng et al., 2025). To expand upon the MLG

results at the genome level, we performed comparative alignments between the nuclear-

phased haplotypes of a proposed somatic hybrid, PstS10, and those of its two hypothetical

parental isolates, PstS0 (Tam *et al.*, 2025) and PstS7. Pairwise whole-genome alignments

were conducted using NUCmer v3.1 as part of the MUMmer package (Marçais et al., 2018)

with default settings. Only alignment blocks with >75% identity and longer than 500 bp were

retained. SNP counts and average identity were computed from 1-to-1 alignments using

MUMmer's DNAdiff v1.3. Dot plots were generated using a script adapted from paf2dotplot.r

(https://github.com/moold/paf2dotplot). The chromosomal locations and allele sizes for

PstS7 and PstS10 were also visualized in Table S2 and Fig. S3, respectively, and calculated

amplicon lengths were compared with SSR genotyping results.

Virulence phenotyping for validation of recent common origin

Successfully recovered migrant isolates from putative source and recipient areas were

virulence phenotyped to provide an independent assay for relatedness. An updated protocol

was followed using a 0-9 disease scoring scale according to (McNeal et al., 1971), where

infection types (IT) from 0-6 were generally considered incompatible (avirulent) and IT 7-9

compatible (virulent) (Thach et al., 2025). The interpretation of virulence/avirulence for each

isolate was generally based on two (or more) independent wheat lines carrying the

corresponding *P. striiformis Yr* resistance gene.

Plausibility of wind-dispersal

Plausibility of wind-dispersal was analysed by reviewing data from previous studies on related

fungal pathogens (Brown and Hovmøller, 2002; Isard et al., 2005; Meyer et al., 2017a; Meyer

et al., 2017b; Prank et al., 2019; Radici et al., 2022; Brodsky et al., 2023; Miedaner and

Garbelotto, 2024), analyzing long-term mean wind patterns, i.e., wind direction and speed for

times of overlapping wheat seasons at source and recipient areas based on the International

Research Institute for Climate and Society

(https://iridl.ldeo.columbia.edu/maproom/Global/Climatologies/Vector_Winds.html),

estimating typical spore survival times (Maddison and Manners, 1972; Rapilly, 1979; Aylor,

1986), and conducting a set of trajectory simulations using HYSPLIT (Stein et al., 2015) with two

different global-scale gridded meteorological datasets [NOAA's GDAS, 3-hourly temporal

resolution and 1 degree horizontal spatial resolution; ECMWF's ERA5, hourly temporal

resolution and 0.25 degree spatiol resolution (Hersbach et al., 2020) (see Fig. S4 for details).

Time-backwards trajectory simulations based on GDAS were conducted from locations in all

recipient areas to investigate origins of air masses that could have transported spores to

recipient areas where they were first detected. Time-forward simulations based on ERA5 were

conducted to analyze the plausibility of stepwise airborne transmission along two selected

pathways (Europe to South America via Northern Africa; and Central Asia to East Africa via the

Middle East).

Results

Genotypic *Pst* diversity at global scales

A total of 3483 samples collected from 41 countries on six continents were genotyped using

previously published procedures and SSR markers (Thach et al., 2025), which were all

polymorphic except for NRJN12. Of these samples, 136 were duplicates of identical

genotypes/races that had been sub-cultured from original bulk samples, and an additional 153

contained two or more genotypes per sample. The 289 duplicate and mixed samples were

excluded, leaving a total of 3180 for further analysis. An additional 60 reference samples

representing comprehensive long-term national rust surveys in Australia, North America and

Europe were included to illustrate genetic diversity at global scale (Fig. 1). A majority of the

samples (91.7%) grouped into previously defined genetic groups (PstS0-PstS18). The remaining

samples consisted of 74 highly diverse MLGs of distinct geographical origins, including areas in

South Asia, where high diversity has been reported previously. Samples from non-wheat hosts

in northern Europe, e.g., barley, wild grasses, and triticale, formed a distinct and diverse branch

in the phylogenetic tree, suggesting a different evolutionary origin. The 20 reference samples

from North America contained six MLGs grouping within PstS0, PstS1_2, PstS18, and two

unnamed groups. The 31 reference samples from Australia were contained in seven MLGs that

grouped within PstS0, PstS1_2, PstS10 and PstS13, and the six reference samples from Europe

and one from East Africa (1960-2006) represented PstS1_2, PstS3 and PstS4.

Fig. 1. Unrooted Neighbor Joining tree (clone-corrected) using Bruvo's distance showing the genetic relationship of 244 multi-locus genotypes (MLGs) representing world-wide sampling of *P. striiformis* in 41 countries on six continents (2009-2023), including reference isolates representing populations in Australia (1979-2021) and North America (1972-2023). Position of named genetic groups (PstS0-PstS18) and two temporary named groups (AF1, PK1) are indicated by different colours, typically consisting of clusters of multiple, closely related MLGs. Additional 74 unnamed, diverse MLGs were shown in greenish blue. Intercontinental migrants indicated by red arrows (within group position indicative).

The within-group diversity largely reflected time since first detection of the group, evolutionary origin, and likely prevalence of the disease and sampling intensity (Table 1; Fig. S2). The highest number of MLGs were observed in PstS1_2, first detected in East Africa in the 1970s, but

relatively high numbers were also observed for groups emerging from populations with high

genetic diversity in Asia, e.g., PstS7, PstS8, and PstS11, despite low diversity at time of incursion

into Europe (PstS7 and PstS8 in 2011) and East Africa (Psts11 in 2016) (data not shown).

Table 1. Diversity within world-wide genetic groups and geographic *Pst* populations defined
according to Table S1. Shaded values represent population, where each genetic group was
first detected.

Genetic group	Overall diversity				Number of MLGs per population[1] (n =population size)						
	Number of samples (n)	Number of MLG/ genetic	Shannon-Weiner index (H)	Frequency of prevalent MLG/ group	NEU (n=1086)	MEA (n=348)	EAF (n=914)	NWA (n=159)	SAS (n=283)	CWA (n=217)	SAM (n=173)
PstS0	57	7	1.95	54,4	5						4
PstS1_2	420	24	3.18	47,9		6	18	4	1	3	
PstS3	1	1	-	-						1	
PstS4	50	3	1.10	72,0	3	1					
PstS5	14	4	1.39	35,7						5	
PstS6	46	4	1.39	91,3			2		2	2	
PstS7	105	19	2.94	41,0	11	3		2		3	1
PstS8	128	15	2.71	51,6	15						
PstS9	139	14	2.64	53,2			1		1	14	
PstS10	600	16	2.77	82,2	15			1			1
PstS11	454	22	3.09	48,5		1	21		2	1	
PstS12	6	1	0.00	100,0	1						
PstS13	279	6	1.79	84,2	5			1			1
PstS14	239	6	1.79	72,0	5[2]	1	2[2]				1
PstS15	13	5	1.61	30,8	5						
PstS16	278	13	2.56	70,1			9		7		
PstS17	87	3	1.10	77,0	2	2	2				
Others	264	74	4.30		10	9	13		58	15	1
Total - named groups	2916	163			62	14	53	8	13	29	8
Grand total	3180	237	5.47		72	23	66	8	71	44	9

[1] Populations: NEU: Northern Europe, MEA: Middle East, EAF: East Africa, NWA: Northwest Africa, SAS: South Asia, CWA: Central-West Asia, SAM: South America.
[2] PstS14 first detected in NEU in 2015 (one single sample), where no surveys were conducted in NWA. PstS14 was highly dominant NWA from 2016 and onwards.

The distributions of isolates within the named genetic groups (PstS0-PstS17) were highly
divergent in the two time periods considered (2009-2016 and 2017-2023, respectively), and
across the considered geographical areas (Northern Europe, Middle East, East Africa,
Northwest Africa, South Asia, Central-West Asia, South America), which suggested dynamic
pathogen changes during time and significant geographic subdivisions since 2009 (Fig. 2).
Note that reference samples from North America and Australia, which represent historic key
isolates from these continents rather than systematic sampling in the two periods, were not
included in these analyses.

Fig. 2. Frequency of *P. striiformis* genetic groups in geographically distinct populations in two periods, a) 2009-2016 (n= 921) and b) 2017-2023 (n= 2259).

The distinct geographical distribution of genetic groups was striking. The aggressive and warm temperature-adapted group (PstS1/PstS2) was prevalent across Africa and the Middle East in the first period (2009-2016), but absent from South America, Europe and South Asia. PstS5 and PstS9 were only detected in Central Asia. PstS0, the prevalent genetic group in Europe up to 2010, was only detected in Europe, South- and North America and Australia. PstS8 was prevalent in northern Europe until 2016 and was not detected in other areas. PstS10 has been the most prevalent group in Europe since 2014 but has not been detected in East Africa and South- and Central Asia. These overall differences were reflected by corresponding F_{st} values based on an analysis where MLGs were grouped according to geographic population and sampling period (Table S2). Additional analyses based on clone-corrected data confirmed these trends but pointed towards additional connectivity between Central-West Asian populations and African and South Asian populations, respectively.

The limited differentiation between Middle East and East Africa/ NW Africa, and between South Asia and East Africa suggests sufficient migration to maintain low differentiation between these populations. The replacement of the prevalent PstS0 population in South America by a population dominated by PstS13 since 2017 represents the most dramatic temporal population shift between the two periods (Fst= 0.341).

The dataset also revealed several cases of emergence of novel, distinct genetic groups. For example, PstS10 containing a single MLG was first detected in Europe in 2012 and became widespread accounting for 82% of PstS10 samples (Table 1). This was followed by the emergence of 14 additional, closely related MLGs in the European population. The emergence of PstS13 and PstS14 in 2015-2016 in the European population represents two additional, distinct genetic groups that were not readily explained by mutation or migration events.

Identification of hypothetical somatic hybrids

Our genotypic data for PstS10, PstS13 and PstS14 were consistent with the hypothesis of somatic hybridization by nuclear reassortment of isolates of pre-existing lineages within the European population. First, allele sizes detected within samples representing MLGs of the first detected cases of PstS10, PstS13 and PstS14 were compared with samples representing all named genetic groups detected within Europe. This revealed that most MLGs and genetic groups were excluded as potential parent(s) due to mismatching allele sizes (Table S3). The next step involved inference of haplotypes of hypothetical parental pairs and hybrid MLGs by alignment with the PstS0 haplotype-phased reference genome, and assignment of the position of SSR markers on a genetic map, which revealed the genome derived SSR haplotype of PstS0 (Table S2, Fig. S3) as well as the haplotype based on observed SSR allele sizes (Table S6). Nucleus haplotype A of prevalent European races in PstS0 provided a 100% match with a nucleus haplotype of PstS10, where the second haplotype (B) provided a 100% match with a nucleus haplotype represented by the most prevalent MLG in PstS7. These results were confirmed by independent genomic data of isolates representing the parental groups of PstS0 and PstS7, and the hybridization resulting in PstS10 (Fig. S5). Similarly, we obtained 100% nucleus haplotype matches for PstS13 and PstS14, revealing that haplotype B in PstS7 was involved in the three hybridization events, however, arising from two different MLGs within PstS7. A schematic representation of the sequence and network of hybridization events is presented in Fig. 3; supplementary dotplots of the pairwise genome alignments and total SNP counts and average identity among the nuclear phased haplotypes of the PstS10 hybrid and the likely parental isolates shown in Fig. S5.

Fig. 3. Schematic representation of three independent hybridization events supported by observed SSR multilocus genotypes, pairwise genome alignments, and derived nuclear haplotypes. PstS10 was first detected in northern Europe (2012) in an area where the respective MLGs of PstS0 (#180) and PstS7 (#321) were prevalent on shared hosts of bread wheat. PstS13 was first detected in 2015, where a single MLG (#21) was detected at multiple locations in Europe, often on Triticale and rarely on winter wheat. The relevant MLG of PstS4 (#38) was detected almost exclusively on Triticale, while the MLG of PstS7 (#330) was detected widespread on triticale as well as bread wheat. PstS14 (MLG#138) was first detected in Europe in 2015/ NW Africa 2016 (not surveyed in 2015), the involved MLGs of PstS7 (MLG#330) and PstS1_2 (MLG#294) being prevalent in southern Europe/NW Africa at that time. First and last sampling years for individual genetic groups in the dataset are indicated (years before 2009 represented by historic reference samples).

Long distance dispersal events

The hypothesis of migration events at intercontinental scale was investigated by analyses of first detection of specific genetic groups and distinct MLGs within these in the seven geographical populations (Fig. 4). In three cases, first detection was associated with novel evolutionary origin by hybridization of pre-existing MLGs of divergent genetic groups within the considered geographical population, i.e., PstS10 (PstS0 and PstS7), PstS13 (PstS4 and

PstS7), and PstS14 (PstS1_2 and PstS7). These were first detected in Europe and NW Africa

between 2012 and 2016 (Fig. 4).

We documented 12 cases of exotic incursions of *Pst* at intercontinental scales (I-XII). Migration

direction and events were hypothesised based on year of first detection / hybridization event

in source and recipient areas, as well as prevalence and diversity within considered genetic

group (Table 1, Table 2, Figure 4).

Figure 4. Documented inter-continental incursions of *P. striiformis* 2009-2023, based on

sequential resampling of multilocus SSR genotypes/pathogen races on different continents.

Dotted circles represent putative source areas, where considered *Pst* group and MLGs

defined by 18 polymorphic SSR markers were first detected, arrows represent putative

transmission routes, colored circles represent first year of detection on the recipient

continent and the corresponding ID of individual MLGs. Details of individual dispersal events

and individual migrant MLGs are presented in Table 2. Dotted arrows: PstS13-MLG #21

present at two continents at time of incursion into Australia. The positions of circles and

arrows on the map are indicative.

Table 2. Inter-continental *Puccinia striiformis* dispersal events (I-XII) within a 15-year period (2009-2023) documented by source and recipient areas of unique multi-locus genotypes (MLGs), genetic groups and race phenotype.

Dispersal event (cf. Fig. 4)	Migrant (MLG #)	SSR genetic group	First detected (population and year)	Reference	Recipient population and year	Race confirmation in source and recipient population)[3]
I	21	PstS13	Europe (2015)	Ali et al. 2017	South America (2017)[2]	**Confirmed** : -,2,-,-,-,6,7,8,9,-,-,-,-,-,-,-,AvS,-
II	134	PstS14	Europe (2015)	Hovmøller et al. 2017[1]	South America (2017)[2]	-
III	21	PstS13	Europe (2015)	Ali et al. 2017	Australia (2018)[2]	-
IV	244	PstS10	Europe (2012)	Ali et al. 2017	Australia (2017)[2]	-
V	8	PstS6	Central Asia (2009)	Ali et al. 2017	East Africa (2010)[2]	**Confirmed:** 1,2,-,-,-,6,7,-,9,-,-,17,-,-,27,-,-,AvS,-
VI	78	PstS11	Central Asia (2012)	Hovmøller et al. 2018[1]	East Africa (2016)[2]	**Confirmed:** -,2,-,(4),-,6,7,8,-,-,-,17,-,-,27,32,-,AvS,-
VII	205	PstS16	South Asia (2017)	Hovmøller et al. 2023[1]	East Africa (2020)[2]	**Confirmed:** 1,2,3,(4),-,6,7,8,9,-,-,17,-,25,27,32,-,AvS,-
VIII	170	PstS17	Middle East (2018)	Hovmøller et al. 2019[1]	Europe (2020)	**Confirmed:** -,2,-,-,-,6,7,8,-,-,-,17,-,-,-,32,(Sp),AvS,Amb
IX	170	PstS17	Middle East (2018)	this paper	East Africa (2022)	**Confirmed:** -,2,-,-,-,6,7,8,-,-,-,17,-,-,-,32,(Sp),AvS,Amb
X	138	PstS14	Northwest Africa (2016)/ Europe (2016)	this paper	Middle East (2019)	-
XI	103	AF1	Central Asia (2012)	this paper	East Africa (2017)	**Confirmed:** 1,2,-,-,[-],6,7,8,9,-,-,17,[-],-,27,-,-,AvS,-
XII	185	PK1	South Asia (2017)	this paper	East Africa (2022)	-

[1] Annual reports from Global Rust Reference Center (accessible: www.wheatrust.org)

[2] High impact event in recipient areas

[3] Figures and symbols designate virulence and avirulence (-) corresponding to Pst resistance genes: *Yr1, Yr2, Yr3, Yr4, Yr5, Yr6, Yr7, Yr8, Yr9, Yr10, Yr15, Yr17, Yr24, Yr25, Yr27, Yr32* , and the resistance specificity of Spalding Prolific (Sp), Avocet S (AvS) and Ambition (Amb).

In all cases, we were able to detect likely geographical source areas, including two specific MLGs spreading from Europe into South America in 2017 (I and II, Table 2), i.e., PstS13 (MLG #21) and PstS14 (MLG #134), MLG #21 spreading into Australia (III, Table 2) and detected the following year (2018). PstS10 (MLG #244), the most common genotype in Europe since 2016, was also detected as an exotic incursion in Australia in 2017 (IV, Table 2). MLGs within three additional and highly divergent genetic groups, PstS6 (MLG #8), PstS11 (MLG #78) and PstS16 (MLG #205), detected in Central/South Asia in 2009, 2012 and 2017, were observed for the first time in East Africa (Ethiopia) in 2010, 2015 and 2020, respectively (V, VI, VII, Table 2). PstS17 (MLG #170), first detected in the Middle East in 2018 and since then prevalent in this region, was first detected at low frequency in northern Europe (Latvia, 2020, VIII, Table 2) and in East Africa (Ethiopia, 2022, IX, Table 2). PstS14 (MLG138), first detected in Europe and NW Africa in 2015/2016, was detected in the Middle East in 2019 (X, Table 2), later becoming dominant in local areas in West Asia, e.g., Iraq, where a single MLG in PstS14 made up 100% of samples from multiple locations and host cultivars (2021, 2022). Two MLGs, which have not yet been assigned a named genetic group, detected in Afghanistan, 2012 (temporary designation AF1, MLG #103) and Pakistan, 2017 (temporary designation PK1, MLG #185), respectively, were observed in East Africa in 2017 (Kenya) and 2022 (Ethiopia), respectively

(XI, XII, Table 2). These genetic groups, including PstS17, have remained at low frequency to

date in recipient areas in Northern Europe and East Africa.

We grouped the dispersal events according to similarities in source and recipient areas,

resulting in seven potential intercontinental atmospheric transport routes (see Table S6).

Results from HYSPLIT trajectory simulations, analyses of mean wind patterns in combination

with wheat growth timing and spore survival times, as well as previous evidence from related

fungal pathogens lead us to hypothesize that (i) wind dispersal was not plausible for the two

dispersal events, i.e., "Europe to Australia" (route 3) and "South America to Australia" (route

2), where accidental human-mediated transmission seems more likely; (ii) wind dispersal was

highly unlikely for direct movement from Europe to South America (route 1) and from

Central/South Asia directly to East Africa (route 4), although the data does not exclude the

possibility of indirect, stepwise airborne transmission (via northern Africa and the Middle

East/Arabian Peninsula, respectively); (iii) wind dispersal was plausible for the three dispersal

routes, Middle East to Europe (route 5), Middle East to East Africa (route 6) and Europe/North

Africa to the Middle East (route 7).

Discussion

The potential impact of rare events of long-distance dispersal (LDD) of plant pathogens on

plant health has been known for decades, e.g., Brown and Hovmøller (2002), Smart and Fry

(2001), Ristaino et al. (2021), Jones (2021). Despite this, in the vast majority of cases the exact

source and recipient areas are not known. The focus has often been on high impact examples,

e.g., overview for *Pst* on wheat by Jin et al. (2020), rather than cases with insignificant impact,

which may be overlooked and thereby contribute to an underestimation of the true

frequency of LDD in crop pathogens.

The present study was based on unique and systematic wheat rust surveillance efforts over

15 years facilitated by the formation of the Borlaug Global Rust Initiative (BGRI) and the

Global Rust Reference Center in 2008. This network approach has taken advantage of the

engagement of multiple collaborators across the world in coordinated wheat rust sampling

efforts (Table S7), the use of standardized pathogen phenotyping and genotyping

methodologies (Thach et al., 2025), as well as application of commonly agreed data

management and analytical tools (Hovmøller, 2011). Altogether, the sampling covered major wheat growing regions in 37 countries across seven broad geographical areas in Africa, Asia, Europe, and South America, and was further connected to North America and Australia through reference samples from major long-term national rust surveillance initiatives. This study may, therefore, represents the largest and most comprehensive spatiotemporal survey of a major crop pathogen to date (Savary et al., 2019).

We confirmed at least 12 intercontinental LDD dispersal events of Pst on wheat, each supported by distinct multi-locus genotype data, and in many cases also phenotypic virulence data, suggesting a recent common evolutionary origin for samples in both source and recipient areas. In seven cases, the dispersal events resulted in severe and widespread disease epidemics in recipient areas (Anibal Carmona et al. (2019) Riella et al. (2024); Ding et al. (2021), whereas five cases had only minor epidemiological impact (i.e., rare detection of the corresponding genetic groups in following years). In three cases, PstS10, PstS13, and PstS14, the first detected and most prevalent MLGs within each of these groups were hypothesized to originate asexually by somatic hybridization and nuclear reassortment. This provided a unique opportunity to identify the evolutionary mechanism and source of migrants, in addition to identifying the geographical area of first detection. Our conclusions concerning the hypotheses of LLD dispersal in the latter three cases, versus an alternative hypothesis of hybridization of "local" strains in recipient areas, were in all cases supported by lack of detection of matching parental isolates of the hypothesized hybrids in recipient areas (Australia and South America), despite extensive and widespread surveillance efforts (www.wheatrust.org; (Anibal Carmona et al., 2019; Ding et al., 2021; Riella et al., 2024).

Additional LDD events were also identified, although the exact source areas were not confirmed. For instance, the emergence of MLG #321 (PstS7) in Argentina in 2018 following first detection in Europe (2011) and Northwest Africa (2013-2014), MLG #247 (PstS10) in Uruguay in 2021, and MLG #175 (PstS17) in Northern Europe and East Africa. However, these observations most likely followed the dispersal paths already established by related MLGs with documented source origin, e.g., PstS7-MLG #321 and PstS10-MLG #247 (route 1), and PstS17-MLG#175 (route 5), but final conclusions would require additional sampling efforts from potential source areas that were not represented in the dataset.

Our findings represent previously undocumented aspects of yellow rust epidemiology. Unlike

other cereal rust fungi, somatic hybridization in *Pst*, involving nuclear reassortment, and

hypothesized parasexual exchange of genetic material has only been reported under

experimental conditions. Already in the 1960s, mixtures ofurediniospore isolates of different

races suggested that somatic hybridization by fusion of hyphae and subsequent nuclear

reassortment was the likely source of new race variants. More recently, cross infection

studies on wheat and barley using paired experimental isolates supported the occurrence of

somatic hybridization, however, these results could only be explained by assuming

chromosome reassortment and crossovers between parental isolates (Lei et al., 2017).

Earlier, Park and Wellings (2012) concluded that somatic hybridization was likely in many rust

species and *formae speciales*, the process being observed in nature within and between rust

species on *Linum*, *Populus*, *Senecio*, several grass species, and leaf rust on wheat. However,

the exact mechanisms were generally not well understood.

Our data suggest that somatic hybridization may be a more significant source of

agriculturally-relevant genetic diversity in yellow rust than sexual recombination. This may

help explain why the pathogen is rarely detected on the alternate host *Berberis vulgaris* (Jin

et al., 2010; Wang et al., 2016), even in the near Himalayan region or elsewhere, despite high

genetic diversity (Duan et al., 2010; Ali et al., 2014a; Thach et al., 2016; Huang et al., 2021).

In the present study, putative hybridization events in *Pst* were identified by positioning SSR

markers and allele sizes on a chromosomal map based on a haplotype-phased genome of a

PstS0 isolate (Tam et al., 2025). This was combined with careful analysis of MLGs generated

using consistently applied methodologies (Thach et al., 2025), which revealed three major

hybridization events giving rise to PstS10, PstS13 and PstS14. Importantly, the hybrid origin of

PstS10 was confirmed by comparing haplotype-phased nuclear assigned genomes of the

potential parents. The prevalence of matching MLG haplotypes in PstS0 and PstS7 suggested

a hybridization event resulting in PstS10 somewhere on bread wheat in Europe. In contrast,

the inferred parental origin of PstS13 was a relatively rare MLG in PstS7 (#330, <10%),

sampled from both bread wheat and Triticale, and a frequent MLG in PstS4 (#38, > 95%),

almost exclusively associated with Triticale. This suggests that hybridization most likely took

place on Triticale. The resulting MLG (# 21) of PstS13 was frequently detected on Triticale,

durum wheat, and spring wheat, but rarely on winter wheat. For PstS14, one of the putative

parents, PstS1_2 MLG #294, was detected only in NW Africa, coinciding with the presence of

a matching second parent (PstS7 MLG #330 in the same region in 2014. Interestingly, PstS14

made up 100% of samples in surveys in three consecutive years in NW Africa since 2016, i.e.,

2016 (six samples), 2017 (38 samples) and 2018 (54 samples), whereas this group was

observed only sporadically elsewhere. This spatial and temporal pattern strongly support the

emergence of PstS14 through somatic hybridization in Northwest Africa. In addition to these

three major hybridization events, our data revealed two additional cases of 100% matches

between hypothesized parental pairs and hybrids, which remained at low prevalence in the

European *Pst* population, and one case of hybridization in the North American population

(data not shown).

In four high-impact and three low-impact cases, the conclusion of recent dispersal events

was supported by identical virulence phenotypes in source and recipient areas based on

assays comprising wheat experimental lines representing more than 25 different host

resistance specificities (Thach et al., 2025). Because mutation from avirulence to virulence in

plant pathogenic fungi, including *Pst,* may be significantly higher than average mutation rates

(Hovmøller and Justesen, 2007), the existence of an exact match in virulence phenotype is an

additional indication of direct connection between such populations in an evolutionary time

perspective.

Based on the 12 confirmed dispersal events, we hypothesized seven putative transmission

routes, where the plausibility of wind dispersal was investigated through HYSPLIT trajectory

simulations based on high-resolution global meteorological data, testing different spore

survival times in the atmosphere, and also through analyses of wheat growing timing, monthly

mean wind patterns during those periods where the empirical data suggested that LDD could

have taken place. This integrative analysis along with recent evidence from related fungal

pathogens, indicates that both human mediated and windborne dispersal playing important

roles in intercontinental dispersal of wheat yellow rust.

Airborne dispersal appears especially plausible between the Middle East and East Africa as well

as Europe. This aligns with observed, genetic similarities between *Pst* populations in the Middle

East and East Africa (Table S3) that also indicates frequent gene flow between these regions

and help explain several of the reported incursion events (Fig. 4, Table 2). The recent detections

of Pst6, Pst11, Pst16, Pst17, AF1 and PK1 in East Africa after previous detections in the Middle

East and Central/South Asia, provide supportive evidence for the previously proposed Rift

Valley Incursion pathway for rust fungi (Meyer et al., 2017a). While direct dispersal from

Central/South Asia to East Africa seems highly unlikely, the possibility of stepwise airborne

dispersal via the Middle East cannot be ruled out. However, the latter scenario is less supported

since PstS6 and PstS16 were not detected in the Middle East, and PstS11 appeared in the region

only three years after first detection in East Africa, although relatively low sampling intensity

in critical areas on the Arabian Peninsula and the Horn of Africa should be taken into account,

as well as the stochastic nature of these LDD events.

Direct airborne dispersal from Europe to South America or from Central/South Asia to East

Africa were highly unlikely, and a more comprehensive modelling study is required to test the

hypothesis of stepwise airborne transmission. Similarly, for the two dispersal events from

Europe to Australia and South America to Australia, wind dispersal was not plausible, indicating

human mediated transport and emphasizing the importance of quantifying risks due to

unintended human-mediated transmission via trade and travel. The trajectory simulations

presented here were intended at developing initial hypotheses about which dispersal

mechanisms might be plausible for each incursion event. Testing as part of future work would

require large-scale modelling efforts incorporating detailed spore dispersion simulations that

include environmental and epidemiological conditions at source and target regions, along with

sensitivity analysis covering spore survival times, releasing times and locations. To advance

such efforts, two key components are essential. First, the collection of more detailed

experimental data on spore survival times during atmospheric transport obtained from large

samples of viable spores, to estimate more reliably the fraction of spores, out of extremely

large numbers released during outbreaks, and a continued extension of global surveillance

networks are essential for further improving our understanding and capacity to predict and

mitigate risks from long-range exotic incursion events.

Altogether, our study has documented a high degree of connectivity world-wide between

populations of one the most epidemic crop pathogens, yellow rust infecting cereals and

grasses, geographically as well as genetically. Incursions from the Himalayas and elsewhere

into Europe and the Mediterranean basin 10-20 years ago (Hubbard et al., 2015; Hovmøller et

al., 2016; Walter et al., 2016) hybridized with isolates of pre-existing genetic groups, e.g.,

PstS0, PstS1_2, PstS4, the new hybrids replaced to a large extent existing population(s),

spreading onwards to new continents (Hovmøller et al., 2023), where the immigrant strains

rapidly became dominant (Anibal Carmona et al., 2019; Ding et al., 2021; Riella et al., 2024).

Such a connectivity between populations of airborne pathogens is probably particularly evident

in global cropping systems like wheat, which is grown on six continents. This widespread

cultivation underscores the need for coordinated pathogen surveillance and early-warning

systems at both regional and global levels. Access to reference laboratories and testing facilities

is crucial as these enable timely assessment of the vulnerability of breeding material and

deployed cultivars to new and aggressive pathogen strains before they become widespread.

We show here that somatic (asexual) hybridization both takes place and seems to be common

in natural populations of *Pst* despite previously only reported under experimental conditions.

The rapid raise of hybrids such as PstS10, PstS13 and PstS14, which to a large extent replaced

pre-existing pathogen populations in Europe, Australia and South America, suggests that new

epidemiological insights into the actual mechanisms of hybridization at both cellular and

genetic levels are required. While not all hybridization events necessarily lead to increased

fitness or persistence, the widespread success of these particular hybrids suggests that these

may be potentially superior in fitness to their parents taking the entire epidemiological disease

cycle into account, which may involve changed interactions with various components and

mechanisms of host resistance.

Our data suggest that the risks and pace of pathogen evolution driven by mutation and somatic

hybridization may correlate positively with the size and areas affected by ongoing epidemics. In

turn, this may influence the probabilities of further hybridization and risks of unintentional

pathogen spread at local, regional and global levels. Thus, there is an imminent and urgent need

to sustain research and development within the area of plant breeding and disease

epidemiology including both experimental as well as modelling approaches to better anticipate

and prevent future epidemics. In parallel, greater awareness is needed around the risks of

unintentional crop pathogen spread via human activity to enhance food security in

agroecological settings with rapidly changing climate and weather patterns and increasing

political instability.

Acknowledgements

This research was supported by multiple funding sources, including a series of grants by the Bill

and Melinda Gates Foundation, the UK Foreign, Commonwealth & Development Office (INV-

048345,　DEWAS; INV-003439, AGG; ID 0PP1133199, DGGW; OPPGD1389: DRRW), the

Strategic Danish Research Council (grant no. 11-116241, RUSTFIGHT), the Danish Ministry of

Food, Agriculture and Fisheries (grant no. 34120902567); a series of grants from

Jordbruksverket (Sweden), agreement no. 21-11857/09, 25-6922/10, 25-2363/12, 4.4.11-

1615/14, 4.4.11-01001/18, 4.4.11-17402/2021, Research Council of Norway (Hveterust, No.

301835), as well as long-term support by Aarhus University. Mareike Möller was supported by

EU Marie Skłodowska-Curie fellowship (R-evolution, grant no. 10119509). The work of Marcel

Meyer was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) – grant no. 529743941. Work at ANU was supported by DP230100941 from the

Australian Research Council. Rita Tam was supported by a Grains Research and Development

Corporation Graduate Research Scholarship. Rust sampling undertaken by Turkey-ICARDA

Regional Cereal Rust Research Center was additionally supported by TAGEM (General

Directorate of Agricultural Research and Policies of Turkey), CRP Wheat (CIMMYT), and FAO-

Turkey. The authors are grateful to Ellen Jørgensen, Janne Holm Hansen, Jakob Sørensen, and

Steen Meier, Aarhus University, for technical assistance during yellow rust genotyping and

virulence phenotyping of 1000s of rust isolates at the Global Rust Reference Center during 15

years, Sambasivam Periyannan, Andrew Milgate, Ashley Jones, Zhenyan Luo, Zhara Chew, Eric

Pereira and Danish Baig for sample characterization and assistance in the genomic research

concerning isolates of PstS0, PstS7 and PstS10 (ANU), and Ezgi Kurtulus, Handan Kavas and Ali

Kadiroğlu for assisting in collection and preparing samples provided by the Turkey-ICARDA

Regional Cereal Rust Research Center. The deposition and access to a reference collection of

Pst isolates from USA up to 2015 by Eugine Milus is highly appreciated, as well as the

contributions of ~ 420 people assisting in disease surveillance, sampling and submission of rust

samples throughout the period 2009-2023, often under difficult and challenging conditions,

without whom this work would not have been possible (for details, see Table S7).

Competing interests

637　None declared

Author contributions

DH and MSH conceived and designed the surveillance program; TT carried out the experimental SSR genotyping work under the supervision of AFJ, including data alignment and quality check and population genetic analyses; VR assisted in genotyping of samples from South America; JRA performed virulence phenotyping of hypothetical migrant isolates and assisted in SSR genotyping ; JGH developed the web-based data management and display system for world-wide prevalence of *Pst* genetic groups; KN supplied raw SSR data for Middle East samples 2018-2020; RP was curator of reference samples representing the Australian *Pst* population 1979-2021, undertook race analysis, and was responsible for associated funding; RT and MM performed the sequencing and genomic data analyses of PstS10 hybrid and parental isolates; BS and JR contributed to genomic data analysis, supervised and secured funding for the genomic work and provided extensive feedback on the manuscript; PS RT and MM performed the sequencing and genomic data analyses of PstS10 hybrid and parental isolates; BS and JR supervised and secured funding for the genomic work; PS supervised the PhD work of VR in South America; MME conducted trajectory simulations of the plausibility of wind dispersal of spores based on meteorological datasets; MSH was curator of samples hosted by the GRRC, supervised project activities anchored at GRRC, wrote the first draft of the manuscript; all authors revised and accepted the final manuscript.

ORCID

Mogens Støvring Hovmøller https://orcid.org/0000-0002-4432-8898

Tine Thach https://orcid.org/0000-0001-8853-694X

Annemarie Fejer Justesen https://orcid.org/0000-0001-7969-8760

Julian Rodriguez-Algaba https://orcid.org/0000-0002-4960-7562

Jens Grønbech Hansen https://orcid.org/0000-0001-7682-5613

John P. Rathjen https://orcid.org/0000-0003-4073-8088

Benjamin Schwessinger https://orcid.org/0000-0002-7194-2922

Mareike Möller https://orcid.org/0000-0002-2146-5507

Rita Tam https://orcid.org/0009-0001-6032-1872

Robert F. Park https://orcid.org/0000-0002-9145-5371

Marcel Meyer https://orcid.org/0000-0002-8457-9608

Kumarse Nazari https://orcid.org/0000-0001-9348-892X

David Hodson https://orcid.org/0000-0002-8641-8568

Venancio Riella https://orcid.org/0000-0001-5017-7857

Paula Silva https://orcid.org/0000-0003-2655-2949

Data availability

676   All raw sequencing data generated in this study have been submitted to NCBI BioProject

677   database under accession number PRJNA1256629. The SSR genotypic data of the 3180 isolates

678   included in the present study are available at https://doi.org/ (pending)

References

681   Aime, M.C., McTaggart, A.R., Mondo, S.J., and Duplessis, S. (2017). "Chapter Seven - Phylogenetics and
682   Phylogenomics of Rust Fungi," in *Advances in Genetics,* eds. J.P. Townsend & Z. Wang.
683   Academic Press), 267-307.
684   Ali, S., Gladieux, P., Leconte, M., Gautier, A., Justesen, A.F., Hovmøller, M.S., et al. (2014a). Origin,
685   migration routes and worldwide population genetic structure of the wheat yellow rust
686   pathogen *Puccinia striiformis* f.sp. *tritici*. *PLOS Pathogens* 10**,** e1003903. doi:
687   doi.org/10.1371/journal.ppat.1003903.
688   Ali, S., Gladieux, P., Rahman, H., Saqib, M.S., Fiaz, M., Ahmed, H., et al. (2014b). Inferring the
689   contribution of sexual reproduction, migration and off-season survival to the temporal
690   maintenance of microbial populations: a case study on the wheat fungal pathogen *Puccinia*
691   *striiformis* f.sp. *tritici*. *Molecular Ecology* 23**,** 603–617.
692   Ali, S., Rodriguez-Algaba, J., Thach, T., Sorensen, C.K., Hansen, J.G., Lassen, P., et al. (2017). Yellow
693   Rust Epidemics Worldwide Were Caused by Pathogen Races from Divergent Genetic Lineages.
694   *Frontiers in Plant Science* 8. doi: 10.3389/fpls.2017.01057.
695   Anibal Carmona, M., Jose Sautua, F., Perez-Hernandez, O., Grosso, C., Vettorello, L., Milanesio, B., et
696   al. (2019). Rapid emergency response to yellow rust epidemics caused by newly introduced
697   lineages of Puccinia striiformis f. sp. tritici in Argentina. *Tropical Plant Pathology* 44(4)**,** 385-
698   391. doi: 10.1007/s40858-019-00295-y.
699   Anikster, Y. (1985). "The Formae Speciales," in *The Cereal Rusts,* eds. A.P. Roelfs & W.R. Bushnell.
700   Academic Press, Inc.), 115-130.
701   Aylor, D.E. (1986). A framework for examining inter-regional aerial transport of fungal spores.
702   *Agricultural and Forest Meteorology* 38(4)**,** 263-288. doi: https://doi.org/10.1016/0168-
703   1923(86)90017-1.

Bebber, D.P., Holmes, T., and Gurr, S.J. (2014). The global spread of crop pests and pathogens. *Global*
Ecology and Biogeography 23(12)**,** 1398-1407. doi: https://doi.org/10.1111/geb.12214.
Brodsky, H.K., Calderón, R., Hamilton, D.S., Li, L., Miles, A., Pavlick, R., et al. (2023). Assessing long-
distance atmospheric transport of soilborne plant pathogens. *Environmental Research Letters*
18(10)**,** 104021. doi: 10.1088/1748-9326/acf50c.
Brown, J.K.M., and Hovmøller, M.S. (2002). Aerial Dispersal of Pathogens on the Global and
Continental Scales and Its Impact on Plant Disease. *Science* 297(5581)**,** 537-541. doi:
10.1126/science.1072678.
Bruvo, R., Michels, N.K., D'Souza, T.G., and Schulenburg, H. (2004). A simple method for the
calculation of microsatellite genotype distances irrespective of ploidy level. *Molecular Ecology*
13(7)**,** 2101-2106. doi: https://doi.org/10.1111/j.1365-294X.2004.02209.x.
Carvajal-Yepes, M., Cardwell, K., Nelson, A., Garrett, K.A., Giovani, B., Saunders, D.G.O., et al. (2019). A
global surveillance system for crop diseases. *Science* 364(6447)**,** 1237-1239. doi:
doi:10.1126/science.aaw1572.
Chavda, V.P., Patel, A.B., and Vaghasiya, D.D. (2022). SARS-CoV-2 variants and vulnerability at the
global level. *Journal of Medical Virology* 94(7)**,** 2986-3005. doi:
https://doi.org/10.1002/jmv.27717.
Cheng, H., Qu, H., McKenzie, S., Lawrence, K.R., Windsor, R., Vella, M., et al. (2025). Efficient near
telomere-to-telomere assembly of Nanopore Simplex reads. *bioRxiv***,** 2025.2004.2014.648685.
doi: 10.1101/2025.04.14.648685.
Dillon, C.F., Dillon, M.B., and Drake, H.L. (2021). Multiscale Airborne Infectious Disease Transmission.
Applied and Environmental Microbiology 87(4)**,** e02314-02320. doi: doi:10.1128/AEM.02314-
20.
Ding, Y., Cuddy, W.S., Wellings, C.R., Zhang, P., Thach, T., Hovmøller, M.S., et al. (2021). Incursions of
divergent genotypes, evolution of virulence and host jumps shape a continental clonal
population of the stripe rust pathogen Puccinia striiformis. *Molecular Ecology* 30(24)**,** 6566-
6584. doi: https://doi.org/10.1111/mec.16182.
Duan, X., Tellier, A., Wan, A., Leconte, M., de Vallavieille-Pope, C., and Enjalbert, J. (2010). *Puccinia*
striiformis f.sp. *tritici* presents high diversity and recombination in the over-summering zone
of Gansu-China. *Mycologia* 102**,** 44-53. doi: 10.385/08-098.
FAO (2024). "Scaling up One Hedalth for better transboundary pest and disease control".
(https://www.fao.org/animal-health/news-events/news/detail/scaling-up-one-health-for-
better-transboundary-pest-and-disease-control/en: Food and Agriculture Organization of the
United Nations).
Flor, H.H. (1971). The current status of the gene-for-gene concept. *Annual Review of Phytopathology*
9**,** 275-296.
Henningsen, E.C., Lewis, D., Nazareno, E.S., Mangelson, H., Sanchez, M., Langford, K., et al. (2024). A
high-resolution haplotype collection uncovers somatic hybridization, recombination and
intercontinental movement in oat crown rust. *PLOS Genetics* 20(11)**,** e1011493. doi:
10.1371/journal.pgen.1011493.
Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., et al. (2020). The
ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society* 146(730)**,** 1999-
2049. doi: https://doi.org/10.1002/qj.3803.
Hovmøller, M.S. (2011). "The increased risk of global wheat rust pandemics: putting yellow rust into
perspective," in *Fungal Diseases: An Emerging Threat to Human, Animal, and Plant Health,* ed.
E.R.C. Eds L. Olsen, D.A. Relman and L. Pray. (Washington, DC, USA: The National Academic
Press), 252–263.
Hovmøller, M.S., and Justesen, A.F. (2007). Rates of evolution of avirulence phenotypes and DNA
markers in a northwest European population of Puccinia striiformis f. sp tritici. *Molecular*
Ecology 16(21)**,** 4637-4647. doi: 10.1111/j.1365-294X.2007.03513.x.

Hovmøller, M.S., Justesen, A.F., and Brown, J.K.M. (2002). Clonality and long-distance migration of Puccinia striiformis f.sp tritici in north-west Europe. *Plant Pathology* 51(1)**,** 24-32.

Hovmøller, M.S., Thach, T., and Justesen, A.F. (2023). Global dispersal and diversity of rust fungi in the context of plant health. *Current Opinion in Microbiology* 71**,** 102243. doi: https://doi.org/10.1016/j.mib.2022.102243.

Hovmøller, M.S., Walter, S., Bayles, R.A., Hubbard, A., Flath, K., Sommerfeldt, N., et al. (2016). Replacement of the European wheat yellow rust population by new races from the centre of diversity in the near-Himalayan region. *Plant Pathology* 65(3)**,** 402-411. doi: 10.1111/ppa.12433.

Hovmøller, M.S., Walter, S., and Justesen, A.F. (2010). Escalating Threat of Wheat Rusts. *Science* 329(5990)**,** 369-369. doi: 10.1126/science.1194925.

Hovmøller, M.S., Yahyaoui, A.H., Milus, E.A., and Justesen, A.F. (2008). Rapid global spread of two aggressive strains of a wheat rust fungus. *Molecular Ecology* 17(17)**,** 3818-3826.

Huang, M., Liu, T., Cao, S., Yuen, J., Zhan, J., Jia, Q., et al. (2021). Analyses of Wheat Yellow Rust Populations Reveal Sexual Recombination and Seasonal Migration Pattern of Puccinia striiformis f. sp. tritici in Gangu, Northwestern China. *Phytopathology®* 111(12)**,** 2268-2277. doi: 10.1094/phyto-12-20-0558-r.

Hubbard, A., Lewis, C., Yoshida, K., Ramirez-Gonzalez, R., de Vallavieille-Pope, C., Thomas, J., et al. (2015). Field pathogenomics reveals the emergence of a diverse wheat yellow rust population. *Genome Biology* 16(1)**,** 23.

Isard, S.A., Gage, S.H., Comtois, P., and Russo, J.M. (2005). Principles of the Atmospheric Pathway for Invasive Species Applied to Soybean Rust. *BioScience* 55(10)**,** 851-861. doi: 10.1641/0006-3568(2005)055[0851:Potapf]2.0.Co;2.

Jeger, M.J., and Pautasso, M. (2008). Plant disease and global change – the importance of long-term data sets. *New Phytologist* 177(1)**,** 8-11. doi: https://doi.org/10.1111/j.1469-8137.2007.02312.x.

Jin, Y., Szabo, L.J., and Carson, M. (2010). Century-old mystery of Puccinia striiformis life history solved with the identification of Berberis as an Alternate Host. *Phytopathology* 100**,** 432-435.

Jin, Y.K., J; Szabo, L.; Rouse, M. N.; Hovmoeller, M. S.; Olivera, P. D.; Singh, R. P.; Bhavani, S.;, , and (2020). "CHAPTER 4: Emergence and Spread of New Races of Wheat Rust Fungi: Continued Threat to Food Security and Prospects of Genetic Control," in *Emerging Plant Diseases and Global Food Security*.), 53-79.

Jones, R.A.C. (2021). Global Plant Virus Disease Pandemics and Epidemics. *Plants (Basel)* 10(2). doi: 10.3390/plants10020233.

Kamvar, Z.N., Brooks, J.C., and Grünwald, N.J. (2015). Novel R tools for analysis of genome-wide population genetic data with emphasis on clonality. *Front Genet* 6**,** 208. doi: 10.3389/fgene.2015.00208.

Kamvar, Z.N., Tabima, J.F., and Grünwald, N.J. (2014). Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* 2**,** e281. doi: 10.7717/peerj.281.

Lei, Y., Wang, M., Wan, A., Xia, C., See, D.R., Zhang, M., et al. (2017). Virulence and Molecular Characterization of Experimental Isolates of the Stripe Rust Pathogen (Puccinia striiformis) Indicate Somatic Recombination. *Phytopathology®* 107(3)**,** 329-344. doi: 10.1094/phyto-07-16-0261-r.

Li, F., Upadhyaya, N.M., Sperschneider, J., Matny, O., Nguyen-Phuc, H., Mago, R., et al. (2019). Emergence of the Ug99 lineage of the wheat stem rust pathogen through somatic hybridisation. *Nature Communications* 10(1)**,** 5068. doi: 10.1038/s41467-019-12927-7.

Line, R.F. (2002). Stripe rust of wheat and barley in North America: A retrospective historical review. *Annual Review of Phytopathology* 40**,** 75-118. doi: 10.1146/annurev.phyto.40.020102.111645.

Little, R., and Manners, J. (1969). Somatic recombination in yellow rust of wheat (Puccinia striiformis)::
II. Germ tube fusions, nuclear number and nuclear size. *Transactions of the British Mycological*
Society 53(2)**,** 259-267.
Little, R., and Manners, J.G. (1967). Production of new physiologic races in Puccinia striiformis (yellow
rust) by heterokaryosis. *Nature* 213(5074)**,** 422-&.
Maddison, A.C., and Manners, J.G. (1972). Sunlight and viability of cereal rust uredospores.
Transactions of the British Mycological Society 59(3)**,** 429-443. doi:
https://doi.org/10.1016/S0007-1536(72)80124-4.
Marçais, G., Delcher, A.L., Phillippy, A.M., Coston, R., Salzberg, S.L., and Zimin, A. (2018). MUMmer4: A
fast and versatile genome alignment system. *PLOS Computational Biology* 14(1)**,** e1005944.
doi: 10.1371/journal.pcbi.1005944.
Markell, S.G., and Milus, E.A. (2008). Emergence of a novel population of Puccinia striiformis f. sp
tritici in eastern United States. *Phytopathology* 98(6)**,** 632-639. doi: 10.1094/phyto-98-6-0632.
McIntosh, R.A., and Pretorius, Z.A. (2011). Borlaug Global Rust Initiative provides momentum for
wheat rust research. *Euphytica* 179**,** 1–2. doi: 10.1007/s10681-011-0389-y.
McNeal, F.H., Konzak, C.F., Smith, E.P., Tate, W.S., and Russel, T.S. (1971). A uniform system for
recording and processing cereal research data. *US Agricultural Research Service* 42**,** 31-121.
Meyer, M., Burgin, L., Hort, M.C., Hodson, D.P., and Gilligan, C.A. (2017a). Large-Scale Atmospheric
Dispersal Simulations Identify Likely Airborne Incursion Routes of Wheat Stem Rust Into
Ethiopia. *Phytopathology®* 107(10)**,** 1175-1186. doi: 10.1094/phyto-01-17-0035-fi.
Meyer, M., Cox, J.A., Hitchings, M.D.T., Burgin, L., Hort, M.C., Hodson, D.P., et al. (2017b). Quantifying
airborne dispersal routes of pathogens over continents to safeguard global wheat supply.
Nature Plants 3(10)**,** 780-786. doi: 10.1038/s41477-017-0017-5.
Miedaner, T., and Garbelotto, M.M. (2024). Human-mediated migration of plants, their pathogens and
parasites. *Journal of Plant Pathology* 106(2)**,** 301-325. doi: 10.1007/s42161-024-01589-0.
Milus, E.A., Lee, K.D., and Brown-Guedira, G. (2015a). Characterization of Stripe Rust Resistance in
Wheat Lines with Resistance Gene Yr17 and Implications for Evaluating Resistance and
Virulence. *Phytopathology* 105(8)**,** 1123-1130. doi: 10.1094/phyto-11-14-0304-r.
Milus, E.A., Moon, D.E., Lee, K.D., and Mason, R.E. (2015b). Race-Specific Adult-Plant Resistance in
Winter Wheat to Stripe Rust and Characterization of Pathogen Virulence Patterns.
Phytopathology 105(8)**,** 1114-1122. doi: 10.1094/phyto-11-14-0305-r.
Olivera, P., Newcomb, M., Szabo, L.J., Rouse, M., Johnson, J., Gale, S., et al. (2015). Phenotypic and
Genotypic Characterization of Race TKTTF of Puccinia graminis f. sp tritici that Caused a Wheat
Stem Rust Epidemic in Southern Ethiopia in 2013-14. *Phytopathology* 105(7)**,** 917-928. doi:
10.1094/phyto-11-14-0302-fi.
Olsen, L., Choffnes, E.R., Relman, D.A., and Pray, L. (2011). *Fungal Diseases: An Emerging Threat to*
Human, Animal, and Plant Health. Workshop Summary. Washington, DC: The National
Academies Press.
Park, R.F. (2015). Long-term surveys of pathogen populations underpin sustained control of the rust
diseases of wheat in Australia. . *Journal of the Royal Society of New South Wales* 148**,** 15-27.
Park, R.F., and Wellings, C.R. (2012). "Somatic Hybridization in the Uredinales," in *Annual Review of*
Phytopathology, Vol 50, eds. N.K. VanAlfen, J.E. Leach & S. Lindow.), 219-239.
Patpour, M., Hovmøller, M.S., Rodriguez-Algaba, J., Randazzo, B., Villegas, D., Shamanin, V.P., et al.
(2022). Wheat Stem Rust Back in Europe: Diversity, Prevalence and Impact on Host
Resistance. *Frontiers in Plant Science* 13. doi: 10.3389/fpls.2022.882440.
Peakall, R., and Smouse, P.E. (2006). genalex 6: genetic analysis in Excel. Population genetic software
for teaching and research. *Molecular Ecology Notes* 6(1)**,** 288-295. doi:
https://doi.org/10.1111/j.1471-8286.2005.01155.x.
Peakall, R., and Smouse, P.E. (2012). GenAlEx 6.5: genetic analysis in Excel. Population genetic
software for teaching and research—an update. *Bioinformatics* 28(19)**,** 2537-2539. doi:
10.1093/bioinformatics/bts460.

Prank, M., Kenaley, S.C., Bergstrom, G.C., Acevedo, M., and Mahowald, N.M. (2019). Climate change
impacts the spread potential of wheat stem rust, a significant crop disease. *Environmental*
Research Letters 14(12)**,** 124053. doi: 10.1088/1748-9326/ab57de.
Radici, A., Martinetti, D., and Bevacqua, D. (2022). Early-detection surveillance for stem rust of wheat:
insights from a global epidemic network based on airborne connectivity and host phenology.
Environmental Research Letters 17. doi: 10.1088/1748-9326/ac73aa.
Rapilly, F. (1979). Yellow rust epidemiology. *Annual Review of Phytopathology* 17**,** 59-73.
Rautiainen, M., Nurk, S., Walenz, B.P., Logsdon, G.A., Porubsky, D., Rhie, A., et al. (2023). Telomere-to-
telomere assembly of diploid chromosomes with Verkko. *Nature Biotechnology* 41(10)**,** 1474-
1482. doi: 10.1038/s41587-023-01662-6.
Riella, V., Rodriguez-Algaba, J., García, R., Pereira, F., Silva, P., Hovmøller, M.S., et al. (2024). New
Races with Wider Virulence Indicate Rapid Evolution of Puccinia striiformis f. sp. tritici in the
Southern Cone of America. *Plant Disease* 108(8)**,** 2454-2461. doi: 10.1094/pdis-02-24-0320-re.
Ristaino, J.B., Anderson, P.K., Bebber, D.P., Brauman, K.A., Cunniffe, N.J., Fedoroff, N.V., et al. (2021).
The persistent threat of emerging plant disease pandemics to global food security.
Proceedings of the National Academy of Sciences 118(23)**,** e2022239118. doi:
doi:10.1073/pnas.2022239118.
Savary, S., Willocquet, L., Pethybridge, S.J., Esker, P., McRoberts, N., and Nelson, A. (2019). The global
burden of pathogens and pests on major food crops. *Nature Ecology & Evolution* 3(3)**,** 430-
439. doi: 10.1038/s41559-018-0793-y.
Schwessinger, B., Sperschneider, J., Cuddy, W.S., Garnica, D.P., Miller, M.E., Taylor, J.M., et al. (2018).
A Near-Complete Haplotype-Phased Genome of the Dikaryotic Wheat Stripe Rust Fungus
Puccinia striiformis f. sp. *tritici* Reveals High Interhaplotype Diversity.
mBio 9(1)**,** e02275-02217. doi: 10.1128/mBio.02275-17.
Shamanin, V., Salina, E., Wanyera, R., Zelenskiy, Y., Olivera, P., and Morgounov, A. (2016). Genetic
diversity of spring wheat from Kazakhstan and Russia for resistance to stem rust Ug99.
Euphytica 212(2)**,** 287-296. doi: 10.1007/s10681-016-1769-0.
Smart, C., and Fry, W. (2001). Invasions by the late blight pathogen: renewed sex and enhanced
fitness. *Biological Invasions* 3**,** 235-243.
Sperschneider, J., Hewitt, T., Lewis, D.C., Periyannan, S., Milgate, A.W., Hickey, L.T., et al. (2023).
Nuclear exchange generates population diversity in the wheat leaf rust pathogen Puccinia
triticina. *Nature Microbiology* 8(11)**,** 2130-2141. doi: 10.1038/s41564-023-01494-9.
Stein, A.F., Draxler, R.R., Rolph, G.D., Stunder, B.J.B., Cohen, M.D., and Ngan, F. (2015). NOAA's
HYSPLIT Atmospheric Transport and Dispersion Modeling System. *Bulletin of the American*
Meteorological Society 96(12)**,** 2059-2077. doi: https://doi.org/10.1175/BAMS-D-14-00110.1.
Tam, R., Möller, M., Luo, R., Luo, Z., Jones, A., Periyannan, S., et al. (2025). Long-read genomics reveal
extensive nuclear-specific evolution and allele-specific expression in a dikaryotic fungus.
Genome Research. doi: 10.1101/gr.280359.124.
Thach, T., Ali, S., de Vallavieille-Pope, C., Justesen, A.F., and Hovmøller, M.S. (2016). Worldwide
population structure of the wheat rust fungus Puccinia striiformis in the past. *Fungal Genetics*
and Biology 87**,** 1-8. doi: https://doi.org/10.1016/j.fgb.2015.12.014.
Thach, T., Justesen, A.F., Rodriguez-Algaba, J., Sørensen, C.K., and Hovmøller, M.S. (2025). "Integrated
Approach for Genotyping and Race Typing of Puccinia striiformis in Wheat," in *Wheat Rusts*
and Resistance Breeding: Methods and Protocols, eds. G.S. Brar & S. Holden. (New York, NY:
Springer US), 75-91.
van Dijk, M., Morley, T., Rau, M.L., and Saghai, Y. (2021). A meta-analysis of projected global food
demand and population at risk of hunger for the period 2010–2050. *Nature Food* 2(7)**,** 494-
501. doi: 10.1038/s43016-021-00322-9.
Walter, S., Ali, S., Kemen, E., Nazari, K., Bahri, B., Enjalbert, J., et al. (2016). Molecular markers for
tracking the origin and worldwide distribution of invasive strains of *Puccinia striiformis*.
Ecology and Evolution doi: 10.1002/ece3.2069.

Wang, Z., Zhao, J., Chen, X., Peng, Y., Ji, J., Zhao, S., et al. (2016). Virulence Variations of Puccinia
906   striiformis f. sp. tritici Isolates Collected from Berberis spp. in China. *Plant Disease* 100(1)**,** 131-
907   138. doi: 10.1094/pdis-12-14-1296-re.
908   Wellings, C.R., and McIntosh, R.A. (1990). Puccinia striiformis f.sp. tritici in Australasia: pathogenic
909   changes during the first 10 years. *Plant Pathology* 39(2)**,** 316-325. doi:
910   https://doi.org/10.1111/j.1365-3059.1990.tb02509.x.
911   Zhan, J., Pettway, R.E., and McDonald, B.A. (2003). The global genetic structure of the wheat pathogen
912   Mycosphaerella graminicola is characterized by high nuclear diversity, low mitochondrial
913   diversity, regular recombination, and gene flow. *Fungal Genetics and Biology* 38(3)**,** 286-297.
914   doi: https://doi.org/10.1016/S1087-1845(02)00538-8.