Mitochondrial protein FgDML1 impacts DON toxin biosynthesis and cyazofamid sensitivity in *Fusarium graminearum* by affecting mitochondrial homeostasis

Chenguang Wang[a †], Xuewei Mao[b †], Weiwei Cong[a], Lin Yang[a], Yiping Hou[a] *

[a] State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Nanjing Agricultural University, Nanjing, Jiangsu, China

[b] College of Plant Protection, Henan Agricultural University, Zhengzhou, Henan, China

[†] These authors contributed equally to this work

*Corresponding author：Yiping Hou; E-mail address: houyiping@njau.edu.cn

Abstract

Fusarium graminearum is a global pathogen responsible for Fusarium head blight (FHB) in wheat, causing substantial yield losses and producing the mycotoxin deoxynivalenol (DON), which poses a threat to both human and animal health. *Drosophila melanogaster* Misato-Like protein (DML1) plays a critical role in regulating mitochondrial function, yet its function in filamentous fungi remains unexplored. In this study, we characterized FgDML1 in *F. graminearum*. FgDML1 interacts with the mitochondrial fission and fusion protein FgDnm1 to maintain mitochondrial stability, thereby positively regulating acetyl-CoA levels and ATP synthesis, which influences toxisome formation and ultimately affects DON toxin biosynthesis. Additionally, FgDML1 is involved in the regulation of toxin biosynthetic enzyme expression. In the ΔFgDML1 mutant, Complex III enzyme activity decreased, overexpression of complex III assembly factors FgQCR2, FgQCR8, and FgQCR9 may induce conformational changes in the Qi-site protein, specifically altering the sensitivity of *F. graminearum* to respiratory inhibitor cyazofamid not Qo-site inhibitor pyraclostrobin and other fungicides. Furthermore, the loss of FgDML1 leads to defects in nutrient utilization, as well as in asexual and sexual reproduction, and pathogenicity. In conclusion, this study identifies a novel regulatory role for FgDML1 in DON toxin biosynthesis and cyazofamid sensitivity in *F. graminearum*. Our study provides a theoretical framework for understanding DON biosynthesis regulation in *F. graminearum* and identifies potential molecular targets for FHB control.

23 **Keywords:** *Fusarium graminearum*, DON production, FgDML1, Cyazofamid

24 Pathogenicity

Introduction

Fusarium head blight (FHB), caused by *Fusarium graminearum*, is a devastating

fungal disease that affects cereal crops, leading to significant yield losses in

wheat-growing regions worldwide(Dean et al., 2012). This disease is particularly

concerning due to the production of various mycotoxins, including trichothecenes and

zearalenone, during fungal infection(Hai et al., 2023). Among these, deoxynivalenol

(DON), a type of trichothecene, is the most commonly detected mycotoxin(Merhej et

al., 2011). The DON toxin poses a serious threat to both human and animal health,

significantly compromising immune and reproductive systems(Knutsen et al., 2017).

Chronic exposure to DON can lead to anorexia, neurological disorders, and immune

dysregulation, while acute high-dose exposure may result in vomiting, diarrhea,

neurological disturbances, and even miscarriage or stillbirth(Sumarah, 2022; Zhang et

al., 2024). Studies have shown that increased DON levels are positively correlated with

the pathogenicity rate of *F. graminearum*(Bönnighausen et al., 2019; Ilgen et al., 2009;

Mudge et al., 2006). As FHB continues to spread and its severity increases, DON

contamination is emerging as a critical global concern(Duan et al., 2020; Liu et al.,

2016). This highlights the need for effective strategies to manage and mitigate the

impact of both the disease and the associated mycotoxin contamination.

The absence of resistant cultivars necessitates reliance on chemical methods as the

primary approach for controlling FHB(Liu et al., 2019). However, the inappropriate

application of fungicides has led to the emergence of resistance, particularly against

carbendazim, tebuconazole, and pyraclostrobin(Andrade et al., 2022; Yi et al., 2023;

Zhou et al., 2024). Notably, carbendazim has been demonstrated to stimulate the

biosynthesis of DON toxin, especially in fungicide-resistant populations of *F.*

graminearum(Li et al., 2019; Zhou et al., 2020). Similarly, pyraclostrobin significantly

promotes DON biosynthesis, with no selectivity between resistant and sensitive strains.

Although tebuconazole has not been directly implicated in inducing DON production, it

is classified as a fungicide with a moderate risk of resistance development(Duan et al.,

2020). Field observations have revealed diverse resistance mechanisms to

tebuconazole, complicating management efforts(Liu et al., 2011; Zhao et al., 2021).

The potential evolution of resistance to tebuconazole is particularly concerning, as it

could undermine its efficacy and exacerbate the challenge of controlling DON

contamination. There is an urgent need for a fungicide that has not yet been used in *F.*

graminearum to address the DON toxin contamination caused by fungicide resistance.

Cyazofamid is a mitochondrial respiratory chain complex III inhibitor with

demonstrated efficacy in controlling fungal diseases. It specifically binds to the Qi site

of complex III, blocking the reduction of ubiquinone and inhibiting ATP synthesis,

thereby exerting its fungicidal effect(Li et al., 2014; Mitani et al., 2001). However, it

remains unclear whether other proteins within the intricate regulatory network of the

respiratory chain play a role in modulating fungal sensitivity to cyazofamid. Further

investigations are needed to elucidate the potential involvement of these regulatory

components.

The synthesis of DON toxin is primarily regulated by genes involved in

trichothecene biosynthesis, known as *FgTRIs*. Many of these *FgTRI* genes have been

well characterized(Merhej et al., 2011). The initial step in DON synthesis is catalyzed

by the trichothecene synthase *FgTri5*, which facilitates the cyclization of farnesyl

pyrophosphate (FPP) to form trichodiene (TDN)(Kimura et al., 2007). However,

before this catalytic process can occur, the organism must first obtain sufficient ATP

from the mitochondria. ATP is critical not only for the basic cellular functions but also

for driving several key reactions in the isoprenoid biosynthesis pathway. Specifically,

ATP is required for the enzymatic conversion of precursors, leading to the production

of FPP, a vital intermediate in the trichothecene biosynthesis pathway. The availability

of ATP therefore directly impacts the efficient synthesis of FPP, which, in turn,

regulates the downstream production of DON toxin. Without an adequate supply of

ATP, these biosynthetic pathways cannot proceed efficiently, thus hindering the

synthesis of both the precursor molecules and the final toxin product(Alexander et al.,

2009). Pathogen-intrinsic ATP homeostasis is recognized as a critical, rate-limiting

determinant for toxin biosynthesis. Previous studies indicate that dual-target inhibition

of ATP synthase (AtpA) and adenine deaminase (Ade) by a specific small-molecule

probe effectively depletes intracellular ATP, consequently suppressing the synthesis of

key virulence factors TcdA and TcdB transcriptionally and translationally(Marreddy et

al., 2024). The systemic toxicity of Anthrax Edema Toxin (ET) is primarily attributed

to its catalytic activity, which depletes the host cell's ATP reservoir, thereby triggering

a bioenergetic collapse that culminates in cell lysis and death(Liu et al., 2025). The

biosynthesis of DON entails a reorganization of the endoplasmic reticulum into a

specialized compartment termed the "toxisome"(Tang et al., 2018). The assembly of

the toxisome coincides with the aggregation of key biosynthetic enzymes, which in

turn enhances the efficiency of DON production. Concurrently, this

compartmentalization serves as a self-defense mechanism, protecting the fungus from

the autotoxicity of TRI pathway intermediates (Boenisch et al., 2017). The proteins

TRI1, TRI4, TRI14, and Hmr1 are confirmed constituents of this structure(Kistler and

Broz, 2015; Menke et al., 2013).

Drosophila melanogaster Misato-Like protein (DML1) encodes a Misato-like

protein in *Saccharomyces cerevisiae*, an essential factor involved in mitochondrial

DNA (mtDNA) inheritance. In addition to its role in mtDNA maintenance, DML is

implicated in mitochondrial compartmentalization and chromosome segregation. The

protein shares structural features with members of the GTPase family, suggesting

potential involvement in GTP-dependent processes(Gurvitz et al., 2002). In *Homo*

sapiens, the DML protein is localized to the mitochondria, where silencing of DML

results in a transition from a networked mitochondrial morphology to a fragmented

form. This indicates that DML plays a critical role in mitochondrial distribution,

morphological fusion, and overall mitochondrial dynamics(Kimura and Okano, 2007).

In prokaryotes, the homologous protein FtsZ, which shares functional similarities with

DML, has been identified as a target for antimicrobial agents(Huang et al., 2007;

Vollmer, 2006). However, despite these insights from studies in the aforementioned

organisms, the exact role of DML in filamentous fungi remains undetermined.

This study systematically characterized the functional roles of DML protein in

F. graminearum(FgDML1), including mycelial growth, asexual development, and

sexual reproduction. More importantly, we found that FgDML1 reduces DON toxin

synthesis by decreasing the enzyme activity of complex III, reducing ATP synthesis,

and lowering acetyl-CoA accumulation. Furthermore, the deletion of FgDML1 results

in compensatory upregulation of complex III assembly factors, which may alter the

conformation of the Qi site protein and subsequently modulates sensitivity to

cyazofamid.

Results

Identification, deletion, and complementation of FgDML1.

The DML1 protein from S. cerevisiae was used as a query for a BLAST search

against the Fusarium genome database, resulting in the identification of the putative

DML1 gene FgDML1 (FGSG_05390) in *F. graminearum*. The FgDML1 gene is 1455

bp in length, contains three introns, and encodes a 484-amino acid protein with 23.88%

homology to its counterpart in S. cerevisiae. Phylogenetic analysis was performed

using DML1 sequences from other species obtained via NCBI, and a phylogenetic

tree was constructed (Fig. 1A). Structural analysis revealed the presence of the

Misat_Tub_SegII and Tubulin_3 domains. Furthermore, subcellular localization

studies confirmed that FgDML1 localizes to mitochondria, as demonstrated by

colocalization with a mitochondria-specific dye MitoTracker Red CMXRos (Fig. 1B).

To investigate the function of FgDML1, ΔFgDML1 deletion mutant was

constructed and validated via PCR and Southern blot analysis (Fig. S2), confirming

the mutant as a single-copy knockout. To ensure that the observed phenotypes were

specifically attributable to the deletion of FgDML1, an in situ complemented mutant,

ΔFgDML1-C, was generated to restore FgDML1.

136 FgDML1 regulates vegetative growth.

To assess the role of FgDML1 in vegetative growth, PH-1, ΔFgDML1, and

ΔFgDML1-C were cultured on various media, including PDA, CM, MM, and V8, to

measure growth rates. The ΔFgDML1 strain displayed a distinct growth phenotype

characterized by retardation in radial growth and the formation of more compact,

denser hyphal networks on all tested media compared to the PH-1 and ΔFgDML-C

strains. (Fig. 2A). Moreover, ΔFgDML1 displayed pronounced hyphal curvature,

which was not observed in PH-1 or ΔFgDML1-C (Fig. 2B). These findings

demonstrate that FgDML1 is a positive regulator of virulence in *F. graminearum.*

145 FgDML1 plays a critical role in both asexual and sexual reproduction.

To examine the role of FgDML1 in asexual reproduction, a series of experiments

were conducted on conidia. Morphological analysis revealed that conidia from the

ΔFgDML1 mutant were thinner and longer than those from the PH-1 and

ΔFgDML1-*C* (Fig. 2C). Furthermore, statistical analysis of septum number indicated

a significant increase in conidia with ≥7 septa in the ΔFgDML1 compared to the

controls (Fig. 2E). Conidial production was significantly reduced in ΔFgDML1, with

fewer spores produced compared to PH-1 and ΔFgDML1-C (Fig. 2F). The

germination rate of ΔFgDML1 conidia was also significantly lower than that of the

controls (Fig. 2G). Regarding sexual reproduction, ΔFgDML1 failed to produce

perithecia or ascospores on both the 7th and 14th days of incubation (Fig. 2D).

FgDML1 is essential for virulence.

To investigate the role of FgDML1 in virulence, pathogenicity assays were

performed on wheat coleoptiles and wheat leaves. The results revealed that ΔFgDML1

caused only restricted infection at the inoculation site on wheat coleoptiles and failed

to induce systemic infection, in contrast to PH-1 and ΔFgDML1-C, which exhibited

extensive colonization (Fig. 3A, B). Similarly, ΔFgDML1 displayed a near-complete

loss of virulence on wheat leaves (Fig. 3C, D). These findings demonstrate that

FgDML1 is a critical positive regulator of virulence in *F. graminearum*.

FgDML1 is essential for the biosynthesis of the DON toxin.

To evaluate whether the deletion of FgDML1 influences DON production, DON

levels were measured under toxin-inducing conditions (28°C, 145 $\times$ g, 7 days

incubation). The results revealed that DON production in ΔFgDML1 was reduced by

nearly 80% compared to PH-1 and significantly lower than that of both PH-1 and

ΔFgDML1-C (Fig. 4A). To further elucidate the regulatory relationship between

FgDML1 and DON biosynthesis, the transcriptional levels of *FgTri5* and *FgTri6* were

analyzed and found to be significantly downregulated in ΔFgDML1 (Fig. 4B).

Moreover, examination of toxisomes in the mutant demonstrated that their formation

was severely impaired, with almost no detectable toxisomes in ΔFgDML1(Fig. 4C).

Western blot analysis of the DON toxin marker protein FgTri1 further conirmed its

markedly reduced levels in ΔFgDML1 compared to PH-1 (Fig. 4D). These findings

suggest that FgDML1 is essential for regulating DON production in *F. graminearum*.

177　The deletion of FgDML1 results in reduced synthesis of both ATP and acetyl-CoA.

The results showed that ATP levels in ΔFgDML1 were significantly lower

compared to PH-1 and ΔFgDML1-C (Fig. 5D). Previous studies have established a

positive correlation between ATP levels and acetyl-CoA. Consistent with this,

acetyl-CoA levels in ΔFgDML1 were also significantly reduced compared to the

controls (Fig. 5C). In conclusion, the deletion of FgDML1 leads to a reduction in both

the key energy molecule ATP and the DON precursor acetyl-CoA.

184　FgDML1 regulates mitochondrial morphology by interacting with FgDnm1.

To assess whether the loss of FgDML1 induces mitochondrial alterations,

mitochondria-specific dyes were utilized, revealing fragmented mitochondria in the

ΔFgDML1 mutant, whereas PH-1 and ΔFgDML1-C displayed a typical network-like

mitochondrial distribution (Fig. 5A). Further analysis via Transmission Electron

Microscope (TEM) identified damaged mitochondrial structures in the ΔFgDML1

mutant, in contrast to the well-preserved mitochondrial morphology observed in the

control strains (Fig. 5B). Given that FgDnm1 is known to be a critical protein for

mitochondrial fission, the interaction between FgDML1 and FgDnm1 was verified

through co-immunoprecipitation (CoIP) assays (Fig. 5E). These findings indicate that

FgDML1 plays a pivotal role in maintaining mitochondrial morphology by interacting

with FgDnm1.

The deletion of FgDML1 reduces Complex III activity, induces compensatory upregulation of FgQCR2, FgQCR8, FgQCR9.

Deletion of FgDML1 reduced *F. graminearum* sensitivity to the Qi-site fungicide cyazofamid, while having no effect on sensitivity to the Qo-site inhibitors pyraclostrobin and trifloxystrobin, nor to other respiratory inhibitors such as pydiflumetofen, fluxapyroxad, and fluopyram (Fig. 6A, B) (Fig. S3). Measurement of Complex III activity in ΔFgDML1 showed a 38.28% reduction in enzyme activity compared to PH-1 (Fig. 6C). To further investigate the reduced sensitivity to cyazofamid, transcription levels of the three subunits (*FgCytb, FgCytC1, FgISP*) and five assembly factors (*FgQCR2, FgQCR6, FgQCR7, FgQCR8, FgQCR9*) of Complex III were analyzed. The results revealed that, upon Cyazofamid treatment, the expression of *FgQCR2*, *FgQCR7*, and *FgQCR8* was significantly upregulated in ΔFgDML1 (Fig. 6D). To further clarify whether the upregulated expression of *FgQCR2*, *FgQCR7*, and *FgQCR8* genes affects their protein expression levels, we measured the protein levels. The results showed that the protein expression levels of FgQCR2, FgQCR7, and FgQCR8 in ΔFgDML1 were higher than those in PH-1(Fig. 6F). Subsequently, we overexpressed *FgQCR2*, *FgQCR7*, and *FgQCR8* in the wild-type background, and the corresponding overexpression mutants exhibited reduced sensitivity to cyazofamid(Fig. 6E).

Overexpression of *FgQCR2*, *FgQCR8*, and *FgQCR9* may alters the conformation of the QI site, resulting in reduced sensitivity to cyazofamid.

Since Complex III is involved in the action of both cyazofamid (targeting the QI

site) and pyraclostrobin (targeting the QO site), the sensitivity of ΔFgDML1 to

cyazofamid and pyraclostrobin was investigated. The results showed that the deletion

of FgDML1 did not alter sensitivity to a pyraclostrobin (Fig. 6B). ΔFgDML1 was

subsequently treated with pyraclostrobin or cyazofamid, and Complex III enzyme

activity was measured. After pyraclostrobin treatment, no significant change in

enzyme activity was observed in ΔFgDML1 compared to PH-1 and ΔFgDML1-C (Fig.

6C). However, when treated with cyazofamid, a significant increase in Complex III

enzyme activity was observed in ΔFgDML1, while the control groups maintained

relatively low enzyme activity (Fig. 6C). These findings, combined with the

upregulation of assembly factors, suggest that the elevated expression of assembly

factors alters the protein conformation at the QI site, modulating sensitivity to

cyazofamid.

FgDML1 regulates sensitivity to various stresses.

F. graminearum encounters various stresses during infection plant, which

significantly influence its ability to infect. To investigate its response to different

stress conditions, the sensitivity of ΔFgDML1 to various environmental factors was

assessed, including osmotic stress (NaCl, KCl, sorbitol), cell wall stress (Congo red),

membrane stress (SDS), as well as high temperature (30°C) and low temperature

(15°C) stresses. The results revealed that ΔFgDML1 displayed significantly reduced

sensitivity to osmotic stress compared to PH-1 and ΔFgDML1-C (Fig. 7A, B). No

significant difference was observed in sensitivity to membrane stress, while a notable

change in sensitivity to cell wall stress was observed (Fig. 7C, D). Additionally,

differences in response to both high and low temperature stress were noted in

ΔFgDML1 when compared to the control strains PH-1 and ΔFgDML1-C (Fig. 7E, F).

Discussion

Misato Protein was originally identified in *Drosophila melanogaster*, where it was

reported to play essential roles in mitochondrial DNA inheritance, cell division, and energy

metabolism(Miklos et al., 1997). The Misato/DML1 protein family is evolutionarily

conserved from yeast to humans and plays a critical role in mitochondrial regulation.

In *S. cerevisiae*, *DML1* is an essential gene; its deletion is lethal, while its

overexpression results in fragmented mitochondrial networks and aberrant cellular

morphology, underscoring its necessity for normal mitochondrial function (Gurvitz et

al., 2002). Similarly, in *Homo sapiens*, the homolog Misato localizes to the

mitochondrial outer membrane, and both its depletion and overexpression are

sufficient to disrupt mitochondrial morphology and distribution (Kimura and Okano,

2007). However, its functional characterization in filamentous fungi remains largely

unexplored. Our study defines a novel mechanism through which FgDML1 governs

mitochondrial homeostasis. We demonstrate that FgDML1 directly interacts with the

key mitochondrial fission regulator FgDnm1 and positively modulates cellular

bioenergetic metabolism, as evidenced by elevated ATP and acetyl-CoA levels (Fig. 8).

In this study, we identified and analyzed the homolog FgDML1 in the

phytopathogenic fungus *F. graminearum*. The FgDML1 protein contains conserved

Misato_Tub_SegII and Tubulin_3 domains, with high homology observed among

species within the *Fusarium* genus but significant genetic divergence from other taxa

(Fig. 1A). In *S. cerevisiae*, DML1 is a critical regulator of mitochondrial fission and

fusion dynamics, maintaining intracellular energy homeostasis by modulating

mitochondrial morphology and distribution(Gurvitz et al., 2002). Our study

demonstrated that FgDML1 localized to mitochondria in *F. graminearum* (Fig. 1B).

Deletion mutants of *FgDML1* exhibited phenotypes indicative of mitochondrial

dysfunction, including fragmented mitochondrial morphology (Fig. 5A). TEM further

confirmed structural and dynamic damage to mitochondria in the ΔFgDML1 (Fig. 5B).

Structural analysis suggests that DML1 shares similarities with GTPase family

proteins, indicating potential GTPase activity(Gurvitz et al., 2002). GTPases primarily

hydrolyze GTP, but their activity is intricately linked to ATP production and

utilization. These enzymes regulate critical cellular processes such as energy

metabolism, signal transduction, protein synthesis, and cytoskeletal remodeling, all of

which influence ATP levels(Mrnjavac and Martin, 2025; Pb et al., 2023; Rogne et al.,

2018). In agreement with this, we observed a significant reduction in ATP content in

the FgDML1 deletion mutant, underscoring its role in mitochondrial energy

metabolism (Fig. 5D). In *H. sapiens*, DML1 also localizes to mitochondria, where its

silencing results in fragmented and aggregated mitochondria with reduced network

continuity and fusion activity (Kimura and Okano, 2007). Our findings indicate that

FgDML1 plays a conserved role in maintaining mitochondrial integrity and energy

homeostasis. Notably, FgDML1 was found to interact with FgDnm1 (Fig. 5E),

FgDnm1 is a key dynamin-related protein mediating mitochondrial fission(Griffin et

al., 2005; Kang et al., 2023), suggesting that FgDML1 may form a complex with

FgDnm1 to regulate mitochondrial fission and fusion processes. To our knowledge,

this is the first report documenting an interaction between DML1 and Dnm in any

fungal species, including model organisms such as *S. cerevisiae*. This novel finding

provides new insights into the molecular mechanisms underlying mitochondrial

dynamics in filamentous fungi.

Secondary metabolite biosynthesis is generally regarded as an energy-intensive

process that is tightly coupled to cellular energy metabolism. ATP serves as the

primary energy currency supporting enzymatic reactions, macromolecule synthesis,

and subcellular organization required for secondary metabolism. Disruption of ATP

generation has been shown to directly impair toxin biosynthesis: for example,

silencing of ATP synthase subunit α (AtpA) significantly reduces ATP synthesis and

inhibits the production of the TcdA and TcdB toxins(Marreddy et al., 2024). Similarly,

in plants, ATP depletion leads to a metabolic shift in which growth and basic

physiological processes are prioritized at the expense of energetically costly

secondary metabolites, including toxins(Xiao et al., 2024). Together, these findings

highlight ATP availability as a key determinant of secondary metabolite production

across biological systems.

In filamentous fungi, mitochondria play a central role in sustaining cellular ATP

levels through oxidative phosphorylation and are therefore critical for biosynthetic

and stress-adaptive processes. In *F. graminearum*, mutants defective in mitochondrial

components, such as the voltage-dependent anion channel (mitochondrial porin),

exhibit aberrant mitochondrial morphology, reduced ATP production, and markedly

decreased DON accumulation and virulence (Han et al., 2022). These observations

establish a direct link between mitochondrial energy metabolism and secondary

metabolite output, supporting the notion that intact mitochondrial function and

adequate ATP supply are prerequisites for robust DON production.

Consistent with this energy-dependent framework, biosynthesis of the mycotoxin

DON in *F. graminearum* requires substantial ATP input. In the present study, ATP

content in the ΔFgDML1 mutant was significantly lower than in the wild-type PH-1

and the complemented strain ΔFgDML1-C, and DON production was concomitantly

reduced (Fig. 4A). Importantly, DON levels were normalized to mycelial dry weight,

indicating that the observed reduction reflects a decreased biosynthetic capacity per

unit biomass rather than a secondary consequence of reduced fungal growth. This

distinction demonstrates that impaired DON production in the ΔFgDML1 mutant

arises primarily from metabolic limitations.

At the cellular level, ATP depletion compromises multiple energy-dependent

steps required for DON biosynthesis. The formation of toxisomes, which are

specialized subcellular structures responsible for the spatial organization of DON

biosynthetic enzymes, is essential for efficient mycotoxin production and is an

ATP-dependent process. Reduced ATP levels disrupt toxisome assembly, and

accordingly, the ΔFgDML1 mutant was unable to form functional toxisomes (Fig. 4C).

In parallel, western blot analysis revealed a marked reduction in the abundance of the

DON biosynthetic enzyme FgTri1 (Fig. 4D). In addition, ATP-dependent processes

are directly involved in the biogenesis of the DON biosynthetic machinery: the

ATPase activity of myosin I (FgMyo1) is required for efficient translation of key

DON biosynthetic enzymes, and disruption of its ATPase function results in reduced

DON production(Tang et al., 2018). These findings further underscore the dependence

of DON biosynthesis on cellular energy status.

DON production is also regulated at the transcriptional level by the TRI gene

cluster, with Tri5 and Tri6 serving as core components of the biosynthetic pathway.

Tri5 encodes trichodiene synthase, which catalyzes the first committed step of DON

biosynthesis. In the ΔFgDML1 mutant, expression levels of FgTri5 and FgTri6 were

significantly downregulated (Fig. 4B), suggesting that impaired energy metabolism

indirectly affects transcription of DON biosynthetic genes. Although no direct

regulatory role of DML family proteins in gene expression has been reported in

Saccharomyces cerevisiae or Drosophila melanogaster, their established functions in

cell division and microtubule organization raise the possibility that FgDML1

indirectly influences gene expression through effects on chromatin organization or

cell-cycle progression(Schulze and Wallrath, 2007).

In addition to reduced ATP levels, deletion of FgDML1 resulted in a significant

decrease in acetyl-CoA content (Fig. 5C), a key precursor for trichothecene

biosynthesis. Acetyl-CoA links central carbon metabolism with secondary metabolite

production, and its depletion further constrains DON biosynthesis by limiting

substrate availability. Broader metabolomic studies support this relationship, showing

that perturbations in TCA cycle intermediates and central carbon metabolism are

closely associated with altered DON production, reinforcing a mechanistic linkage

between energy generation and toxin biosynthesis(Atanasova-Penichon et al., 2018).

Taken together, these results support a model in which FgDML1 influences DON

production indirectly by maintaining mitochondrial energy metabolism. Reduced ATP

availability in the ΔFgDML1 mutant restricts energy-dependent biosynthetic

processes, disrupts toxisome formation, diminishes DON biosynthetic enzyme

abundance and gene expression, and limits precursor supply, ultimately leading to a

substantial reduction in DON biosynthesis that is independent of fungal biomass

effects.

F. graminearum is a highly destructive pathogen that poses a significant threat to

wheat production worldwide. Cyazofamid is an effective fungicide commonly used to

control plant pathogenic fungi. Cyazofamid specifically targets the Qi site of complex

III in the mitochondrial respiratory chain(Bolgunas et al., 2006; Mitani et al., 2001).

Complex III, also known as the cytochrome bc1 complex, plays a critical role in the

electron transport chain by transferring electrons from ubiquinone to cytochrome c

and pumping protons from the mitochondrial matrix into the intermembrane

space(Fernández-Ortuño et al., 2008; Sarewicz et al., 2013). This complex consists of

several core subunits, including cytochrome b, cytochrome c1, and iron-sulfur

proteins, as well as multiple assembly factors. It contains two quinone binding sites:

the Qo site (quinone oxidation site) and the Qi site (quinone reduction site)(Vercellino

and Sazanov, 2022; Zara et al., 2009). Numerous fungicides have been developed to

inhibit the Qo site (e.g., pyraclostrobin, azoxystrobin)(Nuwamanya et al., 2022; Peng

et al., 2022) and the Qi site (e.g., cyazofamid)(Mitani et al., 2001) of the cytochrome

bc1 complex. In our study, we observed that the deletion of *FgDML1* significantly

reduced the sensitivity of *F. graminearum* to cyazofamid (Fig. 6A), while its

sensitivity to pyraclostrobin remained unaffected (Fig. 6B). In the ΔFgDML1 mutant,

we detected a decrease in complex III enzyme activity (Fig. 6C) and noted the

upregulation of assembly factors *FgQCR2*, *FgQCR7*, and *FgQCR8* following

cyazofamid treatment (Fig. 6D). Further enzyme activity assays revealed that after

pyraclostrobin treatment, ΔFgDML1 exhibited a 73.06% reduction in complex III

activity, compared to an 80.38% decrease in the wild-type strain PH-1 (Fig. 6C).

Interestingly, after cyazofamid treatment, the decrease in enzyme activity in

ΔFgDML1 was only 36.31%, while PH-1 showed a dramatic 83.46% reduction in

complex III activity. *FgQCR2*, *FgQCR7*, and *FgQCR8* are critical assembly factors

that play essential roles in the proper assembly of complex III(Zhan et al., 2024).

These factors stabilize the core subunit structure, promoting correct folding and

functional operation of complex III, particularly during the assembly of the Qi site.

This ensures the proper formation and function of the Qi site, thus supporting the

normal operation of the mitochondrial respiratory chain(Wang et al., 2024; Zara et al.,

2007). Previous studies have reported that mutations in the Complex III assembly

factors TTC19, UQCC2, and UQCC3 impair the assembly and activity of Complex III

(Feichtinger et al., 2017; Wanschers et al., 2014). Given the role of FgDML1 in

mitochondrial homeostasis, we further evaluated the sensitivity of the ΔFgDML1

mutant to other commonly used respiratory chain inhibitors, including pydiflumetofen,

fluxapyroxad, fluopyram, and trifloxystrobin. The results indicated that FgDML1

deletion had no impact on the sensitivity of *F. graminearum* to these fungicides (Fig.

S3). In conclusion, our findings suggest that the overexpression of assembly factors

FgQCR2, *FgQCR7*, and *FgQCR8* in ΔFgDML1 potentially modifies the conformation

of the Qi site, which specifically modulates the sensitivity of *F. graminearum* to

cyazofamid.

In *H. sapiens*, mutations or deletions in DML1 are associated with mitochondrial

diseases characterized by early-onset myopathy and cerebellar ataxia(Chen et al.,

2022; Nasca et al., 2017). In our study, the deletion of *FgDML1* in *F. graminearum*

resulted in significant defects in vegetative growth, including a markedly reduced

growth rate (Fig. 2A) and an aberrant, excessively curved hyphal morphology (Fig.

2B). Notably, the loss of FgDML1 rendered *F. graminearum* incapable of completing

sexual reproduction (Fig. 2D), thereby potentially reducing its primary inoculum

sources. Additionally, the deletion of FgDML1 impaired asexual reproduction and

pathogenicity (Fig. 2) (Fig. 3), critical factors for fungal survival and host infection.

Furthermore, the ΔFgDML1 mutant exhibited altered responses to various stress

conditions (Fig. 7), suggesting a broader role for FgDML1 in regulating fungal stress

tolerance.

In the present study, FgDML1 is localized to the mitochondria and interacted

with FgDnm1, a key regulator of mitochondrial morphology, positively regulates ATP

and acetyl-CoA levels, influences toxin body formation, and ultimately leads to a

significant reduction in DON toxin biosynthesis. Furthermore, deletion of FgDML1

impairs the enzymatic activity of Complex III, which subsequently induces the

upregulation of the assembly factors FgQCR2, FgQCR8, and FgQCR9, resulting in

conformational changes at the Qi site of Complex III and decreased sensitivity to

cyazofamid. In addition, the absence of FgDML1 has profound effects on

pathogenicity, fungal development, and mitochondrial morphology. Based on these

findings, we propose a working model delineating the role of FgDML1 in regulating

DON toxin biosynthesis and cyazofamid sensitivity in *F. graminearum* (Fig. 8). This

study not only provides a theoretical basis for understanding the regulatory

mechanisms underlying toxin biosynthesis in *F. graminearum* but also identifies

potential molecular targets for the development of novel agents to control FHB.

Material and methods

Strains, culture conditions, and fungicides.

The wild-type *F. graminearum* strain PH-1 was used as the parental strain for

protoplast transformation experiments. All strains were preserved at -80°C. The tested

strains were cultured in the dark on potato dextrose agar (PDA) at 25°C. To evaluate

vegetative growth, complete medium (CM), minimal medium (MM), and V8 Juice

Agar (V8) media were prepared as described previously(Tang et al., 2020). Mung

bean liquid (MBL) medium was used for conidial production, while carrot agar (CA)

medium was utilized to assess sexual reproduction(Wang et al., 2011). DON toxin

production was measured using trichothecene biosynthesis-inducing (TBI)

medium(Wang et al., 2024).

Cyazofamid (98%, BASF China Ltd.), pyraclostrobin (98%, BASF China Ltd.)

pydiflumetofen (98%, China Syngenta crop Protection Co., Ltd.), trifloxystrobin

(96%, Bayer Crop Science China Co., Ltd.), fluxapyroxad (98%, BASF China Ltd.)

and fluopyram (96%, Bayer Crop Science China Co., Ltd.) were dissolved in dimethyl

sulfoxide (DMSO) at stock concentrations of 50,000 μg/mL and 10,000 μg/mL,

respectively.

Generation of deletion mutants and complemented mutants.

The split-marker method was employed to generate the ΔFgDML1 mutant(Yu et

al., 2004). Specifically, approximately 1.2 kb of the upstream and downstream

flanking regions of FgDML1 were amplified and fused with a dual-selection marker

containing the hygromycin phosphotransferase (*HPH*) gene and the herpes simplex

virus thymidine kinase (*HSV-tk*) gene fragments. The resulting fusion fragment was

transformed into the wild-type *F. graminearum* PH-1 strain via polyethylene glycol

(PEG)-mediated protoplast transformation. Transformants were selected on PDA

plates containing either 100 μg/mL Hygromycin B (Yeasen, Shanghai, China) or 0.2

μmol/mL 5-Fluorouracil 2'-deoxyriboside (F2du) (Solarbio, Beijing, China)(Zhao et

al., 2022). Positive transformants were initially screened by PCR and subsequently

verified through Southern blot analysis. To generate the complemented mutant

ΔFgDML1-C, the full-length coding sequence (CDS) of FgDML1 was introduced into

the ΔFgDML1 mutant, replacing the dual-selection cassette. Furthermore,

FgDML1-GFP and FgDnm1-3×Flag mutants were constructed using previously

established methods for further functional studies(Wu et al., 2022). Specifically,

FgDML1, including its native promoter region and open reading frame (ORF)

(excluding the stop codon), was amplified.The PCR product was then fused with the

*Xho*I-digested pYF11 vector. After transformation into E. coli and sequence

verification, the plasmid was extracted and subsequently introduced into PH-1

protoplasts. For FgDnm1-3×Flag, the 3×Flag tag was added to the C-terminus of

FgDnm1 by PCR, fused with the hygromycin resistance gene and the FgDnm1

downstream arm, and then introduced into PH-1 protoplasts. The overexpression

mutant was constructed according to a previously described method. Specifically, the

ORF of FgDML1 was amplified and the PCR product was ligated into the

*Sac*II-digested pSXS overexpression vector. The resulting plasmid was then

transformed into PH-1 protoplasts (Shi et al., 2023). For the construction of

PH-1::FgTri1+GFP and ΔFgDML1::FgTri1+GFP, the ORF of FgTri1 was amplified

and ligated into the *Xho*I-digested pYF11 vector as described above. The resulting

vectors were then transformed into protoplasts of PH-1 or ΔFgDML1, respectively.

Vegetative growth and conidiation assays.

Fresh mycelial plugs of the tested strains were transferred to PDA, CM, MM, or

V8 plates to evaluate mycelial growth rates. Conidial production was induced by

incubating the strains in MBL medium, and the number of conidia was determined

using a hemocytometer. The septa of conidia were visualized through calcofluor white

(CFW) staining. For germination rate assessment, conidia were collected and

inoculated onto water agar (WA) plates, followed by incubation at 25°C for 2 and 8

hours. Germination rates were then recorded at the respective time points. For each

sample, a total of 200 conidia were counted. The experiment included three biological

replicates with three technical replicates each.

Sexual reproduction assay.

Sexual reproduction experiments were performed on CA medium. Test strains

were incubated on CA medium in complete darkness for 6 days. After incubation, the

surface mycelia were removed using a mycelial scraper and 2.5% Tween 60 solution.

The plates were then further incubated at 25°C under a 12-hour light/12-hour dark

cycle to induce sexual development. Perithecia and ascospores were observed and

recorded on days 7 and 14 using a Nikon SMZ25 fluorescent stereomicroscope and an

Olympus IX-71 inverted fluorescent microscope. Each treatment group contains three

biological replicates.

Microscopic observation.

To determine the subcellular localization of FgDML1, the FgDML1-GFP mutant

was incubated in YEPD medium at 25°C with shaking at $145 \times g$ for 36 hours.

Subcellular localization was analyzed using a Leica TCS SP8 confocal microscope,

with MitoTracker Red CMXRos (Yeasen, Shanghai, China) applied to confirm

mitochondrial targeting through colocalization of fluorescence signals. To investigate

toxisome morphology, the ΔFgDML1::Tri1+GFP mutant was cultured in TBI medium

at 28°C with shaking for 36 hours, and the structural dynamics of toxisomes within

the mycelium were visualized using Tri1-GFP fluorescence under confocal

microscopy. Additionally, mitochondrial morphology was evaluated by culturing

PH-1, ΔFgDML1, and ΔFgDML1-C in YEPD medium under identical conditions,

followed by staining with MitoTracker Red CMXRos and confocal microscopy

analysis to assess mitochondrial structural alterations, distribution, and dynamics. For

ultrastructural observations, mycelial samples were fixed and dehydrated according to

standard protocols, embedded, sectioned, and stained with uranyl acetate and lead

citrate. Mitochondrial ultrastructure was examined using a Tecnai G2 Spirit Bio

electron microscope (Thermo, MA, USA) at an accelerating voltage of 80 kV to

compare the structural integrity and morphological differences among the strains.

Each treatment group contains three biological replicates and three technical

replicates.

Determination of DON production and pathogenicity.

As previously described (Tang et al., 2020; Wang et al., 2025), Specifically,

coleoptiles were inoculated with conidial suspensions and incubated for 14 days,

while leaves were inoculated with fresh mycelial plugs and incubated for 5 days,

followed by observation and quantification of disease symptoms. DON toxin was

measured using a Wise Science ELISA-based kit (Wise Science, Jiangsu, China) (Li

et al., 2019; Zheng et al., 2018). Under toxin-producing conditions (28 °C, 145 rpm),

fungal strains were cultured in TBI medium for 7 days. Cultures were initiated using

freshly grown mycelia. After incubation, mycelia and culture filtrates were separated

by filtration. The culture filtrates were collected for DON determination, while the

mycelia were harvested for biomass analysis. The collected mycelia were washed

with sterile distilled water and dried at 60 °C to constant weight. The dry weight of

mycelia was recorded and used for normalization of DON production. One mycelial

unit was defined as 1 g of dry mycelial biomass. DON concentration in the culture

filtrates was quantified using an enzyme-linked immunosorbent assay (ELISA).

Briefly, 50 μL of culture filtrate or DON standard solution was added to wells of a

96-well microplate pre-coated with DON antigen, followed by the addition of enzyme

conjugate and antibody working solution according to the manufacturer's instructions.

After incubation and washing, color development was achieved using substrate

solution and terminated by stop solution. Absorbance was measured at 450 nm using a

microplate reader. A standard curve was generated using log□□-transformed DON

concentrations of the standards and the corresponding percentage absorbance values.

DON concentrations in the samples were calculated based on the standard curve. Total

DON production was calculated according to the culture volume (30 mL) and

subsequently normalized to mycelial dry weight. DON production was expressed as

μg DON per g dry mycelium. Each treatment group contains three biological

replicates and three technical replicates.

Determination of acetyl-CoA content and ATP concentration.

ATP content and acetyl-CoA levels were determined using the methods described

previously(Hu et al., 2023; Song et al., 2024). The test strains were cultured in yeast

extract peptone dextrose (YEPD) medium at 25°C and 145 × g for 36 hours. Mycelia

were collected by filtration through three layers of lens-cleaning paper, and excess

moisture was removed with filter paper. Subsequently, ATP and acetyl-CoA levels

were quantified using ATP and acetyl-CoA assay kits (Solarbio, Beijing, China),

respectively. Each treatment group contains three biological replicates and three

technical replicates.

Sensitivity determination.

To investigate the sensitivity of ΔFgDML1 to QoI and QiI fungicides, the

wild-type strain PH-1, ΔFgDML1 mutant, and the complemented mutant

ΔFgDML1-C were inoculated on AEA plates supplemented with gradient

concentrations of cyazofamid and pyraclostrobin. The wild-type PH-1 and its

overexpression mutants (PH-1::FgQCR2-OE, PH-1::FgQCR7-OE, and

PH-1::FgQCR8-OE) were assayed on AEA plates amended with cyazofamid and

supplemented with SHAM. All strains were incubated at 25°C in darkness; however,

due to ΔFgDML1 slower growth, the ΔFgDML1 mutant required a 5-day incubation

period compared to the 3 days used for PH-1 and ΔFgDML1-C. The concentration

gradients for each fungicide in the sensitivity assays were set up according to

Supplementary Table S2. To inhibit the alternative respiratory pathway, 50 µg/mL

SHAM was applied. The percentage inhibition (%) of mycelial growth was calculated

using the formula: (mean control diameter - mean diameter at fungicide concentration)

/ (mean control diameter - 5 mm mycelial plug) × 100. To evaluate the sensitivity of

ΔFgDML1 to various stress conditions, the test strains were cultured on PDA plates

supplemented with stress-inducing agents, including NaCl, KCl, sorbitol, Congo red,

and SDS. Plates were incubated at 25°C for 3 days, and colony growth was assessed.

Each treatment group contains three biological replicates.

Measurement of complex III enzyme activity.

The enzyme activity of Complex III was determined in the PH-1, ΔFgDML1,

and ΔFgDML1-C. Fresh mycelial plugs from each strain were inoculated into liquid

AEA medium and incubated at 25°C with shaking at 145 × *g* for 36 hours.

Subsequently, cyazofamid and pyraclostrobin were added to final concentrations of 30

µg/mL and 1.5 µg/mL, respectively, along with 50 µg/mL SHAM. After an additional

12-hour incubation, the mycelia were harvested, and Complex III activity was

measured according to the complex III enzyme activity kit (Solarbio, Beijing, China)

instructions. Briefly, 0.1 g of mycelia was homogenized with 1 mL of extraction

buffer in an ice bath. The homogenate was centrifuged at 600 ×g for 10 min at 4°C.

The resulting supernatant was then subjected to a second centrifugation at 11,100 ×g

for 10 min at 4°C. The pellet was resuspended in 200 µL of extraction buffer and

disrupted by ultrasonication (200 W, 5 s pulses with 10 s intervals, 15 cycles).

Complex III enzyme activity was finally measured by adding the working solution as

per the manufacturer's protocol. Each treatment group contains three biological

replicates and three technical replicates.

RNA extraction and RT-qPCR assays.

The test strains were transferred to liquid AEA medium and incubated at 25°C

for 48 hours, with cyazofamid added to a final concentration of 20 µg/mL. The

control group was not treated with any fungicide. To measure the transcriptional

levels of genes involved in DON biosynthesis, the test strains were transferred to TBI

medium and incubated for 3 days. Mycelium was collected and total RNA was

extracted following the instructions provided by the Total RNA Extraction Kit

(Tiangen, Beijing, China). cDNA synthesis was performed using Evo M-MLV Mix

Kit (Accurate, Hunan, China). reverse transcription. Reverse Transcription

Quantitative Polymerase Chain Reaction (RT-qPCR) was carried out using the

QuantStudio 6 Flex real-time PCR system (Thermo, Fisher Scientific, USA) to assess

the relative expression of three subunits of Complex III (*FgCytb*, *FgCytc1*, *FgISP*),

five assembly factors (*FgQCR2*, *FgQCR6*, *FgQCR7*, *FgQCR8*, *FgQCR9*), and DON

biosynthesis-related genes (*FgTri5* and *FgTri6*). The *FgCox1* gene was used as the

reference gene for *FgCytb*, while *FgGapdh* was used as the reference for the other

genes. Relative transcript levels were calculated using the $2^{-\Delta\Delta CT}$ method. Each

treatment group contains three biological replicates and three technical replicates.

Western Blot assay and Co-IP assay.

Total protein extraction was performed according to previously established

protocols, and the extracted proteins were separated using sodium dodecyl

sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Subsequently, proteins were

transferred onto an Immobilon-P transfer membrane (Millipore, Billerica, USA). The

membrane was probed with GFP antibody D191040 (BBI, Shanghai, China) and Flag

antibody 30501ES60 (Yeasen, Shanghai, China). The FgDnm1-3×Flag fragment was

introduced into PH-1 and FgDML1+GFP protoplasts, respectively, to obtain

single-tagged and double-tagged strains. All tagged mutant strains were subjected to

triple validation, including PCR, sequencing, and Western blot analysis. Total protein

from tagged strains was incubated with ChromoTek GFP-Trap® Magnetic Agarose

(ChromoTek, Planegg-Martinsried, Germany) beads. Proteins eluted from the beads

were further co-incubated with GFP or Flag antibodies for detection. Gapdh antibody

60004-1-Ig (Proteintech, Rosemont, USA) was used as an internal control for protein

normalization. Chemiluminescent detection was performed following incubation with

the corresponding secondary antibody. All antibodies were diluted as follows: primary

antibodies at 1:1000 and secondary antibodies at 1:10000. The experiment was

independently repeated three times.

Statistical analysis.

All data were analyzed using one-way analysis of variance (ANOVA), followed

by the least significant difference (LSD) test with a significance threshold of $p < 0.05$.

The EC_{50} values were calculated using DPS statistical analysis software. Each

experiment was independently repeated three times.

Acknowledgements

CRediT authorship contribution statement

Chenguang Wang: methodology, investigation, editing, and writing original draft.

Xuewei Mao: formal analysis, data curation. Weiwei Cong, Lin Yang.: performing the

partial work of the experiments and analyzing the data. Yiping Hou: conceptualization,

resources, writing–reviewing, supervision, funding acquisition.

Declaration of competing interest

The authors declare no conflict of interest.

References

Alexander NJ, Proctor RH, McCormick SP. 2009. Genes, gene clusters, and biosynthesis of trichothecenes and fumonisins in fusarium. Toxin Reviews **28**:198–215. DOI: https://doi.org/10.1080/15569540903092142

Andrade SMP, Augusti GR, Paiva GF, Feksa HR, Tessmann DJ, Machado FJ, Mizubuti ESG, Del Ponte EM. 2022. Phenotypic and molecular characterization of the resistance to azoxystrobin and pyraclostrobin in Fusarium graminearum populations from Brazil. Plant Pathology **71**:1152–1163. DOI: https://doi.org/10.1111/ppa.13535

Atanasova-Penichon V, Legoahec L, Bernillon S, Deborde C, Maucourt M, Verdal-Bonnin M-N, Pinson-Gadais L, Ponts N, Moing A, Richard-Forget F. 2018. Mycotoxin biosynthesis and central metabolism are two interlinked pathways in fusarium graminearum, as demonstrated by the extensive metabolic changes induced by caffeic acid exposure. Applied and Environmental Microbiology **84**:e01705-17. DOI: https://doi.org/10.1128/AEM.01705-17, PMID: 29427428

Boenisch MJ, Broz KL, Purvine SO, Chrisler WB, Nicora CD, Connolly LR, Freitag M, Baker SE, Kistler HC. 2017. Structural reorganization of the fungal endoplasmic reticulum upon induction of mycotoxin biosynthesis. Scientific Reports **7**:44296. DOI: https://doi.org/10.1038/srep44296, PMID: 28287158

Bolgunas S, Clark DA, Hanna WS, Mauvais PA, Pember SO. 2006. Potent inhibitors of the qi site of the mitochondrial respiration complex III. Journal of Medicinal Chemistry **49**:4762–4766. DOI: https://doi.org/10.1021/jm060408s

Bönnighausen J, Schauer N, Schäfer W, Bormann J. 2019. Metabolic profiling of wheat rachis node infection by Fusarium graminearum – decoding deoxynivalenol-dependent susceptibility. New Phytologist **221**:459–469. DOI: https://doi.org/10.1111/nph.15377

Chen J, Xiao J, Chen G, Xu Q, Wu X, Tian L, Huang Z, Xin C, Zhao Y, Guo Z, Zou Y, Wu Q. 2022. Indentification of novel MSTO1 compound heterozygous mutations in a Chinese family with recessive cerebellar atrophy and ataxia. Frontiers in Neurology **13**:988519. DOI: https://doi.org/10.3389/fneur.2022.988519, PMID: 36468072

Dean R, Van Kan JAL, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD, Rudd JJ, Dickman M, Kahmann R, Ellis J, Foster GD. 2012. The Top 10 fungal pathogens in molecular plant pathology. Molecular Plant Pathology **13**:414–430. DOI: https://doi.org/10.1111/j.1364-3703.2011.00783.x, PMID: 22471698

Duan Y, Lu F, Zhou Z, Zhao H, Zhang J, Mao Y, Li M, Wang J, Zhou M. 2020. Quinone outside inhibitors affect DON biosynthesis, mitochondrial structure and toxisome formation in *fusarium graminearum*. Journal of Hazardous Materials **398**:122908. DOI: https://doi.org/10.1016/j.jhazmat.2020.122908

Feichtinger RG, Brunner-Krainz M, Alhaddad B, Wortmann SB, Kovacs-Nagy R, Stojakovic T, Erwa W, Resch B, Windischhofer W, Verheyen S, Uhrig S, Windpassinger C, Sternberg F, Makowski C, Strom TM, Meitinger T, Prokisch H, Sperl W, Haack TB, Mayr JA. 2017. Combined respiratory chain deficiency and UQCC2 mutations in neonatal encephalomyopathy: defective supercomplex assembly in complex III deficiencies. Oxidative Medicine and Cellular Longevity **2017**:7202589. DOI: https://doi.org/10.1155/2017/7202589

Fernández-Ortuño D, Torés JA, de Vicente A, Pérez-García A. 2008. Mechanisms of
resistance to QoI fungicides in phytopathogenic fungi. International Microbiology:
The Official Journal of the Spanish Society for Microbiology **11**:1–9. PMID:
18683626
Griffin EE, Graumann J, Chan DC. 2005. The WD40 protein Caf4p is a component of the
mitochondrial fission machinery and recruits Dnm1p to mitochondria. The Journal of
Cell Biology **170**:237–248. DOI: https://doi.org/10.1083/jcb.200503148, PMID:
16009724
Gurvitz A, Hartig A, Ruis H, Hamilton B, de Couet HG. 2002. Preliminary characterisation of
DML1, an essential *saccharomyces cerevisiae* gene related to *misato* of *drosophila*
melanogaster. FEMS Yeast Research **2**:123–135. DOI:
https://doi.org/10.1016/S1567-1356(02)00083-1
Hai S, Chen J, Ma L, Wang C, Chen C, Rahman SU, Zhao C, Feng S, Wu J, Wang X. 2023.
Combination of zearalenone and deoxynivalenol induces apoptosis by mitochondrial
pathway in piglet sertoli cells: role of endoplasmic reticulum stress. Toxins **15**:471.
DOI: https://doi.org/10.3390/toxins15070471, PMID: 37505740
Han X, Li Q, Li X, Lv X, Zhang L, Zou S, Yu J, Dong H, Chen L, Liang Y. 2022.
Mitochondrial porin is involved in development, virulence, and autophagy in
fusarium graminearum. Journal of Fungi **8**:936. DOI:
https://doi.org/10.3390/jof8090936
Hu L, Guo C, Chen J, Jia R, Sun Y, Cao S, Xiang P, Wang Y. 2023. Venturicidin a is a
potential fungicide for controlling fusarium head blight by affecting deoxynivalenol
biosynthesis, toxisome formation, and mitochondrial structure. Journal of Agricultural
and Food Chemistry **71**:12440–12451. DOI: https://doi.org/10.1021/acs.jafc.3c02683
Huang P, Yu X, Liu H, Ding M, Wang Z, Xu J-R, Jiang C. 2024. Regulation of TRI5
expression and deoxynivalenol biosynthesis by a long non-coding RNA in fusarium
graminearum. Nature Communications **15**:1216. DOI:
https://doi.org/10.1038/s41467-024-45502-w
Huang Q, Tonge PJ, Slayden RA, Kirikae T, Ojima I. 2007. FtsZ: a novel target for
tuberculosis drug discovery. Current Topics in Medicinal Chemistry **7**:527–543. DOI:
https://doi.org/10.2174/156802607780059790, PMID: 17346197
Ilgen P, Hadeler B, Maier FJ, Schäfer W. 2009. Developing kernel and rachis node induce the
trichothecene pathway of fusarium graminearum during wheat head infection.
Molecular Plant-Microbe Interactions® **22**:899–908. DOI:
https://doi.org/10.1094/MPMI-22-8-0899
Jiang C, Zhang C, Wu C, Sun P, Hou R, Liu H, Wang C, Xu J-R. 2016. TRI6 and TRI10 play
different roles in the regulation of deoxynivalenol (DON) production by cAMP
signalling in fusarium graminearum. Environmental Microbiology **18**:3689–3701.
DOI: https://doi.org/10.1111/1462-2920.13279
Kang J, Zhang J, Liu Y, Song J, Ou J, Tao X, Zhou M, Duan Y. 2023. Mitochondrial dynamics
caused by QoIs and SDHIs fungicides depended on FgDnm1 in *fusarium*
graminearum. Journal of Integrative Agriculture **22**:481–494. DOI:
https://doi.org/10.1016/j.jia.2022.08.118

Kimura M, Anzai H, Yamaguchi I. 2001. Microbial toxins in plant-pathogen interactions:
biosynthesis, resistance mechanisms, and significance. Journal of General and
Applied Microbiology **47**:149–160. DOI: https://doi.org/10.2323/jgam.47.149, PMID:
12483615

Kimura M, Okano Y. 2007. Human misato regulates mitochondrial distribution and
morphology. Experimental Cell Research **313**:1393–1404. DOI:
https://doi.org/10.1016/j.yexcr.2007.02.004

Kimura M, Tokai T, Takahashi-Ando N, Ohsato S, Fujimura M. 2007. Molecular and genetic
studies of fusarium trichothecene biosynthesis: pathways, genes, and evolution.
Bioscience, Biotechnology, and Biochemistry **71**:2105–2123. DOI:
https://doi.org/10.1271/bbb.70183, PMID: 17827683

Kistler HC, Broz K. 2015. Cellular compartmentalization of secondary metabolism. Frontiers
in Microbiology **6**. DOI: https://doi.org/10.3389/fmicb.2015.00068

Knutsen HK, Alexander J, Barregård L, Bignami M, Brüschweiler B, Ceccatelli S, Cottrill B,
Dinovi M, Grasl Kraupp B, Hogstrand C, Hoogenboom L (Ron), Nebbia CS,
Oswald IP, Petersen A, Rose M, Roudot A, Schwerdtle T, Vleminckx C, Vollmer G,
Wallace H, De Saeger S, Eriksen GS, Farmer P, Fremy J, Gong YY, Meyer K, Naegeli
H, Parent Massin D, Rietjens I, van Egmond H, Altieri A, Eskola M, Gergelova P,
Ramos Bordajandi L, Benkova B, Dörr B, Gkrillas A, Gustavsson N, van Manen M,
Edler L. 2017. Risks to human and animal health related to the presence of
deoxynivalenol and its acetylated and modified forms in food and feed. EFSA Journal
15:e04718. DOI: https://doi.org/10.2903/j.efsa.2017.4718, PMID: 32625635

Li H, Zhu X-L, Yang W-C, Yang G-F. 2014. Comparative Kinetics of Qi Site Inhibitors of
Cytochrome bc1 Complex: Picomolar Antimycin and Micromolar Cyazofamid.
Chemical Biology & Drug Design **83**:71–80. DOI:
https://doi.org/10.1111/cbdd.12199

Li J, Duan Y, Bian C, Pan X, Yao C, Wang J, Zhou M. 2019. Effects of validamycin in
controlling fusarium head blight caused by fusarium graminearum: inhibition of DON
biosynthesis and induction of host resistance. Pesticide Biochemistry and Physiology
153:152–160. DOI: https://doi.org/10.1016/j.pestbp.2018.11.012, PMID: 30744889

Liu J, Cao Q, Ewing M, Zuo Z, Kennerdell JR, Finkel T, Leppla SH, Liu S. 2025. ATP
depletion in anthrax edema toxin pathogenesis. PLOS Pathogens **21**:e1013017. DOI:
https://doi.org/10.1371/journal.ppat.1013017, PMID: 40168442

Liu X, Yu F, Schnabel G, Wu J, Wang Z, Ma Z. 2011. Paralogous cyp51 genes in Fusarium
graminearum mediate differential sensitivity to sterol demethylation inhibitors.
Fungal genetics and biology: FG & B **48**:113–123. DOI:
https://doi.org/10.1016/j.fgb.2010.10.004, PMID: 20955812

Liu Y, Lu Y, Wang L, Chang F, Yang L. 2016. Occurrence of deoxynivalenol in wheat, hebei
province, china. Food Chemistry **197**:1271–1274. DOI:
https://doi.org/10.1016/j.foodchem.2015.11.047

Liu Z, Jian Y, Chen Y, Kistler HC, He P, Ma Z, Yin Y. 2019. A phosphorylated transcription
factor regulates sterol biosynthesis in Fusarium graminearum. Nature
Communications **10**:1228. DOI: https://doi.org/10.1038/s41467-019-09145-6, PMID:
30874562

Marreddy RKR, Phelps GA, Churion K, Picker J, Powell R, Cherian PT, Bowling JJ, Stephan
CC, Lee RE, Hurdle JG. 2024. Chemical genetic analysis of enoxolone inhibition of
clostridioides difficile toxin production reveals adenine deaminase and ATP synthase
as antivirulence targets. Journal of Biological Chemistry **300**:107839. DOI:
https://doi.org/10.1016/j.jbc.2024.107839

Menke J, Weber J, Broz K, Kistler HC. 2013. Cellular development associated with induced
mycotoxin synthesis in the filamentous fungus fusarium graminearum. PLOS One
8:e63077. DOI: https://doi.org/10.1371/journal.pone.0063077, PMID: 23667578

Merhej J, Richard-Forget F, Barreau C. 2011. Regulation of trichothecene biosynthesis in
fusarium: recent advances and new insights. Applied Microbiology and
Biotechnology **91**:519–528. DOI: https://doi.org/10.1007/s00253-011-3397-x

Miklos GLG, Yamamoto M-T, Burns RG, Maleszka R. 1997. An essential cell division gene
of drosophila, absent from saccharomyces, encodes an unusual protein with
tubulin-like and myosin-like peptide motifs. Proceedings of the National Academy of
Sciences **94**:5189–5194. DOI: https://doi.org/10.1073/pnas.94.10.5189

Mitani S, Araki S, Takii Y, Ohshima T, Matsuo N, Miyoshi H. 2001. The biochemical mode of
action of the novel selective fungicide cyazofamid: specific inhibition of
mitochondrial complex III in *phythium spinosum*. Pesticide Biochemistry and
Physiology **71**:107–115. DOI: https://doi.org/10.1006/pest.2001.2569

Mrnjavac N, Martin WF. 2025. GTP before ATP: the energy currency at the origin of genes.
Biochimica et Biophysica Acta (BBA) - Bioenergetics **1866**:149514. DOI:
https://doi.org/10.1016/j.bbabio.2024.149514

Mudge AM, Dill-Macky R, Dong Y, Gardiner DM, White RG, Manners JM. 2006. A role for
the mycotoxin deoxynivalenol in stem colonisation during crown rot disease of wheat
caused by *fusarium graminearum* and *fusarium pseudograminearum*. Physiological
and Molecular Plant Pathology **69**:73–85. DOI:
https://doi.org/10.1016/j.pmpp.2007.01.003

Nasca A, Scotton C, Zaharieva I, Neri M, Selvatici R, Magnusson OT, Gal A, Weaver D,
Rossi R, Armaroli A, Pane M, Phadke R, Sarkozy A, Muntoni F, Hughes I, Cecconi A,
Hajnóczky G, Donati A, Mercuri E, Zeviani M, Ferlini A, Ghezzi D. 2017. Recessive
mutations in MSTO1 cause mitochondrial dynamics impairment, leading to myopathy
and ataxia. Human Mutation **38**:970–977. DOI: https://doi.org/10.1002/humu.23262,
PMID: 28544275

Nuwamanya AM, Runo S, Mwangi M. 2022. In-vitro sensitivity of *alternaria solani* isolates
to azoxystrobin and difenoconazole fungicides in Kenya and detection of *cyt b*
mutations associated with azoxystrobin resistance. Crop Protection **158**:106010. DOI:
https://doi.org/10.1016/j.cropro.2022.106010

Pb P, Gc de FS, Gmb L, Plo C, Aln V, Ht F. 2023. Competitive interaction between ATP and
GTP regulates mitochondrial ATP-sensitive potassium channels. Chemico-Biological
Interactions **381**. DOI: https://doi.org/10.1016/j.cbi.2023.110560, PMID: 37244398

Peng Q, Zhao H, Zhao G, Gao X, Miao J, Liu X. 2022. Resistance assessment of
pyraoxystrobin in *magnaporthe oryzae* and the detection of a point mutation in cyt *b*
that confers resistance. Pesticide Biochemistry and Physiology **180**:105006. DOI:
https://doi.org/10.1016/j.pestbp.2021.105006

Rogne P, Rosselin M, Grundström C, Hedberg C, Sauer UH, Wolf-Watz M. 2018. Molecular
mechanism of ATP versus GTP selectivity of adenylate kinase. Proceedings of the
National Academy of Sciences of the United States of America **115**:3012–3017. DOI:
https://doi.org/10.1073/pnas.1721508115, PMID: 29507216

Sarewicz M, Dutka M, Pintscher S, Osyczka A. 2013. Triplet state of the semiquinone-rieske
cluster as an intermediate of electronic bifurcation catalyzed by cytochrome bc1.
Biochemistry **52**:6388–6395. DOI: https://doi.org/10.1021/bi400624m, PMID:
23941428

Schulze SR, Wallrath LL. 2007. Gene regulation by chromatin structure: paradigms
established in drosophila melanogaster. Annual Review of Entomology **52**:171–192.
DOI: https://doi.org/10.1146/annurev.ento.51.110104.151007

Shi D, Wang J, Cao Y, Zhang Z, Li X, Mbadianya JI, Chen C. 2023. Overexpression of
FgPtp3 Is Involved in Fludioxonil Resistance in *Fusarium graminearum* by
Inhibiting the Phosphorylation of FgHog1. Journal of Agricultural and Food
Chemistry **71**:12807–12818. DOI: https://doi.org/10.1021/acs.jafc.3c02663

Song J, Li Y, Zhang Z, Gao X, Li S, Zhang J, Zhou M, Duan Y. 2024. Endoplasmic
reticulum-mitochondrial encounter structure regulates the mitochondrial morphology,
DON biosynthesis and toxisome formation in *fusarium graminearum*.
Microbiological Research **289**:127892. DOI:
https://doi.org/10.1016/j.micres.2024.127892

Sumarah MW. 2022. The deoxynivalenol challenge. Journal of Agricultural and Food
Chemistry **70**:9619–9624. DOI: https://doi.org/10.1021/acs.jafc.2c03690, PMID:
35912482

Tang G, Chen A, Dawood DH, Liang J, Chen Y, Ma Z. 2020. Capping proteins regulate
fungal development, DON□toxisome formation and virulence in *Fusarium*
graminearum. Molecular Plant Pathology **21**:173–187. DOI:
https://doi.org/10.1111/mpp.12887

Tang G, Chen Y, Xu J-R, Kistler HC, Ma Z. 2018. The fungal myosin I is essential for
Fusarium toxisome formation. PLoS pathogens **14**:e1006827. DOI:
https://doi.org/10.1371/journal.ppat.1006827, PMID: 29357387

Vercellino I, Sazanov LA. 2022. The assembly, regulation and function of the mitochondrial
respiratory chain. Nature Reviews. Molecular Cell Biology **23**:141–161. DOI:
https://doi.org/10.1038/s41580-021-00415-0, PMID: 34621061

Vollmer W. 2006. The prokaryotic cytoskeleton: a putative target for inhibitors and antibiotics?
Applied Microbiology and Biotechnology **73**:37–47. DOI:
https://doi.org/10.1007/s00253-006-0586-0

Wang C, Li P, Cong W, Ma N, Zhou M, Hou Y. 2025. The transcription factor FgCreA
modulates ergosterol biosynthesis and sensitivity to DMI fungicides by regulating
transcription of FgCyp51A and FgErg6A in Fusarium graminearum. International
Journal of Biological Macromolecules **284**:137903. DOI:
https://doi.org/10.1016/j.ijbiomac.2024.137903

Wang C, Zhang S, Hou R, Zhao Z, Zheng Q, Xu Q, Zheng D, Wang G, Liu H, Gao X, Ma
J-W, Kistler HC, Kang Z, Xu J-R. 2011. Functional analysis of the kinome of the

wheat scab fungus fusarium graminearum. PLOS Pathogens **7**:e1002460. DOI: https://doi.org/10.1371/journal.ppat.1002460

Wang M, Wu N, Wang H, Liu C, Chen Q, Xu T, Chen Y, Zhao Y, Ma Z. 2024. Overproduction of mycotoxin biosynthetic enzymes triggers fusarium toxisome-shaped structure formation via endoplasmic reticulum remodeling. PLOS Pathogens **20**:e1011913. DOI: https://doi.org/10.1371/journal.ppat.1011913

Wang Y, Shi Y, Li W, Han X, Lin X, Liu D, Lin Y, Shen L. 2024. Knockdown of BRAWNIN minimally affect mitochondrial complex III assembly in human cells. Biochimica Et Biophysica Acta. Molecular Cell Research **1871**:119601. DOI: https://doi.org/10.1016/j.bbamcr.2023.119601, PMID: 37769950

Wanschers BFJ, Szklarczyk R, van den Brand MAM, Jonckheere A, Suijskens J, Smeets R, Rodenburg RJ, Stephan K, Helland IB, Elkamil A, Rootwelt T, Ott M, van den Heuvel L, Nijtmans LG, Huynen MA. 2014. A mutation in the human CBP4 ortholog UQCC3 impairs complex III assembly, activity and cytochrome b stability. Human Molecular Genetics **23**:6356–6365. DOI: https://doi.org/10.1093/hmg/ddu357, PMID: 25008109

Wu L, Yuan Z, Wang P, Mao X, Zhou M, Hou Y. 2022. The plasma membrane H+-ATPase FgPMA1 regulates the development, pathogenicity, and phenamacril sensitivity of Fusarium graminearum by interacting with FgMyo-5 and FgBmh2. Molecular Plant Pathology **23**:489–502. DOI: https://doi.org/10.1111/mpp.13173

Xiao J, Zhou Y, Xie Y, Li T, Su X, He J, Jiang Y, Zhu H, Qu H. 2024. ATP homeostasis and signaling in plants. Plant Communications **5**:100834. DOI: https://doi.org/10.1016/j.xplc.2024.100834, PMID: 38327057

Yi L, Yang M, Waalwijk C, Xu Jin, Xu Jingsheng, Molnár O, Chen W, Feng J, Zhang H. 2023. Dynamics of carbendazim-resistance frequency of pathogens associated with the epidemic of fusarium head blight. Plant Disease **107**:1690–1696. DOI: https://doi.org/10.1094/PDIS-08-22-1998-SR, PMID: 36471466

Yu J-H, Hamari Z, Han K-H, Seo J-A, Reyes-Domínguez Y, Scazzocchio C. 2004. Double-joint PCR: a PCR-based molecular tool for gene manipulations in filamentous fungi. Fungal genetics and biology: FG & B **41**:973–981. DOI: https://doi.org/10.1016/j.fgb.2004.08.001, PMID: 15465386

Zara V, Conte L, Trumpower BL. 2009. Evidence that the assembly of the yeast cytochrome bc1 complex involves the formation of a large core structure in the inner mitochondrial membrane. FEBS Journal **276**:1900–1914. DOI: https://doi.org/10.1111/j.1742-4658.2009.06916.x, PMID: 19236481

Zara V, Conte L, Trumpower BL. 2007. Identification and characterization of cytochrome bc(1) subcomplexes in mitochondria from yeast with single and double deletions of genes encoding cytochrome bc(1) subunits. FEBS Journal **274**:4526–4539. DOI: https://doi.org/10.1111/j.1742-4658.2007.05982.x, PMID: 17680808

Zhan J, Zeher A, Huang R, Tang WK, Jenkins LM, Xia D. 2024. Conformations of Bcs1L undergoing ATP hydrolysis suggest a concerted translocation mechanism for folded iron-sulfur protein substrate. Nature Communications **15**:4655. DOI: https://doi.org/10.1038/s41467-024-49029-y, PMID: 38821922

894   Zhang Y, Ouyang B, Zhang W, Guang C, Xu W, Mu W. 2024. Deoxynivalenol: occurrence,
895   toxicity, and degradation. Food Control **155**:110027. DOI:
896   https://doi.org/10.1016/j.foodcont.2023.110027

897   Zhao H, Tao X, Song W, Xu H, Li M, Cai Y, Wang J, Duan Y, Zhou M. 2022. Mechanism of
898   fusarium graminearum resistance to ergosterol biosynthesis inhibitors: G443S
899   substitution of the drug target FgCYP51A. Journal of Agricultural and Food
900   Chemistry **70**:1788–1798. DOI: https://doi.org/10.1021/acs.jafc.1c07543

901   Zhao Y, Chi M, Sun H, Qian H, Yang J, Huang J. 2021. The FgCYP51B Y123H Mutation
902   Confers Reduced Sensitivity to Prochloraz and Is Important for Conidiation and
903   Ascospore Development in Fusarium graminearum. Phytopathology **111**:1420–1427.
904   DOI: https://doi.org/10.1094/PHYTO-09-20-0431-R, PMID: 33399013

905   Zheng H, Li L, Miao P, Wu C, Chen X, Yuan M, Fang T, Norvienyeku J, Li G, Zheng W,
906   Wang Z, Zhou J. 2018. FgSec2A, a guanine nucleotide exchange factor of FgRab8, is
907   important for polarized growth, pathogenicity and deoxynivalenol production in
908   Fusarium graminearum. Environmental Microbiology **20**:3378–3392. DOI:
909   https://doi.org/10.1111/1462-2920.14373

910   Zhou F, Zhou X, Yan Jiao N, Han A, Zhou H, Chen Z, Li W, Liu R. 2024. Baseline
911   tebuconazole sensitivity and potential resistant risk in Fusarium Graminearum. BMC
912   plant biology **24**:789. DOI: https://doi.org/10.1186/s12870-024-05206-1, PMID:
913   39164633

914   Zhou Z, Duan Y, Zhou M. 2020. Carbendazim-resistance associated β2 -tubulin substitutions
915   increase deoxynivalenol biosynthesis by reducing the interaction between β2 -tubulin
916   and IDH3 in fusarium graminearum. Environmental Microbiology **22**:598–614. DOI:
917   https://doi.org/10.1111/1462-2920.14874, PMID: 31760682

FIG. 1.

Identification and localization of FgDML1 in *Fusarium graminearum*. (A) The phylogenetic tree was constructed based on the amino acid sequences of DML1 homologs from 10 species using the maximum likelihood method in Mega X. Bootstrap values from 1000 replications were displayed on the branches. (B) FgDML1 was localized to the mitochondria.

FIG. 2.

Effects of ΔFgDML1 on asexual and sexual development. (A) Mycelial growth rate.

PH-1, ΔFgDML1, ΔFgDML1-C were cultured for 3 days on PDA, CM, MM, and V8

solid media at 25°C. (B) Mycelial morphology. The mycelial morphology of PH-1,

ΔFgDML1, and ΔFgDML1-C was examined on WA plates. Scale bar = 72 μm. (C)

Conidial length. Conidia of ΔFgDML1 were significantly longer in length compared

to PH-1 and ΔFgDML1-C. Conidia were germinated in MBL medium, stained with

Calcofluor White (CFW), and observed using an Olympus IX-71 microscope. (D)

Sexual reproduction. PH-1 and ΔFgDML1-C produced perithecia and asci containing

ascospores after 7 or 14 days of incubation on CA medium, while the ΔFgDML1

failed to produce these structures under the same conditions. Perithecia and

ascospores were observed using a Nikon SMZ25 fluorescent stereo microscope and

an Olympus IX-71 inverted fluorescence microscope. Top image: perithecium, scale

bar = 1 mm; bottom image: ascospores, scale bar = 50 μm. (E) Proportion of conidial septa. The proportion of conidial septa was assessed in the tested strains. (F) Conidial production. Conidia were harvested after 4 days of incubation in MBL medium, and their concentration was quantified using a hemocytometer. (G) Germination rate. Conidial germination was observed by incubating 300 conidia for 2 or 8 hours on WA plates. Germination rates were compared, with different letters indicating significant differences based on ANOVA followed by the LSD test (p= 0.05).

FIG. 3.

FgDML1 is required for both the virulence of *Fusarium graminearum*. (A) Lesion lengths on wheat coleoptiles caused by different strains. These images were taken at 14 dpi. (B) Dot plot showing the lesion lengths on wheat coleoptiles at 14 dpi tested with different strains. (C) Disease symptoms on wheat leaves inoculated with PH-1, ΔFgDML, and ΔFgDML-C. Images were taken at 5 dpi. (D) Disease area on wheat leaves inoculated with different strains at 5 dpi. Bars with the same letter indicate no significant difference according to a significant difference (LSD) test at $p = 0.05$.

FIG. 4.

FgDML1 is required for the biosynthesis of DON toxin in *Fusarium graminearum*. (A) ΔFgDML1 exhibited decreased DON content. All strains were incubated in trichothecene biosynthesis induction (TBI) medium for 7 days. (B) Relative mRNA expression levels of *FgTRI5*, and *FgTRI6* in the tested strains. Mycelia were harvested after 2 days of cultivation in TBI medium, and mRNA were extracted for quantification using the $2^{-\Delta\Delta CT}$ method. *FgGapdh* was used as the reference gene. (C) FgDML1 impaired the formation of DON toxisomes, as visualized by Tri1-GFP labeling of toxisomes. Strains were cultured in TBI for 36 hours. Bar = 10 μm (D) ΔFgDML1 exhibited reduced expression levels of Tri1-GFP. Strains were cultured in TBI for 36 hours, and mycelia were harvested for western blot analysis. Bars with the

same letter indicate no significant difference according to a significant difference

(LSD) test at $p = 0.05$.

FIG. 5.

FgDML1 modulates mitochondrial morphology via interaction with FgDnm1 and negatively regulates Acetyl-CoA and ATP synthesis. (A) Effects of FgDML1 on mitochondrial morphology. Mycelial plugs of the wild-type strain PH-1, the ΔFgDML1 deletion mutant, and the complemented mutant ΔFgDML1-C were cultured in YEPD medium for 36 hours. Mitochondria were stained using a mitochondria-specific fluorescent dye and observed under a confocal microscope. Additionally, conidial germlings were cultured in MBL medium and subjected to the same staining and imaging procedures. Scale bar = 5 μm. (B) Ultrastructural analysis of mitochondria by transmission electron microscopy (TEM). Mitochondrial ultrastructure was further examined in PH-1, ΔFgDML1, and ΔFgDML1-C using TEM. Mitochondria are indicated by red arrows. Scale bar = 200 nm. (C) FgDML1 Deletion Reduces Acetyl-CoA Levels. Mycelial cultures of PH-1, ΔFgDML1, and ΔFgDML1-C were incubated in YEPD medium for 36 hours. Acetyl-CoA levels were

quantified using a commercial assay kit, following the instructions provided by the commercial assay kit. (D) FgDML1 deletion reduces ATP levels. The ATP content of PH-1, ΔFgDML1, and ΔFgDML1-C was measured following the same culture conditions as in (C), using a commercial ATP assay kit. (E) Co-immunoprecipitation (Co-IP) confirms interaction between FgDML1 and FgDnm1. Total protein extracts from strains expressing tagged versions of FgDML1 and FgDnm1 were separated by SDS-PAGE and subjected to immunoblot analysis. Co-IP was performed using ChromoTek GFP-Trap® magnetic agarose to pull down interacting proteins. Immunodetection was carried out using polyclonal anti-Flag and monoclonal anti-GFP antibodies. Monoclonal anti-GAPDH antibody was used as a reference for protein samples. Bars with the same letter indicate no significant difference according to a significant difference (LSD) test at $p = 0.05$.

FIG. 6.

FgDML1 modulates the sensitivity of *Fusarium graminearum* to cyazofamid. (A) ΔFgDML1 exhibits reduced sensitivity to cyazofamid. PH-1, ΔFgDML1, and ΔFgDML1-C were cultured on AEA medium modified with cyazofamid, with 50 μg/mL SHAM included to inhibit the alternative oxidase pathway. (B) Deletion of *FgDML1* did not affect *F. graminearum* sensitivity to pyraclostrobin. The tested strains were incubated on AEA medium containing pyraclostrobin and 50 μg/mL SHAM to assess their response to the fungicide. (C) FgDML1 deletion reduces complex III enzymatic activity. Mycelial cultures of PH-1, ΔFgDML1, and ΔFgDML1-C were grown in YEPD for 36 hours, then treated with 30 μg/mL cyazofamid, 1.5 μg/mL pyraclostrobin, and 50 μg/mL SHAM for an additional 12 hours. The enzymatic activity of complex III was quantified using a commercial assay kit. (D) mRNA expression levels of complex III subunits and assembly factors in PH-1, ΔFgDML1, and ΔFgDML1-C. PH-1, ΔFgDML1, and ΔFgDML1-C were incubated in liquid AEA medium for 36 hours, followed by treatment with 30 μg/mL

cyazofamid and 50 μg/mL SHAM. The relative mRNA expression levels of complex III subunits and assembly factors were quantified using the $2^{-\Delta\Delta CT}$ method. "T" denotes cyazofamid treatment. Bars with the same letter indicate no significant difference according to a significant difference (LSD) test at $p = 0.05$. (E) Overexpression of *FgQCR2*, *FgQCR7*, and *FgQCR8*, respectively, in the PH-1 background resulted in reduced sensitivity to cyazofamid in the overexpression mutants. (F) Protein expression levels of FgQCR2, FgQCR7, and FgQCR8 in PH-1 and ΔFgDML1. Strains were cultured in AEA Liquid Medium for 36 h, and mycelia were harvested for western blot analysis.

FIG. 7.

Response of the strains to different stresses. (A) Reduced sensitivity to osmotic stress in ΔFgDML1. PH-1, ΔFgDML1 and ΔFgDML1-C were subjected to osmotic stress using NaCl, KCl, and Sorbitol to evaluate their sensitivity. (B) Statistical analysis of growth inhibition under osmotic stress. Growth inhibition rates of all strains under NaCl, KCl, and Sorbitol stress were statistically analyzed. (C) Sensitivity to cell wall

and membrane damaging agents. The tested strains were exposed to cell wall-damaging agent Congo Red (CR) and cell membrane-damaging agent Sodium dodecyl sulfate (SDS) to assess their sensitivity. (D) Statistical analysis of growth inhibition under CR and SDS stress. Growth inhibition rates of all strains under CR and SDS stress were statistically analyzed. (E) Sensitivity to temperature stresses. The sensitivity of the tested strains to different temperature stresses was evaluated. (F) Statistical analysis of growth inhibition under temperature stress. Growth inhibition rates of all strains under various temperature stresses were statistically analyzed. Significant differences were determined using the LSD test at $p = 0.05$. Bars with the same letter indicate no significant difference.

FIG. 8.

Model of FgDML1 regulation of *Fusarium graminearum* sensitivity to DON toxin synthesis and cyazofamid fungicide. FgDML1 interacts with FgDnm1 to modulate mitochondrial morphology, thereby positively regulating acetyl-CoA and ATP levels, which influence toxisome formation and ultimately affect DON production. Loss of FgDML1 reduces complex III enzymatic activity, leading to the upregulation of the assembly factors *FgQCR2*, *FgQCR8*, and *FgQCR9*. This alteration in assembly factor expression changes the conformation of the Qi site, leading to reduced sensitivity to cyazofamid.