Mitochondrial protein FgDML1 regulates DON toxin biosynthesis and

cyazofamid sensitivity in *Fusarium graminearum* by affecting mitochondrial

homeostasis

Chenguang Wang[a †], Xuewei Mao[b †], Weiwei Cong[a], Lin Yang[a], Yiping Hou[a] *

[a]College of Plant Protection, Nanjing Agricultural University, Nanjing, Jiangsu, China

[b]College of Plant Protection, Henan Agricultural University, Zhengzhou, Henan,

China

[†]These authors contributed equally to this work

*Corresponding author：Yiping Hou; E-mail address: houyiping@njau.edu.cn

Abstract

Fusarium graminearum is a global pathogen responsible for Fusarium head blight (FHB) in wheat, causing substantial yield losses and producing the mycotoxin deoxynivalenol (DON), which poses a threat to both human and animal health. *Drosophila melanogaster* Misato-Like protein (DML1) plays a critical role in regulating mitochondrial function, yet its function in filamentous fungi remains unexplored. In this study, we characterized FgDML1 in *F. graminearum*. FgDML1 interacts with the mitochondrial fission and fusion protein FgDnm1 to maintain mitochondrial stability, thereby positively regulating acetyl-CoA levels and ATP synthesis, which influences toxisome formation and ultimately affects DON toxin biosynthesis. Additionally, FgDML1 is involved in the regulation of toxin biosynthetic enzyme expression. In the ΔFgDML1 mutant, Complex III enzyme activity decreased, overexpression of complex III assembly factors *FgQCR2*, *FgQCR8*, and *FgQCR9* may induce conformational changes in the Qi-site protein, specifically altering the sensitivity of *F. graminearum* to respiratory inhibitor cyazofamid not Qo-site inhibitor pyraclostrobin and other fungicides. Furthermore, the loss of FgDML1 leads to defects in nutrient utilization, as well as in asexual and sexual reproduction, and pathogenicity. In conclusion, this study identifies a novel regulatory role for FgDML1 in DON toxin biosynthesis and cyazofamid sensitivity in *F. graminearum*. Our study provides a theoretical framework for understanding DON biosynthesis regulation in *F. graminearum* and identifies potential molecular targets for FHB control.

32 **Keywords:** *Fusarium graminearum*, DON production, FgDML1, Cyazofamid

33 Pathogenicity

Introduction

Fusarium head blight (FHB), caused by *Fusarium graminearum*, is a devastating

fungal disease that affects cereal crops, leading to significant yield losses in

wheat-growing regions worldwide(Dean et al., 2012). This disease is particularly

concerning due to the production of various mycotoxins, including trichothecenes and

zearalenone, during fungal infection(Hai et al., 2023). Among these, deoxynivalenol

(DON), a type of trichothecene, is the most commonly detected mycotoxin(Merhej et

al., 2011a). The DON toxin poses a serious threat to both human and animal health,

significantly compromising immune and reproductive systems(EFSA Panel on

Contaminants in the Food Chain (CONTAM) et al., 2017). Chronic exposure to DON

can lead to anorexia, neurological disorders, and immune dysregulation, while acute

high-dose exposure may result in vomiting, diarrhea, neurological disturbances, and

even miscarriage or stillbirth(Sumarah, 2022; Zhang et al., 2024). Studies have

demonstrated a positive correlation between elevated DON levels and the prevalence of

FHB in infected crops(Bönnighausen et al., 2019; Ilgen et al., 2009; Mudge et al.,

2006). As FHB continues to spread and its severity increases, DON contamination is

emerging as a critical global concern(Duan et al., 2020; Liu et al., 2016). This

highlights the need for effective strategies to manage and mitigate the impact of both

the disease and the associated mycotoxin contamination.

The absence of resistant cultivars necessitates reliance on chemical methods as the

primary approach for controlling FHB(Liu et al., 2019). However, the inappropriate

application of fungicides has led to the emergence of resistance, particularly against

carbendazim, tebuconazole, and pyraclostrobin(Andrade et al., 2022; Yi et al., 2023;

Zhou et al., 2024). Notably, carbendazim has been demonstrated to stimulate the

biosynthesis of deoxynivalenol (DON) toxin, especially in fungicide-resistant

populations of *F. graminearum*(Li et al., 2019; Zhou et al., 2020). Similarly,

pyraclostrobin significantly promotes DON biosynthesis, with no selectivity between

resistant and sensitive strains. Although tebuconazole has not been directly implicated

in inducing DON production, it is classified as a fungicide with a moderate risk of

resistance development(Duan et al., 2020). Field observations have revealed diverse

resistance mechanisms to tebuconazole, complicating management efforts(Liu et al.,

2011; Zhao et al., 2021). The potential evolution of resistance to tebuconazole is

particularly concerning, as it could undermine its efficacy and exacerbate the challenge

of controlling DON contamination. There is an urgent need for a fungicide that has not

yet been used in *F. graminearum* to address the DON toxin contamination caused by

fungicide resistance. Cyazofamid is a mitochondrial respiratory chain complex III

inhibitor with demonstrated efficacy in controlling fungal diseases. It specifically

binds to the Qi site of complex III, blocking the reduction of ubiquinone and

inhibiting ATP synthesis, thereby exerting its fungicidal effect(Li et al., 2014; Mitani

et al., 2001). However, it remains unclear whether other proteins within the intricate

regulatory network of the respiratory chain play a role in modulating fungal

sensitivity to cyazofamid. Further investigations are needed to elucidate the potential

involvement of these regulatory components.

The synthesis of deoxynivalenol (DON) toxin is primarily regulated by genes

involved in trichothecene biosynthesis, known as *FgTRIs*. Many of these *FgTRI* genes

have been well characterized(Merhej et al., 2011b). The initial step in DON synthesis

is catalyzed by the trichothecene synthase *FgTri5*, which facilitates the cyclization of

farnesyl pyrophosphate (FPP) to form trichodiene (TDN)(Kimura et al., 2007).

However, before this catalytic process can occur, the organism must first obtain

sufficient ATP from the mitochondria. ATP is critical not only for the basic cellular

functions but also for driving several key reactions in the isoprenoid biosynthesis

pathway. Specifically, ATP is required for the enzymatic conversion of precursors,

leading to the production of FPP, a vital intermediate in the trichothecene biosynthesis

pathway. The availability of ATP therefore directly impacts the efficient synthesis of

FPP, which, in turn, regulates the downstream production of DON toxin. Without an

adequate supply of ATP, these biosynthetic pathways cannot proceed efficiently, thus

hindering the synthesis of both the precursor molecules and the final toxin

product(Alexander et al., 2009).

Drosophila melanogaster Misato-Like protein (DML1) encodes a Misato-like

protein in *Saccharomyces cerevisiae*, an essential factor involved in mitochondrial

DNA (mtDNA) inheritance. In addition to its role in mtDNA maintenance, DML is

implicated in mitochondrial compartmentalization and chromosome segregation. The

protein shares structural features with members of the GTPase family, suggesting

potential involvement in GTP-dependent processes(Gurvitz et al., 2002). In *Homo*

sapiens, the DML protein is localized to the mitochondria, where silencing of DML

results in a transition from a networked mitochondrial morphology to a fragmented

form. This indicates that DML plays a critical role in mitochondrial distribution,

morphological fusion, and overall mitochondrial dynamics(Kimura and Okano,

2007a). In prokaryotes, the homologous protein FtsZ, which shares functional

similarities with DML, has been identified as a target for antimicrobial agents(Huang

et al., 2007; Vollmer, 2006). However, despite these insights from studies in the

aforementioned organisms, the exact role of DML in filamentous fungi remains

undetermined.

This study systematically characterized the functional roles of *Drosophila*

melanogaster Misato-Like protein (FgDML1) in *F. graminearum*, including mycelial

growth, asexual development, and sexual reproduction. More importantly, we found

that FgDML1 reduces DON toxin synthesis by decreasing the enzyme activity of

complex III, reducing ATP synthesis, and lowering acetyl-CoA accumulation.

Furthermore, the deletion of FgDML1 results in compensatory upregulation of

complex III assembly factors, which may alter the conformation of the Qi site protein

and subsequently modulates sensitivity to cyazofamid.

Results

Identification, deletion, and complementation of FgDML1.

The *Drosophila melanogaster* Misato-Like (DML1) protein from *S. cerevisiae*

was used as a query for a BLAST search against the Fusarium genome database,

resulting in the identification of the putative DML1 gene FgDML1 (FGSG_05390) in

F. graminearum. The FgDML1 gene is 1455 bp in length, contains three introns, and

encodes a 484-amino acid protein with 23.88% homology to its counterpart in *S.*

cerevisiae. Phylogenetic analysis was performed using DML1 sequences from other

species obtained via NCBI, and a phylogenetic tree was constructed (Fig. 1A).

Structural analysis revealed the presence of the Misat_Tub_SegII and Tubulin_3

domains. Furthermore, subcellular localization studies confirmed that FgDML1

localizes to mitochondria, as demonstrated by colocalization with a

mitochondria-specific dye (Fig. 1B).

To investigate the function of FgDML1, ΔFgDML1 deletion mutant was

constructed and validated via PCR and Southern blot analysis (Fig. S2), confirming

the mutant as a single-copy knockout. To ensure that the observed phenotypes were

specifically attributable to the deletion of FgDML1, an in situ complemented mutant,

ΔFgDML1-C, was generated to restore FgDML1.

FgDML1 is essential for virulence.

To investigate the role of FgDML1 in virulence, pathogenicity assays were

performed on wheat coleoptiles and wheat leaves. The results revealed that ΔFgDML1

caused only restricted infection at the inoculation site on wheat coleoptiles and failed

to induce systemic infection, in contrast to PH-1 and ΔFgDML1-C, which exhibited

extensive colonization (Fig. 2A, B). Similarly, ΔFgDML1 displayed a near-complete

loss of virulence on wheat leaves (Fig. 2C, D). These findings demonstrate that

FgDML1 is a critical positive regulator of virulence in *F. graminearum*.

FgDML1 is indispensable for DON toxin synthesis.

To evaluate whether the deletion of FgDML1 influences DON production, DON

levels were measured under toxin-inducing conditions. The results revealed that DON

production in ΔFgDML1 was reduced by nearly 80% compared to PH-1 and

significantly lower than that of both PH-1 and ΔFgDML1-C (Fig. 3A). To further

elucidate the regulatory relationship between FgDML1 and DON biosynthesis, the

transcriptional levels of *FgTri5* and *FgTri6* were analyzed and found to be

significantly downregulated in ΔFgDML1 (Fig. 3B). Moreover, examination of

toxisomes in the mutant demonstrated that their formation was severely impaired,

with almost no detectable toxisomes in ΔFgDML1(Fig. 3C). Western blot analysis of

the DON toxin marker protein FgTri1 further conirmed its markedly reduced levels in

ΔFgDML1 compared to PH-1 (Fig. 3D). These findings suggest that FgDML1 is

essential for regulating DON production in *F. graminearum*.

The deletion of FgDML1 results in reduced synthesis of both ATP and acetyl-CoA.

The results showed that ATP levels in ΔFgDML1 were significantly lower

compared to PH-1 and ΔFgDML1-C (Fig. 4D). Previous studies have established a

positive correlation between ATP levels and acetyl-CoA. Consistent with this,

acetyl-CoA levels in ΔFgDML1 were also significantly reduced compared to the

controls (Fig. 4C). In conclusion, the deletion of FgDML1 leads to a reduction in both

the key energy molecule ATP and the DON precursor acetyl-CoA.

FgDML1 regulates mitochondrial morphology by interacting with FgDnm1.

To assess whether the loss of FgDML1 induces mitochondrial alterations,

mitochondria-specific dyes were utilized, revealing fragmented mitochondria in the

ΔFgDML1 mutant, whereas PH-1 and ΔFgDML1-C displayed a typical network-like

mitochondrial distribution (Fig. 4A). Further analysis via TEM identified damaged

mitochondrial structures in the ΔFgDML1 mutant, in contrast to the well-preserved

mitochondrial morphology observed in the control strains (Fig. 4B). Given that

FgDnm1 is known to be a critical protein for mitochondrial fission, the interaction

between FgDML1 and FgDnm1 was verified through co-immunoprecipitation (CoIP)

assays (Fig. 4E). These findings indicate that FgDML1 plays a pivotal role in

maintaining mitochondrial morphology by interacting with FgDnm1.

The deletion of FgDML1 reduces Complex III activity, induces compensatory

upregulation of *FgQCR2*, *FgQCR8*, *FgQCR9*.

Deletion of FgDML1 reduced *F. graminearum* sensitivity to the Qi-site fungicide

cyazofamid, while having no effect on sensitivity to the Qo-site inhibitors

pyraclostrobin and trifloxystrobin, nor to other respiratory inhibitors such as

pydiflumetofen, fluxapyroxad, and fluopyram (Fig. 5A, B) (Fig. S3). Measurement of

Complex III activity in ΔFgDML1 showed a 38.28% reduction in enzyme activity

compared to PH-1 (Fig. 5C). To further investigate the reduced sensitivity to

cyazofamid, transcription levels of the three subunits (*FgCytb, FgCytC1, FgISP*) and

five assembly factors (*FgQCR2, FgQCR6, FgQCR7, FgQCR8, FgQCR9*) of Complex

III were analyzed. The results revealed that, upon Cyazofamid treatment, the

expression of *FgQCR2*, *FgQCR7*, and *FgQCR8* was significantly upregulated in Δ

FgDML1 (Fig. 5D).

Overexpression of *FgQCR2*, *FgQCR8*, and *FgQCR9* alters the conformation of the

QI site, resulting in reduced sensitivity to cyazofamid.

Since Complex III is involved in the action of both cyazofamid (targeting the QI

site) and pyraclostrobin (targeting the QO site) fungicides, the sensitivity of

ΔFgDML1 to trifloxystrobin was investigated. The results showed that the deletion of

FgDML1 did not alter sensitivity to a pyraclostrobin (Fig. 5B). ΔFgDML1 was

subsequently treated with pyraclostrobin or cyazofamid, and Complex III enzyme

activity was measured. After pyraclostrobin treatment, no significant change in

enzyme activity was observed in ΔFgDML1 compared to PH-1 and ΔFgDML1-C (Fig.

5C). However, when treated with cyazofamid, a significant increase in Complex III

enzyme activity was observed in ΔFgDML1, while the control groups maintained

relatively low enzyme activity (Fig. 5C). These findings, combined with the

upregulation of assembly factors, suggest that the elevated expression of assembly

factors alters the protein conformation at the QI site, modulating sensitivity to

cyazofamid.

FgDML1 regulates vegetative growth.

To assess the role of FgDML1 in vegetative growth, PH-1, ΔFgDML1, and

ΔFgDML1-C were cultured on various media, including PDA, CM, MM, and V8, to

measure growth rates. The ΔFgDML1 mutant exhibited significantly reduced growth

rates on all tested media compared to PH-1 and the complemented mutant

ΔFgDML1-C (Fig. 6A).   Moreover, ΔFgDML1 displayed pronounced hyphal

curvature, which was not observed in PH-1 or ΔFgDML1-C (Fig. 6B). These findings

indicate that FgDML1 is essential for normal vegetative growth in *F. graminearum*.

FgDML1 plays a critical role in both asexual and sexual reproduction.

To examine the role of FgDML1 in asexual reproduction, a series of experiments

were conducted on conidia. Morphological analysis revealed that conidia from the

ΔFgDML1 mutant were thinner and longer than those from the PH-1 and

ΔFgDML1-C (Fig. 6C). Furthermore, statistical analysis of septum number indicated

a significant increase in conidia with ≥7 septa in the ΔFgDML1 compared to the

controls (Fig. 6E). Conidial production was significantly reduced in ΔFgDML1, with

fewer spores produced compared to PH-1 and ΔFgDML1-C (Fig. 6F). The

germination rate of ΔFgDML1 conidia was also significantly lower than that of the

controls (Fig. 6G). Regarding sexual reproduction, ΔFgDML1 failed to produce

perithecia or ascospores on both the 7th and 14th days of incubation (Fig. 6D).

FgDML1 regulates sensitivity to various stresses.

F. graminearum encounters various stresses during infection plant, which

significantly influence its ability to infect. To investigate its response to different

stress conditions, the sensitivity of ΔFgDML1 to various environmental factors was

assessed, including osmotic stress (NaCl, KCl, sorbitol), cell wall stress (Congo red),

membrane stress (SDS), as well as high temperature (30°C) and low temperature

(15°C) stresses. The results revealed that ΔFgDML1 displayed significantly reduced

sensitivity to osmotic stress compared to PH-1 and ΔFgDML1-C (Fig. 7A, B). No

significant difference was observed in sensitivity to membrane stress, while a notable

change in sensitivity to cell wall stress was observed (Fig. 7C, D). Additionally,

differences in response to both high and low temperature stress were noted in

ΔFgDML1 when compared to the control strains PH-1 and ΔFgDML1-C (Fig. 7E, F).

Discussion

Misato Protein was originally identified in *Drosophila melanogaster*, where it

was reported to play essential roles in mitochondrial DNA inheritance, cell division,

and energy metabolism(Miklos et al., 1997). However, its functional characterization

in filamentous fungi remains largely unexplored. In this study, we identified and

analyzed the homolog FgDML1 in the phytopathogenic fungus *F. graminearum*. The

FgDML1 protein contains conserved Misato_Tub_SegII and Tubulin_3 domains, with

high homology observed among species within the *Fusarium* genus but significant

genetic divergence from other taxa (Fig. 1A). In *S. cerevisiae*, DML1 is a critical

regulator of mitochondrial fission and fusion dynamics, maintaining intracellular

energy   homeostasis   by   modulating   mitochondrial   morphology   and

distribution(Gurvitz et al., 2002). Our study demonstrated that FgDML1 localized to

mitochondria in *F. graminearum* (Fig. 1B). Deletion mutants of FgDML1 exhibited

phenotypes   indicative   of   mitochondrial   dysfunction,   including   fragmented

mitochondrial morphology (Fig. 4A). TEM further confirmed structural and dynamic

damage to mitochondria in the ΔFgDML1 (Fig. 4B). Structural analysis suggests that

DML1 shares similarities with GTPase family proteins, indicating potential GTPase

activity(Gurvitz et al., 2002). GTPases primarily hydrolyze GTP, but their activity is

intricately linked to ATP production and utilization. These enzymes regulate critical

cellular processes such as energy metabolism, signal transduction, protein synthesis,

and cytoskeletal remodeling, all of which influence ATP levels(Mrnjavac and Martin,

2025; Pb et al., 2023; Rogne et al., 2018).   In agreement with this, we observed a

significant reduction in ATP content in the FgDML1 deletion mutant, underscoring its

role in mitochondrial energy metabolism (Fig. 4D). In *Homo sapiens*, DML1 also

localizes to mitochondria, where its silencing results in fragmented and aggregated

mitochondria with reduced network continuity and fusion activity(Kimura and Okano,

2007b). Our findings indicate that FgDML1 plays a conserved role in maintaining

mitochondrial integrity and energy homeostasis. Notably, FgDML1 was found to

interact with FgDnm1(Fig. 4E), FgDnm1 is a key dynamin-related protein mediating

mitochondrial fission(Griffin et al., 2005; Kang et al., 2023), suggesting that FgDML1

may form a complex with FgDnm1 to regulate mitochondrial fission and fusion

processes. To our knowledge, this is the first report documenting an interaction

between DML1 and Dnm in any fungal species, including model organisms such as *S.*

cerevisiae. This novel finding provides new insights into the molecular mechanisms

underlying mitochondrial dynamics in filamentous fungi.

The silencing of ATP synthase subunit α (AtpA) has been shown to reduce ATP

synthesis and inhibit the biosynthesis of TcdA and TcdB toxins(Marreddy et al., 2024).

In plants, the production of secondary metabolites often demands substantial energy,

with ATP serving as the primary energy carrier within cells. When ATP levels are

depleted, plants may prioritize growth and development by reducing the synthesis of

certain secondary metabolites, including toxins(Xiao et al., 2024). Similarly, in *F.*

graminearum, deoxynivalenol (DON) toxin, a highly harmful mycotoxin and

secondary metabolite, requires sufficient energy for its biosynthesis. In our study, the

ATP content in the FgDML1 deletion mutant was significantly lower, and DON

production was markedly reduced, compared to the wild-type PH-1 and the

complementary mutant ΔFgDML1-C (Figure 3A). Reduced ATP levels disrupt the

formation of toxisomes, which are critical for DON toxin biosynthesis(Tang et al.,

2018), and the ΔFgDML1 strain is unable to form functional toxisomes (Fig. 3C).

Western blot analysis also confirmed a significant reduction in the protein expression

of FgTri1 (Fig. 3D). DON production is predominantly regulated by Tri genes, with

Tri5 and *Tri6* serving as critical components in the biosynthetic pathway(Kimura et al.,

2001). Tri5 encodes trichodiene synthase, the key enzyme catalyzing the first step of

DON biosynthesis(Huang et al., 2024; Jiang et al., 2016). In our study, the expression

levels of *FgTri5* and *FgTri6* were significantly downregulated in the FgDML1

deletion mutant (Fig. 3B). To our knowledge, no research has yet elucidated a

regulatory role for DML in gene expression in either *S. cerevisiae* or *Drosophila*

melanogaster. However, given its established functions in cell division and

microtubule organization, FgDML1 might indirectly influence gene expression by

modulating chromatin structure or cell cycle progression(Schulze and Wallrath, 2007).

Investigating this hypothesis will be a key focus of our future research. Given this

substantial reduction in DON production, we investigated whether the precursor for

DON biosynthesis, acetyl-CoA, was also affected. Results revealed a significant

decrease in acetyl-CoA levels in the ΔFgDML1 mutant (Fig. 4C). In conclusion,

reduced ATP levels impair toxin body formation, leading to lower toxin production.

Additionally, decreased acetyl-CoA disrupts the availability of essential precursors for

DON toxin synthesis, further decreasing its production. The combined effects of these

factors result in a significant reduction in DON toxin biosynthesis.

F. graminearum is a highly destructive pathogen that poses a significant threat to

wheat production worldwide. Cyazofamid is an effective fungicide commonly used to

control plant pathogenic fungi. Cyazofamid specifically targets the Qi site of complex

III in the mitochondrial respiratory chain(Bolgunas et al., 2006; Mitani et al., 2001).

Complex III, also known as the cytochrome bc1 complex, plays a critical role in the

electron transport chain by transferring electrons from ubiquinone to cytochrome c

and pumping protons from the mitochondrial matrix into the intermembrane

space(Fernández-Ortuño et al., 2008; Sarewicz et al., 2013). This complex consists of

several core subunits, including cytochrome b, cytochrome c1, and iron-sulfur

proteins, as well as multiple assembly factors. It contains two quinone binding sites:

the Qo site (quinone oxidation site) and the Qi site (quinone reduction site)(Vercellino

and Sazanov, 2022; Zara et al., 2009). In our study, we observed that the deletion of

FgDML1 significantly reduced the sensitivity of *F. graminearum* to cyazofamid (Fig.

5A), while its sensitivity to pyraclostrobin remained unaffected (Fig. 5B). In the

ΔFgDML1 mutant, we detected a decrease in complex III enzyme activity (Fig. 5C)

and noted the upregulation of assembly factors *FgQCR2*, *FgQCR7*, and *FgQCR8*

following cyazofamid treatment (Fig. 5D). Further enzyme activity assays revealed

that after pyraclostrobin treatment, ΔFgDML1 exhibited a 73.06% reduction in

complex III activity, compared to an 80.38% decrease in the wild-type strain PH-1

(Fig. 5C). Interestingly, after cyazofamid treatment, the decrease in enzyme activity in

ΔFgDML1 was only 36.31%, while PH-1 showed a dramatic 83.46% reduction in

complex III activity. *FgQCR2*, *FgQCR7*, and *FgQCR8* are critical assembly factors

that play essential roles in the proper assembly of complex III(Zhan et al., 2024).

These factors stabilize the core subunit structure, promoting correct folding and

functional operation of complex III, particularly during the assembly of the Qi site.

This ensures the proper formation and function of the Qi site, thus supporting the

normal operation of the mitochondrial respiratory chain(Y. Wang et al., 2024; Zara et

al., 2007). Given the role of FgDML1 in mitochondrial homeostasis, we further

evaluated the sensitivity of the ΔFgDML1 mutant to other commonly used respiratory

chain inhibitors, including pydiflumetofen, fluxapyroxad, fluopyram, and

trifloxystrobin. The results indicated that FgDML1 deletion had no impact on the

sensitivity of *F. graminearum* to these fungicides (Fig. S3). In conclusion, our

findings suggest that the overexpression of assembly factors *FgQCR2*, *FgQCR7*, and

FgQCR8 in ΔFgDML1 alters the conformation of the Qi site, which specifically

modulates the sensitivity of *F. graminearum* to cyazofamid.

In *H. sapiens*, mutations or deletions in DML1 are associated with mitochondrial

diseases characterized by early-onset myopathy and cerebellar ataxia(Chen et al.,

2022; Nasca et al., 2017). In our study, the deletion of FgDML1 in *F. graminearum*

resulted in significant defects in vegetative growth, including a markedly reduced

growth rate (Fig. 6A) and an aberrant, excessively curved hyphal morphology (Fig.

6B). Notably, the loss of FgDML1 rendered *F. graminearum* incapable of completing

sexual reproduction (Fig. 6D), thereby potentially reducing its primary inoculum

sources. Additionally, the deletion of FgDML1 impaired asexual reproduction and

pathogenicity (Fig. 2) (Fig. 6), critical factors for fungal survival and host infection.

Furthermore, the ΔFgDML1 mutant exhibited altered responses to various stress

conditions (Fig. 7), suggesting a broader role for FgDML1 in regulating fungal stress

tolerance.

In the present study, FgDML1 is localized to the mitochondria and interacted

with FgDnm1, a key regulator of mitochondrial morphology, positively regulates ATP

and acetyl-CoA levels, influences toxin body formation, and ultimately leads to a

significant reduction in DON toxin biosynthesis. Furthermore, deletion of FgDML1

impairs the enzymatic activity of Complex III, which subsequently induces the

upregulation of the assembly factors FgQCR2, FgQCR8, and FgQCR9, resulting in

conformational changes at the Qi site of Complex III and decreased sensitivity to

cyazofamid. In addition, the absence of FgDML1 has profound effects on

pathogenicity, fungal development, and mitochondrial morphology. Based on these

findings, we propose a working model delineating the role of FgDML1 in regulating

DON toxin biosynthesis and cyazofamid sensitivity in *F. graminearum* (Fig. 8). This

study not only provides a theoretical basis for understanding the regulatory

mechanisms underlying toxin biosynthesis in *F. graminearum* but also identifies

potential molecular targets for the development of novel agents to control FHB.

Material and methods

Strains, culture conditions, and fungicides.

The wild-type *F. graminearum* strain PH-1 was used as the parental strain for protoplast transformation experiments. All strains were preserved at -80°C. The tested strains were cultured in the dark on potato dextrose agar (PDA) at 25°C. To evaluate vegetative growth, CM, MM, and V8 media were prepared as described previously(Tang et al., 2020). MBL medium was used for conidial production, while carrot agar (CA) medium was utilized to assess sexual reproduction(Wang et al., 2011). DON toxin production was measured using TBI medium(M. Wang et al., 2024).

Cyazofamid (98%, BASF China Ltd.), pyraclostrobin (98%, BASF China Ltd.) pydiflumetofen (98%, China Syngenta crop Protection Co., Ltd.), trifloxystrobin (96%, Bayer Crop Science China Co., Ltd.), fluxapyroxad (98%, BASF China Ltd.) and fluopyram (96%, Bayer Crop Science China Co., Ltd.) were dissolved in dimethyl sulfoxide (DMSO) at stock concentrations of 50,000 μg/mL and 10,000 μg/mL, respectively.

Generation of deletion mutants and complemented mutants.

The split-marker method was employed to generate the ΔFgDML1 mutant(Yu et al., 2004). Specifically, approximately 1.2 kb of the upstream and downstream flanking regions of FgDML1 were amplified and fused with a dual-selection marker containing the hygromycin phosphotransferase (HPH) gene and the herpes simplex virus thymidine kinase (HSV-tk) gene fragments. The resulting fusion fragment was

transformed into the wild-type *F. graminearum* PH-1 strain via polyethylene glycol

(PEG)-mediated protoplast transformation. Transformants were selected on PDA

plates containing either Hygromycin B (Yeasen, Shanghai, China) or F2du (Solarbio,

Beijing, China). Positive transformants were initially screened by PCR and

subsequently verified through Southern blot analysis. To generate the complemented

mutant ΔFgDML1-C, the full-length coding sequence (CDS) of FgDML1 was

introduced into the ΔFgDML1 mutant, replacing the dual-selection cassette.

Additionally, the FgDML1-GFP mutant and FgDnm1-3×Flag mutant were

constructed using previously established methods to enable further functional

studies(C. Wang et al., 2024).

Vegetative growth and conidiation assays.

Fresh mycelial plugs of the tested strains were transferred to PDA, CM, MM, or

V8 plates to evaluate mycelial growth rates. Conidial production was induced by

incubating the strains in MBL medium, and the number of conidia was determined

using a hemocytometer. The septa of conidia were visualized through calcofluor white

(CFW) staining. For germination rate assessment, conidia were collected and

inoculated onto water agar (WA) plates, followed by incubation at 25°C for 2 and 8

hours. Germination rates were then recorded at the respective time points.

Sexual reproduction assay.

Sexual reproduction experiments were performed on CA medium. Test strains

were incubated on CA medium in complete darkness for 6 days. After incubation, the

surface mycelia were removed using a mycelial scraper and 2.5% Tween 60 solution.

The plates were then further incubated at 25°C under a 12-hour light/12-hour dark

cycle to induce sexual development. Perithecia and ascospores were observed and

recorded on days 7 and 14 using a Nikon SMZ25 fluorescent stereomicroscope and an

Olympus IX-71 inverted fluorescent microscope.

Determination of DON production and pathogenicity.

As previously described, pathogenicity tests were conducted on wheat

coleoptiles and wheat leaves. DON toxin was measured using a commercial

ELISA-based kit. Fresh mycelial plugs of the test strains were placed in TBI medium

under toxin-producing conditions (28°C, 145 $\times g$, 7 days incubation), and DON toxin

was quantified according to the commercial kit instructions after harvesting mycelia

and toxin-containing filtrates.

Determination of acetyl-CoA content and ATP concentration.

ATP content and acetyl-CoA levels were determined using the methods described

previously(Hu et al., 2023; Song et al., 2024). The test strains were cultured in YEPD

medium at 25°C and 145 $\times g$ for 36 hours. Mycelia were collected by filtration

through three layers of lens-cleaning paper, and excess moisture was removed with

filter paper. Subsequently, ATP and acetyl-CoA levels were quantified using ATP and

acetyl-CoA assay kits (Solarbio, Beijing, China), respectively.

Sensitivity determination.

To investigate the sensitivity of ΔFgDML1 to QoI and QiI fungicides, the

wild-type strain PH-1, the ΔFgDML1 mutant, and the complemented mutant

ΔFgDML1-C were inoculated on AEA plates supplemented with gradient

concentrations of cyazofamid and pyraclostrobin. To inhibit the alternative respiratory

pathway, 50 μg/mL SHAM was applied. The percentage inhibition (%) of mycelial

growth was calculated using the formula: (mean control diameter - mean diameter at

fungicide concentration) / (mean control diameter - 5 mm mycelial plug) × 100. To

evaluate the sensitivity of ΔFgDML1 to various stress conditions, the test strains were

cultured on PDA plates supplemented with stress-inducing agents, including NaCl,

KCl, sorbitol, Congo red, and SDS. Plates were incubated at 25°C for 3 days, and

colony growth was assessed.

Measurement of complex III enzyme activity.

The enzyme activity of Complex III was determined in the PH-1, ΔFgDML1,

and ΔFgDML1-C. Fresh mycelial plugs from each strain were inoculated into liquid

AEA medium and incubated at 25°C with shaking at 145 × g for 36 hours.

Subsequently, cyazofamid and pyraclostrobin were added to final concentrations of 30

μg/mL and 1.5 μg/mL, respectively, along with 50 μg/mL SHAM. After an additional

12-hour incubation, the mycelia were harvested, and Complex III activity was

measured according to the complex III enzyme activity kit (Solarbio, Beijing, China)

instructions.

Microscopic observation.

To determine the subcellular localization of FgDML1, the FgDML1-GFP mutant

was incubated in YEPD medium at 25°C with shaking at 145 × g for 36 hours.

Subcellular localization was analyzed using a Leica TCS SP8 confocal microscope,

with MitoTracker Red CMXRos (Yeasen, Shanghai, China) applied to confirm

mitochondrial targeting through colocalization of fluorescence signals. To investigate

toxisome morphology, the ΔFgDML1::Tri1+GFP mutant was cultured in TBI medium

at 28°C with shaking for 36 hours, and the structural dynamics of toxisomes within

the mycelium were visualized using Tri1-GFP fluorescence under confocal

microscopy. Additionally, mitochondrial morphology was evaluated by culturing

PH-1, ΔFgDML1, and ΔFgDML1-C in YEPD medium under identical conditions,

followed by staining with MitoTracker Red CMXRos and confocal microscopy

analysis to assess mitochondrial structural alterations, distribution, and dynamics. For

ultrastructural observations, mycelial samples were fixed and dehydrated according to

standard protocols, embedded, sectioned, and stained with uranyl acetate and lead

citrate. Mitochondrial ultrastructure was examined using a Tecnai G2 Spirit Bio

electron microscope (Thermo, MA, USA) at an accelerating voltage of 80 kV to

compare the structural integrity and morphological differences among the strains.

RNA extraction and RT-qPCR assays.

The test strains were transferred to liquid AEA medium and incubated at 25°C

for 48 hours, with cyazofamid added to a final concentration of 20 μg/mL. The

control group was not treated with any fungicide. To measure the transcriptional

levels of genes involved in DON biosynthesis, the test strains were transferred to TBI

medium and incubated for 3 days. Mycelium was collected and total RNA was

extracted following the instructions provided by the kit (Tiangen, Beijing, China).

cDNA synthesis was performed using Evo M-MLV Mix Kit (Accurate, Hunan, China).

reverse transcription. Quantitative real-time PCR (qRT-PCR) was carried out using

the QuantStudio 6 Flex real-time PCR system (Thermo, Fisher Scientific, USA) to

assess the relative expression of three subunits of Complex III (*FgCytb*, *FgCytc1*,

FgISP), five assembly factors (*FgQCR2*, *FgQCR6*, *FgQCR7*, *FgQCR8*, *FgQCR9*),

and DON biosynthesis-related genes (*FgTri5* and *FgTri6*). The *FgCox1* gene was used

as the reference gene for *FgCytb*, while *FgGapdh* was used as the reference for the

other genes. Relative transcript levels were calculated using the $2^{-\Delta\Delta CT}$ method.

Western Blot assay and Co-IP assay.

Total protein extraction was performed according to previously established

protocols, and the extracted proteins were separated using sodium dodecyl

sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Subsequently, proteins were

transferred onto an Immobilon-P transfer membrane (Millipore, Billerica, USA). The

membrane was probed with GFP antibody D191040 (BBI, Shanghai, China) and Flag

antibody 30501ES60 (Yeasen, Shanghai, China). The FgDnm1-3×Flag construct was

introduced into PH-1 and FgDML1-GFP, respectively. All tagged mutant strains were

subjected to triple validation, including PCR, sequencing, and Western blot analysis.

Total protein from tagged strains was incubated with ChromoTek GFP-Trap®

Magnetic Agarose (ChromoTek, Planegg-Martinsried, Germany) beads. Proteins

eluted from the beads were further co-incubated with GFP or Flag antibodies for

detection. Gapdh antibody 60004-1-Ig (Proteintech, Rosemont, USA) was used as an

internal control for protein normalization. Chemiluminescent detection was performed

following incubation with the corresponding secondary antibody.

Statistical analysis.

All data were analyzed using one-way analysis of variance (ANOVA), followed by the least significant difference (LSD) test with a significance threshold of P < 0.05. The EC_{50} values were calculated using DPS statistical analysis software. Each experiment was independently repeated three times.

Acknowledgements

This study was sponsored by National Natural Science Foundation of China (32272585).

CRediT authorship contribution statement

Chenguang Wang: methodology, investigation, editing, and writing original draft. Xuewei Mao: formal analysis, data curation. Weiwei Cong, Lin Yang.: performing the partial work of the experiments and analyzing the data. Yiping Hou: conceptualization, resources, writing−reviewing, supervision, funding acquisition.

Declaration of competing interest

The authors declare no conflict of interest.

References

Alexander NJ, Proctor RH, McCormick SP. 2009. Genes, gene clusters, and

biosynthesis of trichothecenes and fumonisins in fusarium. *Toxin Rev* **28**:198–

215. doi:10.1080/15569540903092142

Andrade SMP, Augusti GR, Paiva GF, Feksa HR, Tessmann DJ, Machado FJ,

Mizubuti ESG, Del Ponte EM. 2022. Phenotypic and molecular

characterization of the resistance to azoxystrobin and pyraclostrobin in

Fusarium graminearum populations from Brazil. *Plant Pathol* **71**:1152–1163.

doi:10.1111/ppa.13535

Bolgunas S, Clark DA, Hanna WS, Mauvais PA, Pember SO. 2006. Potent inhibitors

of the qi site of the mitochondrial respiration complex III. *J Med Chem*

49:4762–4766. doi:10.1021/jm060408s

Bönnighausen J, Schauer N, Schäfer W, Bormann J. 2019. Metabolic profiling of

wheat rachis node infection by Fusarium graminearum – decoding

deoxynivalenol-dependent susceptibility. *New Phytol* **221**:459–469.

doi:10.1111/nph.15377

Chen J, Xiao J, Chen G, Xu Q, Wu X, Tian L, Huang Z, Xin C, Zhao Y, Guo Z, Zou Y,

Wu Q. 2022. Indentification of novel MSTO1 compound heterozygous

mutations in a Chinese family with recessive cerebellar atrophy and ataxia.

Front Neurol **13**:988519. doi:10.3389/fneur.2022.988519

Dean R, Van Kan JAL, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD,

Rudd JJ, Dickman M, Kahmann R, Ellis J, Foster GD. 2012. The Top 10 fungal

pathogens in molecular plant pathology. *Mol Plant Pathol* **13**:414–430.

doi:10.1111/j.1364-3703.2011.00783.x

Duan Y, Lu F, Zhou Z, Zhao H, Zhang J, Mao Y, Li M, Wang J, Zhou M. 2020.

Quinone outside inhibitors affect DON biosynthesis, mitochondrial structure

and toxisome formation in *fusarium graminearum*. *J Hazard Mater* **398**:122908.

doi:10.1016/j.jhazmat.2020.122908

EFSA Panel on Contaminants in the Food Chain (CONTAM), Knutsen HK, Alexander

J, Barregård L, Bignami M, Brüschweiler B, Ceccatelli S, Cottrill B, Dinovi M,

Grasl-Kraupp B, Hogstrand C, Hoogenboom LR, Nebbia CS, Oswald IP,

Petersen A, Rose M, Roudot A-C, Schwerdtle T, Vleminckx C, Vollmer G,

Wallace H, De Saeger S, Eriksen GS, Farmer P, Fremy J-M, Gong YY, Meyer

K, Naegeli H, Parent-Massin D, Rietjens I, van Egmond H, Altieri A, Eskola M,

Gergelova P, Ramos Bordajandi L, Benkova B, Dörr B, Gkrillas A, Gustavsson

N, van Manen M, Edler L. 2017. Risks to human and animal health related to

the presence of deoxynivalenol and its acetylated and modified forms in food

and feed. *EFSA j, Eur Food Saf Auth* **15**:e04718.

doi:10.2903/j.efsa.2017.4718

Fernández-Ortuño D, Torés JA, de Vicente A, Pérez-García A. 2008. Mechanisms of

resistance to QoI fungicides in phytopathogenic fungi. *Int Microbiol: Off J Span*

Soc Microbiol **11**:1–9.

Griffin EE, Graumann J, Chan DC. 2005. The WD40 protein Caf4p is a component of

the mitochondrial fission machinery and recruits Dnm1p to mitochondria. *J*

Cell Biol **170**:237–248. doi:10.1083/jcb.200503148

Gurvitz A, Hartig A, Ruis H, Hamilton B, de Couet HG. 2002. Preliminary

characterisation of *DML1*, an essential *saccharomyces cerevisiae* gene

related to *misato* of *drosophila melanogaster*. *FEMS Yeast Res* **2**:123–135.

doi:10.1016/S1567-1356(02)00083-1

Hai S, Chen J, Ma L, Wang C, Chen C, Rahman SU, Zhao C, Feng S, Wu J, Wang X.

2023. Combination of zearalenone and deoxynivalenol induces apoptosis by

mitochondrial pathway in piglet sertoli cells: role of endoplasmic reticulum

stress. *Toxins* **15**:471. doi:10.3390/toxins15070471

Hu L, Guo C, Chen J, Jia R, Sun Y, Cao S, Xiang P, Wang Y. 2023. Venturicidin a is a

potential fungicide for controlling fusarium head blight by affecting

deoxynivalenol biosynthesis, toxisome formation, and mitochondrial structure.

J Agric Food Chem **71**:12440–12451. doi:10.1021/acs.jafc.3c02683

Huang P, Yu X, Liu H, Ding M, Wang Z, Xu J-R, Jiang C. 2024. Regulation of TRI5

expression and deoxynivalenol biosynthesis by a long non-coding RNA in

fusarium graminearum. *Nat Commun* **15**:1216.

doi:10.1038/s41467-024-45502-w

Huang Q, Tonge PJ, Slayden RA, Kirikae T, Ojima I. 2007. FtsZ: a novel target for

tuberculosis drug discovery. *Curr Top Med Chem* **7**:527–543.

doi:10.2174/156802607780059790

Ilgen P, Hadeler B, Maier FJ, Schäfer W. 2009. Developing kernel and rachis node

induce the trichothecene pathway of fusarium graminearum during wheat

head infection. *Mol Plant-Microbe Interact* **22**:899–908.

doi:10.1094/MPMI-22-8-0899

Jiang C, Zhang C, Wu C, Sun P, Hou R, Liu H, Wang C, Xu J-R. 2016. TRI6 and

TRI10 play different roles in the regulation of deoxynivalenol (DON) production

by cAMP signalling in fusarium graminearum. *Environ Microbiol* **18**:3689–3701.

doi:10.1111/1462-2920.13279

Kang J, Zhang J, Liu Y, Song J, Ou J, Tao X, Zhou M, Duan Y. 2023. Mitochondrial

dynamics caused by QoIs and SDHIs fungicides depended on FgDnm1 in

fusarium graminearum. J Integr Agric **22**:481–494.

doi:10.1016/j.jia.2022.08.118

Kimura M, Anzai H, Yamaguchi I. 2001. Microbial toxins in plant-pathogen interactions:

biosynthesis, resistance mechanisms, and significance. *J Gen Appl Microbiol*

47:149–160. doi:10.2323/jgam.47.149

Kimura M, Okano Y. 2007a. Human misato regulates mitochondrial distribution and

morphology. *Exp Cell Res* **313**:1393–1404. doi:10.1016/j.yexcr.2007.02.004

Kimura M, Okano Y. 2007b. Human misato regulates mitochondrial distribution and

morphology. *Exp Cell Res* **313**:1393–1404. doi:10.1016/j.yexcr.2007.02.004

Kimura M, Tokai T, Takahashi-Ando N, Ohsato S, Fujimura M. 2007. Molecular and

genetic studies of fusarium trichothecene biosynthesis: pathways, genes, and

evolution. *Biosci, Biotechnol, Biochem* **71**:2105–2123. doi:10.1271/bbb.70183

Li H, Zhu X-L, Yang W-C, Yang G-F. 2014. Comparative Kinetics of Qi Site Inhibitors

of Cytochrome bc1 Complex: Picomolar Antimycin and Micromolar

Cyazofamid. *Chem Biol Drug Des* **83**:71–80. doi:10.1111/cbdd.12199

Li J, Duan Y, Bian C, Pan X, Yao C, Wang J, Zhou M. 2019. Effects of validamycin in

controlling fusarium head blight caused by fusarium graminearum: inhibition of

DON biosynthesis and induction of host resistance. *Pestic Biochem Physiol*

153:152–160. doi:10.1016/j.pestbp.2018.11.012

Liu X, Yu F, Schnabel G, Wu J, Wang Z, Ma Z. 2011. Paralogous cyp51 genes in

Fusarium graminearum mediate differential sensitivity to sterol demethylation

inhibitors. *Fungal genet biol: FG B* **48**:113–123. doi:10.1016/j.fgb.2010.10.004

Liu Y, Lu Y, Wang L, Chang F, Yang L. 2016. Occurrence of deoxynivalenol in wheat,

hebei province, china. *Food Chemistry* **197**:1271–1274.

doi:10.1016/j.foodchem.2015.11.047

Liu Z, Jian Y, Chen Y, Kistler HC, He P, Ma Z, Yin Y. 2019. A phosphorylated

transcription factor regulates sterol biosynthesis in Fusarium graminearum.

Nat Commun **10**:1228. doi:10.1038/s41467-019-09145-6

Marreddy RKR, Phelps GA, Churion K, Picker J, Powell R, Cherian PT, Bowling JJ,

Stephan CC, Lee RE, Hurdle JG. 2024. Chemical genetic analysis of

enoxolone inhibition of *clostridioides difficile* toxin production reveals adenine

deaminase and ATP synthase as antivirulence targets. *J Biol Chem*

300:107839. doi:10.1016/j.jbc.2024.107839

Merhej J, Richard-Forget F, Barreau C. 2011a. Regulation of trichothecene

biosynthesis in fusarium: recent advances and new insights. *Appl Microbiol*

Biotechnol **91**:519–528. doi:10.1007/s00253-011-3397-x

Merhej J, Richard-Forget F, Barreau C. 2011b. Regulation of trichothecene

biosynthesis in fusarium: recent advances and new insights. *Appl Microbiol*

Biotechnol **91**:519–528. doi:10.1007/s00253-011-3397-x

Miklos GLG, Yamamoto M-T, Burns RG, Maleszka R. 1997. An essential cell division

gene of drosophila, absent from saccharomyces, encodes an unusual protein

with   tubulin-like and myosin-like peptide motifs. *Proc Natl Acad Sci* **94**:5189–

5194. doi:10.1073/pnas.94.10.5189

Mitani S, Araki S, Takii Y, Ohshima T, Matsuo N, Miyoshi H. 2001. The biochemical

mode of action of the novel selective fungicide cyazofamid: specific inhibition

of mitochondrial complex III in *phythium spinosum*. *Pesticide Biochemistry and*

Physiology **71**:107–115. doi:10.1006/pest.2001.2569

Mrnjavac N, Martin WF. 2025. GTP before ATP: the energy currency at the origin of

genes. *Biochim Biophys Acta (BBA) - Bioenerg* **1866**:149514.

doi:10.1016/j.bbabio.2024.149514

Mudge AM, Dill-Macky R, Dong Y, Gardiner DM, White RG, Manners JM. 2006. A role

for the mycotoxin deoxynivalenol in stem colonisation during crown rot disease

of wheat caused by *fusarium graminearum* and *fusarium pseudograminearum*.

Physiological and Molecular Plant Pathology **69**:73–85.

doi:10.1016/j.pmpp.2007.01.003

Nasca A, Scotton C, Zaharieva I, Neri M, Selvatici R, Magnusson OT, Gal A, Weaver

D, Rossi R, Armaroli A, Pane M, Phadke R, Sarkozy A, Muntoni F, Hughes I,

Cecconi A, Hajnóczky G, Donati A, Mercuri E, Zeviani M, Ferlini A, Ghezzi D.

2017. Recessive mutations in MSTO1 cause mitochondrial dynamics

impairment, leading to myopathy and ataxia. *Hum Mutat* **38**:970–977.

doi:10.1002/humu.23262

Pb P, Gc   de FS, Gmb L, Plo C, Aln V, Ht F. 2023. Competitive interaction between

ATP and GTP regulates mitochondrial ATP-sensitive potassium channels.

Chem-Biol Interact **381**. doi:10.1016/j.cbi.2023.110560

Rogne P, Rosselin M, Grundström C, Hedberg C, Sauer UH, Wolf-Watz M. 2018.

Molecular mechanism of ATP versus GTP selectivity of adenylate kinase. *Proc*

Natl Acad Sci U S A **115**:3012–3017. doi:10.1073/pnas.1721508115

Sarewicz M, Dutka M, Pintscher S, Osyczka A. 2013. Triplet state of the

semiquinone-rieske cluster as an intermediate of electronic bifurcation

catalyzed by cytochrome bc1. *Biochemistry* **52**:6388–6395.

doi:10.1021/bi400624m

Schulze SR, Wallrath LL. 2007. Gene regulation by chromatin structure: paradigms

established in drosophila melanogaster. *Annu Rev Entomol* **52**:171–192.

doi:10.1146/annurev.ento.51.110104.151007

Song J, Li Y, Zhang Z, Gao X, Li S, Zhang J, Zhou M, Duan Y. 2024. Endoplasmic

reticulum-mitochondrial encounter structure regulates the mitochondrial

morphology, DON biosynthesis and toxisome formation in *fusarium*

graminearum. *Microbiol Res* **289**:127892. doi:10.1016/j.micres.2024.127892

Sumarah MW. 2022. The deoxynivalenol challenge. *J Agric Food Chem* **70**:9619–

9624. doi:10.1021/acs.jafc.2c03690

Tang G, Chen A, Dawood DH, Liang J, Chen Y, Ma Z. 2020. Capping proteins

regulate fungal development, DON-toxisome formation and virulence in

Fusarium graminearum. *Molecular Plant Pathology* **21**:173–187.

doi:10.1111/mpp.12887

Tang G, Chen Y, Xu J-R, Kistler HC, Ma Z. 2018. The fungal myosin I is essential for

Fusarium toxisome formation. *PLoS Pathog* **14**:e1006827.

doi:10.1371/journal.ppat.1006827

Vercellino I, Sazanov LA. 2022. The assembly, regulation and function of the

mitochondrial respiratory chain. *Nat Rev Mol Cell Biol* **23**:141–161.

doi:10.1038/s41580-021-00415-0

Vollmer W. 2006. The prokaryotic cytoskeleton: a putative target for inhibitors and

antibiotics? *Appl Microbiol Biotechnol* **73**:37–47.

doi:10.1007/s00253-006-0586-0

Wang C, Zhang S, Hou R, Zhao Z, Zheng Q, Xu Q, Zheng D, Wang G, Liu H, Gao X,

Ma J-W, Kistler HC, Kang Z, Xu J-R. 2011. Functional analysis of the kinome

of the wheat scab fungus fusarium graminearum. *PLOS Pathog* **7**:e1002460.

doi:10.1371/journal.ppat.1002460

Wang C, Zhao F, Wu Z, Cai X, Zhou M, Hou Y. 2024. Mitochondria-Associated

Protein FgNdk1 Regulates the Development, Pathogenicity, and SDHI

Fungicide Sensitivity of *Fusarium graminearum* by Interacting with Succinate

Dehydrogenase. *J Agric Food Chem* **72**:3913–3925.

doi:10.1021/acs.jafc.3c07934

Wang M, Wu N, Wang H, Liu C, Chen Q, Xu T, Chen Y, Zhao Y, Ma Z. 2024.

Overproduction of mycotoxin biosynthetic enzymes triggers fusarium

toxisome-shaped structure formation via endoplasmic reticulum remodeling.

PLOS Pathog **20**:e1011913. doi:10.1371/journal.ppat.1011913

Wang Y, Shi Y, Li W, Han X, Lin X, Liu D, Lin Y, Shen L. 2024. Knockdown of

BRAWNIN minimally affect mitochondrial complex III assembly in human cells.

Biochim Biophys Acta, Mol Cell Res **1871**:119601.

doi:10.1016/j.bbamcr.2023.119601

Xiao J, Zhou Y, Xie Y, Li T, Su X, He J, Jiang Y, Zhu H, Qu H. 2024. ATP homeostasis

and signaling in plants. *Plant Commun* **5**:100834.

doi:10.1016/j.xplc.2024.100834

Yi L, Yang M, Waalwijk C, Xu Jin, Xu Jingsheng, Molnár O, Chen W, Feng J, Zhang H.

2023. Dynamics of carbendazim-resistance frequency of pathogens

associated with the epidemic of fusarium head blight. *Plant Dis* **107**:1690–

1696. doi:10.1094/PDIS-08-22-1998-SR

Yu J-H, Hamari Z, Han K-H, Seo J-A, Reyes-Domínguez Y, Scazzocchio C. 2004.

Double-joint PCR: a PCR-based molecular tool for gene manipulations in

filamentous fungi. *Fungal Genet Biol* **41**:973–981.

doi:10.1016/j.fgb.2004.08.001

Zara V, Conte L, Trumpower BL. 2009. Evidence that the assembly of the yeast

cytochrome bc1 complex involves the formation of a large core structure in the

inner mitochondrial membrane. *FEBS J* **276**:1900–1914.

doi:10.1111/j.1742-4658.2009.06916.x

Zara V, Conte L, Trumpower BL. 2007. Identification and characterization of

cytochrome bc(1) subcomplexes in mitochondria from yeast with single and

double deletions of genes encoding cytochrome bc(1) subunits. *FEBS J*

274:4526–4539. doi:10.1111/j.1742-4658.2007.05982.x

Zhan J, Zeher A, Huang R, Tang WK, Jenkins LM, Xia D. 2024. Conformations of

Bcs1L undergoing ATP hydrolysis suggest a concerted translocation

mechanism for folded iron-sulfur protein substrate. *Nat Commun* **15**:4655.

doi:10.1038/s41467-024-49029-y

Zhang Y, Ouyang B, Zhang W, Guang C, Xu W, Mu W. 2024. Deoxynivalenol:

occurrence, toxicity, and degradation. *Food Control* **155**:110027.

doi:10.1016/j.foodcont.2023.110027

Zhao Y, Chi M, Sun H, Qian H, Yang J, Huang J. 2021. The FgCYP51B Y123H

Mutation Confers Reduced Sensitivity to Prochloraz and Is Important for

Conidiation and Ascospore Development in Fusarium graminearum.

Phytopathology **111**:1420–1427. doi:10.1094/PHYTO-09-20-0431-R

Zhou F, Zhou X, Yan Jiao N, Han A, Zhou H, Chen Z, Li W, Liu R. 2024. Baseline

tebuconazole sensitivity and potential resistant risk in Fusarium Graminearum.

BMC Plant Biol **24**:789. doi:10.1186/s12870-024-05206-1

Zhou Z, Duan Y, Zhou M. 2020. Carbendazim-resistance associated β2 -tubulin

substitutions increase deoxynivalenol biosynthesis by reducing the interaction

between β2 -tubulin and IDH3 in fusarium graminearum. *Environ Microbiol*

22:598–614. doi:10.1111/1462-2920.14874

A

B

FIG. 1.

Identification and localization of FgDML1 in *Fusarium graminearum*. (A) The phylogenetic tree was constructed based on the amino acid sequences of DML1 homologs from 10 species using the maximum likelihood method in Mega X. Bootstrap values from 1000 replications were displayed on the branches. (B) FgDML1 was localized to the mitochondria.

FIG. 2.

FgDML1 is required for both the virulence of *F. graminearum*. (A) Lesion lengths on wheat coleoptiles caused by different strains. These images were taken at 14 dpi. (B) Dot plot showing the lesion lengths on wheat coleoptiles at 14 dpi tested with different strains. (C) Disease symptoms on wheat leaves inoculated with PH-1, ΔFgDML, and ΔFgDML-C. Images were taken at 5 dpi. (D) Disease area on wheat leaves inoculated with different strains at 5 dpi.

FIG. 3.

FgDML1 is required for the biosynthesis of DON toxin in *F. graminearum*. (A) ΔFgDML1 exhibited decreased DON content. All strains were incubated in trichothecene biosynthesis induction (TBI) medium for 7 days. (B) Relative mRNA expression levels of *FgTRI5*, and *FgTRI6* in the tested strains. Mycelia were harvested after 2 days of cultivation in TBI medium, and mRNA were extracted for quantification using the $2^{-\Delta\Delta CT}$ method. *FgGapdh* was used as the reference gene. (C) FgDML1 impaired the formation of DON toxisomes, as visualized by Tri1-GFP labeling of toxisomes. Strains were cultured in TBI for 36 hours. Bar = 10 μm (D) ΔFgDML1 exhibited reduced expression levels of Tri1-GFP. Strains were cultured in TBI for 36 hours, and mycelia were harvested for western blot analysis. Bars with the

same letter indicate no significant difference according to a significant difference

(LSD) test at P = 0.05.

FIG. 4.

FgDML1 modulates mitochondrial morphology via interaction with FgDnm1 and negatively regulates Acetyl-CoA and ATP synthesis. (A) Effects of FgDML1 on mitochondrial morphology. Mycelial plugs of the wild-type strain PH-1, the ΔFgDML1 deletion mutant, and the complemented mutant ΔFgDML1-C were cultured in YEPD medium for 36 hours. Mitochondria were stained using a mitochondria-specific fluorescent dye and observed under a confocal microscope. Additionally, conidial germlings were cultured in MBL medium and subjected to the same staining and imaging procedures. Scale bar = 5 μm. (B) Ultrastructural analysis of mitochondria by transmission electron microscopy (TEM). Mitochondrial ultrastructure was further examined in PH-1, ΔFgDML1, and ΔFgDML1-C using TEM. Mitochondria are indicated by red arrows. Scale bar = 200 nm. (C) FgDML1 Deletion Reduces Acetyl-CoA Levels. Mycelial cultures of PH-1, ΔFgDML1, and ΔFgDML1-C were incubated in YEPD medium for 36 hours. Acetyl-CoA levels were

quantified using a commercial assay kit, following the instructions provided by the commercial assay kit. (D) FgDML1 deletion reduces ATP levels. The ATP content of PH-1, ΔFgDML1, and ΔFgDML1-C was measured following the same culture conditions as in (C), using a commercial ATP assay kit. (E) Co-immunoprecipitation (Co-IP) confirms interaction between FgDML1 and FgDnm1. Total protein extracts from strains expressing tagged versions of FgDML1 and FgDnm1 were separated by SDS-PAGE and subjected to immunoblot analysis. Co-IP was performed using ChromoTek GFP-Trap® magnetic agarose to pull down interacting proteins. Immunodetection was carried out using polyclonal anti-Flag and monoclonal anti-GFP antibodies. Monoclonal anti-GAPDH antibody was used as a reference for protein samples.

FIG. 5.

FgDML1 modulates the sensitivity of *F. graminearum* to cyazofamid. (A) ΔFgDML1 exhibits reduced sensitivity to cyazofamid. PH-1, ΔFgDML1, and ΔFgDML1-C were cultured on AEA medium modified with cyazofamid, with 50 µg/mL SHAM included to inhibit the alternative oxidase pathway. (B) Deletion of FgDML1 did not affect *F. graminearum* sensitivity to pyraclostrobin. The tested strains were incubated on AEA medium containing pyraclostrobin and 50 µg/mL SHAM to assess their response to the fungicide. (C) FgDML1 deletion reduces complex III enzymatic activity. Mycelial cultures of PH-1, ΔFgDML1, and ΔFgDML1-C were grown in YEPD for 36 hours, then treated with 30 µg/mL cyazofamid, 1.5 µg/mL pyraclostrobin, and 50 µg/mL SHAM for an additional 12 hours. The enzymatic activity of complex III was

quantified using a commercial assay kit. (D) mRNA expression levels of complex III subunits and assembly factors in PH-1, ΔFgDML1, and ΔFgDML1-C. PH-1, ΔFgDML1, and ΔFgDML1-C were incubated in liquid AEA medium for 36 hours, followed by treatment with 30 μg/mL cyazofamid and 50 μg/mL SHAM. The relative mRNA expression levels of complex III subunits and assembly factors were quantified using the $2^{-\Delta\Delta CT}$ method. "T" denotes cyazofamid treatment.

FIG. 6.

Effects of ΔFgDML1 on asexual and sexual development. (A) Mycelial growth rate. PH-1, ΔFgDML1, ΔFgDML1-C were cultured for 3 days on PDA, CM, MM, and V8 solid media at 25°C. (B) Mycelial morphology. The mycelial morphology of PH-1, ΔFgDML1, and ΔFgDML1-C was examined on WA plates. Scale bar = 72 μm. (C) Conidial length. Conidia of ΔFgDML1 were significantly longer in length compared to PH-1 and ΔFgDML1-C. Conidia were germinated in MBL medium, stained with Calcofluor White (CFW), and observed using an Olympus IX-71 microscope. (D) Sexual reproduction. PH-1 and ΔFgDML1-C produced perithecia and asci containing ascospores after 7 or 14 days of incubation on CA medium, while the ΔFgDML1 failed to produce these structures under the same conditions. Perithecia and ascospores were observed using a Nikon SMZ25 fluorescent stereo microscope and an Olympus IX-71 inverted fluorescence microscope. Top image: perithecium, scale

bar = 1 mm; bottom image: ascospores, scale bar = 50 μm. (E) Proportion of conidial septa. The proportion of conidial septa was assessed in the tested strains. (F) Conidial production. Conidia were harvested after 4 days of incubation in MBL medium, and their concentration was quantified using a hemocytometer. (G) Germination rate. Conidial germination was observed by incubating 300 conidia for 2 or 8 hours on WA plates. Germination rates were compared, with different letters indicating significant differences based on ANOVA followed by the LSD test (P = 0.05).

FIG. 7.

Response of the strains to different stresses. (A) Reduced sensitivity to osmotic stress in ΔFgDML1. PH-1, ΔFgDML1 and ΔFgDML1-C were subjected to osmotic stress using NaCl, KCl, and Sorbitol to evaluate their sensitivity. (B) Statistical analysis of growth inhibition under osmotic stress. Growth inhibition rates of all strains under NaCl, KCl, and Sorbitol stress were statistically analyzed. (C) Sensitivity to cell wall

and membrane damaging agents. The tested strains were exposed to cell wall-damaging agent Congo Red (CR) and cell membrane-damaging agent Sodium dodecyl sulfate (SDS) to assess their sensitivity. (D) Statistical analysis of growth inhibition under CR and SDS stress. Growth inhibition rates of all strains under CR and SDS stress were statistically analyzed. (E) Sensitivity to temperature stresses. The sensitivity of the tested strains to different temperature stresses was evaluated. (F) Statistical analysis of growth inhibition under temperature stress. Growth inhibition rates of all strains under various temperature stresses were statistically analyzed. Significant differences were determined using the LSD test at $P = 0.05$. Bars with the same letter indicate no significant difference.

FIG. 8.

Model of FgDML1 regulation of *F. graminearum* sensitivity to DON toxin synthesis and cyazofamid fungicide. FgDML1 interacts with FgDnm1 to modulate mitochondrial morphology, thereby positively regulating acetyl-CoA and ATP levels, which influence toxisome formation and ultimately affect DON production. Loss of FgDML1 reduces complex III enzymatic activity, leading to the upregulation of the assembly factors *FgQCR2*, *FgQCR8*, and *FgQCR9*. This alteration in assembly factor expression changes the conformation of the Qi site, leading to reduced sensitivity to cyazofamid.