Male gut microbiome mediates post-mating sexual selection in *Drosophila*

Komal Maggu*, Abhishek Meena, Alessio N. Nardo, Sonja H. Sbilordo, Jeannine Roy &

Stefan Lüpold*

Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, 8057 Zurich, Switzerland

Correspondence: stefan.luepold@ieu.uzh.ch, komal024maggu@gmail.com

ABSTRACT

In polyandrous systems, male reproductive success is critically influenced by sperm

competition, a process often mediated by ejaculate characteristics that are sensitive to male

condition. While condition dependence is a cornerstone of sexual selection theory, the role of

the gut microbiome in modulating male reproductive fitness under sperm competition remains

unexplored. Here, we investigated the impact of varying gut microbial compositions on sperm

competitive ability in *Drosophila melanogaster*. By inoculating germ-free (axenic) males with

all 32 full-factorial combinations of five common commensal bacteria, we demonstrated that

the presence and diversity of the gut microbiota significantly enhance male sperm offense

ability. Specifically, males associated with a more diverse microbiota exhibited superior

competitive fertilization success, accompanied by increased sperm transfer, higher sperm

viability, and a greater proportion of sperm stored by females. Predictive models further

revealed complex, non-additive interactions among bacterial species in their effects on sperm

competitiveness. These findings suggest that the gut microbiome acts as a critical modulator

of male reproductive fitness, challenging traditional models of sexual selection that overlook

contributions of non-pathogenic bacteria to condition-dependent traits.

INTRODUCTION

In polyandrous species, where females mate with multiple males, sperm competition often

determines fertilization success based on ejaculate size and quality (1, 2). However, producing

large, high-quality ejaculates is costly by diverting resources from other fitness traits (3–8).

The dependence of trait expression on resource availability and allocation between growth,

maintenance, and reproduction is central to sexual selection theory, linking phenotypic quality

to underlying genetic variation: traits that reliably indicate a male's ability to acquire and

allocate limited resources are hypothesized to signal heritable fitness benefits to mates or rivals

(9, 10). This 'good genes' framework assumes that male condition is shaped by genetic variation

in resource acquisition, allocation efficiency, or stress tolerance (11, 12). However, mounting

evidence suggests that the gut microbiome, a dynamic microbial consortium, may decouple

condition from host genetics by enhancing nutrient extraction, metabolic regulation, and

hormone synthesis (7–9). If microbial communities independently modulate resource budgets,

they could fundamentally alter the heritability of condition-dependent traits, challenging a

cornerstone of sexual selection theory.

The gut microbiome directly influences host physiology, including nutrient assimilation

(13–15) and energy allocation (16–17). For example, *Acetobacter* and *Lactobacillus* species in

Drosophila melanogaster differentially regulate host metabolism, with *Acetobacter*-dominated

microbiomes enhancing nutrient absorption and fecundity in females (18, 19). Conversely,

disruptions to the gut microbiome can alter host metabolism, gene expression, or epigenetic

modifications through changes in metabolite, enzyme, and hormone signalling (20–23), which

can impair reproductive functions (23–25) or cause pathologies (26–28). While these effects

are well-documented (29, 30), the role of the gut microbiome in male reproductive

performance, particularly in the context of post-mating sexual selection, remains entirely

unexplored.

In *D. melanogaster*, a model system for host-microbe interactions, the gut microbiome

is simple, tractable (31, 32), and critical for female reproductive traits (33, 34). For example,

microbial composition shapes female fecundity, lifespan, and offspring development (18, 19).

Such knowledge, however, is lacking for males. Limited work in mammals links gut dysbiosis

to impaired testicular development (35) and spermatogenesis (36), while seminal microbes

alter sperm motility in humans (37, 38). Yet, nothing is known about how gut microbes

influence male fertilization success, let alone under sperm competition.

Here, we tested the hypothesis that the gut microbiome modulates male reproductive

fitness under competitive fertilization by enhancing condition-dependent ejaculate traits. Using

D. melanogaster males inoculated with all 32 combinations of five gut bacteria (*Acetobacter*

orientalis, *A. tropicalis*, *A. pasteurianus*, *Lactobacillus brevis*, and *L. plantarum*; (19)), we

quantified competitive fertilization success (P_2) and its mechanistic underpinnings: sperm

viability, transfer, and storage. We predicted that (i) males with diverse microbiomes

outperform axenic or low-diversity males due to improved nutrient acquisition (39, 40), (ii),

synergistic interactions (e.g., *Acetobacter* dominance) amplify competitive advantages (19),

and (iii) microbiome-driven gains in sperm viability and transfer efficiency underpin

competitive fertilization success (41). By resolving how microbial communities shape sperm

competition, this study bridges microbial ecology with sexual selection theory, offering novel

insights into the evolution of condition-dependent traits.

Results

Microbial diversity enhances sperm competitiveness

To test if gut microbes influence male reproductive fitness under post-mating sexual selection, we compared the competitive fertilization success of male *D. melanogaster*, inoculated in all possible combinations between our five strains of gut bacteria, each against either an axenic (germ-free) or a conventional male (i.e., associated with an intact microbiome). These standard competitor males, which mated three days before our wild-type focal males, ubiquitously expressed green fluorescent protein (GFP) to facilitate paternity assignment (42, 43).

In competition against axenic rivals, we found experimental males harbouring any single bacterial species (except *L. brevis*) or consortium to outperform males without a microbiome, with second-male paternity (P_2) increasing with species richness ($t_{30} = 2.18$, $P = 0.037$; Fig. 1A). Being associated with all five species translated into a 27% gain in P_2 compared to the axenic state (mean ± s.d. = 0.85 ± 0.05 vs. 0.67 ± 0.05, respectively). Further, *Acetobacter*-dominated communities conferred greater benefits than those biased toward *Lactobacillus* (Fig. 2A). These patterns persisted when focal males competed against rivals with intact microbiotas (increase with richness: $t_{30} = 2.44$, $P = 0.021$; relative gain for 5-species association: 20% (0.72 ± 0.08 vs. 0.60 ± 0.03)), confirming that microbial benefits are not contingent on competitors being axenic. The contrasting impacts of *Acetobacter* and *Lactobacillus* on P_2 were further supported by bootstrapped correlations (Fig. 3A) showing P_2 varied with the relative abundance (based on CFU counts) of each species across different microbial communities, not just with their presence or absence (Fig. 3A).

Microbial interactions shape competitive outcomes

Mixed-effects models accounting for experimental block revealed that microbial interactions, rather than additive effects, drove variation in P_2. Even though the five-way interaction between

bacterial strains was not statistically significant in competitions against either axenic ($N = 646$,

$t = 0.58$, $P = 0.561$) or conventional rivals ($N = 624$, $t = -0.86$, $P = 0.388$), eight and six of the

10 pairwise interactions, respectively, were statistically significant (all $t \leq -2.00$, $P \leq 0.046$),

with another two trends among the three-way interactions ($t \geq 1.69$, $P \leq 0.092$). Further, when

removing the non-significant five-way interaction, the four-way interaction of $At \times Ap \times Lb \times$

Lp ($t = 2.13$, $P = 0.034$) was also indicative of a more complex contribution of microbial

composition in this scenario.

Predictive modelling showed that averaging mono-association effects between bacterial

strains poorly forecasted outcomes for multi-species communities (e.g., mean of At, Ap, and

Lb to predict the At-Ap-Lb combination), with only eight (30.8%) of the 26 possible

combinations correctly predicted based on 95% confidence intervals (95%CI) including zero

difference in competitive matings against axenic rivals (Fig. 2B). Averaging pairwise

combinations to predict three-way species interactions (e.g., At-Ap, At-Lb and Ap-Lb to predict

At-Ap-Lb) indicated similar accuracy (31.2%: 5 of 16 predicted combinations; Fig. 2B). These

predictions by mono- or di-associations were broadly comparable when focal males competed

against conventional rather than axenic first males (33.0 and 37.5%, respectively). Combined,

these results further support the importance of interactions between bacterial species.

Additionally, pairwise interactions were context-dependent, varying conditionally on the

remaining ("bystander") species as reflected by interactions trending from positive

(synergistic) to negative (antagonistic) between the same two species depending on the

configuration of additional species in the microbiota (Fig. 4A). This dependence on bystanders

also held for three-way interactions between focal species (Fig. 4B).

Mechanistic basis of microbiome-mediated fitness

To dissect how microbiomes enhance competitive fertilization success, we quantified sperm transfer, sperm viability, and female sperm retention and storage patterns, based on prior knowledge (42–45). Due to the similar results between axenic and conventional competitor males, we here restricted our mechanistic investigation to axenic first males. For visualization of sperm, all assays except sperm viability involved standard and experimental males that, respectively, expressed green or red fluorescent protein in sperm heads (44).

While axenic and mono-associated focal males transferred similar sperm quantities, multi-associated males transferred 3–36% more sperm than either of the former categories, with gains scaling with microbial diversity ($t_{30} = 5.63$, $P < 0.0001$; Figs. 1B and 2C,D). Sperm viability also increased with diversity: males with all five species had $94.9 \pm 2.3\%$ viable sperm compared to $54.5 \pm 6.5\%$ in axenic males ($t_{30} = 11.54$, $P < 0.0001$; Figs. 1C and 2E, F). Critically, although freshly remated females did not significantly change the timing of ejecting displaced and excess sperm from first (standard) and second (experimental) males, respectively, in response to second-male microbial diversity ($t_{30} = 1.10$, $P = 0.100$; Figs. 1D and 2G,H), the sperm retained were increasingly biased toward focal males if these had more diverse microbiomes ($t_{30} = 13.10$, $P < 0.0001$; Figs. 1E, 2I,J). Hence, microbiome-enhanced sperm quality or displacement efficiency biased post-mating sperm storage dynamics. Confirming prior work (42, 43, 45, 46), the relative numbers of sperm stored and their viability were significantly correlated with paternity outcomes (Fig. 3B). Further, these phenotypic traits again covaried mostly positively with *Acetobacter* and negatively with *Lactobacillus* when considering relative colony forming units (CFUs) rather than species presence or absence (Fig. 3A).

Similar to P_2, sperm viability and the relative representation of focal-male sperm in storage after female sperm ejection responded to microbial interactions, with significant five-

way interactions in generalized linear mixed models for both sperm viability ($N = 320$, $z = -$

2.03, $P = 0.043$) and the proportion of second-male sperm in storage ($N = 582$, $z = 2.11$, $P =$

0.035). Predictive models for both variables also indicated considerable underestimation of

measured values by both averaged mono- or di-associations (Fig. 2F,J), highlighting the

importance of higher-order interactive effects, whereas this dependence was weaker for both

the number of sperm transferred and time to sperm ejection. For all stages of competitive

fertilization process, the interactive effects between pairs or triplets of microbes were strongly

influenced by bystander species in their contribution to phenotypic variation (Fig. 4C-J).

Discussion

Our study of *Drosophila melanogaster* establishes the gut microbiome as a significant

determinant of male reproductive success competitive fertilization. We demonstrate that

microbial diversity and composition, shaped by complex interspecific interactions, markedly

influence male competitive fertilization success (P_2). A reproductive advantage is directly

linked to improvements in critical ejaculate components, including increased sperm transfer,

substantially enhanced sperm viability (e.g., from ~55% in axenic males to ~95% in males with

the full 5-species consortium), and a greater proportion of sperm stored by females after mating

(Fig. 2). These findings redefine condition-dependent reproductive success by revealing the

gut microbiome as a dynamic mediator of sexual selection. This challenges classical

frameworks that largely attribute condition dependence to host resource status or stress

resilience (9–12) highlighting instead how symbiotic microbial interactions can directly shape

ejaculate traits and competitive outcomes, thereby acting as a critical, previously overlooked

evolutionary driver in reproductive adaptation (47).

Linking microbial ecology to ejaculate quality

The pronounced reproductive deficits in axenic males align with established connections between microbiome absence and physiological dysregulation in *Drosophila* (48). While our study did not directly assay host metabolic states, prior work compellingly shows that axenic flies often exhibit hyperlipidemia and impaired insulin signalling – conditions detrimental to overall health and potentially sperm function (46, 49–53). These metabolic disturbances can be rescued or modulated by specific gut bacteria; for example, *Acetobacter* species are known to restore lipid homeostasis, potentially via acetate production (54), whilst certain *Lactobacillus* species can also influence host metabolism, albeit sometimes with trade-offs like increased oxidative stress (55).

It is important to emphasize that our experimental design involved inoculating males only after adult eclosion, following axenic development. This approach effectively isolates the microbiome's influence on adult physiological processes critical for ejaculate production, maintenance, and competitive performance. However, this focus on adult function means that our estimates of the total contribution of the gut microbiome to male reproductive success are likely to be conservative. Microbes present during larval and pupal development in insects are known to impact host growth rate, final body size, and the maturation of various tissues (56). Had these microbial partners also influenced the developmental trajectory—potentially affecting body size, resource acquisition capacity, or the formation of reproductive organs— the observed effects on adult competitive ability could have been further amplified or otherwise altered. Disentangling these life-stage specific microbial effects remain an important avenue for future research.

Based on our experimental context, we hypothesize that the observed improvements in ejaculate quality and subsequent P_2 gains conferred by species-rich microbial communities stem from their ability to restore or optimize host metabolic homeostasis and nutrient allocation

toward reproduction. The superior performance associated with *Acetobacter*-dominated

communities (Fig. 1) lends support to this hypothesis, given their known efficacy in nutrient

assimilation and influencing host metabolism (48). These microbes may channel resources

more effectively towards energetically costly ejaculate production and maintenance,

potentially mitigating damage from metabolic byproducts. However, the significant non-

additive effects underscore that the *integrated function* of the microbial consortium, rather than

simply the presence of key species, dictates the ultimate physiological impact (Fig. 3),

mirroring the metabolic interplay seen in microbial ecology (57). Direct confirmation of these

proposed metabolic links—identifying specific microbial metabolites (e.g., acetate or lactate)

and tracing their effects on host resource allocation and sperm physiology—awaits future

metabolomic and transcriptomic studies.

Evolutionary implications

Our findings prompt a re-evaluation of core tenets in sexual selection theory. By acting as an

amplifier of condition-dependent reproductive traits, the microbiome introduces complexity to

the "good genes" model, where such traits are assumed to reliably signal host genetic quality

(10, 11). Our results suggest that the microbiome functions as a "hidden mediator," bridging

host genetics and environmental inputs (58), with at least three key evolutionary implications.

First, the microbiome potentially introduces a transmissible component influencing fitness that

is sensitive to environmental factors (diet, social context) and operates outside strict Mendelian

inheritance (58). If microbial consortia are consistently acquired or partially transmitted

(vertically or horizontally), this could contribute to maintaining heritable variation in sexually

selected traits, offering a potential factor in resolving the lek paradox (11). Furthermore, the

microbiome might act as an "environmental sensor," allowing males to buffer reproductive

output against nutritional fluctuations (5), thereby stabilizing their fitness.

Second, the context-dependent microbial interactions (Fig. 3) and potential trade-offs

(e.g., benefits vs. costs like oxidative stress potentially associated with *L. brevis*) suggest

complex coevolutionary dynamics. Selection may favour host mechanisms for managing

microbial partners ("screening" or selectively retaining beneficial symbionts), while microbes

may adapt to host environments, potentially leading to conflicts or mutualisms influencing

reproductive outcomes (59).

Third, the influence of the microbiome clearly extends into the female reproductive tract.

The finding that females preferentially stored sperm from males with more diverse

microbiomes (Figs. 1E and 2I) strongly suggests that microbiome-mediated enhancements to

ejaculate quality (sperm function and viability, possibly seminal fluid composition) interact

with female physiological processes. This opens a potential new dimension for cryptic female

choice (60, 61) with selection acting on microbiome-influenced male phenotypes.

The prevalence of synergistic effects also suggests that microbiome complexity and

functional integration might themselves be targets of selection, favouring males capable of

harbouring stable, high-performing microbial communities. Mechanistically, this could

involve selection for host factors that influence microbial colonization, resource availability

within the host environment, or the establishment of specific microbial interactions. Males with

genotypes promoting a harmonious and efficient microbial ecosystem would likely experience

enhanced physiological performance, ultimately translating to greater reproductive success.

Towards microbiome-integrated sexual selection

Realizing a comprehensive, microbiome-aware framework for sexual selection necessitates

several key avenues of future research. Foremost among these is the need to elucidate the

precise functional mechanisms underlying the observed fitness effects. This will require

integrated 'omics' approaches (e.g., metabolomics, transcriptomics) to map the links between

microbial communities, their metabolites, and key ejaculate features influencing fertilization

success. Alongside understanding the functional basis, it is also critical to determine how these

fitness-enhancing microbiotas are maintained within populations. Future studies should

therefore quantify the relative importance of different routes of transmission—including

vertical passage, coprophagy, social interactions, or seminal transfer—and assess the stability,

or heritability in a broad ecological sense, of beneficial microbial consortia across generations

and varying environmental conditions.

256   Furthermore, exploring the role of females in these interactions would be an important

avenue for future research. Investigations should examine female biases in both pre- and post-

mating choice relative to potential male microbial cues or the enhanced quality of ejaculates

produced by microbiome-rich males. A particularly crucial, yet unexplored, aspect is whether

the female's own resident microbiome interacts with the incoming ejaculate or associated

microbes, potentially influencing sperm storage patterns and biasing competition outcomes

independently of male traits.

263   Finally, the ecological context and broader applicability of our findings remains to be

explored. Research should examine how interactions between diet and the microbiome shape

male reproductive strategies and sperm competition dynamics across different resource

conditions. Assessing the generality of the microbiome-mediated effects on reproductive

fitness in other species, particularly those with diverse mating systems, diets, and modes of

microbial transmission, will be essential for understanding the widespread evolutionary

significance of host-microbiome interactions in sexual selection.

Conclusion

Our research demonstrates that the gut microbiome is an active participant in shaping male reproductive competitiveness. By influencing ejaculate quality and quantity, likely through modulation of host metabolism and resource allocation (6, 62, 63), these microbial symbionts redefine the landscape of condition dependence. Integrating the microbiome into sexual selection theory may fundamentally alter our understanding of how fitness variation is generated, maintained, and acted upon by selection in polyandrous systems and beyond. Our study underscores the importance of considering the hologenome—the host plus its associated microbiota—as the relevant unit of selection when investigating the evolution of complex reproductive traits.

Material and Methods

Experimental populations

Throughout this study, we used *Drosophila melanogaster* of the LH_m strain. These were either derived from an outbred wild-type population or, for standard competitors in the sperm competition assay, from a backcrossed population that had been genetically modified to ubiquitously express green fluorescent protein (GFP), permitting paternity assignment against the focal wild-type males (42, 43). To study biases in female sperm storage, we used focal males whose sperm heads expressed red fluorescent protein (RFP), and our standard competitors with ubiquitous GFP expression, permitting quantification of each competitor's sperm inside the female reproductive tract (42–44). We maintained all fly populations at moderate densities under constant conditions of 25°C, 60% humidity, and a 12h:12h light:dark cycle.

Generating axenic flies

To generate axenic flies, we performed all steps hereafter in a laminar flow chamber to avoid contamination. First, we collected eggs from Petri dishes containing grape-juice agar with a smear of yeast paste that had been placed in four large population cages after a six-hour egg-laying period. After washing these eggs off the plates into a strainer, we dechorionated them with 3.5% bleach for 2 minutes, followed by a rinse in 70% ethanol and three rinses with sterile water. We then transferred these eggs to autoclaved standard fly food supplemented with antibiotics [a cocktail of Ampicillin (50mg/mL), Kanamycin (50mg/mL, Erythromycin (10mg/mL), Tetracycline (10 mg/mL)] in autoclaved glass vials. These vials were then left undisturbed under standard conditions for flies to develop.

Upon fly eclosion, we collected both males and females as virgins, separated them by sex and confirmed their axenic state by plating a homogenized subsample of flies on LB and MRS agar followed by 16S PCR. Briefly, we extracted genomic DNA from 3−5 pooled axenic flies per sample using the ZymoBiomics^{TM} DNA Microprep kit following the instructions provided by the manufacturer. We also extracted DNA from flies with an intact microbiome to serve as positive controls. For DNA amplification, we used bacterial universal primers 27F and 1492R as forward and reverse primers, respectively. We prepared the PCR master mix using 9.5 μl of nuclease free water, 12.5 μl kappa HiFi master mix, 1 μl of forward and reverse primer and 1 μl of template DNA. The PCR cycle included an initial denaturation at 95°C for 90 s, followed by 35 cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s, an extension at 72°C for 90 s, and a final extension at 72°C for 5 mins. We ran the PCR products on a 1% agarose gel and visualized them under ultraviolet light after staining with SYBR Safe (Invitrogen) to confirm the absence of bacteria in axenic individuals and the presence of bacteria in controls.

321   We maintained all females and standard competitors as axenic flies throughout the

322   experiment and supplemented axenic focal males with different bacterial cultures to produce

323   gnotobiotic associations as described below.

Bacterial cultures and gnotobiotic associations

Throughout this study, we used five bacterial species isolated from the gut of wild-type flies

(20): *Acetobacter orientalis*, *A. tropicalis*, *A. pasteurianus*, *Lactobacillus plantarum*, and *L.*

brevis. We associated our focal flies with all 32 possible combinations among these bacterial

strains following the general protocol of Gould et al. (19). To prepare bacterial cultures, we

grew them overnight: *Lactobacillus* spp. in MRS (De Man–Rogosa–Sharpe) broth without

shaking and *Acetobacter* spp. in MRS + mannitol medium at 30°C on a shaker. The next

morning, we checked these bacterial cultures for growth and performed their optical density

(OD) measurements at 600 nm. Then, we spun down these different bacterial cultures at 7000

rpm for 5 minutes and resuspended the resulting pellets to a concentration of 10^8 cells/mL in

sterile phosphate-buffered saline (PBS). Resuspension volume was determined following

established protocols (64, 65). Finally, we added 50 µL (5×10^6 colony forming units, CFUs)

of bacterial culture to sterile food vials, with equal representation of bacterial strains for

combinations with more than one association. This bacterial density was comparable to prior

reports for conventional flies (19, 65).

After adding the bacterial culture to sterile food vials and air-drying these under in laminar

flow hood, we transferred 10 one-day-old axenic virgin males to each vial for inoculation with

the added bacterial strains. We moved these males to fresh food vials with same bacterial

treatment every alternate day for 5 days. Thereafter, we confirmed gnotobiotic associations by

plating together the homogenates of 3-4 adult flies from same treatments and checking their

colony morphologies, and by Sanger sequencing of 16S PCR homogenates. For multi-

associations, we plated the fly homogenates and picked the distinct types of colonies from a

given plate for PCR and performed sequencing to confirm that morphologically similar species

corresponds to same species in all the replicates, thus allowing us to distinguish between

different bacterial species by looking at their colony appearance. Thereafter, we tested the

presence of the expected bacterial species in a particular bacterial combination by plating them

and looking at the colony morphologies.

Bacterial abundance in different combinations

To determine bacterial abundance in different gnotobiotic associations, we counted the CFUs

per fly for each bacterial species across the different gnotobiotic associations. To this end, we

washed males with 70% ethanol, individually homogenized them with a pestle in

microcentrifuge tubes, and then diluted these homogenates to different dilutions with a final

volume of 100 μl to find a dilution with comparable colony counts (approx. 100 CFUs) for

visual quantification. In gnotobiotic cultures with both bacterial genera, we transferred an

aliquot of well-mixed fly homogenate to two plates, supplementing the MRS media of one

plate with 10 μg/ml ampicillin to selectively inhibit the growth of *Lactobacillus*, while

incubating the other in a CO_2 atmosphere to prevent the growth of *Acetobacter*. We achieved

within-genus differentiation of strains by colony morphologies: *A. orientalis* with flat colonies

with ruffled borders, *A. tropicalis* with small tan coloured and highly reflective colonies, and

A. pasteurianus with large tan colonies. Similarly, *L. brevis* produces brownish yellow colonies

and *L. plantarum* milky white colonies (Supplementary Fig. S1). Using 16S PCR amplification

followed by Sanger sequencing, we confirmed specificity of these colony properties,

demonstrating the robustness of our method based on morphological identifications during

experiments. After counting the number of colonies on each plate, we calculated CFU per fly

as CFU = C × D × V / P, where C = number of colonies counted, D = dilution factor, V =

volume of fly homogenate, and P = volume plated (64).

Across all 10 replicate plates combination examined, the expected species were present, and

none of the remaining species. Consistent with Gould et al. (19), total CFU counts increased

with species richness while the species-specific abundances declined (Supplementary Fig. S2).

Sperm competitive ability

To investigate sperm competitiveness of focal males, we quantified their sperm offense ability

(P_2) by mating them as the second males with axenic females that had each mated with a an

axenic GFP male three days earlier. We monitored the occurrence of copulations for 7-8 hours

and transferred remated females to oviposition vials while giving those that did not remate

another two 8-hour remating opportunities on the following days. Flies that did not remate by

the last opportunity (within these 3 days) were discarded. We allowed all remated females to

lay eggs for three days. After all progeny eclosed, we scored them for paternity under a

fluorescence stereoscope, based on the presence/absence of the ubiquitous GFP marker in

standard competitors. The proportion of second-male progeny indicated the focal-male sperm

competitive ability (i.e., sperm offense). We replicated this experiment thrice with 30 replicates

per bacterial treatment each time (total N = 2,880 trials).

Sperm viability

We quantified sperm viability for 10 males per bacterial treatment (total N = 320 males). Once

males were six days old and inoculated, we anaesthetized them on ice and dissected them on a

microscopic slide to isolate the seminal vesicles from their reproductive tissue. We then placed

one of the seminal vesicles in 10 µl of Ringers solution and punctured it with an insect pin to

release the sperm mass for immediate staining with LIVE/DEAD fluorescent dye, following

the principles of Tourmente et al. (41). This two-component dye consists of SYBR-14 ('LIVE'

component emitting green fluorescence), which is actively incorporated into living cells, and

propidium iodide (PI, with red fluorescence) that can only permeate damaged membranes

(typical of dead cells). We prepared an aqueous staining solution by combining 2.4 µL of

SYBR-14 stock solution (1 mM in DMSO) and 4.5 µl of PI stock solution (2.4 mM in water)

in 93.1 µl of Ringer's solution. Then, we added 5 µl of the staining solution to the sperm sample

before covering it with a coverslip and incubating it for 5 minutes in the dark. Under an

Olympus BX50 fluorescence microscope with a GFP/RFP dual filter, we captured images of

randomly chosen areas of each sample to later count at least 200 sperm and calculate the

proportion of viable sperm in ImageJ v. 1.34 (66).

Visualizing sperm competition within female reproductive tract

Since competitive fertilization events ultimately are the result of both male and female effects

(43, 67), we investigated how females bias their sperm storage based on interactions with males

supplemented with different microbiome. For this experiment, we again mated axenic virgin

females first with an axenic GFP male and, three days later, with a focal male as for sperm

competitive ability (above). To visualize sperm of both males, we used focal males expressing

RFP in their sperm heads. Immediately after remating, we transferred females to individual

wells of cell-culture plates, with a coverslip held in place with dots of rubber cement in two

corners. We checked these females every 10 min for up to 6 h for the ejection of the sperm

mass containing both displaced first-male and excess second-male sperm (42, 44). The timing

and extent of sperm ejection have previously been reported to affect relative numbers of first-

and second-male sperm stored by females and, consequently, competitive fertilization success

among males (43).

Upon female sperm ejection, we recorded the interval from mating to ejection before removing

females from the well and freezing them individually in Eppendorf tubes. Thereafter, we

dissected these females and counted the green (standard competitors) and red sperm (focal

males) within the female reproductive tract (paired spermathecae, seminal receptacle and bursa

copulatrix) under a fluorescence microscope. From these counts, we determined S_2, the

proportion of all sperm stored by the female that came from the second (experimental) male.

We also counted the respective sperm numbers in the ejected sperm mass to estimate, combined

with above counts, the total number of sperm transferred by the second male and the number

of first-male sperm still residing in storage at the time of remating. This experiment was

performed in five blocks with 10 replicates per treatment for each block (total $N = 1,600$ trials).

Statistical analyses

We estimated the contributions of microbial interactions to variation in reproductive traits

using mixed-effects models with the five bacterial species and all two- to five-way interactions

as predictors, while accounting for experimental block as a random effect. Models were either

linear (for number of sperm transferred, time to sperm ejection) or generalized with a binomial

error structure (for proportions of viable sperm or of second-male sperm in storage). For the

proportion of second-male offspring, we chose a linear model on logit-transformed P_2 data

instead of a binomial model because the latter showed poor performance, also when adjusting

for dispersion or residual biases by using betabinomial or Tweedie link functions instead of the

default 'binomial' function. Additionally, we quantified the effect of species richness on trait

expression in models using the number of bacterial species per treatment as the fixed effect,

and block, the treatment group and (where applicable) observation as random effects. Here, the

binomial model on P_2 performed well and was preferred.

To better understand the importance of species combinations in the microbiome and to isolate

emergent effects beyond individual species contributions, we used an epistasis-based

framework to elucidate how well the averaged phenotypic effects of single- or pairwise species

associations predict higher-order phenotypes (19). For single-species predictions, we compared

the measured trait values of males associated with two- to five-species microbiotas to values

averaged across mono-associated males that jointly represented the same bacterial

combinations. These differences (averaged prediction minus measured trait value) were then

plotted with bootstrapped 95% confidence intervals to visually determine if predictions

deviated from the measured estimates (i.e., confidence interval excluding zero difference). We

then repeated this procedure to predict phenotypes by averaging the estimates of di-associated

males (for calculations see (19)).

Following the same general principle, we further quantified how the presence or absence of

non-focal ("bystander") community members modulates both pairwise and three-way

microbial interactions by exhaustively enumerating all possible bystander contexts. We then

compared this "contextual" analysis against a "standard" test to reveal how higher-order

community background shapes emergent effects on host traits. Following Gould et al.'s (19)

nomenclature, the "standard" test calculated the interaction coefficient among all species

present in a specific community configuration, whereas the "contextual" tests assessed the

interaction between a focal species pair within the context of additional bystander species. We

computed interaction coefficients for each focal species pair or triplet across all possible

presence/absence combinations of bystander species and plotted them with bootstrapped

standard errors to highlight the varying directions of interactions depending on the bystander

combinations.

In both the single/double-species predictions and these standard/contextual tests, we derived

the coefficients to calculate the deviations from a mixed-effects models with an appropriate

link function for each microbial combination, controlling for block effects. We repeated each

model across 1,000 bootstrapped datasets to calculate the standard errors or 95% confidence

intervals from the distribution of the calculated coefficient or interaction differentials.

Funding

This project was supported by the Swiss National Science Foundation (grant PP00P3_202662

to S.L.).

References

1.   J. L. Fitzpatrick, S. Lüpold, Sexual selection and the evolution of sperm quality. *Mol.*
Hum. Reprod. **20**, 1180–1189 (2014).

2.   J. L. Fitzpatrick, L. W. Simmons, J. P. Evans, Complex patterns of multivariate
selection on the ejaculate of a broadcast spawning marine invertebrate. *Evolution* **66**,
2451–2460 (2012).

3.   D. A. Dewsbury, Ejaculate Cost and Male Choice. *Am. Nat.* **119**, 601–610 (1982).

4.   J. L. Godwin, *et al.*, Experimental evolution reveals that sperm competition intensity
selects for longer, more costly sperm. *Evol. Lett.* **1**, 102–113 (2017).

5.   S. Lüpold, *et al.*, How sexual selection can drive the evolution of costly sperm
ornamentation. *Nature* **533**, 535–538 (2016).

6.   E. L. Macartney, A. J. Crean, S. Nakagawa, R. Bonduriansky, Effects of nutrient
limitation on sperm and seminal fluid: a systematic review and meta-analysis. *Biol. Rev.*
94, 1722–1739 (2019).

7.   M. Olsson, T. Madsen, R. Shine, Is sperm really so cheap? Costs of reproduction in
male adders, Vipera berus. *Proc. R. Soc. Lond. B Biol. Sci.* **264**, 455–459 (1997).

8.   J. C. Perry, C. T. Tse, Extreme costs of mating for male two-spot ladybird beetles. *Plos*
One **8**, e81934 (2013).

9.   Y. Iwasa, A. Pomiankowski, S. Nee, The evolution of costly mate preferences ii. The
"handicap" principle. *Evolution* **45**, 1431–1442 (1991).

10.   A. Grafen, Sexual selection unhandicapped by the fisher process. *J. Theor. Biol.* **144**,
473–516 (1990).

11.   G. E. Hill, Condition-dependent traits as signals of the functionality of vital cellular
processes. *Ecol. Lett.* **14**, 625–634 (2011).

12.   L. Rowe, D. Houle, The lek paradox and the capture of genetic variance by condition
dependent traits. *Proc. R. Soc. Lond. B Biol. Sci.* **263**, 1415–1421 (1996).

13.   S. O. Fetissov, Role of the gut microbiota in host appetite control: bacterial growth to
animal feeding behaviour. *Nat. Rev. Endocrinol.* **13**, 11–25 (2017).

14.   J. Alcock, C. C. Maley, C. A. Aktipis, Is eating behavior manipulated by the
gastrointestinal microbiota? Evolutionary pressures and potential mechanisms.
BioEssays **36**, 940–949 (2014).

15.   A. E. Douglas, The microbial dimension in insect nutritional ecology. *Funct. Ecol.* **23**,
38–47 (2009).

16.   V. Emelianoff, *et al.*, A survival-reproduction trade-off in entomopathogenic nematodes
mediated by their bacterial symbionts. *Evolution* **62**, 932–942 (2008).

17.   A. W. Walters, *et al.*, The microbiota influences the *Drosophila melanogaster* life
history strategy. *Mol. Ecol.* **29**, 639–653 (2020).

18.   M. K. Matthews, J. Malcolm, J. M. Chaston, Microbiota influences fitness and timing of
reproduction in the fruit fly *Drosophila melanogaster*. *Microbiol. Spectr.* **9**, e00034-21
(2021).

19.   A. L. Gould, *et al.*, Microbiome interactions shape host fitness. *Proc. Natl. Acad. Sci.*
115 (2018).

20.   J. M. Baker, L. Al-Nakkash, M. M. Herbst-Kralovetz, Estrogen–gut microbiome axis:
Physiological and clinical implications. *Maturitas* **103**, 45–53 (2017).

21.   D. J. Morrison, T. Preston, Formation of short chain fatty acids by the gut microbiota
521   and their impact on human metabolism. *Gut Microbes* **7**, 189–200 (2016).

522   22.   M. Insenser, *et al.*, Gut microbiota and the polycystic ovary syndrome: influence of sex,
523   sex hormones, and obesity. *J. Clin. Endocrinol. Metab.* **103**, 2552–2562 (2018).

524   23.   F. Zhang, D. Aschenbrenner, J. Y. Yoo, T. Zuo, The gut mycobiome in health, disease,
525   and clinical applications in association with the gut bacterial microbiome assembly.
526   *Lancet Microbe* **3**, e969–e983 (2022).

527   24.   J. M. Fettweis, *et al.*, The vaginal microbiome and preterm birth. *Nat. Med.* **25**, 1012–
528   1021 (2019).

529   25.   F. Pinto, D. A. Medina, J. R. Pérez-Correa, D. Garrido, Modeling metabolic interactions
530   in a consortium of the infant gut microbiome. *Front. Microbiol.* **8**, 2507 (2017).

531   26.   K. Tremellen, K. Pearce, Dysbiosis of Gut Microbiota (DOGMA)–a novel theory for the
532   development of Polycystic Ovarian Syndrome. *Med. Hypotheses* **79**, 104–112 (2012).

533   27.   J. M. Franasiak, R. T. Scott Jr, Reproductive tract microbiome in assisted reproductive
534   technologies. *Fertil. Steril.* **104**, 1364–1371 (2015).

535   28.   K. A. Green, S. M. Zarek, W. H. Catherino, Gynecologic health and disease in relation
536   to the microbiome of the female reproductive tract. *Fertil. Steril.* **104**, 1351–1357
537   (2015).

538   29.   A. E. Douglas, Fundamentals of microbiome science: how microbes shape animal
539   biology (Princeton University Press, 2018).

540   30.   P. Engel, N. A. Moran, The gut microbiota of insects–diversity in structure and function.
541   *FEMS Microbiol. Rev.* **37**, 699–735 (2013).

542   31.   E. V. Ridley, A. C. Wong, A. E. Douglas, Microbe-dependent and nonspecific effects of
543   procedures to eliminate the resident microbiota from *Drosophila melanogaster*. *Appl.*
544   *Environ. Microbiol.* **79**, 3209–3214 (2013).

545   32.   A. C. Wong, J. M. Chaston, A. E. Douglas, The inconstant gut microbiota of *Drosophila*
546   species revealed by 16S rRNA gene analysis. *ISME J.* **7**, 1922–1932 (2013).

547   33.   M. Elgart, *et al.*, Impact of gut microbiota on the fly's germ line. *Nat. Commun.* **7**,
548   11280 (2016).

549   34.   S. Y. Delbare, Y. H. Ahmed-Braimah, M. F. Wolfner, A. G. Clark, Interactions between
550   the microbiome and mating influence the female's transcriptional profile in *Drosophila*
551   *melanogaster. Sci. Rep.* **10**, 18168 (2020).

552   35.   M. Al-Asmakh, *et al.*, The gut microbiota and developmental programming of the testis
553   in mice. *PLoS ONE* **9**, e103809 (2014).

36. N. Ding, *et al.*, Impairment of spermatogenesis and sperm motility by the high-fat diet-
induced dysbiosis of gut microbes. *Gut* **69**, 1608–1619 (2020).

37. M. Alfano, *et al.*, Testicular microbiome in azoospermic men—first evidence of the
impact of an altered microenvironment. *Hum. Reprod.* **33**, 1212–1217 (2018).

38. S.-L. Weng, *et al.*, Bacterial communities in semen from men of infertile couples:
metagenomic sequencing reveals relationships of seminal microbiota to semen quality.
PloS One **9**, e110152 (2014).

39. A. E. Douglas, The microbial dimension in insect nutritional ecology. *Funct. Ecol.* **23**,
38–47 (2009).

40. J. G. McMullen, G. Peters-Schulze, J. Cai, A. D. Patterson, A. E. Douglas, How gut
microbiome interactions affect nutritional traits of *Drosophila melanogaster*. *J. Exp.*
Biol. **223**, jeb227843 (2020).

41. M. Tourmente, C. R. Archer, D. J. Hosken, Complex interactions between sperm
viability and female fertility. *Sci. Rep.* **9**, 15366 (2019).

42. S. Lüpold, *et al.*, How Multivariate Ejaculate Traits Determine Competitive Fertilization
Success in *Drosophila melanogaster*. *Curr. Biol.* **22**, 1667–1672 (2012).

43. S. Lüpold, *et al.*, Female mediation of competitive fertilization success in *Drosophila*
melanogaster. *Proc. Natl. Acad. Sci.* **110**, 10693–10698 (2013).

44. M. K. Manier, *et al.*, Resolving mechanisms of competitive fertilization success in
Drosophila melanogaster. *science* **328**, 354–357 (2010).

45. S. Lüpold, *et al.*, How female× male and male× male interactions influence competitive
fertilization in *Drosophila melanogaster*. *Evol. Lett.* **4**, 416–429 (2020).

46. M. Tourmente, E. R. Roldan, Mass-specific metabolic rate influences sperm
performance through energy production in mammals. *PLoS One* **10**, e0138185 (2015).

47. M. Rowe, L. Veerus, P. Trosvik, A. Buckling, T. Pizzari, The Reproductive
Microbiome: An Emerging Driver of Sexual Selection, Sexual Conflict, Mating
Systems, and Reproductive Isolation. *Trends Ecol. Evol.* **35**, 220–234 (2020).

48. A. C.-N. Wong, A. J. Dobson, A. E. Douglas, Gut microbiota dictates the metabolic
response of *Drosophila* to diet. *J. Exp. Biol.* **217**, 1894–1901 (2014).

49. J. delBarco-Trillo, M. Tourmente, E. R. Roldan, Metabolic rate limits the effect of
sperm competition on mammalian spermatogenesis. *PLoS One* **8**, e76510 (2013).

50. Z. B. Hamad Zubi, H. A. Hamad Alfarisi, Hyperlipidemia and male infertility. *Egypt. J.*
Basic Appl. Sci. **8**, 385–396 (2021).

51. C. C. Maresch, *et al.*, Diabetes-induced hyperglycemia impairs male reproductive
function: a systematic review. *Hum. Reprod. Update* **24**, 86–105 (2018).

52. S. Subhadarshini, V. Mallesh, G. Bharath, Semen quality alterations in men with
hyperlipidemia and diabetes mellitus: a cross-sectional study. *Int J Acad Med Pharm* **6**,
639–644 (2024).

53. X. Zeng, L. Tong, The Impact of Diabetes on Male Silkworm Reproductive Health.
Biology **13**, 557 (2024).

54. S. C. Shin, *et al.*, *Drosophila* microbiome modulates host developmental and metabolic
homeostasis via insulin signaling. *Science* **334**, 670–674 (2011).

55. I. Iatsenko, J.-P. Boquete, B. Lemaitre, Microbiota-derived lactate activates production
of reactive oxygen species by the intestinal NADPH oxidase Nox and shortens
Drosophila lifespan. *Immunity* **49**, 929–942 (2018).

56. G. Storelli, *et al.*, *Lactobacillus plantarum* promotes *Drosophila* systemic growth by
modulating hormonal signals through TOR-dependent nutrient sensing. *Cell Metab.* **14**,
403–414 (2011).

57. A. Zelezniak, *et al.*, Metabolic dependencies drive species co-occurrence in diverse
microbial communities. *Proc. Natl. Acad. Sci.* **112**, 6449–6454 (2015).

58. L. P. Henry, M. Bruijning, S. K. Forsberg, J. F. Ayroles, The microbiome extends host
evolutionary potential. *Nat. Commun.* **12**, 5141 (2021).

59. N. A. Moran, H. Ochman, T. J. Hammer, Evolutionary and ecological consequences of
gut microbial communities. *Annu. Rev. Ecol. Evol. Syst.* **50**, 451–475 (2019).

60. R. C. Firman, C. Gasparini, M. K. Manier, T. Pizzari, Postmating female control: 20
years of cryptic female choice. *Trends Ecol. Evol.* **32**, 368–382 (2017).

61. L. Daupagne, *et al.*, One size does not fit all: female–male interactions on the path to
fertilization. (2025). https://doi.org/10.1530/REP-24-0238.

62. E. L. Macartney, *et al.*, Sperm depletion in relation to developmental nutrition and
genotype in *Drosophila melanogaster*. *Evolution* **75**, 2830–2841 (2021).

63. V. Zeender, S. Pfammatter, B. Roschitzki, S. Dorus, S. Lüpold, Genotype-by-
environment interactions influence the composition of the *Drosophila* seminal
proteome. *Proc. R. Soc. B Biol. Sci.* **290**, 20231313 (2023).

64. M. L. Koyle, *et al.*, Rearing the fruit fly *Drosophila melanogaster* under axenic and
gnotobiotic conditions. *JoVE J. Vis. Exp.* e54219 (2016).

65. P. D. Newell, A. E. Douglas, Interspecies interactions determine the impact of the gut

microbiota on nutrient allocation in *Drosophila melanogaster*. *Appl. Environ.*

Microbiol. **80**, 788–796 (2014).

66. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of

image analysis. *Nat. Methods* **9**, 671–675 (2012).

67. W. Eberhard, Female control: sexual selection by cryptic female choice (Princeton

University Press, 1996).

Figures

A   B   C   D   E

Fig. 1. Associations between the number of bacterial species and the different fitness traits.

Each point indicates the mean of all replicates for a given microbiome consortium (*N* = 32).

The lines and 5% confidence bands are derived from (generalized) linear mixed-effects models

with microbial species composition and block as random effects (for binomial models

additionally also an observation-level random effect to address overdispersion).

Fig. 2. Variation in male fitness traits relative to their microbial community (left) and error in predicting these values for males associated with multiple species based on single- or pairwise species averages (right). In the left panels, colours indicate the different combinations of

Acetobacter and *Lactobacillus* species associated with experimental males. The right panels

depict the deviations in trait predictions from measured traits by averaging the corresponding

coefficients of (generalized) linear mixed-effects models for individual species (orange points)

or species pairs (green diamonds) that jointly represented the same species communities. The

relevant microbial combinations are plotted in the same order across all panels. Error bars

reflect 95% confidence intervals (from raw distributions in left panels, from bootstrapped

prediction differentials in right panels).

Fig. 3. Correlations between the different phenotypic traits and the relative abundance (based

on CFU counts) of each of the five bacterial species (left) and among the traits themselves

(right). Values reflect the averaged correlation coefficients with 95% confidence limits of

pairwise Pearson correlations across bootstrapped group means ($N = 1,000$ samples) among

males with a given microbiome consortium. Relative abundances and proportional traits (sperm

viability, S_2, P_2) were logit-transformed before bootstrapping. Colours indicate the strength and

direction of correlations as reflected by the colour-gradient bar.

Fig. 4. Interactive effects between bacterial species contingent on bystander species, with two-way interactions between pairs of focal of species in the left panels, and three-way interactions between trios of focal species in the right panels. For each code along the x-axes, 1's indicate the species configuration of the interaction and asterisks denote placeholders for bystander species (order within codes from left to right: *A. orientalis*, *A. tropicalis*, *A. pasteurianus*, *L.*

brevis, *L. plantarum*). The binary codes in the legends indicate the configuration of bystanders

in the order, in which they would fill the placeholders (*) in each focal species configuration

(1 = presence, 0 = absence). The right panels distinguish between the "standard" tests (grey

triangles; i.e., global interaction across all community members present) and "contextual" tests

(coloured points; i.e., separate interactions between a focal species pair or triplet for each

bystander combination) for each focal triplet. All coefficients and standard errors were derived

from (generalized) linear mixed-effects models across 1,000 bootstrapped datasets.