Male gut microbiome mediates post-mating sexual selection in *Drosophila*

Komal Maggu*, Abhishek Meena, Alessio N. De Nardo, Sonja H. Sbilordo, Jeannine Roy &

Stefan Lüpold*

Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, 8057 Zurich, Switzerland

Correspondence: komal024maggu@gmail.com, stefan.luepold@uzh.ch

ABSTRACT

In polyandrous systems, male reproductive success is critically influenced by sperm

competition, a process often mediated by ejaculate characteristics that are sensitive to male

condition. While condition dependence is a cornerstone of sexual selection theory, the role of

the gut microbiome in modulating male reproductive fitness under sperm competition remains

unexplored. Here, we investigated the impact of varying gut microbial compositions on sperm

competitive ability in *Drosophila melanogaster*. By inoculating germ-free (axenic) males with

all 32 full-factorial combinations of five common commensal bacteria, we demonstrated that

the presence and diversity of commensal gut microbes significantly enhance male sperm

offense ability. Specifically, males associated with a more diverse microbiota exhibited

superior competitive fertilization success, accompanied by increased sperm transfer, higher

sperm viability, and a greater proportion of sperm stored by females. Predictive models further

revealed complex, non-additive interactions among bacterial species in their effects on sperm

competitiveness. These findings suggest that the gut microbiome acts as a critical modulator

of male reproductive fitness, challenging traditional models of sexual selection that overlook

contributions of non-pathogenic bacteria to condition-dependent traits.

INTRODUCTION

In polyandrous species, where females mate with multiple males, sperm competition often

determines fertilization success based on ejaculate size and quality (1, 2). However, producing

large, high-quality ejaculates is costly by diverting resources from other fitness traits (3–7).

The dependence of trait expression on resource availability and allocation between growth,

maintenance, and reproduction is central to sexual selection theory, linking phenotypic quality

to underlying genetic variation: traits that reliably indicate a male's ability to acquire and

allocate limited resources are hypothesized to signal heritable fitness benefits to mates or rivals

(8, 9). This 'good genes' framework assumes that male condition is shaped by genetic variation

in resource acquisition, allocation efficiency, or stress tolerance (10, 11). However, the gut

microbiome – a dynamic microbial consortium known to modulate host nutrient extraction,

metabolic regulation, and hormone synthesis (12, 13) – represents a potent, yet unexplored,

factor that could decouple male condition from host genetics. Critically, by independently

modulating host resource budgets, microbial communities could fundamentally alter the

heritability of condition-dependent traits, thereby challenging a cornerstone of sexual selection

theory.

The gut microbiome directly modulates host physiology, including nutrient assimilation

(14–16), energy allocation (17–19), and hormone signalling (20), which collectively can

influence reproductive function. Indeed, across diverse taxa, microbiome composition is well-

documented to shape female reproductive biology (17, 21, 22), with functional differences

between key bacterial groups (e.g., *Acetobacter* vs. *Lactobacillus* in *Drosophila*) influencing

metabolic and fecundity outcomes (17, 23). By contrast, the role of the gut microbiome in male

reproductive fitness, particularly its impact on fertilization success under sperm competition,

remains entirely unexplored, despite limited associations between dysbiosis and impaired

testicular function(24, 25) or between seminal microbes to sperm motility (26, 27).

Here, we tested the hypothesis that the gut microbiome modulates male reproductive

fitness under competitive fertilization by enhancing condition-dependent ejaculate traits in

Drosophila melanogaster, a model for tractable host-microbe interactions (28, 29).

Specifically, we inoculated males with all 32 full-factorial combinations of five gut bacteria

(*Acetobacter orientalis*, *A. tropicalis*, *A. pasteurianus*, *Lactobacillus brevis*, and *L. plantarum*).

When present together, these species recapitulate key phenotypic traits of flies with intact

microbiomes (Supplementary Fig. 1) and represent bacteria with established roles in female

fitness (17). We then quantified competitive fertilization success (P_2) and its mechanistic

underpinnings: sperm viability, transfer, and storage. We predicted that (i) males with diverse

microbiomes outperform axenic (i.e., germ-free) or low-diversity males due to improved

nutrient acquisition (14, 30), (ii), competitive advantages are amplified by bacterial identity

(e.g., *Acetobacter* abundance), synergistic interactions, and community composition (e.g.,

Acetobacter-dominance) (17), and (iii) microbiome-driven gains in sperm viability, transfer

efficiency and competitive storage underpin fertilization success (31, 32). By resolving how

microbial communities shape sperm competition, this study bridges microbial ecology with

sexual selection theory, offering novel insights into the evolution of condition-dependent traits.

RESULTS AND DISCUSSION

Microbial diversity enhances sperm competitiveness

To test if gut microbes influence male reproductive fitness under post-mating sexual selection,

we compared the competitive fertilization success of male *D. melanogaster*, inoculated with

gut bacteria in all possible combinations between our five strains, using a full-factorial

combinatorial approach (32 consortia) validated through colony morphology and 16S

sequencing (Methods, Supplementary Figs. S1 and S2). This combinatorial design—building

on established gnotobiotic methodologies (17) but applied here to sperm competition—enabled

high-resolution dissection of species interactions mediating host phenotypes. Standard

competitor males (either axenic or conventional with intact microbiomes), which ubiquitously

expressed green fluorescent protein (GFP) for paternity assignment (32, 33), mated three days

before focal wild-type males, thereby providing distinct competitive contexts to resolve

intrinsic microbiome benefits.

In competition against axenic rivals, we found experimental males harbouring any single

bacterial species (except *L. brevis*) or consortium to outperform males without a microbiome,

with second-male paternity (P_2) increasing with species richness (generalized linear mixed

models (GLMM) repeated across 1,000 bootstrapped datasets, accounting for block, microbial

combination and overdispersion: $N = 646$ across 32 microbial combinations: odds ratio, $OR =$

1.09 [95% confidence limits: 1.00, 0.15]; Fig. 1A). Being associated with all five species

translated into a 27% gain in P_2 compared to the axenic state (mean proportion ± s.d. = 0.85 ±

0.05 vs. 0.67 ± 0.05, respectively). These patterns persisted when focal males competed against

rivals with intact (unmanipulated) microbiotas [increase with richness: $N = 624$, $OR = 1.05$

[1.00, 1.10]; relative gain for 5-species association: 20% (0.72 ± 0.08 vs. 0.60 ± 0.03)],

confirming that microbiome benefits reflect intrinsic physiological enhancement rather than

competitor compromise.

This reproductive deficit in axenic males aligns with established connections between

microbiome absence and physiological dysregulation in *Drosophila* (34). Such microbial

modulation of condition dependence likely extends beyond *Drosophila*, as gut microbiomes

influence nutrient assimilation across the animal tree of life. While our study did not directly

assay host metabolic states, prior work compellingly shows that axenic flies often exhibit

hyperlipidemia and impaired insulin signalling – conditions detrimental to overall health and

potentially sperm function (35, 36). These metabolic disturbances can be rescued or modulated

by specific gut bacteria; for example, *Acetobacter* species are known to restore lipid

homeostasis, potentially via acetate production (37) (which may protect against sperm

oxidative damage), whilst certain *Lactobacillus* species exacerbate oxidative stress (38), which

can directly compromise sperm membranes and DNA integrity (39). In fact, *Acetobacter*-

dominated communities conferred greater benefits than those biased toward *Lactobacillus*,

regardless of the standard competitor (Fig. 2A). The contrasting impacts of *Acetobacter* and

Lactobacillus on P_2 based on their presence and absence were further supported by

bootstrapped correlations between P_2 and the relative abundances (based on CFU counts) of

each species across different microbial communities. These correlations yielded consistently

positive and negative covariation with *Acetobacter* and *Lactobacillus* species, respectively,

indicating a more quantitative pattern than sole presence or absence (Fig. 3A).

Combined, although our results establish that microbial diversity and particularly

Acetobacter dominance directly enhance male competitive fertilization success, challenging

classical frameworks attributing condition dependence solely to host genetics and resource

status or stress resilience (11).

Microbial interactions shape competitive outcomes

Bootstrapped generalized linear mixed-effects models (GLMMs), controlling for experimental

block, indicated that microbial interactions—rather than additive effects—were the primary

drivers of variation in P_2 across the 32 microbial consortia. There was no strong evidence for a

five-way interaction among bacterial strains in sperm competition against either axenic males

($N = 646$, $OR = 0.82$ [0.55, 2.66]) or conventional rivals ($N = 624$, $OR = 0.76$ [0.40, 1.39]).

However, after excluding these highest-order interactions, there was a clear four-way

interaction among $At \times Ap \times Lb \times Lp$ in competitions against axenic males (OR = 1.84 [1.22,

2.77]) and among $Ao \times Ap \times Lb \times Lp$ in those against conventional males (OR = 0.67 [0.49,

0.93]), suggesting complex microbial effects on reproductive outcomes. Additionally, two and

four three-way interactions, as well as six and eight of the ten two-way interactions had

bootstrapped confidence intervals that excluded zero between the two models, respectively,

further supporting a non-additive contribution of microbial composition (Supplementary Figs.

S2 and S3).

Predictive modelling showed that averaging mono-association effects between bacterial

strains poorly forecasted outcomes for multi-species communities (e.g., mean of At, Ap, and

Lb to predict the At-Ap-Lb combination), with only eight (30.8%) of the 26 possible

combinations correctly predicted based on 95% confidence intervals including zero difference

in competitive matings against axenic rivals (Fig. 2B). Averaging pairwise combinations to

predict three-way species interactions (e.g., At-Ap, At-Lb and Ap-Lb to predict At-Ap-Lb)

indicated similar accuracy (31.2%: 5 of 16 predicted combinations; Fig. 2B). These predictions

by mono- or di-associations were broadly comparable when focal males competed against

conventional rather than axenic first males (33.0 and 37.5%, respectively).

Additionally, pairwise interactions were context-dependent, varying conditionally on the

remaining ("bystander") species as reflected by interactions trending from positive

(synergistic) to negative (antagonistic) between the same two species depending on the

configuration of additional species in the microbiota (Fig. 4A). This dependence on bystanders

also held for three-way interactions between focal species (Fig. 4B). The prevalence of these

non-additive effects reveals that microbial consortia function as integrated units—not mere

sums of parts—mirroring ecological dynamics in host-microbe systems (40). Such complexity

suggests microbiome stability itself could be a target of sexual selection, with selection

favouring host genotypes that stabilize high-performance communities (e.g., *Acetobacter*

dominance), potentially explaining a microbiome-mediated resolution of the lek paradox (41).

Mechanistic basis of microbiome-mediated fitness

To dissect how microbiomes enhance competitive fertilization success, we quantified sperm

transfer, sperm viability, and female sperm retention and storage patterns, based on prior

knowledge (31–33, 42). Due to the similar results between axenic and conventional competitor

males, we here restricted our mechanistic investigation to axenic first males. For visualization

of sperm, all assays except sperm viability involved standard and experimental males that,

respectively, expressed green or red fluorescent protein in sperm heads (43).

We found that a more diverse microbiome directly translates into improved ejaculate

quality, providing the mechanistic basis for enhanced competitive fertilization success (Fig. 1).

While axenic and mono-associated focal males transferred similar sperm quantities (Figs. 1B

and 2C), multi-associated males transferred 3–36% more sperm than either of the former

categories, with gains scaling with microbial diversity (bootstrapped linear mixed model

(LMM) controlling for block and microbial combination: $N = 582$, $\beta = 57.14$ [40.68, 76.47];

Figs. 1B and 2C,D). Sperm viability also increased with diversity: males with all five species

had $94.9 \pm 2.3\%$ viable sperm compared to $54.5 \pm 6.5\%$ in axenic males (GLMM: $N = 320$, *OR*

$= 1.62$ [1.52, 1.71]; Figs. 1C and 2E, F). Critically, freshly remated females tended to delay the

ejection of displaced and excess sperm from first (standard) and second (experimental) males,

respectively, if second males were associated with a more diverse microbiome (LMM: $N =$

582, $\beta = 3.73$ [-0.09, 7.39; 90%CI: 0.45, 6.90]; Figs. 1D and 2G,H), thereby increasingly

biasing sperm storage toward these males (GLMM on relative sperm numbers in storage: $N =$

582, *OR* $= 1.40$ [1.34, 1.46]; Figs. 1E, 2I,J). Hence, microbiome-enhanced sperm quality or

displacement efficiency biased post-mating sperm storage dynamics. Confirming prior work

(31–33, 42), the relative numbers of sperm stored and their viability covaried with paternity

outcomes (Fig. 3B). Further, these phenotypic traits again covaried mostly positively with

Acetobacter and negatively with *Lactobacillus* when considering relative colony forming units

(CFUs) rather than species presence or absence (Fig. 3A).

Similar to P_2, the component traits underlying competitive fertilization success were

shaped by higher-order microbial interactions. Bootstrapped GLMMs revealed significant five-

way interactions for both sperm viability ($N = 320$, $OR = 0.29$ [0.13, 0.63], Supplementary Fig.

S4) and the proportion of second-male sperm stored by females ($N = 582$, $OR = 2.34$ [1.38,

4.11], Supplementary Fig. S5). We detected a comparable five-way interaction in an LMM for

time to female sperm ejection ($N = 582$, $\beta = 150.10$ [41.74, 255.22], Supplementary Fig. S6).

While no five-way interaction was found for the number of sperm transferred, a consistent

four-way interaction among $Ao \times Ap \times Lb \times Lp$ emerged ($N = 582$, $\beta = 393.09$ [198.49,

582.51], Supplementary Fig. S7), with nearly all two- and three-way interactions also showing

effect distributions shifted away from zero. Predictive models also indicated considerable

underestimation of observed values for sperm viability and relative sperm storage when using

additive models based on mono- or di-associations (Fig. 2F,J), highlighting the importance of

higher-order interactive effects. This dependence on microbial context appeared weaker for

both the number of sperm transferred and time to sperm ejection. Across all stages of the

competitive fertilization process, interactions among microbial pairs or triplets were often

modulated by the presence of additional "bystander" species, emphasizing the context-

dependent nature of microbial effects (Fig. 4C–J).

Evolutionary implications

The observed improvements in ejaculate quality and subsequent P_2 gains conferred by species-

rich microbial communities likely stem from their ability to restore or optimize host metabolic

homeostasis and nutrient allocation toward reproduction. The superior performance associated

with *Acetobacter*-dominated communities (Fig. 1) lends support to this hypothesis, given their

known efficacy in nutrient assimilation and influencing host metabolism (34). These microbes

may channel resources more effectively towards energetically costly ejaculate production and

maintenance, potentially mitigating damage from metabolic byproducts. However, the

significant non-additive effects underscore that the *integrated function* of the microbial

consortium, rather than simply the presence of key species, dictates the ultimate physiological

impact (Fig. 3). Although metabolite-specific mechanisms await future study, our reductionist

approach definitively isolates microbiome contributions to sperm competitiveness.

Our findings prompt a re-evaluation of core tenets in sexual selection theory. By acting

as an amplifier of condition-dependent reproductive traits, the microbiome introduces

complexity to the "good genes" model, where such traits are assumed to reliably signal host

genetic quality (10, 11). Our results suggest that the microbiome functions as a "hidden

mediator," bridging host genetics and environmental inputs (44), with at least three key

evolutionary implications.

First, the microbiome potentially introduces a transmissible component influencing

fitness that is sensitive to environmental factors (diet, social context) and operates outside strict

Mendelian inheritance (44). If microbial consortia are consistently acquired or partially

transmitted (vertically or horizontally), this could contribute to maintaining heritable variation

in sexually selected traits, offering a potential factor in resolving the lek paradox (41).

Furthermore, the microbiome might act as an "environmental sensor," allowing males to buffer

reproductive output against nutritional fluctuations (7), thereby stabilizing their fitness.

Second, the context-dependent microbial interactions (Fig. 3) and potential trade-offs

(e.g., benefits vs. costs like oxidative stress potentially associated with *L. brevis*) suggest

complex coevolutionary dynamics. Selection may favour host mechanisms for managing

microbial partners ("screening" or selectively retaining beneficial symbionts), while microbes may adapt to host environments, potentially leading to conflicts or mutualisms influencing reproductive outcomes(45).

Third, the influence of the microbiome clearly extends into the female reproductive tract. The finding that females preferentially stored sperm from males with more diverse microbiomes (Figs. 1E and 2I) strongly suggests that microbiome-mediated enhancements to ejaculate quality (sperm function and viability, possibly seminal fluid composition) interact with female physiological processes. This opens a potential new dimension for cryptic female choice (46, 47) with selection acting on microbiome-influenced male phenotypes. Critically, while our *Drosophila* model provides mechanistic tractability, microbiome-mediated condition dependence likely operates universally where sperm competition occurs.

The prevalence of synergistic effects also suggests that microbiome complexity and functional integration might themselves be targets of selection, favouring males capable of harbouring stable, high-performing microbial communities. Mechanistically, this could involve selection for host factors that influence microbial colonization, resource availability within the host environment, or the establishment of specific microbial interactions. Males with genotypes promoting a harmonious and efficient microbial ecosystem would likely experience enhanced physiological performance, ultimately translating to greater reproductive success.

Towards microbiome-integrated sexual selection

Realizing a comprehensive, microbiome-aware framework for sexual selection necessitates several key avenues of future research. Foremost among these is the need to elucidate the precise functional mechanisms underlying the observed fitness effects. This will require integrated 'omics' approaches (e.g., metabolomics, proteomics, transcriptomics) to map the links between microbial communities, their metabolites, and key ejaculate features influencing

fertilization success. Alongside understanding the functional basis, it is also critical to determine how these fitness-enhancing microbiotas are maintained within populations. Future studies should therefore quantify the relative importance of different routes of transmission—including vertical passage, coprophagy, social interactions, or seminal transfer—and assess the stability, or heritability in a broad ecological sense, of beneficial microbial consortia across generations and varying environmental conditions.

Furthermore, exploring the role of females in these interactions would be an important avenue for future research. Investigations should examine female biases in both pre- and post-mating choice relative to potential male microbial cues or the enhanced quality of ejaculates produced by microbiome-rich males. A particularly crucial, yet unexplored, aspect is whether the female's own resident microbiome interacts with the incoming ejaculate or associated microbes, potentially influencing sperm storage patterns and biasing competitive outcomes independently of male traits.

Finally, the ecological context and broader applicability of our findings require exploration. While host-manipulation becomes increasingly challenging in systems with richer microbiotas, our full-factorial study established core principles of microbial interaction hierarchies that likely scale to complex communities. Research should examine how diet-microbiome interactions shape male reproductive strategies under varying resource conditions. Critically, assessing generality in species with diverse mating systems and modes of microbial transmission will reveal the evolutionary scope of these interactions. Although natural microbiomes generally exceed five species, functionally defined 'guilds' (e.g., acetate-producers) can be modelled combinatorially. Given deeply conserved microbiome functions across taxa, our framework provides a blueprint for targeted manipulation of keystone interactions in other tractable systems.

Conclusion

Our research demonstrates that the gut microbiome is an active participant in shaping male reproductive competitiveness. By influencing ejaculate quality and quantity, likely through modulation of host metabolism and resource allocation(7, 48, 49), these microbial symbionts redefine the landscape of condition dependence. Integrating the microbiome into sexual selection theory may fundamentally alter our understanding of how fitness variation is generated, maintained, and acted upon by selection in polyandrous systems and beyond. Our study emphasized the importance of considering the hologenome—the host plus its associated microbiota—as the relevant unit of selection when investigating the evolution of complex reproductive traits.

MATERIAL AND METHODS

Experimental populations

Throughout this study, we used *Drosophila melanogaster* of the LH$_m$ strain. These were either derived from an outbred wild-type population or, for standard competitors in the sperm competition assay, from a backcrossed population that had been genetically modified to ubiquitously express green fluorescent protein (GFP), permitting paternity assignment against the focal wild-type males (32, 33). To study biases in female sperm storage, we used focal males whose sperm heads expressed red fluorescent protein (RFP), and our standard competitors with ubiquitous GFP expression, permitting quantification of each competitor's sperm inside the female reproductive tract (32, 33, 43). We maintained all fly populations at moderate densities under constant conditions of 25°C, 60% humidity, and a 12h:12h light:dark cycle.

Generating axenic flies

To generate axenic flies, we performed all steps hereafter in a laminar flow chamber to avoid contamination. First, we collected eggs from Petri dishes containing grape-juice agar with a smear of yeast paste that had been placed in four large population cages after a six-hour egg-laying period. After washing these eggs off the plates into a strainer, we dechorionated them with 3.5% bleach for 2 minutes, followed by a rinse in 70% ethanol and three rinses with sterile water. We then transferred these eggs to autoclaved standard fly food supplemented with antibiotics (ampicillin, kanamycin, erythromycin, and tetracycline at final concentrations of 50, 50, 10, and 10 μg/ml, respectively) in autoclaved glass vials. These vials were then left undisturbed under standard conditions for flies to develop.

Upon fly eclosion, we collected both males and females as virgins, separated them by sex and confirmed their axenic state by plating a homogenized subsample of flies on LB and MRS agar followed by 16S PCR. Briefly, we extracted genomic DNA from 3−5 pooled axenic flies per sample using the ZymoBiomics^{TM} DNA Microprep kit following the instructions provided by the manufacturer. We also extracted DNA from flies with an intact microbiome to serve as positive controls. For DNA amplification, we used bacterial universal primers 27F and 1492R as forward and reverse primers, respectively. We prepared the PCR master mix using 9.5 μl of nuclease free water, 12.5 μl kappa HiFi master mix, 1 μl of forward and reverse primer and 1 μl of template DNA. The PCR cycle included an initial denaturation at 95°C for 90 s, followed by 35 cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s, an extension at 72°C for 90 s, and a final extension at 72°C for 5 mins. We ran the PCR products on a 1% agarose gel and visualized them under ultraviolet light after staining with SYBR Safe (Invitrogen) to confirm the absence of bacteria in axenic individuals and the presence of bacteria in controls.

We maintained all females and standard competitors as axenic flies throughout the

experiment and supplemented axenic focal males with different bacterial cultures to produce

gnotobiotic associations as described below. All adult food was free of antibiotics.

Bacterial cultures and gnotobiotic associations

Throughout this study, we used five bacterial species isolated from the gut of wild-type flies

(17): *Acetobacter orientalis*, *A. tropicalis*, *A. pasteurianus*, *Lactobacillus plantarum*, and *L.*

brevis. We associated our focal flies with all 32 possible combinations among these bacterial

strains following the general protocol of Gould et al. (17). To prepare bacterial cultures, we

grew them overnight: *Lactobacillus* spp. in MRS (De Man–Rogosa–Sharpe) broth without

shaking and *Acetobacter* spp. in MRS + mannitol medium at 30°C on a shaker. The next

morning, we checked these bacterial cultures for growth and performed their optical density

(OD) measurements at 600 nm. Then, we spun down these different bacterial cultures at 7000

rpm for 5 minutes and resuspended the resulting pellets to a concentration of 10^8 cells/mL in

sterile phosphate-buffered saline (PBS). Resuspension volume was determined following

established protocols (50, 51). Finally, we added a 50 µl (i.e., ca. 5×10^6 colony forming units,

CFUs) of bacterial culture to sterile (but antibiotic-free) fly food vials, adjusting the volume

per bacterial strain in multi-species associations for equal representation within a consistent

total density, which itself was comparable to prior reports for conventional flies (17, 51).

After adding the bacteria to food vials and air-drying these under in laminar flow hood,

we transferred 10 one-day-old axenic virgin males to each vial for inoculation with the added

bacterial strains. We moved these males to fresh food vials with same bacterial treatment every

alternate day for 5 days. Thereafter, we confirmed gnotobiotic associations by plating together

the homogenates of 3-4 adult flies from same treatments and checking their colony

morphologies, and by Sanger sequencing of 16S PCR homogenates. For multi-associations, we

plated the fly homogenates and picked the distinct types of colonies from a given plate for PCR and performed sequencing to confirm that morphologically similar species corresponds to same species in all the replicates, thus allowing us to distinguish between different bacterial species by looking at their colony appearance. Thereafter, we tested the presence of the expected bacterial species in a particular bacterial combination by plating them and looking at the colony morphologies.

Bacterial abundance in different combinations

To determine bacterial abundance in different gnotobiotic associations, we counted the CFUs per fly for each bacterial species across the different gnotobiotic associations. To this end, we washed males with 70% ethanol, individually homogenized them with a pestle in microcentrifuge tubes, and then diluted these homogenates to different dilutions with a final volume of 100 μl to find a dilution with comparable colony counts (approx. 100 CFUs) for visual quantification. In gnotobiotic cultures with both bacterial genera, we transferred an aliquot of well-mixed fly homogenate to two plates, supplementing the MRS media of one plate with 10 μg/ml ampicillin to selectively inhibit the growth of *Lactobacillus*, while incubating the other in a CO_2 atmosphere to prevent the growth of *Acetobacter*. We achieved within-genus differentiation of strains by colony morphologies: *A. orientalis* with flat colonies with ruffled borders, *A. tropicalis* with small tan coloured and highly reflective colonies, and *A. pasteurianus* with large tan colonies. Similarly, *L. brevis* produces brownish yellow colonies and *L. plantarum* milky white colonies (Supplementary Fig. S8). Using 16S PCR amplification followed by Sanger sequencing, we confirmed specificity of these colony properties, demonstrating the robustness of our method based on morphological identifications during experiments. After counting the number of colonies on each plate, we calculated CFU per fly

as CFU = $C \times D \times V / P$, where C = number of colonies counted, D = dilution factor, V =

volume of fly homogenate, and P = volume plated (50).

Across all 10 replicate plates combination examined, the expected species were present,

and none of the remaining species. Consistent with Gould et al. (17), total CFU counts

increased with species richness while the species-specific abundances declined

(Supplementary Fig. S9).

Sperm competitive ability

To investigate sperm competitiveness of focal males, we quantified their sperm offense ability

(P_2) by mating them as the second males with axenic females that had each mated with a an

axenic GFP male three days earlier. We monitored the occurrence of copulations for 7-8 hours

and transferred remated females to oviposition vials while giving those that did not remate

another two 8-hour remating opportunities on the following days. Flies that did not remate by

the last opportunity (within these 3 days) were discarded. We allowed all remated females to

lay eggs for three days. After all progeny eclosed, we scored them for paternity under a

fluorescence stereoscope, based on the presence/absence of the ubiquitous GFP marker in

standard competitors. The proportion of second-male progeny indicated the focal-male sperm

competitive ability (i.e., sperm offense). We replicated this experiment thrice with 30 replicates

per bacterial treatment each time (total N = 2,880 trials).

Sperm viability

We quantified sperm viability for 10 males per bacterial treatment (total N = 320 males). Once

males were six days old and inoculated, we anaesthetized them on ice and dissected them on a

microscopic slide to isolate the seminal vesicles from their reproductive tissue. We then placed

one of the seminal vesicles in 10 µl of Ringers solution and punctured it with an insect pin to

release the sperm mass for immediate staining with LIVE/DEAD fluorescent dye, following

the principles of prior studies on *Drosophila* (31, 52). This two-component dye consists of

SYBR-14 ('LIVE' component emitting green fluorescence), which is actively incorporated

into living cells, and propidium iodide (PI, with red fluorescence) that can only permeate

damaged membranes (typical of dead cells). We prepared an aqueous staining solution by

combining 2.4 µl of SYBR-14 stock solution (1 mM in DMSO) and 4.5 µl of PI stock solution

(2.4 mM in water) in 93.1 µl of Ringer's solution. Then, we added 5 µl of the staining solution

to the sperm sample before covering it with a coverslip and incubating it for 5 minutes in the

dark. Under an Olympus BX50 fluorescence microscope with a GFP/RFP dual filter, we

captured images of randomly chosen areas of each sample to later manually count at least 200

sperm and calculate the proportion of viable sperm in ImageJ v. 1.34(53).

Visualizing sperm competition within female reproductive tract

Since competitive fertilization events ultimately are the result of both male and female effects

(42, 54), we investigated how females bias their sperm storage based on interactions with males

supplemented with different microbiome. For this experiment, we again mated axenic virgin

females first with an axenic GFP male and, three days later, with a focal male as for sperm

competitive ability (above). To visualize sperm of both males, we used focal males expressing

RFP in their sperm heads. Immediately after remating, we transferred females to individual

wells of cell-culture plates, with a coverslip held in place with dots of rubber cement in two

corners. We checked these females every 10 min for up to 6 h for the ejection of the sperm

mass containing both displaced first-male and excess second-male sperm (32, 43). The timing

and extent of sperm ejection have previously been reported to affect relative numbers of first-

and second-male sperm stored by females and, consequently, competitive fertilization success

among males (33, 42).

Upon female sperm ejection, we recorded the interval from mating to ejection before

removing females from the well and freezing them individually in Eppendorf tubes. Thereafter,

we dissected these females and counted the green (standard competitors) and red sperm (focal

males) within the female reproductive tract (paired spermathecae, seminal receptacle and bursa

copulatrix) under a fluorescence microscope. From these counts, we determined S_2, the

proportion of all sperm stored by the female that came from the second (experimental) male.

We also counted the respective sperm numbers in the ejected sperm mass to estimate, combined

with above counts, the total number of sperm transferred by the second male and the number

of first-male sperm still residing in storage at the time of remating. This experiment was

performed in five blocks with 10 replicates per treatment for each block (total $N = 1,600$

remating trials).

Statistical analyses

All analyses were conducted in R v4.4.3 (R Core Team 2025), using bootstrapping procedures

with 1,000 iterations to assess the stability and robustness of model estimates. To evaluate the

effects of microbial diversity on reproductive outcomes, we resampled microbiome

compositions ($N = 32$ unique consortia) and fitted mixed-effects models across all bootstrap

datasets. These models included species richness (number of bacterial species) as a fixed effect

and both experimental block and microbial composition (i.e., consortium identity) as random

effects. Depending on the response variable, models were either linear mixed-effects models

(for traits such as the number of sperm transferred and time to sperm ejection) or generalized

linear mixed-effects models with a binomial error distribution and logit link function (for

proportions of viable sperm, second-male sperm in storage, and second-male progeny).

Binomial models included an observation-level random effect to account for overdispersion.

To examine the contribution of specific microbial interactions, we fitted similar mixed-

effects models using the presence/absence of each bacterial species and their two- to five-way

interactions as fixed effects, with block and, where appropriate, observation identity as random

effects. In this interaction-focused analysis, we bootstrapped at the level of individual

observations, stratified by microbial consortium, to retain the structure of specific

combinations. All subsequent analyses used the same set of bootstrapped datasets for

consistency and comparability.

To better understand the importance of species combinations in the microbiome and to

isolate emergent effects beyond individual species contributions, we used an epistasis-based

framework to elucidate how well the averaged phenotypic effects of single- or pairwise species

associations predict higher-order phenotypes (17). For single-species predictions, we compared

the measured trait values of males associated with two- to five-species microbiotas to values

averaged across mono-associated males that jointly represented the same bacterial

combinations. These differences (averaged prediction minus measured trait value) were then

plotted with bootstrapped 95% confidence intervals to visually determine if predictions

deviated from the measured estimates (i.e., confidence interval excluding zero difference). We

then repeated this procedure to predict phenotypes by averaging the estimates of di-associated

males (for calculations see (17)).

Following the same general principle, we further quantified how the presence or absence

of non-focal ("bystander") community members modulates both pairwise and three-way

microbial interactions by exhaustively enumerating all possible bystander contexts. We then

compared this "contextual" analysis against a "standard" test to reveal how higher-order

community background shapes emergent effects on host traits. Following Gould et al.'s (17)

nomenclature, the "standard" test calculated the interaction coefficient among all species

present in a specific community configuration, whereas the "contextual" tests assessed the

interaction between a focal species pair within the context of additional bystander species. We

computed interaction coefficients for each focal species pair or triplet across all possible

presence/absence combinations of bystander species and plotted them with bootstrapped

standard errors to highlight the varying directions of interactions depending on the bystander

combinations.

In both the single/double-species predictions and these standard/contextual tests, we

derived the coefficients to calculate the deviations from a mixed-effects models with an

appropriate link function for each microbial combination, controlling for block effects. We

repeated each model across 1,000 bootstrapped datasets to calculate the standard errors or 95%

confidence intervals from the distribution of the calculated coefficient or interaction

differentials.

Acknowledgements

We thank W. Ludington for providing the five bacterial strains used in our experiments.

Funding

This project was supported by the Swiss National Science Foundation (grant PP00P3_202662

to S.L.).

Author contributions

K.M.: conceptualization, methodology, investigation, data collection and curation, data

analysis, writing original draft, writing—review and editing, project administration; A.M.:

investigation, data collection, writing—review and editing; A.N.D.N: investigation, data

collection, writing—review and editing; S.H.S: investigation, writing—review and editing;

J.R.: investigation; S.L.: conceptualization, methodology, funding acquisition, data analysis,

project administration, supervision, writing original draft, writing manuscript—review and editing.

All authors gave final approval for publication and have agreed to be held accountable for the work performed therein.

Competing interests

All authors declare that they have no conflict of interest.

REFERENCES

1. L. W. Simmons, J. L. Fitzpatrick, Sperm wars and the evolution of male fertility. *Reproduction* **144**, 519–534 (2012).

2. J. L. Fitzpatrick, S. Lüpold, Sexual selection and the evolution of sperm quality. *Mol. Hum. Reprod.* **20**, 1180–1189 (2014).

3. D. A. Dewsbury, Ejaculate cost and male choice. *Am. Nat.* **119**, 601–610 (1982).

4. S. Lüpold, *et al.*, How sexual selection can drive the evolution of costly sperm ornamentation. *Nature* **533**, 535–538 (2016).

5. M. Olsson, T. Madsen, R. Shine, Is sperm really so cheap? Costs of reproduction in male adders, *Vipera berus*. *Proc. R. Soc. B* **264**, 455–459 (1997).

6. J. C. Perry, C. T. Tse, Extreme costs of mating for male two-spot ladybird beetles. *PLoS ONE* **8**, e81934 (2013).

7. E. L. Macartney, A. J. Crean, S. Nakagawa, R. Bonduriansky, Effects of nutrient limitation on sperm and seminal fluid: a systematic review and meta-analysis. *Biol. Rev.* **94**, 1722–1739 (2019).

8. Y. Iwasa, A. Pomiankowski, S. Nee, The evolution of costly mate preferences. II. The

handicap principle. *Evolution* **45**, 1431–1442 (1991).

9. A. Grafen, Sexual selection unhandicapped by the Fisher process. *J. Theor. Biol.* **144**,

517–546 (1990).

10. L. Rowe, D. Houle, The lek paradox and the capture of genetic variance by condition

dependent traits. *Proc. R. Soc. B* **263**, 1415–1421 (1996).

11. G. E. Hill, Condition-dependent traits as signals of the functionality of vital cellular

processes. *Ecol. Lett.* **14**, 625–634 (2011).

12. E. C. Lindsay, N. B. Metcalfe, M. S. Llewellyn, The potential role of the gut microbiota

in shaping host energetics and metabolic rate. *J. Anim. Ecol.* **89**, 2415–2426 (2020).

13. T. O. Cox, P. Lundgren, K. Nath, C. A. Thaiss, Metabolic control by the microbiome.

Genome Med. **14**, 80 (2022).

14. A. E. Douglas, The microbial dimension in insect nutritional ecology. *Funct. Ecol.* **23**,

38–47 (2009).

15. J. Alcock, C. C. Maley, C. A. Aktipis, Is eating behavior manipulated by the

gastrointestinal microbiota? Evolutionary pressures and potential mechanisms.

BioEssays **36**, 940–949 (2014).

16. S. O. Fetissov, Role of the gut microbiota in host appetite control: bacterial growth to

animal feeding behaviour. *Nat. Rev. Endocrinol.* **13**, 11–25 (2017).

17. A. L. Gould, *et al.*, Microbiome interactions shape host fitness. *Proc. Natl. Acad. Sci. U.*

S. A. **115**, E11951–E11960 (2018).

18. A. W. Walters, *et al.*, The microbiota influences the *Drosophila melanogaster* life history

strategy. *Mol. Ecol.* **29**, 639–653 (2020).

19. V. Emelianoff, *et al.*, A survival-reproduction trade-off in entomopathogenic nematodes

mediated by their bacterial symbionts. *Evolution* **62**, 932–942 (2008).

20. X. Qi, C. Yun, Y. Pang, J. Qiao, The impact of the gut microbiota on the reproductive

and metabolic endocrine system. *Gut Microbes* **13**, 1894070 (2021).

21. P. Comizzoli, M. L. Power, S. L. Bornbusch, C. R. Muletz-Wolz, Interactions between

reproductive biology and microbiomes in wild animal species. *Anim. Microbiome* **3**, 87

(2021).

22. S. B. Chadchan, V. Singh, R. Kommagani, Female reproductive dysfunctions and the gut

microbiota. *J. Mol. Endocrinol.* **69**, R81–R94 (2022).

23. M. K. Matthews, J. Malcolm, J. M. Chaston, Microbiota influences fitness and timing of

reproduction in the fruit fly *Drosophila melanogaster*. *Microbiol. Spectr.* **9**, e0003421

(2021).

24. M. Al-Asmakh, *et al.*, The gut microbiota and developmental programming of the testis

in mice. *PLoS ONE* **9**, e103809 (2014).

25. N. Ding, *et al.*, Impairment of spermatogenesis and sperm motility by the high-fat diet-

induced dysbiosis of gut microbes. *Gut* **69**, 1608–1619 (2020).

26. M. Alfano, *et al.*, Testicular microbiome in azoospermic men—first evidence of the

impact of an altered microenvironment. *Hum. Reprod.* **33**, 1212–1217 (2018).

27. S.-L. Weng, *et al.*, Bacterial communities in semen from men of infertile couples:

metagenomic sequencing reveals relationships of seminal microbiota to semen quality.

PloS One **9**, e110152 (2014).

28. E. V. Ridley, A. C. N. Wong, A. E. Douglas, Microbe-dependent and nonspecific effects

of procedures to eliminate the resident microbiota from *Drosophila melanogaster*. *Appl.*

Environ. Microbiol. **79**, 3209–3214 (2013).

29. A. C.-N. Wong, J. M. Chaston, A. E. Douglas, The inconstant gut microbiota of

Drosophila species revealed by 16S rRNA gene analysis. *ISME J.* **7**, 1922–1932 (2013).

30. J. G. McMullen, G. Peters-Schulze, J. Cai, A. D. Patterson, A. E. Douglas, How gut

microbiome interactions affect nutritional traits of *Drosophila melanogaster*. *J. Exp.*

Biol. **223** (2020).

31. M. Tourmente, C. R. Archer, D. J. Hosken, Complex interactions between sperm viability

and female fertility. *Sci. Rep.* **9**, 15366 (2019).

32. S. Lüpold, *et al.*, How multivariate ejaculate traits determine competitive fertilization

success in *Drosophila melanogaster*. *Curr. Biol.* **22**, 1667–1672 (2012).

33. S. Lüpold, *et al.*, Female mediation of competitive fertilization success in *Drosophila*

melanogaster. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 10693–10698 (2013).

34. A. C.-N. Wong, A. J. Dobson, A. E. Douglas, Gut microbiota dictates the metabolic

response of *Drosophila* to diet. *J. Exp. Biol.* **217**, 1894–1901 (2014).

35. C. C. Maresch, *et al.*, Diabetes-induced hyperglycemia impairs male reproductive

function: a systematic review. *Hum. Reprod. Update* **24**, 86–105 (2018).

36. S. Subhadarshini, V. Mallesh, G. K. Bharath, Semen quality alterations in men with

hyperlipidemia and diabetes mellitus: a cross-sectional study. *Int. J. Acad. Med. Pharm.*

6, 639–644 (2024).

37. S. C. Shin, *et al.*, *Drosophila* microbiome modulates host developmental and metabolic

homeostasis via insulin signaling. *Science* **334**, 670–674 (2011).

38. I. Iatsenko, J.-P. Boquete, B. Lemaitre, Microbiota-derived lactate activates production of

reactive oxygen species by the intestinal NADPH oxidase nox and shortens *Drosophila*

lifespan. *Immunity* **49**, 929-942.e5 (2018).

39. R. J. Aitken, Z. Gibb, M. A. Baker, J. Drevet, P. Gharagozloo, Causes and consequences

of oxidative stress in spermatozoa. *Reprod. Fertil. Dev.* **28**, 1–10 (2016).

40. A. Zelezniak, *et al.*, Metabolic dependencies drive species co-occurrence in diverse

microbial communities. *Proc. Natl. Acad. Sci.* **112**, 6449–6454 (2015).

41. L. Rowe, D. Houle, The lek paradox and the capture of genetic variance by condition

dependent traits. *Proc. R. Soc. Lond. B Biol. Sci.* **263**, 1415–1421 (1996).

42. S. Lüpold, *et al.*, How female × male and male × male interactions influence competitive

fertilization in *Drosophila melanogaster. Evol. Lett.* **4**, 416–429 (2020).

43. M. K. Manier, *et al.*, Resolving mechanisms of competitive fertilization success in

Drosophila melanogaster. Science **328**, 354–357 (2010).

44. L. P. Henry, M. Bruijning, S. K. Forsberg, J. F. Ayroles, The microbiome extends host

evolutionary potential. *Nat. Commun.* **12**, 5141 (2021).

45. N. A. Moran, H. Ochman, T. J. Hammer, Evolutionary and ecological consequences of

gut microbial communities. *Annu. Rev. Ecol. Evol. Syst.* **50**, 451–475 (2019).

46. R. C. Firman, C. Gasparini, M. K. Manier, T. Pizzari, Postmating female control: 20

years of cryptic female choice. *Trends Ecol. Evol.* **32**, 368–382 (2017).

47. L. Daupagne, *et al.*, One size does not fit all: female–male interactions on the path to

fertilization. (2025). https://doi.org/10.1530/REP-24-0238.

48. E. L. Macartney, *et al.*, Sperm depletion in relation to developmental nutrition and

genotype in *Drosophila melanogaster. Evolution* **75**, 2830–2841 (2021).

49. V. Zeender, S. Pfammatter, B. Roschitzki, S. Dorus, S. Lüpold, Genotype-by-

environment interactions influence the composition of the *Drosophila* seminal

proteome. *Proc. R. Soc. B* **290**, 20231313 (2023).

50. M. L. Koyle, *et al.*, Rearing the fruit fly *Drosophila melanogaster* under axenic and

gnotobiotic conditions. *J. Vis. Exp.* **113**, e54219 (2016).

51. P. D. Newell, A. E. Douglas, Interspecies interactions determine the impact of the gut

microbiota on nutrient allocation in *Drosophila melanogaster. Appl. Environ.*

Microbiol. **80**, 788–796 (2014).

52. A. Meena, *et al.*, Fertility loss and recovery dynamics after repeated heat stress across life

stages in male *Drosophila melanogaster*: Patterns and processes. *R. Soc. Open Sci.* **11**,

241082 (2024).

53. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image

analysis. *Nat. Methods* **9**, 671–675 (2012).

54. W. G. Eberhard, *Female Control: Sexual Selection by Cryptic Female Choice* (Princeton

University Press, 1996).

Figures

A   Number of species   B   Number of species   C   Number of species   D   Number of species   E   Number of species

Fig. 1. Associations between the number of bacterial species in the microbiota and the different

fitness traits. Each point indicates the mean of all replicate samples for a given microbiome

consortium (N = 32). The lines and 95% confidence bands are derived from bootstrapped

(generalized) linear mixed-effects models with microbiome composition and experimental

block as random effects (for binomial models additionally also an observation-level random

effect).

Fig. 2. Variation in male fitness traits relative to their microbial community (left) and error in predicting these values for males associated with multiple species based on single- or pairwise species averages (right). In the left panels, colours indicate the different combinations of

Acetobacter and *Lactobacillus* species associated with experimental males. The right panels

depict the deviations in trait predictions from measured traits by averaging the corresponding

coefficients of (generalized) linear mixed-effects models for individual species (orange points)

or species pairs (green diamonds) that jointly represented the same species communities. The

relevant microbial combinations are plotted in the same order across all panels. Error bars

reflect 95% confidence intervals (from raw distributions in left panels, from bootstrapped

prediction differentials in right panels). Sperm transfer in thousands, time to sperm ejection in

hours, other variables in proportions.

Fig. 3. Correlations between the different phenotypic traits and the relative abundance (based on CFU counts) of each of the five bacterial species (left) and among the traits themselves (right). Values reflect the averaged correlation coefficients with 95% confidence limits of pairwise Pearson correlations across bootstrapped group means ($N = 1,000$ samples) among males with a given microbiome consortium. Relative abundances and proportional traits (sperm viability, S_2, P_2) were logit-transformed before bootstrapping. Colours indicate the strength and direction of correlations as reflected by the colour-gradient bar.

Fig. 4. Interactive effects between bacterial species contingent on bystander species, with two-way interactions between pairs of focal of species in the left panels, and three-way interactions between trios of focal species in the right panels. For each code along the x-axes, 1's indicate the species configuration of the interaction and asterisks denote placeholders for bystander

species (order within codes from left to right: *A. orientalis*, *A. tropicalis*, *A. pasteurianus*, *L.*

brevis, *L. plantarum*). The binary codes in the legends indicate the configuration of bystanders

in the order, in which they would fill the placeholders (*) in each focal species configuration

(1 = presence, 0 = absence). The right panels distinguish between the "standard" tests (grey

triangles; i.e., global interaction across all community members present) and "contextual" tests

(coloured points; i.e., separate interactions between a focal species pair or triplet for each

bystander combination) for each focal triplet. All coefficients and propagated standard errors

were derived from (generalized) linear mixed-effects models across 1,000 bootstrapped

datasets.

Supplementary figures
671

672
673
Supplementary Figure S1: Competitive fertilization success, measured as the proportion of second-
male (i.e., focal-male) paternity, against either axenic or conventional competitors (first males to mate).
In either context, males that were associated with the five-species microbial consortium (*Acetobacter*
orientalis, *A. tropicalis*, *A. pasteurianus*, *Lactobacillus brevis* and *L. plantarum*) after axenic
development outperformed consistently germ-free (axenic) males and even matched males with a
conventional (entirely unmanipulated) microbiome in their sperm competitiveness, at least against
germ-free competitors. Error bars represent 95% confidence intervals.
681

682
683
Supplementary Figure S2: Ridge plot showing the distribution of fixed-effect odds ratios across 1,000
bootstrap iterations of a generalized linear mixed-effects model on the proportion of focal-male progeny
in sperm competition against axenic males. The model includes all main effects and two- to four-way
interactions among five bacterial species. Each ridge represents the distribution of bootstrapped
estimates for a given term, and its stability. Ridge colours indicate the direction of the effect (green:
positive; purple: negative), with darker shades denoting effects whose 95% confidence intervals exclude
zero, and lighter shades for those where only the 90% confidence interval excludes zero. The vertical
dashed line indicates no effect.

Supplementary Figure S3: Ridge plot showing the distribution of fixed-effect odds ratios across 1,000
bootstrap iterations of a generalized linear mixed-effects model on the proportion of focal-male progeny
in sperm competition against males with an intact microbiome. The model includes all main effects and
two- to four-way interactions among five bacterial species. Each ridge represents the distribution of
bootstrapped estimates for a given term, and its stability. Ridge colours indicate the direction of the
effect (green: positive; purple: negative), with darker shades denoting effects whose 95% confidence
intervals exclude zero, and lighter shades for those where only the 90% confidence interval excludes
zero. The vertical dashed line indicates no effect.

702
703
Supplementary Figure S4: Ridge plot showing the distribution of fixed-effect odds ratios across 1,000
bootstrap iterations of a generalized linear mixed-effects model on the proportion of viable sperm. The
model includes all main effects and two- to four-way interactions among five bacterial species. Each
ridge represents the distribution of bootstrapped estimates for a given term, and its stability. Ridge
colours indicate the direction of the effect (green: positive; purple: negative) where 95% confidence
intervals exclude zero. The vertical dashed line indicates no effect.
710

Supplementary Figure S5: Ridge plot showing the distribution of fixed-effect odds ratios across 1,000
bootstrap iterations of a generalized linear mixed-effects model on the proportion of focal-male sperm
remaining in female storage after sperm ejection. The model includes all main effects and two- to four-
way interactions among five bacterial species. Each ridge represents the distribution of bootstrapped
estimates for a given term, and its stability. Ridge colours indicate the direction of the effect (green:
positive; purple: negative) where 95% confidence intervals exclude zero. The vertical dashed line
indicates no effect.

720
721
Supplementary Figure S6: Ridge plot showing the distribution of fixed-effect estimates across 1,000
bootstrap iterations of a linear mixed-effects model on the time to female sperm ejection. The model
includes all main effects and two- to four-way interactions among five bacterial species. Each ridge
represents the distribution of bootstrapped estimates for a given term, and its stability. Ridge colours
indicate the direction of the effect (green: positive; purple: negative), with darker shades denoting
effects whose 95% confidence intervals exclude zero, and lighter shades for those where only the 90%
confidence interval excludes zero. The vertical dashed line indicates no effect. Estimates in minutes.

Supplementary Figure S7: Ridge plot showing the distribution of fixed-effect estimates across 1,000

bootstrap iterations of a linear mixed-effects model on the number of sperm transferred by focal males.

The model includes all main effects and two- to four-way interactions among five bacterial species.

Each ridge represents the distribution of bootstrapped estimates for a given term, and its stability. Ridge

colours indicate the direction of the effect (green: positive; purple: negative), with darker shades

denoting effects whose 95% confidence intervals exclude zero, and lighter shades for those where only

the 90% confidence interval excludes zero. The vertical dashed line indicates no effect. Estimates in

raw sperm counts.

739
740
Supplementary Figure S8: Image showing morphological appearances of colonies formed by different
bacterial species. On right side, acetobacter strains are present with *Acetobacter orientalis* represented
by flat colonies with ruffled borders, *Acetobacter tropicalis* with small tan colored and highly reflective
colonies, and *Acetobacter pasteurianus* with large tan colonies. In contrast on left side, *Lactobacillus*
brevis shows brownish yellow colonies and *Lactobacillus plantarum* with milky white colonies.

Supplementary Figure S9: Abundances of the different species (linear scale) for all 32 possible species
combinations ($N = 10$ replicate flies per combination) with increasing species richness. The height of
the bars in the left panel reflect the total microbial load ($\log_{10}$ scale), with the relative abundances of the
constituent species shown by coloration (linear scale). Total increase in total bacterial load with
increasing species richness is shown by increasing bar height from left to right. The right panel depicts
the decline in species-specific colony-forming units (CFUs) counts in response to increasing species
richness. All error bars show 95% confidence intervals.
755