An innovative in vitro model for studying the biology of cardiac fibroblasts originating from the epicardium

Claudia Müller-Sánchez*[1], María Gertrudis Muñiz-Banciella[1,2], Manuel Reina[1], Francesc X. Soriano[1,3] and Ofelia M. Martínez-Estrada*,[1,2]

1. Celltec-UB, Department of Cell Biology, Physiology, and Immunology, Faculty of Biology, University of Barcelona, Barcelona, Spain
2. Institute of Biomedicine (IBUB), University of Barcelona, Barcelona, Spain
3. Institut de Neurociències, Universitat de Barcelona, Barcelona, Spain

Running title: Innovative cardiac fibroblast model

* Correspondence:

Claudia Müller-Sánchez

Department of Cell Biology, Physiology, and Immunology

Faculty of Biology, University of Barcelona

Ave Diagonal 643

E-mail: cmuller@ub.edu

Ofelia M. Martínez-Estrada

Department of Cell Biology, Physiology, and Immunology

Faculty of Biology, University of Barcelona

Ave Diagonal 643

08028 Barcelona, Spain

E-mail: ofeliamartinez@ub.edu

Abstract

Embryonic epicardium is a major source of cardiac fibroblasts (CFs), which play essential roles in heart development and response to heart injury. In this study, we developed a novel mouse model to identify distinct populations of epicardium-derived CFs based on WT1 expression. Our Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP model enables lineage tracing of WT1Cre-labeled (RFP$^+$) fibroblasts and the identification of cells actively expressing WT1 (GFP$^+$). Flow cytometry at early postnatal stages showed that RFP$^+$ cells form a heterogeneous stromal population, with 20.13% co-expressing GFP, indicating persistent WT1 expression in a subset. We successfully immortalized RFP$^+$ cardiac stromal cells, highly enriched in fibroblasts, by excluding other Wt1Cre-active cell types. Through culture condition optimization, we could selectively expand or differentiate specific fibroblast subpopulations, increasing the model's utility. These immortalized cells, carrying an integrated WT1 reporter system, provide a robust in vitro platform to study fibroblast activation, differentiation, and plasticity under defined conditions.

Summary blurb

This study presents a novel in vitro mouse model for investigating the activation, identity, and functional properties of epicardium-derived cardiac fibroblasts.

INTRODUCTION

Cardiac fibroblasts (CFs) are essential non-myocyte cells that play a crucial role in heart development and homeostasis (Tallquist, 2018). They are among the most abundant non-cardiomyocyte cell types in the postnatal heart and are responsible for extensive extracellular matrix (ECM) deposition. Beyond their structural role, CFs produce paracrine factors critical to heart development, myocardial growth, functional maturation, and blood vessel formation (Acharya et al, 2012; Hortells et al, 2020; Ieda et al, 2009). They are also central to the formation of pathological fibrosis, a hallmark of various cardiovascular diseases, by promoting excessive ECM deposition and tissue remodeling (Humeres & Frangogiannis, 2019).

The understanding of CF biology has advanced significantly over the past decade (Tallquist & Molkentin, 2017). Although several origins have been proposed, lineage tracing studies in mice have unequivocally demonstrated that adult resident CFs primarily arise from the embryonic epicardium and endocardium, with the epicardium being the predominant source (Moore-Morris et al, 2014; Tallquist, 2018). Notably, the population of resident CFs of epicardial origin expands in response to adult cardiac fibrosis, such as that induced by pressure overload, fibroelastosis, or myocardial infarction (Forte et al, 2020; Moore-Morris et al, 2014; Ruiz-Villalba et al, 2015; Zhang et al, 2017).

Compared with other cell types, the absence of unique fibroblast markers, coupled with their inherent heterogeneity, has made the isolation of fibroblasts particularly challenging when using conventional techniques such as fluorescence-activated cell sorting (FACS) (Tallquist & Molkentin, 2017). Various genetic mouse models have, therefore, been developed to facilitate fibroblast identification. Some models employ direct fibroblast reporter systems, such as the Pdgfra-GFP and Collagen1a1-GFP reporter lines (Hamilton et al, 2003; Yata et al, 2003). Additionally, Cre transgenic mouse models have significantly improved the identification and characterization of CFs (Acharya et al, 2011; Tallquist & Molkentin, 2017; Wessels et al, 2012).

The WT1 gene encodes a zinc finger protein essential for heart development (Hastie, 2017; Moore et al, 1999). WT1 is highly expressed in the epicardium and epicardial derived cells (EPDCs) during heart development, where it plays a critical role in their formation and function (Martinez-Estrada et al, 2010; Moore et al, 1999). Epicardial derivatives, such as CFs, also express WT1 both during heart development and following myocardial infarction (Braitsch et al, 2013; Perez-Pomares et al, 2002; Zhou et al, 2011). These data, originally

identified by immunostaining, have been validated by several transcriptomic studies (Deng et al, 2025; Feng et al, 2022). Both constitutive and inducible Wt1Cre mouse models have been generated and used to study fibroblast formation and expansion during development (Wessels et al, 2012; Zhou et al, 2008), and these models have become essential tools for investigating resident fibroblasts of epicardial origin in heart diseases (Forte et al, 2020; Moore-Morris et al, 2018; Ruiz-Villalba et al, 2015).

In this study, we used a constitutive BAC Wt1Cre transgenic mouse model, which efficiently labels resident CFs, and generated a new Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mouse model. This includes WT1 lineage cells that are RFP-positive (RFP$^+$), while cells actively expressing WT1 are GFP-positive (GFP$^+$). We subsequently established immortalized CFs from this model, which represent a valuable and versatile in vitro tool for dissecting the molecular mechanisms underlying fibroblast activation, differentiation, and plasticity.

RESULTS

Characterizing cardiac cells from the Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mouse model

Although several studies have used Wt1Cre mouse models to study fibroblast formation during cardiac development and disease states, none have combined lineage tracing with a Wt1 reporter model to identify both Wt1Cre progeny and Wt1-expressing cells simultaneously, which can be used to FACS-isolate different populations of Wt1Cre positive cells (Forte et al, 2020; Moore-Morris et al, 2014; Wessels et al, 2012). We aimed to generate such a model by crossbreeding Wt1Cre;ROSA26-tdRFP mice with Wt1$^{GFP/+}$ mice (Fig. 1A).

In the Wt1$^{GFP/+}$ model, GFP is knocked into exon 1 and driven by the endogenous Wt1 regulatory sequences, providing a GFP reporter that faithfully reflects Wt1 expression (Hosen et al, 2007; Velecela et al, 2019). In the new model, Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mice contain RFP-positive (RFP$^+$) cells that exhibiting Cre recombinase activity, along with their progeny, while GFP-positive (GFP$^+$) cells express abundant WT1 levels. Flow cytometry analysis of ventricular cardiac cells at early postnatal stages showed that RFP$^+$ cells represented 13.75% of the total cardiac cell population. Of these, 20.13% were GFP$^+$, indicating sustained WT1 expression in a cell subset (Fig. 1A, B, Table 1).

To further characterize the RFP$^+$ population, we employed a panel of several antibodies. This showed that 25.83% of RFP$^+$ cells expressed CD31 and 27.25% expressed CD45,

confirming the presence of endothelial and hematopoietic cells within the RFP$^+$ population, consistent with previous reports (Fig. 1C, D, Table 2) (Cano et al, 2016; Pogontke et al, 2022). Notably, 69.61% of the RFP$^+$ CD45$^+$ cells and 76.34% of the RFP$^+$ CD31$^+$ cells co-expressed GFP (Table 3).We also included markers commonly used to identify the stromal cell populations GP38, CD90, and CD39. Among the RFP$^+$ cells, 73.25% were GP38$^+$, 18.03% were CD90$^+$, and 29.03% were CD39$^+$ (Fig. 1C, Table 2). Of note, the percentages of GFP$^+$ cells varied among these populations, accounting for 28.91% of GP38$^+$, 36.46% of CD90$^+$, and 71.04% of CD39$^+$ cells (Table 4).

Taken together, these results demonstrate that the Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mouse model allows the identification of different populations of Wt1Cre-positive stromal cells based on GFP expression.

Immortalization of cardiac cells in the Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mouse model

Most studies using CF cultures rely on selective attachment to isolate the cell population, often without additional enrichment strategies (Garvin & Katwa, 2023). In this study, we FACS-isolated RFP$^+$ ventricular cardiac stromal cells from the Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mouse model, excluding endothelial (RFP$^+$, CD31$^+$, CD45$^-$), hematopoietic (RFP$^+$, CD31$^-$, CD45$^+$), and high GFP-expressing cells, thereby including epicardial cells, among others (Fig. 2A). The sorted RFP$^+$ cells were cultured on poly-L-lysine-coated plates, where they exhibited a flattened, irregular morphology with abundant cytoplasmic protrusions (Fig. 2B). Next, the RFP$^+$ cells were immortalized using a viral vector containing the SV40 large T antigen (SV40 TAg), which is well-recognized for its robust cell immortalization capability. Following puromycin selection, the immortalized cells exhibited a more uniform morphology with abundant cytoplasmic protrusions (Fig. 2C). Immunofluorescence analysis at low passage confirmed uniform expression of vimentin, a key fibroblast-specific marker, in all cells (Fig. 2D).

These results validate the successful immortalization of FACS-sorted RFP$^+$ cardiac stromal cells.

Characterization of cardiac RFP$^+$ immortalized cells derived from the Wt1$^{GFP/+}$;Wt1Cre;R26-tdRFP mouse model

Next, we aimed to characterize the RFP-immortalized cells in detail. Flow cytometry at lower passage confirmed uniform RFP expression across all cells, with 15.49% expressing GFP (Fig. 3A,Table 5). When we included the antibodies used for in vivo analysis, the flow

cytometry revealed that 97.40% of the RFP$^+$ immortalized cells expressed GP38, that 63.12% expressed CD90, and that 5.87% expressed CD39 (Fig. 3A, Table 5). Consistent with in vivo observations, the proportion of GFP$^+$ cells varied among subpopulations: 10.14%, 11.61%, and 29.98% among the RFP$^+$GP38$^+$, RFP$^+$CD90$^+$, and RFP$^+$CD39$^+$ subpopulations, respectively (Fig. 3B, Table 6). Given that none of these markers are specific to fibroblasts, and that previous studies have shown the Wt1Cre line can recombine in other cell types, we assessed the expression of genes enriched in other cardiac cell populations, including *Des* (smooth muscle cells), *Rgs5* (pericytes), and *Myh6* (cardiomyocytes) (Fig. 3C). We also examined the expression of genes highly expressed in the epicardium, such as *Upk3b* and *Msln* (Fig. 3D). The qRT-PCR analysis revealed that immortalized and sorted RFP$^+$ cells exhibited undetectable levels of these non-fibroblast cell markers and abundant vimentin expression (Fig. 3C, D).

Collectively, these findings demonstrate that the RFP$^+$ cells represent a heterogeneous population of CFs.

Immortalized CFs undergo population shifts over time

Generating immortalized cells allows the study of CF signaling, proliferation, and activation with less need to use animals. This can be achieved because the immortalized cells remain proliferative across multiple passages without showing signs of senescence. An important consideration, however, is that these cells comprise various subpopulations, which we intentionally maintained to reflect the in vivo heterogeneity of CFs more accurately.

To evaluate whether the immortalized cells retained their original characteristics over multiple passages, without isolating distinct subsets, we also analyzed higher passage cells using flow cytometry. Antibody analysis then showed that GP38$^+$ and CD39$^+$ cell expression remained largely unchanged, with most cells being GP38$^+$ and a minority being CD39$^+$ (Fig. 4A, Table 7). At lower passages, CD90$^+$ cells accounted for 63.12%, but this proportion decreased to 11.63% at higher passages (Fig. 3A, 4A, Table 7). Furthermore, as the number of CD90$^+$ cells declined, the proportion of CD90$^-$GFP$^-$ cells increased from 34% to 75.5% (Fig. 4B, C).

Collectively, these findings suggest that immortalized CFs undergo population shifts over time, favoring subpopulations with lower CD90 expression.

DISCUSSION

We developed a novel mouse model that enables distinct cardiac stromal cell populations, including resident fibroblasts, to be isolated based on WT1 expression. The Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP model facilitates both the lineage tracing of cells labeled with Wt1Cre (RFP$^+$) and the identification of cells actively expressing WT1 (GFP$^+$). We generated an immortalized population highly enriched in different CF subtypes that retained RFP expression and expressed key fibroblast-specific markers, such as vimentin. The integrated GFP reporter enabled the isolation and functional analysis of distinct subpopulations within the Wt1Cre lineage.

During cardiac development, a subset of epicardial cells undergoes epithelial-to-mesenchymal transition, giving rise to EPDCs. These cells invade the subepicardial matrix before migrating into the myocardium, where they differentiate into various cardiac lineages, including most CFs (von Gise & Pu, 2012). During early postnatal (P) development, CFs undergo significant expansion via cell proliferation, which dramatically decreases after P7 (Ieda et al, 2009). While several mechanisms that regulate the initial formation and differentiation of EPDCs have been elucidated, the molecular pathways controlling CF proliferation and terminal differentiation remain largely unknown (von Gise & Pu, 2012). Given these limitations, the immortalized CFs generated in this study offer a valuable platform for screening candidate pathways involved in fibroblast function. Indeed, such analyses are typically difficult to perform using primary cultures because they contain limited numbers of cells in each subpopulation, hindering in-depth cellular and molecular studies.

The utility of this model also extends beyond developmental studies, because the adult mammalian heart has a limited capacity for regeneration. Healing typically occurs through scar formation, in which resident fibroblasts of epicardial origin play a central role. Although this response is essential for maintaining structural integrity and preventing ventricular rupture following injury, excessive or dysregulated activation can result in adverse cardiac remodeling, reduced contractile function, and ultimately, heart failure (Moore-Morris & Evans, 2023; Quijada et al, 2019). Understanding the regulatory mechanisms that govern the expansion and differentiation of CFs into specific subpopulations could help develop therapeutic strategies that mitigate the effects of pathological fibrosis while preserving essential wound healing processes.

WT1 is a transcription factor essential for development and tissue homeostasis (Hastie, 2017). Recent studies have highlighted its profibrotic role in lung fibrosis; however, WT1

deletion in the liver has also been shown to exacerbate fibrogenesis following injury (Kendall et al, 2019; Sontake et al, 2018). In the heart, studies using a zebrafish model of myocardial injury have shown that *wt1a:GFP⁺* cells upregulate a fibrotic gene program in response to damage. After analyzing different scRNA-seq datasets in mammals, WT1 has been classified as an early transcriptional regulator following myocardial infarction (Patrick et al, 2024; Sanchez-Iranzo et al, 2018). Despite these findings, the contribution of Wt1-expressing fibroblasts to the early stages of fibroblast activation and scar tissue formation after myocardial infarction remains unclear. The in vivo and cellular models generated in this study will allow us to compare the properties of different populations of cardiac fibroblasts and their responses to hypoxia. We hypothesize that this response may differ between WT1-positive and WT1-negative cell populations.

In summary, we have developed a versatile model for studying CFs of epicardial origin. By adjusting the culture conditions, specific fibroblast subpopulations can be selectively promoted to enhance the model's flexibility. The choice of conditions should be guided by the physiologically relevant context and desired subpopulation, which will allow for not only screening GFP expression but also for identifying treatment combinations that can modulate the proportions of different cell subpopulations. In turn, this can provide deeper insights into the functional roles of these cells.

MATERIAL AND METHODS

Animal models

The Wt1$^{GFP/+}$, Wt1Cre (Tg(Wt1-cre)#Jbeb), and ROSA26-tdRFP mice have been described previously (Hosen et al, 2007; Luche et al, 2007; Wessels et al, 2012). To generate Wt1Cre;ROSA26-tdRFP mice, male Wt1Cre mice were mated with ROSA26dtRFP females. To generate Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mice, Wt1Cre;ROSA26-tdRFP males were mated with Wt1$^{GFP/+}$ females. The primers used for genotyping are listed in Table 9.

All animal experiments were carried out in accordance with the regulations of the Animal Experimentation Ethics Committee (CEEA) of the University of Barcelona (ID C121CG2V6), thereby complying with current Spanish and European legislation.

Flow cytometry analysis

Flow cytometry analyses were performed on freshly isolated cells from the cardiac ventricles of mice or on immortalized cells.

For the in vivo analyses, early postnatal heart ventricles from Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mice and littermate negative controls were digested in 1 mg/mL collagenase type I solution (Worthington, 9001-12-1) for 30 minutes at 37°C, using a shaking block set to 1000 rpm (Eppendorf Thermomixer Compact). Collagenase activity was stopped by washing the cells with DMEM (Gibco,11960044) containing 10% fetal bovine serum (FBS; Capricorn scientific, FBS-12B). Subsequently, the cells were then pelleted by centrifugation at 300 xg for 5 minutes and filtered through a 40 µm cell strainer (Corning, 352340) before being resuspended in Hanks' Balanced Salt Solution (HBSS Ca^{2+}, Mg^{2+}; Gibco, 14025092) containing 2% FBS (FACs buffer) (Torres-Cano et al, 2022).

For antibody staining, freshly isolated or immortalized cells were incubated with primary monoclonal or isotype control antibodies at 4°C for 30 minutes (Table 10) (Ramiro-Pareta et al, 2023). Samples were washed twice and resuspended in FACs buffer. Flow cytometry analysis was carried out using a Cytek® Aurora 4L (Cytek) and data were analyzed using FlowJo™ software (v10.10). Cell gating of the corresponding fluorescent protein was carried out using samples negative for the fluorescent transgenic protein and isotype control antibodies.

Immortalization of RFP+ cardiac cells

RFP$^+$ cells were isolated by FACS (FACSAria™ Fusion I, BD) from the enzymatically digested hearts of Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mice, excluding endothelial (RFP$^+$, CD31$^+$, CD45$^-$), hematopoietic (RFP$^+$, CD31$^-$, CD45$^+$), and epicardial (RFP$^+$, GFP^{++}) cell populations. The sorted RFP$^+$ cells were subsequently cultured on 24-well plates coated with poly-L-lysine (Sigma, P7886) in completed cell culture medium (DMEM high-glucose; Gibco, 11960044) supplemented with L-glutamine (Corning, 25-005-CV), 100 µg/mL streptomycin, 100 U/mL penicillin (Gibco, 15140122), and 10% FBS (Capricorn Scientific, FBS-12B) until reaching approximately 70% confluence.

To induce immortalization, cells were incubated overnight at 37°C with 1.2 µL of SV40 viral vector supernatant (GeneCopoeia, LP721-025). After 24 hours, the supernatant was removed and fresh culture medium was added. Cells were then cultured for an additional 21 days in presence of puromycin (Sigma, P8833). Non-transduced cells served as controls for the immortalization process. Once confluence was achieved, cells were trypsinized and designated as passage one (p1).

The immortalized cells were maintained in complete DMEM culture medium and passaged

every 4 to 5 days. To characterize the cells and assess the stability of marker expression over time, two passage ranges were established: low passage (p6–p9) and high passage (p20–p35).

Isolation of RNA and real-time PCR

Total RNA was isolated using a PureLink RNA Mini Kit (Invitrogen, 12183025) according to the manufacturer's instructions. The purified RNA was reverse transcribed into cDNA using UltraScript® Reverse Transcriptase (PCR Biosystems, PB30.11-10). Relative gene expression levels were quantified using SYBR Master Mix (Promega, A6002) and real-time PCR was performed on 96-well QuantStudio 3 (AppliedBiosystems) plates using the Connect analysis software (Thermo Fisher). The primers used are listed in Table 11.

Immunofluorescence

Immortalized cells were seeded on 12 mm glass coverslips (6,000 cells/cm²) and cultured for 48 hours in complete DMEM culture medium or fibroblast growth medium. Cells were then fixed with 4% paraformaldehyde (Sigma, 158127) for 15 minutes, washed with phosphate buffered saline (PBS), permeabilized with 0.1% Triton X-100 (Sigma, 9036-19-5) for 5 minutes, washed again with PBS, and blocked with PBS supplemented with 1% bovine serum albumin (Sigma, 810664) for 1 hour at room temperature. After blocking, cells were incubated overnight at 4°C with primary antibodies against vimentin (ab8978) (Table 12). After 1 day, the cells were incubated with fluorochrome-conjugated secondary antibodies (Table 12) for 1 hour at 37°C. Cell nuclei were then stained with DAPI (Sigma, D9542). Immunofluorescence images were obtained using a fluorescence microscope (Leica Thunder).

Figure legends

Figure 1. Generation and characterization of freshly isolated cardiac cells from the Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mouse model.

(A) Overview of the protocol for enzymatic digestion and flow cytometry analysis of heart ventricles from Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP. **(B)** Flow cytometry analysis of enzymatically digested P2 heart ventricles from Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mice reveals that a subset of RFP$^+$ cells retains sustained GFP expression. Littermate negative mice were used to establish gating parameters. **(C, D)** Representative flow cytometry plots for identification of RFP$^+$ subsets and corresponding quantification. Data are presented as mean ± s.e.m. (n = 4).

Figure 2. Generation of immortalized cardiac cells from Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mice.

(A) Workflow for the strategy used to generate and immortalize postnatal RFP$^+$ cardiac fibroblasts from Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mice excluding endothelial cells (RFP$^+$,CD31$^+$, CD45$^-$), hematopoietic cells (RFP$^+$CD31$^-$,CD45$^+$) , and cells with the highest GFP expression (RFP$^+$,GFP^{++}). **(B, C)** Bright-field image of primary (B) and immortalized (C) cardiac fibroblasts. **(D)** Immunofluorescence staining for the intermediate filament vimentin, demonstrating abundant vimentin expression in immortalized cells. Scale bars: 200 µm in (B, C); 50 µm in (D).

Figure 3. Characterization of immortalized cardiac fibroblasts from Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mouse model.

(A) Representative flow cytometry plots showing the identification of RFP$^+$ cardiac fibroblast subsets at early passage. **(B)** Representative flow cytometry plots for identification of GFP$^+$ subsets. **(C, D)** qRT-PCR analysis of the indicated genes in immortalized cardiac cells from Wt1$^{GFP/+}$;Wt1Cre;ROSA26-tdRFP mice. Compared with total heart tissue or immortalized epicardial cells (Epi), immortalized fibroblasts exhibited undetectable expression of markers specific to smooth muscle cells, pericytes, cardiomyocytes, and epicardial cells. Data are presented as mean ± s.e.m. (n = 3); P < 0.05, two-tailed student *t*-test.

Figure 4. Immortalized fibroblast populations over cell passages.

(A) Representative flow cytometry plots showing the identification of RFP$^+$ cardiac fibroblast subsets at late passage. **(B)** Representative flow cytometry plots for identification of GFP$^+$ subsets. (**C**) Quantification of the indicated CD90/GFP fibroblast subsets (CD90$^+$GFP$^+$,

CD90$^+$GFP$^-$, CD90$^-$GFP$^+$, and CD90$^-$GFP$^-$) over time in culture at low (L) and high (H) passage. Values are presented as mean ± s.e.m. (n = 3).

Acknowledgments

We are grateful for the assistance provided by the Centres Científics i Tecnològics (CCiTUB) of the University of Barcelona for the use of advanced microscopy, flow cytometry, and animal facilities.

Funding

This work was supported by the Ministerio de Ciencia e Innovación/Agencia Estatal de Investigación with grants to O.M.M.-E. (PID2023-148793NB-I00) and F.X.S. (PID2023-149249NB-I00).

Conflict of interest

The authors declare no conflicts of interest.

References

Acharya A, Baek ST, Banfi S, Eskiocak B, Tallquist MD (2011) Efficient inducible Cre-mediated recombination in Tcf21 cell lineages in the heart and kidney. *Genesis* **49:** 870-877

Acharya A, Baek ST, Huang G, Eskiocak B, Goetsch S, Sung CY, Banfi S, Sauer MF, Olsen GS, Duffield JS, Olson EN, Tallquist MD (2012) The bHLH transcription factor Tcf21 is required for lineage-specific EMT of cardiac fibroblast progenitors. *Development* **139:** 2139-2149

Braitsch CM, Kanisicak O, van Berlo JH, Molkentin JD, Yutzey KE (2013) Differential expression of embryonic epicardial progenitor markers and localization of cardiac fibrosis in adult ischemic injury and hypertensive heart disease. *Journal of molecular and cellular cardiology* **65:** 108-119

Cano E, Carmona R, Ruiz-Villalba A, Rojas A, Chau YY, Wagner KD, Wagner N, Hastie ND, Munoz-Chapuli R, Perez-Pomares JM (2016) Extracardiac septum transversum/proepicardial endothelial cells pattern embryonic coronary arterio-venous connections. *Proceedings of the National Academy of Sciences of the United States of America* **113:** 656-661

Deng Y, He Y, Xu J, He H, Zhang M, Li G (2025) Cardiac Fibroblasts regulate myocardium and coronary vasculature development via the collagen signaling pathway. *bioRxiv : the preprint server for biology*

Feng W, Bais A, He H, Rios C, Jiang S, Xu J, Chang C, Kostka D, Li G (2022) Single-cell transcriptomic analysis identifies murine heart molecular features at embryonic and neonatal stages. *Nature communications* **13:** 7960

Forte E, Skelly DA, Chen M, Daigle S, Morelli KA, Hon O, Philip VM, Costa MW, Rosenthal NA, Furtado MB (2020) Dynamic Interstitial Cell Response during Myocardial Infarction Predicts Resilience to Rupture in Genetically Diverse Mice. *Cell reports* **30:** 3149-3163 e3146

Garvin AM, Katwa LC (2023) Primary cardiac fibroblast cell culture: methodological considerations for physiologically relevant conditions. *American journal of physiology Heart and circulatory physiology* **325:** H869-H881

Hamilton TG, Klinghoffer RA, Corrin PD, Soriano P (2003) Evolutionary divergence of platelet-derived growth factor alpha receptor signaling mechanisms. *Molecular and cellular biology* **23:** 4013-4025

Hastie ND (2017) Wilms' tumour 1 (WT1) in development, homeostasis and disease. *Development* **144:** 2862-2872

Hortells L, Valiente-Alandi I, Thomas ZM, Agnew EJ, Schnell DJ, York AJ, Vagnozzi RJ, Meyer EC, Molkentin JD, Yutzey KE (2020) A specialized population of Periostin-expressing cardiac fibroblasts contributes to postnatal cardiomyocyte maturation and innervation. *Proceedings of the National Academy of Sciences of the United States of America* **117:** 21469-21479

Hosen N, Shirakata T, Nishida S, Yanagihara M, Tsuboi A, Kawakami M, Oji Y, Oka Y, Okabe M, Tan B, Sugiyama H, Weissman IL (2007) The Wilms' tumor gene WT1-GFP knock-in mouse reveals the dynamic regulation of WT1 expression in normal and leukemic hematopoiesis. *Leukemia* **21:** 1783-1791

Humeres C, Frangogiannis NG (2019) Fibroblasts in the Infarcted, Remodeling, and Failing Heart. *JACC Basic to translational science* **4:** 449-467

Ieda M, Tsuchihashi T, Ivey KN, Ross RS, Hong TT, Shaw RM, Srivastava D (2009) Cardiac fibroblasts regulate myocardial proliferation through beta1 integrin signaling. *Developmental cell* **16:** 233-244

Kendall TJ, Duff CM, Boulter L, Wilson DH, Freyer E, Aitken S, Forbes SJ, Iredale JP, Hastie ND (2019) Embryonic mesothelial-derived hepatic lineage of quiescent and heterogenous scar-orchestrating cells defined but suppressed by WT1. *Nature communications* **10:** 4688

Luche H, Weber O, Nageswara Rao T, Blum C, Fehling HJ (2007) Faithful activation of an extra-bright red fluorescent protein in "knock-in" Cre-reporter mice ideally suited for lineage tracing studies. *European journal of immunology* **37:** 43-53

Martinez-Estrada OM, Lettice LA, Essafi A, Guadix JA, Slight J, Velecela V, Hall E, Reichmann J, Devenney PS, Hohenstein P, Hosen N, Hill RE, Munoz-Chapuli R, Hastie ND (2010) Wt1 is required for cardiovascular progenitor cell formation through transcriptional control of Snail and E-cadherin. *Nature genetics* **42:** 89-93

Moore-Morris T, Cattaneo P, Guimaraes-Camboa N, Bogomolovas J, Cedenilla M, Banerjee I, Ricote M, Kisseleva T, Zhang L, Gu Y, Dalton ND, Peterson KL, Chen J, Puceat M, Evans SM (2018) Infarct Fibroblasts Do Not Derive From Bone Marrow Lineages. *Circulation research* **122:** 583-590

Moore-Morris T, Evans SM (2023) Cardiac Pericyte Diversity in Infarct Remodeling: Not Just Vascular Support Cells? *Circulation* **148:** 899-902

Moore-Morris T, Guimaraes-Camboa N, Banerjee I, Zambon AC, Kisseleva T, Velayoudon A, Stallcup WB, Gu Y, Dalton ND, Cedenilla M, Gomez-Amaro R, Zhou B, Brenner DA, Peterson KL, Chen J, Evans SM (2014) Resident fibroblast lineages mediate pressure overload-induced cardiac fibrosis. *The Journal of clinical investigation* **124:** 2921-2934

Moore AW, McInnes L, Kreidberg J, Hastie ND, Schedl A (1999) YAC complementation shows a requirement for Wt1 in the development of epicardium, adrenal gland and throughout nephrogenesis. *Development* **126:** 1845-1857

Patrick R, Janbandhu V, Tallapragada V, Tan SSM, McKinna EE, Contreras O, Ghazanfar S, Humphreys DT, Murray NJ, Tran YTH, Hume RD, Chong JJH, Harvey RP (2024) Integration mapping of cardiac fibroblast single-cell transcriptomes elucidates cellular principles of fibrosis in diverse pathologies. *Science advances* **10:** eadk8501

Perez-Pomares JM, Phelps A, Sedmerova M, Carmona R, Gonzalez-Iriarte M, Munoz-Chapuli R, Wessels A (2002) Experimental studies on the spatiotemporal expression of WT1 and RALDH2 in the embryonic avian heart: a model for the regulation of myocardial and valvuloseptal development by epicardially derived cells (EPDCs). *Developmental biology* **247:** 307-326

Pogontke C, Guadix JA, Sanchez-Tevar AM, Munoz-Chapuli R, Ruiz-Villalba A, Perez-Pomares JM (2022) Dynamic Epicardial Contribution to Cardiac Interstitial c-Kit and Sca1 Cellular Fractions. *Frontiers in cell and developmental biology* **10:** 864765

Quijada P, Misra A, Velasquez LS, Burke RM, Lighthouse JK, Mickelsen DM, Dirkx RA, Jr., Small EM (2019) Pre-existing fibroblasts of epicardial origin are the primary source of pathological fibrosis in cardiac ischemia and aging. *Journal of molecular and cellular cardiology* **129:** 92-104

Ramiro-Pareta M, Muller-Sanchez C, Portella-Fortuny R, Soler-Botija C, Torres-Cano A, Esteve-Codina A, Bayes-Genis A, Reina M, Soriano FX, Montanez E, Martinez-Estrada OM (2023) Endothelial deletion of Wt1 disrupts coronary angiogenesis and myocardium development. *Development* **150**

Ruiz-Villalba A, Simon AM, Pogontke C, Castillo MI, Abizanda G, Pelacho B, Sanchez-Dominguez R, Segovia JC, Prosper F, Perez-Pomares JM (2015) Interacting resident epicardium-derived fibroblasts and recruited bone marrow cells form myocardial infarction scar. *Journal of the American College of Cardiology* **65:** 2057-2066

Sanchez-Iranzo H, Galardi-Castilla M, Sanz-Morejon A, Gonzalez-Rosa JM, Costa R, Ernst A, Sainz de Aja J, Langa X, Mercader N (2018) Transient fibrosis resolves via fibroblast inactivation in the regenerating zebrafish heart. *Proceedings of the National Academy of Sciences of the United States of America* **115:** 4188-4193

Sontake V, Kasam RK, Sinner D, Korfhagen TR, Reddy GB, White ES, Jegga AG, Madala SK (2018) Wilms' tumor 1 drives fibroproliferation and myofibroblast transformation in severe fibrotic lung disease. *JCI insight* **3**

Tallquist MD (2018) Cardiac fibroblasts: from origin to injury. *Current opinion in physiology* **1:** 75-79

Tallquist MD, Molkentin JD (2017) Redefining the identity of cardiac fibroblasts. *Nature reviews Cardiology* **14:** 484-491

Torres-Cano A, Portella-Fortuny R, Muller-Sanchez C, Porras-Marfil S, Ramiro-Pareta M, Chau YY, Reina M, Soriano FX, Martinez-Estrada OM (2022) Deletion of Wt1 during early gonadogenesis leads to differences of sex development in male and female adult mice. *PLoS genetics* **18:** e1010240

Velecela V, Torres-Cano A, Garcia-Melero A, Ramiro-Pareta M, Muller-Sanchez C, Segarra-Mondejar M, Chau YY, Campos-Bonilla B, Reina M, Soriano FX, Hastie ND, Martinez FO, Martinez-Estrada OM (2019) Epicardial cell shape and maturation are regulated by Wt1 via transcriptional control of Bmp4. *Development* **146**

von Gise A, Pu WT (2012) Endocardial and epicardial epithelial to mesenchymal transitions in heart development and disease. *Circulation research* **110:** 1628-1645

Wessels A, van den Hoff MJ, Adamo RF, Phelps AL, Lockhart MM, Sauls K, Briggs LE, Norris RA, van Wijk B, Perez-Pomares JM, Dettman RW, Burch JB (2012) Epicardially derived fibroblasts preferentially contribute to the parietal leaflets of the atrioventricular valves in the murine heart. *Developmental biology* **366:** 111-124

Yata Y, Scanga A, Gillan A, Yang L, Reif S, Breindl M, Brenner DA, Rippe RA (2003) DNase I-hypersensitive sites enhance alpha1(I) collagen gene expression in hepatic stellate cells. *Hepatology* **37:** 267-276

Zhang H, Huang X, Liu K, Tang J, He L, Pu W, Liu Q, Li Y, Tian X, Wang Y, Zhang L, Yu Y, Wang H, Hu R, Wang F, Chen T, Wang QD, Qiao Z, Zhang L, Lui KO, Zhou B (2017) Fibroblasts in an endocardial fibroelastosis disease model mainly originate from mesenchymal derivatives of epicardium. *Cell research* **27:** 1157-1177

Zhou B, Honor LB, He H, Ma Q, Oh JH, Butterfield C, Lin RZ, Melero-Martin JM, Dolmatova E, Duffy HS, Gise A, Zhou P, Hu YW, Wang G, Zhang B, Wang L, Hall JL, Moses MA, McGowan FX, Pu WT (2011) Adult mouse epicardium modulates myocardial injury by secreting paracrine factors. *The Journal of clinical investigation* **121:** 1894-1904

Zhou B, Ma Q, Rajagopal S, Wu SM, Domian I, Rivera-Feliciano J, Jiang D, von Gise A, Ikeda S, Chien KR, Pu WT (2008) Epicardial progenitors contribute to the cardiomyocyte lineage in the developing heart. *Nature* **454:** 109-113

Figure 1

Figure 2

A

Wt1^GFP/+; Wt1Cre; ROSA26-tdRFP

Heart ventricle   collagenase

Antibody staining

Immortalization

waste   colection

RFP⁺,CD31+, CD45⁻
RFP⁺,CD31⁻ CD45⁺
RFP⁺,GFP⁺⁺
RFP⁻

RFP⁺   Puromycin selection

B

C

D

Figure 3

Figure 4

Table 1. Percentage of RFP-positive cells relative to the total number of cells, determined by flow cytometry. Data are mean ± s.e.m. (n=4).

	% RFP/total cells
r1	16,65
r2	12,25
r3	12,85
r4	13,25
Mean	**13,75**
SD	**1,98**

Table 2. Percentage of cells expressing the indicated markers within the RFP-positive cell population, determined by flow cytometry. Data are mean ± s.e.m. (n=4).

	RFP+ cells					
	GFP	**GP38**	**CD90**	**CD39**	**CD31**	**CD45**
r1	20,5	79,90	19,4	30,50	22,60	24,50
r2	24,1	69,50	19,9	30,90	24,70	24,50
r3	23,8	69,20	15,8	31,30	24,50	25,40
r4	12,10	74,40	17	23,40	31,50	34,60
Mean	**20,13**	**73,25**	**18,03**	**29,03**	**25,83**	**27,25**
SD	**5,59**	**5,03**	**1,95**	**3,76**	**3,90**	**4,92**

Table 3. Percentage of GFP cells within the RFP+ CD31+ and RFP+ CD45+ cell population, determined by flow cytometry. Data are mean ± s.e.m. (n=4).

	% GFP+ cells	
	RFP+/CD31+	**RFP+/CD45+**
r1	73,38	64,69
r2	79,96	76,50
r3	76,45	69,65
r4	75,58	67,59
Mean	**76,34**	**69,61**
SD	**2,74**	**5,03**

Table 4. Percentage of cells expressing the indicated markers within the RFP-positive cell population, determined by flow cytometry. Data are mean ± s.e.m. (n=4).

	RFP+/GP38+/			RFP+/CD90+/			RFP+/CD39+/		
	GFP	CD90	CD39	GFP	GP38	CD39	GFP	GP38	CD90
r1	29,69	22,92	27,87	38,94	85,86	34,43	70,97	77,76	21,82
r2	37,09	23,18	24,90	46,52	76,14	45,66	78,65	57,96	8,68
r3	30,81	18,69	24,54	32,06	75,82	28,65	77,70	58,93	9,44
r4	18,06	20,23	16,95	28,33	80,06	38,28	56,84	57,67	18,73
Mean	**28,91**	**21,25**	**23,56**	**36,46**	**79,47**	**36,76**	**71,04**	**63,08**	**14,67**
SD	**7,93**	**2,17**	**4,65**	**8,02**	**4,68**	**7,13**	**10,07**	**9,80**	**6,61**

Table 5. Percentage of immortalized cells at low passages expressing the indicated markers within the RFP-positive cell population, determined by flow cytometry. Data are mean ± s.e.m. (n=3).

	Immortalized cells low passages RFP+			
	GFP	GP38	CD90	CD39
r1	18,62	98,04	64,44	7,79
r2	14,87	97,33	63,95	5,67
r3	12,99	96,84	60,97	4,14
Mean	**15,49**	**97,40**	**63,12**	**5,87**
SD	**2,87**	**0,60**	**1,88**	**1,83**

Table 6. Percentage of immortalized cells at low passages expressing the indicated markers within the RFP-positive cell population, determined by flow cytometry. Data are mean ± s.e.m. (n=3).

	RFP+/GFP+/			RFP+/GP38+/			RFP+/CD90+/			RFP+/CD39+/		
	GP38	CD90	CD39	GFP	CD90	CD39	GFP	GP38	CD39	GFP	GP38	CD90
r1	99,83	78,25	30,27	12,07	66,05	6,63	14,06	99,44	10,96	42,82	89,44	82,18
r2	99,12	70,50	17,16	9,75	66,01	4,69	10,23	99,01	8,14	26,83	91,19	82,60
r3	99,42	79,51	12,27	8,59	63,25	4,06	10,52	99,00	6,26	20,28	88,58	74,09
Mean	**99,46**	**76,09**	**19,90**	**10,14**	**65,11**	**5,13**	**11,61**	**99,15**	**8,45**	**29,98**	**89,73**	**79,62**
SD	**0,36**	**4,88**	**9,31**	**1,77**	**1,61**	**1,34**	**2,13**	**0,25**	**2,36**	**11,59**	**1,33**	**4,80**

Table 7. Percentage of immortalized cells at high passages expressing the indicated markers within the RFP-positive cell population, determined by flow cytometry. Data are mean ± s.e.m. (n=3).

	Immortalized cells high passages RFP+			
	GFP	GP38	CD90	CD39
r1	16,50	99,10	11,40	2,92
r2	19,70	98,70	12,40	1,40
r3	17,00	96,60	11,10	1,19
Mean	17,73	98,13	11,63	1,84
SD	1,72	1,34	0,68	0,94

Table 8. Percentage of immortalized cells at high passages expressing the indicated markers within the RFP-positive cell population, determined by flow cytometry. Data are mean ± s.e.m. (n=3).

	RFP+/GFP+/			RFP+/GP38+/			RFP+/CD90+/			RFP+/CD39+/		
	GP38	CD90	CD39	GFP	CD90	CD39	GFP	GP38	CD39	GFP	GP38	CD90
r1	99,31	32,01	8,08	15,91	9,96	2,33	48,41	99,99	15,05	44,71	99,82	54,61
r2	98,91	17,69	3,74	19,13	10,60	1,04	30,03	99,97	6,63	51,77	99,44	53,90
r3	96,89	18,45	3,26	16,19	9,63	0,87	30,07	99,95	5,57	46,22	99,66	47,90
Mean	98,37	22,72	5,03	17,08	10,06	1,41	36,17	99,97	9,09	47,57	99,64	52,14
SD	1,29	8,06	2,66	1,79	0,49	0,80	10,60	0,02	5,19	3,72	0,19	3,69

Table 9. List of PCR primers used for genotyping

Gene	Primers (5' - … - 3')
Cre	Cre F: GCATTACCGGTCGATGCAACGAGTGATGAG Cre R: GAGTGAACGAACCTGGTCGAAATCAGTGCG Fabpi200 F: TGGACAGGACTGGACCTCTGCTTTCCTAGA Fabpi200 R: TAGAGCTTTGCCACATCACAGGTCATTCAG
Wt1^{GFP}	Wt1-GFP F: GCCTGAAGAACGAGATCAGC Wt1-GFP F2: AGCCTGAAGCTGCTCACATCC Wt1-GFP R: GGCAGCTTGAATTCCTCTCA
RFP	R26g2 F: TGTTATCAGTAAGGGAGCT R26g2 R mut: AAGACCGCGAAGAGTTTGT R26g2 R wt: CACACCAGGTTAGCCTTTA

Table 10. List of antibodies used for flow cytometry analysis

Antibody	Manufacturer	Cat. Number	Dilution
APC Rat anti-mouse CD31 (Clone: MEC 13.3)	BD Pharmigen™	561815	1:1000
PE-Cy™7 Rat anti-mouse CD45 (Clone: 30-F11)	BD Pharmigen™	552848	1:1000
Brilliant Violet 510™ Rat anti-mouse CD90.2 (Thy-1.2)	BioLegend	140319	2 µl per million cells in 100 µl
Brilliant Violet 421™ Syrian Hamster anti-mouse Podoplanin	BioLegend	127423	≤ 0.25 µg per million cells in 100 µl
PE-Cy™7 Rat anti-mouse CD39	BioLegend	143805	≤ 1.0 µg per million cells in 100 µl

Table 11. qRT-PCR primer list

Gene	Primers (5' - … - 3')
Desmin	F: GCGTGACCAACCTGATAGACG R: GTTGGATTTCCTCCTGTAGTT
Rgs5	F: AAGTTGGGAATTCTCCTCCAG R: TTCCTCACTGAATTCAGACTTC
Msln	F: CTGCAGACCCAGACTACAA R: GAGGCCTGTGGGGAGACT
Upk3b	F: CCTGTCCAGATGTGGGATCT R: TCCGAGGATAGTTTGAGAGCA
Vim	F: CCAACCTTTTCTTCCCTGAA R: TGAGTGGGTGTCAACCAGAG
Myh6	F: TGCTCAGAGCTCAAGAAGGAT R: CCCAGCCATCTCCTCTGTTA

Table 12. List of primary and secondary antibodies used for immunofluorescence.

Antibody	Manufacturer	Cat. Number	Species	Dilution
Anti-Vimentin	Abcam	ab8978	Mouse	1:100
Anti-mouse, AlexaFluor™ 546	Invitrogen	A11003	Goat	1:400