Xcr1[+] type 1 conventional dendritic cells are essential mediators for atherosclerosis progression

Running tittle: Targeting of cDC1 cells attenuates atherosclerosis

Tianhan Li[1#], Liaoxun Lu[2#], Juanjuan Qiu[2], Xin Dong[2], Le Yang[2], Kexin He[2], Yanrong Gu[2], Binhui Zhou[2], Tingting Jia[2], Toby Lawrence[3], Marie Malissen[3], Guixue Wang[4], Rong Huang[2], Hui Wang[2], Bernard Malissen[3*], Yinming Liang[2,5*] Lichen Zhang[2*]

[1]*School of Basic Medicine, Xinxiang Medical University, Xinxiang, China.*

[2]*School of Medical Technology, Xinxiang Medical University, Xinxiang, China.*

[3]*Centre d'Immunologie de Marseille-Luminy, INSERM, CNRS, Aix-Marseille Université, Marseille, France*

[4]*Key Laboratory for Biorheological Science and Technology of Ministry of Education, State and Local Joint Engineering Laboratory for Vascular Implants, Bioengineering College of Chongqing University, Chongqing, China*

[5]*Center of Disease Model and Immunology, Hunan Academy of Chinese Medicine, Changsha, China.*

[#]*These authors contributed equally to this work.*

[*]Correspondence: yinming.liang@gris.org.cn (Y L); zhanglichen@gris.org.cn (L Z).

bernardm@ciml.univ-mrs.fr (B M).

Abstract:

Background:

Atherosclerosis is characterized by lipid accumulation within plaques, leading to foam cell

formation and an inflammatory response within the aortic lesions. Lipid disorders have been extensively investigated, however the cellular and molecular mechanisms that trigger the inflammatory response in atherosclerotic plaques remain far from being fully understood. $Xcr1^+$ cDC1 cells are newly identified antigen-presenting cells in activating immune cells. However, the role of cDC1 cells in the development of atherosclerosis remains highly controversial.

Results:

We first confirmed the presence of cDC1 within human atherosclerotic plaques and discovered a significant association between the increasing cDC1 numbers and atherosclerosis progression in mice. Subsequently, we established $Xcr1^{Cre\text{-}Gfp}$ $Rosa26^{LSL\text{-}DTA}$ $ApoE^{-/-}$ mice, a novel and complex genetic model, in which cDC1 was constitutively depleted *in vivo* during atherosclerosis development. Intriguingly, we observed a notable reduction in atherosclerotic lesions in hyperlipidemic mice, alongside suppressed T cell activation of both $CD4^+$ and $CD8^+$ subsets in the aortic plaques. Notably, aortic macrophages and serum lipid levels were not significantly changed in the cDC1-depleted mice. Single-cell RNA sequencing revealed heterogeneity of $Xcr1^+$ cDC1 cells across the aorta and lymphoid organs under hyperlipidemic conditions. As Xcr1 is the sole receptor for Xcl1, we next explored to target $Xcr1^+$ cDC1 cells via Xcl1 by establishing Xcl1 and ApoE deficient mice. Xcl1 knockout mice exhibited decreased atherosclerotic plaque formation and reduced aortic cDC1 accumulation, indicating that Xcl1 contributes to cDC1-mediated atherosclerotic lesion development.

Conclusions:

Our results reveal crucial roles of cDC1 in atherosclerosis progression and provide insights into the development of immunotherapies by targeting cDC1 through Xcl1.

Introduction

Atherosclerosis is a chronic inflammatory disease that affects the large and medium-sized arteries, and it is the leading cause of cardiovascular mortality and morbidity worldwide [1]. Atherosclerosis is a complex pathological process during which lipid-laden phagocytes such as macrophages, along with lymphocytes and dendritic cells, accumulate in the plaque [2]. Mounting evidence from animal models and human studies demonstrates involvement of both the innate and adaptive immunity in the development of atherosclerosis [3-5]. In healthy arteries, DCs are typically present in the subendothelial space and work in the front line of immune surveillance [6, 7], and small portion of resident vascular DCs are localized in the adventitia. Various studies have demonstrated that DCs accumulated in the advanced atherosclerotic lesions in both humans and animal models [8-11]. Recently, several phenotypically and functionally distinct vascular DCs have been identified in mouse and human atherosclerotic lesions [12]. Xcr1$^+$ type 1 conventional dendritic cells (Xcr1$^+$ cDC1) (also known as CD8$^+$ DC or CD103$^+$ DCs) has been recognized for its role in cross-presentation and activating naive CD8$^+$ T cells, and are critical for anti-viral and anti-tumor immune responses [13, 14]. Owing to the challenges associated with obtaining a model for specific cDC1 depletion *in vivo*, previous studies investigating the role of cDC1 cells in atherosclerosis using various knockout mice have produced conflicting results [15-19]. It is important to note that previous Cre lines developed to achieve specific Cre expression in cDC1s still led to leakiness in recombinase activity outside of cDC1s, as demonstrated in previous publications [20, 21]. Moreover, scant mechanistic understanding regarding the regulation of cDC1 activity during atherosclerosis has been reported. Therefore, developing a more specific mouse model for cDC1 is essential for elucidating its precise role in atherosclerosis. Here, we established a novel Cre line Xcr1$^{Cre-Gfp}$ mice

and did not observe any leakiness of Cre activity, and further experiments revealed that specific

depletion cDC1 attenuates atherosclerosis development by inhibiting both $CD4^+$ and $CD8^+$ T cell

activation in hyperlipidemic Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice.

The origin of phagocytes within the atherosclerotic plaques has been controversial. Studies have

shown that the major contribution of macrophages in the aortic plaque is recruited from the

circulation, while other studies have emphasized the significance of local proliferation [22, 23].

Tissue-resident dendritic cells inside the plaques have been well documented [24, 25], regardless of

the fact that the contribution and origin of Xcr1$^+$ cDC1 have not been appropriately addressed. To

understand the origin of Xcr1$^+$ cDC1 inside the plaque, we performed bone marrow transplantation

using Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ mice as donors and found that in ApoE$^{-/-}$ recipient mice placed on

a high-fat diet (HFD), cDC1 cells were completely absent in the aortas. Therefore, our experiments

have elucidated that cDC1 cells in the atherosclerotic plaques originate from the bone marrow.

Further single cell RNA sequencing illustrated that notable differences in Xcr1$^+$ cDC1 cells are

discovered among the aorta, spleen, and lymph nodes. *Ccr2*, *Sept3* and *Cldnd1* were highly

expressed in Xcr1$^+$ cDC1 cells from aorta compared with that from spleen and lymph nodes.

We further aimed to ascertain whether cDC1 can be modulated by targeting the chemokine Xcl1,

which is a ligand of Xcr1. Considering the vital role that Xcr1$^+$ cDC1 plays in the progression of

atherosclerosis and the therapeutic potential of manipulating their functions via chemokines[26],

comprehending the feasibility of targeting Xcl1 holds considerable promise for developing novel

therapeutic approaches to halt the advancement of atherosclerosis. We conducted further studies

using Xcl1$^{-/-}$ ApoE$^{-/-}$ double knockout mice to assess the pathological grade of atherosclerotic

lesions. Our data indicated that the knockout of Xcl1 led to a mild yet significant reduction of cDC1

cells in the aorta. However, strikingly, when compared to the ApoE$^{-/-}$ control mice, the Xcl1$^{-/-}$

ApoE$^{-/-}$ mice exhibited a markedly decreased severity of the atherosclerotic lesions, to an extent

comparable to that observed in cDC1-depleted Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice.

Therefore, our study which employed genetic models to deplete Xcr1^{+} cDC1 *in vivo* in

hyperlipidemic mice, has unequivocally determined the essential role of this specific cell type in the

development of atherosclerosis. Besides their well-established role in cross-presentation to CD8^{+} T

cells, recent studies have shown Xcr1^{+} cDC1 cells also engage with CD4^{+} T cells to augment CD8^{+}

T cell responses [20, 27]. Our data showing Xcr1^{+} cDC1-dependent CD4 and CD8 T cell activation

in atherosclerotic plaques is in line with the new paradigm for cDC1s as a platform for both CD4

and CD8 T cell responses in the novel context of atherosclerosis.

Most importantly, we have successfully demonstrated that targeting the chemokine Xcl1, the ligand

of Xcr1, can effectively inhibit atherosclerosis. These novel findings hold the potential to forge a

novel trajectory for therapeutic interventions within the field of atherosclerosis, presenting

promising prospects for better understanding and treatment of this disorder.

Results:

Progressive accumulation of Xcr1^{+} cDC1 cells in advanced atherosclerotic lesions in human

and mice

Dendritic cells are present in healthy arteries and accumulate within atherosclerotic lesions,

participating in various pathogenic and protective mechanisms during atherogenesis [28, 29].

However, the intra-plaque dynamics of cDC1 and their contribution to atherosclerosis remain

undetermined. Using the archived specimens of paraffin-embedded human atherosclerotic arteries

from our previous study [30], we examined the Xcr1$^+$ dendritic cells in human atherosclerotic

plaques. As shown in **Figure 1A and 1B**, Xcr1$^+$ dendritic cells were significantly more abundant in

the plaque area of the human femoral artery compared to that in the plaque-free area. Similarly, the

immunofluorescence staining results using specific antibodies for Xcr1 and CD11c revealed that

Xcr1$^+$CD11c$^+$ dendritic cells accumulated in the plaques of the aortic root in ApoE$^{-/-}$ mice that were

placed on a HFD for 16 weeks (**Figure 1C and 1D**). Sample preparations entailed sectioning and

appropriate fixation of the aortic root tissues, which were described in greater detail in the Methods

and Materials section. To further elucidate the dynamics of Xcr1$^+$ cDC1 cells within the plaque

during the progression of atherosclerosis, we carried out immunohistochemical analyses of these

cells in the lesioned aortic root (**Figure 1E**). The area positive for Xcr1 expression was quantified

at 8-week on a normal diet (ND), 8-week on HFD, 12-week on HFD and 16-week on HFD,

respectively. Quantitative analysis was conducted using image processing software, revealing a

continuous accumulation of Xcr1$^+$ cDC1 cells in the lesioned area as the disease progressed (**Figure**

1F). Therefore, our experiments provided data indicating the enrichment of Xcr1$^+$ cDC1 cells in

aortic lesions in both humans and mice, and demonstrated that the accumulation of Xcr1$^+$ cDC1

cells was significantly greater during the progression of atherosclerosis.

Xcr1$^{Cre\text{-}Gfp}$ is selectively expressed in cDC1

To understand the role of Xcr1$^+$ cDC1 in regulating the development of atherosclerosis, we

established the Xcr1$^{Cre\text{-}Gfp}$ ApoE$^{-/-}$ mice in which the Cre recombinase and eGFP was co-expressed

under the control of endogenous Xcr1 promoter following start codon (**Figure S1**). This new Cre

line differs from a previous design in which Cre recombinase was expressed via IRES linkage at the

3' UTR and leaked expression of Cre was frequently observed [20, 31]. To verify the fidelity of Cre

expression under hyperlipidemic condition, we used hyperlipidemic ApoE$^{-/-}$ mice as recipient mice

and used bone marrow cells from Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-tdRFP}$ mice as donors for the bone marrow

transplantation (**Figure 2A**). In Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-tdRFP}$ mice, the presence of Cre protein would

turn on the expression of tdRFP as a reporter for the activity of Cre recombinase [32]. Remarkably,

we found that the expression of tdRFP was exclusively present in Xcr1$^+$ cDC1 when mice were fed

a HFD for 16 weeks in the bone marrow transplanted mice, and the Cre recombinase activity was

not detectable in non-cDC1 cells (**Figure 2B and 2C**).

To determine the specificity of GFP expression originating from Xcr1$^{Cre-Gfp}$ genetic modification,

we examined the GFP levels in three populations of DCs, namely pDC, cDC1, and cDC2 cells, from

the spleen of Xcr1$^{Cre-Gfp}$ ApoE$^{-/-}$ mice that were fed a HFD for 16 weeks. As expected, we found

that the GFP expression was stringently restricted to cDC1 cells (**Figure 2D and 2E**). In parallel,

we compared the GFP expression in Xcr1$^+$ cDC1-depleted mice that were fed a HFD for 16 weeks.

Such cDC1-depleted Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice were obtained by crossing Xcr1$^{Cre-Gfp}$

ApoE$^{-/-}$ mice with Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice. Strikingly, we observed a complete loss of GFP-

positive cells and Xcr1$^+$ cDC1 cells in Xcr1$^+$ cDC1-depleted mice (**Figure 2D and 2F**). We also

examined DCs, including pDCs, cDC1, and cDC2, in the lymph nodes and spleens of Xcr1$^{Cre-Gfp}$

Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ and ApoE$^{-/-}$ mice maintained on a ND for 7 weeks. The flow cytometric

analyses also confirmed the absence of Xcr1$^+$ cDC1 population in Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE^{-}

$^{/-}$ mice (**Figure S2**). Collectively, these data suggest that Cre recombinase activity was selectively

confined to cDC1 cells, and no leakiness was observed in lineages other than the Xcr1$^+$ cDC1

population. We also demonstrated that Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice constitute a desirable

model for exploring the role of cDC1 in atherosclerosis.

The specific depletion of Xcr1$^+$ cDC1 cells significantly alleviates atherosclerosis without

influencing the lipid profile and the number of intra-plaque macrophages

To investigate the role of cDC1 cells in atherosclerosis, both Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ and

the control ApoE$^{-/-}$ mice were subjected to a HFD for 16 weeks. The ORO staining results

demonstrated that the size of lesion in both descending aorta and aortic root was significantly

smaller in Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice compared with ApoE$^{-/-}$ control mice (12.82% *vs*

6.01% in descending aorta; 39.25% *vs* 33.57% in ORO-stained aortic root) (**Figure 3A-D**). Given

that macrophages play a pivotal role in the development and progression of atherosclerotic lesions,

particularly within the necrotic core [33, 34], we next investigated macrophage involvement in the

aorta using IHC and flow cytometry. Interestingly, no significant differences were observed in the

necrotic core area of the aortic root between Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice and ApoE$^{-/-}$

control mice (**Figure 3E and 3F**). Similarly, subsequent flow cytometry analyses and IHC results

revealed comparable proportions of F4/80$^+$ CD11b$^+$ macrophages in the aorta, as well as similar

percentages of CD68-positive macrophages in the aortic lesions between the two groups of mice

(**Figure 3G-J**). Furthermore, since dyslipidemia is a key characteristic of HFD-treated ApoE$^{-/-}$ mice

[35, 36], it is critical to assess body weight and lipid profiles in both groups. Our results indicated

no significant differences in body weight or levels of total cholesterol (TC), triglycerides (TG), low-

density lipoprotein (LDL), and high-density lipoprotein (HDL) between the two groups (**Figure**

S3A-B). Additionally, ORO staining of the liver revealed no significant differences either (**Figure**

S3C-D). Taken together, these results suggested that specific depletion of cDC1 attenuated

atherogenesis without affecting lipid status or macrophagic accumulation in aortic lesions.

Specific depletion of cDC1 attenuates the development of atherosclerosis by modulating the activation of T cell in ApoE$^{-/-}$ mice

T cells are among the critical drivers of the pathogenesis of atherosclerosis, and therefore, the triggers of T cell activation in aortic plaques are of great interest for controlling disease progression [2, 37, 38]. We initially analyzed the cDC1 cells through flow cytometry in both Xcr1$^+$ cDC1 depleted mice and control mice. In the Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice that were fed a HFD for 16 weeks, we observed a total absence of Xcr1$^+$ cDC1 in the aorta, lymph nodes, and spleen (**Figure 4A-C and Figure S4**). Subsequently, we assessed T cell phenotype in the two groups of mice. While neither the frequencies nor absolute counts of aortic CD4$^+$ and CD8$^+$ T cells differed significantly between two groups of mice (**Figure 4D-F**), CD69 frequency and CD44 MFI (Mean Fluorescence Intensity), the T cell activation markers, were significantly reduced in both CD4$^+$ and CD8$^+$ T cells from Xcr1$^+$ cDC1 depleted mice compared to controls (**Figure 4G** and **H**). Interestingly, such decreased frequencies of CD69$^+$ T cells were only observed in aortic lesions but not in lymphoid organs such as lymph nodes and the spleen in the Xcr1$^+$ cDC1 depleted mice (**Figure S5A-B and S5D-E**). We further analyzed the absolute numbers of CD8$^+$ and CD4$^+$ T cells in the spleen and lymph nodes and found that the absolute numbers of CD8$^+$ and CD4$^+$ T cells were markedly reduced in the spleen but not in the lymph nodes in Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice compared to those in ApoE$^{-/-}$ mice (**Figure S5C and S5F**). Absence of cDC1 also resulted in reduced frequencies of CD8$^+$ T cells in both spleen and lymph nodes in hyperlipidemic mice. We also analyzed T cell populations in the lymph nodes and spleen in two groups of mice that were fed a ND and found that the frequencies of CD8$^+$ T cells were significantly decreased in the lymph nodes of Xcr1$^+$ cDC1 depleted mice (**Figure S6A-B**). More importantly, the frequencies of CD44^{high}

CD62L^{low} activated memory cells within the CD8$^+$ subsets in both the spleen and lymph nodes were

significantly lower than those in the ApoE$^{-/-}$ control mice (**Figure S6A-D**). However, in the CD4$^+$

subsets, such frequencies of CD44^{high} CD62L^{low} activated cells were comparable in the lymph nodes

but marginally lower in the spleen (**Figure S6A-D**). It is important to note that the decreased

absolute number of CD8$^+$ T cells in the spleen in Xcr1$^+$ cDC1 depleted mice was only observed in

HFD groups but not in ND groups (**Figure S5F and Figure S6F**). Our data indicate that in aortic

plaques, Xcr1$^+$ cDC1 cells are implicated in activating CD4$^+$ and CD8$^+$ T cells, and the loss of the

cDC1 population led to a decreased absolute count of CD8$^+$ T cells in the lymph nodes of HFD

treated mice. Whereas in mice treated with a ND, the loss of the cDC1 population resulted in reduced

frequencies of activated memory T cells but did not affect the absolute counts of T cells. Attenuated

T cell activation might account for the mitigated atherosclerosis in Xcr1$^+$ cDC1 depleted mice.

Characterization of aortic Xcr1$^+$ cDC1 cells in atherosclerotic lesions

Although the majority of cDC1 cells derive from bone marrow, other mechanisms such as local

proliferation of existing DCs can also contribute to the cDC1 population in inflammatory diseases

[39, 40]. It remains unclear whether cDC1 cells in lymphoid organs such as spleen and lymph nodes

represent a different status compared to cDC1 cells in atherosclerotic lesions. To elucidate the source

and status of Xcr1$^+$ cDC1 in atherosclerotic lesions, we performed bone marrow transplantation and

single-cell RNA sequencing (sc-RNA seq) of cDC1 cells from aortic plaques and lymphoid organs.

We conducted bone marrow transplantation using Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ or WT mice that carry

the CD45.2 congenic markers as donors, and CD45.1$^+$ CD45.2$^+$ ApoE$^{-/-}$ mice as recipients (**Figure**

S7A). Our data showed that body weight, lipid profile, and hepatic lipid accumulation were

comparable between two groups of bone marrow transplanted mice after being fed with a HFD for

16 weeks (**Figure S7B-D**). Interestingly, flow cytometric analysis of the aorta displayed almost

complete depletion of cDC1 cells in ApoE$^{-/-}$ recipient mice transplanted with Xcr1$^{Cre-Gfp}$ Rosa26^{LSL-}

^{DTA} bone marrow mice (**Figure S7E**). Strikingly we did not find observe significant changes in

pDCs and CD64$^+$F4/80$^+$ aortic macrophages (**Figure S7E-F**). Therefore, our bone marrow

transplantation experiments further confirmed fidelity of the Cre recombinase expression in cDC1

cells in mice.

Next, we carried out sc-RNA seq to characterize Xcr1$^+$ cDC1 cells from multiple organs of

hyperlipidemic mice. Due to the scarcity of Xcr1$^+$ cDC1 cells that can be sorted by FACS and

processed for sc-RNA seq, we pooled FACS sorted Xcr1$^+$ cDC1 cells from three organs of

hyperlipidemic ApoE-deficient mice fed a HFD for 20 weeks, including the aorta, spleen, and lymph

nodes, and Xcr1$^+$ cDC1 cells from aorta and splenocytes of ApoE-deficient mice fed a ND. Libraries

for sc-RNA seq were prepared using barcode oligos conjugated to anti-CD45 antibodies for surface

staining so that samples from each type of organ could be compared in sequential analyses.

Therefore, we had a total of five samples, and our focus was to understand the differences between

Xcr1$^+$ cDC1 cells from three different organs under hyperlipidemic conditions. The cells were sorted

by gating Xcr1-positive cells out of the MHCII$^+$CD11c$^+$ conventional dendritic cells (**Figure S8**).

Firstly, we obtained the UMAP plot of 10 clusters of cDC1 cells using the Seurat 4.0 package

(**Figure 5A**). In this analysis, 3830 quality-controlled Xcr1$^+$ cDC1 cells were from three types of

organs in hyperlipidemic mice, and an additional 2081 cells were from cDC1 cells sorted from the

spleen and aorta of ApoE deficient mice fed on a ND. The clusters were differentiated according to

top 10 differentially expressed genes (**Supplementary Table 2**), and the heatmap shows marker

genes and heterogeneity among the 10 clusters of cDC1 cells (**Figure 5B**). The purpose of this

experiment was to determine whether cDC1 cells from different organs possess unique features and

whether cDC1 cells from aortic lesions can be characterized by the expression of marker genes. In

hyperlipidemic mice, cDC1 cells from the spleen and lymph nodes differed quite dramatically in

Cluster 2 (*Stk17b* and *Gpr171* highly expressing cells), where 28.9% of cDC1 cells from the spleen

versus 1.0% of cDC1 cells from the lymph nodes were located. Conversely, 3.15% of cDC1 cells

from the spleen versus 32.4% of cDC1 cells from the lymph nodes fell in Cluster 3 (*Ccl5* and *Epsti1*

highly expressing cells), and 4.6% of cDC1 cells from the spleen versus 25.9% of cDC1 cells from

the lymph nodes fell in Cluster 4 (*Slfn5* and *Ifi44* highly expressing cells). Despite the fact that we

had only transcriptomic data for 68 Xcr1$^+$ cDC1 cells from aortic lesions, we were surprised to find

that such aortic cDC1 mainly fell into Cluster 1, 4 and 5, comprising 80.8% of the cells analyzed

(**Figure 5C**). In Cluster 1 (*Hspa1b* and *Dnajb1* highly expressing cells), where aortic cDC1 is

dominantly enriched, the cDC1 cells sorted from lymph nodes were essentially absent, while in

Cluster 4, aortic cDC1 cells were highly enriched and splenic cDC1 was obviously lower in

percentage (**Figure 5C**). Cluster 7 (*Fabp5* and *S100a4* highly expressing cells) is exclusively

comprised of cDC1 cells sourced from lymph nodes, with a striking absence of cDC1 cells from the

spleen and aorta (**Figure 5C**). To better visualize the enrichment of aortic Xcr1$^+$ cDC1 cells

distributed in a few clusters, we separately compared the three samples from the spleen, lymph

nodes, and aorta of hyperlipidemic mice using density plots. Interestingly, the Xcr1$^+$ cDC1 cells

from the spleen and lymph nodes displayed an obvious difference in their distribution, and aortic

Xcr1$^+$ cDC1 cells appeared more enriched in a few clusters (**Figure 5D-E**). Therefore, our data

suggest that Xcr1$^+$ cDC1 cells from the spleen and lymph nodes can differ markedly in heterogeneity,

and aortic Xcr1$^+$ cDC1 cells are enriched in a few clusters that express marker genes, *Ccr2*, *Sept3*

and *Cldnd1* (**Supplementary Table 3**). Using the scRNA seq data of Xcr1$^+$ cDC1 sorted from

hyperlipidemic mice, we annotated the 10 populations as shown in Figure S9A, following the

methodology from a previous study [41]. *Ccr7*$^+$ mature cDC1s (Cluster 3, 7 and 9) and *Ccr7-*

immature cDC1s (remaining clusters) were identified across cDC1 cells sorted from aorta, spleen

and lymph nodes (Figure S9B). Further stratification based on marker genes reveals that Cluster 10

is the pre-cDC1, with high expression level of *CD62L (Sell)* and low expression level of *CD8a*

(Figure S9C). Cluster 6 and 8 are the proliferating cDC1s, which express high level of cell cycling

genes *Stmn1* and *Top2a* (Figure S9D). Cluster 1 and 4 are early immature cDC1s, and cluster 2 and

5 are late immature cDC1s, according to the expression pattern of *Itgae*, *Nr4a2* (Figure S9E). Cluster

9 cells are early mature cDC1s, with elevated expression of *Cxcl9* and *Cxcl10* (Figure S9F). Cluster

3 and 7 as late mature cDC1s, characterized by the expression of *Cd63* and *Fscn1* (Figure S9G). As

shown in Figure 5C and Figure S9, the 10 populations displayed a major difference of aortic cDC1

cells that lack in pre-cDC1s (cluster 10) and mature cells (cluster 3, 7 and 9). Interestingly, in

hyperlipidemic mice splenic cDC1 possess only Cluster 3 as the late mature cells while the lymph

node cDC1 cells have two late mature populations namely Cluster 3 and Cluster 7. In further

analysis, we also compared splenic cDC1 cells from HFD mice to those from ND mice. As shown

in Figure S10, HFD appears to impact early immature cDC1-1 cells (Cluster 1) and increases the

abundance of late immature cDC1 cells (Cluster 2 and 5), regardless of the fact that all 10

populations are present in two origins of samples. We also found that *Tnfaip3* and *Serinc3* are among

the most upregulated genes, while *Apol7c* and *Tifab* are downregulated in splenic cDC1 cells sorted

from HFD mice.

Together, these data implied that Xcr1$^+$ cDC1 in the aorta originate from bone marrow, and Xcr1$^+$

cDC1 cells exhibit strong heterogeneity among aorta, spleen and lymph nodes.

Knockout of Xcl1 attenuates the development of atherosclerosis in ApoE$^{-/-}$ mice

Xcl1 is the major chemokine involved in cDC1 recruitment [42, 43], and plays critical role in

immune response with diverse physical and pathological implication through its interaction with the

sole receptor, Xcr1 [44, 45]. We found the presence of Xcl1 in the lesion area in both human and

mouse atherosclerotic lesions (data not shown). To elucidate the causal relationship between Xcl1

and Xcr1$^+$ cDC1 in the development of atherosclerosis, and more importantly to experimentally

verify Xcl1 as a potential therapeutic target for atherosclerosis treatment, we established the Xcl1

and ApoE double knockout mice. After being fed with a HFD for 19 weeks, both Xcl1$^{-/-}$ ApoE$^{-/-}$

and ApoE$^{-/-}$ mice were sacrificed to assess the severity of atherosclerosis. As shown in **Figure 6A-**

D, the size of lesion at the descending aorta and aortic root was significantly smaller in Xcl1$^{-/-}$

ApoE$^{-/-}$ compared to that in ApoE$^{-/-}$ mice. Importantly, ORO staining of the liver, body weight and

blood lipid profiles revealed no significant differences between the two groups (**Figure 6E-H**).

Moreover, the IHC experiment result show the percentage of CD68 positive macrophages in aortic

lesion was similar between two groups (**Figure 6I-J**).

Previous study identified that Xcl1 serves as a potent chemokine, selectively acting on cDC1 cells

through its exclusive receptor, Xcr1 [43]. Consequently, we examined the DCs and the activation of

T cells in different organs, including aorta, lymph node and spleen. The flow cytometric results

illustrated that both frequencies and absolute counts of Xcr1$^+$ cDC1 cells in the aorta were

significantly reduced, but cDCs and cDC2 cells from Xcl1$^{-/-}$ ApoE$^{-/-}$ were comparable with that

from ApoE$^{-/-}$ (**Figure 7A-C**). Moreover, in both lymph node and spleen, the absolute numbers of

pDC, cDC1 and cDC2 from Xcl1$^{-/-}$ ApoE$^{-/-}$ were comparable with that from ApoE$^{-/-}$ (**Figure S11**).

Crucially, CD69$^+$ frequency and CD44 MFI remained comparable in both aortic CD4$^+$ and CD8$^+$ T cells between two groups (**Figure 7D-F**). However, aortic CD8$^+$ T cells exhibited reduced frequency and absolute count, while CD4$^+$ T cells showed increased frequency but unchanged counts in Xcl1$^{-/-}$ ApoE$^{-/-}$ mouse versus controls (**Figure 7G** and **H**). Furthermore, the absolute numbers of CD4$^+$, CD8$^+$, CD4$^+$ CD69$^+$ and CD8$^+$ CD69$^+$ T cells in both lymph node and spleen from Xcl1$^{-/-}$ ApoE$^{-/-}$ mouse were comparable with that from ApoE$^{-/-}$ mouse (**Figure S12**).

Collectively, these data suggest that Xcl1 promotes atherosclerosis by recruiting Xcr1$^+$ cDC1 cells, and facilitating CD8$^+$ T cell accumulation in lesions. More importantly, *in vivo* data demonstrated the promising potential of targeting Xcl1 to reduce the inflammatory response in atherosclerotic lesions. Our work paves the way to discover novel solutions for the treatment of atherosclerosis by targeting Xcl1 and Xcr1$^+$ cDC1 cells.

Discussion

Atherosclerosis is a complex disorder, and understanding the role of the immune system, particularly dendritic cells (DCs), is of paramount importance [46, 47]. The marked diversity of DC subsets and their distinct, and at times contradictory functions pose challenges in clarifying the role of each specific type of DC in the context of atherosclerosis [48-54]. Moreover, the restricted characterization of each subset of DCs in atherosclerosis, especially when compared to that in well-studied organs such as the skin and lymph nodes [13], has been severely hindered by the limited application of appropriate genetic models and *in vivo* data. Equally significant is that dissecting the functions of each subset of DCs *in vivo* relies on the development of genetic tools validated for lineage specificity. Recently, Xcr1$^+$ cDC1 has been extensively investigated using the Cre line under the promoter of the Xcr1, despite well-documented expression leakiness [20, 31]. Compelling

evidence from tumor and viral infection models [14, 42, 55-57], as well as metabolic disorders like

non-alcoholic steatohepatitis [58], has demonstrated the critical role of Xcr1$^+$ cDC1 cells in diseases.

Nevertheless, the precise role of Xcr1$^+$ cDC1 cells in atherosclerosis remains undetermined.

The precise characterization of Xcr1$^+$ cDC1 and its functions within the atherosclerotic milieu could

provide valuable insights into the immunopathogenesis of the disease. By delineating the specific

contributions of Xcr1$^+$ cDC1, we may identify novel therapeutic targets and develop more targeted

and effective strategies for the prevention and treatment of atherosclerosis. Therefore, investigation

to understand the roles of Xcr1$^+$ cDC1 in the context of atherosclerosis and to identify therapeutic

targets is highly warranted. However, at least two major obstacles impede the *in vivo* study of Xcr1$^+$

cDC1 in atherosclerotic mice. First, Xcr1$^+$ cDC1 cells are extremely scarce in number, and it is quite

challenging to obtain highly purified Xcr1$^+$ cDC1 cells from aortic plaque in hyperlipidemic mice.

Second, there are very few appropriate genetic model systems that permit functional analyses of

Xcr1$^+$ cDC1 *in vivo*. Earlier studies on the role of Xcr1$^+$ cDC1 cells in atherosclerosis have yielded

conflicting results, largely due to the use of different mouse models. For instance, the deletion of

Batf3, Irf8, Dngr1, or Flt3 decreased the cDC1 cells. Nevertheless, Batf3 deficiency had no impact

on atherosclerosis in hyperlipidemic mice. In contrast, Irf8 or Dngr1 deficiency reduced the

development of atherosclerosis, while Flt3 deficiency exacerbated the condition [15-19]. The

recently developed Xcr1-Cre was found to have leakiness of recombinase activity in other immune

cells [20]. Here, we optimized the design for the construction of the Cre line using the endogenous

promoter of Xcr1 and validated its fidelity under hyperlipidemic conditions. We crossed the novel

knock-in line termed Xcr1$^{Cre\text{-}Gfp}$ with Rosa26$^{LSL\text{-}tdRFP}$ Cre reporter mice and used the resulting F1

hybrids as bone marrow donors. The bone marrow cells were then transferred into ApoE$^{-/-}$ mice,

which were subsequently fed on a HFD. We monitored the Cre recombinase activity, which was

reflected by the tdRFP expression of this novel mouse line. Under hyperlipidemic conditions, we

did not observe any leakiness of Cre activity, as tdRFP was strictly restricted to cDC1 cells. To the

best of our knowledge, this novel Cre line is the first of its kind. We further crossed the Xcr1$^{Cre-Gfp}$

and Rosa26$^{LSL-DTA}$ mice with ApoE$^{-/-}$ mice respectively. Subsequently, by further crossing, we

obtained Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice to create a complex genetic model, in which Xcr1^{+}

cDC1 is depleted *in vivo* in the context of atherosclerosis. Very interestingly, in the Xcr1^{+} cDC1

depleted mice, atherosclerosis was significantly ameliorated. Coupled with the data we obtained

from murine and human samples, where Xcr1^{+} cDC1 cells exist in the aortic plaque and their

accumulation is correlated with disease progression, we conclude that Xcr1^{+} cDC1 is essential for

the development of atherosclerosis.

Despite the fact that Xcr1^{+} cDC1 cells are extremely rare in number, we were inclined to understand

if there were any differences among Xcr1^{+} cDC1 cells from different organs. Firstly, we determined

that the Xcr1^{+} cDC1 in the aortic plaque is replenished by circulating precursors using the bone

marrow transfer model. Then we compared, under hyperlipidemic conditions, the Xcr1^{+} cDC1 cells

sorted from the spleen and lymph nodes, supplemented by a small number of but well-purified Xcr1^{+}

cDC1 cells from aortic lesions. To our surprise, the clusters of Xcr1^{+} cDC1 displayed obvious

distinctions among different organs. When the Xcr1^{+} cDC1 cells from aortic lesions were projected

to the clusters, we found that the major subsets present in the lymph nodes, characterized by the

expression of markers *Ccl5 and Fabp5*, and in the spleen, marked by *Stk17b* and *Gpr171*, were

basically absent in the aorta. Meanwhile, compared with spleen and lymph nodes, the aortic Xcr1^{+}

cDC1 cells are highly expressed *Ccr2*, which is an extremely important chemokine receptor

dominating monocyte homing [59-61].

This pronounced tissue-specific compartmentalization of Xcr1$^+$ cDC1 subsets may related to

multiple mechanisms including developmental imprinting that instructs precursor differentiation

into transcriptionally distinct subpopulations [62], and microenvironmental filtering through organ-

specific chemokine axes (e.g., CCL2/CCR2 in spleen) selectively recruits receptor-matched subsets

[63, 64]. This spatial specialization optimizes pathogen surveillance for local immunological

challenges. Based on the maturation analysis of the cDC1 scRNA seq data [41], our findings suggest

that the aortic cDC1 cells display a major difference from those of spleen and lymph nodes by

lacking the mature clusters, whereas lymph node cDC1 cells contain an additional *Fabp5$^+$ S100a4$^+$*

late mature Cluster. Our results also suggest that hyperlipidemia contributes to alteration in early

immature cDC1 and in the abundance of late immature cDC1 cells, which was associated with

dramatic change in gene expression of *Tnfaip3*, *Serinc3*, *Apol7c* and *Tifab*.

In other disease models, Xcr1$^+$ cDC1 cells are crucial for activating T cells [56, 65], and we

confirmed that in hyperlipidemic mice, the depletion of cDC1 substantially attenuated T cell

activation responses. It is important to note that Xcr1$^+$ cDC1 cells are known for well-established

role in cross-presentation to CD8$^+$ T cells, recent studies have shown Xcr1$^+$ cDC1 cells also engage

with CD4$^+$ T cells [20, 27]. Our data showing Xcr1$^+$ cDC1-dependent CD4 and CD8 T cell

activation in atherosclerotic plaques is in line with the new paradigm for cDC1s as a platform for

both CD4 and CD8 T cell responses in the novel context of atherosclerosis. Considering the

multifaceted roles of macrophages in lipid metabolism, inflammation, and plaque dynamics during

the development of atherosclerosis [66], we evaluated the presence of macrophages in

atherosclerotic lesions using IHC and flow cytometry. Our results indicated comparable proportions

of macrophages in the aorta between Xcr1$^+$ cDC1 depleted mice and ApoE$^{-/-}$ control mice.

Additionally, our results also demonstrated that the loss of Xcr1$^+$ cDC1 cells did not affect lipid

status.

Since we elucidated the critical role of Xcr1$^+$ cDC1 cells in atherosclerosis, our aim was to test the

therapeutic potential of inhibiting this specific cell type during disease progression. As Xcr1 is the

sole receptor for Xcl1, and Xcl1 selectively attracts Xcr1$^+$ cDC1 cells in both mice and humans [45,

67], we performed Xcl1 knockout on the background of ApoE deficient mice and assessed the

impact on atherosclerosis development. We found that the Xcl1-Xcr1 axis plays a crucial role in the

progression of atherosclerosis. Our results demonstrated that the loss of Xcl1 in hyperlipidemic

ApoE$^{-/-}$ mice significantly reduced atherosclerotic lesion formation, decreased the frequencies and

absolute coutns of Xcr1$^+$ cDC1 cells and CD8$^+$ T cells in the aorta, without affecting lipid status or

the number of aortic macrophages. Notably, while complete ablation of Xcr1$^+$ cDC1s impaired T

cell activation, reduction of Xcr1$^+$ cDC1 recruitment via Xcl1 deletion did not significantly

compromise this process. This discrepancy may arise through compensatory

mechanisms: alternative chemokine axes (e.g., CCL5/CCR5, CXCL9/CXCR3, BCL9/BCL9L) may

partially rescue Xcr1$^+$ cDC1 homing [13, 68, 69], while non-cDC1 antigen-presenting cells (e.g.,

monocytes, cDC2s) may sustain T cell activation [55, 70]. Furthermore, tissue-specific

microenvironment factors could potentially modulate its role in other diseases. In summary, our

findings identify Xcl1 as a potential therapeutic target for atherosclerosis therapy, though its cellular

origins and regulation of lesional Xcr1$^+$ cDC1 and T cells dynamics require further studies.

Materials and Methods

Animal experiments and patients

All mice were raised in a Specific Pathogen-Free facility and the experiments were conducted in accordance with the guidelines approved by the Animal Ethics Committee of Xinxiang Medical University (China) (Reference No. XYLL-2016S001). The sections of the human femoral artery were obtained in accordance with our previous study [30] and approved by the Ethics Committee of the Third Military Medical University (Project identification code SYXK-PLA-20170005). Rosa26$^{LSL-RFP}$ knock-in mice were generously provided by Hervé Luche at University Clinics Ulm in Germany [32]. C57BL/6 and ApoE$^{-/-}$ mice were purchased from Beijing Vital River Laboratory Animal Technology (China). Rosa26$^{LSL-DTA}$ knock-in mice (Stock Number 009669) were purchased from the Jackson Laboratory (United States of America). Both HFD (contained 20% protein, 20% carbohydrate, 40% fat and 1.25% cholesterol, Cat: XT108C), and ND (contained 20% protein, 64% carbohydrate, 20% fat, Cat: AIN-93G) were purchased from Jiangsu Xietong Pharmaceutical Bioengineering Co., Ltd. 7-8 weeks old mice were used to establish atherosclerosis model through fed on a HFD for 8-20 weeks. On harvest, each sample was assigned an analytical code that was irrelevant to its genotype and processed and analyzed by a researcher blinded to its origin.

Generation of Xcr1$^{Cre-Gfp}$ knock-in mice

The Xcr1$^{Cre-Gfp}$ mouse line was generated by inserting a Cre-P2A-GFP-P2A cassette immediately upstream of the endogenous *Xcr1* translational start codon (ATG). This design enables co-expression of Cre recombinase and GFP under the control of the native *Xcr1* promoter. A 3.2-kb repair template was synthesized (GenScript, Nanjing, China), consisting of a 0.62-kb 5' homology

arm (HA), a codon-optimized *Cre* sequence fused to a P2A peptide, an enhanced GFP reporter

linked to a second P2A site, and a 0.67-kb 3' homology arm. For targeted integration, CRISPR/Cas9-

mediated homology-directed repair (HDR) was performed by co-injecting Cas9 mRNA, a chimeric

sgRNA (sgXcr1-KI), and the repair template into fertilized C57BL/6 zygotes at the one-cell stage,

following established protocols[71, 72]. Founders (F0) harboring the correct knock-in allele were

identified by PCR and Sanger sequencing, then used to establish heterozygous lines. The sgRNA

sequence is provided in **Supplementary Table 1**.

Breeding strategy of Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$

To generate Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, we performed the following crosses: Xcr1^{Cre-}

^{Gfp} mice were crossed with ApoE$^{-/-}$ mice, and offspring heterozygous for both alleles were

intercrossed to obtain homozygous Xcr1$^{Cre-Gfp}$ ApoE$^{-/-}$ mice. Rosa26$^{LSL-DTA}$ mice were crossed with

ApoE$^{-/-}$ mice, and progeny heterozygous for both alleles were intercrossed to generate homozygous

Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice. Finally, homozygous Xcr1$^{Cre-Gfp}$ ApoE$^{-/-}$ mice were crossed with

homozygous Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice to produce Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ offspring.

Generation of Xcl1 knockout mice

The Xcl1 knockout mouse line was generated using CRISPR/Cas9 genome editing through the

microinjection of a mixture containing Cas9 mRNA and two sgRNAs into fertilized C57BL/6

zygotes at the one-cell stage, as previously described [73]. The genotypes were confirmed using

PCR and Sanger sequencing. Subsequently, Xcl1 and ApoE double knockout mice were obtained

by crossing the Xcl1$^{-/-}$ mice with ApoE$^{-/-}$ mice. The sequences of the sgRNA are reported in

Supplementary Table 1.

Bone marrow transplantation

After one week of antibiotic treatment, the recipient $CD45.1^+ CD45.2^+ ApoE^{-/-}$ mice were injected

intraperitoneally with busulfan at a dosage of 30 μg/g every two days for one week. Subsequently,

bone marrow from 8-week-old $CD45.2^+ Xcr1^{Cre-Gfp} Rosa26^{LSL-RFP}$, $CD45.2^+ Xcr1^{Cre-Gfp} Rosa26^{LSL-}$

^{DTA} or WT ($CD45.2^+$ wild-type) donor mice was intravenously transferred to each recipient mouse,

with two million cells administered per recipient. The efficiency of transplantation was assessed

four-week post-transplantation, after which all mice were fed a HFD for 16 weeks. Finally, the mice

were sacrificed for *en face* analysis of the descending aorta, hematoxylin and eosin (H&E) staining,

Oil Red O (ORO) staining of the aortic root, and flow cytometry analysis, as previously described

[74].

Immunofluorescence

Frozen sections of the aortic root from mice were incubated at 60 °C for 2 minutes and then fixed

in 4% paraformaldehyde for 30 minutes. The samples were subsequently washed with PBS and

permeabilized/blocked with 0.1% Triton X-100 in 5% BSA. Next, the primary antibodies, including

anti-Xcr1 (BD Biosciences, Cat. 148212) and anti-CD11c (BD Biosciences, Cat. 12-0114-83), were

applied in a wet box at 4 °C overnight. After washing with PBST (PBS containing 0.1% Tween-20)

for five times, the samples were incubated with secondary antibodies for 1 hour, followed by a 10-

minute incubation with DAPI at room temperature (protected from light). The fluorescent signals

of the sections were detected using a Nikon C2 confocal microscope (Nikon, Japan).

Morphometric and immunohistochemical analyses

After modeling, the heart and aorta tissue were obtained from mice. The *en face* descending aortas

were fixed with 4% paraformaldehyde and subsequently stained with ORO. Additionally, frozen

sections of the aortic roots were stained with ORO and H&E to assess the severity of atherosclerosis,

and frozen sections of livers were stained with ORO to assess the severity of lipid accumulation, in

accordance with our previous study [74]. Paraffin-embedded sections of the aortic root were

dewaxed in xylene twice for 10 minutes, rehydrated through descending grades of ethanol, and

subjected to sodium citrate-induced antigen retrieval for 10 minutes at 95 °C. The sections were

then incubated with 3% H_2O_2 for 10 minutes at room temperature. Following this, the sections were

blocked and incubated with primary antibodies, either anti-Xcr1 (NBP1-02343) or CD68

(FB113109). After five washes with DPBS, the sections were incubated with secondary antibodies

for 30 minutes and stained with DAB at room temperature. Subsequently, the sections were

counterstained with hematoxylin, dehydrated in ethanol, and mounted with neutral balsam. Images

were captured using the digital pathology system (Pannoramic MIDI, Hungary), viewed using

CaseViewer2.4 software and analyzed using ImageJ software.

Flow cytometry analysis

To obtain the single-cell suspension, aortas, lymph nodes and spleens were collected from each

mouse. Spleens were directly dissociated using the Gentle MACS Dissociators (Miltenyi Biotec).

Aortas and lymph nodes were cut into small pieces and then dissociated using the Gentle MACS

Dissociators. Dissociated tissues digested with digestive enzymes (Collagenase IV, Cat: V900893

from Sigma; DNase I, Cat: DN25 from Sigma; Liberase DH, Cat: 5401054001 from Roche) for 30

minutes at 37 °C using a shaker set to 300 rpm or a heating block. The resulting cell suspension was

filtered through 75 μm strainers and subsequently blocked with 2.4G2 for 20 minutes at 4 °C,

followed by washing with stain buffer (BD, Cat: 554656). Then samples were stained with

monoclonal antibody mixes for 30 minutes at 4 °C followed by washing with stain buffer. Finally,

cells were resuspended in stain buffer containing 200 nM Sytox blue (ThermoFisher Scientific, Cat:

S34857) and acquired using a flow cytometer (ThermoFisher Scientific, Invitrogen Attune NxT

Flow Cytometer), and the FACS data were analyzed using FlowJo software. The following

antibodies purchased from BioLegend, BD or eBioscience were used to detect surface markers by

flow cytometry: anti-CD45.2 (104) conjugated to eFluor 450, FITC or PE-Cy7; anti-CD45.1 (A20)

conjugated to PE or eFluor450; anti-CD45 (30-F11) conjugated to PE; anti-CD11b (M1/70)

conjugated to super Bright 600, BUV395 or biotin; anti-CD3e (145-2C11) conjugated to PE-Cy5.5

or PE-Cy7; anti-CD3e (eBio500A2) conjugated to Alexa Fluor700; anti-CD19 (eBio1D3)

conjugated to PE-Cy5.5 or biotin; anti-Xcr1 (ZET) conjugated to APC or BV650; anti-Ly6G (1A8)

conjugated to Alexa Fluor 700; anti-MHCII (M5/114.15.2) conjugated to APC-eFluor780; anti-

CD11c (N418) conjugated to PE or PE-Cy5.5; anti-Siglec H (eBio440c) conjugated to PE-Cy7 or

PE; anti-CD8b (eBioH35-17.2) conjugated to eFluor 450; anti-CD44 (IM7) conjugated to BV605;

anti-CD4 (RM4-5) conjugated to PE-Cy5.5; anti-CD69 (H1.2F3) conjugated to PE-Cy7; anti-

CD62L (MEL-14) conjugated to APC; anti-CD5 (53-7.3) conjugated to APC-eFluor780 or biotin;

anti-Siglec F (E50-2440) conjugated to BV421; anti-F4/80 (BM8) conjugated to PE-Cy7; anti-

CD64 (X54-5/7.1) conjugated to APC; anti-NK1.1 (PK136) conjugated to Alexa Fluor 700; anti-

CD172α (P84) conjugated to FITC.

Single cell RNA sequencing

Eight-week-old male ApoE$^{-/-}$ mice were fed with 20-week ND or HFD. The single cell suspension

of spleen, lymph node and aorta were obtained according to the flow cytometry analysis section.

Single cell suspension from spleens and lymph nodes were firstly removed the CD5^{+}, CD19^{+} and

CD11b$^+$ cells via SAV conjugated beads. Then two million live cells from spleen and lymph node

were stained with mix1 (CD45.2-PE-Cy7, CD11c-PE-Cy5.5, CD172α-FITC, MHCII- APC-

eFluor780 and Xcr1-APC), and all cells from aorta were stained with mix2 (CD45.2-PE-Cy7,

CD11b-super bright 600, CD11c-PE-Cy5.5, MHCII- FITC, CD3- Alexa Fluor 700, CD19- Alexa

Fluor 700, Ly6G- Alexa Fluor 700 and Xcr1-APC) for 30 minutes at 4 °C. After washing with 1 mL

FACS buffer two times, cells from different organs were stained with different anti-CD45 antibodies

conjugated with barcode oligos from BD™ Ms Single Cell Sample Multiplexing Kit (BD

Bioscience, Cat: 633793). Finally, cells were resuspended with Sytox blue and CD11c$^+$ MHCII$^+$

Xcr1$^+$ cells were sorted and subjected to the BD Rhapsody Express system. Then cDNA and sample

tag libraries were built with BD Rhapsody whole transcriptome amplification (WTA) reagent kit

(BD Bioscience, Cat: 633733, 633773, 633801, 664887) following manufacturer's instructions, and

sequenced on an illumine Novaseq 600 sequencer. Pair-end Fastq files of sample tag and WTA data

were processed via BD Rhapsody™ analysis pipeline v2.0, and the resultant dataset was mainly

analysis using SeqGeq™ software, which contain Lex-BDSMK and Seurat v4.04 plugin

components. Raw data and processed data were uploaded into GEO (Accession number:

GSE279370).

Statistical analysis

All data were presented as means ± SEM (Standard Error of the Mean), reflecting the average values

of biological measurements across multiple samples. Data analysis was performed using GraphPad

Prism software (version 8.0). Statistical significance was assessed by both parametric (unpaired

Student's *t*-test) and nonparametric (Mann–Whitney test) methods when comparing two groups.

The assumptions of the tests were checked to ensure appropriate application. Significant differences

between different groups were set at * $P<0.05$, ** $P<0.01$, *** $P<0.001$ and **** $P<0.0001$.

Graphical representations of the data will include bar graphs with error bars denoting SEM.

Acknowledgements

548　We thank the animal facility and flow cytometry facility of GRIS in Xinxiang Medical University

549　for providing technical support and assistance. Generation of murine genetic models were supported

550　by 111 program (D20036).

Author Contributions

Lichen Zhang and **Yinming Liang** conceived the project. **Tianhan Li** and **Liaoxun Lu** wrote the

manuscript and carried out the majority of the experiments. **Bernard Malissen** and **Lichen Zhang**

designed the Xcr1Cre-Gfp knock-in mice and Xcl1 knockout mice. **Juanjuan Qiu**, **Xin Dong**, **Le**

Yang, **Kexin He**, and **Yanrong Gu** assisted in the experimental procedures. **Binhui Zhou**

performed cell sorting for scRNA sequencing. **Tingting Jia** and **Rong Huang** contributed to data

curation and validation. **Toby Lawrence**, **Marie Malissen**, and **Hui Wang** participated in the

manuscript review and editing. **Guixue Wang** assisted in the preparation of human samples.

Yinming Liang, **Lichen Zhang**, and **Tianhan Li** contributed to obtaining the funding for the

project.

Sources of Funding

562　This research was funded by grants from Science and Technology Innovation Program of Hunan

563　Province [2023DK2005], [2024RC1071] and [2022RC1223], and NSFC grants 32170879 and

564　82301972, and Science and Technology Department of Henan Province 242102310030.

Declaration of competing interest

The authors declare no competing interests.

Figure legends

Figure 1. Progressive accumulation of Xcr1$^+$ cDC1 cells in human and mouse advanced atherosclerotic lesions

A, Representative images of immunohistochemical staining demonstrate Xcr1 expression in normal areas and plaque areas of the femoral artery from three cardiovascular disease patients. **B**, Quantitative analysis of the Xcr1-positive area in normal area and plaque regions. **C**, Representative immunofluorescence images depict DAPI (blue), Xcr1 (green) and CD11c (red) within the lesions of aortic root of ApoE$^{-/-}$ mice fed with 16-week ND or HFD. **D**, Quantitative analysis of Xcr1 and CD11c double positive area in lesion area (n = 6). Scale bars, 100 μm. **E**, Representative images depicting immunohistochemical staining of Xcr1 in lesions of the aortic root of ApoE$^{-/-}$ mice fed a HFD for varying durations. **F**, Quantitative analysis of the Xcr1-positive area in the lesion regions. (8-week HFD, n = 11; 12-week HFD, n = 15; 16-week HFD, n = 29). Data represent as mean ± SEM. * $P<0.05$. *** $P<0.001$.**** $P<0.0001$.

Figure 2. Selective expression of Xcr1$^{Cre-Gfp}$ in cDC1 cells.

A, Diagram illustrating the bone marrow transfer process. **B** and **C**, Representative flow cytometric analysis and corresponding quantification of GFP$^+$RFP$^+$ in cDC1 and non-cDC1 populations within the cDC1 and non-cDC1 populations of the spleen. These data were obtained from ApoE$^{-/-}$ mice that had received bone marrow transplants from either WT or Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-RFP}$ donors and were maintained on a HFD for 16 weeks (WT, n = 4; Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-RFP}$, n = 3). **D** and **E**, Representative flow cytometric analysis and quantification of the percentage of GFP$^+$ cells among pDC, cDC1 and cDC2 cells populations in the spleen. **F**, Quantification of the percentage of pDC,

cDC1 and cDC2 cells in spleen. (D-F, ApoE$^{-/-}$, Xcr1$^{Cre-Gfp}$ ApoE$^{-/-}$ and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$

ApoE$^{-/-}$ mice fed with a HFD for 16 weeks, n = 3). Data represent as mean ± SEM. **** $P<0.0001$;

NS, non-significant.

Figure 3. Specific depletion of Xcr1$^+$ cDC1 cells in ApoE$^{-/-}$ mice reduces atherosclerosis

progression with no effect on macrophages.

A, ORO staining of the descending aortas. Scale bar, 5 mm. **B**, Quantification of the lesion area in

the descending aorta (ApoE$^{-/-}$ mice, n = 9; Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 6). **C**, ORO

and H&E staining of the aortic roots of representative mice from each group. **D**, Quantification of

lesion area in the aortic root. (ApoE$^{-/-}$ mice, n = 7; Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 8).

Scale bar, 200 μm. **E** and **F**, H&E staining and quantification of necrotic core area in the lesions of

aortic root. (ApoE$^{-/-}$ mice, n = 7; Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 8). Scale bar, 200 μm.

G and **H**, Representative flow cytometric analysis and quantification of macrophages in the aorta.

(ApoE$^{-/-}$ mice, n = 8; Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 6). **I** and **J**, Representative

immunohistochemical staining images of CD68 and quantification of CD68 positive area in the

lesions of the aortic root. (ApoE$^{-/-}$ mice, n = 6; Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 8). Scale

bar, 100 μm. ApoE$^{-/-}$ and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice were fed on a 16-week HFD to

develop atherosclerosis. Data represent as mean ± SEM. NS, non-significant. ** $P<0.01$. ***

$P<0.001$.

Figure 4. Specific depletion of Xcr1$^+$ cDC1 cells in ApoE$^{-/-}$ mice decreases T cell activation

within aorta.

Figure 5. Sc-RNA sequencing analysis of the aorta, lymph nodes and spleen in ApoE$^{-/-}$ mice

fed with a 20-week HFD.

A, UMAP plot delineates 10 annotated cell types of Xcr1$^+$ cDC1 cells from aorta, lymph node and

spleen in ApoE$^{-/-}$ mice maintained on a HFD or ND for 20 weeks. **B**, Heatmap statistic map displays

the top ten up-regulated genes across various clusters. **C**, The proportion of 10 clusters among

groups. **D** and **E**, UMAP plot representing Xcr1$^+$ cDC1 cells from aorta, lymph nodes, spleen, and

a merged plot in ApoE$^{-/-}$ mice maintained on a HFD for 20 weeks.

Figure 6. Xcl1 deficiency reduces atherosclerotic lesion formation in ApoE$^{-/-}$ mice.

A, ORO staining of the descending aortas. Scale bar, 5 mm. **B**, Quantification of the lesion area in

aortas. **C**, ORO and H&E staining of aortic roots. Scale bar, 200 μm. **D**, Quantification of the lesion

area in aortic roots. **E** and **F**, Representative liver images and quantification of ORO positive area

in livers (n = 24 regions from 8 mice, 3 regions per mice). Scale bar, 100 μm. **G**, The concentrations

of TC, TG, LDL and HDL in the serum. **H**, Body weight of mice before and after feeding with 19-

634 week HFD. **I**, Representative immunohistochemical staining of CD68 in aortic roots. **J**,

635 Quantification of CD68 positive area in the lesions of aortic roots (n = 24 lesions from 8 mice, 3

636 lesions per mice). Scale bar, 100 μm. ApoE$^{-/-}$ mice and Xcl1$^{-/-}$ ApoE$^{-/-}$ mice fed a HFD for 19

637 weeks (n = 8 per group). Data represent as mean ± SEM. ** *P*<0.01, *** *P*<0.001, NS, non-

638 significant.

640 **Figure 7. Xcl1 deficiency reduces CD8$^+$ T cells in the aortas from ApoE$^{-/-}$ mice.**

641 **A**, Representative flow cytometric analysis of cDC, cDC1 and cDC2 cells in aortas. **B** and **C**,

642 Quantification of the frequencies and absolute counts of cDC, cDC1 and cDC2 cells in aortas (n =

643 7). **D**, Representative flow cytometric analysis of T cell subsets in aortas, including CD3e$^+$ CD5$^+$,

644 CD4$^+$, CD4$^+$ CD69$^+$, CD8$^+$ and CD8$^+$ CD69$^+$ T cells. **E**, Quantification of the frequencies of

645 CD4$^+$CD69$^+$ and CD8$^+$ CD69$^+$ in aortas. **F**, Quantification of the MFI of CD44 in CD4$^+$ and CD8$^+$

646 T cells in aortas. **G** and **H**, Quantification of the frequencies and absolute counts of CD3e$^+$ CD5$^+$,

647 CD4$^+$ and CD8$^+$ cells in aortas. (D-H, ApoE$^{-/-}$ mice (n = 5) and Xcl1$^{-/-}$ApoE$^{-/-}$ mice (n = 6) fed

648 with a 16-week HFD). Data represent as mean ± SEM. * *P*<0.05, NS, non-significant.

650 **Figure S1. Schematic diagram illustrating the knock-in of the 5'HA-iCre-P2A-EGFP-P2A-**

651 **3'HA vector into the mouse Xcr1 locus.**

653 **Figure S2. Flow cytometric analysis of cDC1 cells in the lymph nodes and spleens of ApoE$^{-/-}$**

654 **mice and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice fed a 7-weeks ND.**

655 **A**, Representative flow cytometric analysis of cDC1 cells in lymph nodes. **B**, Quantification of the

frequencies and number of cDC1 cell in lymph nodes. **C**, Representative flow cytometric analysis

cDC1 cells in the spleens. **D**, Quantification of the frequencies and number of cDC1 cell in the

spleens. ApoE$^{-/-}$ mice and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice fed with 7-week ND, n = 7. Data

represent the mean ± SEM. *** *P*<0.001.

Figure S3. Lipid profile, body weight and ORO staining of the liver in ApoE$^{-/-}$ mice and

Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice fed a 16-week HFD.

A, Concentrations of TC, TG, LDL and HDL in the serum. **B**, Body weight measurements of both

groups of mice. **C**, ORO staining of liver from each group. Scale bar, 100 μm. **D**, Quantification of

ORO positive area in liver. ApoE$^{-/-}$ mice (n =7) and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice (n =8)

were fed on a 16-week HFD to develop atherosclerosis. Data represent the mean ± SEM. NS, non-

significant.

Figure S4. Flow cytometric analysis of cDC1 cells in the lymph nodes and spleens of ApoE$^{-/-}$

and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice fed a 16-week HFD.

A, Representative flow cytometric analysis of cDC1 cells in lymph nodes. **B**, Quantification of the

frequencies and numbers of cDC1 cells in lymph nodes. **C**, Representative flow cytometric analysis

of cDC1 cells in the spleens. **D**, Quantification of the frequencies and numbers of cDC1 cells in the

spleens. ApoE$^{-/-}$ mice (n =7) and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice (n =8) were fed on a 16-

week HFD. Data represent the mean ± SEM. *** *P*<0.001.

Figure S5. Flow cytometric analysis of T cells in the lymph nodes and spleens of ApoE$^{-/-}$ mice

and Xcr1^{Cre-Gfp} Rosa26^{LSL-DTA} ApoE^{−/−} mice fed a 16-week HFD.

A, Representative flow cytometric analysis of $CD4^+$, $CD8^+$, $CD4^+CD69^+$ and $CD8^+CD69^+$ T cells

in lymph nodes. **B** and **C**, Quantification of the frequencies and numbers of $CD4^+$, $CD8^+$, $CD4^+$

$CD69^+$ and $CD8^+ CD69^+$ T cells in lymph nodes. **D**, Representative flow cytometric analysis and

quantification of $CD4^+$, $CD8^+$, $CD4^+ CD69^+$ and $CD8^+ CD69^+$ T cells in the spleens. **E** and **F**,

Quantification of the frequencies and numbers of $CD4^+$, $CD8^+$, $CD4^+ CD69^+$ and $CD8^+CD69^+$ T

cells in the spleens. ApoE^{−/−} mice (n =7) and Xcr1^{Cre-Gfp} Rosa26^{LSL-DTA} ApoE^{−/−} mice (n =8) were

fed on a 16-week HFD. Data represent the mean ± SEM. * $P<0.05$, ** $P<0.01$, *** $P<0.001$, NS,

non-significant.

Figure S6. T cells analysis in the lymph nodes and spleens of ApoE^{−/−} mice and Xcr1^{Cre-Gfp}

Rosa26^{LSL-DTA} ApoE^{−/−} mice fed a 7-week ND.

A, Representative flow cytometric analysis and quantification of $CD4^+$, $CD4^+$ memory, $CD8^+$ and

$CD8^+$ memory T cells in lymph nodes. **B**, Quantification of the frequencies of $CD4^+$, $CD8^+$,

$CD4^+CD69^+$ and $CD8^+CD69^+$ T cells in the lymph nodes. **C**, Representative flow cytometric

analysis and quantification of $CD4^+$, $CD4^+$ memory, $CD8^+$ and $CD8^+$ memory T cells in the spleens.

D, Quantification of the frequencies of $CD4^+$, $CD8^+$, $CD4^+ CD69^+$ and $CD8^+ CD69^+$ T cells in the

spleens. **E** and **F**, Quantification of the numbers of $CD4^+$ and $CD8^+$ T cells in the lymph nodes and

spleens. ApoE^{−/−} mice and Xcr1^{Cre-Gfp} Rosa26^{LSL-DTA} ApoE^{−/−} mice fed with a 7-week ND (n = 6).

Data represent the mean ± SEM. * $P<0.05$, ** $P<0.01$, NS, non-significant.

Figure S7. Depletion of Xcr1⁺ cDC1 cells of bone marrow reduces cDC1 cells without affecting macrophages in atherosclerotic lesion of ApoE⁻/⁻ mice.

A, Diagram of the bone marrow transfer process. ApoE⁻/⁻ mice transplanted with bone marrow from WT or Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ donors (n = 7). **B**, Body weight of each group. **C**, Serum concentrations of TC, TG, LDL and HDL. **D**, Representative ORO staining images of liver sections and quantification of the ORO positive area percentage in the liver. Scale bar, 50 μm. **E**, Representative flow cytometric analysis of pDC, cDC1 and cDC2 cells in the aortas, Lin⁻ means CD3⁻CD19⁻Ly6G⁻NK1.1⁻. **F**, Flow cytometric analysis of macrophages in the aortas. Data represent as mean ± SEM. *** $P<0.001$, NS, non-significant.

Figure S8. The single-cell RNA sequencing process.

Figure S9. Sc-RNA sequencing analysis of the aorta, lymph nodes and spleen in ApoE⁻/⁻ mice.
A, UMAP plot delineates 10 annotated cell types of Xcr1⁺ cDC1 cells from aorta, lymph node and spleen in ApoE⁻/⁻ mice maintained on a HFD or ND for 20 weeks. **B-G**, Heatmap statistic map showing marker genes in different clusters. *Ccr7* is the mature gene marker (B). *Sell*⁺*CD8a*⁻ cDC1s are defined as Pre-cDC1s (C). *Stmn1* and *Top2a* are the proliferating markers (D). *Itgae* and *Nr4a2* genes marked the transition from early to late immature cDC1s (E). *Cxcl9* and *Cxcl10* genes marked the transition from late immature to early mature cDC1s (F). *Cd63* and *Fscn1* are the late mature markers (G).

Figure S10. Single-cell RNA sequencing of splenic cDC1s from both the HFD and ND groups.

, UMAP plot of splenic cDC1s showing annotated clusters. **B**, The proportions of cDC1

maturation states across diet groups. **C-H**, Heatmap statistic map depicting representative markers

defining cDC1 maturation clusters. *Ccr7* is the mature gene marker (C). *Sell*$^+$*CD8a*$^-$ cDC1s are

defined as Pre-cDC1s (D). *Top2a*, *Stmn1* and *Mki67* are the proliferating markers (E). *Itgae*, *Nr4a2*,

Nr4a3 and *Egr3* genes marked the transition from early to late immature cDC1s (F). *Cxcl9* and

Cxcl10 genes marked the transition from late immature to early mature cDC1s (G). *Cd63* and *Fscn1*

are the late mature markers (H). **I-J**, Heatmap statistic map of upregulated genes

(*Tnfaip3* and *Serinc3*), and downregulated genes (*Apol7c* and *Tifab*) in the HFD spleen compared

to the ND spleen.

Figure S11. DC cells analysis in lymph nodes and spleens of ApoE$^{-/-}$ and Xcl1$^{-/-}$ApoE$^{-/-}$ mice

fed with 19-week HFD.

A, Representative flow cytometric analysis and quantification of pDC, cDCs, cDC1 and cDC2 cells

in lymph nodes. **B**, Quantification of absolute counts of pDC, cDCs, cDC1 and cDC2 cells in lymph

nodes. **C**, Representative flow cytometric analysis of pDC, cDC1 and cDC2 cells in spleens. **D**,

Quantification of absolute counts of pDC, cDCs, cDC1 and cDC2 cells in spleens. ApoE$^{-/-}$ (n = 9)

and Xcl1$^{-/-}$ApoE$^{-/-}$ mice (n = 7 or 8) fed a HFD for 19 weeks. Data represent the mean ± SEM. NS,

non-significant.

Figure S12. T cells analysis in lymph node and spleen of ApoE$^{-/-}$ and Xcl1$^{-/-}$ApoE$^{-/-}$ mice fed

with 16-week HFD.

A and **B**, Representative flow cytometric analysis and quantification of absolute counts of CD4$^+$,

CD8$^+$, CD4$^+$CD69$^+$ and CD8$^+$CD69$^+$ T cells in lymph nodes (ApoE$^{-/-}$, n = 9; Xcl1$^{-/-}$ ApoE$^{-/-}$, n =

7). **C** and **D**, Representative flow cytometric analysis and quantification of absolute counts of CD4$^+$,

CD8$^+$, CD4$^+$CD69$^+$ and CD8$^+$CD69$^+$ T cells in spleens (ApoE$^{-/-}$, n = 10; Xcl1$^{-/-}$ ApoE$^{-/-}$, n = 7).

Data represent the mean ± SEM. NS, non-significant.

Supplementary table 1: **The sequences of the sgRNA and primers used for generation and**

validation of Xcr1$^{Cre-Gfp}$ knock-in mice and Xcl1 knockout mice

Supplementary table 2: **Up regulated genes in Xcr1$^+$ cDC1 cells from aorta vs spleen, and**

aorta vs lymph node in ApoE-deficient mice fed a HFD for 20 weeks

Supplementary table 3: **The top ten high-expressed marker genes in ten clusters**

References

1.   Bjorkegren JLM, Lusis AJ. Atherosclerosis: Recent developments. Cell. 2022;185(10):1630-45. doi:
10.1016/j.cell.2022.04.004. PubMed PMID: 35504280; PubMed Central PMCID: PMC9119695.
2.   Hansson GK, Hermansson A. The immune system in atherosclerosis. Nat Immunol.
2011;12(3):204-12. Epub 2011/02/16. doi: 10.1038/ni.2001. PubMed PMID: 21321594.
3.   Lusta KA, Poznyak AV, Sukhorukov VN, Eremin, II, Nadelyaeva, II, Orekhov AN. Hypotheses on
Atherogenesis Triggering: Does the Infectious Nature of Atherosclerosis Development Have a
Substruction? Cells. 2023;12(5). doi: 10.3390/cells12050707. PubMed PMID: 36899843; PubMed
Central PMCID: PMC10001176.
4.   Hansson GK, Robertson AK, Soderberg-Naucler C. Inflammation and atherosclerosis. Annu Rev
Pathol. 2006;1:297-329. Epub 2007/11/28. doi: 10.1146/annurev.pathol.1.110304.100100. PubMed
PMID: 18039117.
5.   Hasham SN, Pillarisetti S. Vascular lipases, inflammation and atherosclerosis. Clin Chim Acta.
2006;372(1-2):179-83. Epub 2006/06/13. doi: 10.1016/j.cca.2006.04.020. PubMed PMID: 16765928.
6.   Steinman RM, Hawiger D, Liu K, Bonifaz L, Bonnyay D, Mahnke K, et al. Dendritic cell function
in vivo during the steady state: a role in peripheral tolerance. Ann N Y Acad Sci. 2003;987:15-25. Epub

2003/05/03. doi: 10.1111/j.1749-6632.2003.tb06029.x. PubMed PMID: 12727620.

7.   Ma-Krupa W, Jeon MS, Spoerl S, Tedder TF, Goronzy JJ, Weyand CM. Activation of arterial wall dendritic cells and breakdown of self-tolerance in giant cell arteritis. J Exp Med. 2004;199(2):173-83. Epub 2004/01/22. doi: 10.1084/jem.20030850. PubMed PMID: 14734523; PubMed Central PMCID: PMCPMC2211768.

8.   Liu P, Yu YR, Spencer JA, Johnson AE, Vallanat CT, Fong AM, et al. CX3CR1 deficiency impairs dendritic cell accumulation in arterial intima and reduces atherosclerotic burden. Arterioscler Thromb Vasc Biol. 2008;28(2):243-50. Epub 2007/12/15. doi: 10.1161/ATVBAHA.107.158675. PubMed PMID: 18079406.

9.   Jongstra-Bilen J, Haidari M, Zhu SN, Chen M, Guha D, Cybulsky MI. Low-grade chronic inflammation in regions of the normal mouse arterial intima predisposed to atherosclerosis. J Exp Med. 2006;203(9):2073-83. Epub 2006/08/09. doi: 10.1084/jem.20060245. PubMed PMID: 16894012; PubMed Central PMCID: PMCPMC2118407.

10.   Bobryshev YV, Lord RS. Ultrastructural recognition of cells with dendritic cell morphology in human aortic intima. Contacting interactions of Vascular Dendritic Cells in athero-resistant and athero-prone areas of the normal aorta. Arch Histol Cytol. 1995;58(3):307-22. Epub 1995/08/01. doi: 10.1679/aohc.58.307. PubMed PMID: 8527238.

11.   Kawahara I, Kitagawa N, Tsutsumi K, Nagata I, Hayashi T, Koji T. The expression of vascular dendritic cells in human atherosclerotic carotid plaques. Hum Pathol. 2007;38(9):1378-85. Epub 2007/06/09. doi: 10.1016/j.humpath.2007.02.004. PubMed PMID: 17555794.

12.   Zernecke A, Erhard F, Weinberger T, Schulz C, Ley K, Saliba AE, et al. Integrated single-cell analysis based classification of vascular mononuclear phagocytes in mouse and human atherosclerosis. Cardiovasc Res. 2022. Epub 2022/10/04. doi: 10.1093/cvr/cvac161. PubMed PMID: 36190844.

13.   Eisenbarth SC. Dendritic cell subsets in T cell programming: location dictates function. Nat Rev Immunol. 2019;19(2):89-103. Epub 2018/11/23. doi: 10.1038/s41577-018-0088-1. PubMed PMID: 30464294; PubMed Central PMCID: PMCPMC7755085.

14.   Bayerl F, Meiser P, Donakonda S, Hirschberger A, Lacher SB, Pedde AM, et al. Tumor-derived prostaglandin E2 programs cDC1 dysfunction to impair intratumoral orchestration of anti-cancer T cell responses. Immunity. 2023;56(6):1341-58 e11. Epub 2023/06/15. doi: 10.1016/j.immuni.2023.05.011. PubMed PMID: 37315536.

15.   Legein B, Janssen EM, Theelen TL, Gijbels MJ, Walraven J, Klarquist JS, et al. Ablation of CD8alpha(+) dendritic cell mediated cross-presentation does not impact atherosclerosis in hyperlipidemic mice. Scientific reports. 2015;5:15414. Epub 20151021. doi: 10.1038/srep15414. PubMed PMID: 26486587; PubMed Central PMCID: PMCPMC4614009.

16.   Gil-Pulido J, Cochain C, Lippert MA, Schneider N, Butt E, Amezaga N, et al. Deletion of Batf3-dependent antigen-presenting cells does not affect atherosclerotic lesion formation in mice. PLoS One. 2017;12(8):e0181947. Epub 20170803. doi: 10.1371/journal.pone.0181947. PubMed PMID: 28771609; PubMed Central PMCID: PMCPMC5542449.

17.   Choi JH, Cheong C, Dandamudi DB, Park CG, Rodriguez A, Mehandru S, et al. Flt3 signaling-dependent dendritic cells protect against atherosclerosis. Immunity. 2011;35(5):819-31. Epub 2011/11/15. doi: 10.1016/j.immuni.2011.09.014. PubMed PMID: 22078798.

18.   Clement M, Haddad Y, Raffort J, Lareyre F, Newland SA, Master L, et al. Deletion of IRF8 (Interferon Regulatory Factor 8)-Dependent Dendritic Cells Abrogates Proatherogenic Adaptive Immunity. Circ Res. 2018;122(6):813-20. Epub 2018/02/14. doi: 10.1161/CIRCRESAHA.118.312713.

PubMed PMID: 29436389.

19. Haddad Y, Lahoute C, Clement M, Laurans L, Metghalchi S, Zeboudj L, et al. The Dendritic Cell Receptor DNGR-1 Promotes the Development of Atherosclerosis in Mice. Circ Res. 2017;121(3):234-43. Epub 2017/06/14. doi: 10.1161/CIRCRESAHA.117.310960. PubMed PMID: 28607102.

20. Wohn C, Le Guen V, Voluzan O, Fiore F, Henri S, Malissen B. Absence of MHC class II on cDC1 dendritic cells triggers fatal autoimmunity to a cross-presented self-antigen. Sci Immunol. 2020;5(45). Epub 2020/03/15. doi: 10.1126/sciimmunol.aba1896. PubMed PMID: 32169954.

21. Mundt S, Mrdjen D, Utz SG, Greter M, Schreiner B, Becher B. Conventional DCs sample and present myelin antigens in the healthy CNS and allow parenchymal T cell entry to initiate neuroinflammation. Sci Immunol. 2019;4(31). doi: 10.1126/sciimmunol.aau8380. PubMed PMID: 30679199.

22. Robbins CS, Hilgendorf I, Weber GF, Theurl I, Iwamoto Y, Figueiredo JL, et al. Local proliferation dominates lesional macrophage accumulation in atherosclerosis. Nat Med. 2013;19(9):1166-72. Epub 2013/08/13. doi: 10.1038/nm.3258. PubMed PMID: 23933982; PubMed Central PMCID: PMCPMC3769444.

23. Potteaux S, Gautier EL, Hutchison SB, van Rooijen N, Rader DJ, Thomas MJ, et al. Suppressed monocyte recruitment drives macrophage removal from atherosclerotic plaques of Apoe-/- mice during disease regression. J Clin Invest. 2011;121(5):2025-36. Epub 2011/04/21. doi: 10.1172/JCI43802. PubMed PMID: 21505265; PubMed Central PMCID: PMCPMC3083793.

24. Yilmaz A, Weber J, Cicha I, Stumpf C, Klein M, Raithel D, et al. Decrease in circulating myeloid dendritic cell precursors in coronary artery disease. J Am Coll Cardiol. 2006;48(1):70-80. Epub 2006/07/04. doi: 10.1016/j.jacc.2006.01.078. PubMed PMID: 16814651.

25. Alberts-Grill N, Denning TL, Rezvan A, Jo H. The role of the vascular dendritic cell network in atherosclerosis. Am J Physiol Cell Physiol. 2013;305(1):C1-21. Epub 2013/04/05. doi: 10.1152/ajpcell.00017.2013. PubMed PMID: 23552284; PubMed Central PMCID: PMCPMC3725520.

26. Yan Y, Thakur M, van der Vorst EPC, Weber C, Doring Y. Targeting the chemokine network in atherosclerosis. Atherosclerosis. 2021;330:95-106. Epub 2021/07/13. doi: 10.1016/j.atherosclerosis.2021.06.912. PubMed PMID: 34247863.

27. Eickhoff S, Brewitz A, Gerner MY, Klauschen F, Komander K, Hemmi H, et al. Robust Anti-viral Immunity Requires Multiple Distinct T Cell-Dendritic Cell Interactions. Cell. 2015;162(6):1322-37. Epub 2015/08/25. doi: 10.1016/j.cell.2015.08.004. PubMed PMID: 26296422; PubMed Central PMCID: PMCPMC4567961.

28. Weber C, Zernecke A, Libby P. The multifaceted contributions of leukocyte subsets to atherosclerosis: lessons from mouse models. Nat Rev Immunol. 2008;8(10):802-15. Epub 2008/10/01. doi: 10.1038/nri2415. PubMed PMID: 18825131.

29. Worbs T, Hammerschmidt SI, Forster R. Dendritic cell migration in health and disease. Nat Rev Immunol. 2017;17(1):30-48. Epub 2016/11/29. doi: 10.1038/nri.2016.116. PubMed PMID: 27890914.

30. Li T, Safitri M, Zhang K, Wang Y, Huang L, Zhu Y, et al. Downregulation of G3BP2 reduces atherosclerotic lesions in ApoE(-/-) mice. Atherosclerosis. 2020;310:64-74. Epub 2020/09/13. doi: 10.1016/j.atherosclerosis.2020.08.003. PubMed PMID: 32919187.

31. Mattiuz R, Wohn C, Ghilas S, Ambrosini M, Alexandre YO, Sanchez C, et al. Novel Cre-Expressing Mouse Strains Permitting to Selectively Track and Edit Type 1 Conventional Dendritic Cells Facilitate Disentangling Their Complexity in vivo. Front Immunol. 2018;9:2805. Epub 2018/12/20. doi: 10.3389/fimmu.2018.02805. PubMed PMID: 30564233; PubMed Central PMCID: PMCPMC6288293.

32. Luche H, Weber O, Nageswara Rao T, Blum C, Fehling HJ. Faithful activation of an extra-bright red fluorescent protein in "knock-in" Cre-reporter mice ideally suited for lineage tracing studies. European journal of immunology. 2007;37(1):43-53. Epub 2006/12/16. doi: 10.1002/eji.200636745. PubMed PMID: 17171761.

33. De Meyer GRY, Zurek M, Puylaert P, Martinet W. Programmed death of macrophages in atherosclerosis: mechanisms and therapeutic targets. Nat Rev Cardiol. 2024. Epub 2024/01/02. doi: 10.1038/s41569-023-00957-0. PubMed PMID: 38163815.

34. Yu EPK, Reinhold J, Yu H, Starks L, Uryga AK, Foote K, et al. Mitochondrial Respiration Is Reduced in Atherosclerosis, Promoting Necrotic Core Formation and Reducing Relative Fibrous Cap Thickness. Arterioscler Thromb Vasc Biol. 2017;37(12):2322-32. Epub 2017/10/04. doi: 10.1161/ATVBAHA.117.310042. PubMed PMID: 28970293; PubMed Central PMCID: PMCPMC5701734.

35. Piedrahita JA, Zhang SH, Hagaman JR, Oliver PM, Maeda N. Generation of mice carrying a mutant apolipoprotein E gene inactivated by gene targeting in embryonic stem cells. Proc Natl Acad Sci U S A. 1992;89(10):4471-5. Epub 1992/05/15. doi: 10.1073/pnas.89.10.4471. PubMed PMID: 1584779; PubMed Central PMCID: PMCPMC49104.

36. Plump AS, Smith JD, Hayek T, Aalto-Setala K, Walsh A, Verstuyft JG, et al. Severe hypercholesterolemia and atherosclerosis in apolipoprotein E-deficient mice created by homologous recombination in ES cells. Cell. 1992;71(2):343-53. Epub 1992/10/16. doi: 10.1016/0092-8674(92)90362-g. PubMed PMID: 1423598.

37. Fernandez DM, Rahman AH, Fernandez NF, Chudnovskiy A, Amir ED, Amadori L, et al. Single-cell immune landscape of human atherosclerotic plaques. Nat Med. 2019;25(10):1576-88. Epub 2019/10/09. doi: 10.1038/s41591-019-0590-4. PubMed PMID: 31591603; PubMed Central PMCID: PMCPMC7318784.

38. Afek A, Harats D, Roth A, Keren G, George J. Evidence for the involvement of T cell costimulation through the B-7/CD28 pathway in atherosclerotic plaques from apolipoprotein E knockout mice. Exp Mol Pathol. 2004;76(3):219-23. Epub 2004/05/06. doi: 10.1016/j.yexmp.2003.12.001. PubMed PMID: 15126104.

39. Cancel JC, Crozat K, Dalod M, Mattiuz R. Are Conventional Type 1 Dendritic Cells Critical for Protective Antitumor Immunity and How? Front Immunol. 2019;10:9. Epub 2019/02/28. doi: 10.3389/fimmu.2019.00009. PubMed PMID: 30809220; PubMed Central PMCID: PMCPMC6379659.

40. Li N, Steiger S, Fei L, Li C, Shi C, Salei N, et al. IRF8-Dependent Type I Conventional Dendritic Cells (cDC1s) Control Post-Ischemic Inflammation and Mildly Protect Against Post-Ischemic Acute Kidney Injury and Disease. Front Immunol. 2021;12:685559. Epub 2021/07/09. doi: 10.3389/fimmu.2021.685559. PubMed PMID: 34234783; PubMed Central PMCID: PMCPMC8255684.

41. Bosteels V, Marechal S, De Nolf C, Rennen S, Maelfait J, Tavernier SJ, et al. LXR signaling controls homeostatic dendritic cell maturation. Sci Immunol. 2023;8(83):eadd3955. Epub 2023/05/12. doi: 10.1126/sciimmunol.add3955. PubMed PMID: 37172103.

42. Bottcher JP, Bonavita E, Chakravarty P, Blees H, Cabeza-Cabrerizo M, Sammicheli S, et al. NK Cells Stimulate Recruitment of cDC1 into the Tumor Microenvironment Promoting Cancer Immune Control. Cell. 2018;172(5):1022-37 e14. doi: 10.1016/j.cell.2018.01.004. PubMed PMID: 29429633; PubMed Central PMCID: PMC5847168.

43. Dorner BG, Dorner MB, Zhou X, Opitz C, Mora A, Guttler S, et al. Selective expression of the chemokine receptor XCR1 on cross-presenting dendritic cells determines cooperation with CD8+ T cells.

Immunity. 2009;31(5):823-33. doi: 10.1016/j.immuni.2009.08.027. PubMed PMID: 19913446.

44. Xu F, He D, Ning R, Zeng B, Thompson CW, Li Y, et al. Genetic diversity of chemokine XCL1 and its receptor XCR1 in murine rodents. Developmental and comparative immunology. 2019;98:80-8. doi: 10.1016/j.dci.2019.04.008. PubMed PMID: 31026469.

45. Yoshida T, Izawa D, Nakayama T, Nakahara K, Kakizaki M, Imai T, et al. Molecular cloning of mXCR1, the murine SCM-1/lymphotactin receptor. FEBS letters. 1999;458(1):37-40. doi: 10.1016/s0014-5793(99)01114-x. PubMed PMID: 10518929.

46. Tall AR, Bornfeldt KE. Inflammasomes and Atherosclerosis: a Mixed Picture. Circ Res. 2023;132(11):1505-20. doi: 10.1161/CIRCRESAHA.123.321637. PubMed PMID: 37228237; PubMed Central PMCID: PMC10213995.

47. Vallejo J, Cochain C, Zernecke A, Ley K. Heterogeneity of immune cells in human atherosclerosis revealed by scRNA-Seq. Cardiovasc Res. 2021;117(13):2537-43. doi: 10.1093/cvr/cvab260. PubMed PMID: 34343272; PubMed Central PMCID: PMC8921647.

48. Guo X, Zhou Y, Wu T, Zhu X, Lai W, Wu L. Generation of mouse and human dendritic cells in vitro. J Immunol Methods. 2016;432:24-9. Epub 2016/02/16. doi: 10.1016/j.jim.2016.02.011. PubMed PMID: 26876301.

49. Anselmi G, Helft J, Guermonprez P. Development and function of human dendritic cells in humanized mice models. Mol Immunol. 2020;125:151-61. Epub 2020/07/21. doi: 10.1016/j.molimm.2020.07.005. PubMed PMID: 32688117.

50. Blanco P, Palucka AK, Pascual V, Banchereau J. Dendritic cells and cytokines in human inflammatory and autoimmune diseases. Cytokine Growth Factor Rev. 2008;19(1):41-52. Epub 2008/02/09. doi: 10.1016/j.cytogfr.2007.10.004. PubMed PMID: 18258476; PubMed Central PMCID: PMCPMC2413068.

51. Steinman RM. Dendritic cells: understanding immunogenicity. European journal of immunology. 2007;37 Suppl 1:S53-60. Epub 2007/11/01. doi: 10.1002/eji.200737400. PubMed PMID: 17972346.

52. Steinman RM, Adams JC, Cohn ZA. Identification of a novel cell type in peripheral lymphoid organs of mice. IV. Identification and distribution in mouse spleen. J Exp Med. 1975;141(4):804-20. Epub 1975/04/01. PubMed PMID: 1127378; PubMed Central PMCID: PMCPMC2189754.

53. Steinman RM, Kaplan G, Witmer MD, Cohn ZA. Identification of a novel cell type in peripheral lymphoid organs of mice. V. Purification of spleen dendritic cells, new surface markers, and maintenance in vitro. J Exp Med. 1979;149(1):1-16. Epub 1979/01/01. doi: 10.1084/jem.149.1.1. PubMed PMID: 762493; PubMed Central PMCID: PMCPMC2184752.

54. Clark GJ, Silveira PA, Hogarth PM, Hart DNJ. The cell surface phenotype of human dendritic cells. Semin Cell Dev Biol. 2019;86:3-14. Epub 2018/03/03. doi: 10.1016/j.semcdb.2018.02.013. PubMed PMID: 29499385.

55. Brewitz A, Eickhoff S, Dahling S, Quast T, Bedoui S, Kroczek RA, et al. CD8(+) T Cells Orchestrate pDC-XCR1(+) Dendritic Cell Spatial and Functional Cooperativity to Optimize Priming. Immunity. 2017;46(2):205-19. Epub 2017/02/14. doi: 10.1016/j.immuni.2017.01.003. PubMed PMID: 28190711; PubMed Central PMCID: PMCPMC5362251.

56. Alexandre YO, Ghilas S, Sanchez C, Le Bon A, Crozat K, Dalod M. XCR1+ dendritic cells promote memory CD8+ T cell recall upon secondary infections with Listeria monocytogenes or certain viruses. J Exp Med. 2016;213(1):75-92. Epub 2015/12/24. doi: 10.1084/jem.20142350. PubMed PMID: 26694969; PubMed Central PMCID: PMCPMC4710197.

57. Domenjo-Vila E, Casella V, Iwabuchi R, Fossum E, Pedragosa M, Castellvi Q, et al. XCR1+ DCs

are critical for T cell-mediated immunotherapy of chronic viral infections. Cell Rep. 2023;42(2):112123. Epub 2023/02/17. doi: 10.1016/j.celrep.2023.112123. PubMed PMID: 36795562.

58. Deczkowska A, David E, Ramadori P, Pfister D, Safran M, Li B, et al. XCR1(+) type 1 conventional dendritic cells drive liver pathology in non-alcoholic steatohepatitis. Nat Med. 2021;27(6):1043-54. Epub 2021/05/22. doi: 10.1038/s41591-021-01344-3. PubMed PMID: 34017133.

59. Mysore V, Tahir S, Furuhashi K, Arora J, Rosetti F, Cullere X, et al. Monocytes transition to macrophages within the inflamed vasculature via monocyte CCR2 and endothelial TNFR2. J Exp Med. 2022;219(5). Epub 2022/04/12. doi: 10.1084/jem.20210562. PubMed PMID: 35404389; PubMed Central PMCID: PMCPMC9006314.

60. Liu Z, Liao F, Zhu J, Zhou D, Heo GS, Leuhmann HP, et al. Reprogramming alveolar macrophage responses to TGF-beta reveals CCR2+ monocyte activity that promotes bronchiolitis obliterans syndrome. J Clin Invest. 2022;132(19). Epub 2022/10/04. doi: 10.1172/JCI159229. PubMed PMID: 36189800; PubMed Central PMCID: PMCPMC9525120.

61. Kim D, An L, Moon J, Maymi VI, McGurk AI, Rudd BD, et al. Ccr2+ Monocyte-Derived Macrophages Influence Trajectories of Acquired Therapy Resistance in Braf-Mutant Melanoma. Cancer Res. 2023;83(14):2328-44. Epub 2023/05/17. doi: 10.1158/0008-5472.CAN-22-2841. PubMed PMID: 37195124; PubMed Central PMCID: PMCPMC10478295.

62. Liu Z, Gu Y, Chakarov S, Bleriot C, Kwok I, Chen X, et al. Fate Mapping via Ms4a3-Expression History Traces Monocyte-Derived Cells. Cell. 2019;178(6):1509-25.e19. Epub 2019/09/07. doi: 10.1016/j.cell.2019.08.009. PubMed PMID: 31491389.

63. Bosmans LA, van Tiel CM, Aarts S, Willemsen L, Baardman J, van Os BW, et al. Myeloid CD40 deficiency reduces atherosclerosis by impairing macrophages' transition into a pro-inflammatory state. Cardiovasc Res. 2023;119(5):1146-60. Epub 2022/05/20. doi: 10.1093/cvr/cvac084. PubMed PMID: 35587037; PubMed Central PMCID: PMCPMC10202633.

64. Mildner A, Schonheit J, Giladi A, David E, Lara-Astiaso D, Lorenzo-Vivas E, et al. Genomic Characterization of Murine Monocytes Reveals C/EBPbeta Transcription Factor Dependence of Ly6C(-) Cells. Immunity. 2017;46(5):849-62 e7. Epub 2017/05/18. doi: 10.1016/j.immuni.2017.04.018. PubMed PMID: 28514690.

65. Inui H, Nishida M, Ichii M, Nakaoka H, Asaji M, Ide S, et al. XCR1(+) conventional dendritic cell-induced CD4(+) T helper 1 cell activation exacerbates cardiac remodeling after ischemic myocardial injury. J Mol Cell Cardiol. 2023;176:68-83. Epub 2023/02/06. doi: 10.1016/j.yjmcc.2023.01.011. PubMed PMID: 36739942.

66. Moore KJ, Tabas I. Macrophages in the pathogenesis of atherosclerosis. Cell. 2011;145(3):341-55. Epub 2011/05/03. doi: 10.1016/j.cell.2011.04.005. PubMed PMID: 21529710; PubMed Central PMCID: PMCPMC3111065.

67. Yoshida T, Imai T, Kakizaki M, Nishimura M, Takagi S, Yoshie O. Identification of single C motif-1/lymphotactin receptor XCR1. The Journal of biological chemistry. 1998;273(26):16551-4. doi: 10.1074/jbc.273.26.16551. PubMed PMID: 9632725.

68. de Oliveira CE, Gasparoto TH, Pinheiro CR, Amor NG, Nogueira MRS, Kaneno R, et al. CCR5-Dependent Homing of T Regulatory Cells to the Tumor Microenvironment Contributes to Skin Squamous Cell Carcinoma Development. Mol Cancer Ther. 2017;16(12):2871-80. Epub 2017/09/15. doi: 10.1158/1535-7163.MCT-17-0341. PubMed PMID: 28904130.

69. He F, Wu Z, Liu C, Zhu Y, Zhou Y, Tian E, et al. Targeting BCL9/BCL9L enhances antigen presentation by promoting conventional type 1 dendritic cell (cDC1) activation and tumor infiltration.

Signal Transduct Target Ther. 2024;9(1):139. Epub 2024/05/30. doi: 10.1038/s41392-024-01838-9. PubMed PMID: 38811552; PubMed Central PMCID: PMCPMC11137111.

70. Bottcher JP, Beyer M, Meissner F, Abdullah Z, Sander J, Hochst B, et al. Functional classification of memory CD8(+) T cells by CX3CR1 expression. Nat Commun. 2015;6:8306. Epub 2015/09/26. doi: 10.1038/ncomms9306. PubMed PMID: 26404698; PubMed Central PMCID: PMCPMC4667439.

71. Chu VT, Weber T, Graf R, Sommermann T, Petsch K, Sack U, et al. Efficient generation of Rosa26 knock-in mice using CRISPR/Cas9 in C57BL/6 zygotes. BMC Biotechnol. 2016;16:4. Epub 2016/01/17. doi: 10.1186/s12896-016-0234-4. PubMed PMID: 26772810; PubMed Central PMCID: PMCPMC4715285.

72. Larson MA. Pronuclear Microinjection of One-Cell Embryos. Methods in molecular biology. 2020;2066:27-33. doi: 10.1007/978-1-4939-9837-1_2. PubMed PMID: 31512204.

73. Voisinne G, Kersse K, Chaoui K, Lu L, Chaix J, Zhang L, et al. Quantitative interactomics in primary T cells unveils TCR signal diversification extent and dynamics. Nat Immunol. 2019;20(11):1530-41. Epub 2019/10/09. doi: 10.1038/s41590-019-0489-8. PubMed PMID: 31591574; PubMed Central PMCID: PMCPMC6859066.

74. Huang R, Guo G, Lu L, Fu R, Luo J, Liu Z, et al. The three members of the Vav family proteins form complexes that concur to foam cell formation and atherosclerosis. J Lipid Res. 2019;60(12):2006-19. Epub 2019/10/02. doi: 10.1194/jlr.M094771. PubMed PMID: 31570505; PubMed Central PMCID: PMCPMC6889716.

1013   Figure 1

1014
1015

1016   Figure 2

1017
1018

1019   Figure 3

1020
1021

1022   Figure 4

1023
1024

1025   Figure 5

1026
1027

1028   Figure 6

1029

1030

Figure 7