Xcr1[+] type 1 conventional dendritic cells are essential mediators for atherosclerosis progression

Running tittle: Targeting of cDC1 cells attenuates atherosclerosis

Tianhan Li[1#], Liaoxun Lu[2#], Juanjuan Qiu[2], Xin Dong[2], Le Yang[2], Kexin He[2], Yanrong Gu[2], Binhui Zhou[2], Tingting Jia[2], Toby Lawrence[3], Marie Malissen[3], Guixue Wang[4], Rong Huang[2], Hui Wang[2], Bernard Malissen[3*], Yinming Liang[2,5*] Lichen Zhang[2*]

[1]*School of Basic Medicine, Xinxiang Medical University, Xinxiang, China.*

[2]*School of Medical Technology, Xinxiang Medical University, Xinxiang, China.*

[3]*Centre d'Immunologie de Marseille-Luminy, INSERM, CNRS, Aix-Marseille Université, Marseille, France*

[4]*Key Laboratory for Biorheological Science and Technology of Ministry of Education, State and Local Joint Engineering Laboratory for Vascular Implants, Bioengineering College of Chongqing University, Chongqing, China*

[5]*Center of Disease Model and Immunology, Hunan Academy of Chinese Medicine, Changsha, China.*

[#]*These authors contributed equally to this work.*

[*]Correspondence: yinming.liang@gris.org.cn (Y L); zhanglichen@gris.org.cn (L Z).

bernardm@ciml.univ-mrs.fr (B M).

Abstract:

Background:

Atherosclerosis is characterized by lipid accumulation within plaques, leading to foam cell

formation and an inflammatory response within the aortic lesions. Lipid disorders have been

extensively investigated, however the cellular and molecular mechanisms that trigger the

inflammatory response in atherosclerotic plaques remain far from being fully understood. $Xcr1^+$

cDC1 cells are newly identified antigen-presenting cells in activating immune cells. However, the

role of cDC1 cells in the development of atherosclerosis remains highly controversial.

Methods and Results:

We first confirmed the presence of cDC1 within human atherosclerotic plaques and discovered a

significant association between the increasing cDC1 numbers and atherosclerosis progression in

mice. Subsequently, we established $Xcr1^{Cre-Gfp}$ $Rosa26^{LSL-DTA}$ $ApoE^{-/-}$ mice, a novel and complex

genetic model, in which cDC1 was selectively depleted in vivo during atherosclerosis

development. Intriguingly, we observed a notable reduction in atherosclerotic lesions in

hyperlipidemic mice, alongside suppressed T cell activation of both $CD4^+$ and $CD8^+$ subsets in the

aortic plaques. Notably, aortic macrophages and serum lipid levels were not significantly changed

in the cDC1-depleted mice. Single-cell RNA sequencing revealed heterogeneity of $Xcr1^+$ cDC1

cells across the aorta and lymphoid organs under hyperlipidemic conditions. As Xcr1 is the sole

receptor for Xcl1, we next explored to target $Xcr1^+$ cDC1 cells via Xcl1 by establishing Xcl1 and

ApoE deficient mice. Our data indicate that atherosclerotic lesions mediated by cDC1 are

dependent on Xcl1.

Conclusions:

Our results reveal crucial roles of cDC1 in atherosclerosis progression and provide insights into

the development of immunotherapies by targeting cDC1 through Xcl1.

Introduction

Atherosclerosis is a chronic inflammatory disease that affects the large and medium-sized arteries,

and it is the leading cause of cardiovascular mortality and morbidity worldwide [1].

Atherosclerosis is a complex pathological process during which lipid-laden phagocytes such as

macrophages, along with lymphocytes and dendritic cells, accumulate in the plaque [2]. Mounting

evidence from animal models and human studies demonstrates involvement of both the innate and

adaptive immunity in the development of atherosclerosis [3-5]. In healthy arteries, DCs are

typically present in the subendothelial space and work in the front line of immune surveillance [6,

7], and small portion of resident vascular DCs are localized in the adventitia. Various studies have

demonstrated that DCs accumulated in the advanced atherosclerotic lesions in both humans and

animal models [8-11]. Recently, several phenotypically and functionally distinct vascular DCs

have been identified in mouse and human atherosclerotic lesions [12]. Xcr1$^+$ type 1 conventional

dendritic cells (Xcr1$^+$cDC1) (also known as CD8$^+$ DC or CD103$^+$ DCs) has been recognized for

its role in cross-presentation and activating naive CD8$^+$ T cells, and are critical for anti-viral and

anti-tumor immune responses [13, 14]. Owing to the challenges associated with obtaining a model

for specific cDC1 depletion *in vivo*, previous studies investigating the role of cDC1 cells in

atherosclerosis using various knockout mice have produced conflicting results [15-19]. It is

important to note that previous Cre lines developed to achieve specific Cre expression in cDC1s

still led to leakiness in recombinase activity outside of cDC1s, as demonstrated in previous

publications [20, 21]. Moreover, scant mechanistic understanding regarding the regulation of

cDC1 activity during atherosclerosis has been reported. Therefore, developing a more specific

mouse model for cDC1 is essential for elucidating its precise role in atherosclerosis. Here, we

established a novel Cre line Xcr1$^{Cre-Gfp}$ mice and did not observe any leakiness of Cre activity, and

further experiments revealed that specific depletion cDC1 attenuates atherosclerosis development

by inhibiting both $CD4^+$ and $CD8^+$ T cell activation in hyperlipidemic Xcr1$^{Cre\text{-}Gfp}$ Rosa26$^{LSL\text{-}DTA}$

ApoE$^{-/-}$ mice.

The origin of phagocytes within the atherosclerotic plaques has been controversial. Studies have

shown that the major contribution of macrophages in the aortic plaque is recruited from the

circulation, while other studies have emphasized the significance of local proliferation [22, 23].

Tissue-resident dendritic cells inside the plaques have been well documented [24, 25], regardless

of the fact that the contribution and origin of Xcr1$^+$ cDC1 have not been appropriately addressed.

To understand the origin of Xcr1$^+$ cDC1 inside the plaque, we performed bone marrow

transplantation using Xcr1$^{Cre\text{-}Gfp}$ Rosa26$^{LSL\text{-}DTA}$ mice as donors and found that in ApoE$^{-/-}$ recipient

mice placed on a high-fat diet (HFD), cDC1 cells were completely absent in the aortas. Therefore,

our experiments have elucidated that cDC1 cells in the atherosclerotic plaques originate from the

bone marrow. Further single cell RNA sequencing illustrated that notable differences in Xcr1$^+$

cDC1 cells are discovered among the aorta, spleen, and lymph nodes. *Ccr2*, *Sept3* and *Cldnd1*

were highly expressed in Xcr1$^+$ cDC1 cells from aorta compared with that from spleen and lymph

nodes.

We further aimed to ascertain whether cDC1 can be modulated by targeting the chemokine Xcl1,

which is a ligand of Xcr1. Considering the vital role that Xcr1$^+$ cDC1 plays in the progression of

atherosclerosis and the therapeutic potential of manipulating their functions via chemokines[26],

comprehending the feasibility of targeting Xcl1 holds considerable promise for developing novel

therapeutic approaches to halt the advancement of atherosclerosis. We conducted further studies

using Xcl1$^{-/-}$ ApoE$^{-/-}$ double knockout mice to assess the pathological grade of atherosclerotic

lesions. Our data indicated that the knockout of Xcl1 led to a mild yet significant reduction of

cDC1 cells in the aorta. However, strikingly, when compared to the ApoE$^{-/-}$ control mice, the

Xcl1$^{-/-}$ ApoE$^{-/-}$ mice exhibited a markedly decreased severity of the atherosclerotic lesions, to an

extent comparable to that observed in cDC1-depleted Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice.

Therefore, our study which employed genetic models to deplete Xcr1^{+} cDC1 *in vivo* in

hyperlipidemic mice, has unequivocally determined the essential role of this specific cell type in

the development of atherosclerosis. Besides their well-established role in cross-presentation to

CD8^{+} T cells, recent studies have shown Xcr1^{+} cDC1 cells also engage with CD4^{+} T cells to

augment CD8^{+} T cell responses [20, 27]. Our data showing Xcr1^{+} cDC1-dependent CD4 and CD8

T cell activation in atherosclerotic plaques is in line with the new paradigm for cDC1s as a

platform for both CD4 and CD8 T cell responses in the novel context of atherosclerosis.

Most importantly, we have successfully demonstrated that targeting the chemokine Xcl1, the

ligand of Xcr1, can effectively inhibit atherosclerosis. These novel findings hold the potential to

forge a novel trajectory for therapeutic interventions within the field of atherosclerosis, presenting

promising prospects for better understanding and treatment of this disorder.

Results:

Progressive accumulation of Xcr1^{+} cDC1 cells in advanced atherosclerotic lesions in human

and mice

Dendritic cells are present in healthy arteries and accumulate within atherosclerotic lesions,

participating in various pathogenic and protective mechanisms during atherogenesis [28, 29].

However, the intra-plaque dynamics of cDC1 and their contribution to atherosclerosis remain

undetermined. Using the archived specimens of paraffin-embedded human atherosclerotic arteries

from our previous study [30], we examined the Xcr1$^+$ dendritic cells in human atherosclerotic

plaques. As shown in **Figure 1A and 1B**, Xcr1$^+$ dendritic cells were significantly more abundant

in the plaque area of the human femoral artery compared to that in the plaque-free area. Similarly,

the immunofluorescence staining results using specific antibodies for Xcr1 and CD11c revealed

that Xcr1$^+$CD11c$^+$ dendritic cells accumulated in the plaques of the aortic root in ApoE$^{-/-}$ mice

that were placed on a HFD for 16 weeks (**Figure 1C and 1D**). Sample preparations entailed

sectioning and appropriate fixation of the aortic root tissues, which were described in greater

detail in the Methods and Materials section. To further elucidate the dynamics of Xcr1$^+$ cDC1

cells within the plaque during the progression of atherosclerosis, we carried out

immunohistochemical analyses of these cells in the lesioned aortic root (**Figure 1E**). The area

positive for Xcr1 expression was quantified at 8-week on a chow diet, 8-week on HFD, 12-week

on HFD and 16-week on HFD, respectively. Quantitative analysis was conducted using image

processing software, revealing a continuous accumulation of Xcr1$^+$ cDC1 cells in the lesioned area

as the disease progressed (**Figure 1F**). Therefore, our experiments provided data indicating the

enrichment of Xcr1$^+$ cDC1 cells in aortic lesions in both humans and mice, and demonstrated that

the accumulation of Xcr1$^+$ cDC1 cells was significantly greater during the progression of

atherosclerosis.

Xcr1$^{\text{Cre-Gfp}}$ is selectively expressed in cDC1

To understand the role of Xcr1$^+$ cDC1 in regulating the development of atherosclerosis, we

established the Xcr1$^{\text{Cre-Gfp}}$ApoE$^{-/-}$ mice in which the Cre recombinase and eGFP was co-expressed

under the control of endogenous Xcr1 promoter following start codon (**Figure S1**). This new Cre

line differs from a previous design in which Cre recombinase was expressed via IRES linkage at

the 3' UTR and leaked expression of Cre was frequently observed [20, 31]. To verify the fidelity

of Cre expression under hyperlipidemic condition, we used hyperlipidemic ApoE$^{-/-}$ mice as

recipient mice and used bone marrow cells from Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-tdRFP}$ mice as donors for the

bone marrow transplantation (**Figure 2A**). In Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-tdRFP}$ mice, the presence of Cre

protein would turn on the expression of tdRFP as a reporter for the activity of Cre recombinase

[32]. Remarkably, we found that the expression of tdRFP was exclusively present in Xcr1^{+} cDC1

when mice were fed a HFD for 16 weeks in the bone marrow transplanted mice, and the Cre

recombinase activity was not detectable in non-cDC1 cells (**Figure 2B and 2C**).

To determine the specificity of GFP expression originating from Xcr1$^{Cre-Gfp}$ genetic modification,

we examined the GFP levels in three populations of DCs, namely pDC, cDC1, and cDC2 cells,

from the spleen of Xcr1$^{Cre-Gfp}$ ApoE$^{-/-}$ mice that were fed a HFD for 16 weeks. As expected, we

found that the GFP expression was stringently restricted to cDC1 cells (**Figure 2D and 2E**). In

parallel, we compared the GFP expression in Xcr1^{+} cDC1-depleted mice that were fed a HFD for

16 weeks. Such cDC1-depleted Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice were obtained by crossing

Xcr1$^{Cre-Gfp}$ ApoE$^{-/-}$ mice with Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice. Strikingly, we observed a complete

loss of GFP-positive cells and Xcr1^{+} cDC1 cells in Xcr1^{+} cDC1-depleted mice (**Figure 2D and**

2F). We also examined DCs, including pDCs, cDC1, and cDC2, in the lymph nodes and spleens of

Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ and ApoE$^{-/-}$ mice maintained on a chow diet for 7 weeks. The

flow cytometric analyses also confirmed the absence of Xcr1^{+} cDC1 population in Xcr1$^{Cre-Gfp}$

Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice (**Figure S2**). Collectively, these data suggest that Cre recombinase

activity was selectively confined to cDC1 cells, and no leakiness was observed in lineages other

than the $Xcr1^+$ cDC1 population. We also demonstrated that $Xcr1^{Cre-Gfp}$ $Rosa26^{LSL-DTA}$ $ApoE^{-/-}$

mice constitute a desirable model for exploring the role of cDC1 in atherosclerosis.

The specific depletion of $Xcr1^+$ cDC1 cells significantly alleviates atherosclerosis without

influencing the lipid profile and the number of intra-plaque macrophages

To investigate the role of cDC1 cells in atherosclerosis, both $Xcr1^{Cre-Gfp}$ $Rosa26^{LSL-DTA}$ $ApoE^{-/-}$ and

the control $ApoE^{-/-}$ mice were subjected to a HFD for 16 weeks. The ORO staining results

demonstrated that the size of lesion in both descending aorta and aortic root was significantly

smaller in $Xcr1^{Cre-Gfp}$ $Rosa26^{LSL-DTA}$ $ApoE^{-/-}$ mice compared with $ApoE^{-/-}$ control mice (12.82% *vs*

6.01% in descending aorta, $p < 0.001$; 39.25% *vs* 33.57% in aortic root, $p < 0.01$) (**Figure 3A-D**).

Given that macrophages play a pivotal role in the development and progression of atherosclerotic

lesions, particularly within the necrotic core [33, 34], we next investigated macrophage

involvement in the aorta using IHC and flow cytometry. Interestingly, no significant differences

were observed in the necrotic core area of the aortic root between $Xcr1^{Cre-Gfp}$ $Rosa26^{LSL-DTA}$ $ApoE^-$

$^{/-}$ mice and $ApoE^{-/-}$ control mice (**Figure 3E and 3F**). Similarly, subsequent flow cytometry

analyses and IHC results revealed comparable proportions of $F4/80^+$ $CD11b^+$ macrophages in the

aorta, as well as similar percentages of CD68-positive macrophages in the aortic lesions between

the two groups of mice (**Figure 3G-I**). Furthermore, since dyslipidemia is a key characteristic of

HFD-treated $ApoE^{-/-}$ mice [35, 36], it is critical to assess body weight and lipid profiles in both

groups. Our results indicated no significant differences in body weight or levels of total

cholesterol (TC), triglycerides (TG), low-density lipoprotein (LDL), and high-density lipoprotein

(HDL) between the two groups (**Figure S3A-B**). Additionally, ORO staining of the liver revealed

no significant differences either (**Figure S3C-D**). Taken together, these results suggested that

specific depletion of cDC1 attenuated atherogenesis without affecting lipid status or macrophagic

accumulation in aortic lesions.

Specific depletion of cDC1 attenuates the development of atherosclerosis by modulating the

activation of T cell in ApoE$^{-/-}$ mice

T cells are among the critical drivers of the pathogenesis of atherosclerosis, and therefore, the

triggers of T cell activation in aortic plaques are of great interest for controlling disease

progression [2, 37, 38]. We initially analyzed the cDC1 cells through flow cytometry in both

Xcr1$^+$ cDC1 depleted mice and control mice. In the Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice that

were fed a HFD for 16 weeks, we observed a total absence of Xcr1$^+$ cDC1 in the aorta, lymph

nodes, and spleen (**Figure 4A-B and Figure S4**). Subsequently, we aimed to explore the T cell

phenotype in the two groups of mice, and found that the activation of aortic CD4$^+$ and CD8$^+$ T

cells were both significantly inhibited. The expression of CD69 in aortic CD4$^+$ and CD8$^+$ T cells

was significantly lower in the Xcr1$^+$ cDC1 depleted mice after being placed on a HFD for 16

weeks (**Figure 4C-D**). Interestingly, such decreased frequencies of CD69$^+$ T cells were only

observed in aortic lesions but not in lymphoid organs such as lymph nodes and the spleen in the

Xcr1$^+$ cDC1 depleted mice (**Figure S5A-B and S5D-E**). We further analyzed the absolute

numbers of CD8$^+$ and CD4$^+$ T cells in the spleen and lymph nodes and found that the absolute

numbers of CD8$^+$ and CD4$^+$ T cells were markedly reduced in the spleen but not in the lymph

nodes in Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice compared to those in ApoE$^{-/-}$ mice (**Figure S5C**

and S5F). Absence of cDC1 also resulted in reduced frequencies of CD8$^+$ T cells in both spleen

and lymph nodes in hyperlipidemic mice. We also analyzed T cell populations in the lymph nodes

and spleen in two groups of mice that were fed a chow diet and found that the frequencies of

CD8$^+$ T cells were significantly decreased in the lymph nodes of Xcr1$^+$ cDC1 depleted mice

(**Figure S6A-B**). More importantly, the frequencies of CD44^{high} CD62L^{low} activated memory cells

within the CD8$^+$ subsets in both the spleen and lymph nodes were significantly lower than those in

the ApoE$^{-/-}$ control mice (**Figure S6A-D**). However, in the CD4$^+$ subsets, such frequencies of

CD44^{high} CD62L^{low} activated cells were comparable in the lymph nodes but marginally lower in

the spleen (**Figure S6A-D**). It is important to note that the decreased absolute number of CD8$^+$ T

cells in the spleen in Xcr1$^+$ cDC1 depleted mice was only observed in HFD groups but not in

chow diet groups (**Figure S5F and Figure S6F**). Our data indicate that in aortic plaques, Xcr1$^+$

cDC1 cells are implicated in activating CD4$^+$ and CD8$^+$ T cells, and the loss of the cDC1

population led to a decreased absolute count of CD8$^+$ T cells in the lymph nodes of HFD treated

mice. Whereas in mice treated with a chow diet, the loss of the cDC1 population resulted in

reduced frequencies of activated memory T cells but did not affect the absolute counts of T cells.

Attenuated T cell activation might account for the mitigated atherosclerosis in Xcr1$^+$ cDC1

depleted mice.

Characterization of aortic Xcr1$^+$ cDC1 cells in atherosclerotic lesions

Although the majority of cDC1 cells derive from bone marrow, other mechanisms such as local

proliferation of existing DCs can also contribute to the cDC1 population in inflammatory diseases

[39, 40]. It remains unclear whether cDC1 cells in lymphoid organs such as spleen and lymph

nodes represent a different status compared to cDC1 cells in atherosclerotic lesions. To elucidate

the source and status of Xcr1$^+$ cDC1 in atherosclerotic lesions, we performed bone marrow

transplantation and single-cell RNA sequencing (sc-RNA seq) of cDC1 cells from aortic plaques

and lymphoid organs. We conducted bone marrow transplantation using Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$

or WT mice that carry the CD45.2 congenic markers as donors, and CD45.1$^+$ CD45.2$^+$ ApoE$^{-/-}$

mice as recipients (**Figure S7A**). Our data showed that body weight, lipid profile, hepatic lipid

accumulation were comparable between two groups of bone marrow transplanted mice after being

fed with a HFD for 16 weeks (**Figure S7B-D**). Interestingly, flow cytometric analysis of the aorta

displayed almost complete depletion of cDC1 cells in ApoE$^{-/-}$ recipient mice transplanted with

Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ bone marrow mice (**Figure S7E**). Strikingly we did not find observe

significant changes in pDCs and CD64$^+$F4/80$^+$ aortic macrophages (**Figure S7E-F**). Therefore, our

bone marrow transplantation experiments further confirmed fidelity of the Cre recombinase

expression in cDC1 cells in mice.

Next, we carried out sc-RNA seq to characterize Xcr1$^+$ cDC1 cells from multiple organs of

hyperlipidemic mice. Due to the scarcity of Xcr1$^+$ cDC1 cells that can be sorted by FACS and

processed for sc-RNA seq, we pooled FACS sorted Xcr1$^+$ cDC1 cells from three organs of

hyperlipidemic ApoE-deficient mice fed a HFD for 20 weeks, including the aorta, spleen, and

lymph nodes, and Xcr1$^+$ cDC1 cells from aorta and splenocytes of ApoE-deficient mice fed a

chow diet. Libraries for sc-RNA seq were prepared using barcode oligos conjugated to anti-CD45

antibodies for surface staining so that samples from each type of organ could be compared in

sequential analyses. Therefore, we had a total of five samples, and our focus was to understand the

differences between Xcr1$^+$ cDC1 cells from three different organs under hyperlipidemic

conditions. The cells were sorted by gating Xcr1-positive cells out of the MHCII$^+$CD11c$^+$

conventional dendritic cells (**Figure S8**). Firstly, we obtained the UMAP plot of 10 clusters of

cDC1 cells using the Seurat 4.0 package (**Figure 5A**). In this analysis, 3830 quality-controlled

Xcr1$^+$ cDC1 cells were from three types of organs in hyperlipidemic mice, and an additional 2081

cells were from cDC1 cells sorted from the spleen and aorta of ApoE deficient mice fed on a chow

diet. The clusters were differentiated according to top 10 differentially expressed genes

(**Supplementary Table 2**), and the heatmap shows marker genes and heterogeneity among the 10

clusters of cDC1 cells (**Figure 5B**). The purpose of this experiment was to determine whether

cDC1 cells from different organs possess unique features and whether cDC1 cells from aortic

lesions can be characterized by the expression of marker genes. In hyperlipidemic mice, cDC1

cells from the spleen and lymph nodes differed quite dramatically in Cluster 2 (*Stk17b* and

Gpr171 highly expressing cells), where 28.9% of cDC1 cells from the spleen versus 1.0% of

cDC1 cells from the lymph nodes were located. Conversely, 3.15% of cDC1 cells from the spleen

versus 32.4% of cDC1 cells from the lymph nodes fell in Cluster 3 (*Ccl5* and *Epsti1* highly

expressing cells), and 4.6% of cDC1 cells from the spleen versus 25.9% of cDC1 cells from the

lymph nodes fell in Cluster 4 (*Slfn5* and *Ifi44* highly expressing cells). Despite the fact that we had

only transcriptomic data for 68 Xcr1$^+$ cDC1 cells from aortic lesions, we were surprised to find

that such aortic cDC1 mainly fell into Cluster 1, 4 and 5, comprising 80.8% of the cells analyzed

(**Figure 5C**). In Cluster 1 (*Hspa1b* and *Dnajb1* highly expressing cells), where aortic cDC1 is

dominantly enriched, the cDC1 cells sorted from lymph nodes were essentially absent, while in

Cluster 4, aortic cDC1 cells were highly enriched and splenic cDC1 was obviously lower in

percentage (**Figure 5C**). Cluster 7 (*Fabp5* and *S100a4* highly expressing cells) is exclusively

comprised of cDC1 cells sourced from lymph nodes, with a striking absence of cDC1 cells from

the spleen and aorta (**Figure 5C**). To better visualize the enrichment of aortic Xcr1$^+$ cDC1 cells

distributed in a few clusters, we separately compared the three samples from the spleen, lymph

nodes, and aorta of hyperlipidemic mice using density plots. Interestingly, the Xcr1$^+$ cDC1 cells

from the spleen and lymph nodes displayed an obvious difference in their distribution, and aortic

Xcr1$^+$ cDC1 cells appeared more enriched in a few clusters (**Figure 5D-E**). Therefore, our data

suggest that Xcr1$^+$ cDC1 cells from the spleen and lymph nodes can differ markedly in

heterogeneity, and aortic Xcr1$^+$ cDC1 cells are enriched in a few clusters that express marker

genes, *Ccr2*, *Sept3* and *Cldnd1* (**Supplementary Table 3**).

Together, these data implied that Xcr1$^+$ cDC1 in the aorta originate from bone marrow, and Xcr1$^+$

cDC1 cells exhibit strong heterogeneity among aorta, spleen and lymph nodes.

Knockout of Xcl1 attenuates the development of atherosclerosis in ApoE$^{-/-}$ mice

Xcl1 is the major chemokine involved in cDC1 recruitment [41, 42], and plays critical role in

immune response with diverse physical and pathological implication through its interaction with

the sole receptor, Xcr1 [43, 44]. We found the presence of Xcl1 in the lesion area in both human

and mouse atherosclerotic lesions (data not shown). To elucidate the causal relationship between

Xcl1 and Xcr1$^+$ cDC1 in the development of atherosclerosis, and more importantly to

experimentally verify Xcl1 as a potential therapeutic target for atherosclerosis treatment, we

established the Xcl1 and ApoE double knockout mice. After being fed with a HFD for 19 weeks,

both Xcl1$^{-/-}$ApoE$^{-/-}$ and ApoE$^{-/-}$ mice were sacrificed to assess the severity of atherosclerosis. As

shown in **Figure 6A-D**, the size of lesion at the descending aorta and aortic root was significantly

smaller in Xcl1$^{-/-}$ApoE$^{-/-}$ compared to that in ApoE$^{-/-}$ mice. Importantly, ORO staining of the

liver, body weight and blood lipid profiles revealed no significant differences between the two

groups (**Figure 6E-H**). Moreover, the IHC experiment result show the percentage of CD68

positive macrophages in aortic lesion was similar between two groups (**Figure 6I-J**).

Previous study identified that Xcl1 serves as a potent chemokine, selectively acting on cDC1 cells

through its exclusive receptor, Xcr1 [42]. Consequently, we examined the DCs and the activation

of T cells in different organs, including aorta, lymph node and spleen. The flow cytometric results

illustrated that the frequencies of $Xcr1^+$ cDC1 cells in the aorta were significantly reduced, but the

frequencies of pDC and cDC2 cells from $Xcl1^{-/-} ApoE^{-/-}$ were comparable with that from $ApoE^{-/-}$

(**Figure 7A-B**). Moreover, in both lymph node and spleen, the absolute numbers of pDC, cDC1

and cDC2 from $Xcl1^{-/-} ApoE^{-/-}$ were comparable with that from $ApoE^{-/-}$ (**Figure S9**). The

frequencies of $CD8b^+$ T cells were also significantly lower, and $CD4^+$ T cells were significantly

higher, but the $CD4^+CD69^+$ and $CD8^+CD69^+$ T cells in the aorta from $Xcl1^{-/-} ApoE^{-/-}$ mouse were

comparable with that from $ApoE^{-/-}$ mouse (**Figure 7C-D**). Furthermore, the absolute numbers of

$CD4^+$, $CD8^+$, $CD4^+CD69^+$ and $CD8^+CD69^+$ T cells in both lymph node and spleen from $Xcl1^{-/-}$

$ApoE^{-/-}$ mouse were comparable with that from $ApoE^{-/-}$ mouse (**Figure S10**).

Collectively, these data suggested that Xcl1 could attenuate the development of atherosclerosis via

reducing the recruitment of $Xcr1^+$ cDC1 cells and $CD8^+$ T cells in atherosclerotic lesions. More

importantly, *in vivo* data demonstrated the promising potential of targeting Xcl1 to reduce the

inflammatory response in atherosclerotic lesions. Our work paves the way to discover novel

solutions for the treatment of atherosclerosis by targeting Xcl1 and $Xcr1^+$ cDC1 cells.

Discussion

Atherosclerosis is a complex disorder, and understanding the role of the immune system,

particularly dendritic cells (DCs), is of paramount importance [45, 46]. The marked diversity of

DC subsets and their distinct, and at times contradictory functions pose challenges in clarifying

the role of each specific type of DC in the context of atherosclerosis [47-53]. Moreover, the

restricted characterization of each subset of DCs in atherosclerosis, especially when compared to

that in well-studied organs such as the skin and lymph nodes [13], has been severely hindered by

the limited application of appropriate genetic models and *in vivo* data. Equally significant is that

dissecting the functions of each subset of DCs *in vivo* relies on the development of genetic tools

validated for lineage specificity. Recently, Xcr1$^+$ cDC1 has been extensively investigated using the

Cre line under the promoter of the Xcr1, despite well-documented expression leakiness [20, 31].

Compelling evidence from tumor and viral infection models [14, 41, 54-56], as well as metabolic

disorders like non-alcoholic steatohepatitis [57], has demonstrated the critical role of Xcr1$^+$cDC1

cells in diseases. Nevertheless, the precise role of Xcr1$^+$ cDC1 cells in atherosclerosis remains

undetermined.

The precise characterization of Xcr1$^+$ cDC1 and its functions within the atherosclerotic milieu

could provide valuable insights into the immunopathogenesis of the disease. By delineating the

specific contributions of Xcr1$^+$ cDC1, we may identify novel therapeutic targets and develop more

targeted and effective strategies for the prevention and treatment of atherosclerosis. Therefore,

investigation to understand the roles of Xcr1$^+$ cDC1 in the context of atherosclerosis and to

identify therapeutic targets is highly warranted. However, at least two major obstacles impede the

in vivo study of Xcr1$^+$ cDC1 in atherosclerotic mice. First, Xcr1$^+$ cDC1 cells are extremely scarce

in number, and it is quite challenging to obtain highly purified Xcr1$^+$ cDC1 cells from aortic

plaque in hyperlipidemic mice. Second, there are very few appropriate genetic model systems that

permit functional analyses of Xcr1$^+$ cDC1 *in vivo*. Earlier studies on the role of Xcr1$^+$ cDC1 cells

in atherosclerosis have yielded conflicting results, largely due to the use of different mouse models.

For instance, the deletion of Batf3, Irf8, Dngr1, or Flt3 decreased the cDC1 cells. Nevertheless,

Batf3 deficiency had no impact on atherosclerosis in hyperlipidemic mice. In contrast, Irf8 or

Dngr1 deficiency reduced the development of atherosclerosis, while Flt3 deficiency exacerbated

the condition [15-19]. The recently developed Xcr1-Cre was found to have leakiness of

recombinase activity in other immune cells [20]. Here, we optimized the design for the

construction of the Cre line using the endogenous promoter of Xcr1 and validated its fidelity

under hyperlipidemic conditions. We crossed the novel knock-in line termed Xcr1[Cre-Gfp] with

Rosa26[LSL-tdRFP] Cre reporter mice and used the resulting F1 hybrids as bone marrow donors. The

bone marrow cells were then transferred into ApoE$^{-/-}$ mice, which were subsequently fed on a

HFD. We monitored the Cre recombinase activity, which was reflected by the tdRFP expression of

this novel mouse line. Under hyperlipidemic conditions, we did not observe any leakiness of Cre

activity, as tdRFP was strictly restricted to cDC1 cells. To the best of our knowledge, this novel

Cre line is the first of its kind. We further crossed the Xcr1[Cre-Gfp] and Rosa26[LSL-DTA] mice with

ApoE$^{-/-}$ mice respectively. Subsequently, by further crossing, we obtained Xcr1[Cre-Gfp]

Rosa26[LSL-DTA] ApoE$^{-/-}$ mice to create a complex genetic model, in which Xcr1^{+} cDC1 is depleted

in vivo in the context of atherosclerosis. Very interestingly, in the Xcr1^{+} cDC1 depleted mice,

atherosclerosis was significantly ameliorated. Coupled with the data we obtained from murine and

human samples, where Xcr1^{+} cDC1 cells exist in the aortic plaque and their accumulation is

correlated with disease progression, we conclude that Xcr1^{+} cDC1 is essential for the development

of atherosclerosis.

Despite the fact that Xcr1^{+} cDC1 cells are extremely rare in number, we were inclined to

understand if there were any differences among Xcr1^{+} cDC1 cells from different organs. We

compared, under hyperlipidemic conditions, the Xcr1^{+} cDC1 cells sorted from the spleen and

lymph nodes, supplemented by a small number of but well-purified Xcr1[+]cDC1 cells from aortic

lesions. To our surprise, the clusters of Xcr1[+] cDC1 displayed obvious distinctions among

different organs. When the Xcr1[+] cDC1 cells from aortic lesions were projected to the clusters, we

found that the major subsets present in the lymph nodes, characterized by the expression of

markers *Ccl5 and Fabp5*, and in the spleen, marked by *Stk17b* and *Gpr171*, were basically absent

in the aorta. Meanwhile, compared with spleen and lymph nodes, the aortic Xcr1[+] cDC1 cells are

highly expressed Ccr2, which is an extremely important chemokine receptor dominating monocyte

homing [58-60]. Using the bone marrow transfer model, we also determined that the Xcr1[+] cDC1

in the aortic plaque is replenished by circulating precursors.

In other disease models, Xcr1[+] cDC1 cells are crucial for activating T cells [55, 61], and we

confirmed that in hyperlipidemic mice, the depletion of cDC1 led to a significant inhibition of T

cell activation. It is important to note that Xcr1[+] cDC1 cells are known for well-established role in

cross-presentation to CD8[+] T cells, recent studies have shown Xcr1[+] cDC1 cells also engage with

CD4[+] T cells [20, 27]. Our data showing Xcr1[+] cDC1-dependent CD4 and CD8 T cell activation

in atherosclerotic plaques is in line with the new paradigm for cDC1s as a platform for both CD4

and CD8 T cell responses in the novel context of atherosclerosis. Considering the multifaceted

roles of macrophages in lipid metabolism, inflammation, and plaque dynamics during the

development of atherosclerosis [62], we evaluated the presence of macrophages in atherosclerotic

lesions using IHC and flow cytometry. Our results indicated comparable proportions of

macrophages in the aorta between Xcr1[+] cDC1 depleted mice and ApoE[−/−] control mice.

Additionally, our results also demonstrated that the loss of Xcr1[+] cDC1 cells did not affect lipid

status.

Since we elucidated the critical role of $Xcr1^+$ cDC1 cells in atherosclerosis, our aim was to test the

therapeutic potential of inhibiting this specific cell type during disease progression. As Xcr1 is the

sole receptor for Xcl1, and Xcl1 selectively attracts $Xcr1^+$ cDC1 cells in both mice and humans

[44, 63], we performed Xcl1 knockout on the background of ApoE deficient mice and assessed the

impact on atherosclerosis development. We found that the Xcl1-Xcr1 axis plays a crucial role in

the progression of atherosclerosis. Our results demonstrated that the loss of Xcl1 in

hyperlipidemic $ApoE^{-/-}$ mice significantly reduced atherosclerotic lesion formation, decreased the

frequencies of $Xcr1^+$ cDC1 cells and $CD8^+$ T cells in the aorta, without affecting lipid status or the

number of aortic macrophages. Our findings indicate that Xcl1 could potentially serve as an

effective target for the treatment of atherosclerosis.

Materials and Methods

Animal experiments and patients

All mice were raised in a Specific Pathogen-Free facility and the experiments were conducted in

accordance with the guidelines approved by the Animal Ethics Committee of Xinxiang Medical

University (China) (Reference No. XYLL-2016S001). The sections of the human femoral artery

were obtained in accordance with our previous study [30] and approved by the Ethics Committee

of the Third Military Medical University (Project identification code SYXK-PLA-20170005).

$Rosa26^{LSL-RFP}$ knock-in mice were generously provided by Hervé Luche at University Clinics Ulm

in Germany [32]. C57BL/6 and $ApoE^{-/-}$ mice were purchased from Beijing Vital River Laboratory

Animal Technology (China). $Rosa26^{LSL-DTA}$ knock-in mice (Stock Number 009669) were

purchased from the Jackson Laboratory (United States of America). Both HFD (contained 20%

protein, 20% carbohydrate, 40% fat and 1.25% cholesterol, Cat: XT108C), and chow diet

(contained 20% protein, 64% carbohydrate, 20% fat, Cat: AIN-93G) were purchased from Jiangsu

Xietong Pharmaceutical Bioengineering Co., Ltd. 7-8 weeks old mice were used to establish

atherosclerosis model through fed on a HFD for 8-20 weeks. On harvest, each sample was

assigned an analytical code that was irrelevant to its genotype and processed and analyzed by a

researcher blinded to its origin.

Generation of Xcr1$^{Cre-Gfp}$ knock-in mice

The Xcr1$^{Cre-Gfp}$ mouse line was generated by inserting a Cre-P2A-GFP-P2A cassette immediately

upstream of the endogenous *Xcr1* translational start codon (ATG). This design enables

co-expression of Cre recombinase and GFP under the control of the native *Xcr1* promoter. A

3.2-kb repair template was synthesized (GenScript, Nanjing, China), consisting of a 0.62-kb 5'

homology arm (HA), a codon-optimized *Cre* sequence fused to a P2A peptide, an enhanced GFP

reporter linked to a second P2A site, and a 0.67-kb 3' homology arm. For targeted integration,

CRISPR/Cas9-mediated homology-directed repair (HDR) was performed by co-injecting Cas9

mRNA, a chimeric sgRNA (sgXcr1-KI), and the repair template into fertilized C57BL/6 zygotes

at the one-cell stage, following established protocols[64, 65]. Founders (F0) harboring the correct

knock-in allele were identified by PCR and Sanger sequencing, then used to establish

heterozygous lines. The sgRNA sequence is provided in **Supplementary Table 1**.

Breeding strategy of Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$

To generate Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, we performed the following crosses:

Xcr1$^{Cre-Gfp}$ mice were crossed with ApoE$^{-/-}$ mice, and offspring heterozygous for both alleles were

intercrossed to obtain homozygous $Xcr1^{Cre-Gfp} ApoE^{-/-}$ mice. $Rosa26^{LSL-DTA}$ mice were crossed

with $ApoE^{-/-}$ mice, and progeny heterozygous for both alleles were intercrossed to generate

homozygous $Rosa26^{LSL-DTA} ApoE^{-/-}$ mice. Finally, homozygous $Xcr1^{Cre-Gfp} ApoE^{-/-}$ mice were

crossed with homozygous $Rosa26^{LSL-DTA} ApoE^{-/-}$ mice to produce $Xcr1^{Cre-Gfp} Rosa26^{LSL-DTA}$

$ApoE^{-/-}$ offspring.

Generation of Xcl1 knockout mice

The Xcl1 knockout mouse line was generated using CRISPR/Cas9 genome editing through the

microinjection of a mixture containing Cas9 mRNA and two sgRNAs into fertilized C57BL/6

zygotes at the one-cell stage, as previously described [66]. The genotypes were confirmed using

PCR and Sanger sequencing. Subsequently, Xcl1 and ApoE double knockout mice were obtained

by crossing the $Xcl1^{-/-}$ mice with $ApoE^{-/-}$ mice. The sequences of the sgRNA are reported in

Supplementary Table 1.

Bone marrow transplantation

After one week of antibiotic treatment, the recipient $CD45.1^+ CD45.2^+ ApoE^{-/-}$ mice were injected

intraperitoneally with busulfan at a dosage of 30 μg/g every two days for one week. Subsequently,

bone marrow from 8-week-old $CD45.2^+ Xcr1^{Cre-Gfp} Rosa26^{LSL-RFP}$, $CD45.2^+ Xcr1^{Cre-Gfp}$

$Rosa26^{LSL-DTA}$ or WT ($CD45.2^+$ wild-type) donor mice was intravenously transferred to each

recipient mouse, with two million cells administered per recipient. The efficiency of

transplantation was assessed four-week post-transplantation, after which all mice were fed a HFD

for 16 weeks. Finally, the mice were sacrificed for *en face* analysis of the descending aorta,

hematoxylin and eosin (H&E) staining, Oil Red O (ORO) staining of the aortic root, and flow

cytometry analysis, as previously described [67].

Immunofluorescence

Frozen sections of the aortic root from mice were incubated at 60 °C for 2 minutes and then fixed

in 4% paraformaldehyde for 30 minutes. The samples were subsequently washed with PBS and

permeabilized/blocked with 0.1% Triton X-100 in 5% BSA. Next, the primary antibodies,

including anti-Xcr1 (BD Biosciences, Cat. 148212) and anti-CD11c (BD Biosciences, Cat.

12-0114-83), were applied in a wet box at 4 °C overnight. After washing with PBST (PBS

containing 0.1% Tween-20) for five times, the samples were incubated with secondary antibodies

for 1 hour, followed by a 10-minute incubation with DAPI at room temperature (protected from

light). The fluorescent signals of the sections were detected using a Nikon C2 confocal

microscope (Nikon, Japan).

Morphometric and immunohistochemical analyses

After modeling, the heart and aorta tissue were obtained from mice. The *en face* descending aortas

were fixed with 4% paraformaldehyde and subsequently stained with ORO. Additionally, frozen

sections of the aortic roots were stained with ORO and H&E to assess the severity of

atherosclerosis, and frozen sections of livers were stained with ORO to assess the severity of lipid

accumulation, in accordance with our previous study [67]. Paraffin-embedded sections of the

aortic root were dewaxed in xylene twice for 10 minutes, rehydrated through descending grades of

ethanol, and subjected to sodium citrate-induced antigen retrieval for 10 minutes at 95 °C. The

sections were then incubated with 3% H_2O_2 for 10 minutes at room temperature. Following this,

the sections were blocked and incubated with primary antibodies, either anti-Xcr1 (NBP1-02343)

or CD68 (FB113109). After five washes with DPBS, the sections were incubated with secondary

antibodies for 30 minutes and stained with DAB at room temperature. Subsequently, the sections

were counterstained with hematoxylin, dehydrated in ethanol, and mounted with neutral balsam.

Images were captured using the digital pathology system (Pannoramic MIDI, Hungary), viewed

using CaseViewer2.4 software and analyzed using ImageJ software.

Flow cytometry analysis

To obtain the single-cell suspension, aortas, lymph nodes and spleens were collected from each

mouse. Spleens were directly dissociated using the Gentle MACS Dissociators (Miltenyi Biotec).

Aortas and lymph nodes were cut into small pieces and then dissociated using the Gentle MACS

Dissociators. Dissociated tissues digested with digestive enzymes (Collagenase IV, Cat: V900893

from Sigma; DNase I, Cat: DN25 from Sigma; Liberase DH, Cat: 5401054001 from Roche) for 30

minutes at 37 °C using a shaker set to 300 rpm or a heating block. The resulting cell suspension

was filtered through 75 μm strainers and subsequently blocked with 2.4G2 for 20 minutes at 4 °C,

followed by washing with stain buffer (BD, Cat: 554656). Then samples were stained with

monoclonal antibody mixes for 30 minutes at 4 °C followed by washing with stain buffer. Finally,

cells were resuspended in stain buffer containing 200 nM Sytox blue (ThermoFisher Scientific,

Cat: S34857) and acquired using a flow cytometer (ThermoFisher Scientific, Invitrogen Attune

NxT Flow Cytometer), and the FACS data were analyzed using FlowJo software. The following

antibodies purchased from BioLegend, BD or eBioscience were used to detect surface markers by

flow cytometry: anti-CD45.2 (104) conjugated to eFluor 450, FITC or PE-Cy7; anti-CD45.1 (A20)

conjugated to PE or eFluor450; anti-CD45 (30-F11) conjugated to PE; anti-CD11b (M1/70)

conjugated to super Bright 600, BUV395 or biotin; anti-CD3e (145-2C11) conjugated to

PE-Cy5.5 or PE-Cy7; anti-CD3e (eBio500A2) conjugated to Alexa Fluor700; anti-CD19

(eBio1D3) conjugated to PE-Cy5.5 or biotin; anti-Xcr1 (ZET) conjugated to APC or BV650;

anti-Ly6G (1A8) conjugated to Alexa Fluor 700; anti-MHCII (M5/114.15.2) conjugated to

APC-eFluor780; anti-CD11c (N418) conjugated to PE or PE-Cy5.5; anti-Siglec H (eBio440c)

conjugated to PE-Cy7 or PE; anti-CD8b (eBioH35-17.2) conjugated to eFluor 450; anti-CD44

(IM7) conjugated to BV605; anti-CD4 (RM4-5) conjugated to PE-Cy5.5; anti-CD69 (H1.2F3)

conjugated to PE-Cy7; anti-CD62L (MEL-14) conjugated to APC; anti-CD5 (53-7.3) conjugated

to APC-eFluor780 or biotin; anti-Siglec F (E50-2440) conjugated to BV421; anti-F4/80 (BM8)

conjugated to PE-Cy7; anti-CD64 (X54-5/7.1) conjugated to APC; anti-NK1.1 (PK136)

conjugated to Alexa Fluor 700; anti-CD172α (P84) conjugated to FITC.

Single cell RNA sequencing

Eight-week-old male ApoE$^{-/-}$ mice were fed with 20-week chow diet or HFD. The single cell

suspension of spleen, lymph node and aorta were obtained according to the flow cytometry

analysis section. Single cell suspension from spleens and lymph nodes were firstly removed the

CD5^{+}, CD19^{+} and CD11b^{+} cells via SAV conjugated beads. Then two million live cells from

spleen and lymph node were stained with mix1 (CD45.2-PE-Cy7, CD11c-PE-Cy5.5,

CD172α-FITC, MHCII- APC-eFluor780 and Xcr1-APC), and all cells from aorta were stained

with mix2 (CD45.2-PE-Cy7, CD11b-super bright 600, CD11c-PE-Cy5.5, MHCII- FITC, CD3-

Alexa Fluor 700, CD19- Alexa Fluor 700, Ly6G- Alexa Fluor 700 and Xcr1-APC) for 30 minutes

at 4 °C. After washing with 1 mL FACS buffer two times, cells from different organs were stained

with different anti-CD45 antibodies conjugated with barcode oligos from BD™ Ms Single Cell

Sample Multiplexing Kit (BD Bioscience, Cat: 633793). Finally, cells were resuspended with

Sytox blue and CD11c$^+$ MHCII$^+$ Xcr1$^+$ cells were sorted and subjected to the BD Rhapsody

Express system. Then cDNA and sample tag libraries were built with BD Rhapsody whole

transcriptome amplification (WTA) reagent kit (BD Bioscience, Cat: 633733, 633773, 633801,

664887) following manufacturer's instructions, and sequenced on an illumine Novaseq 600

sequencer. Pair-end Fastq files of sample tag and WTA data were processed via BD Rhapsody^{TM}

analysis pipeline v2.0, and the resultant dataset was mainly analysis using SeqGeq^{TM} software,

which contain Lex-BDSMK and Seurat v4.04 plugin components. Raw data and processed data

were uploaded into GEO (Accession number: GSE279370).

Statistical analysis

All data were presented as means ± SEM (Standard Error of the Mean), reflecting the average

values of biological measurements across multiple samples. Data analysis was performed using

GraphPad Prism software (version 8.0). Statistical significance was assessed by both parametric

(unpaired Student's *t*-test) and nonparametric (Mann–Whitney test) methods when comparing two

groups. The assumptions of the tests were checked to ensure appropriate application. Significant

differences between different groups were set at * $P<0.05$, ** $P<0.01$, *** $P<0.001$ and ****

$P<0.0001$. Graphical representations of the data will include bar graphs with error bars denoting

SEM.

Acknowledgements

We thank the animal facility and flow cytometry facility of GRIS in Xinxiang Medical University

for providing technical support and assistance. Generation of murine genetic models were

supported by 111 program (D20036).

Author Contributions

Lichen Zhang and **Yinming Liang** conceived the project. **Tianhan Li** and **Liaoxun Lu** wrote the manuscript and carried out the majority of the experiments. **Bernard Malissen** and **Lichen Zhang** designed the Xcr1Cre-Gfp knock-in mice and Xcl1 knockout mice. **Juanjuan Qiu**, **Xin Dong**, **Le Yang**, **Kexin He**, and **Yanrong Gu** assisted in the experimental procedures. **Binhui Zhou** performed cell sorting for scRNA sequencing. **Tingting Jia** and **Rong Huang** contributed to data curation and validation. **Toby Lawrence**, **Marie Malissen**, and **Hui Wang** participated in the manuscript review and editing. **Guixue Wang** assisted in the preparation of human samples. **Yinming Liang**, **Lichen Zhang**, and **Tianhan Li** contributed to obtaining the funding for the project.

Sources of Funding

This research was funded by projects 2022WZ1021 and 2023DK2005 to YL, 2022RC1223 to LZ, 242102310030 to TL, and the NSFC grants 82301972 and 32170879 to TL and LZ.

Declaration of competing interest

The authors declare no competing interests.

 Figure legends

 Figure 1. Progressive accumulation of Xcr1$^+$ cDC1 cells in human and mouse advanced

 atherosclerotic lesions

 A, Representative images of immunohistochemical staining demonstrate Xcr1 expression in

 normal areas and plaque areas of the femoral artery from three cardiovascular disease patients. **B**,

 Quantitative analysis of the Xcr1-positive area in normal area and plaque regions. **C**,

 Representative immunofluorescence images depict DAPI (blue), Xcr1 (green) and CD11c (red)

 within the lesions of aortic root of ApoE$^{-/-}$ mice fed with 16-week chow diet or HFD. **D**,

 Quantitative analysis of Xcr1 and CD11c double positive area in lesion area (n = 6). Scale bars,

 100 μm. **E**, Representative images depicting immunohistochemical staining of Xcr1 in lesions of

 the aortic root of ApoE$^{-/-}$ mice fed a HFD for varying durations. **F**, Quantitative analysis of the

 Xcr1-positive area in the lesion regions. (HFD-8W mice, n = 11; HFD-12W mice, n = 15;

 HFD-16W mice, n = 29). Data represent as mean ± SEM. *** *P*<0.001.

 Figure 2. Selective expression of Xcr1$^{\text{Cre-Gfp}}$ in cDC1 cells.

 A, Diagram illustrating the bone marrow transfer process. **B** and **C**, Representative flow

 cytometric analysis and corresponding quantification of GFP$^+$RFP$^+$ in cDC1 and non-cDC1

 populations within the cDC1 and non-cDC1 populations of the spleen. These data were obtained

 from ApoE$^{-/-}$ mice that had received bone marrow transplants from either WT or Xcr1$^{\text{Cre-Gfp}}$

 Rosa26$^{\text{LSL-RFP}}$ donors and were maintained on a HFD for 16 weeks (WT, n = 4; Xcr1$^{\text{Cre-Gfp}}$

 Rosa26$^{\text{LSL-RFP}}$, n = 3). **D** through **F**, Eight-week-old ApoE$^{-/-}$, Xcr1$^{\text{Cre-Gfp}}$ ApoE$^{-/-}$ and Xcr1$^{\text{Cre-Gfp}}$

 Rosa26$^{\text{LSL-DTA}}$ ApoE$^{-/-}$ mice fed with a HFD for 16 weeks (n = 3). **D** and **E**, Representative flow

cytometric analysis and quantification of the percentage of GFP$^+$ cells among pDC, cDC1 and

cDC2 cells populations in the spleen. **F**, Quantification of the percentage of pDC, cDC1 and cDC2

cells in spleen. Data represent as mean ± SEM. **** P<0.0001; NS, non-significant.

Figure 3. Specific depletion of Xcr1$^+$ cDC1 cells in ApoE$^{-/-}$ mice reduces atherosclerosis

progression with no effect on macrophages.

A through **I**, Eight-week-old ApoE$^{-/-}$ mice and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice were fed

on a 16-week HFD to develop atherosclerosis. **A**, ORO staining of the descending aortas. Scale

bar, 5 mm. **B**, Quantification of the lesion area in the descending aorta (ApoE$^{-/-}$ mice, n = 9;

Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 6). **C**, ORO and H&E staining of the aortic roots of

representative mice from each group. **D**, Quantification of lesion area in the aortic root. (ApoE$^{-/-}$

mice, n = 7; Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 8). Scale bar, 200 μm. **E** and **F**, H&E

staining and quantification of necrotic core area in the lesions of aortic root. (ApoE$^{-/-}$ mice, n = 7;

Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 8). Scale bar, 200 μm. **G** and **H**, Representative flow

cytometric analysis and quantification of macrophages in the aorta. (ApoE$^{-/-}$ mice, n = 8;

Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 6). **I** and **J,** Representative immunohistochemical

staining images of CD68 and quantification of CD68 positive area in the lesions of the aortic root.

(ApoE$^{-/-}$ mice, n = 6; Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice, n = 8). Scale bar, 100 μm. Data

represent as mean ± SEM. NS, non-significant.

Figure 4. Specific depletion of Xcr1$^+$ cDC1 cells in ApoE$^{-/-}$ mice decreases T cell activation

within aorta.

A through **D**, Eight-week-old ApoE$^{-/-}$ mice and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice fed a HFD

for 16 weeks. **A**, Representative flow cytometric analysis cDC1 cells in aortas. **B**, Quantification

of cDC1 cells in aorta (n = 4). **C**, Representative flow cytometric analysis T cells in aortas **D**,

Quantification of T cell populations, including CD4$^+$, CD4$^+$CD69L$^+$, CD8b$^+$ and CD8b$^+$CD69L$^+$ T

cells in aortas (n = 7). Data represent as mean ± SEM. * $P<0.05$, ** $P<0.01$, *** $P<0.001$, ns,

non-significant.

Figure 5. Sc-RNA sequencing analysis of the aorta, lymph nodes and spleen in ApoE$^{-/-}$ mice

fed with a 20-week HFD.

A, UMAP plot delineates 10 annotated cell types of CD11c$^+$MHCII$^+$Xcr1$^+$cDC1 cells from aorta,

lymph node and spleen in ApoE$^{-/-}$ mice maintained on a HFD for 20 weeks. **B**, Heatmap statistic

map displays the top ten up-regulated genes across various clusters. **C**, Histogram plot illustrates

the proportion of different cell types among groups. **D** and **E**, UMAP plot representing CD11c$^+$

MHCII$^+$Xcr1$^+$cDC1 cells from aorta, lymph nodes, spleen, and a merged plot in ApoE$^{-/-}$ mice

maintained on a HFD for 20 weeks.

Figure 6. Xcl1 deficiency reduces atherosclerotic lesion formation in ApoE$^{-/-}$ mice.

A through **I**, Eight-week-old ApoE$^{-/-}$ mice and Xcl1$^{-/-}$ ApoE$^{-/-}$ mice fed a HFD for 19 weeks (n =

8 per group). **A**, ORO staining of the descending aortas. Scale bar, 5 mm. **B**, Quantification of the

lesion area in aortas. **C**, ORO and H&E staining of aortic roots. Scale bar, 200 μm. **D**,

Quantification of the lesion area in aortic roots. **E** and **F**, Representative liver images and

quantification of ORO positive area in livers (n = 24 regions from 8 mice, 3 regions per mice).

Scale bar, 100 μm. **G**, The concentrations of TC, TG, LDL and HDL in the serum. **H**, Body weight

of mice before and after feeding with 19-week HFD. **I**, Representative immunohistochemical

staining of CD68 in aortic roots. **J**, Quantification of CD68 positive area in the lesions of aortic

roots (n = 24 lesions from 8 mice, 3 lesions per mice). Scale bar, 100 μm. Data represent as mean

± SEM. ** *P*<0.01, *** *P*<0.001, NS, non-significant.

Figure 7. Xcl1 deficiency reduces CD8$^+$ T cells frequency in the aorta from.

A through **D**, Eight-week-old ApoE$^{-/-}$ mice and Xcl1$^{-/-}$ApoE$^{-/-}$ mice fed with a 16-week HFD. (n

= 7). **A** and **B**, Representative flow cytometric analysis and quantification of pDC, cDC1 and

cDC2 cells in aortas. **C** and **D**, Representative flow cytometric analysis and quantification of T

cell subsets, including CD4$^+$, CD4$^+$CD69L$^+$, CD8b$^+$ and CD8b$^+$CD69L$^+$ T cells in aortas. Data

represent as mean ± SEM. * *P*<0.05, NS, non-significant.

Figure   **S1.**   **Schematic**   **diagram**   **illustrating**   **the**   **knock-in**   **of**   **the**

5'HA-iCre-P2A-EGFP-P2A-3'HA vector into the mouse Xcr1 locus.

Figure S2. Flow cytometric analysis of cDC1 cells in the lymph nodes and spleens of ApoE$^{-/-}$

mice and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ApoE$^{-/-}$ mice fed a 7-weeks chow diet.

A through **D**, ApoE$^{-/-}$ mice and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ApoE$^{-/-}$ mice fed with a 7-week chow

diet. (n = 7). **A**, Representative flow cytometric analysis of cDC1 cells in lymph nodes. **B**,

Quantification of the frequencies and number of cDC1 cell in lymph nodes. **C**, Representative

flow cytometric analysis cDC1 cells in the spleens. **D**, Quantification of the frequencies and

number of cDC1 cell in the spleens. Data represent the mean ± SEM. *** $P<0.001$.

Figure S3. Lipid profile, body weight and ORO staining of the liver in ApoE$^{-/-}$ mice and

Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice fed a 16-week HFD.

A through **D**, Eight-week-old ApoE$^{-/-}$ mice (n =7) and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice (n

=8) were fed on a 16-week HFD to develop atherosclerosis. **A**, Concentrations of TC, TG, LDL

and HDL in the serum. **B**, Body weight measurements of both groups of mice. **C**, ORO staining of

liver from each group. Scale bar, 100 μm. **D**, Quantification of ORO positive area in liver. Data

represent the mean ± SEM. ** $P<0.01$, *** $P<0.001$, NS, non-significant.

Figure S4. Flow cytometric analysis of cDC1 cells in the lymph nodes and spleens of ApoE$^{-/-}$

and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice fed a 16-week HFD.

A through **D**, Eight-week-old ApoE$^{-/-}$ mice (n =7) and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice (n

=8) were fed on a 16-week HFD to develop atherosclerosis. **A**, Representative flow cytometric

analysis of cDC1 cells in lymph nodes. **B**, Quantification of the frequencies and numbers of cDC1

cells in lymph nodes. **C**, Representative flow cytometric analysis of cDC1 cells in the spleens. **D**,

Quantification of the frequencies and numbers of cDC1 cells in the spleens. Data represent the

mean ± SEM. ** $P<0.01$, *** $P<0.001$, NS, non-significant.

Figure S5. Flow cytometric analysis of T cells in the lymph nodes and spleens of ApoE$^{-/-}$ mice

and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice fed a 16-week HFD.

A through **F**, Eight-week-old ApoE$^{-/-}$ mice (n =7) and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ ApoE$^{-/-}$ mice (n

=8) were fed on a 16-week HFD to develop atherosclerosis. **A**, Representative flow cytometric

analysis of $CD4^+$, $CD8^+$, $CD4^+CD69^+$ and $CD8^+CD69^+$ T cells in lymph nodes. **B** and **C**,

Quantification of the frequencies and numbers of $CD4^+$, $CD8^+$, $CD4^+CD69^+$ and $CD8^+CD69^+$ T

cells in lymph nodes. **D**, Representative flow cytometric analysis and quantification of $CD4^+$,

$CD8^+$, $CD4^+CD69^+$ and $CD8^+CD69^+$ T cells in the spleens. **E** and **F**, Quantification of the

frequencies and numbers of $CD4^+$, $CD8^+$, $CD4^+CD69^+$ and $CD8^+CD69^+$ T cells in the spleens.

Data represent the mean ± SEM. * $P<0.05$, ** $P<0.01$, *** $P<0.001$, NS, non-significant.

Figure S6. T cells analysis in the lymph nodes and spleens of ApoE$^{-/-}$ mice and Xcr1$^{Cre-Gfp}$

Rosa26$^{LSL-DTA}$ApoE$^{-/-}$ mice fed a 7-week chow diet.

A through **F**, ApoE$^{-/-}$ mice and Xcr1$^{Cre-Gfp}$ Rosa26$^{LSL-DTA}$ApoE$^{-/-}$ mice fed with a 7-week chow

diet. (n = 7). **A**, Representative flow cytometric analysis and quantification of $CD4^+$, $CD4^+CD44^+$

$CD62L^-$, $CD8^+$ and $CD8^+CD44^+CD62L^-$ T cells in lymph nodes. **B**, Quantification of the

frequencies of $CD4^+$, $CD8^+$, $CD4^+CD69^+$ and $CD8^+CD69^+$ T cells in the lymph nodes. **C**,

Representative flow cytometric analysis and quantification of $CD4^+$, $CD4^+CD44^+CD62L^-$, $CD8^+$

and $CD8^+CD44^+CD62L^-$ T cells in the spleens. **D**, Quantification of the frequencies of $CD4^+$,

$CD8^+$, $CD4^+CD69^+$ and $CD8^+CD69^+$ T cells in the spleens. **E** and **F**, Quantification of the

numbers of $CD4^+$ and $CD8^+$ T cells in the lymph nodes and spleens. Data represent the mean ±

SEM. * $P<0.05$, ** $P<0.01$, NS, non-significant.

Figure S7. Depletion of Xcr1$^+$ cDC1 cells of bone marrow reduces cDC1 cells without

affecting macrophages in atherosclerotic lesion of ApoE$^{-/-}$ mice.

A, Diagram of the bone marrow transfer process. ApoE$^{-/-}$ mice transplanted with bone marrow

from WT or Xcr1$^{Cre\text{-}Gfp}$ Rosa26$^{LSL\text{-}DTA}$ donors before and after 16 weeks on a HFD (n = 7). **B**,

Body weight of each group. **C**, Serum concentrations of TC, TG, LDL and HDL. **D**,

Representative ORO staining images of liver sections and quantification of the ORO positive area

percentage in the liver. Scale bar, 50 μm. **E**, Representative flow cytometric analysis of pDC,

cDC1 and cDC2 cells in the aortas, Lin$^-$ means CD3$^-$ CD19$^-$ Ly6G$^-$ NK1.1$^-$. **F**, Representative flow

cytometric analysis of macrophages in the aortas of each group. Data represent as mean ± SEM.

*** *P*<0.001, NS, non-significant.

Figure S8. The single-cell RNA sequencing process.

Figure S9. DC cells analysis in lymph nodes and spleens of ApoE$^{-/-}$ and Xcl1$^{-/-}$ApoE$^{-/-}$ mice

fed with 19-week HFD.

A through **D**, Eight-week-old ApoE$^{-/-}$ (n = 9) and Xcl1$^{-/-}$ApoE$^{-/-}$ mice (n = 7) fed a HFD for 19

weeks. **A**, Representative flow cytometric analysis and quantification of pDC, cDCs, cDC1 and

cDC2 cells in lymph nodes. **B**, Quantification of absolute counts of pDC, cDCs, cDC1 and cDC2

cells in lymph nodes. **C**, Representative flow cytometric analysis of pDC, cDC1 and cDC2 cells in

spleens. **D**, Quantification of absolute counts of pDC, cDCs, cDC1 and cDC2 cells in spleens.

Data represent the mean ± SEM. NS, non-significant.

Figure S10. T cells analysis in lymph node and spleen of ApoE$^{-/-}$ and Xcl1$^{-/-}$ApoE$^{-/-}$ mice fed

with 16-week HFD.

A through **D**, Eight-week-old ApoE$^{-/-}$ (n = 9) and Xcl1$^{-/-}$ApoE$^{-/-}$ mice (n = 7) fed a HFD for 19

weeks. **A** and **B**, Representative flow cytometric analysis and quantification of absolute counts of

CD4$^+$, CD8$^+$, CD4$^+$CD69$^+$ and CD8$^+$CD69$^+$ T cells in lymph nodes. **C** and **D**, Representative flow

cytometric analysis and quantification of absolute counts of CD4$^+$, CD8$^+$, CD4$^+$CD69$^+$ and

CD8$^+$CD69$^+$ T cells in spleens. Data represent the mean ± SEM. NS, non-significant.

Supplementary table 1: **The sequences of the sgRNA and primers used for generation and**

validation of Xcr1$^{Cre-Gfp}$ knock-in mice and Xcl1 knockout mice

Supplementary table 2: **Up regulated genes in Xcr1$^+$ cDC1 cells from aorta vs spleen, and**

aorta vs lymph node in ApoE-deficient mice fed a HFD for 20 weeks

Supplementary table 3: **The top ten high-expressed marker genes in ten clusters**

References

1.   Bjorkegren   JLM,   Lusis   AJ.   Atherosclerosis: Recent   developments.   Cell.

2022;185(10):1630-45. doi: 10.1016/j.cell.2022.04.004. PubMed PMID: 35504280; PubMed

Central PMCID: PMC9119695.

2.   Hansson GK, Hermansson A. The immune system in atherosclerosis. Nat Immunol.

2011;12(3):204-12. Epub 2011/02/16. doi: 10.1038/ni.2001. PubMed PMID: 21321594.

3.   Lusta KA, Poznyak AV, Sukhorukov VN, Eremin, II, Nadelyaeva, II, Orekhov AN.

Hypotheses on Atherogenesis Triggering: Does the Infectious Nature of Atherosclerosis

Development Have a Substruction? Cells. 2023;12(5). doi: 10.3390/cells12050707. PubMed

PMID: 36899843; PubMed Central PMCID: PMC10001176.

4.   Hansson GK, Robertson AK, Soderberg-Naucler C. Inflammation and atherosclerosis.

Annu   Rev   Pathol.   2006;1:297-329.   Epub   2007/11/28.   doi:

10.1146/annurev.pathol.1.110304.100100. PubMed PMID: 18039117.

5.   Hasham SN, Pillarisetti S. Vascular lipases, inflammation and atherosclerosis. Clin Chim

Acta. 2006;372(1-2):179-83. Epub 2006/06/13. doi: 10.1016/j.cca.2006.04.020. PubMed PMID:

16765928.

6.   Steinman RM, Hawiger D, Liu K, Bonifaz L, Bonnyay D, Mahnke K, et al. Dendritic cell

function in vivo during the steady state: a role in peripheral tolerance. Ann N Y Acad Sci.

2003;987:15-25. Epub 2003/05/03. doi: 10.1111/j.1749-6632.2003.tb06029.x. PubMed PMID:

12727620.

7.   Ma-Krupa W, Jeon MS, Spoerl S, Tedder TF, Goronzy JJ, Weyand CM. Activation of

arterial wall dendritic cells and breakdown of self-tolerance in giant cell arteritis. J Exp Med.

2004;199(2):173-83. Epub 2004/01/22. doi: 10.1084/jem.20030850. PubMed PMID: 14734523;

PubMed Central PMCID: PMCPMC2211768.

8.   Liu P, Yu YR, Spencer JA, Johnson AE, Vallanat CT, Fong AM, et al. CX3CR1 deficiency

impairs dendritic cell accumulation in arterial intima and reduces atherosclerotic burden.

Arterioscler   Thromb   Vasc   Biol.   2008;28(2):243-50.   Epub   2007/12/15.   doi:

10.1161/ATVBAHA.107.158675. PubMed PMID: 18079406.

9.   Jongstra-Bilen J, Haidari M, Zhu SN, Chen M, Guha D, Cybulsky MI. Low-grade chronic

inflammation in regions of the normal mouse arterial intima predisposed to atherosclerosis. J

Exp Med. 2006;203(9):2073-83. Epub 2006/08/09. doi: 10.1084/jem.20060245. PubMed PMID:

16894012; PubMed Central PMCID: PMCPMC2118407.

10. Bobryshev YV, Lord RS. Ultrastructural recognition of cells with dendritic cell morphology

in human aortic intima. Contacting interactions of Vascular Dendritic Cells in athero-resistant

and athero-prone areas of the normal aorta. Arch Histol Cytol. 1995;58(3):307-22. Epub

1995/08/01. doi: 10.1679/aohc.58.307. PubMed PMID: 8527238.

11. Kawahara I, Kitagawa N, Tsutsumi K, Nagata I, Hayashi T, Koji T. The expression of

vascular dendritic cells in human atherosclerotic carotid plaques. Hum Pathol.

2007;38(9):1378-85. Epub 2007/06/09. doi: 10.1016/j.humpath.2007.02.004. PubMed PMID:

17555794.

12. Zernecke A, Erhard F, Weinberger T, Schulz C, Ley K, Saliba AE, et al. Integrated

single-cell analysis based classification of vascular mononuclear phagocytes in mouse and

human atherosclerosis. Cardiovasc Res. 2022. Epub 2022/10/04. doi: 10.1093/cvr/cvac161.

PubMed PMID: 36190844.

13. Eisenbarth SC. Dendritic cell subsets in T cell programming: location dictates function.

Nat Rev Immunol. 2019;19(2):89-103. Epub 2018/11/23. doi: 10.1038/s41577-018-0088-1.

PubMed PMID: 30464294; PubMed Central PMCID: PMCPMC7755085.

14. Bayerl F, Meiser P, Donakonda S, Hirschberger A, Lacher SB, Pedde AM, et al.

Tumor-derived prostaglandin E2 programs cDC1 dysfunction to impair intratumoral

orchestration of anti-cancer T cell responses. Immunity. 2023;56(6):1341-58 e11. Epub

2023/06/15. doi: 10.1016/j.immuni.2023.05.011. PubMed PMID: 37315536.

15. Legein B, Janssen EM, Theelen TL, Gijbels MJ, Walraven J, Klarquist JS, et al. Ablation

of CD8alpha(+) dendritic cell mediated cross-presentation does not impact atherosclerosis in

hyperlipidemic mice. Scientific reports. 2015;5:15414. Epub 20151021. doi:

10.1038/srep15414. PubMed PMID: 26486587; PubMed Central PMCID: PMCPMC4614009.

16. Gil-Pulido J, Cochain C, Lippert MA, Schneider N, Butt E, Amezaga N, et al. Deletion of

Batf3-dependent antigen-presenting cells does not affect atherosclerotic lesion formation in

mice. PLoS One. 2017;12(8):e0181947. Epub 20170803. doi: 10.1371/journal.pone.0181947.

PubMed PMID: 28771609; PubMed Central PMCID: PMCPMC5542449.

17. Choi JH, Cheong C, Dandamudi DB, Park CG, Rodriguez A, Mehandru S, et al. Flt3

signaling-dependent dendritic cells protect against atherosclerosis. Immunity.

2011;35(5):819-31. Epub 2011/11/15. doi: 10.1016/j.immuni.2011.09.014. PubMed PMID:

22078798.

18. Clement M, Haddad Y, Raffort J, Lareyre F, Newland SA, Master L, et al. Deletion of IRF8

(Interferon Regulatory Factor 8)-Dependent Dendritic Cells Abrogates Proatherogenic

Adaptive Immunity. Circ Res. 2018;122(6):813-20. Epub 2018/02/14. doi:

10.1161/CIRCRESAHA.118.312713. PubMed PMID: 29436389.

19. Haddad Y, Lahoute C, Clement M, Laurans L, Metghalchi S, Zeboudj L, et al. The

Dendritic Cell Receptor DNGR-1 Promotes the Development of Atherosclerosis in Mice. Circ

Res. 2017;121(3):234-43. Epub 2017/06/14. doi: 10.1161/CIRCRESAHA.117.310960.

PubMed PMID: 28607102.

20. Wohn C, Le Guen V, Voluzan O, Fiore F, Henri S, Malissen B. Absence of MHC class II

on cDC1 dendritic cells triggers fatal autoimmunity to a cross-presented self-antigen. Sci

Immunol. 2020;5(45). Epub 2020/03/15. doi: 10.1126/sciimmunol.aba1896. PubMed PMID:

32169954.

21. Mundt S, Mrdjen D, Utz SG, Greter M, Schreiner B, Becher B. Conventional DCs sample

and present myelin antigens in the healthy CNS and allow parenchymal T cell entry to initiate

neuroinflammation. Sci Immunol. 2019;4(31). doi: 10.1126/sciimmunol.aau8380. PubMed

PMID: 30679199.

22. Robbins CS, Hilgendorf I, Weber GF, Theurl I, Iwamoto Y, Figueiredo JL, et al. Local

proliferation dominates lesional macrophage accumulation in atherosclerosis. Nat Med.

2013;19(9):1166-72. Epub 2013/08/13. doi: 10.1038/nm.3258. PubMed PMID: 23933982;

PubMed Central PMCID: PMCPMC3769444.

23. Potteaux S, Gautier EL, Hutchison SB, van Rooijen N, Rader DJ, Thomas MJ, et al.

Suppressed monocyte recruitment drives macrophage removal from atherosclerotic plaques

of Apoe-/- mice during disease regression. J Clin Invest. 2011;121(5):2025-36. Epub

2011/04/21. doi: 10.1172/JCI43802. PubMed PMID: 21505265; PubMed Central PMCID:

PMCPMC3083793.

24. Yilmaz A, Weber J, Cicha I, Stumpf C, Klein M, Raithel D, et al. Decrease in circulating

myeloid dendritic cell precursors in coronary artery disease. J Am Coll Cardiol.

2006;48(1):70-80. Epub 2006/07/04. doi: 10.1016/j.jacc.2006.01.078. PubMed PMID:

16814651.

25. Alberts-Grill N, Denning TL, Rezvan A, Jo H. The role of the vascular dendritic cell

network in atherosclerosis. Am J Physiol Cell Physiol. 2013;305(1):C1-21. Epub 2013/04/05.

doi: 10.1152/ajpcell.00017.2013. PubMed PMID: 23552284; PubMed Central PMCID:

PMCPMC3725520.

26. Yan Y, Thakur M, van der Vorst EPC, Weber C, Doring Y. Targeting the chemokine

network in atherosclerosis. Atherosclerosis. 2021;330:95-106. Epub 2021/07/13. doi:

10.1016/j.atherosclerosis.2021.06.912. PubMed PMID: 34247863.

27. Eickhoff S, Brewitz A, Gerner MY, Klauschen F, Komander K, Hemmi H, et al. Robust

Anti-viral Immunity Requires Multiple Distinct T Cell-Dendritic Cell Interactions. Cell.

2015;162(6):1322-37. Epub 2015/08/25. doi: 10.1016/j.cell.2015.08.004. PubMed PMID:

26296422; PubMed Central PMCID: PMCPMC4567961.

28. Weber C, Zernecke A, Libby P. The multifaceted contributions of leukocyte subsets to

atherosclerosis: lessons from mouse models. Nat Rev Immunol. 2008;8(10):802-15. Epub

2008/10/01. doi: 10.1038/nri2415. PubMed PMID: 18825131.

29. Worbs T, Hammerschmidt SI, Forster R. Dendritic cell migration in health and disease.

Nat Rev Immunol. 2017;17(1):30-48. Epub 2016/11/29. doi: 10.1038/nri.2016.116. PubMed

PMID: 27890914.

30. Li T, Safitri M, Zhang K, Wang Y, Huang L, Zhu Y, et al. Downregulation of G3BP2

reduces atherosclerotic lesions in ApoE(-/-) mice. Atherosclerosis. 2020;310:64-74. Epub

2020/09/13. doi: 10.1016/j.atherosclerosis.2020.08.003. PubMed PMID: 32919187.

31. Mattiuz R, Wohn C, Ghilas S, Ambrosini M, Alexandre YO, Sanchez C, et al. Novel

Cre-Expressing Mouse Strains Permitting to Selectively Track and Edit Type 1 Conventional

Dendritic Cells Facilitate Disentangling Their Complexity in vivo. Front Immunol. 2018;9:2805.

Epub 2018/12/20. doi: 10.3389/fimmu.2018.02805. PubMed PMID: 30564233; PubMed

Central PMCID: PMCPMC6288293.

32. Luche H, Weber O, Nageswara Rao T, Blum C, Fehling HJ. Faithful activation of an

extra-bright red fluorescent protein in "knock-in" Cre-reporter mice ideally suited for lineage

tracing studies. European journal of immunology. 2007;37(1):43-53. Epub 2006/12/16. doi:

10.1002/eji.200636745. PubMed PMID: 17171761.

33. De Meyer GRY, Zurek M, Puylaert P, Martinet W. Programmed death of macrophages in

atherosclerosis: mechanisms and therapeutic targets. Nat Rev Cardiol. 2024. Epub

2024/01/02. doi: 10.1038/s41569-023-00957-0. PubMed PMID: 38163815.

34. Yu EPK, Reinhold J, Yu H, Starks L, Uryga AK, Foote K, et al. Mitochondrial Respiration

Is Reduced in Atherosclerosis, Promoting Necrotic Core Formation and Reducing Relative

Fibrous Cap Thickness. Arterioscler Thromb Vasc Biol. 2017;37(12):2322-32. Epub

2017/10/04. doi: 10.1161/ATVBAHA.117.310042. PubMed PMID: 28970293; PubMed Central

PMCID: PMCPMC5701734.

35. Piedrahita JA, Zhang SH, Hagaman JR, Oliver PM, Maeda N. Generation of mice

carrying a mutant apolipoprotein E gene inactivated by gene targeting in embryonic stem cells.

Proc Natl Acad Sci U S A. 1992;89(10):4471-5. Epub 1992/05/15. doi:

10.1073/pnas.89.10.4471. PubMed PMID: 1584779; PubMed Central PMCID:

PMCPMC49104.

36. Plump AS, Smith JD, Hayek T, Aalto-Setala K, Walsh A, Verstuyft JG, et al. Severe

hypercholesterolemia and atherosclerosis in apolipoprotein E-deficient mice created by

homologous recombination in ES cells. Cell. 1992;71(2):343-53. Epub 1992/10/16. doi:

10.1016/0092-8674(92)90362-g. PubMed PMID: 1423598.

37.   Fernandez DM, Rahman AH, Fernandez NF, Chudnovskiy A, Amir ED, Amadori L, et al.

Single-cell   immune   landscape   of   human   atherosclerotic   plaques.   Nat   Med.

2019;25(10):1576-88. Epub 2019/10/09. doi: 10.1038/s41591-019-0590-4. PubMed PMID:

31591603; PubMed Central PMCID: PMCPMC7318784.

38.   Afek A, Harats D, Roth A, Keren G, George J. Evidence for the involvement of T cell

costimulation through the B-7/CD28 pathway in atherosclerotic plaques from apolipoprotein E

knockout   mice.   Exp   Mol   Pathol.   2004;76(3):219-23.   Epub   2004/05/06.   doi:

10.1016/j.yexmp.2003.12.001. PubMed PMID: 15126104.

39.   Cancel JC, Crozat K, Dalod M, Mattiuz R. Are Conventional Type 1 Dendritic Cells Critical

for Protective Antitumor Immunity and How? Front Immunol. 2019;10:9. Epub 2019/02/28. doi:

10.3389/fimmu.2019.00009.   PubMed   PMID: 30809220;   PubMed   Central   PMCID:

PMCPMC6379659.

40.   Li N, Steiger S, Fei L, Li C, Shi C, Salei N, et al. IRF8-Dependent Type I Conventional

Dendritic Cells (cDC1s) Control Post-Ischemic Inflammation and Mildly Protect Against

Post-Ischemic Acute Kidney Injury and Disease. Front Immunol. 2021;12:685559. Epub

2021/07/09. doi: 10.3389/fimmu.2021.685559. PubMed PMID: 34234783; PubMed Central

PMCID: PMCPMC8255684.

41.   Bottcher JP, Bonavita E, Chakravarty P, Blees H, Cabeza-Cabrerizo M, Sammicheli S, et

al. NK Cells Stimulate Recruitment of cDC1 into the Tumor Microenvironment Promoting

Cancer   Immune   Control.   Cell.   2018;172(5):1022-37   e14.   doi: 10.1016/j.cell.2018.01.004.

PubMed PMID: 29429633; PubMed Central PMCID: PMC5847168.

42.   Dorner BG, Dorner MB, Zhou X, Opitz C, Mora A, Guttler S, et al. Selective expression of

the chemokine receptor XCR1 on cross-presenting dendritic cells determines cooperation with

CD8+ T cells. Immunity. 2009;31(5):823-33. doi: 10.1016/j.immuni.2009.08.027. PubMed

PMID: 19913446.

43. Xu F, He D, Ning R, Zeng B, Thompson CW, Li Y, et al. Genetic diversity of chemokine

XCL1 and its receptor XCR1 in murine rodents. Developmental and comparative immunology.

2019;98:80-8. doi: 10.1016/j.dci.2019.04.008. PubMed PMID: 31026469.

44. Yoshida T, Izawa D, Nakayama T, Nakahara K, Kakizaki M, Imai T, et al. Molecular

cloning of mXCR1, the murine SCM-1/lymphotactin receptor. FEBS letters. 1999;458(1):37-40.

doi: 10.1016/s0014-5793(99)01114-x. PubMed PMID: 10518929.

45. Tall AR, Bornfeldt KE. Inflammasomes and Atherosclerosis: a Mixed Picture. Circ Res.

2023;132(11):1505-20. doi: 10.1161/CIRCRESAHA.123.321637. PubMed PMID: 37228237;

PubMed Central PMCID: PMC10213995.

46. Vallejo J, Cochain C, Zernecke A, Ley K. Heterogeneity of immune cells in human

atherosclerosis revealed by scRNA-Seq. Cardiovasc Res. 2021;117(13):2537-43. doi:

10.1093/cvr/cvab260. PubMed PMID: 34343272; PubMed Central PMCID: PMC8921647.

47. Guo X, Zhou Y, Wu T, Zhu X, Lai W, Wu L. Generation of mouse and human dendritic

cells in vitro. J Immunol Methods. 2016;432:24-9. Epub 2016/02/16. doi:

10.1016/j.jim.2016.02.011. PubMed PMID: 26876301.

48. Anselmi G, Helft J, Guermonprez P. Development and function of human dendritic cells in

humanized mice models. Mol Immunol. 2020;125:151-61. Epub 2020/07/21. doi:

10.1016/j.molimm.2020.07.005. PubMed PMID: 32688117.

49. Blanco P, Palucka AK, Pascual V, Banchereau J. Dendritic cells and cytokines in human

inflammatory and autoimmune diseases. Cytokine Growth Factor Rev. 2008;19(1):41-52.

Epub 2008/02/09. doi: 10.1016/j.cytogfr.2007.10.004. PubMed PMID: 18258476; PubMed

Central PMCID: PMCPMC2413068.

50. Steinman RM. Dendritic cells: understanding immunogenicity. European journal of

immunology. 2007;37 Suppl 1:S53-60. Epub 2007/11/01. doi: 10.1002/eji.200737400.

PubMed PMID: 17972346.

51. Steinman RM, Adams JC, Cohn ZA. Identification of a novel cell type in peripheral

lymphoid organs of mice. IV. Identification and distribution in mouse spleen. J Exp Med.

1975;141(4):804-20. Epub 1975/04/01. PubMed PMID: 1127378; PubMed Central PMCID:

PMCPMC2189754.

52. Steinman RM, Kaplan G, Witmer MD, Cohn ZA. Identification of a novel cell type in

peripheral lymphoid organs of mice. V. Purification of spleen dendritic cells, new surface

markers, and maintenance in vitro. J Exp Med. 1979;149(1):1-16. Epub 1979/01/01. doi:

10.1084/jem.149.1.1. PubMed PMID: 762493; PubMed Central PMCID: PMCPMC2184752.

53. Clark GJ, Silveira PA, Hogarth PM, Hart DNJ. The cell surface phenotype of human

dendritic cells. Semin Cell Dev Biol. 2019;86:3-14. Epub 2018/03/03. doi:

10.1016/j.semcdb.2018.02.013. PubMed PMID: 29499385.

54. Brewitz A, Eickhoff S, Dahling S, Quast T, Bedoui S, Kroczek RA, et al. CD8(+) T Cells

Orchestrate pDC-XCR1(+) Dendritic Cell Spatial and Functional Cooperativity to Optimize

Priming. Immunity. 2017;46(2):205-19. Epub 2017/02/14. doi: 10.1016/j.immuni.2017.01.003.

PubMed PMID: 28190711; PubMed Central PMCID: PMCPMC5362251.

55. Alexandre YO, Ghilas S, Sanchez C, Le Bon A, Crozat K, Dalod M. XCR1+ dendritic cells

promote memory CD8+ T cell recall upon secondary infections with Listeria monocytogenes or

certain viruses. J Exp Med. 2016;213(1):75-92. Epub 2015/12/24. doi: 10.1084/jem.20142350.

PubMed PMID: 26694969; PubMed Central PMCID: PMCPMC4710197.

56. Domenjo-Vila E, Casella V, Iwabuchi R, Fossum E, Pedragosa M, Castellvi Q, et al.

XCR1+ DCs are critical for T cell-mediated immunotherapy of chronic viral infections. Cell Rep.

2023;42(2):112123. Epub 2023/02/17. doi: 10.1016/j.celrep.2023.112123. PubMed PMID:

36795562.

57. Deczkowska A, David E, Ramadori P, Pfister D, Safran M, Li B, et al. XCR1(+) type 1

conventional dendritic cells drive liver pathology in non-alcoholic steatohepatitis. Nat Med.

2021;27(6):1043-54. Epub 2021/05/22. doi: 10.1038/s41591-021-01344-3. PubMed PMID:

34017133.

58. Mysore V, Tahir S, Furuhashi K, Arora J, Rosetti F, Cullere X, et al. Monocytes transition

to macrophages within the inflamed vasculature via monocyte CCR2 and endothelial TNFR2.

J Exp Med. 2022;219(5). Epub 2022/04/12. doi: 10.1084/jem.20210562. PubMed PMID:

35404389; PubMed Central PMCID: PMCPMC9006314.

59. Liu Z, Liao F, Zhu J, Zhou D, Heo GS, Leuhmann HP, et al. Reprogramming alveolar

macrophage responses to TGF-beta reveals CCR2+ monocyte activity that promotes

bronchiolitis obliterans syndrome. J Clin Invest. 2022;132(19). Epub 2022/10/04. doi:

10.1172/JCI159229. PubMed PMID: 36189800; PubMed Central PMCID: PMCPMC9525120.

60. Kim D, An L, Moon J, Maymi VI, McGurk AI, Rudd BD, et al. Ccr2+ Monocyte-Derived

Macrophages Influence Trajectories of Acquired Therapy Resistance in Braf-Mutant

Melanoma. Cancer Res. 2023;83(14):2328-44. Epub 2023/05/17. doi:

10.1158/0008-5472.CAN-22-2841. PubMed PMID: 37195124; PubMed Central PMCID:

PMCPMC10478295.

61. Inui H, Nishida M, Ichii M, Nakaoka H, Asaji M, Ide S, et al. XCR1(+) conventional

dendritic cell-induced CD4(+) T helper 1 cell activation exacerbates cardiac remodeling after

ischemic myocardial injury. J Mol Cell Cardiol. 2023;176:68-83. Epub 2023/02/06. doi:

10.1016/j.yjmcc.2023.01.011. PubMed PMID: 36739942.

62. Moore KJ, Tabas I. Macrophages in the pathogenesis of atherosclerosis. Cell.

2011;145(3):341-55. Epub 2011/05/03. doi: 10.1016/j.cell.2011.04.005. PubMed PMID:

21529710; PubMed Central PMCID: PMCPMC3111065.

63. Yoshida T, Imai T, Kakizaki M, Nishimura M, Takagi S, Yoshie O. Identification of single C

motif-1/lymphotactin receptor XCR1. The Journal of biological chemistry.

1998;273(26):16551-4. doi: 10.1074/jbc.273.26.16551. PubMed PMID: 9632725.

64. Chu VT, Weber T, Graf R, Sommermann T, Petsch K, Sack U, et al. Efficient generation

of Rosa26 knock-in mice using CRISPR/Cas9 in C57BL/6 zygotes. BMC biotechnology.

2016;16:4. doi: 10.1186/s12896-016-0234-4. PubMed PMID: 26772810; PubMed Central

PMCID: PMC4715285.

65. Larson MA. Pronuclear Microinjection of One-Cell Embryos. Methods in molecular biology.

2020;2066:27-33. doi: 10.1007/978-1-4939-9837-1_2. PubMed PMID: 31512204.

66. Voisinne G, Kersse K, Chaoui K, Lu L, Chaix J, Zhang L, et al. Quantitative interactomics

in primary T cells unveils TCR signal diversification extent and dynamics. Nat Immunol.

2019;20(11):1530-41. Epub 2019/10/09. doi: 10.1038/s41590-019-0489-8. PubMed PMID:

31591574; PubMed Central PMCID: PMCPMC6859066.

67. Huang R, Guo G, Lu L, Fu R, Luo J, Liu Z, et al. The three members of the Vav family

proteins form complexes that concur to foam cell formation and atherosclerosis. Journal of

lipid research. 2019;60(12):2006-19. doi: 10.1194/jlr.M094771. PubMed PMID: 31570505;

PubMed Central PMCID: PMC6889716.

Human femoral artery

A Normal area

lumen

Plaque area

lumen

B

Xcr1+ area
(% total lesion area)

***

Normal Plaque

Mouse aortic root

C DAPI/Xcr1/CD11c

Chow diet

lumen

HFD

lumen

D

Xcr1+CD11c+ cell
(% total cells in lesion/16 weeks)

***

Chow diet HFD

E Chow diet (8 weeks)

lumen

HFD (8 weeks)

lumen

HFD (12 weeks)

lumen

HFD (16 weeks)

lumen

F

Xcr1+ area
(% total lesion area)

****

***

*

8   8   12   16   weeks
Chow diet   HFD

Wild-type allele

Chr9

5' UTR

Exon 2

3' UTR

5' UTR

Xcr1 Targeting vector

Cre   P2A   EGFP   P2A

5' HA
0.62 kb

3' HA
0.67 kb

HDR

Targeted allele

Chr9

5' UTR

Cre   P2A   EGFP   P2A   Exon 2

3' UTR

Single-cell RNA sequencing process

| Sample Collection | Cell Labeling | Cell Sorting | Single Cell Capture | Library Preparation | Sequencing |

Sample Collection

ApoE$^{-/-}$ Chow Diet (n=4) — Spleen, Aorta

ApoE$^{-/-}$ High-fat Diet (n=4) — Spleen, Lymph nodes, Aorta

Cell Labeling
Mouse Immune Single-Cell Multiplexing Kit (5 Sample Tags)

Fluorescent Antibody mix

Cell Sorting
Xcr1$^+$ cells sorted (5 Samples from all tissues)

Single Cell Capture
BD Rhapsody™ Single-Cell Analysis System

Library Preparation
BD Rhapsody™ Mouse Immune Response Panel

Sequencing
Next Generation Sequencing Novaseq 6000 (Illumina)

| Analysis pepline | Data analysis using seqGeq™ |

Analysis pepline

R1 FASTQ, R2 FASTQ, Reference

Read quality filtering

Identify cell ID and molecular ID

Align to reference (mRNA)

Raw read counts per bioproduce per cell

Molecule counts per bioproduct per cell error correction

Identify putative cells

Identify per cell sample of origin (sample multiplexing)

Output Files:
- Molecule and read counts per cell
- Sample tag calls
- Metrics
- BAM alignment

Quality Control
nFeature_RNA   nCount_RNA   perent_mito

Differentially expressed genes

Seurat clustering