1 **Pre-dauer starvation decouples somatic from germline development with lifelong**
2 **reproductive consequences in *C. elegans***
3

4 Fred A. Koitz[1], Camille P. Miller[1], Kacy Lynn Gordon[1,2]
5

6 [1]Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, North
7 Carolina 27599
8 [2]To whom correspondence should be addressed, kacy.gordon@unc.edu

9
10 **Running title:** Starvation and reproduction
11
12 **Keywords:** *C. elegans,* dauer, starvation, LAG-2, germline, germline stem cell niche
13

14
15
ABSTRACT

Early life stresses impact reproductive outcomes in many organisms. In response to
crowding and starvation, *C. elegans* nematodes form dauer larvae, in which
development arrests until conditions improve. We discovered dramatic differences in
gonad size and germ cell number among dauers that form under different conditions.
We used live cell imaging of fluorescent markers in otherwise wild-type and mutant
animals combined with food-removal, recovery, and brood-size assays to investigate
the causes and consequences of this germline difference. Pre-dauer feeding, but not
nutrient sensing via the DAF-2/insulin-like signaling receptor or DAF-7/TGF-β, is
required for plasticity in gonad size. Gonad differences in dauer have lifelong
reproductive consequences; worms with small dauer gonads recover to have smaller
broods. Somatic and germline development are decoupled as pre-dauer starvation
induces germline quiescence while the soma continues its development until dauer is
formed. A rapid return to germline Notch-dependence and an increase in presentation
by the germline stem cell niche of the Notch ligand LAG-2–regulated at the protein and
not transcript level–are among the earliest events of dauer recovery.

32
33
34 **SUMMARY STATEMENT**

35 Pre-dauer starvation limits the size of the dauer gonad, reduces presentation of the
36 germline niche cue LAG-2, and diminishes future reproductive success.

37
38
39

INTRODUCTION

Lifelong development requires environmental responsiveness within the developmental program. A special case of developmental response to difficult environmental conditions is the diapause state, in which development arrests in unfavorable conditions and resumes when conditions improve (Harsimran Kaur Gill et al., 2017; Renfree and Fenelon, 2017). While diapause states can be obligate - that is, built into an animal's life cycle in anticipation of poor conditions - other diapause states are facultative and responsive to an organism's present environment (Hand et al., 2016; Wilsterman et al., 2021). One well-studied facultative diapause state is the *C. elegans* dauer (Cassada and Russell, 1975), formed in response to starvation and crowding sensed by pheromone (Golden and Riddle, 1984a) and enhanced by higher temperatures (Ailion and Thomas, 2000). Dauer larvae withstand poor environmental conditions (Cassada and Russell, 1975; Golden and Riddle, 1984a) for many times the typical lifespan of a worm (Klass and Hirsh, 1976). Improved environmental conditions trigger exit from dauer, resumption of development, and a (mostly) normal reproductive life (Cassada and Russell, 1975; Klass and Hirsh, 1976). Famously, time spent in the dauer larval stage does not affect adult lifespan (Klass and Hirsh, 1976).

It is the preservation of fertility rather than increased lifespan alone that makes dauer adaptive (Schluter et al., 1991). Previous studies disagree about the long-term effects of dauer for future reproduction (Kim and Paik, 2008; Klass and Hirsh, 1976; Ow et al., 2018; Webster et al., 2018). These studies focused on recovery and did not investigate differences of the cells of the dauer reproductive system itself. We set out to investigate the cells of the gonad and germline during the dauer larval period to look for developmental antecedents of later reproductive differences.

To introduce our study, we present brief summaries of the parallel processes of the dauer decision and early reproductive system development. The *C. elegans* life cycle goes from embryonic development through four larval stages (L1-L4) punctuated by molting before a terminal molt into the adult reproductive form (Corsi et al., 2015). The dauer is an alternate third larval stage. Entry into the dauer developmental pathway is specified in the first larval stage (L1) and is followed by a second stage called L2d (Golden and Riddle, 1984a). In persistently harsh conditions, the L2d molts into a dauer larva with a thick cuticle, arrested development, and altered metabolism (Cassada and Russell, 1975; Erkut and Kurzchalia, 2015; Golden and Riddle, 1984a; Wadsworth and Riddle, 1989).

The *C. elegans* hermaphrodite reproductive system develops post-embryonically from a

4-cell primordium of two somatic blast cells and two primordial germ cells (Kimble and

Hirsh, 1979). In the L1 stage, a germline stem cell niche, the distal tip cell (DTC) is born

from each of the somatic blast cells and the primordial germ cells make a first division or

two in the L1 larval stage (Kimble and Hirsh, 1979). The germline relies on DTC

expression of the stemness cue LAG-2, a Delta/Serrate/LAG-2 (DSL) ligand of the

Notch signaling pathway to maintain germ cells in an undifferentiated state throughout

development (Kimble and Crittenden, 2005). By the end of the L2 stage a population of

20-30 germ cells (Kimble and White, 1981) is supported by two DTCs and a 10-cell

somatic gonad primordium (Kimble and Hirsh, 1979). The developmental arrest of

gonad and germ cell lineages in the dauer larva have been investigated genetically

(Narbonne and Roy, 2006; Tenen and Greenwald, 2019), primarily using genetic

mutants that constitutively form dauers at high temperature.

Among these are mutations to genes in the insulin-IGF-1 signaling (IIS) pathway,

including the gene encoding the sole *C. elegans* IIS receptor *daf-2* (Kimura et al., 1997).

IIS integrates nutritional status with metabolism, developmental control, reproduction,

and aging in *C. elegans* (Murphy and Hu, 2013) and in other organisms, including

humans (Barbieri et al., 2003; Claeys et al., 2002; Das and Arur, 2017). A reduction-of-

function allele, *daf-2(e1370),* is commonly used to induce dauer larvae at high

temperature in experimental settings (Karp, 2018). Another reduction-of-function allele,

daf-7(e1372), affecting a parallel pathway leading to dauer entry–TGF-β signaling–also

forms dauers constitutively at high temperature (Karp, 2018). The TGF-β pathway links

external cues to the reproductive system (Dalfó et al., 2012; Pekar et al., 2017). The

germline is especially sensitive to starvation throughout life (Angelo and Van Gilst,

2009; Dalfó et al., 2012; Pekar et al., 2017; Seidel and Kimble, 2011; Webster et al.,

2022), so we hypothesized that pre-dauer nutrition may cause differences in the dauer

reproductive system that impact post-dauer recovery of fertility.

We discovered that *C. elegans* enter dauer with different numbers of germline cells

depending on feeding prior to dauer entry. The most severely starved worms have

many times fewer cells and roughly half the brood size upon recovery compared to

those that fed the most during pre-dauer development. Surprisingly, the DAF-2/IIS

receptor is dispensable for this difference, suggesting that the fed state itself, rather

than nutrient sensing via IIS, regulates pre-dauer germline growth. Because the somatic

lineage is invariant in *C. elegans* (Kimble and Hirsh, 1979; Sulston and Horvitz, 1977;

Sulston et al., 1983)*,* early life starvation decouples germline from somatic

development, pausing germline development while the soma continues to the "safe

harbor" state of dauer diapause. The mechanism of this decoupling involves reduced
presentation of the stemness cue LAG-2 by the germline stem cell niche and the
induction of a Notch-independent quiescent state of the germline. This state is rapidly
reversed in the DTC and germline upon removal from dauer-forming conditions, and
appears to be regulated post-translationally. Altogether, this work reports heretofore
unrecognized differences in dauer gonads mediated by pre-dauer nutrient stress effects
on niche and stem cell maintenance that predict differential reproductive recovery from
dauer.
120
121   **RESULTS**
122   **Dauers that form constitutively have longer gonads than those that form after**
123   **progressive crowding and starvation**
124
We examined the gonad in dauer animals under two dauer-induction regimes that are
commonly used in the field–facultative dauer entry after starvation and crowding and
constitutive dauer forming mutants (Karp, 2018). First, we progressively starved
crowded worm growth plates (see Methods) and assayed both "early dauers" on such a
plate within the first week of starvation, as well as "late dauers" that were recovered
after >4 weeks. We used otherwise wild-type worms expressing markers of the germline
(*mex-5p::H2B::mCherry::nos-2 3'UTR*, (Linden et al., 2017)) and an endogenously
tagged integrin alpha subunit *ina-1(qy23[ina-1::mNG])*(Jayadev et al., 2019)) that aids
visualization of the somatic gonad.
134
We also used the field-standard dauer-formation constitutive (daf-c) mutants carrying
reduction-of-function alleles of *daf-2(e1370)* and *daf-7(e1372)*, which enter dauer
constitutively at high temperature (Karp, 2018). It is thought that DAF-2/IGFR (insulin-
like growth factor receptor) or DAF-7/TGF-β function is impaired, respectively, at all
rearing temperatures, as non-dauer-related loss-of-function phenotypes are observed at
both low and high temperatures (Blackwell et al., 2015; Gems et al., 1998; Greer et al.,
2008; Kenyon et al., 1993; McGehee, 2019; Trent et al., 1983; Tullet et al., 2008).
Indeed, dauer itself is triggered by high-temperature rearing in wild-type worms (Ailion
and Thomas, 2000), so these daf-c mutants are considered to be reduction-of-function
alleles that are sensitized for dauer formation at a higher temperature (Golden and
Riddle, 1984b).
146
This experimental regimen allowed us to compare between early and late dauers,
among genotypes after progressive starvation (control with gonad fluorescent markers
compared to *daf-2(e1370)* and *daf-7(e1372)*), and between modes of dauer induction
for both mutant genotypes (progressive starvation vs. constitutive entry, for both early
and late dauers).

Under progressive starvation, early dauers of all three genotypes had longer gonads
than late dauers, by nearly 2x on average (Fig. 1A,B). This raised the possibility that
dauer gonads change over time. We tested this hypothesis by using the dauer
constitutive mutants to form a population of constitutive, synchronous dauers and
measuring both early and late dauers for gonad size. Neither mutant showed a time-
related change of gonad length over a month in dauer (Fig. 1C,D), suggesting that time
spent in dauer was not responsible for the smaller gonads seen in control late dauers.
We therefore concluded that neither genotype (*daf-2(e1370)* or *daf-7(e1372)* vs. control)
nor time in dauer (over one month) is responsible for the gonad size difference between
early and late dauers formed by progressive starvation.

Worms with longer gonads have better post-dauer reproductive recovery than
worms with shorter gonads, regardless of genotype

We next asked if these same dauer-forming conditions for the same genotypes
considered above had functional consequences for post-dauer reproductive success.
We recovered dauers (Methods) and tracked individual worm recovery and eventual
brood size (Fig. 1E). For each genotype, we included a never-dauer, continuously-fed
control, since brood size can vary across strains. Otherwise wild-type worms with
fluorescent markers (see next section) show a normal brood size (of ~244 ± 31) in
never-dauer controls. Early dauers induced by progressive starvation (which have long
gonads, Fig. 1A) recover to produce a normal brood (271 ± 47) (Fig. 1F Early). In
contrast, late dauers (which have small gonads, Fig. 1A) recover to produce a
substantially decreased brood size compared to the early dauers (171 ± 112) (Fig. 1F
Late). The same pattern is observed for the *daf-2(e1370)* mutant genotype after
progressive starvation: early dauers (long gonads, Fig. 1A,B) produce normal broods
relative to the never-dauer control, while late dauers (short gonads, Fig. 1C,D) have
worse reproductive recovery (Fig. 1G).

When mutant *daf-2(e1370)* animals with the germ cell nuclear marker constitutively
entered dauer (larger germlines, Fig. 1C,D) both early and late dauers recovered
equally well, both producing broods equivalent to the never-dauer control of that
genotype (Fig.1G). Therefore, time spent in dauer alone did not reduce the brood of
subsequently recovered dauer animals. From these brood size assays, we concluded
that dauer gonad size was best correlated with post-dauer reproductive success. We
next asked what determines dauer gonad size.

Two of our prior experiments and one published finding (Cassada and Russell, 1975)
implicate pre-dauer feeding as a potential determinant of gonad size in dauer. First,

when we used daf-c alleles to induce constitutive dauer formation at high temperature,
the worms feed continuously until right before dauer entry and had uniformly large
gonads and brood. Second, for all genotypes exposed to progressive starvation, early
dauers from newly starved plates had large gonads and broods, and may have had
access to more food during pre-dauer development than late dauers. Finally, dauer
body length differs proportionally with the titre of bacterial food that worms ate prior to
dauer (Cassada and Russell, 1975). We hypothesized that pre-dauer feeding
determines gonad size in dauer.
200
201   **Starvation in the second larval stage decouples germline development from**
202   **somatic development independent of insulin-like signaling**
203
To test the hypothesis that feeding is responsible for dauer gonad length, we
manipulated *daf-2(e1370)* daf-c mutants expressing a germ cell nuclear marker (*mex-*
5p::H2B::mCherry::nos-2 3'UTR, (Linden et al., 2017)) in a series of food removal
experiments. The use of a daf-c mutant at 25°C ensured that we could vary food
exposure while ensuring every animal was in the L2d dauer entry program. Worms were
reared on growth plates with a lawn of bacterial food for the specified number of hours
at 25°C (Fig. 2A), removed from food at 25°C, and maintained until the endpoint at
which a population of majority dauers was noted.
212
213   We assayed germ cell number directly upon removal from food and again within a day
214   of dauer entry to test the relationship between time spent feeding and germ cell number,
215   and how much germ cell number changed between the cessation of feeding and entry
216   into dauer (Fig. 2B). We found a proportional relationship between time spent feeding
217   and germ cell number at the time of food removal and in dauer.
218
Between removal from food at the 12, 16, and 20 hours fed timepoints, it appears that
cells undergo one additional round of division before dauer entry (average germ cell
numbers approximately double, Fig. 2B,C). After 24 hours or longer of L2d feeding, the
increase in cell number before dauer entry is less dramatic, likely reflecting that late L2d
animals begin to slow germ cell division in preparation for dauer entry ((Narbonne and
Roy, 2006), also in *daf-2(e1370)* mutants). Germ cells will complete an average of one
additional cell division after DTC laser ablation in the L2-L3 stages of development
(Kimble and White, 1981), and genetic ablation of the stemness receptor GLP-1/Notch
allows germ cells to complete their current mitotic cell cycle and make a terminal
division before differentiating (Fox and Schedl, 2015). Our finding that germ cells
complete one round of division after food removal (Fig. 2B,D) tracks with our
established understanding of cell cycle constraints on stem-like germ cell proliferative
dynamics.

Comparing the rate of germline growth among these treatments revealed an even more
dramatic response to starvation than that of germ cell number alone (Fig. 2D). Animals
that fed (12h fed+24h fed) gained an average of about one germ cell per hour, while
animals that starved (12h fed+24h starved) added only one germ cell in that 24 hour
period (Fig. 2D, gray shaded box). In response to starvation, animals are able to arrest
germline development while slowing but not arresting somatic development until the
dauer larva is formed (Fig. 2E and (Cassada and Russell, 1975)), demonstrating a
decoupling of germline development from somatic development during the L2d period.
Because these experiments were done in *daf-2(e1370)* mutants, another major
conclusion from this experiment is that DAF-2/IGFR-mediated nutrient sensing itself is
not required for worms to modulate gonad size in response to food removal.

Dauer gonad size differences are driven by germline cell number and somatic cell
size

In the otherwise wild-type strain expressing *ina-1::mNG* and the germ cell nuclear
marker, early dauers had not only longer gonads, but more germ cells than late dauers
(Fig. 3A,B), demonstrating that the trend we observe in *daf-2(e1370)* mutants (Fig.2B,D)
holds. Gonad length is highly correlated to germ cell number (Fig. 3C). Late dauer germ
cells had slightly larger nuclei (Fig. 3D), and all dauer germ cell nuclei had a compacted
chromatin appearance (Fig. 3A, inset) characteristic of germ cell quiescence that has
been observed in starvation-induced L1 arrest (Belew et al., 2021).

We next asked if germ cell number was the only factor contributing to gonad size
differences. After progressive starvation, early wild-type (N2) dauers (Fig. 3E) displayed
greater gonad length (~1.7x) (Fig. 3F) and body width (~1.3x) (Fig. 3G) than late
dauers. However, the ratio of gonad length to body width also differs, with the early
dauer gonads being proportionally larger compared to the body width (~4.5:1) than
those of late dauer gonads (~3.5:1) (Fig. 3H), violating an assumption of simple
isometric scaling between body and gonad size. This suggests that growth is regulated
differently in the gonad vs. the rest of the body.

Unlike the germline, in which cell divisions vary, the somatic gonad cell lineage is
invariant (Kimble and Hirsh, 1979; Sulston and Horvitz, 1977) (Fig. 3I). When we
examine a somatic gonad nuclear marker (Shaffer and Greenwald, 2022), early and late
dauers have the same number of somatic gonad cells (two DTCs, four sheath-
spermathecal (SS) cells, and six uterine blast cells Fig. 3J). We further quantified the
size of the somatic gonad as the distance between these nuclei. Early dauers have

longer individual gonad arms (DTC-SS) than late dauers in addition to an overall greater
gonad length (DTC-DTC) (Fig. 3K). The SS-SS measurements represent the length of
the bulk of the somatic gonad blast cells found between the two gonad arms that will
give rise to the somatic gonad sheath, spermatheca, and uterus; early dauers were also
larger than late dauers by this measure (Fig. 3K). Therefore, late dauers have a more
compact somatic gonad than early dauers.

The only somatic gonad cell to be terminally differentiated at this stage of development
is the DTC, which is the germline stem cell niche that maintains germ cells in an
undifferentiated, stem-like state (Crittenden et al., 2006; Hubbard, 2007; Kimble and
White, 1981; Kimble, 2014). Late dauers had smaller DTCs than early dauers, though
both had markedly smaller DTC cell size than a continuously-fed L2 control (Fig. 3L,M).
Thus, we find two separate mechanisms responsible for the smaller size of the gonad of
late dauers, corresponding to their respective cell lineage limitations: while the germline
is reduced in cell number, the somatic gonad appears to be more compact with the
same number of cells, and the DTCis smaller in cross-section. Because the DTC is
required for germ cell proliferation, we next examined the role of this cell in supporting
the dauer germline.

The dauer germline is maintained in a Notch-independent, quiescent state

One necessary factor regulating germline growth is LAG-2 ligand expressed by the DTC
activating GLP-1/Notch signaling in the distal germline (Henderson et al., 1994). Under
replete conditions, niche signaling via LAG-2 to the GLP-1/Notch receptor in the germ
cells is required for their maintenance in a mitotic, undifferentiated state at all
developmental stages; both DTC ablation (Kimble and White, 1981) and genetic
ablation of Notch signaling (Fox and Schedl, 2015; Kodoyianni et al., 1992) are
sufficient to trigger meiosis within hours. However, starvation has been shown to induce
a G2-arrested, Notch-independent quiescent state in transiently starved adults (Seidel
and Kimble, 2011). We found that primordial germ cells in L1 arrested animals, which
are known to be G2-arrested and quiescent (Belew et al., 2021), are also Notch-
independent (n=15/15 recovered *glp-1(bn18)* L1 arrested animals were fertile after 24h
at 25°C).

We hypothesized that the germline of both early and late dauers, which are in a *de facto*
starved state and known to be G2-arrested (Narbonne and Roy, 2006), is similarly
Notch-independent. We investigated this question using worms carrying a temperature
sensitive allele of *glp-1*/Notch*, glp-1(bn18),* which at the permissive temperature (16°C)
have adequate Notch signaling to remain fertile, but upon shifting to the restrictive
temperature (25°C) early in development become sterile due to failure of germline

induction or irreversible meiotic entry of germ cells (Fox and Schedl, 2015; Kodoyianni
et al., 1992). In adults, signs of meiotic entry appear in distal adult germ cells within 6
hrs (Fox and Schedl, 2015; Kodoyianni et al., 1992).
314
When continuously-fed populations of *glp-1(bn18)* mutants were shifted to the restrictive
temperature for 24 hours (Fig. 4A, and see Methods), nearly all animals singled after
the treatment were sterile when shifted back to the permissive temperature (Fig. 4B), as
expected. In comparison, when *glp-1(bn18)* dauers–both early and late–were shifted to
the restrictive temperature for 24 hours, all recovered as fertile adults at the permissive
temperature (Fig. 4B). From this, we conclude that both the early and late dauer
germline is in a Notch-independent, quiescent state.
322
323   **Starvation reduces presentation of the stemness cue by the germline stem cell**
324   **niche prior to dauer entry**
325
It is not known whether or how LAG-2 protein abundance or localization changes under
conditions that induce a Notch-independent, quiescent germline. We used a strain co-
expressing an endogenously tagged LAG-2::mNeonGreen (Gordon et al., 2019) and
DTC membrane marker (*lag-2p::myrTdTomato,* (Byrd et al., 2014)) to quantify LAG-2
abundance in an L2 control compared to both early and late dauers. Both early and late
dauers had severely diminished (~10x) LAG-2::mNG signal in the DTC compared to the
L2 control but did not significantly differ from one another (Fig. 4C-F). The
transcriptional reporter is less active in both dauers compared to the L2, but not by the
same magnitude as the LAG-2::mNG (Fig. 4B and 4G). While dauer DTCs produce less
lag-2 gene product than fed L2s, the absolute amount of LAG-2 signaling protein in
early vs. late dauer cannot explain the different germline sizes of these groups.
337
We next turned to the dynamics with which LAG-2 expression is diminished during the
pre-dauer period, the time during which feeding changes germline growth (shown in Fig.
2B,C). We combined the LAG-2::mNG endogenous protein tag and the *lag-*
2p::myrTdTomato transcriptional reporter with the *daf-2(e1370)* allele and performed a
timed food removal assay on L2d daf-c animals at 25°C (Fig. 4H, as in Fig. 2). We
found that LAG-2::mNG protein abundance is highly responsive to the onset of
starvation in L2d. LAG-2::mNG drops from the 20h fed baseline within 5 hours of food
removal (Fig 4I-L). Transcription of *lag-2* is known to be lower in late larval animals in
unfavorable conditions (low food, high pheromone) that cause reduced DAF-7 signaling
(Dalfó et al., 2012; Pekar et al., 2017), and we find that animals in L2d have
dramatically less transcription from this reporter than control L2s (Fig. 4M). However,
the transcriptional reporter stays fairly steady upon food removal in L2d (Fig. 4M). Thus,
reduced *lag-2* transcription cannot be the entire mechanism behind the drop in LAG-

2::mNG abundance upon food removal in L2d; diminished LAG-2::mNG protein stability
(at least in comparison to the myristoylated TdTomato reporter) must also play a role.
353
We conclude that dauer germline is maintained in a Notch-independent state with low
levels of the stemness cue LAG-2 expressed by the DTC. The amount of LAG-2 being
presented to the germ cells does not differ between early and late dauers, however,
suggesting that niche limitations in dauer do not regulate the different gonad sizes
between these two cohorts. Instead, the timing at which LAG-2 drops is acutely driven
(within 5 hours) by the termination of pre-dauer feeding. We next examined the
dynamics of Notch signaling during dauer exit.
361
362 **Post-dauer Notch-dependence is restored in tandem with a transcription-**
363 **independent burst of LAG-2 ligand presentation by the germline stem cell niche**
364
We hypothesized that the post-dauer germline would rapidly return to a Notch-
dependent state upon germ cell-cycle re-entry during dauer recovery, as proliferating *C.*
elegans germ cells are Notch-dependent in other life stages (Austin and Kimble, 1987;
Cinquin et al., 2010). We predicted that LAG-2 protein abundance would rapidly
increase to meet a post-dauer requirement for Notch signaling. We measured
expression in an otherwise wild-type strain of the LAG-2::mNG protein and a *lag-2-*
promoter-driven transcriptional reporter upon dauer recovery (see Methods).
372
At 0 hours of recovery, late dauers had detectable signal of both the LAG-2::mNG
protein and the *lag-2* transcriptional reporter (Fig. 5A-D), suggesting that the dauer DTC
continues to express the *lag-2* locus and localize LAG-2 protein to the membrane.
Dauer expression of *lag-2* reporters in the DTC has been reported previously in *daf-*
2(e1370) mutant dauers (Narbonne and Roy, 2006), and here we show that this
expression endures through several weeks of dauer. During the first two hours of dauer
recovery, LAG-2::mNG protein abundance increases 2-3-fold (Fig. 5A). Surprisingly, we
did not observe a corresponding increase in signal from the co-expressed *lag-*
2p::myrTdTomato transcriptional reporter (Fig. 5B). Such dynamics suggest that the
immediate restoration of LAG-2 on the DTC–much like its initial downregulation upon
starvation (Fig. 4I-M)–is post-transcriptionally regulated.
384
Before drawing this conclusion, we needed to rule out experimental artifacts that could
complicate the analysis of the transcriptional reporter. We considered that slow folding
of TdTomato fluorescent protein could cause a lag in its transcriptional readout (see
Methods), or that post-transcriptional regulation of *lag-2* mRNA could uniquely affect
LAG-2 production. We therefore examined the expression of another pair of DTC-
expressed markers: a faster-folding, membrane-localized *lag-*

2p::mNeonGreen::PLCδ1^PH (*cpIs122,* (Linden et al., 2017)) and a histone
H2B::mTurquoise2 knocked into the endogenous *lag-2* locus with a 2A peptide
cleavage site (Medwig-Kinney et al., 2022). This second element will generate a
polycistronic mRNA, the translation of which will produce one LAG-2 protein and one
H2B::mTurquoise2, meaning that the histone reporter is under the same transcriptional
and post-transcriptional, transcript-based regulation (e.g. by 3' UTR-mediated
repression by microRNAs or RNA binding proteins, transcript decay, etc.) as the
endogenous *lag-2* gene. In the two hours after recovery from dauer, we observe no
change in the levels of expression of either the transcriptional membrane (Fig. 5E,G) or
polycistronic nuclear reporter (Fig. 5F,H). We conclude that the *lag-2* locus is
transcriptionally active in the DTC during dauer–despite the germline being maintained
in a Notch-independent state–and LAG-2 is under protein-specific negative regulation
that is rapidly alleviated upon dauer exit.
404
Given the rapid return of LAG-2 to the DTC surface after dauer exit, we predicted that
the germline would become Notch-dependent in the first hours of recovery from dauer.
We performed temperature shift experiments during dauer recovery for *glp-1(bn18)*
mutant worms (Fig. 5I). Dauer *glp-1(bn18)* mutant animals all recover fertility after an
excursion to the restrictive temperature (Fig. 4B). If worms are kept at the restrictive
temperature for the first 4 hours of recovery from dauer, the stem-like germ cells remain
Notch-independent and the animals recover as fertile adults (Fig. 5J, top, n=39/39).
After 6 hours of recovery from dauer at 25°C, ~20% of worms become sterile in
adulthood (Fig. 5J, middle, n=7/34). After 8 hours of recovery from dauer at 25°C, more
than half of worms will recover as sterile adults (Fig. 5J, bottom, n=25/43). Taken
together we see a rebound of LAG-2 protein on the DTC within two hours of recovery
from dauer, preceding a return of Notch dependence in the germline between 4-8 hours
of recovery. Because dauer exit is asynchronous among individuals (Cassada and
Russell, 1975), the precise timing of these events with respect to dauer exit and to one
another may be even more tightly coordinated. Other early features of dauer exit–loss of
the buccal plug and resumption of pharyngeal pumping–also occur within a two-hour
window of removal from dauer conditions (Cassada and Russell, 1975), making the
return of LAG-2 on the DTC surface as rapid a response to the end of dauer as any
other measured event. This correlation suggests that systemic recovery is more
integrated across body systems than pre-dauer changes, which were able to be
uncoupled. Our findings establish germline regulation as a central, early feature of
dauer recovery.
427
DISCUSSION
Here we report that changes in the germline stem cell niche cue LAG-2 are among the
earliest responses both to starvation preceding dauer entry and to environmental cues

that allow dauer exit. Reduction of pre-dauer feeding reduces the amount of LAG-2
presented by the germline stem cell niche and the proliferation of germline cells.
Starved pre-dauer germ cells swiftly become quiescent and complete a single terminal
division before dauer entry. The response at the level of germ cell number is dramatic–
the most severely starved dauers (wild-type late dauers and *daf-2(e1370)* mutants
starved early in L2d) have as few cells in the germline (~5-8) as worms with severe
loss-of-function alleles of *glp-1*/Notch receptor gene (Austin and Kimble, 1987), and
even fewer germ cells than worms have after early laser ablation of the DTC (Kimble
and White, 1981).
440
441   While reduced food intake also slowed the rate of somatic progression towards the
442   L2d/dauer molt, it does not alter somatic cell numbers in the somatic gonad (Fig. 3).
443   Instead, the soma as a whole continues its development through L2d and into dauer
444   while the germline arrests, both in terms of absolute germ cell number and–even more
445   dramatically–rate of germline grown over time (Fig. 2).
446
Differences in germline and somatic response to the starvation-induced L1 larval arrest
stage have previously been observed, both in differential transcriptional responses
(Webster et al., 2022), and chromatin compaction in the germline (Belew et al., 2021;
Morao and Ercan, 2021). The L1 arrest state is triggered by complete starvation after
hatching (Baugh, 2013; Johnson et al., 1984). Several checkpoints triggered by later
starvation arrest somatic development at a later larval stage transition (Schindler et al.,
2014), and the somatic gonad relays environmental information to the germline
throughout life (Gordon, 2020). We observe that the soma and germline can decouple
their development plastically and rapidly in response to changing environmental
conditions in the middle of a larval stage.
457
Notch signaling via *lag-2* in the nervous system is required both during dauer and for
dauer recovery (Ouellet et al., 2008). LAG-2 protein downregulation in the DTC has
been observed during worm aging (Aprison et al., 2024 preprint; Singh et al., 2024), and
lag-2 transcription is downregulated via reduced TGF-β signaling after food removal in
late larval worms (Dalfó et al., 2012; Pekar et al., 2017). The transition of the germline
from a Notch-dependent to a Notch-independent state has been reported during
starvation (transient adult starvation (Seidel and Kimble, 2011) and L1 arrest (this
paper)), and it should be investigated whether DTC-expressed LAG-2 is also reduced in
those contexts. The question remains whether a reduction in LAG-2 signal in response
to starvation is the proximal cause of germline quiescence, or whether germline
quiescence is a necessary precondition for the germline to tolerate a reduction in LAG-2
as a byproduct of starvation.
470

Notably, expression of both endogenously tagged LAG-2 protein and *lag-2* promoter-
driven transgenes (Fig. 5 and (Narbonne and Roy, 2006)) remain detectable in dauer–
even after an an extended duration of dauer diapause–so the downregulation of *lag-2*
during starvation is relative and not absolute. The regulation of LAG-2 in the DTC in our
experiments also appears to entail post-translational downregulation upon starvation
and during dauer which rapidly ceases at dauer exit.
477
It has previously been shown that the TGF-β pathway regulates *lag-2* expression in the
DTC as part of environmental sensing, including as a response to dauer pheromone
and low food, though this work was done in late larvae that had bypassed the dauer
decision (Pekar et al., 2017). TGF-β signaling is restored during dauer exit (Ren et al.,
1996; Schackwitz et al., 1996), so we predict that *lag-2* transcription eventually rises
during recovery from dauer, but apparently not on the rapid time scale of the recovery of
LAG-2 protein in the DTC.
485
While the precise molecular mechanisms linking nutrition to LAG-2 protein expression in
the DTC remain to be found, we establish that they proceed even with the field-standard
daf-2(e1370) class 2 loss-of-function allele of the sole *C. elegans* IIS receptor.
Not only is starvation-triggered LAG-2 downregulation DAF-2-independent, but the
reproductive consequences of pre-dauer food intake are also the same in *daf-2(e1370)*
animals as they are in wild-type worms. Worms that fed before dauer (both early dauers
of both genotypes and mutant worms induced to form dauer at high temperature) have
normal brood sizes after recovery.
494
The germline is the only lineage in *C. elegans* with an indeterminate division pattern and
variable cell number. It is therefore the only cell population that can respond to
starvation by limiting the number of cells it produces. We have demonstrated that this
unique flexibility allows germline development to be decoupled from the rest of the
dauer entry program earlier than was previously thought. In genetic backgrounds in
which animals are able to feed before genetically-induced dauer entry, it has been
established that the germline and somatic gonad require AMPK (*lbk-1* and *aak-2*)
signaling and DAF-18/PTEN–independent of the DAF-2 IIS receptor–to enter a
quiescent state (Narbonne and Roy, 2006; Tenen and Greenwald, 2019). Prior to dauer
entry, the temperature-sensitive *glp-1* allele in *daf-2(e1370)* genetic background shifted
to the restrictive temperature causes germ cell pachytene arrest during dauer
(Narbonne and Roy, 2006). Since these *daf-2(e1370)* mutants feed before entering
dauer, this shift to the restrictive temperature occurred during what our work suggests is
a Notch-dependent phase of pre-dauer germline growth. We find that after starvation
prior to dauer entry, the germline undergoes a reduction in the stemness cue presented

by the somatic gonad (also independent of DAF-2) before becoming Notch-independent
in dauer.
512
Early life stresses are known to affect later life outcomes in a range of organisms and
contexts with potential mechanisms ranging from epigenetics to inflammation to
hormone signaling to lasting variation in central nervous system development and
function (Taylor, 2010). In *C. elegans,* there is precedent for differential reproductive
recovery after dauer depending on whether starvation or crowding (high levels of the
dauer pheromone) was the primary trigger to dauer entry (Ow et al., 2018). Our
progressive starvation protocol involves both starvation and crowding, and future work
will probe the contributions of both factors to pre-dauer germline quiescence.
521
Taken together, we propose that germline regulation is a key response to both the cues
that induce dauer and the withdrawal of those cues in recovery from dauer. Future work
will focus on how the germline transitions from a Notch-independent state back to
Notch-fueled germline proliferation. One possible driver is nucleotide levels, which have
been shown to limit cell division but not cell growth in unicellular organisms (Diehl et al.,
2022) and to affect germline growth specifically in non-dauer *C. elegans* (Chi et al.,
2016). The rapid restoration of LAG-2 protein to the DTC surface upon withdrawal of
dauer-maintenance conditions (Fig. 5) certainly suggests that reestablishing a stem cell
niche capable of supporting germline growth is an early and crucial feature of dauer
recovery. Our analyses of endogenously tagged LAG-2::mNeonGreen protein recovery
vs. transcriptional reporters and a polycistronic reporter altogether suggest that the
ligand's increase at the membrane is regulated post-translationally (Fig. 5). Notch
signaling is regulated at the protein level in multiple ways (Fortini, 2009; Suarez
Rodriguez et al., 2023; Tian et al., 2004; Weinmaster and Fischer, 2011; Wu et al.,
2016; Yamamoto et al., 2010), and future work will delve into this aspect of its regulation
upon starvation and during dauer exit. Tellingly, in our experiment, LAG-2 increases
upon recovery from dauer in the absence of food, suggesting that LAG-2 is actively
inhibited by the dauer pheromone rather than actively promoted by nutritional cues.
540
Such a mechanism of rapid ligand protein up-regulation by means of alleviated
repression could theoretically yield a faster response than upregulation that requires
new transcription initiation. However, continuous transcription, translation, and LAG-2
protein degradation comes with an energy cost that a dauer worm would only incur if the
alternative–slower restoration of LAG-2 protein on the stem cell niche surface–carried
an even greater cost of reduced fertility upon return of the germline to Notch-
dependence.
548
549

MATERIALS AND METHODS

Sections of this text are adapted from prior Gordon lab publications (Li et al., 2022; Singh et al., 2024), as they describe our standard laboratory practices and equipment.

Strains

We used Wormbase (Sternberg et al., 2024) while conducting this study. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440) and are to be requested directly from CGC. The following *C. elegans* strains were obtained from the CGC: N2 (Brenner, 1974), CB1370 *daf-2(e1370)* III (Kimura et al., 1997), CB1372 *daf-7(e1372)* III (Vowels and Thomas, 1992), DG2389 *glp-1(bn18)* III (Kodoyianni et al., 1992).

The following *C. elegans* strains were generously shared by community members or generated previously in our lab:
NK2517 *qIs154(lag-2p:: myr::tdTomato); lag-2(cp193[lag-2:: mNeonGreenˆ3xFlag])* V) (Gordon et al., 2019), KLG034 *cpIs122(lag-2p::mNeonGreen:: PLCδ1^{PH})*II*; lag-2(bmd202 [lag-2*::P2A::H2B::mTurquoise2^lox511I^2xHA]) V. (Singh et al., 2024), GS9692 *arTi435(rps-27p::2xnls::gfp(flexon)::unc-54 3' UTR* I); *arTi237(ckb-3p::Cre(opti)::tbb-2 3' UTR* X) (Shaffer and Greenwald, 2022).

The following *C. elegans* strains were generated for this paper by crossing alleles from sources that are cited after each strain description: KLG047 *ina-1(qy23[ina-1::mNeonGreen]);naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR)* II (Jayadev et al., 2019; Linden et al., 2017), KLG050 *daf-2(e1370)* III*; naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR)* II, KLG051 *daf-2(e1370)* III*; qIs154(lag-2p:: myr::tdTomato)* V*; lag-2(cp193[lag-2:: mNeonGreenˆ3xFlag])* V.

Strain maintenance and synchronization

Worm strains were maintained on nematode growth media (NGM) at 16°C unless otherwise specified. Populations were synchronized by bleaching according to a standard egg prep protocol (Stiernagle, 2006).

Dauer formation by progressive starvation and crowding

To more precisely time the formation of facultative dauers, a large, mixed population of well-fed worms was transferred from an uncrowded growth plate to NGM plates seeded with ~70µL of an overnight culture of *E. coli* OP50 bacterial food. Plates were sealed with parafilm and placed at 25°C for the non-temperature-sensitive strains and 16°C for strains containing *daf-2(e1370), daf-7(e1372),* or *glp-1(bn18)* temperature-sensitive alleles. Plates were monitored daily for evidence of starvation (bacterial lawn depleted and worms dispersed). At 25°C this took ~2-4 days and at 16°C this took ~5-7 days

The first day starvation was noted became day 0 of starvation. Early dauers were
defined as coming from plates that were 5 days or fewer starved, while late dauers were
defined as coming from plates that were at least 28 days starved.

Brood size assays

All brood size assays were conducted at 16°C to match the necessary conditions of the
temperature sensitive strains. After experimental treatment (see below), worms were
singly picked to NGM plates seeded with ~70µL *E. coli* OP50. Egg-laying adults were
subsequently passaged to a new plate once daily until egg laying stopped. Plates with
progeny were kept at 16°C until the oldest worm on the plate had reached L4, after
which they were counted. Schematic illustrating this method in Fig. 1E.

For never-dauer control brood assays, L4 animals of each strain were singled on Day 0.
For brood after recovery from constitutive dauer formation, healthy plates of adults were
egg prepped; eggs were rolled in M9 overnight at room temperature to obtain a
population of arrested L1 larvae. Animals were dropped onto NGM plates seeded with
OP50. Plates were sealed with parafilm and put at the restrictive temperature (25°C) to
induce dauer formation. Early dauers that were constitutively formed came from plates 5
days or fewer after dauer formation; late dauers came from plates at least 28 days after
dauer formation. For brood after recovery from progressive starvation, the above
protocol was followed to induce dauer. Worms were then isolated by SDS treatment
(see next section) and dauers were singly recovered to NGM plates seeded with OP50
at 16°C.

Dauer isolation protocol

For plates with mixed populations, dauers were isolated with 1% SDS (Karp, 2018) After
SDS treatment, the population of recovered animals was placed onto an unseeded
NGM plate and the excess liquid was allowed to dry, after which animals were picked to
a slide for imaging, were kept on the unseeded plate for the specified number of hours
to evaluate short-term recovery, or picked to an NGM plate seeded with OP50 for
recovery and brood measurement.

Food removal experiments

A synchronized population of *daf-2(e1370); naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR
II)* L1s was added to NGM plates seeded with OP50 and were put at 25°C to induce
constitutive dauer formation. After the specified number of hours (Fig. 2A), animals were
rinsed off the seeded plate with M9, recovered to a 15mL conical tube, and were
subsequently washed in M9 3-5 times and then rolled in M9 at room temperature for 20
minutes before being washed an additional time in M9 to clear bacteria. Washed
animals were then transferred to peptone-free plates (recipe from (Eustice et al., 2022))

and returned to 25°C to continue to develop. Baseline imaging was done immediately
after animals were washed off the food plates. Subsequent imaging done at the
specified times (Fig. 2B). For Fig. 2B, animals were subjected to feeding/starvation for
the indicated times and then kept at 25°C until the plate visibly had >50% dauers.
Animals that appeared to be dauers under a dissecting scope were picked to image for
the dauer endpoint of the experiment; only animals with dauer morphology were
included in the final dataset. While removal of food in the L1 stage prior to the L1-L2d
transition caused L1 developmental arrest, as expected, we found that removal of food
after the L2d molt did not inhibit *daf-2(e1370)* animals from progressing through L2d or
entering dauer.
638
***glp-1(ts)* temperature shift experiments**
Two temperature shift experiments used the temperature-sensitive allele of the Notch
receptor *glp-1(bn18)*. To test for *glp-1* dependence during dauer (Fig. 4 A,B), two
populations were used. As a control, an unstarved, uncrowded population was reared at
16°C, shifted to the restrictive temperature of 25°C for 24 hours, larval worms were
singly picked to NGM plates seeded with *E. coli* OP50, and these plates were kept at
16°C until adulthood. As two test groups, worms that had been subjected to progressive
starvation and crowding at 16°C (both early and late) were shifted to 25°C for 24 hours.
After 24 hours, dauers were recovered by SDS isolation and singled to NGM plates
seeded with *E. coli* OP50 and kept at 16°C until adulthood. Each plate was
subsequently assayed for the presence of progeny.
650
To test for when *glp-1* dependence returns after dauer, *glp-1(bn18)* worms were put into
dauer by progressive starvation at 16°C, shifted to 25°C overnight, and then SDS
isolated, transferred to NGM plates without food and placed back at 25°C for an
additional 4, 6, or 8 hours. At that time, the developmentally oldest/largest worms on the
plate (the worms that exited dauer first, since dauer exit is asynchronous) were picked
as single animals to NGM plates seeded with *E. coli* OP50 food and kept at 16°C until
adulthood. After 4 days of recovery at 16°C, each plate was assayed for live larvae, and
those that had offspring were scored as fertile. Plates that lacked larvae were imaged at
600x magnification in DIC and scored for the presence of embryos in the uterus (fertile)
or an absence of gametes (sterile). Embryo retention appeared to be a phenotype of
glp-1(bn18).
662
Confocal imaging
All images were acquired at room temperature on a Leica DMI8 with an xLIGHT V3
confocal spinning disk head (89 North) with a 63× Plan-Apochromat (1.4 NA) objective
and an ORCAFusion GenIII sCMOS camera (Hamamatsu Photonics) controlled by
microManager. RFPs were excited with a 555 nm laser; GFP and mNG were excited

with a 488 nm laser; mTurquoise was excited with a 445 nm laser; DAPI was excited
with a 405 nm laser. Z-stacks through the gonad were acquired with a z-step size of 0.3
or 0.5 µm as noted. Worms were mounted on agar pads in M9 buffer with 0.01 M
sodium azide (VWR (Avantor) Catalog Number 26628-22-8). Some samples were
acquired with a 1.6X optical zoom, as indicated in the figure legends. Samples used for
fluorescence quantification were acquired with the same laser power, exposure time,
and z-step size within the datasets to be compared.
675
676 **Image analysis**
677 Images were processed in FIJI89 (Version: 2.14.1/1.54f).
678
Gonad length and body width measurements
From DIC confocal z-stacks, gonad length was measured from the distal most end of
one gonad arm to the distal most end of the other gonad arm using the segmented line
tool in FIJI following the general shape of the gonad. Body width was measured from
cuticle edge to cuticle edge perpendicular to the gonad using the straight line tool in FIJI
across the z-slice that captured the widest part of the body.
685
Somatic gonad marker measurements
Somatic gonad cell number (Fig. 3J-K): The strain GS9692 expressing *arTi435(rps-*
27p::2xnls::gfp(flexon)::unc-54 3' UTR); *arTi237(ckb-3p::Cre(opti)::tbb-2 3' UTR)*
(Shaffer and Greenwald, 2022) was specifically designed to activate enduring GFP
expression in the cells of the somatic gonad. We examined this strain in early and late
dauers after inducing dauer with progressive starvation and crowding (see above). We
always see the expected two DTCs and 10 primordial gonad cells in dauers of any age.
Variation in expression levels/tissue depth can sometimes complicate detection of the
cell at the center of the gonad primordium (more orange cells in Fig. 3J). Landmarks
were selected based on relative position with DTCs at the gonad tips and the next-most-
distal fluorescent cells being the SS cells used for measuring.
697
DTC measurement (Fig. 3L-M): Strain NK2517 *qIs154(lag-2p:: myr::tdTomato); lag-*
2(cp193[lag-2:: mNeonGreenˆ3xFlag] V*)* was imaged for the DTC membrane marker
only at the stages specified in Fig. 3M. A sum projection through ten 0.3 µm z-steps
capturing the superficial surface of the DTC was made, traced by hand in FIJI, and the
area measured. We chose an L2 control for this experiment because, while the dauer
larva is an alternative third larval stage, the worm's reproductive system undergoes
morphogenetic changes and changes in cell number during the normal L3 stage. These
developmental events are arrested in dauer (Tenen and Greenwald, 2019), making L3 a
confounding control for the dauer larvae reproductive system.
707

Dauer recovery experiments

To measure gene expression during recovery, worms were isolated as described above (Dauer isolation protocol), and subsequently split such that some worms were imaged at 0 hours of recovery, and other worms were recovered to NGM plates without food (to isolate withdrawal of dauer maintenance conditions from food-specific cues) for the specified length of time.

Reporters of *lag-2* measurement: Endogenously tagged LAG-2::mNG with coexpressed membrane TdTomato

Strain NK2517 *qIs154(lag-2p:: myr::tdTomato); lag-2(cp193[lag-2:: mNeonGreenˆ3xFlag]) V* was used to measure endogenously tagged LAG-2::mNG protein and a coexpressed transcriptional reporter of the *lag-2* promoter. All worms, including controls, were reared at 25°C. TdTomato has a half-time to maturation of one hour at 37°C (Shaner et al., 2004), while mNeonGreen has a half-time to maturation of fewer than ten minutes at 37°C (Shaner et al., 2013) (both will fold slower at worm-rearing temperatures). TdTomato signal is visible in adjacent neurons (Fig. 4 and Fig. 5) and the autofluorescent gut granules of the worm are visible in the channel of the endogenously tagged LAG-2::mNG. These structures were avoided as much as possible during projection, tracing, and measurement of signal.

Confocal z-stacks were acquired with a 0.3 μm step size. A sum intensity z-projection was created in FIJI consisting of 10 slices representing the superficial half of the cell (from the superficial cell surface to the cross-section of the nucleus). Either one or both DTCs from each worm were analyzed depending on if the gut obscured one. The DTC was hand-traced on the membrane marker RFP channel; that ROI was subsequently used to measure both the RFP (membrane) and GFP (endogenously tagged protein) channels, and slid off the DTC to the adjacent body of the worm to measure the background for both channels. Background subtracted measurements were thus obtained for both Mean Gray Value (mean pixel value in the ROI) and Integrated Density (total pixel value in the region = MGV*area). Integrated density for all samples was normalized to the mean integrated density at the 0h timepoint, so fold-change at 2h time point is shown.

Reporters of *lag-2* measurement: Polycistronic histone reporter with coexpressed membrane mNeonGreen

We examined another pair of DTC-expressed markers: a faster-folding, membrane-localized *lag-2p::mNeonGreen::PLCδ1^{PH}* (*cpIs122,* (Linden et al., 2017)) and a histone H2B::mTurquoise2 (half-time to maturation of 33.5 minutes, (Goedhart et al., 2012)) that was knocked into the endogenous *lag-2* locus with a 2A peptide cleavage site (Medwig-Kinney et al., 2022). This second element will generate a polycistronic mRNA, the

translation of which will produce one LAG-2 protein and one H2B::mTurquoise2,
meaning that the histone reporter is under the same transcriptional and post-
transcriptional, transcript-based regulation (e.g. by 3' UTR-mediated repression by
microRNAs or RNA binding proteins, transcript decay, etc.) as the endogenous *lag-2*
gene. The membrane-localized mNeonGreen reporter, on the other hand, will simply
come on when the *lag-2* promoter (including the TGF-β-responsive element described
by (Pekar et al., 2017)) is active.
755
Confocal z-stacks were acquired with a 0.3 µm step size. A sum intensity z-projection
through a 6-7 µm depth of the DTC (to capture the entire nucleus in z) was generated in
FIJI, the DTC was hand-traced, and membrane fluorescence was measured for the
GFP channel in that ROI. Both DTCs from each animal were analyzed in separate
projections. A background measurement was made by sliding the DTC ROI onto the
adjacent body of the worm. The nuclear histone signal was obtained using an ellipse
tool tight around the nucleus in the same z-projection to measure on the CFP channel,
and background for the nuclear signal was made by sliding that ROI onto the adjacent
body of the worm. Background subtracted measurements were thus obtained for both
Mean Gray Value (mean pixel value in the ROI) and Integrated Density (total pixel value
in the region, MGV*area). Integrated density for all samples was normalized to the
mean integrated density at the 0h timepoint, so fold-change at 2h time point is shown.
768

Statistical analysis
All statistical analysis was done using GraphPad Prism version 10.4.2 for Windows,
GraphPad Software, Boston, Massachusetts USA, www.graphpad.com. Specific tests
specified in the relevant figure legends.
Statistical analysis of fold change shown in 5A, B, E, F; statistical analysis (unpaired t-
test) of non-normalized integrated densities for these datasets show the same pattern of
significance (that is, only LAG-2::mNG signal changes between 0h and 2h) as the fold
change analysis.
777

Author Contributions
779   Conceptualization: F.A.K. and K.L.G.; Formal analysis: F.A.K.; Investigation: F.A.K. and
780   C.P.M.; Writing – original draft preparation: F.A.K. and K.L.G.; Writing – review and
781   editing: F.A.K., C.P.M., and K.L.G.; Visualization: F.A.K.; Supervision: K.L.G.; Project
782   administration: K.L.G.; Funding acquisition: F.A.K. and K.L.G.
783

Acknowledgements
785   We thank members of the Gordon Lab and F.A.K. dissertation committee members
786   R.H.D. and D.R.M. for thoughtful comments on the manuscript.
787

Funding
Some strains were provided by the CGC, which is funded by National Institutes of
Health Office of Research Infrastructure Programs (P40 OD010440) and are to be
requested directly from CGC. Funded by National Institute of General Medical Sciences
grant R35GM147704 to K.L.G. and NIGMS Diversity Supplement 3R35GM147704-
02S1 to support the work of F.A.K.

Data and resource availability
All relevant data and details of resources can be found within the article.

FIGURE LEGENDS
**Figure 1: Dauer larvae that form after prolonged starvation and crowding have
smaller gonads and recover to have smaller broods than dauer larvae that form
constitutively**.
(A) Gonad length in facultative dauers of control *ina-1::mNG; naSi2(mex-
5p::H2B::mCherry::nos-2 3'UTR)* early (n=24) and late (n=28), *daf-2(e1370);naSi2(mex-
5p::H2B::mCherry::nos-2 3'UTR)* early (n=19) and late (n=19), *and daf-7(e1372)* early
(n=33) and late (n=15). Kruskal-Wallis test statistic=109.2, p<0.0001 with Dunn's
correction for multiple comparisons.
(B) Representative differential interference contrast (DIC) microscopy images of
facultative mutant early and late dauer gonads. Gonads outlined.
(C) Gonad length in worms reared at 25°C. Control dataset same as in (1A), *daf-
2(e1370);naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR)* constitutive early (n=22) and late
(n=29) dauers, and *daf-7(e1372)* constitutive early (n=31) and late (n=32) dauers.
Kruskal-Wallis test statistic=109.1, p<0.0001 with Dunn's correction for multiple
comparisons.
(D) Representative DIC images of constitutive mutant early and late dauer gonads.
Gonads outlined.
(E) Schematic for brood size counts in (1F,G).
(F) Brood size of control strain. Never-dauer control (n=15), recovered early (n=21), and
recovered late (n=17). Includes animals that died after recovery and animals that were
fully sterile, does not include animals that failed to recover from dauer. Kruskal-Wallis
test statistic=10.89, p=0.0043 with Dunn's correction for multiple comparisons.
(G) Brood size of *daf-2(e1370);naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR)*. Never-
dauer control (n=14), light shaded background indicates constitutive early (n=14) and
late (n=12) recovered animals, dark shaded background indicates facultative early

(n=14) and late (n=17) recovered animals. Includes animals that died after recovery and
animals that were fully sterile but not animals that failed to recover from dauer. Kruskal-
Wallis test statistic=27.76, $p < 0.0001$ with Dunn's correction for multiple comparisons.
All scale bars 10 µM. Significance indicated as $p < 0.0001$ ****; $p < 0.005$ **.

Figure 2: Withdrawal of food in the pre-dauer period arrests germ cell
proliferation after a single round of division.
(A) Schematic illustrating food removal experiments. Worms moved off food (dashed
lines, imaged "Baseline" (2B,C)) and reared until dauer entry (stars, imaged "Dauer"
(2B,C). All experiments in Fig. 2 were conducted at 25°C.
(B) Results of experiment detailed in (2A), showing number of germ cell nuclei in *daf-*
2(e1370);naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR). Sample sizes: 12h fed baseline
n=31, dauers n=22; 16h fed baseline n=16, dauers n=8; 20h fed baseline n=18, dauers
n=13; 24h baseline n=23, dauer=24; continuously fed dauers n=33.
(C) Representative images of *daf-2(e1370);naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR)*
(magenta) measured for (2B) at time of food removal (baseline, left) and in dauer (right).
Gross structure of the gonad outlined in white.
(D) Experiment as in (2A) with additional +24h endpoint. Darker dots indicate animals in
dauer diapause at the time of counting (see 2E). "12h fed" n=13; "12h fed, 24h starved"
n=16; "12h fed, 24h fed" n=14, refers to animals that were washed at 12h and returned
to food. "12h fed, 72h starved" n=16. Kruskal-Wallis test statistic=39.12; $p < 0.0001$ with
Dunn's correction for multiple comparisons.
(E) Percentage of worms that had reached the dauer stage under each condition in 2D.
In 2D,E, a grey shaded box marks the samples quantified in parallel after 24 hours
under experimental conditions, both on and off food.
All scale bars 10 µM. Significance indicated as $p < 0.0001$ ****; $p < 0.005$ **.

Figure 3: Early and late dauers differ in somatic gonad cell size and germline cell
number
(A) Representative images of early and late dauers expressing (green) *ina-1::mNG*, an
alpha integrin subunit and (magenta) *naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR)*
transgene marking germ cell nuclei. Gonads outlined. Inset image of a germ cell
nucleus showing characteristic compacted chromatin morphology of a quiescent germ
cell (Belew et al., 2021), scale bar 2 µM.
(B) Germ cell nuclei per animal in *ina-1::mNG;naSi2(mex-5p::H2B::mCherry::nos-2*
3'UTR) early (n=24) and late (n=28) dauers. Mann-Whitney test statistic U=0, $p < 0.0001$.
(C) Germ cell number (3B) is highly correlated to the length of the gonad (same
samples as in Fig. 1A) for early (light) and late (dark) dauers. $R^2 = 0.9317$.

(D) Nuclear size of early (n=35 nuclei from 3 worms) and late (n=27 nuclei from 4

worms) dauers shown in 3A,B. Unpaired t-test statistic t=2.591, df=60; p=0.0120.

(E) Representative DIC images of wild type early and late dauer gonads (outlined).

(F) Gonad length of wild type early (n=35) and late (n=40) dauers. Mann Whitney test

statistic U=18, p<0.0001.

(G) Body width of wild type early (n=35) and late (n=40) dauers (same samples as in

3F). Mann Whitney test statistic U=17, p<0.0001.

(H) Test for isometric scaling of gonad with body. Ratio of gonad length to body width of

the same early (n=35) and late (n=40) dauers in 3F,G. Mann-Whitney test statistic

U=111, p<0.0001.

(I) Illustration of the 12 somatic gonad cells based on (Lints and Hall, 2004), L2/L3

molt). We observe the same number of cells in the dauer (K, L). Distal tip cells (DTC),

sheath-spermathecal (SS), dorsal uterine (DU), ventral uterine (VU), anchor cell (AC).

Germ cells (GC) are separated into two gonad arms. Brackets at top indicate

measurements used in (3K).

(J) Representative images of early (top) and late (bottom) dauer gonads (outlined) with

a somatic gonad-specific nuclear marker *arTi435(rps-27p::2xnls::gfp(flexon)::unc-54 3'*

UTR); arTi237(ckb-3p::Cre(opti)::tbb-2 3' UTR) (Shaffer and Greenwald, 2022)). Single-

slice images colored by fire LUT.

(K) Distance between three sets of cells of the somatic gonad in early and late dauers of

the strain in (3J). DTC-DTC, DTC-closest SS cell in that same arm, and SS-SS cell

across the central region. DTC-DTC, Mann-Whitney test statistic U=31, p<0.0001; DTC-

SS, Mann-Whitney test statistic U=193, p<0.0001; SS-SS, Mann-Whitney test statistic

U=21, p<0.0001.

(L) Area of trace around sum projection of DTC as in (M) in early and late dauers and

an L2 control. Welch's ANOVA test statistic=91.91 (2.000, 24.57), p<0.0001 with

Dunnett's T3 post-hoc multiple comparisons test.

(M) Sum projection through ten 0.3 μm z-slices of the superficial surface of a DTC

expressing a membrane-localized transgenic reporter *qIs154(lag-2p:: myr::tdTomato)*

for stages described at top.

All scale bars (except inset) 10 μM. Significance indicated as p<0.0001 ****; p<0.005 **;

p<0.05 *

Figure 4: Dauer germ cells are arrested in a Notch independent state with a

reduced DTC presentation of the LAG-2 stemness cue

(A) Schematic illustrating temperature shift experiment testing germline *glp-1*/Notch-

independence in dauer. Well-fed control *glp-1(bn18)* mutants were compared dauers of

that same genotype from starved and crowded plates. The experiment was repeated for

early (<5d in dauer) and late (>4 weeks) dauers.

(B) Worm fertility after (4A). Control (never-dauer population temp shift in parallel with
early dauer) n=19, Recovered (early) n=14, Control (temp shift in parallel with late
dauer) n=9, Recovered (late) n=15.
(C-E) Sum projection through ten 0.3 μm z-slices of the superficial surface of a DTC
expressing endogenously tagged LAG-2 protein (green, *lag-2(cp193[lag-2::*
mNeonGreen])) and a membrane-localized transgenic reporter (magenta, *qIs154(lag-*
2p:: myr::tdTomato)) for stages described at top. Each fluorescence channel is shown
with identical scaling between treatments, meaning apparent differences in brightness
reflect actual differences in signal intensity.
(C-C") Fed L2 control.
(D-D") Early dauer.
(E-E") Late dauer.
(F-G) Integrated density measured for DTCs described in 4C-E. (F) LAG-2::mNG in
early (n=15 DTCs) and late (n=13 DTCs) dauers and L2 control (n=14 DTCs). Welch's
ANOVA test statistic W=43.65 (2.000, 22.50), p<0.0001 with Dunnett's T3 multiple
comparisons.
(G) *lag-2p::myrTdTomato* transcriptional reporter in early (n=15 DTCs) and late (n=13
DTCs) dauers and L2 control (n=14 DTCs). Ordinary one-way ANOVA test statistic
F=12.27, p<0.0001 with Tukey's multiple comparisons test.
(H) Schematic of experiment performed for 4I-K. The transgenes shown in 4C-E were
crossed into a *daf-2(e1370)* mutant background. Worms were raised on food at 25°C for
20h, imaged, moved off of food, and imaged 5h later.
(I-K) Representative images of worms described in (4H). Sum projection through ten 0.3
μm z-slices of the superficial surface of the DTC coexpressing endogenously tagged
LAG-2 protein (green) with DTC membrane-localized transcriptional reporter (magenta).
(I-I") Fed L2 control animal reared at 25°C.
(J-J") L2d fed for 20h at 25°C.
(K-K") Same as in (J) but 5h after food removal at 25°C.
(L-M) Integrated density measured for DTCs described in 4H-K. (L) LAG-2::mNG, 20h
L2d fed baseline (n=8) 5h starved (n=12), L2 control (same dataset as shown in (4F)).
Welch's ANOVA test statistic W (DFn, DFd)=61.10 (2.000, 17.93), p<0.0001 with
Dunnett's T3 multiple comparisons.
(M) Transcriptional reporter *lag-2p:: myr::tdTomato,* 20 h L2d fed baseline (n=8 DTCs)
and 5h starved (n=12), L2 control (same dataset as in (4G)). Kruskal-Wallis test
statistic=25.41, p<0.0001 with Dunn's post-hoc correction for multiple comparisons.
All scale bars 10 μM. DTCs outlined in yellow. Significance indicated as p<0.0001 ****;
p<0.0005 ***; p<0.005 **.
945
946

Figure 5: Removal from dauer maintenance conditions triggers a burst of LAG-2
presentation on the DTC niche that precedes a return to Notch dependence
(A-D) An otherwise wild-type strain coexpressing the endogenously tagged LAG-2
protein *lag-2(cp193[lag-2:: mNeonGreen])* (green in 5C,D) with a membrane-localized
transgenic reporter *qIs154(lag-2p:: myr::tdTomato)* (magenta in 5C,D) in dauer (same
dataset at "late" in Fig. 4F,G as this was an extension of that experiment and 2 h after
removal from dauer maintenance conditions.
(A) Fold change of LAG-2::mNG relative to mean at 0h (0h n=13; 2h n=15). Two high
outliers shown at non-proportional distance from the bulk of the data. 2h mean (solid),
median (dashed). Mann-Whitney test statistic U=22, p=0.0001.
(B) Fold change of the *lag-2p::myrTdTomato* transcriptional reporter for samples in (5A).
Unpaired t-test statistic t=0.6645, df=27; p=0.5120 (ns).
(C-D) Sum projection through ten 0.3 µm z-slices of the superficial surface of a DTC for
stages described at top. For 5C,D, each fluorescence channel is shown with identical
scaling between time points, meaning apparent differences in brightness reflect actual
differences in signal intensity.
(C-C") 0h of recovery.
(D-D") 2h of recovery.
(E-H) Otherwise wild-type strain co-expressing a *lag-2(bmd202 [lag-*
2::P2A::H2B::mTurquoise2]) polycistronic histone reporter knocked into the endogenous
lag-2 locus (magenta in 5G-H) with a *lag-2p::mNeonGreen::PLCδ1^{PH}* membrane-
localized transcriptional reporter (green in 5G-H), same treatment at A-D.
(E) Fold change of polycistronic reporter *lag-2(bmd202 [lag-2*::P2A::H2B::mTurquoise2])
relative to mean at 0h (0h n=46; 2h n=48). Unpaired t-test statistic t=0.5458, df=92,
p=0.5865 (ns).
(F) Fold change of *lag-2p::mNeonGreen::PLCδ1^{PH}* transcriptional reporter relative to
mean at 0h (0h n=46; 2n=48). Mann-Whitney test statistic U=1031, p=0.5851 (ns).
(G-H) Sum projections through ~20 0.3 µm Z-slices from superficial surface of DTC to
deep bottom of nucleus. Acquired with a 1.6x optical zoom.
(G-G") 0h of recovery.
(H-H") 2h of recovery.
(I) Schematic of dauer recovery *glp-1(bn18)* temperature-shift experiment testing the
return of Notch-dependence to the germline shown in 5J.
(J) Percentage of animals recovering as fertile vs. sterile results after removal from
dauer-maintenance conditions at 25°C for the first 4 h (top, n=39), 6 h (middle, n=34))
and 8 h (bottom, n=43) of recovery.
DTCs outlined in yellow. All scale bars 10 µM.
984
985

References

Ailion, M., Thomas, J.H., 2000. Dauer formation induced by high temperatures in Caenorhabditis elegans. Genetics 156, 1047–1067. https://doi.org/10.1093/genetics/156.3.1047.

Angelo, G., Van Gilst, M.R., 2009. Starvation protects germline stem cells and extends reproductive longevity in C. elegans. Science 326, 954–958. https://doi.org/10.1126/science.1178343.

Aprison, E.Z., Dzitoyeva, S., Ruvinsky, I., 2024. The roles of TGFβ and serotonin signaling in regulating proliferation of oocyte precursors and germline aging. bioRxiv.

Austin, J., Kimble, J., 1987. glp-1 is required in the germ line for regulation of the decision between mitosis and meiosis in C. elegans. Cell 51, 589–599. https://doi.org/10.1016/0092-8674(87)90128-0.

Barbieri, M., Bonafè, M., Franceschi, C., Paolisso, G., 2003. Insulin/IGF-I-signaling pathway: an evolutionarily conserved mechanism of longevity from yeast to humans. Am. J. Physiol. Endocrinol. Metab. 285, E1064-71. https://doi.org/10.1152/ajpendo.00296.2003.

Baugh, L.R., 2013. To grow or not to grow: nutritional control of development during Caenorhabditis elegans L1 arrest. Genetics 194, 539–555. https://doi.org/10.1534/genetics.113.150847.

Belew, M.D., Chien, E., Wong, M., Michael, W.M., 2021. A global chromatin compaction pathway that represses germline gene expression during starvation. J. Cell Biol. 220. https://doi.org/10.1083/jcb.202009197.

Blackwell, T.K., Steinbaugh, M.J., Hourihan, J.M., Ewald, C.Y., Isik, M., 2015. SKN-1/Nrf, stress responses, and aging in Caenorhabditis elegans. Free Radic. Biol. Med. 88, 290–301. https://doi.org/10.1016/j.freeradbiomed.2015.06.008.

Brenner, S., 1974. THE GENETICS OF *caenorhabditis elegans*. Genetics 77, 71–94. https://doi.org/10.1093/genetics/77.1.71.

Byrd, D.T., Knobel, K., Affeldt, K., Crittenden, S.L., Kimble, J., 2014. A DTC niche plexus surrounds the germline stem cell pool in Caenorhabditis elegans. PLoS ONE 9, e88372. https://doi.org/10.1371/journal.pone.0088372.

Cassada, R.C., Russell, R.L., 1975. The dauerlarva, a post-embryonic developmental variant of the nematode Caenorhabditis elegans. Dev. Biol. 46, 326–342. https://doi.org/10.1016/0012-1606(75)90109-8.

Chi, C., Ronai, D., Than, M.T., Walker, C.J., Sewell, A.K., Han, M., 2016. Nucleotide levels regulate germline proliferation through modulating GLP-1/Notch signaling in C. elegans. Genes Dev. 30, 307–320. https://doi.org/10.1101/gad.275107.115.

Cinquin, O., Crittenden, S.L., Morgan, D.E., Kimble, J., 2010. Progression from a stem cell-like state to early differentiation in the C. elegans germ line. Proc Natl Acad Sci USA 107, 2048–2053. https://doi.org/10.1073/pnas.0912704107.

Claeys, I., Simonet, G., Poels, J., Van Loy, T., Vercammen, L., De Loof, A., Vanden Broeck, J., 2002. Insulin-related peptides and their conserved signal transduction pathway. Peptides 23, 807–816. https://doi.org/10.1016/s0196-9781(01)00666-0.

Corsi, A.K., Wightman, B., Chalfie, M., 2015. A Transparent Window into Biology: A Primer on Caenorhabditis elegans. Genetics 200, 387–407. https://doi.org/10.1534/genetics.115.176099.

Crittenden, S.L., Leonhard, K.A., Byrd, D.T., Kimble, J., 2006. Cellular analyses of the mitotic
region in the Caenorhabditis elegans adult germ line. Mol. Biol. Cell 17, 3051–3061.
https://doi.org/10.1091/mbc.E06-03-0170.
Dalfó, D., Michaelson, D., Hubbard, E.J.A., 2012. Sensory regulation of the C. elegans germline
through TGF-β-dependent signaling in the niche. Curr. Biol. 22, 712–719.
https://doi.org/10.1016/j.cub.2012.02.064.
Das, D., Arur, S., 2017. Conserved insulin signaling in the regulation of oocyte growth,
development, and maturation. Mol. Reprod. Dev. 84, 444–459.
https://doi.org/10.1002/mrd.22806.
Diehl, F.F., Miettinen, T.P., Elbashir, R., Nabel, C.S., Darnell, A.M., Do, B.T., Manalis, S.R.,
Lewis, C.A., Vander Heiden, M.G., 2022. Nucleotide imbalance decouples cell growth from
cell proliferation. Nat. Cell Biol. 24, 1252–1264. https://doi.org/10.1038/s41556-022-00965-
1.
Erkut, C., Kurzchalia, T.V., 2015. The C. elegans dauer larva as a paradigm to study metabolic
suppression and desiccation tolerance. Planta 242, 389–396.
https://doi.org/10.1007/s00425-015-2300-x.
Eustice, M., Konzman, D., Reece, J.M., Ghosh, S., Alston, J., Hansen, T., Golden, A., Bond,
M.R., Abramowitz, L.K., Hanover, J.A., 2022. Nutrient sensing pathways regulating adult
reproductive diapause in C. elegans. PLoS ONE 17, e0274076.
https://doi.org/10.1371/journal.pone.0274076.
Fortini, M.E., 2009. Notch signaling: the core pathway and its posttranslational regulation. Dev.
Cell 16, 633–647. https://doi.org/10.1016/j.devcel.2009.03.010.
Fox, P.M., Schedl, T., 2015. Analysis of Germline Stem Cell Differentiation Following Loss of
GLP-1 Notch Activity in Caenorhabditis elegans. Genetics 201, 167–184.
https://doi.org/10.1534/genetics.115.178061.
Gems, D., Sutton, A.J., Sundermeyer, M.L., Albert, P.S., King, K.V., Edgley, M.L., Larsen, P.L.,
Riddle, D.L., 1998. Two pleiotropic classes of daf-2 mutation affect larval arrest, adult
behavior, reproduction and longevity in Caenorhabditis elegans. Genetics 150, 129–155.
https://doi.org/10.1093/genetics/150.1.129.
Goedhart, J., von Stetten, D., Noirclerc-Savoye, M., Lelimousin, M., Joosen, L., Hink, M.A., van
Weeren, L., Gadella, T.W.J., Royant, A., 2012. Structure-guided evolution of cyan
fluorescent proteins towards a quantum yield of 93%. Nat. Commun. 3, 751.
https://doi.org/10.1038/ncomms1738.
Golden, J.W., Riddle, D.L., 1984a. The Caenorhabditis elegans dauer larva: developmental
effects of pheromone, food, and temperature. Dev. Biol. 102, 368–378.
https://doi.org/10.1016/0012-1606(84)90201-X.
Golden, J.W., Riddle, D.L., 1984b. A pheromone-induced developmental switch in
Caenorhabditis elegans: Temperature-sensitive mutants reveal a wild-type temperature-
dependent process. Proc Natl Acad Sci USA 81, 819–823.
https://doi.org/10.1073/pnas.81.3.819.
Gordon, K.L., Payne, S.G., Linden-High, L.M., Pani, A.M., Goldstein, B., Hubbard, E.J.A.,
Sherwood, D.R., 2019. Ectopic Germ Cells Can Induce Niche-like Enwrapment by
Neighboring Body Wall Muscle. Curr. Biol. 29, 823-833.e5.
https://doi.org/10.1016/j.cub.2019.01.056.

Gordon, K., 2020. Recent Advances in the Genetic, Anatomical, and Environmental Regulation
of the C. elegans Germ Line Progenitor Zone. J. Dev. Biol. 8.
https://doi.org/10.3390/jdb8030014.
Greer, E.R., Pérez, C.L., Van Gilst, M.R., Lee, B.H., Ashrafi, K., 2008. Neural and molecular
dissection of a C. elegans sensory circuit that regulates fat and feeding. Cell Metab. 8,
118–131. https://doi.org/10.1016/j.cmet.2008.06.005.
Hand, S.C., Denlinger, D.L., Podrabsky, J.E., Roy, R., 2016. Mechanisms of animal diapause:
recent developments from nematodes, crustaceans, insects, and fish. Am. J. Physiol.
Regul. Integr. Comp. Physiol. 310, R1193-211. https://doi.org/10.1152/ajpregu.00250.2015.
Harsimran Kaur Gill, Gaurav Goyal, Gurminder Chahil, 2017. Insect diapause: A review. JAST-
A 7. https://doi.org/10.17265/2161-6256/2017.07.002.
Henderson, S.T., Gao, D., Lambie, E.J., Kimble, J., 1994. lag-2 may encode a signaling ligand
for the GLP-1 and LIN-12 receptors of C. elegans. Development 120, 2913–2924.
https://doi.org/10.1242/dev.120.10.2913.
Hubbard, E.J.A., 2007. Caenorhabditis elegans germ line: a model for stem cell biology. Dev.
Dyn. 236, 3343–3357. https://doi.org/10.1002/dvdy.21335.
Jayadev, R., Chi, Q., Keeley, D.P., Hastie, E.L., Kelley, L.C., Sherwood, D.R., 2019. α-Integrins
dictate distinct modes of type IV collagen recruitment to basement membranes. J. Cell Biol.
218, 3098–3116. https://doi.org/10.1083/jcb.201903124.
Johnson, T.E., Mitchell, D.H., Kline, S., Kemal, R., Foy, J., 1984. Arresting development arrests
aging in the nematode Caenorhabditis elegans. Mech. Ageing Dev. 28, 23–40.
https://doi.org/10.1016/0047-6374(84)90150-7.
Karp, X., 2018. Working with dauer larvae. WormBook 2018, 1–19.
https://doi.org/10.1895/wormbook.1.180.1.
Kenyon, C., Chang, J., Gensch, E., Rudner, A., Tabtiang, R., 1993. A C. elegans mutant that
lives twice as long as wild type. Nature 366, 461–464. https://doi.org/10.1038/366461a0.
Kimble, J., Crittenden, S.L., 2005. Germline proliferation and its control. WormBook 1–14.
https://doi.org/10.1895/wormbook.1.13.1.
Kimble, J., Hirsh, D., 1979. The postembryonic cell lineages of the hermaphrodite and male
gonads in Caenorhabditis elegans. Dev. Biol. 70, 396–417. https://doi.org/10.1016/0012-
1606(79)90035-6.
Kimble, J.E., White, J.G., 1981. On the control of germ cell development in Caenorhabditis
elegans. Dev. Biol. 81, 208–219. https://doi.org/10.1016/0012-1606(81)90284-0.
Kimble, J., 2014. C. elegans germline stem cells and their niche. StemBook.
https://doi.org/10.3824/stembook.1.95.1.
Kimura, K.D., Tissenbaum, H.A., Liu, Y., Ruvkun, G., 1997. daf-2, an insulin receptor-like gene
that regulates longevity and diapause in Caenorhabditis elegans. Science 277, 942–946.
https://doi.org/10.1126/science.277.5328.942.
Kim, S., Paik, Y.-K., 2008. Developmental and reproductive consequences of prolonged non-
aging dauer in Caenorhabditis elegans. Biochem. Biophys. Res. Commun. 368, 588–592.
https://doi.org/10.1016/j.bbrc.2008.01.131.
Klass, M., Hirsh, D., 1976. Non-ageing developmental variant of Caenorhabditis elegans.
Nature 260, 523–525. https://doi.org/10.1038/260523a0.
Kodoyianni, V., Maine, E.M., Kimble, J., 1992. Molecular basis of loss-of-function mutations in

the glp-1 gene of Caenorhabditis elegans. Mol. Biol. Cell 3, 1199–1213.
https://doi.org/10.1091/mbc.3.11.1199.
Linden, L.M., Gordon, K.L., Pani, A.M., Payne, S.G., Garde, A., Burkholder, D., Chi, Q.,
Goldstein, B., Sherwood, D.R., 2017. Identification of regulators of germ stem cell
enwrapment by its niche in C. elegans. Dev. Biol. 429, 271–284.
https://doi.org/10.1016/j.ydbio.2017.06.019.
Lints, R., Hall, D.H., 2004. WormAtlas Hermaphrodite Handbook - Reproductive System -
Somatic Gonad. WormAtlas. https://doi.org/10.3908/wormatlas.1.22.
Li, X., Singh, N., Miller, C., Washington, I., Sosseh, B., Gordon, K.L., 2022. The C. elegans
gonadal sheath Sh1 cells extend asymmetrically over a differentiating germ cell population
in the proliferative zone. eLife 11. https://doi.org/10.7554/eLife.75497.
McGehee, A., 2019. The GLR-1 phenotypes of the daf-7(e1372) allele are not temperature
sensitive. MicroPubl. Biol. 2019. https://doi.org/10.17912/micropub.biology.000158.
Medwig-Kinney, T.N., Sirota, S.S., Gibney, T.V., Pani, A.M., Matus, D.Q., 2022. An in vivo
toolkit to visualize endogenous LAG-2/Delta and LIN-12/Notch signaling in C. elegans.
MicroPubl. Biol. 2022. https://doi.org/10.17912/micropub.biology.000602.
Morao, A.K., Ercan, S., 2021. Hatched and starved: Two chromatin compaction mechanisms
join forces to silence germ cell genome. J. Cell Biol. 220.
https://doi.org/10.1083/jcb.202107026.
Murphy, C.T., Hu, P.J., 2013. Insulin/insulin-like growth factor signaling in C. elegans.
WormBook 1–43. https://doi.org/10.1895/wormbook.1.164.1.
Narbonne, P., Roy, R., 2006. Inhibition of germline proliferation during C. elegans dauer
development requires PTEN, LKB1 and AMPK signalling. Development 133, 611–619.
https://doi.org/10.1242/dev.02232.
Ouellet, J., Li, S., Roy, R., 2008. Notch signalling is required for both dauer maintenance and
recovery in C. elegans. Development 135, 2583–2592. https://doi.org/10.1242/dev.012435.
Ow, M.C., Borziak, K., Nichitean, A.M., Dorus, S., Hall, S.E., 2018. Early experiences mediate
distinct adult gene expression and reproductive programs in Caenorhabditis elegans. PLoS
Genet. 14, e1007219. https://doi.org/10.1371/journal.pgen.1007219.
Pekar, O., Ow, M.C., Hui, K.Y., Noyes, M.B., Hall, S.E., Hubbard, E.J.A., 2017. Linking the
environment, DAF-7/TGFβ signaling and LAG-2/DSL ligand expression in the germline
stem cell niche. Development 144, 2896–2906. https://doi.org/10.1242/dev.147660.
Renfree, M.B., Fenelon, J.C., 2017. The enigma of embryonic diapause. Development 144,
3199–3210. https://doi.org/10.1242/dev.148213.
Ren, P., Lim, C.S., Johnsen, R., Albert, P.S., Pilgrim, D., Riddle, D.L., 1996. Control of C.
elegans larval development by neuronal expression of a TGF-beta homolog. Science 274,
1389–1391. https://doi.org/10.1126/science.274.5291.1389.
Schackwitz, W.S., Inoue, T., Thomas, J.H., 1996. Chemosensory neurons function in parallel to
mediate a pheromone response in C. elegans. Neuron 17, 719–728.
https://doi.org/10.1016/s0896-6273(00)80203-2.
Schindler, A.J., Baugh, L.R., Sherwood, D.R., 2014. Identification of late larval stage
developmental checkpoints in Caenorhabditis elegans regulated by insulin/IGF and steroid
hormone signaling pathways. PLoS Genet. 10, e1004426.
https://doi.org/10.1371/journal.pgen.1004426.

Schluter, D., Price, T.D., Rowe, L., 1991. Conflicting selection pressures and life history trade-offs. Proc. R. Soc. Lond. B 246, 11–17. https://doi.org/10.1098/rspb.1991.0118.

Seidel, H.S., Kimble, J., 2011. The oogenic germline starvation response in C. elegans. PLoS ONE 6, e28074. https://doi.org/10.1371/journal.pone.0028074.

Shaffer, J.M., Greenwald, I., 2022. Floxed exon (Flexon): A flexibly positioned stop cassette for recombinase-mediated conditional gene expression. Proc Natl Acad Sci USA 119. https://doi.org/10.1073/pnas.2117451119.

Shaner, N.C., Campbell, R.E., Steinbach, P.A., Giepmans, B.N.G., Palmer, A.E., Tsien, R.Y., 2004. Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma sp. red fluorescent protein. Nat. Biotechnol. 22, 1567–1572. https://doi.org/10.1038/nbt1037.

Shaner, N.C., Lambert, G.G., Chammas, A., Ni, Y., Cranfill, P.J., Baird, M.A., Sell, B.R., Allen, J.R., Day, R.N., Israelsson, M., Davidson, M.W., Wang, J., 2013. A bright monomeric green fluorescent protein derived from Branchiostoma lanceolatum. Nat. Methods 10, 407–409. https://doi.org/10.1038/nmeth.2413.

Singh, N., Zhang, P., Li, K.J., Gordon, K.L., 2024. The Rac pathway prevents cell fragmentation in a nonprotrusively migrating leader cell during C. elegans gonad organogenesis. Curr. Biol. 34, 2387-2402.e5. https://doi.org/10.1016/j.cub.2024.04.073.

Sternberg, P.W., Van Auken, K., Wang, Q., Wright, A., Yook, K., Zarowiecki, M., Arnaboldi, V., Becerra, A., Brown, S., Cain, S., Chan, J., Chen, W.J., Cho, J., Davis, P., Diamantakis, S., Dyer, S., Grigoriadis, D., Grove, C.A., Harris, T., Howe, K., Stein, L., 2024. WormBase 2024: status and transitioning to Alliance infrastructure. Genetics 227. https://doi.org/10.1093/genetics/iyae050.

Stiernagle, T., 2006. Maintenance of C. elegans. WormBook 1–11. https://doi.org/10.1895/wormbook.1.101.1.

Suarez Rodriguez, F., Sanlidag, S., Sahlgren, C., 2023. Mechanical regulation of the Notch signaling pathway. Curr. Opin. Cell Biol. 85, 102244. https://doi.org/10.1016/j.ceb.2023.102244.

Sulston, J.E., Horvitz, H.R., 1977. Post-embryonic cell lineages of the nematode, Caenorhabditis elegans. Dev. Biol. 56, 110–156. https://doi.org/10.1016/0012-1606(77)90158-0.

Sulston, J.E., Schierenberg, E., White, J.G., Thomson, J.N., 1983. The embryonic cell lineage of the nematode Caenorhabditis elegans. Dev. Biol. 100, 64–119. https://doi.org/10.1016/0012-1606(83)90201-4.

Taylor, S.E., 2010. Mechanisms linking early life stress to adult health outcomes. Proc Natl Acad Sci USA 107, 8507–8512. https://doi.org/10.1073/pnas.1003890107.

Tenen, C.C., Greenwald, I., 2019. Cell Non-autonomous Function of daf-18/PTEN in the Somatic Gonad Coordinates Somatic Gonad and Germline Development in C. elegans Dauer Larvae. Curr. Biol. 29, 1064-1072.e8. https://doi.org/10.1016/j.cub.2019.01.076.

Tian, X., Hansen, D., Schedl, T., Skeath, J.B., 2004. Epsin potentiates Notch pathway activity in Drosophila and C. elegans. Development 131, 5807–5815. https://doi.org/10.1242/dev.01459.

Trent, C., Tsuing, N., Horvitz, H.R., 1983. Egg-laying defective mutants of the nematode Caenorhabditis elegans. Genetics 104, 619–647.

https://doi.org/10.1093/genetics/104.4.619.

Tullet, J.M.A., Hertweck, M., An, J.H., Baker, J., Hwang, J.Y., Liu, S., Oliveira, R.P., Baumeister,
R., Blackwell, T.K., 2008. Direct inhibition of the longevity-promoting factor SKN-1 by
insulin-like signaling in C. elegans. Cell 132, 1025–1038.
https://doi.org/10.1016/j.cell.2008.01.030.

Vowels, J.J., Thomas, J.H., 1992. Genetic analysis of chemosensory control of dauer formation
in Caenorhabditis elegans. Genetics 130, 105–123.
https://doi.org/10.1093/genetics/130.1.105.

Wadsworth, W.G., Riddle, D.L., 1989. Developmental regulation of energy metabolism in
Caenorhabditis elegans. Dev. Biol. 132, 167–173. https://doi.org/10.1016/0012-
1606(89)90214-5.

Webster, A.K., Chitrakar, R., Taylor, S.M., Baugh, L.R., 2022. Alternative somatic and germline
gene-regulatory strategies during starvation-induced developmental arrest. Cell Rep. 41,
111473. https://doi.org/10.1016/j.celrep.2022.111473.

Webster, A.K., Jordan, J.M., Hibshman, J.D., Chitrakar, R., Baugh, L.R., 2018.
Transgenerational Effects of Extended Dauer Diapause on Starvation Survival and Gene
Expression Plasticity in Caenorhabditis elegans. Genetics 210, 263–274.
https://doi.org/10.1534/genetics.118.301250.

Weinmaster, G., Fischer, J.A., 2011. Notch ligand ubiquitylation: what is it good for? Dev. Cell
21, 134–144. https://doi.org/10.1016/j.devcel.2011.06.006.

Wilsterman, K., Ballinger, M.A., Williams, C.M., 2021. A unifying, eco-physiological framework
for animal dormancy. Funct. Ecol. 35, 11–31. https://doi.org/10.1111/1365-2435.13718.

Wu, X., Fleming, A., Ricketts, T., Pavel, M., Virgin, H., Menzies, F.M., Rubinsztein, D.C., 2016.
Autophagy regulates Notch degradation and modulates stem cell development and
neurogenesis. Nat. Commun. 7, 10533. https://doi.org/10.1038/ncomms10533.

Yamamoto, S., Charng, W.-L., Bellen, H.J., 2010. Endocytosis and intracellular trafficking of
Notch and its ligands. Curr. Top. Dev. Biol. 92, 165–200. https://doi.org/10.1016/S0070-
2153(10)92005-X.

Figure 1:

Figure 2:

Figure 3:

Figure 4:

Figure 5: