1 **Pre-dauer starvation rapidly and reversibly reduces niche proliferative signaling**
2 **to the *C. elegans* germ line**
3
4

5 Fred A. Koitz[1], Camille P. Miller[1], Brian Kinney[1], Kacy Lynn Gordon[1,2]
6

7 [1]Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, North
8 Carolina 27599
9 [2]To whom correspondence should be addressed, kacy.gordon@unc.edu

10
11 **Running title:** Starvation affects the germ line
12
13 **Keywords:** *C. elegans,* dauer, starvation, LAG-2, Notch, germ line, germline stem cell niche
14

ABSTRACT

Early life stresses impact reproductive outcomes in many organisms. In response to
crowding and starvation, *C. elegans* nematodes form dauer larvae, in which
development arrests until conditions improve. We discovered dramatic differences in
gonad size and germ cell number among dauers that form under different conditions.
We used live cell imaging of fluorescent proteins in otherwise wild-type and mutant
animals combined with food-removal, recovery, and brood-size assays to investigate
the causes and consequences of this germline difference. Pre-dauer feeding, but not
nutrient sensing via the DAF-2/insulin-like signaling receptor or DAF-7/TGF-β, is
required for plasticity in gonad size. Gonad differences in dauer have lifelong
reproductive consequences; severely starved worms make small dauer gonads and
have small broods. Pre-dauer starvation induces germline quiescence and near-
instantaneous reduction of the Notch ligand LAG-2 on the germline stem cell niche. A
rapid return to germline Notch dependence and an increase in presentation by the
germline stem cell niche of LAG-2–independent of *lag-2* transcriptional upregulation–are
among the earliest events of dauer recovery.

SUMMARY STATEMENT
33 Pre-dauer starvation arrests gonad growth, reduces presentation of the germline niche
34 cue LAG-2, and diminishes future reproductive success. Dauer recovery rapidly
35 restores LAG-2 ligand and germline Notch dependence.

INTRODUCTION

Lifelong development requires environmental responsiveness within the developmental program. A special case of developmental response to difficult environmental conditions is the diapause state, in which development arrests in unfavorable conditions and resumes when conditions improve (Harsimran Kaur Gill et al., 2017; Renfree and Fenelon, 2017). Obligate diapause states are built into an animal's life cycle in anticipation of poor conditions, while facultative diapause states respond to an organism's environment (Hand et al., 2016; Wilsterman et al., 2021). One well-studied facultative diapause state is the *C. elegans* dauer (Cassada and Russell, 1975). Dauer larvae can withstand poor environmental conditions (Cassada and Russell, 1975; Golden and Riddle, 1984) for many times the typical lifespan of a worm (Klass and Hirsh, 1976). Improved environmental conditions trigger exit from dauer, resumption of development, and a (mostly) normal reproductive life (Cassada and Russell, 1975; Klass and Hirsh, 1976). Famously, time spent in the dauer larval stage does not negatively affect adult lifespan (Hirsh et al., 1976; Klass and Hirsh, 1976).

It is the preservation of fertility rather than increased total lifespan alone that makes dauer adaptive. Previous studies discovered different long-term effects on post-dauer reproduction depending on the dauer-inducing conditions worms experienced (Gimond et al., 2025; Hall et al., 2010; Kim and Paik, 2008; Klass and Hirsh, 1976; Ow et al., 2018; Webster et al., 2018), focused on recovery. We set out to investigate the cells of the gonad and germ line during the dauer larval period itself to look for developmental antecedents of later reproductive differences.

C. elegans goes through four larval stages (L1-L4) before a terminal molt into the reproductive adult (Corsi et al., 2015). The dauer is an alternate third larval stage. Entry into the dauer developmental pathway is specified in the L1 larval stage by low food and high pheromone, and is followed by L2d, which lasts longer than the standard L2 (Golden and Riddle, 1984). In persistently poor conditions, the L2d molts into a non-feeding dauer larva with a thick cuticle, arrested development including absence of germ cell proliferation, and altered metabolism (Cassada and Russell, 1975; Erkut and Kurzchalia, 2015; Golden and Riddle, 1984; Wadsworth and Riddle, 1989).

The *C. elegans* hermaphrodite gonad develops post-embryonically from a four-cell primordium of two somatic blast cells and two primordial germ cells (Kimble and Hirsh, 1979). In the L1, a distal tip cell (DTC) is born from each of the somatic lineages and the

primordial germ cells begin to divide (Kimble and Hirsh, 1979). The germ line relies on
DTC expression of LAG-2, a Delta/Serrate/LAG-2 (DSL) ligand of the Notch signaling
pathway, to maintain distal germ cells in an undifferentiated state, making the DTC the
germline stem cell niche (Kimble and Crittenden, 2005). The developmental arrest of
gonad and germ cell lineages in the dauer larva have been investigated genetically
(Narbonne and Roy, 2006; Tenen and Greenwald, 2019), primarily using genetic
mutants that constitutively form dauers at high temperature, called daf-c mutants.
80
Among these are mutations to genes in the insulin-IGF-1 signaling (IIS) pathway,
including the gene encoding the sole *C. elegans* IIS receptor *daf-2* (Kimura et al., 1997).
IIS integrates nutritional status with metabolism, developmental control, reproduction,
and aging in *C. elegans* (Ewald et al., 2018; Murphy and Hu, 2013) and in other
organisms, including humans (Barbieri et al., 2003; Claeys et al., 2002; Das and Arur,
2017). A "class 2" reduction-of-function allele, *daf-2(e1370),* is experimentally used to
induce dauer larvae at high temperature (Gems et al., 1998; Karp, 2018; Kenyon et al.,
1993). Another reduction-of-function allele, *daf-7(e1372)*/TGF-β, affecting a parallel
pathway leading to dauer entry, also causes constitutive dauer formation at high
temperature (Karp, 2018; Pierce et al., 2001; Ren et al., 1996). The TGF-β pathway
links external cues to the reproductive system (Dalfó et al., 2012; Park et al., 2010,
2021; Pekar et al., 2017).
93
The germ line is especially sensitive to starvation throughout life (Angelo and Van Gilst,
2009; Dalfó et al., 2012; Pekar et al., 2017; Seidel and Kimble, 2011; Webster et al.,
2022), so we hypothesized that pre-dauer nutrition may cause differences in the dauer
reproductive system that impact post-dauer recovery of fertility. We discovered that *C.*
elegans enter dauer with different numbers of germline cells depending on pre-dauer
feeding. More severely starved worms have fewer cells and a smaller brood size upon
recovery than those that fed more before dauer. Surprisingly, the DAF-2/IIS receptor
and the DAF-7/TGF-β ligand were dispensable for this nutritional response. The onset
of starvation rapidly diminished presentation of the stemness cue LAG-2 by the
germline stem cell niche and induced a Notch-independent quiescent state in the germ
line. These DTC and germline changes were reversed within hours of recovery from
dauer, including exponential recovery of LAG-2 on the DTC that appears to be
regulated post-translationally. Altogether, this work reports that differences in recovery
of fertility after dauer and dauer gonad variation are mediated by pre-dauer nutrient
access, with severely starved worms having small gonads and broods. Pre-dauer

feeding affects niche and stem cell maintenance via dynamic regulation of the LAG-2
niche signal at the onset of and recovery from starvation.
111
112 **RESULTS**
113 **Dauers that form in the presence of food have more germ cells and recover with**
114 **larger broods than those that form after abject starvation**
115
Different dauer induction regimes lead to differential reproductive recovery (Ow et al.,
2018). We compared dauer gonads produced by various regimes. Two of these induce
dauer under conditions of abject starvation: dauer entry after food depletion by mixed
populations on parafilmed plates (Karp, 2018) and starvation in liquid culture of a
synchronized population (Hibshman et al., 2021). Two other methods produce dauers in
the presence of food using crowding (high density plating, (Ow and Hall, 2015)), or
isolation of the first-formed dauers from crowded growth plates (this study, see
Methods). We used an otherwise wild-type marker control strain expressing a germ cell
nuclear marker (*mex-5p::H2B::mCherry::nos-2 3'UTR*) and an endogenously tagged
integrin alpha subunit *ina-1(qy23[ina-1::mNG])* that aids visualization of the somatic
gonad.
127
Worms that starved severely before dauer had about half to a third as many germ cells
(average of 10.9 and 10.8 germ cells for starved plate and liquid culture, respectively)
than those that had access to food (average of 22.2 and 37.8 germ cells for high density
and first-formed, respectively, Fig. 1A,B), and we recapitulated the finding that dauers
induced by high-density plating recover to have larger broods than those that starved
before dauer (Fig. 1C). Brood size of first-formed dauers was also higher than starved
dauers (Fig. 1C and Table S1A). Different dauer induction protocols changed overall
gonad size but did not alter somatic cell numbers in the dauer gonad, only their
proximity to one another (Fig. S1A-H), demonstrating that dauer gonads can vary in
germ cell number and somatic cell size. The gonad displayed allometric scaling with
body width and was relatively smaller in starved dauers compared to first-formed dauers
(Fig. S1E), further demonstrating different growth dynamics in the soma and germ line.
140
We then used the field-standard dauer-formation constitutive (daf-c) mutants carrying
reduction-of-function alleles encoding *daf-2(e1370)* (the sole IIS-like signaling receptor)
or *daf-7(e1372)* (a TGF-β ligand), which enter dauer constitutively at high temperature
even while feeding (Karp, 2018; Kenyon et al., 1993; Kimura et al., 1997; Ren et al.,
1996; Thomas et al., 1993; Vowels and Thomas, 1992)). These are not thought to be
true "temperature sensitive" mutations in which a gene product is functional at low
temperature and nonfunctional at higher temperature, rather it is thought these are
reduction-of-function alleles at all temperatures that are sensitized for dauer formation

at high temperatures (Dr. Collin Ewald, ETH Zürich, personal communication, Ewald et
al., 2018; Gems et al., 1998; Greer et al., 2008; Kenyon et al., 1993; McGehee, 2019;
Trent et al., 1983; Tullet et al., 2008). High temperatures also increase the likelihood of
dauer formation in wild-type worms (Ailion and Thomas, 2000). Into these genetic
backgrounds we crossed the germ cell nuclear marker shown in Fig. 1A.
154
At low temperatures, daf-c mutants can form dauers using the same protocols that
induce dauer in wild-type worms (Fig. 1D-F), and when they do, they display the same
patterns as wild-type worms: few germ cells when severely starved before dauer, more
germ cells when able to feed more before dauer (Fig. 1D-F). When *daf-2(e1370)* and
daf-7(e1372) mutants are reared at high temperatures and constitutively form dauer
despite feeding *ad libitum*, they have many germ cells (Fig. 1D-F). Thus daf-c worms,
like wild-type worms, have different germ cell numbers in dauer depending on the dauer
induction method.
163
Finally, we measured brood sizes of the *daf-2(e1370)* strain after recovery from dauer
induced by different methods. Recovered, starvation-induced *daf-2(e1370)* dauers had
lower brood size than recovered crowding-induced dauers, first-formed dauers, and
constitutive dauers of that genotype, which all had the same brood size as the never-
dauer control *daf-2(e1370)* (Fig. 1G). Our experiments implicate pre-dauer feeding as a
potential determinant of germ cell number in dauer and brood size after recovery from
dauer, and wild type alleles of neither *daf-2* nor *daf-7* are required for the response to
pre-dauer feeding.
172
173 **The germ line grows exponentially with pre-dauer feeding in mutants with**
174 **defective IIS and TGF-β signaling**
175
To further test the hypothesis that pre-dauer feeding determines dauer germ cell
number, we performed a series of food removal experiments on *daf-2(e1370)* and *daf-*
7(e1372) daf-c mutants expressing the germ cell nuclear marker. Using daf-c mutants at
25°C ensured that we could vary food exposure while keeping every animal in the L2d
dauer entry program. Worms were reared on plates with bacterial food for the specified
number of hours at 25°C (Fig. 2A), removed from food at 25°C, and maintained until a
population of majority dauers was noted (see Methods).
183
We removed worms from food during the L2d stage at 12, 16, 20, 24 hours, and allowed
others to feed continuously until dauer (48+ hours (Karp, 2018)). For each duration of
feeding, we assayed germ cell number directly upon removal from food (Fig. 2
"Baseline") and again within a day after dauer entry (Fig. 2 "Dauer", cuticular alae
visible, Fig. S2A,A'). These comparisons allowed us to determine the relationship

between time spent feeding and germ cell number at both stages, and how much germ
cell numbers changed between the cessation of feeding and dauer entry (Figs 2A-D,
S2A). Baseline germ cell numbers increased with time spent feeding, and germ cell
numbers at terminal gonad size in dauer increased exponentially for both *daf-2(e1370)*
and *daf-7(e1372)* (Figs 2B,C, S2B,C). Thus we conclude it is not the different methods
of inducing dauer that lead to differences in dauer gonad size, but the fact that worms
feed more or less before entering dauer via these different methods.
196
While actively feeding (comparing among "Baseline" measures in Fig. 2B,C), the
gonads of both *daf-2(e1370)* and *daf-7(e1372)* worms grow very little during the interval
between 12 and 20 hours (Fig. 2B,C), however feeding during this stage sets up
significant differences in terminal gonad size in dauer (for *daf-2(e1370)* $mean_{dauer\ fed\ 12}$
$_h$=6.68, $mean_{dauer\ fed\ 20\ h}$=10.38, Mann-Whitney U=16.50, p<0.0001; for *daf-7(e1372)*
$mean_{dauer\ fed\ 12\ h}$=7.16, $mean_{dauer\ fed\ 20\ h}$=17.04, Welch's t-test t=11.45, p<0.0001).
Comparing the rate of germline growth for *daf-2(e1370)* animals revealed an even more
dramatic response to starvation than that of germ cell number alone (Fig. 2E). Animals
gained an average of one germ cell per hour while feeding (12 h fed+24 h fed), while
animals that starved (12 h fed+24 h starved) added on average only one germ cell total
in that 24 hour period (Fig. 2E, gray shaded box) and entered dauer more slowly (Fig.
2F, gray shaded box). Our findings reveal pre-dauer germline responsiveness to
nutrition is surprisingly robust to mutation in IIS and TGF-β signaling genes.
210
Average germ cell numbers approximately double in both genetic backgrounds between
the food removal baseline and dauer entry at each timepoint (Fig. 2B,C), with the
exception of 24 h of L2d feeding for *daf-2(e1370)* animals, which are known to slow
germ cell division in late L2d in preparation for dauer entry (Narbonne and Roy, 2006).
This slowing may also explain why terminal growth plateaus between 24 h dauers and
continuously fed dauers. Germ cell doubling means cells undergo one additional round
of division before dauer. Germ cells will complete an average of one additional cell
division after DTC laser ablation in the L2-L3 stages of development (Kimble and White,
1981), and genetic ablation of the stemness receptor GLP-1/Notch allows adult germ
cells to complete their current mitotic cell cycle to make a terminal division before
differentiating (Fox and Schedl, 2015). Based on these established cell cycle constraints
on stem-like germ cell proliferation dynamics, our finding that germ cells complete one
round of division after food removal suggests that the withdrawal of food effectuates the
rapid termination of pro-proliferative signaling to the L2d germ line. The pro-proliferative
signal to the germ line is the DTC-expressed LAG-2 ligand, so we next asked whether
this signal was down-regulated upon food removal.
227

The DTC stemness cue LAG-2 requires *daf-2* and *daf-7* in fed L2s, but not for dramatic downregulation upon food removal in L2d

The pro-proliferative, anti-differentiation signal to the germ line is the DTC-expressed LAG-2 ligand activating GLP-1/Notch signaling in the distal germ cells (Henderson et al., 1994). We wanted to test whether shifting worms off food in the L2d sensitive window in which feeding changes germline growth (shown in Fig. 2) also alters the LAG-2 signal in the DTC. We used an endogenously tagged LAG-2::mNeonGreen and an integrated *lag-2* transcriptional reporter that drives a DTC membrane marker (*qIs154(lag-2p::myrTdTomato)*) to quantify DTC LAG-2 abundance and *lag-2* transcriptional activity.

Otherwise wild-type dauer worms have notably less expression of both LAG-2::mNG and *lag-2* transcriptional reporter than fed L2 controls, though the magnitude of transcriptional reporter reduction in dauer (25-30%) cannot alone explain the reduction in LAG-2::mNG protein signal (~30 fold) (Fig. 3A-D). Comparing otherwise wild-type worms to *daf-2(e1370)* and *daf-7(e1372)* reveals that both IIS and TGF-β are required for normal expression levels of of *lag-2* promoter activity and LAG-2::mNG protein in the DTC in fed L2 animals (Fig. 3E-I); this concords with what is known about the positive transcriptional regulation of the *lag-2* promoter by *daf-7* later in continuous, non-dauer development (Pekar et al., 2017).

We could now compare expression of LAG-2::mNG for each daf-c genotype in a food removal assay in L2d (Fig. 3J). Endogenous LAG-2::mNG protein abundance is highly responsive to the onset of starvation in L2d in both genotypes (Fig. 3K-M). LAG-2::mNG drops from the L2d 20 h fed baseline within one hour of food removal in both *daf-2(e1370)* (-27.7%) and *daf-7(e1372)* (-48.1%) mutants, and continues to fall in an exponential decay (Fig. 3M). The abundance of most proteins is governed by first-order kinetics, leading to exponential decay in the absence of new protein synthesis ((McShane et al., 2016) and references therein). Thus our results suggest that LAG-2 deposition on the membrane in the DTC arrests nearly instantaneously upon food removal and existing protein is actively depleted thereafter. The starvation-induced arrest of germ cell proliferation that we observed in Fig. 2 is accompanied by a rapid starvation-induced downregulation of of LAG-2::mNG in the germline stem cell niche, and both processes are robust to deficits in *daf-2* and *daf-7* signaling, because we observe them in both reduction-of-function mutants.

This result distinguishes the process observed here from the *daf-7*-dependent downregulation of *lag-2* transcriptional reporters observed in L4 and young adult animals reared in unfavorable conditions (Dalfó et al., 2012; Pekar et al., 2017). Further

comparisons of *lag-2* promoter activity and protein abundance in *daf-7(e1372)* mutants
reinforced our conclusions that the role of *daf-7* in regulating *lag-2* in pre-dauer and
dauer stages (Fig. S3) is distinct from the simple positive relationship reported during
continuous development by Pekar et al. (2017). Across stages and conditions, *daf-*
7/TGF-β signaling regulates *lag-2* in the DTC in a complex manner, but wildtype *daf-7* is
not required for rapid downregulation of LAG-2 in the DTC after the acute onset of
starvation (Fig. 3L,M).
275
276   **The dauer germ line is maintained in a Notch-independent, quiescent state, and**
277   **exits that state within hours of dauer recovery**
278
We next asked how the germ line tolerates diminished LAG-2 niche signal in dauer.
Under continuous feeding, niche signaling via LAG-2 to the GLP-1/Notch receptor in the
germ cells is required for their maintenance in a mitotic, undifferentiated state at all
developmental stages (Austin and Kimble, 1989, 1987; Hansen and Schedl, 2006;
Henderson et al., 1994; Kimble and Crittenden, 2007; Yochem and Greenwald, 1989).
Both DTC ablation (Kimble and White, 1981) and genetic ablation of Notch signaling
with a temperature-sensitive allele of *glp-1* (Cinquin et al., 2010; Fox and Schedl, 2015;
Kodoyianni et al., 1992) are sufficient to trigger meiosis in all previously-mitotic distal
germ cells within hours. However, transient adult starvation induces a G2-arrested,
Notch-independent quiescent state in which genetic ablation of Notch signaling does not
cause germ cells to differentiate; instead, germ cells remain mitotic upon subsequent
recovery of Notch signaling at the permissive temperature (Seidel and Kimble, 2015).
Once in the dauer stage, worms are in a *de facto* starved state, and dauer germ cells
are G2-arrested (Narbonne and Roy, 2006). We therefore hypothesized that the germ
line of dauers, like that of transiently starved adults, is Notch-independent for the
maintenance of an undifferentiated, stem-like population of germ cells.
295
We investigated dauer germ cell Notch-dependence using worms carrying a
temperature sensitive allele of *glp-1*/Notch*, glp-1(bn18).* At the permissive temperature
(16°C), *glp-1(ts)* mutants have adequate Notch signaling to remain fertile, but upon
shifting to the restrictive temperature (25°C), they lose active Notch signaling and thus
lose the germline stem cell population to differentiation (Kodoyianni et al., 1992). In
adults, signs of meiotic entry appear in distal germ cells within six hours of temperature
upshift (Fox and Schedl, 2015), so the germ cell fate response to lost Notch-signaling is
rapid.
304
We first shifted continuously-fed control populations of *glp-1(bn18)* mutants to the
restrictive temperature for 24 hours (green dashed path in Fig. 4A, and see Methods),
then recovered single individuals of various larval stages to plates at the permissive

temperature. Nearly all animals recovered as sterile adults (1/28 had a few offspring,
Fig. 4B), demonstrating the expected Notch dependence of the fed larval germ line. If
the germ line of dauer worms is likewise Notch-dependent, sterility should also occur
when *glp-1(bn18)* dauers are shifted to the restrictive temperature.
312
This is not what we find. When we caused *glp-1(bn18)* mutant worms to form dauers at
the permissive temperature and then shifted them to the restrictive temperature for 24
hours to abolish Notch signaling (purple solid path in Fig. 4A, and see Methods),
isolated single dauer individuals to plates with food at the permissive temperature, and
allowed the worms to recover, all of the worms recovered as fertile adults (n=29/29, Fig.
4B). Thus, the dauer germ line is in a Notch-independent, G2-arrested state and can
maintain germline stem cells even in the absence of niche signaling. Because germ
cells cannot directly transition from mitotic G2 to prophase I of the meiotic cell cycle
without completing mitotic M phase (Fox and Schedl, 2015), cell-cycle quiescence
protects dauer germ cells, like those in transiently starved adults (Seidel and Kimble,
2015), from differentiating in the absence of Notch activity.
324
We hypothesized that the germ line must return to a Notch-dependent state during
dauer recovery. We performed temperature downshift experiments during dauer
recovery for *glp-1(bn18)* mutant worms (Fig. 4C, and see Methods). When *glp-1(bn18)*
mutant dauers were raised to 25°C, isolated, kept at that restrictive temperature for the
first four hours of recovery from dauer, and then shifted to the permissive temperature
to complete development, all animals recovered as fertile adults (Fig. 4D, n=15/15),
meaning that germ cells remain Notch independent during early dauer recovery. After
six hours of recovery from dauer at 25°C, a third of worms became sterile in adulthood
(Fig. 4D, n=5/15). After eight hours of recovery from dauer at 25°C, more than half of
worms recovered as sterile adults (Fig. 4D, n=8/15). The return of Notch dependence
occurs at the earliest in the 4-6 hour window after recovery from dauer and is prevalent
by eight hours of recovery.
337
Finally, we find that germ cell proliferation recommences in this same window, with a
first doubling observed by eight hours, and modeled at ~7 hours by our nonlinear
regression (Fig. 4E). This proliferation rate accords with the conservative estimate of
doubling in continuously fed L3 germ lines and those recovering from adult reproductive
diapause (average of ~9 hours, (Roy et al., 2016)). Mitotic figures reappear in the germ
line during this time period (Fig. 4F,G). The germline mitotic index we observe (2.26% at
four hours and 1.75% at eight hours) is intermediate between the mitotic indices
reported for wild-type fed larvae and adults (Roy et al., 2016). The exit of G2-arrested
quiescence and resumption of germ cell cycling temporally matches the return of Notch

dependence in the germ line. We next asked when during recovery the Notch pathway
ligand LAG-2 returns to high levels in the DTC.
349
350 **Dauer recovery initiates a transcription-independent burst of LAG-2 ligand**
351 **presentation by the germline stem cell niche**
352
We predicted that LAG-2 protein abundance would rapidly increase to meet the post-
dauer requirement for germline Notch signaling. Immediately after dauer isolation ("0 h"
of recovery, Fig. 5A-C), otherwise wild-type dauers had low LAG-2::mNG protein and
lag-2 transcriptional reporter signal. During the first two hours of dauer recovery, LAG-
2::mNG protein abundance increased ~4-fold, and continued to increase exponentially,
reaching a 38-fold increase by eight hours post dauer recovery (Figs 5A,C, S4A, S5).
We did not observe a corresponding increase in signal from the *lag-2p::myrTdTomato*
transcriptional reporter (Figs 5B,C, S4B), suggesting that the rapid restoration of LAG-2
protein on the DTC may not be regulated transcriptionally.
362
To further test this hypothesis, we examined another pair of *lag-2* reporters. To rule out
potential artifacts caused by the multi-copy transgene or slow maturation of
myrTdTomato fluorescence signal, we used a single-copy transcriptional reporter
encoding a faster-folding fluorescent protein (*cpIs122(lag-2p::mNeonGreen::PLCδ1^{PH}*).
To rule out potential artifacts caused by the selection of the 3kb upstream promoter
region for these transgenes and their lack of endogenous transcript regulation, we used
a histone H2B::mTurquoise2 knocked into the endogenous *lag-2* locus with a P2A
peptide cleavage site (*lag-2(bmd202*[*lag-2*::P2A::H2B::mTurquoise2^lox511I^2xHA])).
This element will generate a polycistronic mRNA (Medwig-Kinney et al., 2022), each
translation of which will produce one LAG-2 protein and one H2B::mTurquoise2 that
separate at translation due to ribosome skipping at the P2A site (Donnelly et al., 2001;
reviewed in de Lima and Lanza, 2021; Donnelly et al., 2001). The H2B::mTurquoise2 is
therefore subject to endogenous *lag-2* transcript-based regulation (e.g. by
transcriptional regulation, 3' UTR-mediated translational repression by microRNAs or
RNA binding proteins, transcript decay, etc.). RNAi against *lag-2* was used to validate
this feature of the reagent (Fig. S6).
379
Up to eight hours after recovery from dauer, we observe no significant change in the
levels of expression vs. baseline of either the faster-folding single copy transcriptional
membrane reporter or polycistronic nuclear reporter (Figs 5D-F, S4C,D). Given that the
germ line of 53% of animals is Notch dependent by eight hours of dauer recovery (Fig.
4D), transcription-based upregulation of *lag-2* would not appear to be sufficient to
increase Notch ligand on the timeline necessary for dauer recovery.
386

Indeed, the only way to see an increase of endogenously tagged LAG-2::mNG but not
an increase in the H2B::mTurquoise2 from the polycistronic knock-in is for the two
proteins to be turned over at different rates. We ruled out that H2B::mTurquoise2 is
particularly long-lived (see Methods and Fig. S7). This leaves only LAG-2 protein
regulation as a source for their different recovery dynamics. We therefore conclude that
the *lag-2* locus is transcriptionally active in the DTC during dauer–despite the germ line
being maintained in a Notch-independent state–and LAG-2 is under protein-specific
negative regulation that is rapidly alleviated upon dauer exit.
395
Taken together we see an exponential rebound of LAG-2 protein on the DTC begin
within two hours of recovery from dauer, preceding a return of Notch dependence and
proliferation in the germ line between 4-8 hours of recovery. Because dauer exit is
asynchronous among individuals (Cassada and Russell, 1975), the precise timing of
these events with respect to dauer exit and to one another may be even more tightly
coordinated. Other early features of dauer exit–loss of the buccal plug/SDS resistance
and resumption of pharyngeal pumping–also occur within a two-hour window of removal
from dauer conditions (Cassada and Russell, 1975), making the return of LAG-2 on the
DTC surface as rapid a response to the end of dauer as any other measured event. Our
findings establish upregulation of germline stem cell niche signaling as a central, early
feature of dauer recovery.
407
DISCUSSION
The germ line is the only lineage in *C. elegans* with an indeterminate division pattern
and variable cell number, while the somatic lineage is invariant (Kimble and Hirsh, 1979;
Sulston and Horvitz, 1977; Sulston et al., 1983)*.* This unique flexibility allows the germ
line to be the only cell population that can respond to starvation by limiting the number
of cells it produces, while the soma continues to the "safe harbor" state of dauer
diapause. Starvation induces a precipitate IIS- and TGF-β-independent halt in pro-
proliferative niche signaling and germ cell proliferation, which is swiftly reversible upon
dauer recovery. That speed is particularly important as germ cells quickly exit the Notch
independence of dauer and require active Notch signaling to maintain stemness as they
resume cycling during recovery from dauer.
419
Rapid changes in protein levels and discordance of protein and transcriptional dynamics
both suggest post-translational regulation (Liu et al., 2016). Exponential decay of LAG-
2::mNG signal after food removal in L2d in daf-c worms suggests active degradation of
LAG-2 without further production. The exponential recovery of LAG-2::mNG upon dauer
exit suggests the cessation of that degradation process. We propose a model by which
steady but fairly low levels of *lag-2* transcription and translation in dauer are coupled

with LAG-2 protein degradation triggered by starvation. Upon dauer recovery, protein
degradation ceases, leading to the rapid accumulation of LAG-2 protein in the DTC.
428
Such a regulatory paradigm famously governs the p53 family of tumor suppressor
proteins, which are transcribed, translated, ubiquitinated, and degraded under normal
growth conditions when p53 is not active. DNA damage or other stresses repress its
ubiquitin ligase, allowing the rapid accumulation, nuclear translocation, and activity of
p53 (Abuetabh et al., 2022). Recently, cyclin D has been discovered to be regulated by
a similar process (Chaikovsky et al., 2021; Maiani et al., 2021; Simoneschi et al., 2021).
Other proteins that mediate rapid response to changing environmental conditions are
also regulated by steady state production and degradation in the absence of induction,
like the hypoxia inducible transcription factor subunit HIF-α (reviewed in Flick and
Kaiser, 2012 and in Weidemann and Johnson, 2008) and the Nrf2 transcriptional
regulator of heme oxygenase HO-1 (Stewart et al., 2003; reviewed in Flick and Kaiser,
2012).
441
Notch signaling is regulated at the protein level in multiple ways (Fortini, 2009; Suarez
Rodriguez et al., 2023; Tian et al., 2004; Wu et al., 2016), including by E3-ubiquitin-
ligase (Fortini, 2009; Yamamoto et al., 2010). In *Drosophila* neuroblast stem cells,
Notch signaling regulates a nutritionally-responsive entry into and exit from quiescence
that is driven by amino acid availability, though this mechanism is insulin dependent
(Sipe and Siegrist, 2017; Sood et al., 2022). Future work will investigate regulation of
LAG-2 in the DTC by the ubiquitin-proteasome pathway, to test if it is regulated in a
similar way to p53 upon starvation and during dauer exit. Such a regulatory mechanism
"takes the foot off the brake" to rapidly induce a regulator under conditions in which its
activity is required, priming a cell to respond to a high-stakes change in state. Protection
of the germ line during starvation and recovery is a high-stakes regulatory event in the
life of a worm.
454
Upon food removal, pre-dauer germ cells complete a single terminal division before
dauer entry. The most severely starved dauers have as few germ cells (~5-8) as worms
with severe loss-of-function alleles of *glp-1*/Notch receptor gene (Austin and Kimble,
1987), and even fewer germ cells than worms have after early laser ablation of the DTC
(Kimble and White, 1981), suggesting the cessation of pro-proliferative cues upon
starvation is near-immediate. Thus we interpret the observation that germ cell number
during recovery from dauer differs depending on how dauer was induced (Ow et al.,
2021, 2018) to reflect differences in initial dauer germ cell number induced by these
same treatments.
464

The question remains whether a reduction in DTC LAG-2 signal in response to
starvation is the proximal cause of germline quiescence, or whether germline
quiescence is a necessary precondition for the germ line to tolerate a reduction in DTC
LAG-2 as a byproduct of starvation. Previous work reported that the dauer germ line is
quiescent for germ cell division even with constitutively active Notch signaling
(Narbonne and Roy, 2006), suggesting that downregulation of LAG-2 in the DTC is not
necessary to prevent germ cell division in dauer itself, though the pre-dauer period we
focus on here was not tested. Dauer germ cells are quiescent for proliferation and also
for differentiation; meiotic germ cells in *glp-1(ts)* mutant dauers will not progress through
spermatogenesis (Narbonne and Roy, 2006). This quiescence depends on AMPK
signaling (*lbk-1* and *aak-2*) independent of the DAF-2 IIS receptor (Narbonne and Roy,
2006). The somatic gonad requires DAF-18/PTEN–independent of the DAF-2 IIS
receptor or DAF-7/TGF-β–to enter a quiescent state and to mediate germline
quiescence in dauer itself, but does not affect quiescence in the pre-dauer period that is
the focus of our study (Tenen and Greenwald, 2019).
480
Expression of both endogenously tagged LAG-2 protein and *lag-2* promoter-driven
transgenes (Fig. 5 and (Narbonne and Roy, 2006)) remain detectable in dauer, so the
downregulation of *lag-2* during starvation is relative, not absolute. Downregulation of
lag-2 transcriptional reporters in response to starvation or pheromone occurs in later
larval stages (Pekar et al., 2017), and the degree of that observed transcriptional
difference (~50%) is on the order of what we observe with transcriptional reporters in
this study in dauers compared to fed L2s (~30%). In contrast, LAG-2::mNG protein is
reduced by *~30 fold* in otherwise wild-type dauers compared to fed L2s and then
increases by a similar magnitude within hours of recovery from dauer, a window of time
in which the signal from transcriptional reporters does not rise at all. Notch signaling via
lag-2 is important in dauer and in germline regulation in many contexts (reviewed in
Gordon, 2020). Expression of *lag-2* in the nervous system is required both during dauer
and for dauer recovery (Ouellet et al., 2008), and *lag-2* reporter activation in the vulval
precursor cells differs between continuous L2 vs. L2d and L3 vs. dauer (Karp and
Greenwald, 2013). LAG-2 protein downregulation in the DTC has been observed during
worm aging (Aprison et al., 2024 preprint; Singh et al., 2024), and *lag-2* transcription
responds to male pheromone signals (Aprison et al., 2024 preprint). All levels of *lag-2*
regulation–from transcriptional to post-translational–sculpt its dynamics in a range of
cells to govern a number of processes, notably those regulating environmental
response in the reproductive system.
501
502 While the precise molecular mechanisms linking nutrition to LAG-2 protein abundance
503 in the DTC remain to be found, we establish that they proceed even with the field-
504 standard *daf-7(e1372)* reduction of TGF-β signal and the *daf-2(e1370)* reduction of-

function allele of the sole *C. elegans* IIS receptor. The reproductive consequences of pre-dauer food intake are also the same in *daf-2(e1370)* animals as they are in wild-type worms (that is, pre-dauer starvation leads to reduced brood size). In addition to seeking the molecular mechanisms of starvation-induced LAG-2 depletion in the DTC, future work will also aim to identify the systemic cues that trigger that depletion.

Early life stresses are known to affect later life outcomes in a range of organisms and contexts with potential mechanisms ranging from epigenetics to inflammation to hormone signaling to lasting variation in central nervous system development and function (Taylor, 2010). In *C. elegans,* primordial germ cells are kept in a quiescent state by the somatic gonad precursors (McIntyre and Nance, 2023). Arrested vs. fed L1 worms have differential transcriptional responses in the soma and germ line (Webster et al., 2022), and display chromatin compaction in the germ line (Belew et al., 2021; Morao and Ercan, 2021). The L1 arrest state is triggered by complete starvation after hatching (Baugh and Hu, 2020; Baugh, 2013; Johnson et al., 1984) and several checkpoints triggered by later starvation can arrest somatic development (Baugh and Hu, 2020; Schindler et al., 2014).

Future work will focus on how the germ line transitions from a Notch-independent state back to Notch-fueled germline proliferation. One possible driver is nucleotide levels, which have been shown to limit cell division but not cell growth in unicellular organisms (Diehl et al., 2022) and to affect germline growth in non-dauer *C. elegans* (Chi et al., 2016). Considering nucleotide levels and the limits imposed by starvation upon cellular energy prompts the observation that continuous transcription, translation, and LAG-2 protein degradation during dauer comes with an energy cost. A dauer worm would only pay this price if the alternative–slower restoration of LAG-2 protein to the germline stem cell niche surface–carried an even greater cost of reduced fertility upon return of the germ line to Notch dependence.

MATERIALS AND METHODS
Sections of this text are adapted from prior Gordon lab publications (Li et al., 2022; Singh et al., 2024), as they describe our standard laboratory practices and equipment.

Strains
We used Wormbase (Sternberg et al., 2024) while conducting this study. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440) and are to be requested directly from CGC. The following *C. elegans* strains were obtained from the CGC: N2 (Brenner, 1974), CB1370 *daf-2(e1370)* III (Kenyon et al., 1993; Kimura et al., 1997), CB1372 *daf-7(e1372)* III (Pierce et al., 2001; Ren et al., 1996), DG2389 *glp-1(bn18)* III (Kodoyianni et al., 1992).

The following *C. elegans* strains were generously shared by community members or
generated previously in our lab (with sources given for each strain and allele):
NK2517 (Gordon et al., 2019) *qIs154(lag-2p:: myr::tdTomato) (Byrd et al., 2014); lag-*
2(cp193[lag-2:: mNeonGreenˆ3xFlag]) V)
KLG034 (Gordon et al., 2019) *cpIs122(lag-2p::mNeonGreen:: PLCδ1^{PH})*II (Linden et al.,
2017)*; lag-2(bmd202 [lag-2*::P2A::H2B::mTurquoise2^lox511I^2xHA]) V (Medwig-
Kinney et al., 2022),
GS9692 *arTi435(rps-27p::2xnls::gfp(flexon)::unc-54 3' UTR* I); *arTi237(ckb-*
3p::Cre(opti)::tbb-2 3' UTR X) (Shaffer and Greenwald, 2022).
554
The following *C. elegans* strains were generated for this paper by crossing existing
alleles and markers (with sources given for alleles not previously attributed above):
KLG047 *ina-1(qy23[ina-1::mNeonGreen]) (Jayadev et al., 2019)*; *naSi2(mex-*
5p::H2B::mCherry::nos-2 3'UTR) II (Linden et al., 2017). We call this the "marker control
strain" in the text and refer to the *nasi2* transgene as "germ cell histone marker".
KLG050 *daf-2(e1370)* III*; naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR)* II,
KLG051 *daf-2(e1370)* III*; qIs154(lag-2p:: myr::tdTomato)* V*; lag-2(cp193[lag-2::*
mNeonGreenˆ3xFlag]) V.
KLG055 *daf-7(e1372)* III*;nasi2(mex-5p::H2B::mCherry::nos-2 3'UTR)* II,
KLG056 *daf-2(e1372)* III*; qIs154(lag-2p:: myr::tdTomato)* V*; lag-2(cp193[lag-2::*
mNeonGreenˆ3xFlag]) V.
566
567 **Strain maintenance and synchronization**
568 Worm strains were maintained on nematode growth media (NGM) at 16°C unless
569 otherwise specified.
570
Dauer formation by starvation on parafilmed plates
To more precisely time the formation of facultative dauers, a large, mixed population of
well-fed worms was transferred from an uncrowded growth plate to NGM plates seeded
with ~70 μL of an overnight culture of *E. coli* OP50 bacterial food. Plates were sealed
with parafilm and placed at 25°C for the non-temperature-sensitive strains and 16°C for
strains containing *daf-2(e1370), daf-7(e1372),* or *glp-1(bn18)* temperature-sensitive
alleles. Plates were monitored daily for evidence of starvation (bacterial lawn depleted
and worms dispersed). At 25°C this took ~2-4 days and at 16°C this took ~5-7 days.
Plates were analyzed after 1+ weeks of starvation.
580
581 **Recovery of first-formed dauers**
582 As above, evaluated <4 days after food exhaustion was noted, before a robust
583 population of dauers are visible on the plate. Dauers were isolated with a treatment of
584 1% SDS for 20 minutes (see Dauer isolation, below).

Dauer formation by liquid culture

For dauer formation by liquid culture, the protocol from Hibshman et al., 2021 was used. Briefly, animals were egg prepped (Stiernagle, 2006) from 4-5 well-populated NGM plates seeded with OP50 and embryos were allowed to hatch out in S-complete medium overnight such that a synchronized population of L1s was generated. Animals were concentrated to 5 worms/µL in S-complete and added to glass test-tubes or 25 mL Erlenmeyer flasks, to which *E. coli* HB101 bacteria was added from a concentrated stock such that the final concentration of bacteria in S-complete was 1 mg/mL. Cultures were placed on a shaker (180 rpm). Deviating from the original protocol, animals were incubated at 25°C for 4-5 days and then treated with SDS to match the dauer isolation protocol used in our other methods.

Dauer formation by high density plating

For high-density plating conditions, the protocol described in Ow & Hall 2015 (Ow and Hall, 2015) and summarized by (Karp, 2018) was used. Briefly, mixed populations of worms were chunked to four 100 mm plates seeded with 1 mL *E. coli* OP50 or 6-8 60 mm plates seeded with the same titre of food. Plates were grown until densely populated but not depleted of bacterial food. Worms were washed from plates in M9 buffer and allowed to gravity-settle, after which the supernatant (containing younger worms) was removed from the pellet of adults, which was retained for transfer to a single 35 mm plate seeded with *E. coli* OP50. We seeded the 35 mm plates with more bacterial food (100-200 uL of 20x concentrated *E. coli* OP50) than the original protocol (50 uL of an overnight culture) to ensure that the animals had abundant food available prior to dauer formation. The culture plates were then blanketed with a cooked, pasteurized egg white mixture as per the original protocol. To these plates, the pelleted adults were added. For daf-c strains at 16°C, plates were incubated for longer than the 72 h specified by the original protocol, as worm development is slower at that temperature. To recover dauers, plates were SDS-treated when there were dauers visible in a sample of the egg white mixture.

Dauer isolation protocols

For plates with mixed populations, dauers were isolated with 1% SDS (Karp, 2018). After SDS treatment, the population of recovered animals was placed onto an unseeded NGM plate and the excess liquid was allowed to dry, after which animals were picked to a slide for imaging, were kept on the unseeded plate for the specified number of hours to evaluate short-term recovery, or picked to an NGM plate seeded with OP50 for recovery and brood measurement. For synchronous daf-c populations in dauer, worms were picked from all-dauer populations and dauer morphology was ascertained for each specimen by microscopy.

Daf-c mutant constitutive dauer formation

Constitutive dauer formation of daf-c mutants (*daf-2(e1370)* and *daf-7(e1372)*) was achieved by collecting embryos via a standard egg prep (Stiernagle, 2006) with embryos hatching into L1s in M9 to create a synchronized population. This population was concentrated and added to NGM plates seeded with *E. coli* OP50. The mutant daf-c strains used in this study were the reduction of function alleles of the insulin-like receptor, class II allele *daf-2(e1370),* and the TGF-β ligand, *daf-7(e1372)* that are field-standard genetic backgrounds for studying dauer (Karp, 2018).

Brood size assays

All brood size assays were conducted at 16°C to match the necessary conditions of the daf-c strains. After experimental treatment (see below), worms were singly picked to NGM plates seeded with ~70 µL *E. coli* OP50. Egg-laying adults were subsequently passaged to a new plate once daily until egg laying stopped. Plates with progeny were kept at 16°C until the oldest worm on the plate had reached L4, after which they were counted. Brood totals include broods of animals that died during recovery and animals that were fully sterile, does not include animals that failed to recover from dauer to adulthood.

For never-dauer control brood assays, fed L4 animals of each strain were singled from mixed populations on Day 0. For brood after recovery from constitutive dauer formation, healthy plates of adults were egg prepped; eggs were rolled in M9 overnight at room temperature to obtain a population of arrested L1 larvae. Animals were dropped onto NGM plates seeded with OP50. Plates were sealed with parafilm and put at the restrictive temperature (25°C) to induce dauer formation. Worms were then isolated by SDS treatment (see next section) and dauers were singly recovered to NGM plates seeded with OP50 at 16°C to recover to reproductive adulthood.

Brood size was not measured for *daf-7(e1372)* mutant strains because these worms have a high rate of bagging, even at the permissive temperature (Karp, 2018; Shaw et al., 2007; Trent et al., 1983).

Food removal experiments

A synchronized population of *daf-2(e1370); naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR II)* L1s or *daf-7(e1372); naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR II)* was obtained via an egg prep of at least eight 60 mm NGM plates seeded with *E. coli OP50.* Synchronized L1s were added to NGM plates seeded with OP50 and were put at 25°C to induce constitutive dauer formation. After the specified number of hours (Fig. 2A), animals were rinsed off the seeded plate with M9, recovered to a 15 mL conical tube, and were subsequently washed in M9 3-5 times and then rolled in an excess of M9 at

665   room temperature for 20 minutes before being washed an additional time in M9 to clear
666   bacteria from the skin and gut. To circumvent worms sticking to the side of the conical
667   tube, tubes were briefly vortexed between spins as needed.
668
Washed animals were transferred to peptone-free plates (recipe from (Eustice et al.,
2022)) and returned to 25°C to continue to develop. Imaging was done immediately
after animals were washed off the food plates (baseline), and parallel populations were
kept at 25°C until the plate visibly had >50% dauers (the rest being pre-dauers; these
strains do not progress past dauer at 25°C). Animals that appeared to be dauers under
a dissecting scope were picked to image for the dauer endpoint of the experiment; only
animals with dauer morphology (recognized by radial constriction and the presence of
alae) were measured for germ cell number. While removal of food in the L1 stage prior
to the L1-L2d transition caused L1 developmental arrest, as expected, we found that
removal of food after the L2d molt did not inhibit *daf-2(e1370)* or *daf-7(e1372)* animals
from progressing through L2d or entering dauer.
680
At this temperature, these strains experience a ~100% developmental arrest in the
dauer stage and will not molt beyond dauer unless returned to lower temperatures
(Gems et al., 1998; Karp, 2018; Kenyon et al., 1993; Ren et al., 1996). We used this
endpoint rather than a set clock-time because the duration of the pre-dauer period of
daf-c mutants at 25°C is much longer than the duration of the L1 and L2 stages of fed
animals. The *daf-2(e1370)* mutant takes approximately 80 hours to enter dauer, and
daf-7(e1372) takes approximately 48 hours (Karp, 2018), compared to just over 24
hours for continuously fed animals to reach L3 (Byerly et al., 1976). The pre-dauer
period is also extended in wild-type animals that feed on a lower titre of bacterial food
(Cassada and Russell, 1975).
691
***glp-1(ts)* temperature shift experiments**
Temperature shift experiments used the temperature-sensitive allele of the Notch
receptor *glp-1(bn18)*. Loss of Notch signaling causes failure of germline induction (if the
temperature shift happens in the embryonic period) or irreversible loss of the stem-like
cell fate and meiotic entry of all germ cells (Fox and Schedl, 2015; Kodoyianni et al.,
1992). To test for *glp-1* dependence during dauer (Fig. 4A,B), two populations were
used. As a control, an unstarved, uncrowded population was reared at 16°C, shifted to
the restrictive temperature of 25°C for 24 hours, larval worms were singly picked to
NGM plates seeded with *E. coli* OP50, and these plates were kept at 16°C until
adulthood. The experimental group was made of worms that had been starved on plates
at 16°C to form dauers, which were shifted to 25°C for 24 hours. After 24 hours, dauers
were recovered by SDS isolation and singled to NGM plates seeded with *E. coli* OP50

and kept at 16°C until adulthood. Each plate was subsequently assayed for the
presence of progeny.
706
To test for when *glp-1* dependence returns after dauer (Fig. 4C,D), *glp-1(bn18)* worms
were put into dauer on starved plates at 16°C (n=14 first formed, and n=15 starved),
shifted to 25°C for a minimum of 12 hours, SDS isolated to trigger dauer exit, and
transferred to NGM plates with food and placed back at 25°C for an additional four, six,
or eight hours. At that time, the developmentally oldest/largest worms on the plate (the
worms that exited dauer first, since dauer exit is asynchronous) were picked as single
animals to NGM plates seeded with *E. coli* OP50 food and kept at 16°C until adulthood.
After four days of recovery at 16°C, each plate was assayed for live larval offspring, and
those that had offspring were scored as fertile. Parental worms on plates that lacked
larvae were imaged at 60x magnification and scored for the presence of embryos in the
uterus (fertile) or an absence of gametes (sterile). Embryo retention appeared to be a
phenotype of *glp-1(bn18)*.
719

Confocal imaging

All images were acquired at room temperature on a Leica DMI8 with an xLIGHT V3
confocal spinning disk head (89 North) with a 63× Plan-Apochromat (1.4 NA) objective
and an ORCAFusion GenIII sCMOS camera (Hamamatsu Photonics) controlled by
microManager. RFPs were excited with a 555 nm laser; GFP and mNG were excited
with a 488 nm laser; mTurquoise was excited with a 445 nm laser. Z-stacks through the
gonad were acquired with a z-step size of 0.3 or 0.5 μm as noted. Worms were
mounted on agar pads in M9 buffer with 0.01-0.02 M sodium azide paralytic (VWR
(Avantor) Catalog Number 26628-22-8). Samples used for fluorescence quantification
were acquired with the same laser power, exposure time, and z-step size within the
datasets to be compared, with histograms monitored to ensure proper exposure.
Images with poor body placement, debris on the slide, or other image quality concerns
were not analyzed.
733

Image analysis software

735　Images were processed in FIJI89 (Version: 2.14.1/1.54f).
736

Dauer recovery experiments

738　To measure fluorescent protein expression and germ cell proliferation during dauer
739　recovery, worms were isolated as described above (Dauer isolation protocol), and
740　subsequently split such that some worms were imaged immediately ("0 hours of
741　recovery", Figs 4,5), and other worms were recovered to *E. coli* OP50-seeded NGM
742　plates for the specified length of time.
743

Germer cell proliferation during dauer recovery

Germ cell proliferation during dauer recovery
Our otherwise wild-type marker control strain coexpressing *ina-1(qy23[ina-*
1::mNeonGreen]); naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR) II was put into dauer by
starvation on plates, and dauers were isolated as above (Dauer isolation protocol).
Worms were imaged at that 0 h starting point, and parallel populations were recovered
to plates seeded with *E. coli* OP50 bacterial food. Worms were imaged at four and eight
hours after plating on food. Germ cells were counted manually and scored for mitotic
figures (metaphase and anaphase chromatin condensations visible in the H2B::mCherry
signal, Fig. 4G. We plotted germ cell numbers over time and the percentage of worms
for which we observed mitotic figures in Fig. 4F. Mitotic indices were calculated as
number of mitotic cells/total cells for each specimen as in (Roy et al., 2016) and
averages for each timepoint are reported in the text.
756
Fluorescence intensity measurement of reporters of *lag-2*: Endogenously tagged
LAG-2::mNG with coexpressed membrane TdTomato
Strain NK2517 *qIs154(lag-2p:: myr::tdTomato); lag-2(cp193[lag-2::*
mNeonGreenˆ3xFlag]) V was used to measure endogenously tagged LAG-2::mNG
protein (Gordon et al., 2019) and a coexpressed multicopy integrated transcriptional
reporter of the *lag-2* ~3 kb upstream promoter with an *unc-54* 3' UTR (Byrd et al., 2014).
All worms, including controls, were reared at 25°C. TdTomato has a half-time to
maturation of one hour at 37°C (Shaner et al., 2004), while mNeonGreen has a half-
time to maturation of fewer than ten minutes at 37°C (Shaner et al., 2013) (both will fold
slower at worm-rearing temperatures). TdTomato signal is visible in adjacent neurons
(Fig. 3 and Fig. 5) and the autofluorescent gut granules of the worm are visible in the
channel of the endogenously tagged LAG-2::mNG. These structures were avoided as
much as possible during projection, tracing, and measurement of signal.
770
771 We chose an L2 control for this experiment because, while the dauer larva is an
772 alternative third larval stage, the worm's reproductive system undergoes morphogenetic
773 changes and changes in cell number during the normal L3 stage. These developmental
774 events are arrested in dauer (Tenen and Greenwald, 2019), making L3 a confounding
775 control for the dauer larvae reproductive system.
776
Confocal z-stacks were acquired with a 0.3 µm step size. A sum intensity z-projection
was created in FIJI consisting of eleven slices representing the superficial half of the cell
(from the superficial cell surface to the cross-section of the nucleus, which is visible as a
dark void in the center of the cell body). Usually one DTC from each worm was
analyzed, as the gut often obscures one. The DTC was hand-traced on the membrane
marker RFP channel; that ROI was subsequently used to measure both the RFP
(membrane) and GFP (endogenously tagged protein) channels, and slid off the DTC to

the adjacent body of the worm to measure the background for both channels.
Background subtracted Integrated Density measurements were thus obtained. We then
normalized all integrated density measurements to the mean integrated density at the 0
h timepoint for that fluorescent protein, so fold-change values and statistics are shown
in Fig. 5A,B,D,E, with raw values shown in Fig. S4. Kruskal-Wallis test with follow up
Dunn's correction for multiple comparisons of the raw, non-normalized integrated
densities show the same pattern of significance (that is, only LAG-2::mNG signal
increases relative to baseline) as the fold change analysis (Fig. S4). Integrated density
was used for this measure instead of raw integrated density so that size differences in
the DTC did not influence our measurements, as we noted that dauer DTCs are smaller
than non-dauer L2 DTCs (Fig. S1I-J). To display the ~40x range of LAG-2::mNG
intensity with the same image scaling, some pixels are saturated in display images
(Fig.5C). Saturation was not observed in the raw images from which measurements
were made (Fig. S5).
798
Fluorescence intensity measurement of reporters of *lag-2*: Polycistronic histone
reporter with coexpressed membrane mNeonGreen
We considered the ~3 kb upstream *lag-2* promoter fragment used in the transcriptional
reporter transgenes could insufficiently capture endogenous transcriptional regulation,
or that post-transcriptional regulation of *lag-2* mRNA could uniquely affect LAG-2
production. The membrane-localized transcriptional reporters that we use from (Byrd et
al., 2014; Linden et al., 2017) like those of (Blelloch et al., 1999; Henderson et al., 1994;
Pekar et al., 2017) are driven by ~3 kb upstream of the *lag-2* transcription start site
(including the TGF-β-responsive element described by (Pekar et al., 2017). This ~3 kb
element is sufficient to drive expression in DTCs, however (Karp and Greenwald, 2013)
identified regulatory elements of *lag-2* as far away as ~6.4 kb upstream, raising the
possibility that ~3 kb *lag-2* reporters are missing relevant regulatory information that is
important for DTC expression under certain, previously unexamined conditions.
812
We therefore examined a histone H2B::mTurquoise2 (half-time to maturation of 33.5
minutes, (Goedhart et al., 2012) that was knocked into the endogenous *lag-2* locus with
a P2A peptide (Medwig-Kinney et al., 2022). This element will generate a polycistronic
mRNA, the translation of which will produce one LAG-2 protein and one
H2B::mTurquoise2, meaning that the histone reporter is under the same transcriptional
and post-transcriptional, transcript-based regulation (e.g. by 3' UTR-mediated
repression by microRNAs or RNA binding proteins, transcript decay, etc.) as the
endogenous *lag-2* gene. It is coexpressed with another DTC-expressed transcriptional
reporter, a faster-folding, membrane-localized *lag-2p::mNeonGreen::PLCδ1[PH]* driven by
the ~3 kb upstream *lag-2* promoter with a *let-858* 3' terminator sequence, integrated into

the MoscI site on Ch. II (*cpIs122, (Linden et al., 2017)*). This strain was previously
analyzed in (Singh et al., 2024) and (Li and Gordon, 2025).
825
Confocal z-stacks were acquired with a 0.3 μm step size. A sum intensity z-projection
through a 6-7 μm depth of the DTC (to capture the entire nucleus in z, ~20 slices) was
generated in FIJI, the DTC was hand-traced, and membrane fluorescence was
measured for the GFP channel in that ROI. A background measurement was made by
sliding the DTC ROI onto the adjacent body of the worm. The nuclear histone signal
was obtained using an ellipse tool tight around the nucleus in the same z-projection to
measure on the CFP channel, and background for the nuclear signal was made by
sliding that ROI onto the adjacent body of the worm. Background subtracted Integrated
Density measurements were thus obtained. We then normalized all integrated density
measurements to the mean integrated density at the 0 h timepoint for that fluorescent
protein, so fold-change values and statistics are shown in Fig. 5A,B,D,E, with raw
values shown in Fig. S4. Kruskal-Wallis test with follow up Dunn's correction for multiple
comparisons of the raw, non-normalized integrated densities show the same pattern of
significance (that is, no pairwise difference relative to baseline for either fluorescent
protein) as the fold change analysis (Fig. S4).
841
To test whether perdurance of H2B::mTurquoise2 could mask the addition of new
protein upon transcriptional upregulation (Fig. 5D), we examined its dynamics in the Z1
and Z4 lineages in early larval stages (Fig. S7). Late L1 and early L2 worms (12 h after
release from L1 arrest at 25°C), and later L2 (15 h) worms were imaged as above and
sum projections were made through ~3.3 microns (eleven slices) of the superficial
gonad arm (and for one sample, for both arms). This captured the DTC (Z1.aa or Z4.pp)
and its sister cell (Z1.ap or Z4.pa), or else caught the parent of these cells (Z1.a or
Z4.p) right before it divides. We did not note any systematic difference between Z1 and
Z4-lineage cells for expression of this marker despite notably different membrane
fluorescence in these cells driven by the ~3 kb upstream *lag-2* promoter (Singh et al.,
2025). For each gonad arm, we used an elliptical ROI to measure H2B::mTurquoise2
signal in the nuclei of these cells, as well as a background region in the middle of the
gonad where no nucleus expresses the marker. The nucleus of the sister cell was
identified for measurement as a void in the GFP channel if no histone signal could
easily be noted. Background-subtracted measurements were made and compared to
quantify perdurance of the fluorescent histone signal.
858
lag-2 RNAi
To verify that the H2B::mTurquoise2 is indeed coregulated at the transcript level with
endogenous *lag-2*, we measured the effect of *lag-2* RNAi on H2B::mTurquoise2 signal
(Fig. S6). Single-colony overnight cultures were made of *E. coli* HT115 containing either

the empty vector L4440 or clone Y73C8B.4 from the Ahringer library (Kamath and
Ahringer, 2003; Kamath et al., 2003), which is complementary to >900 nt of the *lag-2*
mRNA extending from just upstream of the start codon through exon 1, intron 1, and
some of exon 2. Cultures contained ampicillin (100 µg/mL, VWR (Avantor), Catalog no.
76204-346) and were grown at 37°C, expression was induced with 1 mM IPTG (Apex
BioResearch Products, cat# 20-109) for 1 h at 37°C, and the culture was plated and
allowed to grow at least overnight at room temperature on NGM plates prepared with
IPTG and ampicillin. L2 larval stage worms were picked onto NGM plates without food
and allowed to move around for at least 30 minutes to prevent bacterial carryover.
Animals were then hand-picked either onto *lag-2* RNAi or empty L4440 vector control
plates, cultured for 24 h at 16°C to the L4 stage, imaged, and background-subtracted
fluorescence intensity was measured as above.
875
Somatic gonad marker measurements
Somatic gonad cell number (Fig. S1G): The strain GS9692 expressing *arTi435(rps-*
27p::2xnls::gfp(flexon)::unc-54 3′ UTR); *arTi237(ckb-3p::Cre(opti)::tbb-2 3′ UTR)*
(Shaffer and Greenwald, 2022) was specifically designed to activate enduring GFP
expression in the cells of the somatic gonad. We manually counted gonad cells in this
strain after inducing first-formed and starved plate dauers (Fig. S1F-H). We always see
the expected two DTCs and 10 gonad blast cells in dauers of any age. Variation in
expression levels/tissue depth can sometimes complicate detection of the cell at the
center of the gonad primordium (more orange cells in Fig. S1G). Landmarks were
selected based on relative position with DTCs at the gonad tips and the next-most-distal
fluorescent cells being the SS cells used for measuring.
887
888   **Statistical analysis**
889   All statistical analysis was done using GraphPad Prism version 10.4.2 for Windows,
890   GraphPad Software, Boston, Massachusetts USA, www.graphpad.com. All tests and
891   sample sizes are specified in the relevant figure legends.
892
Before any statistical tests were conducted on data, the data was first evaluated for
normal distribution using the Shapiro-Wilk test for normality. Statistical tests were
selected based on the appropriate test given the normality or non-normality of the data.
If the data were normally distributed, Welch's t-test (two groups) or Welch's ANOVA
(three or more groups) was used because it is more robust against Type I error rates
than Student's t-test or an ordinary one-way ANOVA (Delacre et al., 2019; Derrick et al.,
2016). When data violated the assumption of normality, non-parametric Mann-Whitney
(two groups) or Kruskal-Wallis (three or more groups) tests were used. Post-hoc
analyses for pairwise comparisons were selected based on appropriateness for the test
used.

For curve fitting in Figs 3M, 4E, 5A, S2B,C, nonlinear regression models were
compared to null-hypothesis linear regression models to test which was a better fit.
Curves were fit to the means of the data at each timepoint. For exponential decay of
LAG-2 protein, the plateau of the model was not constrained.
907

Author Contributions
Conceptualization: F.A.K. and K.L.G.; Formal analysis: F.A.K.; Investigation: F.A.K.,
C.P.M., Data collection assistance: B.K.; Writing – original draft preparation: F.A.K. and
K.L.G.; Writing – review and editing: F.A.K., C.P.M., and K.L.G.; Visualization: F.A.K.;
Supervision: K.L.G.; Project administration: K.L.G.; Funding acquisition: F.A.K. and
K.L.G.
914

915 **Acknowledgements**
916 We thank members of the Gordon Lab and F.A.K. dissertation committee members
917 R.H.D. and D.R.M. for thoughtful comments on the manuscript.
918

Funding
Some strains were provided by the CGC, which is funded by National Institutes of
Health Office of Research Infrastructure Programs (P40 OD010440) and are to be
requested directly from CGC. Research reported in this publication was supported by
the National Institute Of General Medical Sciences of the National Institutes of Health
under Award Number R35GM147704 to K.L.G. and NIGMS Diversity Supplement
3R35GM147704-02S1 to support the work of F.A.K. Support was also provided by the
National Science Foundation under NSF CAREER 2442303. The content is solely the
responsibility of the authors and does not necessarily represent the official views of the
National Institutes of Health or the National Science Foundation.
929

930 **Data and resource availability**
931 All relevant data and details of resources can be found within the article. All *C. elegans*
932 strains are available upon request, except those sourced from the CGC (above).
933

Conflicts of Interest
The authors declare no conflicts of interest.

FIGURE LEGENDS
Figure 1: Dauer larvae that form after severe starvation have smaller gonads and
recover to have smaller broods than dauer larvae that form after feeding.
(A) Representative images of dauers induced with given methods. Gonads outlined.
(B) Germ cell numbers in dauers of marker control strain induced by different methods.
Starved plate (n=59), liquid culture (n=46), high density (n=61), and first-formed (n=24).
Kruskal-Wallis test statistic H=111.0, p<0.0001.
(C) Brood size of marker control strain. Never-dauer control (n=15), recovered dauers
from starved plate (n=17), high-density plating (n=19), and first-formed (n=21). Kruskal-
Wallis test statistic H=27.19, p<0.0001.
(D-E) Representative fluorescence images of germ cells in dauers induced with given
methods for (D) *daf-2(e1370)* and (E) *daf-7(e1372)* expressing germ cell nuclear
marker. Gonads outlined.
(F) Germ cell numbers in *daf-2(e1370)* and *daf-7(e1372)* dauers induced with four
methods. Starved plate (n_{daf-2}=52, n_{daf-7}=47), high density (n_{daf-2}=37, n_{daf-7}=21), first-
formed (n_{daf-2}=19, n_{daf-7}=22), and constitutive (n_{daf-2}=33, n_{daf-7}=27). Kruskal-Wallis test
statistic H=175.3 p<0.0001.
(G) Brood size of *daf-2(e1370);naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR)*. Never-
dauer control (n=14), recovered dauers from starved plate (n=17), high density plating
(n=15), first-formed (n=14), and constitutive late (n=12) recovered animals. Kruskal-
Wallis test statistic=30.51, p<0.0001.
All scale bars 10 μm. Dunn's correction for multiple comparisons was used post-hoc to
determine statistical significance of pairwise differences for relevant comparisons,
asterisks on graph indicate these at p<0.0001 ****; p<0.005 **. Full results of statistical
analyses in Supplemental Table S1.
961
Figure 2: Withdrawal of food in the pre-dauer period arrests germ cell
proliferation after a single round of division.
(A) Schematic illustrating food removal experiments. Worms moved off food (dashed
lines, imaged "Baseline" (2B-D)) and reared until dauer entry (stars, imaged "Dauer"
(2B-D). All experiments in Fig. 2 were conducted at 25°C. Both strains express germ
cell nuclear marker.
(B) Results of experiment detailed in (2A), showing number of germ cell nuclei in *daf-*
2(e1370). Sample sizes: 12 h fed $n_{baseline}$=31, n_{dauer}=22; 16 h fed $n_{baseline}$=16, n_{dauer}=8;
20 h fed $n_{baseline}$=18, n_{dauer}=13; 24 h $n_{baseline}$=23, n_{dauer}=24; continuously fed dauer n=33.
(C) Results of experiment detailed in (2A), showing number of germ cell nuclei in *daf-*
7(e1372). Sample sizes: 12 h fed $n_{baseline}$=20, n_{dauer}=17; 16 h fed $n_{baseline}$=20, n_{dauer}
n=17; 20 h fed $n_{baseline}$=20, n_{dauer}=22; 24 h fed $n_{baseline}$=18, n_{dauer}=12; continuously fed
dauer n=26.

(D) Representative images of *daf-2(e1370)* measured for (2B) at time of food removal
(baseline, left) and in dauer (right). Gonad outlined in white. *daf-7(e1372)* shown in Fig.
S2. All scale bars 10 µm.
(E) Experiment with *daf-2(e1370)* as in (2A) with additional +24 h endpoint. Darker dots
indicate animals in dauer diapause at the time of counting (see 2E). $n_{12\,h\,fed}=13$; $n_{12\,h\,fed,\,24\,h\,starved}=16$; $n_{12\,h\,fed,\,24\,h\,fed}=14$, refers to animals that were washed at 12 h and returned
to food. $n_{12\,h\,fed,\,72\,h\,starved}=16$. Kruskal-Wallis test statistic=39.12; p<0.0001 with
significance from Dunn's correction for multiple comparisons indicated as p<0.0001 ****;
p<0.005 **.
(F) Percentage of worms that had reached the dauer stage under each condition in 2D.
(E-F) Grey shaded box marks the samples quantified in parallel after 24 hours under
experimental conditions, both on and off food.
987
Figure 3: LAG-2 expression is higher in fed animals than in dauer and drops
rapidly upon food removal in L2d, with *daf-2* and *daf-7* required for the former but
not the latter.
(A-B) Sum projections through eleven 0.3 µm z-slices of the superficial surface of a
DTC expressing endogenously tagged LAG-2 protein (fire lookup table, *lag-2(cp193[lag-*
2:: mNeonGreen])) and a membrane-localized transgenic reporter (grayscale,
qIs154(lag-2p:: myr::tdTomato)) in fed L2 controls (n=19) and starved plate dauers
(n=21). Each fluorescence channel is shown with identical scaling between treatments.
(C-D) Integrated density measured for DTCs described in (A-B). (C) LAG-2::mNG, note
log_{10} scale Mann-Whitney test statistic U=0, p<0.0001, and (D) *qIs154(lag-2p::*
myr::tdTomato). Unpaired t-test with Welch's correction t=3.491, p=0.0012.
(E-G) Representative images of the transgenes in (A) in fed L2 worms that are (E)
otherwise wild-type (n=19), (F) *daf-2(e1370)* mutants (n=20), (G) *daf-7(e1372)* mutants
(n=18).
(H-I) Integrated density measured for DTCs with sample sizes given in (E-G). (H) LAG-
2::mNG, note log_{10} scale. Welch's ANOVA test statistic W=89.99 (2.000, 18.76)
p<0.0001 with Dunnett's T3 multiple comparisons, and (I) *qIs154(lag-2p::*
myr::tdTomato). Otherwise wild-type control, same L2 dataset as in (B-C). Welch's
ANOVA test statistic W=139.9 (2.000, 29.43) p<0.0001 with Dunnett's T3 multiple
comparisons.
(J) Schematic of experiment performed for (K-M). Worms carrying the transgenes in the
daf-2(e1370) and *daf-7(e1372)* mutant backgrounds were raised on food at 25°C for 20
h, imaged, moved off of food, and imaged one and two hours later.
(K-L) Representative images of worms described in (J). Sum projection through eleven
0.3 µm z-slices of the superficial surface of the DTC at timepoints shown above for (K)
daf-2(e1370) ($n_{20\,h\,fed}=9$, $n_{+1\,h\,starved}=16$, $n_{+2\,h\,starved}=21$) and (L) *daf-7(e1372)* mutants (n_{20}

h fe=16, n+1 h starved=18, n+2 h starved=20) coexpressing endogenously tagged LAG-2 protein
(fire lookup table) with DTC membrane-localized transcriptional reporter (grayscale).
(M) Integrated density measured for LAG-2::mNG in DTCs with sample sizes given in K-
L. Kruskal-Wallis test statistic H=53.78, p<0.0001 with Dunn's correction for multiple
comparisons; pairwise comparisons were made between 20 h fed and the two food-
removal timepoints per genotype. Regression and R^2 values displayed on graph.
All scale bars 10 µm. DTCs outlined in yellow. Significance of pairwise comparisons
after post-hoc corrections indicated as p<0.0001 ****; p<0.001 ***; p<0.01 **.
1022
Figure 4: Dauer germ cells are arrested in a Notch-independent state and rapidly
regain Notch dependence and cell cycling upon dauer exit
(A) Schematic illustrating temperature shift experiment testing germline *glp-1*/Notch
dependence in dauer. (B) Worm fertility after (A). Control (never-dauer population temp
shift in parallel with dauer) n=28, Recovered dauers n=29.
(C) Schematic of dauer recovery *glp-1(bn18)* temperature-shift experiment testing the
return of Notch dependence to the germ line. (D) Percentage of animals recovering as
fertile vs. sterile results after recovery from dauer at 25°C for the first 4 h (n=15), 6 h
(n=15) and 8 h (n=15) of dauer recovery.
(E) Number of germ cells after specified hours of dauer recovery in the marker control
strain n0 h=35, n4 h=29, n8 h=34.
(F) Number of mitotic figures observed in each sample in E.
(G) Representative images of gonads at 0 h, 4 h, and 8 h of recovery. Note 8 h sample
has large and small nuclei typical of cells just before and after division.
1037
Figure 5: Recovery from dauer triggers a burst of LAG-2 presentation on the DTC
niche without a coincident rise in *lag-2* transcription
(A-D) An otherwise wild-type strain coexpressing the endogenously tagged LAG-2
protein *lag-2(cp193[lag-2:: mNeonGreen])* (green in C) with a membrane-localized
transgenic reporter *qIs154(lag-2p:: myr::tdTomato)* (magenta in C) recovering from
dauer.
(A) Fold change of LAG-2::mNG relative to mean at 0 h (n0 h =30; n2 h =29; n4 h=18; n8
h=20). Note Y-axis scale compared to other graphs. Kruskal-Wallis test statistic
H=67.26, p<0.0001. Dunn's correction for multiple comparisons show all pairwise
comparisons are significantly different.
(B) Fold change of the *lag-2p::myrTdTomato* transcriptional reporter for samples in A.
Kruskal-Wallis test statistic H=8.905, p=0.0306.
(C) Sum projection through eleven 0.3 µm z-slices of the superficial surface of a DTC
for the time points shown above. Each fluorescence channel is shown with identical
scaling between time points.

(D-F) Otherwise wild-type strain co-expressing a *lag-2(bmd202 [lag-*
2::P2A::H2B::mTurquoise2]) polycistronic histone reporter knocked into the endogenous
lag-2 locus (blue in F) with a *lag-2p::mNeonGreen::PLCδ1^{PH}* membrane-localized
transcriptional reporter (yellow in F), same treatment as A-C.
(D) Fold change of mTurquoise2 relative to mean at 0 h ($n_{0\,h}$=29; $n_{2\,h}$=13; $n_{4\,h}$=10; n_8
$_h$=19). Kruskal-Wallis test statistic H=13.30m, p=0.0040, with Dunn's correction for
multiple comparisons showing that none of the recovery timepoints is significantly
different from 0 h baseline.
(E) Fold change of *lag-2p::mNeonGreen::PLCδ1^{PH}* transcriptional reporter relative to
mean at 0h ($n_{0\,h}$=29; $n_{2\,h}$=13; $n_{4\,h}$=10; $n_{8\,h}$=19). Kruskal-Wallis test statistic H=17.41,
p=0.0006, with Dunn's correction for multiple comparisons showing that none of the
recovery timepoints is significantly different from 0 h baseline.
(F) Sum projections through 0.3 μm Z-slices from superficial surface of DTC to bottom
of nucleus.
Non-normalized fluorescence measures and statistical analyses shown in Fig. S4.
DTCs outlined. All scale bars 10 μm.
1069
Supplemental Material
Table S1. Statistical analyses for Fig. 1.
Figure S1. Somatic gonad cell number is unaffected by method of dauer induction.
Figure S2. *daf-7* and *daf-2* are dispensable for nutritionally-sensitive exponential gonad
growth before dauer.
Figure S3. Expression dynamics of *lag-2* transcriptional reporter and LAG-2::mNG
protein vary across stages in *daf-2(e1370)* and *daf-7(e1372)* mutants.
Figure S4. Non-normalized expression of LAG-2::mNG and *lag-2* reporters during
dauer recovery show no increase in transcriptional activity.
Figure S5. Image scaling for display of qualitative comparison creates image saturation
that is not present in the original data.
Figure S6. Knockdown of *lag-2(bmd202[lag-2::P2A::H2B::mTurquoise2])* signal by *lag-*
2 RNAi verifies H2B::mTurquoise2 transcript-based coregulation with *lag-2.*
Figure S7. Signal decay of *lag-2(bmd202 [lag-2::P2A::H2B::mTurquoise2])* reveals
effective detection of continuous vs. previous expression.

References:

Abuetabh, Y., Wu, H.H., Chai, C., Al Yousef, H., Persad, S., Sergi, C.M., Leng, R., 2022. DNA damage response revisited: the p53 family and its regulators provide endless cancer therapy opportunities. Exp. Mol. Med. 54, 1658–1669. https://doi.org/10.1038/s12276-022-00863-4.

Ailion, M., Thomas, J.H., 2000. Dauer formation induced by high temperatures in Caenorhabditis elegans. Genetics 156, 1047–1067. https://doi.org/10.1093/genetics/156.3.1047.

Angelo, G., Van Gilst, M.R., 2009. Starvation protects germline stem cells and extends reproductive longevity in C. elegans. Science 326, 954–958. https://doi.org/10.1126/science.1178343.

Aprison, E.Z., Dzitoyeva, S., Ruvinsky, I., 2024. The roles of TGFβ and serotonin signaling in regulating proliferation of oocyte precursors and germline aging. bioRxiv.

Austin, J., Kimble, J, 1989. Transcript analysis of glp-1 and lin-12, homologous genes required for cell interactions during development of C. elegans. Cell 58, 565–571. https://doi.org/10.1016/0092-8674(89)90437-6.

Austin, J., Kimble, J, 1987. glp-1 is required in the germ line for regulation of the decision between mitosis and meiosis in C. elegans. Cell 51, 589–599. https://doi.org/10.1016/0092-8674(87)90128-0.

Barbieri, M., Bonafè, M., Franceschi, C., Paolisso, G., 2003. Insulin/IGF-I-signaling pathway: an evolutionarily conserved mechanism of longevity from yeast to humans. Am. J. Physiol. Endocrinol. Metab. 285, E1064-71. https://doi.org/10.1152/ajpendo.00296.2003.

Baugh, L.R., Hu, P.J., 2020. Starvation responses throughout the caenorhabditiselegans life cycle. Genetics 216, 837–878. https://doi.org/10.1534/genetics.120.303565.

Baugh, L.R., 2013. To grow or not to grow: nutritional control of development during Caenorhabditis elegans L1 arrest. Genetics 194, 539–555. https://doi.org/10.1534/genetics.113.150847.

Belew, M.D., Chien, E., Wong, M., Michael, W.M., 2021. A global chromatin compaction pathway that represses germline gene expression during starvation. J. Cell Biol. 220. https://doi.org/10.1083/jcb.202009197.

Blelloch, R., Anna-Arriola, S.S., Gao, D., Li, Y., Hodgkin, J., Kimble, J., 1999. The gon-1 gene is required for gonadal morphogenesis in Caenorhabditis elegans. Dev. Biol. 216, 382–393. https://doi.org/10.1006/dbio.1999.9491.

Brenner, S., 1974. THE GENETICS OF *caenorhabditis elegans*. Genetics 77, 71–94. https://doi.org/10.1093/genetics/77.1.71.

Byerly, L., Cassada, R.C., Russell, R.L., 1976. The life cycle of the nematode Caenorhabditis elegans. I. Wild-type growth and reproduction. Dev. Biol. 51, 23–33. https://doi.org/10.1016/0012-1606(76)90119-6.

Byrd, D.T., Knobel, K., Affeldt, K., Crittenden, S.L., Kimble, J., 2014. A DTC niche plexus surrounds the germline stem cell pool in Caenorhabditis elegans. PLoS ONE 9, e88372. https://doi.org/10.1371/journal.pone.0088372.

Cassada, R.C., Russell, R.L., 1975. The dauerlarva, a post-embryonic developmental variant of the nematode Caenorhabditis elegans. Dev. Biol. 46, 326–342.

https://doi.org/10.1016/0012-1606(75)90109-8.

Chaikovsky, A.C., Li, C., Jeng, E.E., Loebell, S., Lee, M.C., Murray, C.W., Cheng, R., Demeter,
J., Swaney, D.L., Chen, S.H., Newton, B.W., Johnson, J.R., Drainas, A.P., Shue, Y.T.,
Seoane, J.A., Srinivasan, P., He, A., Yoshida, A., Hipkins, S.Q., McCrea, E., Sage, J.,
2021. The AMBRA1 E3 ligase adaptor regulates the stability of cyclin D. Nature 592, 794–
798. https://doi.org/10.1038/s41586-021-03474-7.

Chi, C., Ronai, D., Than, M.T., Walker, C.J., Sewell, A.K., Han, M., 2016. Nucleotide levels
regulate germline proliferation through modulating GLP-1/Notch signaling in C. elegans.
Genes Dev. 30, 307–320. https://doi.org/10.1101/gad.275107.115.

Cinquin, O., Crittenden, S.L., Morgan, D.E., Kimble, J., 2010. Progression from a stem cell-like
state to early differentiation in the C. elegans germ line. Proc Natl Acad Sci USA 107,
2048–2053. https://doi.org/10.1073/pnas.0912704107.

Claeys, I., Simonet, G., Poels, J., Van Loy, T., Vercammen, L., De Loof, A., Vanden Broeck, J.,
2002. Insulin-related peptides and their conserved signal transduction pathway. Peptides
23, 807–816. https://doi.org/10.1016/s0196-9781(01)00666-0.

Corsi, A.K., Wightman, B., Chalfie, M., 2015. A Transparent Window into Biology: A Primer on
Caenorhabditis elegans. Genetics 200, 387–407.
https://doi.org/10.1534/genetics.115.176099.

Dalfó, D., Michaelson, D., Hubbard, E.J.A., 2012. Sensory regulation of the C. elegans germline
through TGF-β-dependent signaling in the niche. Curr. Biol. 22, 712–719.
https://doi.org/10.1016/j.cub.2012.02.064.

Das, D., Arur, S., 2017. Conserved insulin signaling in the regulation of oocyte growth,
development, and maturation. Mol. Reprod. Dev. 84, 444–459.
https://doi.org/10.1002/mrd.22806.

Delacre, M., Leys, C., Mora, Y.L., Lakens, D., 2019. Taking Parametric Assumptions Seriously:
Arguments for the Use of Welch's F-test instead of the Classical F-test in One-Way
ANOVA. rips 32, 13. https://doi.org/10.5334/irsp.198.

Derrick, B., Toher, D., White, P., 2016. Why Welch's test is Type I error robust. TQMP 12, 30–
38. https://doi.org/10.20982/tqmp.12.1.p030.

de Lima, J.G.S., Lanza, D.C.F., 2021. 2A and 2A-like Sequences: Distribution in Different Virus
Species and Applications in Biotechnology. Viruses 13. https://doi.org/10.3390/v13112160.

Diehl, F.F., Miettinen, T.P., Elbashir, R., Nabel, C.S., Darnell, A.M., Do, B.T., Manalis, S.R.,
Lewis, C.A., Vander Heiden, M.G., 2022. Nucleotide imbalance decouples cell growth from
cell proliferation. Nat. Cell Biol. 24, 1252–1264. https://doi.org/10.1038/s41556-022-00965-
1.

Donnelly, M.L.L., Luke, G., Mehrotra, A., Li, X., Hughes, L.E., Gani, D., Ryan, M.D., 2001.
Analysis of the aphthovirus 2A/2B polyprotein "cleavage" mechanism indicates not a
proteolytic reaction, but a novel translational effect: a putative ribosomal "skip". J. Gen.
Virol. 82, 1013–1025. https://doi.org/10.1099/0022-1317-82-5-1013.

Erkut, C., Kurzchalia, T.V., 2015. The C. elegans dauer larva as a paradigm to study metabolic
suppression and desiccation tolerance. Planta 242, 389–396.
https://doi.org/10.1007/s00425-015-2300-x.

Eustice, M., Konzman, D., Reece, J.M., Ghosh, S., Alston, J., Hansen, T., Golden, A., Bond,
M.R., Abramowitz, L.K., Hanover, J.A., 2022. Nutrient sensing pathways regulating adult

reproductive diapause in C. elegans. PLoS ONE 17, e0274076.

https://doi.org/10.1371/journal.pone.0274076.

Ewald, C.Y., Castillo-Quan, J.I., Blackwell, T.K., 2018. Untangling Longevity, Dauer, and

Healthspan in Caenorhabditis elegans Insulin/IGF-1-Signalling. Gerontology 64, 96–104.

https://doi.org/10.1159/000480504.

Flick, K., Kaiser, P., 2012. Protein degradation and the stress response. Semin. Cell Dev. Biol.

23, 515–522. https://doi.org/10.1016/j.semcdb.2012.01.019.

Fortini, M.E., 2009. Notch signaling: the core pathway and its posttranslational regulation. Dev.

Cell 16, 633–647. https://doi.org/10.1016/j.devcel.2009.03.010.

Fox, P.M., Schedl, T., 2015. Analysis of Germline Stem Cell Differentiation Following Loss of

GLP-1 Notch Activity in Caenorhabditis elegans. Genetics 201, 167–184.

https://doi.org/10.1534/genetics.115.178061.

Gems, D., Sutton, A.J., Sundermeyer, M.L., Albert, P.S., King, K.V., Edgley, M.L., Larsen, P.L.,

Riddle, D.L., 1998. Two pleiotropic classes of daf-2 mutation affect larval arrest, adult

behavior, reproduction and longevity in Caenorhabditis elegans. Genetics 150, 129–155.

https://doi.org/10.1093/genetics/150.1.129.

Gimond, C., Poullet, N., Vielle, A., Demoinet, E., Braendle, C., 2025. Developmental plasticity of

hermaphrodite sperm production across environments in Caenorhabditis elegans. PLoS

ONE 20, e0336162. https://doi.org/10.1371/journal.pone.0336162.

Goedhart, J., von Stetten, D., Noirclerc-Savoye, M., Lelimousin, M., Joosen, L., Hink, M.A., van

Weeren, L., Gadella, T.W.J., Royant, A., 2012. Structure-guided evolution of cyan

fluorescent proteins towards a quantum yield of 93%. Nat. Commun. 3, 751.

https://doi.org/10.1038/ncomms1738.

Golden, J.W., Riddle, D.L., 1984. The Caenorhabditis elegans dauer larva: developmental

effects of pheromone, food, and temperature. Dev. Biol. 102, 368–378.

https://doi.org/10.1016/0012-1606(84)90201-X.

Gordon, K.L., Payne, S.G., Linden-High, L.M., Pani, A.M., Goldstein, B., Hubbard, E.J.A.,

Sherwood, D.R., 2019. Ectopic Germ Cells Can Induce Niche-like Enwrapment by

Neighboring Body Wall Muscle. Curr. Biol. 29, 823-833.e5.

https://doi.org/10.1016/j.cub.2019.01.056.

Gordon, K., 2020. Recent Advances in the Genetic, Anatomical, and Environmental Regulation

of the C. elegans Germ Line Progenitor Zone. J. Dev. Biol. 8.

https://doi.org/10.3390/jdb8030014.

Greer, E.R., Pérez, C.L., Van Gilst, M.R., Lee, B.H., Ashrafi, K., 2008. Neural and molecular

dissection of a C. elegans sensory circuit that regulates fat and feeding. Cell Metab. 8,

118–131. https://doi.org/10.1016/j.cmet.2008.06.005.

Hall, S.E., Beverly, M., Russ, C., Nusbaum, C., Sengupta, P., 2010. A cellular memory of

developmental history generates phenotypic diversity in C. elegans. Curr. Biol. 20, 149–

155. https://doi.org/10.1016/j.cub.2009.11.035.

Hand, S.C., Denlinger, D.L., Podrabsky, J.E., Roy, R., 2016. Mechanisms of animal diapause:

recent developments from nematodes, crustaceans, insects, and fish. Am. J. Physiol.

Regul. Integr. Comp. Physiol. 310, R1193-211. https://doi.org/10.1152/ajpregu.00250.2015.

Hansen, D., Schedl, T., 2006. The Regulatory Network Controlling the Proliferation–Meiotic

Entry Decision in the Caenorhabditis elegans Germ Line, in: Current Topics in

Developmental Biology. Elsevier, pp. 185–215. https://doi.org/10.1016/S0070-
2153(06)76006-9.
Harsimran Kaur Gill, Gaurav Goyal, Gurminder Chahil, 2017. Insect diapause: A review. JAST-
A 7. https://doi.org/10.17265/2161-6256/2017.07.002.
Henderson, S.T., Gao, D., Lambie, E.J., Kimble, J., 1994. lag-2 may encode a signaling ligand
for the GLP-1 and LIN-12 receptors of C. elegans. Development 120, 2913–2924.
https://doi.org/10.1242/dev.120.10.2913.
Hibshman, J.D., Webster, A.K., Baugh, L.R., 2021. Liquid-culture protocols for synchronous
starvation, growth, dauer formation, and dietary restriction of Caenorhabditis elegans.
STAR Protocols 2, 100276. https://doi.org/10.1016/j.xpro.2020.100276.
Hirsh, D., Oppenheim, D., Klass, M., 1976. Development of the reproductive system of
Caenorhabditis elegans. Dev. Biol. 49, 200–219. https://doi.org/10.1016/0012-
1606(76)90267-0.
Jayadev, R., Chi, Q., Keeley, D.P., Hastie, E.L., Kelley, L.C., Sherwood, D.R., 2019. α-Integrins
dictate distinct modes of type IV collagen recruitment to basement membranes. J. Cell Biol.
218, 3098–3116. https://doi.org/10.1083/jcb.201903124.
Johnson, T.E., Mitchell, D.H., Kline, S., Kemal, R., Foy, J., 1984. Arresting development arrests
aging in the nematode Caenorhabditis elegans. Mech. Ageing Dev. 28, 23–40.
https://doi.org/10.1016/0047-6374(84)90150-7.
Kamath, R.S., Ahringer, J., 2003. Genome-wide RNAi screening in Caenorhabditis elegans.
Methods 30, 313–321. https://doi.org/10.1016/s1046-2023(03)00050-1.
Kamath, R.S., Fraser, A.G., Dong, Y., Poulin, G., Durbin, R., Gotta, M., Kanapin, A., Le Bot, N.,
Moreno, S., Sohrmann, M., Welchman, D.P., Zipperlen, P., Ahringer, J., 2003. Systematic
functional analysis of the Caenorhabditis elegans genome using RNAi. Nature 421, 231–
237. https://doi.org/10.1038/nature01278.
Karp, X., Greenwald, I., 2013. Control of cell-fate plasticity and maintenance of multipotency by
DAF-16/FoxO in quiescent Caenorhabditis elegans. Proc Natl Acad Sci USA 110, 2181–
2186. https://doi.org/10.1073/pnas.1222377110.
Karp, X., 2018. Working with dauer larvae. WormBook 2018, 1–19.
https://doi.org/10.1895/wormbook.1.180.1.
Kenyon, C., Chang, J., Gensch, E., Rudner, A., Tabtiang, R., 1993. A C. elegans mutant that
lives twice as long as wild type. Nature 366, 461–464. https://doi.org/10.1038/366461a0.
Kimble, J., Crittenden, S.L., 2007. Controls of germline stem cells, entry into meiosis, and the
sperm/oocyte decision in Caenorhabditis elegans. Annu. Rev. Cell Dev. Biol. 23, 405–433.
https://doi.org/10.1146/annurev.cellbio.23.090506.123326.
Kimble, J., Crittenden, S.L., 2005. Germline proliferation and its control. WormBook 1–14.
https://doi.org/10.1895/wormbook.1.13.1.
Kimble, J., Hirsh, D., 1979. The postembryonic cell lineages of the hermaphrodite and male
gonads in Caenorhabditis elegans. Dev. Biol. 70, 396–417. https://doi.org/10.1016/0012-
1606(79)90035-6.
Kimble, J.E., White, J.G., 1981. On the control of germ cell development in Caenorhabditis
elegans. Dev. Biol. 81, 208–219. https://doi.org/10.1016/0012-1606(81)90284-0.
Kimura, K.D., Tissenbaum, H.A., Liu, Y., Ruvkun, G., 1997. daf-2, an insulin receptor-like gene
that regulates longevity and diapause in Caenorhabditis elegans. Science 277, 942–946.

https://doi.org/10.1126/science.277.5328.942.

Kim, S., Paik, Y.-K., 2008. Developmental and reproductive consequences of prolonged non-
aging dauer in Caenorhabditis elegans. Biochem. Biophys. Res. Commun. 368, 588–592.
https://doi.org/10.1016/j.bbrc.2008.01.131.

Klass, M., Hirsh, D., 1976. Non-ageing developmental variant of Caenorhabditis elegans.
Nature 260, 523–525. https://doi.org/10.1038/260523a0.

Kodoyianni, V., Maine, E.M., Kimble, J., 1992. Molecular basis of loss-of-function mutations in
the glp-1 gene of Caenorhabditis elegans. Mol. Biol. Cell 3, 1199–1213.
https://doi.org/10.1091/mbc.3.11.1199.

Linden, L.M., Gordon, K.L., Pani, A.M., Payne, S.G., Garde, A., Burkholder, D., Chi, Q.,
Goldstein, B., Sherwood, D.R., 2017. Identification of regulators of germ stem cell
enwrapment by its niche in C. elegans. Dev. Biol. 429, 271–284.
https://doi.org/10.1016/j.ydbio.2017.06.019.

Liu, Y., Beyer, A., Aebersold, R., 2016. On the Dependency of Cellular Protein Levels on mRNA
Abundance. Cell 165, 535–550. https://doi.org/10.1016/j.cell.2016.03.014.

Li, X., Gordon, K.L., 2025. MIG-21 interacts with Wnt and Netrin signaling in gonad migration in
C. elegans. PLoS Genet. 21, e1011866. https://doi.org/10.1371/journal.pgen.1011866.

Li, X., Singh, N., Miller, C., Washington, I., Sosseh, B., Gordon, K.L., 2022. The C. elegans
gonadal sheath Sh1 cells extend asymmetrically over a differentiating germ cell population
in the proliferative zone. eLife 11. https://doi.org/10.7554/eLife.75497.

Maiani, E., Milletti, G., Nazio, F., Holdgaard, S.G., Bartkova, J., Rizza, S., Cianfanelli, V.,
Lorente, M., Simoneschi, D., Di Marco, M., D'Acunzo, P., Di Leo, L., Rasmussen, R.,
Montagna, C., Raciti, M., De Stefanis, C., Gabicagogeascoa, E., Rona, G., Salvador, N.,
Pupo, E., Cecconi, F., 2021. AMBRA1 regulates cyclin D to guard S-phase entry and
genomic integrity. Nature 592, 799–803. https://doi.org/10.1038/s41586-021-03422-5.

McGehee, A., 2019. The GLR-1 phenotypes of the daf-7(e1372) allele are not temperature
sensitive. MicroPubl. Biol. 2019. https://doi.org/10.17912/micropub.biology.000158.

McIntyre, D.C., Nance, J., 2023. Niche cells regulate primordial germ cell quiescence in
response to basement membrane signaling. Development 150.
https://doi.org/10.1242/dev.201640.

McShane, E., Sin, C., Zauber, H., Wells, J.N., Donnelly, N., Wang, X., Hou, J., Chen, W.,
Storchova, Z., Marsh, J.A., Valleriani, A., Selbach, M., 2016. Kinetic Analysis of Protein
Stability Reveals Age-Dependent Degradation. Cell 167, 803-815.e21.
https://doi.org/10.1016/j.cell.2016.09.015.

Medwig-Kinney, T.N., Sirota, S.S., Gibney, T.V., Pani, A.M., Matus, D.Q., 2022. An in vivo
toolkit to visualize endogenous LAG-2/Delta and LIN-12/Notch signaling in C. elegans.
MicroPubl. Biol. 2022. https://doi.org/10.17912/micropub.biology.000602.

Morao, A.K., Ercan, S., 2021. Hatched and starved: Two chromatin compaction mechanisms
join forces to silence germ cell genome. J. Cell Biol. 220.
https://doi.org/10.1083/jcb.202107026.

Murphy, C.T., Hu, P.J., 2013. Insulin/insulin-like growth factor signaling in C. elegans.
WormBook 1–43. https://doi.org/10.1895/wormbook.1.164.1.

Narbonne, P., Roy, R., 2006. Inhibition of germline proliferation during C. elegans dauer
development requires PTEN, LKB1 and AMPK signalling. Development 133, 611–619.

https://doi.org/10.1242/dev.02232.

Ouellet, J., Li, S., Roy, R., 2008. Notch signalling is required for both dauer maintenance and
recovery in C. elegans. Development 135, 2583–2592. https://doi.org/10.1242/dev.012435.

Ow, M.C., Borziak, K., Nichitean, A.M., Dorus, S., Hall, S.E., 2018. Early experiences mediate
distinct adult gene expression and reproductive programs in Caenorhabditis elegans. PLoS
Genet. 14, e1007219. https://doi.org/10.1371/journal.pgen.1007219.

Ow, M.C., Hall, S.E., 2015. A Method for Obtaining Large Populations of Synchronized
Caenorhabditis elegans Dauer Larvae. Methods Mol. Biol. 1327, 209–219.
https://doi.org/10.1007/978-1-4939-2842-2_15.

Ow, M.C., Nichitean, A.M., Hall, S.E., 2021. Somatic aging pathways regulate reproductive
plasticity in Caenorhabditis elegans. eLife 10. https://doi.org/10.7554/eLife.61459.

Park, D., Estevez, A., Riddle, D.L., 2010. Antagonistic Smad transcription factors control the
dauer/non-dauer switch in C. elegans. Development 137, 477–485.
https://doi.org/10.1242/dev.043752.

Park, J., Oh, H., Kim, D.-Y., Cheon, Y., Park, Y.-J., Hwang, H., Neal, S.J., Dar, A.R., Butcher,
R.A., Sengupta, P., Kim, D.-W., Kim, K., 2021. CREB mediates the C. elegans dauer
polyphenism through direct and cell-autonomous regulation of TGF-β expression. PLoS
Genet. 17, e1009678. https://doi.org/10.1371/journal.pgen.1009678.

Pekar, O., Ow, M.C., Hui, K.Y., Noyes, M.B., Hall, S.E., Hubbard, E.J.A., 2017. Linking the
environment, DAF-7/TGFβ signaling and LAG-2/DSL ligand expression in the germline
stem cell niche. Development 144, 2896–2906. https://doi.org/10.1242/dev.147660.

Pierce, S.B., Costa, M., Wisotzkey, R., Devadhar, S., Homburger, S.A., Buchman, A.R.,
Ferguson, K.C., Heller, J., Platt, D.M., Pasquinelli, A.A., Liu, L.X., Doberstein, S.K.,
Ruvkun, G., 2001. Regulation of DAF-2 receptor signaling by human insulin and ins-1, a
member of the unusually large and diverse C. elegans insulin gene family. Genes Dev. 15,
672–686. https://doi.org/10.1101/gad.867301.

Renfree, M.B., Fenelon, J.C., 2017. The enigma of embryonic diapause. Development 144,
3199–3210. https://doi.org/10.1242/dev.148213.

Ren, P., Lim, C.S., Johnsen, R., Albert, P.S., Pilgrim, D., Riddle, D.L., 1996. Control of C.
elegans larval development by neuronal expression of a TGF-beta homolog. Science 274,
1389–1391. https://doi.org/10.1126/science.274.5291.1389.

Roy, D., Michaelson, D., Hochman, T., Santella, A., Bao, Z., Goldberg, J.D., Hubbard, E.J.A.,
2016. Cell cycle features of C. elegans germline stem/progenitor cells vary temporally and
spatially. Dev. Biol. 409, 261–271. https://doi.org/10.1016/j.ydbio.2015.10.031.

Schindler, A.J., Baugh, L.R., Sherwood, D.R., 2014. Identification of late larval stage
developmental checkpoints in Caenorhabditis elegans regulated by insulin/IGF and steroid
hormone signaling pathways. PLoS Genet. 10, e1004426.
https://doi.org/10.1371/journal.pgen.1004426.

Seidel, H.S., Kimble, J., 2015. Cell-cycle quiescence maintains Caenorhabditis elegans
germline stem cells independent of GLP-1/Notch. eLife 4.
https://doi.org/10.7554/eLife.10832.

Seidel, H.S., Kimble, J., 2011. The oogenic germline starvation response in C. elegans. PLoS
ONE 6, e28074. https://doi.org/10.1371/journal.pone.0028074.

Shaffer, J.M., Greenwald, I., 2022. Floxed exon (Flexon): A flexibly positioned stop cassette for

recombinase-mediated conditional gene expression. Proc Natl Acad Sci USA 119.
https://doi.org/10.1073/pnas.2117451119.
Shaner, N.C., Campbell, R.E., Steinbach, P.A., Giepmans, B.N.G., Palmer, A.E., Tsien, R.Y.,
2004. Improved monomeric red, orange and yellow fluorescent proteins derived from
Discosoma sp. red fluorescent protein. Nat. Biotechnol. 22, 1567–1572.
https://doi.org/10.1038/nbt1037.
Shaner, N.C., Lambert, G.G., Chammas, A., Ni, Y., Cranfill, P.J., Baird, M.A., Sell, B.R., Allen,
J.R., Day, R.N., Israelsson, M., Davidson, M.W., Wang, J., 2013. A bright monomeric green
fluorescent protein derived from Branchiostoma lanceolatum. Nat. Methods 10, 407–409.
https://doi.org/10.1038/nmeth.2413.
Shaw, W.M., Luo, S., Landis, J., Ashraf, J., Murphy, C.T., 2007. The C. elegans TGF-beta
Dauer pathway regulates longevity via insulin signaling. Curr. Biol. 17, 1635–1645.
https://doi.org/10.1016/j.cub.2007.08.058.
Simoneschi, D., Rona, G., Zhou, N., Jeong, Y.T., Jiang, S., Milletti, G., Arbini, A.A., O'Sullivan,
A., Wang, A.A., Nithikasem, S., Keegan, S., Siu, Y., Cianfanelli, V., Maiani, E., Nazio, F.,
Cecconi, F., Boccalatte, F., Fenyö, D., Jones, D.R., Busino, L., Pagano, M., 2021.
CRL4AMBRA1 is a master regulator of D-type cyclins. Nature 592, 789–793.
https://doi.org/10.1038/s41586-021-03445-y.
Singh, N., Scott, K., Proctor, J., Gordon, K.L., 2025. An RNAi screen of Rab GTPase genes in
Caenorhabditis elegans reveals that morphogenesis has a higher demand than stem cell
niche maintenance for rab-1 in the somatic cells of the reproductive system. G3 (Bethesda)
15. https://doi.org/10.1093/g3journal/jkaf085.
Singh, N., Zhang, P., Li, K.J., Gordon, K.L., 2024. The Rac pathway prevents cell fragmentation
in a nonprotrusively migrating leader cell during C. elegans gonad organogenesis. Curr.
Biol. 34, 2387-2402.e5. https://doi.org/10.1016/j.cub.2024.04.073.
Sipe, C.W., Siegrist, S.E., 2017. Eyeless uncouples mushroom body neuroblast proliferation
from dietary amino acids in Drosophila. eLife 6. https://doi.org/10.7554/eLife.26343.
Sood, C., Justis, V.T., Doyle, S.E., Siegrist, S.E., 2022. Notch signaling regulates neural stem
cell quiescence entry and exit in Drosophila. Development 149.
https://doi.org/10.1242/dev.200275.
Sternberg, P.W., Van Auken, K., Wang, Q., Wright, A., Yook, K., Zarowiecki, M., Arnaboldi, V.,
Becerra, A., Brown, S., Cain, S., Chan, J., Chen, W.J., Cho, J., Davis, P., Diamantakis, S.,
Dyer, S., Grigoriadis, D., Grove, C.A., Harris, T., Howe, K., Stein, L., 2024. WormBase
2024: status and transitioning to Alliance infrastructure. Genetics 227.
https://doi.org/10.1093/genetics/iyae050.
Stewart, D., Killeen, E., Naquin, R., Alam, S., Alam, J., 2003. Degradation of transcription factor
Nrf2 via the ubiquitin-proteasome pathway and stabilization by cadmium. J. Biol. Chem.
278, 2396–2402. https://doi.org/10.1074/jbc.M209195200.
Stiernagle, T., 2006. Maintenance of C. elegans. WormBook 1–11.
https://doi.org/10.1895/wormbook.1.101.1.
Suarez Rodriguez, F., Sanlidag, S., Sahlgren, C., 2023. Mechanical regulation of the Notch
signaling pathway. Curr. Opin. Cell Biol. 85, 102244.
https://doi.org/10.1016/j.ceb.2023.102244.
Sulston, J.E., Horvitz, H.R., 1977. Post-embryonic cell lineages of the nematode,

Caenorhabditis elegans. Dev. Biol. 56, 110–156. https://doi.org/10.1016/0012-1606(77)90158-0.

Sulston, J.E., Schierenberg, E., White, J.G., Thomson, J.N., 1983. The embryonic cell lineage of the nematode Caenorhabditis elegans. Dev. Biol. 100, 64–119. https://doi.org/10.1016/0012-1606(83)90201-4.

Taylor, S.E., 2010. Mechanisms linking early life stress to adult health outcomes. Proc Natl Acad Sci USA 107, 8507–8512. https://doi.org/10.1073/pnas.1003890107.

Tenen, C.C., Greenwald, I., 2019. Cell Non-autonomous Function of daf-18/PTEN in the Somatic Gonad Coordinates Somatic Gonad and Germline Development in C. elegans Dauer Larvae. Curr. Biol. 29, 1064-1072.e8. https://doi.org/10.1016/j.cub.2019.01.076.

Thomas, J.H., Birnby, D.A., Vowels, J.J., 1993. Evidence for parallel processing of sensory information controlling dauer formation in Caenorhabditis elegans. Genetics 134, 1105–1117.

Tian, X., Hansen, D., Schedl, T., Skeath, J.B., 2004. Epsin potentiates Notch pathway activity in Drosophila and C. elegans. Development 131, 5807–5815. https://doi.org/10.1242/dev.01459.

Trent, C., Tsuing, N., Horvitz, H.R., 1983. Egg-laying defective mutants of the nematode Caenorhabditis elegans. Genetics 104, 619–647. https://doi.org/10.1093/genetics/104.4.619.

Tullet, J.M.A., Hertweck, M., An, J.H., Baker, J., Hwang, J.Y., Liu, S., Oliveira, R.P., Baumeister, R., Blackwell, T.K., 2008. Direct inhibition of the longevity-promoting factor SKN-1 by insulin-like signaling in C. elegans. Cell 132, 1025–1038. https://doi.org/10.1016/j.cell.2008.01.030.

Vowels, J.J., Thomas, J.H., 1992. Genetic analysis of chemosensory control of dauer formation in Caenorhabditis elegans. Genetics 130, 105–123. https://doi.org/10.1093/genetics/130.1.105.

Wadsworth, W.G., Riddle, D.L., 1989. Developmental regulation of energy metabolism in Caenorhabditis elegans. Dev. Biol. 132, 167–173. https://doi.org/10.1016/0012-1606(89)90214-5.

Webster, A.K., Chitrakar, R., Taylor, S.M., Baugh, L.R., 2022. Alternative somatic and germline gene-regulatory strategies during starvation-induced developmental arrest. Cell Rep. 41, 111473. https://doi.org/10.1016/j.celrep.2022.111473.

Webster, A.K., Jordan, J.M., Hibshman, J.D., Chitrakar, R., Baugh, L.R., 2018. Transgenerational Effects of Extended Dauer Diapause on Starvation Survival and Gene Expression Plasticity in Caenorhabditis elegans. Genetics 210, 263–274. https://doi.org/10.1534/genetics.118.301250.

Weidemann, A., Johnson, R.S., 2008. Biology of HIF-1alpha. Cell Death Differ. 15, 621–627. https://doi.org/10.1038/cdd.2008.12.

Wilsterman, K., Ballinger, M.A., Williams, C.M., 2021. A unifying, eco- physiological framework for animal dormancy. Funct. Ecol. 35, 11–31. https://doi.org/10.1111/1365-2435.13718.

Wu, X., Fleming, A., Ricketts, T., Pavel, M., Virgin, H., Menzies, F.M., Rubinsztein, D.C., 2016. Autophagy regulates Notch degradation and modulates stem cell development and neurogenesis. Nat. Commun. 7, 10533. https://doi.org/10.1038/ncomms10533.

Yamamoto, S., Charng, W.-L., Bellen, H.J., 2010. Endocytosis and intracellular trafficking of

1436   Notch and its ligands. Curr. Top. Dev. Biol. 92, 165–200. https://doi.org/10.1016/S0070-
1437   2153(10)92005-X.
1438   Yochem, J., Greenwald, I., 1989. glp-1 and lin-12, genes implicated in distinct cell-cell
1439   interactions in C. elegans, encode similar transmembrane proteins. Cell 58, 553–563.
1440   https://doi.org/10.1016/0092-8674(89)90436-4.

A LAG-2::mNG (endogenous protein)

$f(t)=1.674e^{.3952t}$
$R^2=0.9992$

39.53x ****

8.06x ****

4.21x ***

Fold change over baseline average
Hours after dauer recovery

B *lag-2p::myrTdTomato* (transcriptional, membrane)

ns
*
ns

Fold change over baseline average
Hours after dauer recovery

C 0h recovered | 2h recovered | 4h recovered | 8h recovered
Merge
LAG-2::mNG
Membrane

D LAG-2::P2A::H2B::mTurquoise2 (polycistronic, nucleus)

ns
ns
ns

Fold change over baseline average
Hours after dauer recovery

E *lag-2p::mNeonGreen::PLCδ1^{PH}* (transcriptional, membrane)

ns
ns
ns

Fold change over baseline average
Hours after dauer recovery

F 0h recovered | 2h recovered | 4h recovered | 8h recovered
Merge
Nucleus
Membrane

CONFIDENTIAL