Differential expression of mitochondria-associated genes in clinical samples of *Plasmodium falciparum* showing severe manifestations

Sukriti Gujarati[1], Dinesh Gupta[2], Sanjay Kumar Kochar[3], Dhanpat Kumar Kochar[3], Ashis Das[1]*

[1]Molecular Parasitology and Systems Biology Laboratory, Department of Biological

Sciences, Birla Institute of Technology and Science (BITS), Pilani campus, Rajasthan, India

[2]Translational Bioinformatics Group, International Centre for Genetic Engineering and

Biotechnology (ICGEB), New Delhi, India

[3]Department of Medicine, S.P. Medical College, Bikaner, Rajasthan, India

* Correspondence: Ashis Das adas@pilani.bits-pilani.ac.in, ashisd28@gmail.com

Abstract

The malaria parasite mitochondrial proteins are critical targets for antimalarial drugs, however, the emergence of drug resistance against the existing protein targets necessitates novel therapeutic approaches. In this study, using custom-designed, multi-probe microarrays and pooled RNA sequencing experiments in patient samples of *Plasmodium falciparum,* we identified disease-specific sense and Natural Antisense Transcripts (NATs) encoded by the mitochondrial genome, along with those encoded in the nucleus and targeted to the parasite mitochondria. These findings span two clinical manifestations of severe malaria - hepatic dysfunction (HD) and cerebral malaria (CM). Although mitochondrial activity is known to be reduced in blood-stage malaria, upregulation of genes linked to the TCA cycle in the CM cluster indicates disrupted mitochondrial bioenergetics which might contribute to severe disease pathogenesis. Profiling of sense and antisense mitochondrial transcripts reveal a striking correlated expression of sense-antisense transcript pairs in both the disease manifestations, indicating a potential regulatory role of NATs in mitochondrial function. This study provides direct evidence for the origin of NATs from the mitochondrial genome. It further provides a dynamic gene regulation profile of key sense and antisense transcripts responsible for mitochondrial translation and bioenergetics. We propose these differentially regulated NATs against protein-coding genes specific to the observed disease severity as potential novel targets to disrupt key mitochondrial functions of the parasite.

Keywords: Mitochondria, Malaria, *Plasmodium falciparum*, Natural Antisense Transcripts (NATs), Hepatic Dysfunction, Cerebral Malaria

Introduction

Malaria is an acute febrile illness caused by the *Plasmodium* parasite and transmitted through the bite of the female *Anopheles* mosquito. *Plasmodium* infection can be broadly classified as uncomplicated or severe. Severe malaria (SM) occurs when infections are complicated by vital

organ dysfunctions or aberrant metabolism [1-2]. Infection begins with the parasite inhabiting

the red blood cells and traversing through different asexual blood stages involving active DNA

and organelle replication. It expands its ER, mitochondria, and apicoplast to establish an

intraerythrocytic cycle within 48 hours [3].

The ultrastructure of the parasite cell has revealed the division of the *Plasmodium* mitochondria

from a narrow, tubular structure in early ring stages to getting extensively branched on the

onset of schizogony [3-4]. Known to play a 'minimal' role during blood stages, *Plasmodium*

mitochondrion spans a highly conserved genome of 6 kb encoding for only three protein-coding

genes. The mitochondrial genome also encodes multiple discrete fragments of large and small

rRNA subunits, which combine with nuclear-encoded ribosomal proteins to form

mitoribosomes [5-7]. Thus, the parasite mitochondria rely significantly on the influx of proteins

and RNA molecules from the nucleus for diverse biological functions, including heme and

pyrimidine biosynthesis, maintenance of transmembrane potential for electron transport, and

oxidative phosphorylation. Mitochondria also can replicate their DNA and transcribe it with

the help of nuclear-encoded molecules. [8-9]. This diverse functionality of the parasite

mitochondrion makes this organelle an attractive target for various antimalarial drugs [10-12].

However, experimental evidence of increased drug resistance against these conventional

"druggable" mitochondrial protein targets has been widely reported [13-14]. This calls for new,

effective antimalarial approaches that can influence disease pathogenesis by manipulating gene

expression.

Advances in transcriptomics have identified non-coding RNAs, especially Natural Antisense

Transcripts (NATs), for their role in manipulating gene expression, suggesting their use as

potential 'non-conventional' drug targets [15-16]. Natural antisense transcripts (NATs) are

transcribed from the DNA strand complementary to a region harbouring a sense transcript of

either protein-coding or non-coding genes, influencing the upregulation or downregulation of

the sense transcripts [17-18]. The emergence of natural antisense transcripts and their potential

role in regulating gene expression has been widely reported in the malaria parasites [19-21].

We have previously reported the presence of NATs in the nuclear transcriptome of clinical

isolates of *P. falciparum* [22]. This study focuses on both sense and NATs that are either

encoded or targeted to the parasite mitochondria across two severe malaria manifestations -

Hepatic dysfunction and cerebral malaria [23-24] and reports the *P. falciparum* mitochondria

transcriptional profile.

Materials and Methods

Consolidating genes with potential mitochondrial localization

A systematic attempt was made to consolidate a list of *P. falciparum* mitochondrial genes using

web-based prediction tools, functional annotation, and experimental validation as published in

the literature. Due to the lack of a single, well-characterized mitochondrial localization signal

in *Plasmodium* and the lack of a single error-free mitochondrial localization prediction tool,

the nuclear-encoded and mitochondria-targeted genes were predicted using multiple web tools

based on different algorithms and trained on different datasets as suggested by Sun and

Habermann [25]. Four major web-based prediction tools used were - TargetP, based on the

mitochondrial transit peptides (mTp) enriched with arginine, leucine, and serine, responsible

for targeting proteins in the mitochondrial matrix [26], Cello2GO utilizing the BLAST

homology searching approaches [27], MitoFate, that uses mitochondrial pre-sequence motifs

and other classical factors for predicting mitochondrial proteome [28] and PFMpred, which

primarily predicts mitochondrial proteins based on amino acid and dipeptide compositions

[29]. Apart from these prediction tools, genes from PlasmoDB showing a mitochondrial

targeting domain are also taken into account. The final gene list was prepared by considering

their predicted occurrence from at least a consensus of two or more tools to increase the

reliability of the potential mitochondria-targeted genes [30]. A comprehensive literature survey

was also done to include genes experimentally validated and reported as part of the

mitochondrial proteome [32-34, 39, 44].

Blood sample processing

Venous blood samples were collected from *P. falciparum-infected* adult patients admitted to

S.P. Medical College, Bikaner, India, and passed through a histopaque density gradient to

enrich for infected erythrocytes. The team of clinicians at S.P. Medical College, Bikaner,

collected patient samples with informed consent according to hospital guidelines as per

approval. Infection with *P. falciparum* was confirmed by slide microscopy and RDTs. The

mono-infection by *P. falciparum* was further confirmed by PCR diagnosis [69-70]. Patient

samples were grouped into 3 categories, as per the WHO-defined clinical guidelines, as

tabulated in Table S1, and hybridized individually on single-color custom-designed 8*60K

microarrays.

Microarray Hybridization

The RNA from blood samples was extracted as per standard protocols [22] and used after the

concentration and purity of the RNA were evaluated using the Nanodrop Spectrophotometer

(Thermo Scientific; 2000) and Qubit (Thermo Scientific, USA) assay. The integrity of the RNA

was analyzed on Tapestation (Agilent). The microarray hybridization and scanning were

performed at the Agilent-certified microarray facility of Genotypic Technology Pvt. Ltd.,

Bengaluru, India. The samples were labeled using an Agilent Quick-Amp labeling Kit

(p/n5190-0442). Total RNA was reverse transcribed at 40°C using an oligo-dT primer tagged

to a T7 polymerase promoter and converted to double-stranded cDNA. Synthesized double-

stranded cDNA was used as the template for cRNA generation. cRNA was generated by in

vitro transcription, and the dye Cy3 CTP(Agilent) was incorporated during this step. The cDNA

synthesis and in vitro transcription steps were carried out at 40°C. Labeled cRNA was cleaned

using Qiagen RNeasy columns (Qiagen, Cat No: 74106), and quality was assessed for yields

and specific activity using the Nanodrop ND-1000. Labeled cRNA samples were fragmented

at 60°C and hybridized onto an Agilent Custom *Plasmodium falciparum* Gene Expression

Microarray 8X60K. Fragmentation of labeled cRNA and hybridization were done using the

Gene Expression Hybridization kit (Agilent Technologies, In situ Hybridization kit, Part

Number 5190-0404). Hybridization was carried out in Agilent's SureHyb Chambers at 65°C

for 16 hours. The hybridized slides were washed using Agilent Gene Expression wash buffers

(Agilent Technologies, Part Number 5188-5327) and scanned using the Agilent Microarray

Scanner (Agilent Technologies, Part Number G2600D). Raw data extraction from images was

obtained using Agilent Feature Extraction software.

Microarray design

60K sense-antisense microarray for *P. falciparum* was designed using annotations of the *P.*

falciparum 3D7 genome from PlasmoDBv57 (www.plasmodb.org). These arrays comprise 60-

mer oligonucleotide probes, covering ORFs from the nuclear, apicoplast, and mitochondrial

genomes. Most ORFs had five probes each for sense and antisense orientation. These probes

were checked against the human transcriptome, and all cross-hybridizing probes were removed.

The probes for the plasmodial mitochondrial sequence were also scanned against the *Homo*

sapiens mitochondrial genome to check for cross-hybridization. The arrays were designed and

printed on the Agilent platform. Probes from our earlier validated 244K microarray have been

incorporated into the current version of the 60K array [22]. Here, we report only the transcripts

detected by the hybridization, which are either encoded by the mitochondrial genome itself or

targeted to the mitochondria.

Microarray Data Analysis

Feature-extracted raw image data, two-fold above background intensity, was obtained and
analyzed using Agilent GeneSpring GX Software. The data was normalized in GeneSpring GX
using the 75th percentile shift method. To make the analysis highly stringent, only the probes
detecting transcripts in at least 40% of samples in each condition (including complicated and
uncomplicated) were taken further into analysis. Further, probes showing uncorrelated
expression for the same transcript were filtered out by calculating the Pearson Correlation
Coefficient with a cutoff of >=0.8. Fold change values were obtained by comparing test
samples (patients showing hepatic dysfunction and cerebral malaria manifestations) with
respect to the control samples (patients showing uncomplicated disease manifestations).
Average fold expression values were calculated for each transcript for the two test groups.
Significant genes upregulated with fold change >=0.8 (logbase2) and downregulated with fold
change <=-0.8 (logbase2) in the test samples with respect to the control group were identified.

Strand-specific RNA Sequencing

Strand-specific RNA sequencing libraries were prepared using Illumina-compatible
NEBNext® Ultra™ II Directional RNA Library Preparation Kit (New England BioLabs, MA,
USA) at Genotypic Technology Pvt. Ltd., Bangalore, India. Briefly, 500 ng of total pooled
RNA was taken for mRNA isolation, fragmentation, and priming. Two such pools of RNA
were sequenced, representing uncomplicated and severe disease complications (denoted with
* symbol in Table S1). Fragmented and primed mRNA was further subjected to first-strand
synthesis followed by second-strand synthesis. The double-stranded cDNA was purified using
JetSeq Beads (Bioline, Cat #BIO-68031). Purified cDNA was end-repaired, adenylated, and
ligated to Illumina multiplex barcode adapters as per NEBNext® Ultra™ II Directional RNA
Library Prep protocol followed by second strand excision using USER enzyme at 37˚C for 15
minutes. Adapter-ligated cDNA was purified and subjected to 11 cycles for Indexing (98˚C for
30 sec, cycling (98˚C for 10 sec, 65˚C for 75 sec) and 65˚C for 5min) to enrich the adapter-
ligated fragments. Final PCR products (sequencing libraries) were quantified by a Qubit
fluorometer after a quality control check (Thermo Fisher Scientific, MA, USA). The libraries
were paired-end sequenced on Illumina HiSeq X Ten sequencer (Illumina, San Diego, USA)
for 150 cycles following the manufacturer's instructions.

RNA Sequencing Data Analysis

Transcriptome analysis was performed by processing the raw data to remove low-quality bases

(<Q30) and adapter sequences. The high-quality reads were considered for alignment with the

human reference *Homo sapiens* (GRCh38.p13) genome using Hisat2. Unaligned reads against

the human genome were mapped against the *Plasmodium falciparum 3D7* reference genome.

Alignment against *Plasmodium* was carried out at two levels:

(I) R1 reads were aligned against reference mRNA sequences.

(ii) R2 reads were aligned against the reverse complement of mRNA sequences.

Further, read abundance was calculated for the aligned reads to be used for downstream

analysis. Differential gene expression analysis was performed for control and treatment

conditions using the R package, DEseq. Sequencing (uneven library size/depth) bias among

the samples was removed by library normalization using size factor calculation in DESeq.

DESeq normalized expression values were used to calculate fold change for a given transcript.

The differential expression calculation at log2fold change cutoff +/- 0.8 is considered.

Functional enrichment by Gene Ontology

Functional enrichment analysis of differentially expressed sense and antisense transcripts in

the two disease cohorts was performed individually using DAVID online tools (DAVIDv6.8;

http://david.ncifcrf.gov/) against a background list of the consolidated mitochondrial gene list.

The GO terms were classified into three categories: Biological process (BP), cellular

component (CC), and molecular function (MF), and p<0.05 (EASE score) was considered to

indicate a statistically significant difference.

Results

Mitochondrial associated genes

A list of 886 genes was prepared which included 42 genes encoded by the mitochondrial

genome, 18 encoded by the apicoplast genome, targeted to mitochondria; and 826 encoded in

the nucleus and targeted to the mitochondria for their molecular function (Supplementary

Sheet1). Out of the total number of genes reported from various prediction tools, 47 genes from

TargetP, 463 genes from Cello2GO, 187 genes from MitoFate, 401 genes from PFMpred, and

398 genes from PlasmoDB were shortlisted to be part of the mitochondrial proteome. The

criteria used for selection was for a gene to be predicted by at least two of the tools. We further

performed literature screening of *P. falciparum* mitochondria associated genes to include them

in the study. There appears to be a good overlap between the genes predicted from the literature

sources and those from the bioinformatics tools. This comprehensive list of 886 genes

(Supplementary Sheet1) therefore comprises 581 genes with published experimental evidence
[32-34,39,44] and 305 predicted-only genes, and forms the basis for the transcriptomic analyses
presented here.

Identification of sense-antisense transcript pairs specific to disease manifestation
Out of the 886 genes analysed for expression in the CM and HD cohort, 662 genes with sense,
and 480 genes with antisense transcripts were found to be expressed in the HD cohort, while
711 sense and 505 antisense transcripts were expressed in the CM cohort. Based on microarray
expression data, the breakup of differentially expressed genes in the three disease
manifestations is tabulated in table 1. There appears to be a much higher number of both sense
and antisense transcripts upregulated in the cerebral malaria cohort (227 upregulated sense
genes and 60 downregulated antisense genes) compared to the hepatic dysfunction disease
manifestations (34 upregulated sense genes and 20 upregulated antisense genes). The
expression profile of these upregulated genes (with average a fold value across samples >=0.8)
in the individual disease cluster is given in Supplementary Sheet2.

Out of the 886 shortlisted mitochondria associated genes, 660 sense transcripts, 448 antisense
transcripts, and 412 sense-antisense transcript pairs expressed are common between the HD
and CM datasets (Figure 1; refer to Supplementary Sheet 3 for transcript IDs). We identified
29 and 59 sense-antisense transcript pairs uniquely expressed in the hepatic dysfunction and
cerebral malaria cases, respectively (Supplementary Sheet 3). Much of the sense–antisense
transcript pairs in the two disease conditions are positively correlated (70% of S-AS pairs in
HD and 50% of S-AS pairs in CM) but there are some genes with a negative correlation
between their sense and antisense transcript pairs (0.68% in hepatic dysfunction and 2.54% in
cerebral malaria cases), i.e., if sense transcript is upregulated, antisense transcript is
downregulated and vice versa. None of the negatively correlated sense-antisense transcript
pairs are common between the two disease conditions. A heatmap expression of these
transcripts with the gene description is depicted in Figure 2. Except for the gene
PF3D7_0923200 (NOS, nitric oxide synthase, putative), all negatively correlated antisense
transcripts are downregulated in the HD and CM manifestations.

Data validation using RNA sequencing
To validate the gene expression data from microarray experimentation, an RNA sequencing
experiment was carried out by pooling 2 severe samples, one each representing the severe HD

and CM complication against an uncomplicated pool of sample (marked with * in Table S1).

Comparing the expression profile of the 412 common expressed S-AS transcript pairs between

the microarray and RNA sequencing analysis was done by calculating the Pearson correlation

coefficient (R) to reflect linearity and concordance between the two experimental approaches,

where values closer to 1 depict total positive correlation; -1 depicts total negative correlation;

0 depicts no linear correlation. The correlation plot shows r =0.82 for the expression of sense

transcripts, while antisense expression data shows r=0.6 (*pval* in both cases <0.05); reflecting

a positive correlation between the two experimental gene expression datasets (Figure 3,

Supplementary Sheet4).

Gene Ontology-based enrichment of genes with upregulated antisense transcripts

The gene ontology analysis for genes expressing upregulated and downregulated antisense

transcripts (pval / EASE score<0.05) in the respective datasets has identified several biological

processes unique to individual disease types as depicted in figure 4. In the HD manifestation,

GO terms enriched with antisense upregulated genes included intracellular protein transport

(GO:0006886) and cellular macromolecule localization (GO:0070727), suggesting a role in

intracellular molecular trafficking. Conversely, antisense downregulated genes were

significantly enriched in Carbon and pyruvate metabolism (pfa01200 and pfa00620). Key

molecular functions included oxidoreductase activity (GO:0016491) and flavin adenine

dinucleotide (FAD) binding (GO:0050660). Pathway analysis revealed significant enrichment

in TCA cycle (pfa00020), pyruvate metabolism (pfa00620), and biosynthesis of secondary

metabolites (pfa01110), indicating reduced role of antisense transcripts in electron transport

chain.

A similar GO enrichment analysis for the differentially expressed antisense transcripts in the

CM cohort show that antisense-upregulated genes were significantly enriched for ATP

synthesis coupled electron transport (GO:0042773), and oxidative phosphorylation

(GO:0006119). Molecular function enrichment includes DNA binding (GO:0003677) and

active transmembrane transporter activity (GO:0022804), suggesting an influence on nuclear

processes and membrane transport. Antisense-downregulated genes in this condition were

predominantly associated with ribosomal function and RNA processing with the most

overrepresented terms being cytosolic ribosome (GO:0022626), ribosome biogenesis

(GO:0042254), and RNA binding (GO:0003723). These results suggest a potential suppression

of protein synthesis machinery through antisense-mediated downregulation of ribosome-

associated genes. The observed differences between the two conditions indicate context-

specific expression of antisense transcripts in cellular metabolism and gene expression
regulation.

Discussion

A comparative analysis of 23 species of *Plasmodium* has determined the mitochondrial (mt)
genome of the *P. falciparum* to be the smallest [9]. It comprises a 6 kb tandemly arranged,
linear, and conserved segment of DNA, encoding only 3 protein-coding genes - *cox1*,
(PF3D7_MIT02100), *cox3* (PF3D7_MIT01400) and *cytb* (PF3D7_MIT02300). The remainder
encodes several small rRNAs transcribed from both the strands of the mtDNA and ranges from
23-190 nt in length [35]. In total, the mitochondrial genome of *P. falciparum* comprises 42
genes, of which 20 were sufficiently long to be included in the 60-mer probe design in both
sense and antisense orientations; the remaining ORFs were too short (<60 nucleotides) for
effective probe design and were therefore, excluded from the microarray.

Although the prevalence of NATs encoded from the nuclear genome of the parasite has been
well documented [22,36], the transcription profile of the NATs from the mitochondrial genome
of the parasite has been less clear. An overlap of 274 and 285 NATs in HD and CM cluster
respectively has been identified after comparing our mitochondrial dataset with the genome
wide NATs assembled by Subudhi et. al. [22]. Comparison of our NATs expression data with
that of time-based NATs catalogue of laboratory strain of asexual blood stage parasite 3D7
[36] revealed 222 antisense RNAs that appear exclusively in the clinical dataset analysed here
(Supplementary Sheet5). This study further detected antisense transcripts for all 20
mitochondrial genes included in the array. Upon investigation of the protein coding genes of
the mtDNA, we observe an enhanced expression of the sense mRNA of the three respiratory
chain-associated genes: *cox1*, *cox3*, and *cytb*, in both the disease manifestations. These three
genes form the core components of the parasite's mitochondrial electron transport chain
(mETC), essential for ATP production and parasite survival, and is critical to the parasite
transmission from human host to mosquito [35]. The enhanced expression of these genes in the
sense orientation (with a corresponding decrease in antisense levels) in severe malaria
conditions might reflect the parasite's increased ATP requirement associated with rapid
proliferation, high parasitemia, and the immune-evasive demands of severe host-pathogen
interactions during severe malaria infection. Cerebral malaria is often associated with higher
parasitemia, increased metabolic needs, and the parasite's attempt to evade the host immune

response [24]; thus, bolstering the efficiency of the ETC could be advantageous for rapid proliferation and survival in this stringent environment. This is further confirmed by the enrichment of biological processes like – *oxidative phosphorylation* and *ATP synthesis coupled electron transport chain* associated with upregulated sense transcripts in CM manifestation. However, further functional studies are needed to confirm this correlation.

The mtETC has been a critical target for malaria treatment [50]. One prominent example is the antimalarial drug atovaquone, which effectively blocks parasite growth by inhibiting the mitochondrial protein cytb during the asexual blood stage of infection [51]. However, resistance against atovaquone has emerged due to point mutations in the cytb gene. Antisense oligonucleotide (ASO)-mediated gene silencing offers a promising alternative therapeutic strategy capable of overcoming such drug resistance by specifically reducing target mRNA expression levels, thus circumventing mutation-based resistance mechanisms [52]. The observed low antisense transcript levels for these critical ETC genes suggest limited transcriptional regulation of these genes by their complementary NATs under severe malaria conditions, potentially facilitating effective ASO-based therapeutic intervention to disrupt parasite proliferation. In addition to mitochondrial encoded ETC genes (*cox1, cox3, and cytb*), nuclear-encoded, apicomplexan-specific ETC subunits (PfCOX6A, PfNDUFA4, PfCOX4, ApiCOX24) also exhibit minimal to no antisense RNA expression, suggesting a conserved pattern of minimal antisense regulation for critical ETC components. This observation underscores the potential for these genes as attractive targets for antisense-based therapeutic approaches, and can be functionally investigated for disrupting essential metabolic processes in malaria parasites.

Despite encoding for three protein-coding genes, plasmodial mitochondria is a validated drug target and has been reported to perform several critical functions for the parasite viz, shuttling electrons for oxidative phosphorylation, iron-sulfur cluster biosynthesis, pyrimidine metabolism, and de novo biosynthesis of heme [8,39]. Owing to its diversified functions, the parasite mitochondria heavily depend on the trafficking of molecules from the nucleus. We have divided these nuclear-encoded transcripts with putative mitochondrial functions into the distinct functional roles and investigated their expression for and antisense transcripts across the two disease conditions (Figure 5). The presence of a functional TCA cycle in the blood stages of malaria has remained debatable [9,47]. We have identified several key transcripts of the TCA cycle and mETC, upregulated in cerebral malaria patients (Figure 5). In addition to this, the enrichment of '*Oxidative phosphorylation*' (pfa00190) pathway highlights the

potential functionality of these genes in driving the parasite's energy metabolism during blood-

stage infection. The genes for the TCA cycle showed expression in parasites in uncomplicated

disease conditions. However, they were upregulated in case of cerebral malaria. The

identification of upregulated key transcripts of the TCA cycle and mitochondrial ETC in

cerebral malaria adds a valuable layer to understanding the parasite's metabolic adaptations

under severe disease conditions. Studying the protein abundance of these genes would provide

conclusive evidence of the extent to which the parasite brings its oxidative phosphorylation

into effect in the vertebrate host.

Plasmodium mitochondria encode 27 rRNA 'minigenes' which form the structural component

of the parasite's mitochondrial ribosome (mitoribosome), enabling mitochondrial protein

synthesis to occur [7]. The pivotal role of mitoribosomes has been characterized in translating

genes encoded in the mtDNA, and a conditional knockdown of these transcripts has shown

reduced activity of the cytochrome bc1 complex [40]. Our expression dataset has detected the

prevalence of 12 ncRNA genes encoding for ribosomal rRNA fragment in both sense and

antisense orientation in all the clinical isolates hybridized under investigation. The inability to

detect the expression of the remaining ncRNAs could be attributed to experimental limitations

of microarray and short-read RNAseq and emerging technologies employing longer-read

RNAseq could help capture the complete transcriptome for these ncRNAs [53]. However, the

expression of both sense and antisense transcripts for the genes involved in mitoribosome

assembly across both complicated and uncomplicated disease states warrants the fact that the

Plasmodium mitochondria have a functionally active protein translational machinery in asexual

ring stage of development. Another molecule, ICT1 (PF3D7_1117600), that is perpetually

expressed across all disease states, encodes for peptidyl-tRNA hydrolase enzyme to catalyze

hydrolytic cleavage of prematurely terminated peptidyl-tRNA moieties in stalled

mitoribosomes from the mitochondrial ORFs of humans [41-42]. Its critical role in gene

expression in organellar translation identifies this as an important antimalarial target [54],

warranting further investigation. Transcriptional interference of these ribosomal RNAs by

complementation with their respective antisense RNAs could be a plausible mechanism to

control gene regulation in the parasite by stalling mitoribosome assembly. However, further

experimental validation is necessary to confirm this regulatory model.

Mitochondrial gene regulation is increasingly recognized as an important contributor to the cell

physiology [55-56]. NATs have been recognized for their roles in controlling gene expression

at multiple levels, including transcriptional interference and chromatin remodeling [57-59]. In

the context of malaria disease biology, these patterns of sense/antisense expression in the

parasite's mitochondrion may have functional significance. We have observed discordant (high

sense and low antisense) gene expression of fifteen S-AS transcripts in the two clinical

manifestations investigated (Figure 2). Four of these (PF3D7_1235100, PF3D7_1336600,

PF3D7_0821000, PF3D7_0912700) encode for conserved *Plasmodium* proteins and are

important targets for further characterization.

While most of these negatively regulated gene pairs show high sense and low antisense

transcription, we observed that only one gene, NOS (PF3D7_0923200, nitric oxide synthase),

exhibited inverse expression pattern with elevated levels of antisense transcripts alongside

reduced levels of its corresponding sense transcript. This pattern suggests the potential

involvement of antisense RNA in the negative regulation of NOS expression, thereby

modulating nitric oxide (NO) production. Low NO levels have been associated with more

severe disease [60-61]. In an experimental cerebral malaria model, inhaled nitric oxide reduced

inflammation and improved survival of the infected mice [62]. Given that host-derived NO

exerts antiparasitic effects [63], the parasite's own control over NO synthase expression could

be critical. If the parasite were to produce large amounts of NO through its own NOS activity,

this might be detrimental to its survival. In this context, antisense regulation could serve as a

safeguard, maintaining NO production at levels that are not self-damaging to the parasite. This

finding supports previous reports that highlight the importance of finely tuning NO production

during malaria infections [64].

We have further presented a comprehensive list of upregulated antisense transcripts in

individual disease manifestations (Supplementary Sheet 2). Four of them are encoded by the

nucleus (PF3D7_0410800, AKIT3; PF3D7_0524700, TOM22; PF3D7_0719100, ATP

synthase F0 subunit a-like protein and PF3D7_1037300, AAC1) and consistently exhibit

elevated antisense transcription in both the disease manifestations. These genes are critical

components of the mitochondrial protein import machinery in *P. falciparum*. Specifically,

AAC1 (PF3D7_1037300 - ADP/ATP carrier protein 1) which functions as an ATP/ADP

antiporter, plays an essential role in the energy metabolism of the parasite [43]. Transcriptomic

profiling of antisense genes by [36], from culture strain of *P. falciparum* detected low

abundance of this gene across all four time points in contrast to the transcriptome reported in

the current study. This discrepancy emphasizes the need for both in vitro and ex vivo datasets

for a comprehensive and accurate understanding of parasite gene expression. TOM22

(PF3D7_0524700) is another key gene, encoding the mitochondrial import receptor subunit

TOM22, which is translocated to the outer membrane of mitochondria for the assembly of the

TOM complex (translocase of the outer membrane of mitochondria) for importing

mitochondrial proteins [44-45]. Collectively, these results support the idea that mitochondrial

transport genes are promising targets for further investigation in severe *P. falciparum* malaria,

for several reasons: i) their functional assembly mediates the uptake of solutes essential for

parasite proliferation and survival; hence, perturbing their function can compromise critical

mitochondrial processes; (ii) *P. falciparum* exhibits resistance against various conventional

transporters [46], signalling the need for new therapeutic strategies less prone to resistance;

and (iii) the consistent abundance of antisense transcripts for these genes in both the disease

clusters (by both microarray and RNA sequencing) may suggest their regulation under control

by sense/antisense interaction. Further, direct human homolog of these genes shows negligible

similarity by standard BLAST searches, enhancing the clinical potential of these targets. A

functionally analogous receptor (TOMM22) does exist in humans, but does not share high

sequence identity. The upregulated NATs associated with these genes may form part of a

network that ensures timely production and assembly of the corresponding protein-coding

transcripts [65-66], serving as important control points in parasite biology. Elevated NATs can

significantly perturb the translational potential of their targets [65-67]. Future work involving

targeted knockdowns, CRISPR interference, or other functional approaches against these genes

is warranted to validate their precise regulatory roles and clinical relevance as potential

biomarkers in severe malaria.

Studies on non-coding RNAs of *P. falciparum* have only recently been taken up by multiple

groups [36,57,68]. A constant challenge in understanding parasite disease biology is that most

of these studies work in culture systems. We have earlier presented the profile of NATs clinical

datasets of *P. falciparum* [22]. This work further presents a comprehensive transcriptomic

dataset of the parasite mitochondria from 22 clinical isolates, with their differential regulation

across two of the most severe disease complications of malaria. We have identified sense-

antisense pairs specific to respective disease manifestations. Given the minimal role of

mitochondria in blood-stage malaria [9], the extensive presence of these NATs from

mitochondrial ORFs in severe malaria patient isolates presses upon their regulatory functions

in parasite biology. Since the mitochondrial genome is arranged in tandem repeats, detecting

both sense and antisense transcripts hints at active post-transcriptional regulation in the

mitochondrion. This work expands our understanding of *Plasmodium* mitochondrial biology,

highlights the interplay of these sense and antisense transcripts in the context of mitochondrial microenvironment of the parasite under host pressure, emphasizing key genes whose investigation could provide us with potential targets for novel therapeutic interventions against *P. falciparum* malaria. Although further mechanistic work—such as dissecting how precisely these antisense transcripts engage with the sense mRNAs, and whether they are processed by parasite-specific RNA regulatory pathways—remains to be done, the current findings will facilitate deeper investigations into *RNA*-mediated mitochondrial gene regulation in *Plasmodium* infection.

Acknowledgments

We thank all the patients and technical workers for participating in and supporting this project. This work is funded by the Indian Council of Medical Research (ICMR) under the Extramural Ad-hoc Scheme (PID: 2019-1121, a multicentric project with AD as PI and DG and SKK as Co-PI). We thank Birla Institute of Technology (BITS), Pilani (Pilani campus), International Centre for Genetic Engineering and Biotechnology (ICGEB), New Delhi, and SP Medical College, Bikaner, for providing the necessary facilities required for this work. We also thank Genotypic Technology Pvt. Ltd., Bangalore, India, for the microarray hybridization and RNA sequencing experimentation and consultation regarding data analysis.

Availability of data and materials

All microarray results have been deposited in the Gene Expression Omnibus database (GSE272963). All RNAseq data corresponding to the genes investigated under this study are provided in the supplementary data files. All patient meta-data included in this article are included in Supplementary Sheet1 (Table S1). Reprint requests should be addressed to AD.

Declaration of interest

The authors declare no competing interests.

Ethics statement

Ethics Approval - Sample collection was earlier approved by SP Medical College Hospital's Ethics Committee (No. F (Acad)SPMC/2003/2395). Permission to use these samples for further studies was given through IERB approval No. F29(Acad)SPMC/2020/3151 dated 05.09.2020.

and fresh samples were collected according to EC/SPMC/ECA/06/28-02-2019. We thank all the

patients for their voluntary consent and participation.

Author Contributions

AD conceived the study, acquired funding from ICMR, India, and networked to facilitate the

outcome. SG was involved in performing the experimentation, analysis, and data interpretation,

designing the figures, and drafting the manuscript under the constant guidance of AD. DG has

been closely involved in consultations on data analysis and related discussions since the study's

inception. DKK and SKK supervised sample collection and clinical characterization. All authors

read and approved the manuscript.

References

[1] World Malaria Report 2021. Geneva: World Health Organization; 2021. License: CC BY-

NC-SA 3.0 IGO. Available from https://www.who.int/teams/global-malaria-

programme/reports/world-malaria-report-2021.

[2] The World Health Organization- Severe Malaria TM & IH; 2014. 19,7-131. 2014;

Available from: https://onlinelibrary.wiley.com/doi/10.1111/tmi.12313_2.

[3] Krungkrai S, Leangaramgul P, Kudan S. et al, Mitochondrial Heterogeneity in Human

Malarial Parasite Plasmodium falciparum. ScienceAsia. 1999;77-83.

[4] Van Dooren GG, Marti M, Tonkin CJ, et al. Development of the endoplasmic reticulum,

mitochondrion, and apicoplast during the asexual life cycle of Plasmodium falciparum. Mol

Microbiol. 2005; 57:405–419.

[5] Vaidya AB, Akella R, Suplick K. Sequences similar to genes for two mitochondrial proteins

and portions of ribosomal RNA in tandemly arrayed 6-kilobase-pair DNA of a malarial

parasite. Mol Biochem Parasitol. 1989; 15;35(2):97-107. doi: 10.1016/0166-6851(89)90112-

6.

[6] Gupta A, Shah P, Haider A, Gupta K, Siddiqi MI, Ralph SA, Habib S. Reduced ribosomes

of the apicoplast and mitochondrion of Plasmodium spp. and predicted interactions with

antibiotics. Open Biol. 2014; 5:140045. doi: 10.1098/rsob.140045.

[7] Ling L, Mulaka M, Munro J, Dass S, et. al. Genetic ablation of the mitoribosome in the

malaria parasite Plasmodium falciparum sensitizes it to antimalarials that target mitochondrial

functions. J Biol Chem. 2020; 295(21):7235-7248. doi: 10.1074/jbc.RA120.012646.

[8] Krungkrai J. The multiple roles of the mitochondrion of the malarial parasite. Parasitology.

2004; 129(Pt 5):511-24. doi: 10.1017/s0031182004005888.

[9] Vaidya AB, Mather MW. Mitochondrial evolution and functions in malaria parasites. Annu

Rev Microbiol. 2009; 63:249-67. doi: 10.1146/annurev.micro.091208.073424.

[10] Fry M, Pudney M. Site of action of the antimalarial hydroxynaphthoquinone, 2-[trans-4-

(4'-chlorophenyl) cyclohexyl]-3-hydroxy-1,4-naphthoquinone (566C80). Biochem Pharmacol.

1992; 43(7):1545-53. doi: 10.1016/0006-2952(92)90213-3.

[11] Srivastava IK, Vaidya AB. A Mechanism for the Synergistic Antimalarial Action of Atovaquone and Proguanil. Antimicrob Agents Chemother. 1999; 43(6):1334-9. doi: 10.1128/AAC.43.6.1334.

[12] Painter HJ, Morrisey JM, Vaidya AB. Mitochondrial electron transport inhibition and viability of intraerythrocytic Plasmodium falciparum. Antimicrob Agents Chemother. 2010; 54:5281–5287. doi: 10.1128/AAC.00937-10.

[13] Vaidya AB, Mather MW. Atovaquone resistance in malaria parasites. Drug Resist Updat. 2000; 3(5):283-287. doi: 10.1054/drup.2000.0157.

[14] Fivelman QL, Butcher GA, Adagu IS et al. Malarone treatment failure and in vitro confirmation of resistance of Plasmodium falciparum isolate from Lagos, Nigeria. Malar J. 2002;1:1. doi: 10.1186/1475-2875-1-1.2002.

[15] Noonpakdee W, Pothikasikorn J, Nimitsantiwong W, et al. Inhibition of Plasmodium falciparum proliferation in vitro by antisense oligodeoxynucleotides against malarial topoisomerase II. Biochem Biophys Res Commun. 2003; 302:659–664. doi: 10.1016/s0006-291x(03)00246-8.

[16] Wahlestedt C. Natural antisense and noncoding RNA transcripts as potential drug targets. Drug Discov Today. 2006 11(11-12):503-8. doi: 10.1016/j.drudis.2006.04.013.

[17] Lodde V, Floris M, Muroni MR, et al. Non-coding RNAs in malaria infection. Wiley Interdiscip Rev RNA. 2022; 13(3):e1697. doi: 10.1002/wrna.1697.

[18] Werner A, Berdal A. Natural antisense transcripts: sound or silence? Physiol Genomics. 2005; 23(2):125–131.

[19] Militello KT, Patel V, Chessler AD, et al. RNA polymerase II synthesizes antisense RNA in Plasmodium falciparum. RNA. 2005; 11(4):365-70. doi: 10.1261/rna.7940705.

[20] Broadbent KM, Broadbent JC, Ribacke U, et al. Strand-specific RNA sequencing in Plasmodium falciparum malaria identifies developmentally regulated long non-coding RNA and circular RNA. BMC Genomics. 2015; 16(1):454. doi: 10.1186/s12864-015-1603-4.

[21] Boopathi PA, Subudhi AK, Garg S, et al. Revealing natural antisense transcripts from Plasmodium vivax isolates: Evidence of genome regulation in complicated malaria. Infect Genet Evol. 2013; 20:428-43. doi: 10.1016/j.meegid.2013.09.026.

[22] Subudhi AK, Boopathi PA, Garg S, et al. Natural antisense transcripts in Plasmodium falciparum isolates from patients with complicated malaria. Exp Parasitol. 2014; 141:39-54. doi: 10.1016/j.exppara.2014.03.008.

[23] Severe falciparum malaria. Geneva: World Health Organization, Communicable Disease Cluster. Transactions of the Royal Society of Tropical Medicine and Hygiene. 2000; 94: Supplement I.

[24] Muppidi P, Wright E, Wassmer SC and Gupta H. Diagnosis of cerebral malaria: Tools to reduce Plasmodium falciparum associated mortality. Front. Cell. Infect. Microbiol. 2023; 13:1090013. doi: 10.3389/fcimb.2023.1090013.

[25] Sun S, Habermann BH. A guide to computational methods for predicting mitochondrial localization. Methods Mol Biol. 2017; 1567:1-14. doi: 10.1007/978-1-4939-6824-4_1.

[26] Almagro Armenteros JJ, Salvatore M, Emanuelsson O, et al. Detecting sequence signals in targeting peptides using deep learning. Life Sci Alliance. 2019; 2(5):e201900429. doi: 10.26508/lsa.201900429.

[27] Yu CS, Cheng CW, Su WC, et al. CELLO2GO: a web server for protein subCELlular LOcalization prediction with functional gene ontology annotation. PLoS One. 2014; 9(6):e99368. doi: 10.1371/journal.pone.0099368.

[28] Fukasawa Y, Tsuji J, Fu SC, et al. MitoFates: Improved prediction of mitochondrial targeting sequences and their cleavage sites. Mol Cell Proteomics. 2015; 14(4):1113-26. doi: 10.1074/mcp.M114.043083.

[29] Verma R, Varshney GC, Raghava GPS. PFMpred: Predicting mitochondrial proteins of P. falciparum. Amino Acids. 2009;10.1007/s00726-009-0381-1.

[30] Claros' MG, Vincens P. Computational method to predict mitochondrially imported proteins and their targeting sequences. Eur. J. Biochem. 1996; 241(3):779-86. doi: 10.1111/j.1432-1033.1996.00779.x.

[31] Management of severe malaria: a practical handbook – 3rd ed. World Health Organization; 2012. WHO Library Cataloguing-in-Publication Data.

[32] Evers F, Cabrera-Orefice A, Elurbe DM, et al. Composition and stage dynamics of mitochondrial complexes in Plasmodium falciparum. Nat Commun. 2021; 12(1):3820. doi: 10.1038/s41467-021-23919-x.

[33] Lamb IM, Rios KT, Shukla A, et al. Mitochondrially targeted proximity biotinylation and proteomic analysis in Plasmodium falciparum. PLoS One. 2022; 17 (8):e0273357. doi: 10.1371/journal.pone.0273357.

[34] van Esveld SL, Meerstein-Kessel L, Boshoven C, et al. A Prioritized and Validated Resource of Mitochondrial Proteins in Plasmodium Identifies Unique Biology. mSphere. 2021; 6(5):e0061421. doi: 10.1128/mSphere.00614-21.

[35] Hikosaka K, Watanabe Y ichi, Kobayashi F, et al. Highly conserved gene arrangement of the mitochondrial genomes of 23 Plasmodium species. Parasitol Int. 2011; 60(2):175-80. doi: 10.1016/j.parint.2011.02.001.

[36] Siegel TN, Hon CC, Zhang Q, et al. Strand-specific RNA-Seq reveals widespread and developmentally regulated transcription of natural antisense transcripts in Plasmodium falciparum. BMC Genomics. 2014; 15(1):150. doi: 10.1186/1471-2164-15-150.

[37] Feagin JE, Harrell MI, Lee JC, et al. The fragmented mitochondrial ribosomal RNAs of plasmodium falciparum. PLoS One. 2012; 7(6):e38320. doi: 10.1371/journal.pone.0038320.

[38] Rehkopf DH, Gillespie DE, Harrell MI, et al. Transcriptional mapping and RNA processing of the Plasmodium falciparum mitochondrial mRNAs. Mol Biochem Parasitol. 2000; 105(1):91-103. doi: 10.1016/s0166-6851(99)00170-x.

[39] Van Dooren GG, Stimmler LM, McFadden GI. Metabolic maps and functions of the Plasmodium mitochondrion. FEMS Microbiol Rev. 2006; 30(4):596-630. doi: 10.1111/j.1574-6976.2006.00027.x.

[40] Dass S, Mather MW, Morrisey JM, et al. Transcriptional changes in Plasmodium falciparum upon conditional knockdown of mitochondrial ribosomal proteins RSM22 and L23. PLoS One. 2022; 17(10):e0274993. doi: 10.1371/journal.pone.0274993.

[41] Richter R, Rorbach J, Pajak A, et al. A functional peptidyl-tRNA hydrolase, ICT1, has been recruited into the human mitochondrial ribosome. EMBO Journal. 2010; 29:1116–1125.

[42] Vaishya S, Kumar V, Gupta A, et al. Polypeptide release factors and stop codon recognition in the apicoplast and mitochondrion of Plasmodium falciparum. Mol Microbiol. 2016; 100:1080–1095.

[43] Hatin I, Jaureguiberry G. Molecular Characterisation of the ADP/ATP-Transporter cDNA from the Human Malaria Parasite Plasmodium Falciparum. Eur J Biochem. 1995; 228(1):86–91. doi: 10.1111/j.1432-1033.1995.0086o.x.

[44] Deponte M. Mitochondrial Protein Import in Malaria Parasites. 2013. In: Hommel, M., Kremsner, P. (eds) Encyclopedia of Malaria. Springer, New York, NY. https://doi.org/10.1007/978-1-4614-8757-9_95-1

[45] Wunderlich J. Updated List of Transport Proteins in Plasmodium falciparum. Front Cell Infect Microbiol. 2022. 12:926541. doi: 10.3389/fcimb.2022.926541.

[46] Kim J, Tan YZ, Wicht KJ, et. al. Structure and drug resistance of the Plasmodium falciparum transporter PfCRT. Nature. 2019;576(7786):315-320. doi: 10.1038/s41586-019-1795-x.

[47] Ke H, Lewis IA, Morrisey JM, et. al. Genetic investigation of tricarboxylic acid metabolism during the Plasmodium falciparum life cycle. Cell Rep. 2015;11(1):164-74. doi: 10.1016/j.celrep.2015.03.011.

[50] Stocks PA, Barton V, Antoine T, Biagini BA, Ward SA, O'neill PM. Novel inhibitors of the plasmodium falciparum electron transport chain. *Parasitology*. 2014;141(1):50-65. doi:10.1017/S0031182013001571.

[51] Painter, H. J., Morrisey, J. M., Mather, M. W. & Vaidya, A. B. Specific role of mitochondrial electron transport in blood-stage *Plasmodium falciparum*. *Nature* **446**, 88–91 (2007).

[52] Verma A. Recent Advances in Antisense Oligonucleotide Therapy in Genetic Neuromuscular Diseases. Ann Indian Acad Neurol. 2018 Jan-Mar;21(1):3-8. doi: 10.4103/aian.AIAN_298_17. PMID: 29720791; PMCID: PMC5909143.

[53] Ren, Bingbing et al. Emerging functions of mitochondria-encoded noncoding RNAs. Trends in Genetics, Volume 39, Issue 2, 125 – 139.

[54] Vaishya, S., Kumar, V., Gupta, A., Siddiqi, M. I., & Habib, S. (2016). Polypeptide release factors and stop codon recognition in the apicoplast and mitochondrion of *Plasmodium falciparum*. *Molecular Microbiology, 100*(6), 1080–1095. https://doi.org/10.1111/mmi.13369.

[55] Front. Microbiol., 15 October 2021 Sec. Evolutionary and Genomic Microbiology, Volume 12 - 2021 | https://doi.org/10.3389/fmicb.2021.751775

[56] Sun W, Lu Y, Zhang H, Zhang J, Fang X, Wang J, Li M. Mitochondrial Non-Coding RNAs Are Potential Mediators of Mitochondrial Homeostasis. Biomolecules. 2022 Dec 13;12(12):1863. doi: 10.3390/biom12121863. PMID: 36551291; PMCID: PMC9775270.

[57] Vembar SS, Scherf A, Siegel TN. Noncoding RNAs as emerging regulators of Plasmodium falciparum virulence gene expression. Curr Opin Microbiol. 2014 Aug;20:153-61. doi: 10.1016/j.mib.2014.06.013. Epub 2014 Jul 12.

[58] Zhao S, Zhang X, Chen S, Zhang S. Natural antisense transcripts in the biological hallmarks of cancer: powerful regulators hidden in the dark. J Exp Clin Cancer Res. 2020 Sep 14;39(1):187. doi: 10.1186/s13046-020-01700-0.

[59] Novel insights on the positive correlation between sense and antisense pairs on gene expression, Subhadeep Das, Maria Paula Zea Rojas, Elizabeth J. Tran. First published: 01 August 2024 https://doi.org/10.1002/wrna.1864.

[60] Liu X, Huang Y, Pokreisz P, Vermeersch P, Marsboom G, et al. (2007) Nitric oxide inhalation improves microvascular flow and decreases infraction size after myocardial ischemia and reperfusion. J Am Coll Cardiol 50: 808–817.

[61] Weinberg JB, Lopansri BK, Mwaikambo E, Granger DL (2008) Arginine, nitric oxide, carbon monoxide, and endothelial function in severe malaria. Curr Opin Infect Dis 21: 468–475.

[62] Inhaled Nitric Oxide Reduces Endothelial Activation and Parasite Accumulation in the Brain, and Enhances Survival in Experimental Cerebral Malaria, Published: November 16, 2011, https://doi.org/10.1371/journal.pone.0027714.

[63] Simon A Cobbold, Manuel Llinás, Kiaran Kirk. Sequestration and metabolism of host cell arginine by the intraerythrocytic malaria parasite Plasmodium falciparum. https://doi.org/10.1111/cmi.12552.

[64] Kotepui KU, Mahittikorn A, Wilairatana P, Masangkay FR, Kotepui M. Association between *Plasmodium* Infection and Nitric Oxide Levels: A Systematic Review and Meta-Analysis. Antioxidants (Basel). 2023 Oct 16;12(10):1868. doi: 10.3390/antiox12101868.

[65] Faghihi MA, Wahlestedt C. Regulatory roles of natural antisense transcripts. Nat Rev Mol Cell Biol. 2009 Sep;10(9):637-43. doi: 10.1038/nrm2738. Epub 2009 Jul 29. PMID: 19638999; PMCID: PMC2850559.

[66] Liang XH, Nichols JG, Hsu CW, Vickers TA, Crooke ST. mRNA levels can be reduced by antisense oligonucleotides via no-go decay pathway. Nucleic Acids Res. 2019 Jul 26;47(13):6900-6916. doi: 10.1093/nar/gkz500. PMID: 31165876; PMCID: PMC6649848.

[67] Crooke ST, Liang XH, Baker BF, Crooke RM. Antisense technology: A review. J Biol Chem. 2021; 296:100416. doi: 10.1016/j.jbc.2021.100416. Epub 2021 Feb 16.

[68] Thompson, Trevor A. et al.The role of long noncoding RNAs in malaria parasites. Trends in Parasitology, Volume 39, Issue 7, 517 – 531.

[69] Pakalapati D, Garg S, Middha S, Kochar A, Subudhi AK, Arunachalam BP, Kochar SK, Saxena V, Pareek RP, Acharya J, Kochar DK, Das A. Comparative evaluation of microscopy, OptiMAL(®) and 18S rRNA gene based multiplex PCR for detection of Plasmodium falciparum & Plasmodium vivax from field isolates of Bikaner, India. Asian Pac J Trop Med. 2013 May 13;6(5):346-51. doi: 10.1016/S1995-7645(13)60037-1. PMID: 23608372.

[70] Pakalapati D, Garg S, Middha S, Acharya J, Subudhi AK, Boopathi AP, Saxena V, Kochar SK, Kochar DK, Das A. Development and evaluation of a 28S rRNA gene-based nested PCR assay for P. falciparum and P. vivax. Pathog Glob Health. 2013 Jun;107(4):180-8. doi: 10.1179/2047773213Y.0000000090. PMID: 23816509; PMCID: PMC4001468.

Transcript type	Hepatic Dysfunction			Cerebral Malaria		
	UP	DOWN	NDE	UP	DOWN	NDE
Sense transcripts	34	82	546	227	148	336
Antisense transcripts	20	112	348	60	112	333

Table 1 Log2 Fold change value of >=0.8 for upregulation, <=-0.8 for downregulation and between 0.8 to -0.8 for non-differential expression of transcripts.

Figure 1 Venn diagram showing the overlap of the number of sense-antisense transcript pairs across two severe manifestations: HD: Hepatic dysfunction; CM: Cerebral Malaria; S: Sense transcripts; AS: Antisense transcripts

Gene ID	Product description
PF3D7_1014700	prohibitin 2
PF3D7_1235100	conserved protein, unknown function
PF3D7_0908800	mitochondrial carrier protein, putative
PF3D7_0604600	DNA helicase, putative
PF3D7_1336600	conserved protein, unknown function
PF3D7_1362000	cytochrome c oxidase subunit ApiCOX24, putative
PF3D7_0934300	ATP synthase-associated protein, putative
PF3D7_0107400	ATP synthase-associated protein, putative
PF3D7_1459700	pyridoxal 5'-phosphate synthase, putative
PF3D7_0824300	GTP-binding protein Obg2, putative
PF3D7_1239700	ATP-dependent zinc metalloprotease FTSH 1
PF3D7_0821000	conserved Plasmodium protein, unknown function
PF3D7_0912700	conserved Plasmodium protein, unknown function
PF3D7_1340800	mitochondrial pyruvate carrier protein 1, putative
PF3D7_0923200	nitric oxide synthase, putative
PF3D7_0306400	FAD-dependent glycerol-3-phosphate dehydrogenase, putative
PF3D7_0923800	thioredoxin reductase

Figure 2 Heatmap diagram showing the gene expression values of sense-antisense transcript pairs that are negatively correlated in the two disease cohorts- HD and CM. individually The adjoining table corresponds to the description of product these genes as per PlasmoDBv65.

HD: Hepatic Dysfunction: CM: Cerebral malaria

Figure 3 The sample correlation coefficient (R) for antisense transcripts is 0.6, while that for sense transcripts is 0.82 between microarray and RNA sequencing pool of samples (p-value < 0.05).

S-RNAseq: Sense transcript expression in RNA seq experiment | AS-RNAseq: Antisense transcript expression in RNA seq experiment | Avg S-pool microarray: Average fold expression of sense transcripts for the pooled samples used in RNAseq experimentation | Avg AS-pool microarray: Average fold expression of antisense transcripts for the pooled samples used in RNAseq experimentation

Figure 4 Significant Biological processes and KEGG pathways enriched for differentially expressed sense and antisense transcripts across the two disease types HD: Hepatic Dysfunction: CM: Cerebral malaria (pval <0.05).

Figure 5 Heatmap expression of sense and antisense transcripts. The transcripts are categorized by their functions, which are indicated by the colored tiles on the left side of the heatmap. White represents no transcript detected corresponding to that gene for the disease group. HD: Hepatic Dysfunction: CM: Cerebral malaria.