Advanced neural activity mapping in brain organoids via field potential imaging with ultra-high-density CMOS microelectrodes.

Remi Yokoi[1], Naoki Matsuda[1], Yuto Ishibashi[1], Ikuro Suzuki[1*]

[1]Department of Electronics, Graduate School of Engineering, Tohoku Institute of Technology, 35-1
Yagiyama Kasumicho, Taihaku-ku, Sendai, Japan

* Correspondence:
Ikuro Suzuki
i-suzuki@tohtech.ac.jp

Keywords: organoid[1], assembloid[2], cerebral[3], midbrain[4], striatum[5], micro-electrode array[6],
complementary metal-oxide semiconductor[7]

Abstract

Human iPSC-derived brain organoids and assembloids have emerged as promising in vitro models
for recapitulating human brain development, neurological disorders, and drug responses. However,
detailed analysis of their electrophysiological properties requires advanced measurement techniques.
Here, we present a novel analytical approach utilizing ultra-high-density (UHD) CMOS
microelectrode arrays (MEAs) containing 236,880 electrodes (10.52 μm × 10.52 μm each) distributed
over a broad sensing area of 32.45 mm² for field potential imaging (FPI) of brain organoids.
Neuronal activity was recorded simultaneously from over 46,000 electrodes interfaced with brain
organoids, allowing for the identification of single-cell firing events and the assessment of neuronal
network connectivity based on individual spikes. In midbrain organoids, administration of L-DOPA
revealed both excitatory and inhibitory cellular responses, with a dose-dependent increase in the
proportion of excitatory responses, suggesting enhanced network connectivity. Capitalizing on the
spatial and temporal resolution of UHD-CMOS-MEAs, we introduced new endpoints for network
activity: propagation speed and propagation area. In cortical organoids, application of the GABA$_A$
receptor antagonist picrotoxin led to increased propagation speed, whereas the NMDA receptor
antagonist MK-801 resulted in a broad reduction of propagation area, along with localized increases.
As FPI enables direct recording of electrical potential waveforms, frequency-domain analyses were
also conducted. Spontaneous activity in cortical organoids exhibited region-specific frequency
distributions, with gamma-band activity displaying distinct patterns compared to other frequency
bands. Additionally, in midbrain–striatal assembloids, electrophysiological activity was observed in
both regions. Connectivity analysis showed that treatment with 4-aminopyridine enhanced inter-
organoid connection strength. This large-scale, single-cell-resolved recording approach using UHD-
CMOS-MEAs facilitates comprehensive analysis of network connectivity, propagation velocity and
propagation area, and frequency characteristics. It represents a powerful platform for advancing our
understanding of the electrophysiological functions of brain organoids and assembloids, and holds
significant potential for drug screening and disease modeling in human neuroscience research.

1   Introduction

The human brain exhibits remarkably complex structures and functions, and the diverse dynamics of
neural activity are intricately involved in development, cognition, and disease. Brain organoids have
recently attracted attention as in vitro models that enable the reconstruction and analysis of these

brain functions. They are three-dimensional cultured tissues derived from pluripotent stem cells that self-organize into neural structures.(Lancaster et al., 2013; Lancaster and Knoblich, 2014; Giandomenico et al., 2021) To date, region-specific brain organoids mimicking the cerebral cortex (Lancaster et al., 2013), striatum (Miura et al., 2020), midbrain (Jo et al., 2016), and hippocampus (Sakaguchi et al., 2015) have been developed, allowing for the recapitulation of developmental processes and functional assessment of neural networks. In the context of compound screening, brain organoids are expected to serve as novel platforms that enable more precise prediction and evaluation of drug efficacy and toxicity than conventional two-dimensional cultures or animal models. Furthermore, because brain organoids can be generated from patient-derived induced pluripotent stem cells (iPSCs), they provide an opportunity to model human-specific disease phenotypes that are difficult to reproduce in animal systems. Indeed, various disease-related features have been reported using brain organoids, including amyloid-β and tau pathology in Alzheimer's disease (Raja et al., 2016; Gonzalez et al., 2018; Yan et al., 2018), overproduction of GABAergic neurons in autism spectrum disorder (Mariani et al., 2015), reduced neural and increased neurovascular cell populations in schizophrenia (Notaras et al., 2022), and disease-specific frequency characteristics and responses to contraindicated drugs in Dravet syndrome (Yokoi et al., 2023).

To construct more structurally complex in vitro models, increasing efforts have been directed toward generating assembloids by fusing organoids derived from different brain regions. Various types of assembloids have been developed to date, including cortical–striatal (Miura et al., 2020), midbrain–striatal (Ozgun et al., 2024), and midbrain–striatal–cortical assembloids (Reumann et al., 2023). More recently, in 2025, a model of the ascending neural sensory pathway was established by connecting organoids representing the somatosensory system, spinal cord, thalamus, and cortex(Kim et al., 2025).

As a functional application of brain organoids, their potential use in bio-computing systems has attracted growing interest. In 2022, the concept of "Organoid Intelligence" was proposed as an emerging interdisciplinary field that aims to harness the self-organizing capabilities of brain organoids to perform learning and memory functions within bio-computing platforms (Smirnova et al., 2023).

To achieve these goals, it is essential to establish detailed methods for evaluating the electrical activity of brain organoids. Currently, electrophysiological functional assessments of organoids employ techniques such as patch-clamp recordings, calcium imaging, and microelectrode array (MEA) recordings. MEA-based approaches allow for long-term, noninvasive monitoring of the electrical activity of brain organoids and assembloids, enabling the observation of time-dependent changes in network dynamics and the evaluation of pharmacological responses (Tasnim and Liu, 2022; Lv et al., 2023; Pan et al., 2024). Moreover, analysis of low-frequency signals (below 500 Hz) has been shown to be effective for assessing pharmacological responses in brain organoids (Yokoi et al., 2021). In addition, disease-specific frequency characteristics and responses to contraindicated compounds have also been successfully detected (Yokoi et al., 2023). Recently, the development of high-density (HD) CMOS-based MEAs utilizing complementary metal-oxide semiconductor (CMOS) technology has further improved the precision of neuronal network activity evaluation. Recordings from brain organoids using HD CMOS MEAs have enabled the quantification of dynamic single-cell firing rates and the calculation of propagation velocities (Schröter et al., 2022; Sharf et al., 2022). However, these evaluations have generally been limited to small recording areas and a restricted number of electrodes, falling short of capturing activity across the entire organoid.

In this study, we present an analytical approach for detailed evaluation of electrical activity in brain
organoids using field potential imaging (FPI) with an ultra-high-density (UHD) CMOS
microelectrode array (MEA) comprising 236,880 microelectrodes (10.52 μm × 10.52 μm) across a
wide sensing area of 32.45 mm² (Suzuki et al., 2023). Specifically, we conducted (1) single-cell spike
identification across the entire organoid and evaluation of neuronal network connectivity based on
spike activity; (2) calculation of propagation velocity and propagation area as novel endpoints of
network activity; and (3) spatial analysis of frequency-specific signal characteristics. Furthermore,
we applied the UHD-CMOS MEA to midbrain–striatal assembloids to record electrical activity from
both tissues and analyze inter-regional connectivity.

92 2   Materials and Methods

93 2.1   Culture of Human iPS Cells

Human induced pluripotent stem cells (hiPSCs) derived from a healthy donor (201B7 line, obtained
from RIKEN) were cultured in 6-well plates pre-coated with Vitronectin (07180, STEMCELL
Technologies) using mTeSR™ Plus medium (100-0276, STEMCELL Technologies). The culture
medium was partially replaced (50%) every 3 to 4 days. Once the cells reached confluence, they were
dissociated using Gentle Cell Dissociation Reagent (07174, STEMCELL Technologies) and
subsequently used for organoid generation.

100 2.2   Culture of Midbrain Organoids

101 Midbrain organoids used in this study were pre-generated and supplied by STEMCELL
102 Technologies. After acquisition, the organoids were cultured for four months using the STEMdiff™
103 Neural Organoid Maintenance Kit (100-0120, STEMCELL Technologies). The culture medium was
104 then replaced with BrainPhys™ Neuronal Medium (05792, STEMCELL Technologies), and
105 complete medium changes were carried out every 3 to 4 days to maintain the organoids.

106 2.3   Culture of Cerebral Organoids

Cortical organoids were generated using the STEMdiff™ Cerebral Organoid Kit (08570,
STEMCELL Technologies). Human iPS cells were seeded at a density of 9.0×10^3 cells per well,
using EB seeding medium. On days 2 and 4 of culture, 100 μL of EB formation medium was added
to each well. On day 5, the medium was replaced with induction medium, and the cultures were
incubated for 2 days. On day 7, organoids were embedded in Matrigel (354277, Corning) and
incubated in expansion medium for 3 days. From day 10 onward, the expansion medium was
replaced with maturation medium, and the organoids were cultured on an orbital shaker (COSH6, AS
ONE Corporation). Organoids were maintained in maturation medium for 3 months, with medium
changes carried out every 3 to 4 days. After 3 months, the culture medium was switched to
BrainPhys™ Neuronal Medium (05792, STEMCELL Technologies) for long-term maintenance.
Details of the media components and kit reagents are provided in Supplementary Table 1.

118 2.4   Culture of Midbrain–Striatal Assembloids

Midbrain–striatal assembloids were generated using the STEMdiff™ Midbrain Organoid
Differentiation Kit (100-1096, STEMCELL Technologies) and the STEMdiff™ Dorsal Forebrain
Organoid Differentiation Kit (08620, STEMCELL Technologies). To induce striatal differentiation,
organoids were cultured in medium supplemented with Activin A, IWP-2, and SR11237, following
the manufacturer's recommended conditions. Human iPSCs were seeded at a density of 3×10^6 cells

per well in Organoid Formation Medium supplemented with 10 μM Y-27632, using
AggreWell™800 24-well plates (34811, STEMCELL Technologies) pretreated with AggreWell
Rinsing Solution (07010, STEMCELL Technologies). From days 2 to 5, half-medium changes were
performed. On day 6, the culture medium was replaced with Midbrain Organoid Expansion Medium
for midbrain organoids and Striatal Organoid Expansion Medium for striatal organoids. Full medium
changes were performed every other day until day 25. On day 25, the respective Organoid
Differentiation Media were introduced, and the same schedule of complete medium changes
continued until day 43. From day 43 onward, cultures were maintained in Organoid Maintenance
Medium. On day 50 (midbrain) and day 60 (striatal), one organoid of each type was placed together
in a well of a 24-well plate and co-cultured on an orbital shaker (COSH6, AS ONE Corporation) to
form midbrain–striatal assembloids. These assembloids were maintained in Maintenance Medium for
an additional four months, with medium changes every 3 to 4 days. Details of the media components
and kit reagents are provided in Supplementary Table 2.

137   2.5   Immunocytochemistry

Cultured brain organoids were fixed in 4% paraformaldehyde (PFA) in phosphate-buffered saline
(PBS). After fixation, the organoids were embedded in Optimal Cutting Temperature (OCT)
compound (45833, Sakura Finetek Japan), and 10 μm-thick cryosections were prepared using a
cryostat (CM1950, Leica). The sections were first permeabilized with 0.2% Triton X-100 for 10 min
and then incubated in pre-blocking buffer (PBS containing 0.05% Triton X-100 and 5% goat serum)
at 4 °C for 60 min. Primary antibody incubation was carried out in pre-blocking buffer at 4 °C for 12
hours using anti-MAP2 antibody (ab5392, Abcam) and anti-tyrosine hydroxylase (TH) antibody
(ab137869, Abcam). For fluorescence detection, Alexa Fluor 488-conjugated anti-rabbit IgG
(A11040, Invitrogen) and Alexa Fluor 546-conjugated anti-chicken IgG (A11008, Invitrogen) were
diluted 1:1000 in pre-blocking buffer and incubated at room temperature for 1 hour. Nuclear staining
was performed with 1 μg/mL Hoechst 33258 (343-07961, Dojindo) for 1 hour at room temperature.
Fluorescently labeled organoid sections were imaged using a confocal laser scanning microscope
(AX R, Nikon), and image analysis was performed using ImageJ software (NIH).

151   2.6   Ultra-high-density CMOS MEA Recording

In this study, the electrical activity of brain organoids was recorded using the field potential imaging
(FPI) technique with an ultra-high-density CMOS-based microelectrode array (UHD-CMOS MEA)
system (Suzuki et al., 2023). The MEA chip comprised 236,880 electrodes, each measuring 10.52 μm
× 10.52 μm (110.67 μm²), with a total active recording area of 32.45 mm². Cerebral organoids,
midbrain organoids, and midbrain–striatal assembloids were individually mounted on the CMOS-
MEA system (Sony Semiconductor Solutions), and recordings were conducted under controlled
conditions (5% CO_2, 37 °C). Electrical signals were acquired at a sampling rate of either 2 kHz or
10 kHz. Baseline drift was removed using least-squares detrending applied to each 500-sample
segment, followed by offset correction to stabilize the signal.

161   2.7   Pharmacological Assays

Pharmacological experiments were conducted to evaluate the responsiveness of brain organoids to
neurotransmitter-related agents. In 4-month-old midbrain organoids, 3-(3,4-dihydroxyphenyl)-L-
alanine (L-DOPA; D0600, Tokyo Chemical Industry), a dopamine precursor, was applied, and
spontaneous electrical activity was recorded for 2 min at each concentration. In 5-month-old cerebral
organoids, the GABA$_A$ receptor antagonist picrotoxin (PTX; 168-17961, FUJIFILM Wako) and the
NMDA receptor antagonist MK-801 (M107-50MG, Sigma) were administered separately, and

spontaneous activity was recorded for 1 min at each concentration. Additionally, to assess changes in inter-regional connectivity within midbrain–striatal assembloids, 4-aminopyridine (4-AP; 016-02781, FUJIFILM Wako) was applied to 4-month-old assembloids, and spontaneous activity was recorded for 2 min at each concentration following 4-AP application.

2.8 Spike Detection and Soma identification

Spike detection was performed on voltage signals processed with a 100 Hz high-pass filter, using a threshold of $\pm 5.0\sigma$, where σ denotes the standard deviation of baseline noise during a quiescent period. Events exceeding this threshold were defined as spikes. For each detected spike, a waveform segment comprising 20 samples before and after the spike time point (41 samples in total) was extracted. Waveforms with a standard deviation below 13 were considered noise and excluded from further analysis. Soma identification was then performed based on spike timing correlations between electrode pairs. For all electrode pairs in which spikes were detected, inter-spike intervals (ISIs) were calculated. If the ISI between spikes on two electrodes was less than 10 ms, it was counted as a same-count event. Electrode pairs with ≥ 5 same-counts that also constituted $\geq 15\%$ of all possible spike pairs were designated as putative somatic electrode pairs. If the distance between electrodes in a putative pair was less than 50 µm, they were considered to originate from the same neuronal soma, and the location was identified as a somatic candidate site. In the case of midbrain organoids, spike events occurring during network bursts were excluded from soma identification to minimize false positives.

2.9 Z-score Analysis

For each neuronal pair, a synchronous spike was defined as an event in which a spike from one cell was followed by a spike from the other cell within 100 ms. The total number of synchronous spikes was counted for each pair. To assess statistical significance, 100 surrogate spike trains were generated by randomly shuffling the original spike time sequences while preserving each cell's inter-spike interval (ISI) distribution. For each surrogate dataset, the number of synchronous spikes was calculated using the same criteria. A z-score was calculated using the mean and standard deviation of the synchronous spike counts across the surrogate datasets, based on the synchronous spike count in the actual data. A z-score ≥ 3 was considered to indicate significant spike synchrony between the two cells, and such pairs were defined as having a functional connection. Based on the results of the z-score analysis, the number of significantly connected partner cells was counted for each neuron to determine its connection degree.

2.10 Propagation Analysis

Oscillatory activity was detected in voltage waveforms recorded using the UHD-CMOS MEA. Electrodes that exhibited signal amplitudes ≥ 30 µV during these periods were defined as propagating electrodes, while all others were treated as noise electrodes and excluded from the analysis. For each propagating electrode, the peak time—defined as the interval from the onset of oscillation ($t = 0$) to the point of maximum amplitude—was calculated. A time-series dataset was then constructed by counting the number of electrodes that reached their peak within each time bin. The propagation speed of network activity was estimated by computing the temporal derivative of this time series. Propagation speed was calculated for each oscillatory event detected in individual traces, and the average of these values was reported as the final result.

2.11 Frequency Band Power Analysis of Oscillatory Waveforms

Detected oscillatory waveforms were processed using zero-phase filtering with a finite impulse
response (FIR) bandpass filter to isolate signals within distinct frequency bands. The potential
amplitude for each frequency band was calculated from the filtered waveforms. The five frequency
bands analyzed were defined as follows: delta (0.5–3 Hz), theta (4–7 Hz), alpha (8–11 Hz), beta (12–
29 Hz), and gamma (30–100 Hz). For each band, power values were standardized using the mean and
standard deviation across all electrodes. The resulting normalized data were subjected to
unsupervised clustering using the k-means algorithm implemented in MATLAB (MathWorks) to
classify patterns of frequency-specific potential dynamics.

2.12 Analysis of Connection Strength in Midbrain–Striatal Assembloids

For each neuronal pair, a synchronous spike was defined as a spike from one cell followed by a spike
from the other cell within 100 ms. The number of synchronous spikes was counted for each pair. To
minimize the influence of differences in firing rates between cells, the synchronous spike count was
normalized by the mean spike count of the respective cell pair. The resulting value was defined as the
connection strength. Connection strength was calculated separately for three groups: striatal
organoids, midbrain organoids, and midbrain–striatal assembloids.

3 Results

3.1 Single-Cell Firing Rate and Intercellular Connectivity Analysis of Midbrain Organoids
Following L-DOPA Administration

Immunofluorescence staining of midbrain organoids derived from human iPSCs and cultured for 4
months confirmed the expression of MAP2, a marker for mature neurons, and TH, a marker for
dopaminergic neurons (Figure. 1A). In addition, non-TH-positive neurons were present. The
midbrain organoid was mounted on a UHD-CMOS MEA (Figure. 1B), and spontaneous electrical
activity was recorded across the entire organoid, resulting in the identification of 404 individual cells.
A raster plot and histogram of the spontaneous spike activity over a 2-minute period from these
identified cells are shown in Figure. 1C. Network bursts, in which almost all cells spiked
simultaneously, were observed (indicated by peaks in the histogram and vertical black lines in the
raster plot in Figure. 1C), suggesting the formation of functional neuronal circuits within the
midbrain organoid. Figure 1D presents a spatial map of total spike counts over 2 minutes for the 404
identified cells, with dot size representing the number of spikes. Cells whose spike count increased to
≥150% following administration of L-DOPA (30 μM) compared to before treatment are shown in
red, while those with a decrease to ≤50% are shown in blue. The proportion of cells with increased
activity was 19.3% at 0.3 μM, 39.4% at 3 μM, and 59.7% at 30 μM. Conversely, the proportion of
cells with decreased activity was 10.1%, 2.23%, and 0.495%, respectively, while the proportions of
non-responsive cells were 70.5%, 58.4%, and 39.9%, respectively (Figure. 1E). The single-cell
resolution afforded by the UHD-CMOS MEA enabled the detection not only of increased spike
activity following L-DOPA treatment but also of cells that were non-responsive or showed
suppressed activity at lower doses. To evaluate network activity in the midbrain organoid, z-scores
were calculated for cell pairs that spiked within 100 ms. In Figure. 2A, black lines indicate cell pairs
with a z-score ≥ 3 before L-DOPA administration, while blue lines indicate pairs whose z-scores
increased after treatment. L-DOPA administration led to a dose-dependent increase in connection
strength (Figure. 2B). The distribution of z-scores also shifted to the right in accordance with L-
DOPA concentration (Figure. 2C), suggesting enhanced synchrony within the network. To
quantitatively assess changes in connection strength, cell pairs with z-scores ≥ 3 were defined as
connected, and the average number of connected cells per cell was calculated. The number of
connections per cell was 333.2 ± 84.6 at baseline, 335.8 ± 79.2 at 0.3 μM, 380.5 ± 42.8 at 3 μM, and

382.7 ± 45.2 at 30 µM, with significant increases observed at 3 µM and 30 µM compared to baseline
(Figure. 2D). These findings indicate that the UHD-CMOS MEA enables both single-cell spike
analysis and intercellular connectivity evaluation in brain organoids.

3.2   Analysis of Propagation Velocity and Propagation Area in Cortical Organoids

The ultra-high-density (UHD) CMOS MEA, with its high temporal and spatial resolution, enabled
the derivation of novel endpoints for evaluating network activity in brain organoids—specifically,
propagation velocity and propagation area. To assess propagation velocity, the GABA_A receptor
antagonist picrotoxin (PTX) was applied to 5-month-old cerebral organoids. Figure 3A shows
representative propagation waveforms recorded during spontaneous oscillations, with the
corresponding electrode positions indicated by red arrows in the top-left panel of Figure. 3B
(Before). Peak timing maps of oscillatory waveforms across electrodes, recorded before and after
PTX administration, are presented in Figure. 3B, illustrating that neural activity propagated
throughout the entire organoid. Propagation velocity was calculated by differentiating the number of
electrodes reaching peak voltage per unit time (Figure. 3C). The baseline propagation velocity was
0.0166 ± 0.000715 mm²/ms, which significantly increased to 0.0202 ± 0.000523, 0.0193 ± 0.000394,
and 0.0196 ± 0.000257 mm²/ms following administration of 0.1, 1, and 10 µM PTX, respectively
(Figure. 3C). Propagation area was assessed before and after treatment with the NMDA receptor
antagonist MK-801. In the maps shown in Figure. 3D, electrodes exhibiting propagation activity are
indicated in black. At 0.1 µM MK-801, the propagation area increased from 0.851 mm² to 1.05 mm²,
although the change was not statistically significant (Figure. 3E). By contrast, at 1 µM MK-801, the
propagation area decreased markedly across most of the organoid, with a localized increase observed
only in the lower right region (Figure. 3D). At the whole-organoid level, synaptic transmission
blockade resulted in a significant reduction of the propagation area to 0.384 mm² (Figure. 3E). These
findings indicate that the analysis of propagation velocity and propagation area using UHD-CMOS
MEAs provides an effective approach for evaluating compound-induced responses related to synaptic
transmission in brain organoids.

3.3   Frequency Characteristics of Cerebral Organoids

Field potential imaging (FPI) using the UHD-CMOS MEA provides voltage waveforms, making it
suitable for frequency analysis in a manner analogous to conventional MEA recordings. Figure 4A
displays a representative raw waveform recorded from a single electrode in a 5-month-old cerebral
organoid, alongside corresponding signals filtered into delta (0.5–3 Hz), theta (4–7 Hz), alpha (8–
11 Hz), beta (12–29 Hz), and gamma (30–100 Hz) bands using a finite impulse response (FIR)
bandpass filter. When a network burst occurred across the entire cortical organoid, slow-wave field
potentials were observed, as shown in Figure 4A. Field potentials were acquired from 46,630
electrodes, enabling frequency analysis at the whole-organoid level (Figure 4B). Figure 4B shows
heatmaps of standardized potential intensities for each frequency band across all electrodes. Delta to
beta bands exhibited a gradient of increasing intensity from the outer edge toward the center, with the
strongest signals observed in the lower-right region. In contrast, gamma-band activity showed a
distinct pattern, characterized by weaker signals in the lower half and stronger signals in the upper
half of the organoid. To delineate frequency-specific spatial domains within the organoid, k-means
clustering analysis was performed, segmenting the organoid into nine distinct clusters (Figure. 4C).
Comparison of the frequency intensity profiles across clusters revealed that Clusters 1–3 exhibited
low intensity across all bands, whereas Clusters 5, 7, and 8 showed uniformly high intensity. Clusters
6 and 9 exhibited high intensity in the delta to beta bands but relatively low gamma-band activity
(Figure. 4D). Frequency analysis of activity waveforms detected from 46,630 electrodes on the

UHD-CMOS MEA demonstrated that region-specific frequency characteristics can be extracted from
the brain organoid.

3.4 Inter-Regional Connection Analysis in Midbrain–Striatal Assembloids

Using the UHD-CMOS MEA, we recorded the electrical activity of midbrain–striatal assembloids
and evaluated inter-regional connection strength based on single-cell firing activity. Figure 5A shows
a heatmap of peak voltage distribution over a one-minute recording period, where the upper part
corresponds to the striatum and the lower part to the midbrain organoid. Electrical activity was
detected in both regions, indicating that each tissue was functionally active. To assess the effects of
pharmacological intervention, 30 µM of 4-aminopyridine (4-AP), a potassium channel blocker, was
administered to the assembloid, and changes in firing frequency and inter-regional connectivity were
evaluated. Figure 5B presents raster plots of individual cells (red: striatum, yellow: midbrain).
Following 4-AP administration, spike counts increased in both the midbrain and striatum (Figure.
5C). Connection strength was calculated by normalizing the number of synchronized spikes (defined
as spikes occurring within 100 ms between neuron pairs) by the average firing rate of each pair.
Figure 5D shows the connection strength within the striatum, within the midbrain, and between the
midbrain and striatum before and after 4-AP treatment. The connection strength increased from
0.0360 to 0.0946 within the striatum and from 0.181 to 0.377 within the midbrain. Furthermore, the
inter-regional connection strength between the midbrain and striatum also increased from 0.0305 to
0.0632. Single-cell-based network connectivity analysis using the UHD-CMOS MEA revealed that
changes in inter-organoid information transmission within assembloids can be detected following
compound administration.

4   Discussion

In this study, we performed large-scale recordings of brain organoids using field potential imaging
(FPI) with an ultra-high-density (UHD) CMOS microelectrode array (MEA) comprising 236,880
microelectrodes (each measuring 10.52 µm × 10.52 µm) and covering a broad sensing area of
32.45 mm², in order to precisely characterize the detailed electrical activity of brain organoids. Based
on single-cell activity, we analyzed neuronal network connectivity, propagation velocity, propagation
area, and frequency characteristics.

L-Dopa, a precursor to dopamine, is taken up by neurons and decarboxylated by Aromatic L-Amino
Acid Decarboxylase (AADC), resulting in the production of dopamine, which is subsequently
released into the extracellular space. When L-Dopa was administered to midbrain organoids, a
decrease in activity was observed in 10.1% of cells at a concentration of 0.3 µM (Figure 1E). TH-
positive dopaminergic neurons are known to express D2 autoreceptors (Beaulieu and Gainetdinov,
2011), and this suppression of activity is thought to be due to D2 receptor activation following
dopamine release. However, at higher concentrations, this suppression was no longer observed
(Figure. 1E), and instead, an enhancement of network activity was seen throughout the midbrain
organoids (Figure. 2A). The midbrain organoids used in this study contained not only TH-positive
neurons (Figure. 1A) but also likely included vglut2-positive glutamatergic neurons, which are
known to be present in the midbrain, as well as cells expressing D1-type dopamine receptors
(Yamaguchi et al., 2007; Yamaguchi et al., 2011; Morales and Root, 2014). It is possible that
excitatory input from these cells indirectly disinhibited or activated dopaminergic neurons, leading to
increased activity at the single-cell level (Figure. 1E) and strengthened network connectivity across
the entire organoid (Figure. 2A). The use of UHD CMOS MEA enabled quantitative evaluation of

both the firing characteristics of individual cells and the neural network activity based on single-cell
firing patterns within the brain organoids.

Based on the capabilities of the UHD-CMOS-MEA, novel endpoints for the propagation speed and
area of network activity were established. In cortical organoids, the propagation speed of network
activity increased following administration of picrotoxin (PTX), a GABA$_A$ receptor antagonist. This
observation aligns with previous reports showing that administration of bicuculline methiodide—
another GABA$_A$ receptor antagonist—increased propagation speed in cortical slices derived from
human epilepsy patients (Covelo et al., 2025). Although the compounds differ, the results may reflect
similar physiological effects mediated by GABA$_A$ receptor inhibition. Following administration of
MK-801, an NMDA receptor antagonist, a reduction in the propagation area was observed in most
regions of the organoids (Figure 3D, E). Similar reductions in activity propagation caused by NMDA
receptor antagonists have been reported in in vivo EEG recordings from rats and in brain slice
experiments (Morales-Villagrán et al., 1996; Fukuda et al., 1998), suggesting that the responses
observed in this study may reflect physiological phenomena in the intact brain. Notably, although the
propagation area decreased across the majority of the organoid, a localized enhancement in
propagation was detected in the lower-right region following treatment with 1 µM MK-801 (Figure
3D). This focal increase in activity may be attributable to disinhibition resulting from impaired
function of GABAergic neurons. Previous studies have described both direct suppression of
inhibitory neuron activity via NMDA receptor blockade and indirect effects through reduced
excitatory input (Li et al., 2002; Homayoun and Moghaddam, 2007). These mechanisms are
considered to underlie the excitotoxic effects of NMDA receptor antagonists. Due to self-
organization, brain organoids likely exhibit regional variability in the ratio of excitatory to inhibitory
neurons. Consequently, in areas with a lower proportion of inhibitory neurons, the loss of inhibitory
control induced by MK-801 may be more pronounced, potentially leading to enhanced excitatory
activity mediated by AMPA receptors. Analysis of propagation velocity and propagation area using
the UHD-CMOS MEA demonstrated that this method is effective for evaluating compound responses
related to synaptic transmission in brain organoids. Moving forward, integrating analyses of
propagation pathways may enable the identification of regions exhibiting disease-related activity
abnormalities or neural circuits serving as initiation sites for compound responses.

Since FPI represents voltage waveforms, frequency analysis can be conducted similarly to
conventional MEA measurements. Traditional frequency analyses of brain organoids using standard
MEAs were limited by the small number of electrodes, typically ranging from 16 to 64. In this study,
activity was acquired from 46,630 electrodes using the UHD CMOS MEA, which enabled a
comprehensive analysis of frequency distributions across the entire organoid. Delta band potentials
increased from the periphery toward the center. Alpha to beta band activity also intensified toward
the center, with particularly strong signals observed in the lower-right region of the organoid. In
contrast, gamma band activity displayed a distinct pattern—being stronger in the upper region and
weaker in the lower region (Figure 4B). Clustering analysis enabled visualization of the organoid's
structural features based on frequency-specific characteristics (Figure 4C). However, because the
organoids were measured without being anchored to the electrodes, the variability in adhesion
between the organoid and electrodes must be considered. The periphery of the organoid likely had
fewer cells adhering to the electrodes, which may explain the consistently weak potentials observed
in clusters 1, 2, and 3 across all frequency bands. In contrast, clusters 6, 7, 8, and 9, located deeper
within the organoid, exhibited variations in gamma band potentials (Figure 4C), indicating that in
regions with better electrode contact, the intrinsic structural frequency properties of the organoid
were more accurately reflected. The distribution of gamma band activity may be associated with the
expression of PV-positive GABAergic neurons (Cardin et al., 2009), which are known to be essential

for gamma wave generation in the cortex. In clusters 6 and 9, where gamma activity was weak while
other frequency bands were strong, it is likely that PV-positive GABAergic neurons were either
absent or functionally immature. Previous studies have shown that the expression of GABAergic
neurons in cortical organoids increases after six months of culture (Trujillo et al., 2019). Given that
the organoids in this study were five months old, it is plausible that some regions lacked sufficient
expression or functional maturity of PV-positive GABAergic neurons. Comparing the
electrophysiological frequency characteristics of organoids with cell type identity and the
histochemically organized circuit architecture within the organoids remains an important challenge
for future studies. Moreover, samples that exhibited global oscillatory activity throughout the entire
organoid, as observed here, were extremely rare. Even when derived from the same iPSC dish, some
organoids exhibited oscillations only in localized areas, while others showed no detectable electrical
activity. Therefore, establishing reproducible methods for reliably generating organoids with
consistent, whole-organoid activity remains a critical goal. Despite these challenges, this study
demonstrates that UHD CMOS MEA technology enables comprehensive frequency distribution
analysis of entire brain organoids. Importantly, the ability to detect spatial characteristics of gamma
band activity—crucial for the evaluation of neurological disorders such as epilepsy, Alzheimer's
disease, and autism—highlights the utility of this method for assessing frequency-specific disease
models (Herrmann and Demiralp, 2005; Başar, 2013; McNally and McCarley, 2016; Guan et al.,
2022).

Currently, the evaluation of assembloids is primarily performed using techniques such as optogenetic
observation with adeno-associated viruses (AAV), calcium imaging, patch-clamp recordings, and
conventional multi-electrode arrays (MEAs) (Andersen et al., 2020; Miura et al., 2020; Reumann et
al., 2023; Ozgun et al., 2024). However, these methods have limitations in terms of temporal and
spatial resolution, as well as invasiveness to the cells. In contrast, the UHD CMOS MEA enabled the
non-invasive detection of electrical activity at single-cell resolution across the entire assembloid, with
both high temporal and spatial precision (Figure 5A). When 4-Aminopyridine (4-AP), a potassium
channel blocker, was applied to midbrain–striatal assembloids, increased firing activity was observed
in both the midbrain and striatal regions (Figure 5B). Furthermore, an enhancement of inter-regional
connectivity was also detected (Figure 5C). A current limitation of this experimental setup is the
inability to selectively apply compounds to individual tissues, leading to simultaneous exposure of
both regions. This complicates the mechanistic analysis of inter-regional connection enhancement.
One potential solution is to combine UHD CMOS MEA recordings with optogenetic techniques,
such as AAV-mediated gene delivery, enabling targeted stimulation of specific cell types and
subsequent recording of their responses. Additionally, integrating UHD CMOS MEA with
Microphysiological System (MPS) devices containing microfluidic channels would allow for region-
specific compound administration. MPS platforms seed organoids into individual chambers
connected via microfluidic pathways, offering both structural control and the ability to measure not
only tissue-specific activity but also axonal signals transmitted through the connecting channels. A
system that combines these technologies with UHD CMOS MEA for assessing inter-organ
connectivity in assembloids may be particularly valuable for investigating projection deficits caused
by neurological disorders. It may also be applied to disease models involving peripheral nervous
system damage, such as those using sensory neuron–spinal cord assembloids.

In this study, we established a methodology for large-scale recording of brain organoids using UHD-
CMOS MEA, along with analytical techniques to assess network connectivity, propagation speed,
propagation area, and frequency dynamics at the single-cell level. While further refinement is needed
to achieve stable organoid generation and recording techniques, and to elucidate the relationship
between electrophysiological activity and structural characteristics, this approach holds promise for

advancing our understanding of the functional electrophysiology of brain organoids and assembloids.
Moreover, it is expected to be valuable for compound screening and the evaluation of human
neurological disease models.

5 Conflict of Interest

441 The authors declare no conflict of interest.

6 Author Contributions

443 Conceptualization, R.Y. and I.S.; writing—original draft preparation, R.Y. and I.S.; writing—review
444 and editing R.Y., I.S., Y.I. and N.M.; Formal analysis, R.Y., Y.I. and N.M.; cell line culture, R.Y.;
445 immunostaining, R.Y.; funding acquisition, I.S. All authors have read and agreed to the published
446 version of the manuscript.

7 Funding

448 This study was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI
449 (Grant Number 23H03726 and 25K21561).

8 Acknowledgments

451 This study was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI
452 Grant Number 23H03726 and 25K21561. This study was also supported by the UHD-CMOS-MEA
453 system with Sony semiconductor solutions Inc.

9 Data Availability Statement

455 The data and scripts that support the findings of this study are available from the corresponding
456 author upon reasonable request.

10 Reference

Andersen, J., Revah, O., Miura, Y., Thom, N., Amin, N.D., Kelley, K.W., et al. (2020). Generation of
Functional Human 3D Cortico-Motor Assembloids. *Cell* 183(7)**,** 1913-1929.e1926. doi:
10.1016/j.cell.2020.11.017.

Başar, E. (2013). A review of gamma oscillations in healthy subjects and in cognitive impairment.
International Journal of Psychophysiology 90(2)**,** 99-117. doi:
https://doi.org/10.1016/j.ijpsycho.2013.07.005.

Beaulieu, J.M., and Gainetdinov, R.R. (2011). The physiology, signaling, and pharmacology of
dopamine receptors. *Pharmacol Rev* 63(1)**,** 182-217. doi: 10.1124/pr.110.002642.

Cardin, J.A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., et al. (2009). Driving
fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature* 459(7247)**,**
663-667. doi: 10.1038/nature08002.

Covelo, J., Camassa, A., Sanchez-Sanchez, J.M., Manasanch, A., Porta, L.D., Cancino-Fuentes, N.,
et al. (2025). Spatiotemporal network dynamics and structural correlates in the human
cerebral cortex in vitro. *Progress in Neurobiology* 246**,** 102719. doi:
https://doi.org/10.1016/j.pneurobio.2025.102719.

Fukuda, M., Hata, Y., Ohshima, M., and Tsumoto, T. (1998). Role of NMDA receptors in the
propagation of excitation in rat visual cortex as studied by optical imaging. *Neuroscience*
Research 31(1)**,** 9-21. doi: https://doi.org/10.1016/S0168-0102(98)00018-2.

Giandomenico, S.L., Sutcliffe, M., and Lancaster, M.A. (2021). Generation and long-term culture of
advanced cerebral organoids for studying later stages of neural development. *Nat Protoc*
16(2)**,** 579-602. doi: 10.1038/s41596-020-00433-w.

Gonzalez, C., Armijo, E., Bravo-Alegria, J., Becerra-Calixto, A., Mays, C.E., and Soto, C. (2018).
Modeling amyloid beta and tau pathology in human cerebral organoids. *Molecular Psychiatry*
23(12)**,** 2363-2374. doi: 10.1038/s41380-018-0229-8.

Guan, A., Wang, S., Huang, A., Qiu, C., Li, Y., Li, X., et al. (2022). The role of gamma oscillations
in central nervous system diseases: Mechanism and treatment. *Front Cell Neurosci* 16**,**
962957. doi: 10.3389/fncel.2022.962957.

Herrmann, C.S., and Demiralp, T. (2005). Human EEG gamma oscillations in neuropsychiatric
disorders. *Clinical Neurophysiology* 116(12)**,** 2719-2733. doi:
https://doi.org/10.1016/j.clinph.2005.07.007.

Homayoun, H., and Moghaddam, B. (2007). NMDA receptor hypofunction produces opposite effects
on prefrontal cortex interneurons and pyramidal neurons. *J Neurosci* 27(43)**,** 11496-11500.
doi: 10.1523/jneurosci.2213-07.2007.

Jo, J., Xiao, Y., Sun, A.X., Cukuroglu, E., Tran, H.D., Göke, J., et al. (2016). Midbrain-like
Organoids from Human Pluripotent Stem Cells Contain Functional Dopaminergic and
Neuromelanin-Producing Neurons. *Cell Stem Cell* 19(2)**,** 248-257. doi:
10.1016/j.stem.2016.07.005.

Kim, J.-i., Imaizumi, K., Jurjuț, O., Kelley, K.W., Wang, D., Thete, M.V., et al. (2025). Human
assembloid model of the ascending neural sensory pathway. *Nature*. doi: 10.1038/s41586-
025-08808-3.

Lancaster, M.A., and Knoblich, J.A. (2014). Generation of cerebral organoids from human
pluripotent stem cells. *Nat Protoc* 9(10)**,** 2329-2340. doi: 10.1038/nprot.2014.158.

Lancaster, M.A., Renner, M., Martin, C.A., Wenzel, D., Bicknell, L.S., Hurles, M.E., et al. (2013).
Cerebral organoids model human brain development and microcephaly. *Nature* 501(7467)**,**
373-379. doi: 10.1038/nature12517.

Li, Q., Clark, S., Lewis, D.V., and Wilson, W.A. (2002). NMDA receptor antagonists disinhibit rat
posterior cingulate and retrosplenial cortices: a potential mechanism of neurotoxicity. *J*
Neurosci 22(8)**,** 3070-3080. doi: 10.1523/jneurosci.22-08-03070.2002.

Lv, S., He, E., Luo, J., Liu, Y., Liang, W., Xu, S., et al. (2023). Using Human-Induced Pluripotent
Stem Cell Derived Neurons on Microelectrode Arrays to Model Neurological Disease: A
Review. *Adv Sci (Weinh)* 10(33)**,** e2301828. doi: 10.1002/advs.202301828.

Mariani, J., Coppola, G., Zhang, P., Abyzov, A., Provini, L., Tomasini, L., et al. (2015). FOXG1-
Dependent Dysregulation of GABA/Glutamate Neuron Differentiation in Autism Spectrum
Disorders. *Cell* 162(2)**,** 375-390. doi: 10.1016/j.cell.2015.06.034.

McNally, J.M., and McCarley, R.W. (2016). Gamma band oscillations: a key to understanding
schizophrenia symptoms and neural circuit abnormalities. *Curr Opin Psychiatry* 29(3)**,** 202-
210. doi: 10.1097/yco.0000000000000244.

Miura, Y., Li, M.Y., Birey, F., Ikeda, K., Revah, O., Thete, M.V., et al. (2020). Generation of human
striatal organoids and cortico-striatal assembloids from human pluripotent stem cells. *Nat*
Biotechnol 38(12)**,** 1421-1430. doi: 10.1038/s41587-020-00763-w.

Morales-Villagrán, A., Ureña-Guerrero, M.E., and Tapia, R. (1996). Protection by NMDA receptor
antagonists against seizures induced by intracerebral administration of 4-aminopyridine.
European Journal of Pharmacology 305(1)**,** 87-93. doi: https://doi.org/10.1016/0014-
2999(96)00157-4.

Morales, M., and Root, D.H. (2014). Glutamate neurons within the midbrain dopamine regions.
Neuroscience 282**,** 60-68. doi: 10.1016/j.neuroscience.2014.05.032.

Notaras, M., Lodhi, A., Dündar, F., Collier, P., Sayles, N.M., Tilgner, H., et al. (2022). Schizophrenia
is defined by cell-specific neuropathology and multiple neurodevelopmental mechanisms in
patient-derived cerebral organoids. *Mol Psychiatry* 27(3)**,** 1416-1434. doi: 10.1038/s41380-
021-01316-6.

Ozgun, A., Lomboni, D.J., Aylsworth, A., Macdonald, A., Staines, W.A., Martina, M., et al. (2024).
Unraveling the assembloid: Real-time monitoring of dopaminergic neurites in an inter-
organoid pathway connecting midbrain and striatal regions. *Mater Today Bio* 25**,** 100992. doi:
10.1016/j.mtbio.2024.100992.

Pan, T., Jaklic, D.C., Vaid, S., Lin, G., VanHeyningen, D., and Dang, L.T. (2024). A Multi-Electrode
Array Platform for Modeling Epilepsy Using Human Pluripotent Stem Cell-Derived Brain
Assembloids. *J Vis Exp* (211). doi: 10.3791/67396.

Raja, W.K., Mungenast, A.E., Lin, Y.T., Ko, T., Abdurrob, F., Seo, J., et al. (2016). Self-Organizing
3D Human Neural Tissue Derived from Induced Pluripotent Stem Cells Recapitulate
Alzheimer's Disease Phenotypes. *PLoS One* 11(9)**,** e0161969. doi:
10.1371/journal.pone.0161969.

Reumann, D., Krauditsch, C., Novatchkova, M., Sozzi, E., Wong, S.N., Zabolocki, M., et al. (2023).
In vitro modeling of the human dopaminergic system using spatially arranged ventral
midbrain-striatum-cortex assembloids. *Nat Methods* 20(12)**,** 2034-2047. doi: 10.1038/s41592-
023-02080-x.

Sakaguchi, H., Kadoshima, T., Soen, M., Narii, N., Ishida, Y., Ohgushi, M., et al. (2015). Generation
of functional hippocampal neurons from self-organizing human embryonic stem cell-derived
dorsomedial telencephalic tissue. *Nat Commun* 6**,** 8896. doi: 10.1038/ncomms9896.

Schröter, M., Wang, C., Terrigno, M., Hornauer, P., Huang, Z., Jagasia, R., et al. (2022). Functional
imaging of brain organoids using high-density microelectrode arrays. *MRS Bull* 47(6)**,** 530-
544. doi: 10.1557/s43577-022-00282-w.

Sharf, T., van der Molen, T., Glasauer, S.M.K., Guzman, E., Buccino, A.P., Luna, G., et al. (2022).
Functional neuronal circuitry and oscillatory dynamics in human brain organoids. *Nature*
Communications 13(1)**,** 4403. doi: 10.1038/s41467-022-32115-4.

Smirnova, L., Morales Pantoja, I.E., and Hartung, T. (2023). Organoid intelligence (OI) - The
ultimate functionality of a brain microphysiological system. *Altex* 40(2)**,** 191-203. doi:
10.14573/altex.2303261.

Suzuki, I., Matsuda, N., Han, X., Noji, S., Shibata, M., Nagafuku, N., et al. (2023). Large-Area Field
Potential Imaging Having Single Neuron Resolution Using 236 880 Electrodes CMOS-MEA
Technology. *Adv Sci (Weinh)* 10(20)**,** e2207732. doi: 10.1002/advs.202207732.

Tasnim, K., and Liu, J. (2022). Emerging Bioelectronics for Brain Organoid Electrophysiology. *J*
Mol Biol 434(3)**,** 167165. doi: 10.1016/j.jmb.2021.167165.

Trujillo, C.A., Gao, R., Negraes, P.D., Gu, J., Buchanan, J., Preissl, S., et al. (2019). Complex
Oscillatory Waves Emerging from Cortical Organoids Model Early Human Brain Network
Development. *Cell Stem Cell* 25(4)**,** 558-569.e557. doi: 10.1016/j.stem.2019.08.002.

Yamaguchi, T., Sheen, W., and Morales, M. (2007). Glutamatergic neurons are present in the rat
ventral tegmental area. *Eur J Neurosci* 25(1)**,** 106-118. doi: 10.1111/j.1460-
9568.2006.05263.x.

Yamaguchi, T., Wang, H.L., Li, X., Ng, T.H., and Morales, M. (2011). Mesocorticolimbic
glutamatergic pathway. *J Neurosci* 31(23)**,** 8476-8490. doi: 10.1523/jneurosci.1598-11.2011.

Yan, Y., Song, L., Bejoy, J., Zhao, J., Kanekiyo, T., Bu, G., et al. (2018). Modeling
Neurodegenerative Microenvironment Using Cortical Organoids Derived from Human Stem
Cells. *Tissue Eng Part A* 24(13-14)**,** 1125-1137. doi: 10.1089/ten.TEA.2017.0423.

Yokoi, R., Nagafuku, N., Ishibashi, Y., Matsuda, N., and Suzuki, I. (2023). Contraindicated Drug
Responses in Dravet Syndrome Brain Organoids Utilizing Micro Electrode Array Assessment
Methods. *Organoids* 2(4)**,** 177-191.

Yokoi, R., Shibata, M., Odawara, A., Ishibashi, Y., Nagafuku, N., Matsuda, N., et al. (2021).
Analysis of signal components < 500 Hz in brain organoids coupled to microelectrode arrays:
A reliable test-bed for preclinical seizure liability assessment of drugs and screening of
antiepileptic drugs. *Biochem Biophys Rep* 28**,** 101148. doi: 10.1016/j.bbrep.2021.101148.

Figure 1. Single-cell spike analysis of midbrain organoids. (A) Immunofluorescence image of a
human midbrain organoid cultured for 4 months. Green: MAP2; red: TH; blue: Hoechst 33258. Scale
bar = 200 μm. (B) Midbrain organoid mounted on a UHD-CMOS MEA. The sensing area is
indicated by a yellow square. Scale bar = 1 mm. (C) Raster plot and histogram of spontaneous
neuronal spikes over a 2-minute period before and after L-DOPA administration. Scale bar = 30 sec.
(D) Spatial map of spontaneous spikes over a 2-minute period from 404 detected cells. Each dot
represents a cell body, and dot size reflects spike frequency. Red: ≥150% vs. before; blue: ≤50% vs.
before. (E) Change in spike frequency following L-DOPA administration. The proportion of cells
showing altered activity relative to before administration is indicated. Red: ≥150% vs. before; blue:
≤50% vs. before; black: no change.

Figure 2. Neural network analysis of midbrain organoids. Changes in connection strength between
cell pairs before and after L-DOPA administration. To evaluate network activity, z-scores were
calculated for cell pairs that spiked within 100 ms. (A) Cell connection map. Black lines indicate cell
pairs with a z-score $\geq$ 3; blue lines indicate pairs whose z-score increased after L-DOPA treatment.
(B) Heatmap of z-scores for 404 detected cells. (C) Distribution of z-scores. (D) Change in the
number of connected cells per individual cell.

Figure 3. Propagation analysis of cortical organoids. (A) Spontaneous activity propagation waveform color-coded by the timing of voltage peaks. The electrode used for pickup is indicated by a red arrow in (B) Before. (B) Delay maps of oscillation peak timing before and after picrotoxin (PTX) administration. (C) Propagation velocity before and after PTX administration. The average propagation velocity was calculated per oscillation event (before: n = 4; 0.1 μM: n = 2; 1 μM: n = 6; 10 μM: n = 4). Data were analyzed using one-way ANOVA followed by post hoc Dunnett's test (**p < 0.01 vs. before). (D) Electrode maps showing propagation before and after MK-801 administration; black dots indicate propagation electrodes. (E) Propagation area before and after MK-801 administration. The propagation area was calculated per oscillation event (before: n = 6; 0.1 μM: n = 6; 1 μM: n = 1). Data were analyzed using one-way ANOVA followed by post hoc Dunnett's test (*p < 0.05 vs. before).

Figure 4. Frequency distribution characteristics of cortical organoids. (A) Spontaneous oscillation
waveforms for each frequency band. From top to bottom: raw waveform, delta, theta, alpha, beta,
gamma. (B) Spatial distribution maps of normalized signal intensity for each frequency band. (C)
Clustering map showing nine groups classified based on the intensity profiles of five frequency
bands. (D) Heatmap of frequency band intensities for each cluster.

Figure 5. Analysis of inter-regional connectivity in midbrain–striatal assembloids. (A) Peak potential
map of spontaneous activity over a 1-minute recording. Upper region: striatum; lower region:
midbrain. (B) Raster plot of spontaneous spikes from individual cells over 1 minute. Red: striatum;
yellow: midbrain. Scale bar = 15 sec. (C) Total spike count before and after 4-AP administration. The
total number of spikes per minute is shown for each region. Gray: striatum; black: midbrain. (D)
Connection strength before and after 4-AP administration. Strength was calculated by normalizing
the number of synchronous spikes (within 100 ms) by the mean spike count of each cell pair. Gray:
within striatum; black: within midbrain; white: between midbrain and striatum.