The landscape and consequences of transcription stress

Osama Hidmi[1], Sara Oster Flayshman[1], Diala Shatleh[1], Jonathan Monin[1] and Rami I. Aqeilan[1,2,✉]

[1]The Concern Foundation Laboratories, The Lautenberg Center for Immunology and Cancer Research, Department of Immunology and Cancer Research-IMRIC, Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel
[2]Lead contact

✉ ramiaq@mail.huji.ac.il (R.I.A.)

Abstract

Cancer is characterized by uncontrolled proliferation and hypertranscription of oncogenes, leading to transcription stress, a major source of DNA double-strand breaks (DSBs) that threaten genomic stability. However, the landscape and consequences of transcription stress remain incompletely understood. Here, using genome-wide maps of DSBs identified by sBLISS alongside transcription stress markers, we reveal that transcription stress regions are highly enriched at the super-enhancer (SE) regulatory landscape. Transcriptional induction through SEs markedly amplifies DSB accumulation compared to non-SE-driven transcription. γH2AX mapping revealed selective enrichment at transcription stress sites, with only a subset of DSB-enriched genes showing strong γH2AX marking. Notably, genes with high DSBs and strong γH2AX signals exhibited significantly higher DSB turnover and repair than γH2AX-low genes, correlating with increased vulnerability to mutagenesis. These findings identify SE-driven transcription as a major determinant of the transcription stress landscape in cancer and highlight a key mechanism threatening the genomic stability of oncogenes.

Introduction

The most fundamental hallmark of cancer cells is their ability to sustain proliferation, which they acquire through genetic alterations during neoplastic transformation[1]. Simultaneously, dysregulation of the transcriptional program emerges to accommodate the cell's increased demand for cellular factors essential for proliferation, growth, and survival[2,3]. Consequently, genes encoding these factors, i.e., oncogenes, are hypertranscribed to fulfill the needs of cancer cells[4].

A well-established mechanism by which cancer cells upregulate oncogenes involves the formation of oncogenic super-enhancers (SEs)[5–12]. In normal cells, SEs drive the expression of cell-type specific genes that encode master transcription factors (MTFs), which shape cellular identity and the transcriptional program[2]. In cancer cells, SEs drive the expression of cancer-type specific oncogenes and oncogenic MTFs that dictate their transcriptional programs and support their proliferation[2,12]. SEs can become dysregulated in cancer through various mechanisms: 1) overexpression of enhancer transcription factors renders SEs much more effective at driving the expression of their associated genes; 2) translocations can relocate SEs near an inactive oncogene in the corresponding normal cell, and 3) focal amplifications of SEs can lead to the increased expression of target genes[12]. These mechanisms contribute to the activation of proto-oncogenes by SEs, which drive the transcription of these genes to an exceptionally high level termed "hypertranscription."

Transcription is increasingly viewed as a source of DNA damage[13–18]. The most damaging type of DNA lesions produced during transcription is DNA double-strand breaks (DSBs)[19]. DSBs resulting from transcription are primarily facilitated by the formation of pathogenic co-transcriptional R-loops and the aberrant activity of topoisomerase 1 (TOP1)[13,20]. R-loops tend to form more frequently at highly expressed genes, and their formation causes TOP1 to become trapped in its cleavage complex form (TOP1cc)[13].

Although transcription is a highly regulated process, it can occasionally become abortive and result in DSBs. These transcriptional aberrations occur more frequently as transcription levels increase, in a phenomenon known as "transcription stress"[21]. Characterizing the "hotspots" of transcriptional stress is crucial because transcription-related DSBs can lead to cell death, genomic instability, and mutations[22,23]. Moreover, since cancer cells upregulate genes vital for their proliferation and survival, a deeper understanding of transcriptional stress and transcription-

associated DSBs linked to these oncogenes can aid in exploiting their vulnerabilities to target

cancer cells more effectively. Nevertheless, while transcription stress has been increasingly

studied[24–26], its landscape and consequences, particularly in cancer, remain incompletely

characterized.

DSBs are impediments to transcription[27,28] and must, therefore, be efficiently repaired to sustain

continuous high transcriptional levels. However, it is not understood how regions of transcription

stress are equipped to handle these deleterious DSBs.

In this study, we characterize the landscape of transcription stress in cancer cells and reveal

how these regions are structured to promote the rapid and efficient repair of transcriptional DSBs

that sustain hypertranscription.

Results

Super-enhancer-regulated oncogenes are major transcription stress sites.

Pathogenic R-loops preferentially form at highly expressed genes[29], facilitating aberrant TOP1

activity and TOP1cc trapping[13]. The high TOP1 occupancy, TOP1cc trapping, and co-

transcriptional R-loops all contribute to the formation of transcription-associated DSBs [20,30].

Therefore, we reasoned that these features, along with genome-wide maps of endogenous

DSBs (hereafter referred to as "Breakome"), could serve as markers of transcriptional stress. To

determine transcription stress sites in the genome, we analyzed sBLISS and ChIP-seq data for

the breast cancer cell line MCF7, and we applied a Hidden Markov Model (HMM)-based

approach to identify regions in the genome with co-occurrence across the four datasets (sBLISS,

DRIP-seq, TOP1cc, and TOP1 ChIP-seq). As expected, the majority of these regions were genic

(Figure S1A), while the rest were enriched mainly at intergenic super-enhancer (SEs) (Figure

S1B and C). To investigate which genes serve as transcription stress sites, we scored each

gene based on TOP1, TOP1cc, R-loops, and DSBs scores (see Methods). This scoring yielded

a shortlist of genes, many of which have well-established oncogenic functions (Figure 1A). Upon

closer analysis of this list, we noted that many of the genes were regulated by SEs in MCF7

cells, including *CCND1*, *NEAT1*, *miR21*, and *MYC* (SE-regulated genes were downloaded from

SEdb 2.0[31]) (Figure 1A). To statistically validate this observation, we systematically assessed

the representation of multiple gene sets among transcription stress sites by performing gene set

enrichment analysis (Figure 1B; see Methods). SE-regulated genes were the most significantly

enriched category, with SE-regulated oncogenes exhibiting the highest enrichment (~122-fold

over expectation under a random model, $P = 1.85 \times 10^{-15}$; hypergeometric test). Similar results

were obtained when quantifying the overlap between each gene set and the co-marked regions

identified by HMM (Figure S1D). Notably, the enrichment of SE-regulated genes was

substantially higher than that of typical-enhancer (TE)-regulated genes or highly expressed

genes lacking enhancer regulation (Figure 1B), despite having similar expression levels (Figure

S1E). To control for factors that might confound this analysis, we performed multivariate logistic

regression using SE-status, expression level, GC content, and gene length as predictors for

transcription stress at genes. The results of this model demonstrate that SE-status was the most

statistically significant independent predictor of transcription stress (Figure S1F), indicating

specificity to genes regulated by SEs.

DSBs at these genes coincide with TOP1cc and R-loops (Figure 1C). Analysis of sBLISS data

for MCF7 cells depleted for TOP1 and R-loops showed that most of these genes have

significantly fewer breaks upon TOP1 and R-loops depletion (Figure S2A). Additionally, DSB are

enriched at SE-regulated genes in both cycling and G1-synchronized cells (Figure S2B-D),

validating their transcriptional origin and independence from replication.

The relationship between transcription and DSBs was previously established[13–15]. However, our

results suggest that this association may be more pronounced in SE-regulated genes. This

prompted us to further investigate the discrepancies in the relationship between transcription

and DSBs in SE-regulated versus non-SE-regulated genes. Interestingly, the strong correlation

between gene expression and DSBs is observed only in SE-regulated genes and not in others

(Figure S2E-G). Moreover, highly expressed genes enriched with DSBs showed significantly

greater enrichment in TOP1, TOP1cc, and R-loops than similarly expressed genes lacking DSBs

(Figure S3A-E). Additionally, SE-regulated genes were significantly more prevalent among

highly expressed genes enriched with DSBs (Figure S3F). These findings indicate that the

association between transcription and DSBs is primarily attributed to SE-regulated genes.

Since SE activity is significantly more potent in cancer cells than in normal cells, we next aimed

to test whether the enrichment of DSBs at SE-regulated genes is a phenomenon specific to

cancer. To investigate this, we mapped the Breakome in 10 different cell lines, which included

three normal cell lines, one pre-malignant cell line, and six cancer cell lines. We found that SE-

regulated genes are more enriched with DSBs in cancer cells than in their normal counterparts

(Figure 1D), demonstrating the specificity of transcriptional stress to active SE-regulated

oncogenes in cancer cells.

The super-enhancer landscape shapes transcription stress sites in cancer.

The landscape of SEs is cell-type specific, enhancing the transcription of different genes based

on cell identity[12]. If SEs cause transcription stress in their target genes, one would predict that

various cancer cell types exhibit distinct transcription stress sites reflective of their SE

landscapes. To test this, we mapped the breakome of multiple cancer cell lines, focusing on

each cell line's break-prone SE-regulated genes. Interestingly, SE-regulated genes were

enriched in DSBs, predominantly in the cell type where they are regulated by an SE (Figure 2A).

This pattern was also apparent when comparing cell types with very similar characteristics

(MCF7 and T47D, both of which are ER+ luminal A breast cancer cell types), yet they exhibit

differing SE landscapes, suggesting that transcription stress sites correspond to the SE

landscape in cancer.

If SE activity indeed causes the accumulation of DSBs at transcription stress sites, then inhibiting

its activity would abolish SE-mediated DSB accumulation, while conversely, activating it would

enhance DSB accumulation at target genes. To test this hypothesis, we employed two

strategies. First, we inhibited SE activity in MCF7 and HeLa cells through two approaches: using

the BRD4 degron dBET6 and the CDK9 inhibitor (CDK9i) NVP-2, both of which have been shown

to lead to SE collapse and repress SE-mediated transcription[32,33]. Examining the breakome after

a 4-hour incubation with these inhibitors, which effectively suppressed SE-regulated

transcriptional activity as measured by spike-in real-time quantitative PCR (RT qPCR) (Figure

S2G-H), revealed that SE inhibition significantly reduced DSBs at SE-regulated genes in both

cell lines (Figure 2B). Furthermore, SE inhibition using either dBET6 or CDK9i eliminated the

cell-type-specific effect of the SE landscape on the breakome (Figure 2C). Secondly, we utilized

the fact that MCF7 cells are estrogen receptor-positive and that many of their SEs are estrogen-

responsive, and analyzed the breakome of these cells before and after estrogen addition. We

specifically focused on genes regulated by estrogen receptor-occupied SE-regulated genes

(ERSE-genes), which were previously characterized[34]. Consistently, ERSE induction

significantly increased DSBs in target genes but not random genes (Figure 2 D-E), indicating

that SE activity promotes the accumulation of DSBs at SE-regulated genes. Overall, these data demonstrate that the landscape of SE activity in cancer cells shapes the landscape of transcription stress in a cell-type and cancer-type-specific manner.

Specificity of transcriptional stress at super-enhancer-regulated genes

While multiple studies have demonstrated the formation of DSBs as a consequence of transcription, independent of SE regulation [35–37], our results imply that this phenomenon is markedly amplified when transcription is driven by SEs (Figure 1B; Figure S1E–F). To further support this observation, we employed two complementary approaches. First, we examined predictors for the genomic locations of transcription stress sites. Using SE-regulated genes, typical enhancer (TE)-regulated genes, and generally highly expressed genes as predictors, multivariate LASSO regression revealed that SE regulation was the strongest predictor of transcription stress sites (odds ratio = 12.61; Figure 3A), even though the highly expressed gene group exhibited the highest transcriptional levels overall (Figure 3B). Notably, SE regulation was also the best predictor for genomic regions enriched in DSBs (Figure 3C).

Second, to experimentally assess whether transcriptional activation through SEs promotes greater transcription stress and DSB accumulation, we conducted a deeper analysis of the MCF7 breakome before and after estradiol treatment. Estrogen activates transcription via both promoter-proximal binding and estrogen-responsive SEs[38], enabling us to compare the impacts of these two mechanisms. We grouped genes by their level of transcriptional upregulation (based on GRO-seq fold change) and further subdivided each group into SE-regulated and non-SE-regulated categories. Consistently, SE-regulated genes accumulated significantly more DSBs than their non-SE counterparts. This effect was especially prominent in the highest induction group (fold change > 10; Figure 3D). A similar pattern was observed upon CDK9 inhibition, where SE-regulated genes exhibited a significantly greater reduction in DSBs compared to non-SE-regulated genes with matched transcriptional downregulation (measured by RNA-seq; Figure 3E).

Together, these results demonstrate that SE-mediated transcription is a major determinant of transcription stress sites and results in increased DSB accumulation compared to non-SE transcriptional activity, even at equivalent expression levels.[35]

γH2AX marks transcription stress sites

Since DSBs are transcription-blocking lesions[28], transcription stress sites must be rapidly and efficiently repaired to maintain high expression levels. γH2AX is a marker for DDR activation and is among the first responders to DSBs, as it is phosphorylated within a few minutes after DSB formation[39]. To further explore the role of DDR activation in the signaling of DSB repair at transcription stress sites, we conducted ChIP-seq for γH2AX on untreated MCF7 cells and analyzed high-quality available γH2AX ChIP-seq data for HeLa and IMR90 cell lines.

In both MCF7 and HeLa cells, γH2AX signal is predominantly localized to gene bodies, while in IMR90 cells, the distribution is more balanced between genic and intergenic regions (Figure S4A). This pattern is consistent with a greater burden of transcription-associated stress in cancer cells compared to non-transformed cells. Moreover, γH2AX is distributed across the gene bodies of top targets (Figure S4B-D). Ranking genes based on their γH2AX enrichment revealed that most exhibit relatively low γH2AX levels, whereas a subset displays high to very high γH2AX density (Figure 4A). This finding indicates a non-uniform distribution of γH2AX, with preferential enrichment at specific genes. Interestingly, all transcription stress sites are among the top 8% of γH2AX targets (Figure 4A-B). This enrichment in the top γH2AX genes was considerably less for the general high DSB genes (Figure 4C). Additionally, top γH2AX targets are enriched in SE-regulated genes across each respective cell line (P= $2.3e^{-41}$, $2.9e^{-39}$, $8.5e^{-8}$, for MCF7, HeLa, and IMR90 respectively; hypergeometric test) (Figure 4D-F).

The observed γH2AX enrichment across the three cell lines was consistent for several oncogenes, including *miR21, NEAT1, MALAT1,* and *ACTG1* (Figure 4G). Interestingly, many prominent γH2AX targets were unique to specific cell lines, most with established activity in their respective cell types (Figure 4H). For example, two top γH2AX targets in MCF7, *TFF1* and *GATA3*, are well-known oncogenic transcription factors primarily active in breast cancer and were exclusively enriched in γH2AX in MCF7 cells. *KRT7* and *CCAT1*, which are overexpressed and linked to poor prognosis in cervical cancer, are enriched in γH2AX only in HeLa cells. Consequently, we questioned whether these variations in γH2AX enrichment stem from cell-type-specific SEs. To investigate this, we compared γH2AX enrichment of top targets that are regulated by a cell-type specific SE in each respective cell line. Indeed, SE-regulated γH2AX

targets are predominantly enriched in the cell line where they are regulated by an SE (Figure 4

I), underscoring that γH2AX marks cell type-specific transcriptional stress sites.

γH2AX levels vary significantly across genes enriched with high endogenous DSBs.

Next, we examined the correlation between DSBs and γH2AX in genes. Interestingly, analyzing

the correlation between each gene's breaks and its γH2AX density revealed significant variability

of γH2AX among genes with high DSBs (Figure 4 J-K and Figure S5A-E). Some genes enriched

with breaks are marked by γH2AX, while others are not in both MCF7 and HeLa cell lines (Figure

S5F-G). This discrepancy in γH2AX enrichment at various DSB-enriched genes might suggest

that regions enriched with endogenous DSBs in the genome are signaled with varying efficiency

and potentially repaired unevenly, depending on their context.

γH2AX preferentially marks endogenous DSBs of highly active chromatin

Although genes exhibit a moderate correlation between DSB density and γH2AX density

(Spearman's correlation = 0.57), non-linear regression analysis using Generalized Additive

Models (GAM) indicates that the transcription stress score explains a greater proportion of the

variance in γH2AX levels (R^2 = 0.73) than any individual component (Figure 5A and Figure S6).

To further investigate the underlying factors contributing to variability in γH2AX at high DSBs

genes, we defined two groups of genes with similar break density but different γH2AX levels

(Figure 4J): DSBs^{high} γH2AX^{high} genes and DSBs^{high} γH2AX^{low} genes. DSBs^{high} γH2AX^{high} genes

are more expressed and enriched with transcription stress markers (TOP1, TOP1cc, and R-

loops) (Figure S7A-E). Furthermore, 64 of the 65 transcriptional stress sites were among

DSBs^{high} γH2AX^{high} genes (Figure 4J and Figure 5B). This group also included all SE-regulated

transcription stress sites and is significantly enriched with active SE-regulated genes compared

to DSBs^{high} γH2AX^{low} genes, which lack active SE-regulated genes and transcriptional stress

hotspots (Figure 5C and Figure S7F).

Chromatin context was previously shown to impact the phosphorylation of H2AX, specifically,

heterochromatin was shown to be refractory to H2AX phosphorylation[40]. Therefore, we

wondered whether the observed DSBs^{high} γH2AX^{low} genes are close to, or within,

heterochromatic regions. To gain further insights into whether γH2AX preferentially marks

endogenous DSBs based on the chromatin context, we defined two groups of individual DSB
sites across the genome according to their γH2AX signal (Figure 5D and Figure S5G) and
mapped them to the different chromatin states of MCF7 and HeLa. γH2AX^{high} DSBs were biased
toward active chromatin states, while γH2AX^{low} DSBs were biased toward silent regions (Figure
5E and Figure S5H). These results made us question whether the differential γH2AX marking at
DSB-enriched genes was due to their chromatin context. Indeed, the DSBs^{high} γH2AX^{low} group,
but not the DSBs^{high} γH2AX^{high} group, is enriched with DSB-enriched genes that overlap with
heterochromatin and repressed chromatin states (Figure 5F and Figure S5I). Additionally, an
analysis of the chromatin state contexts of DSBs^{high} γH2AX^{high} and DSBs^{high} γH2AX^{low} in both
MCF7 and HeLa showed that DSBs^{high} γH2AX^{high} genes primarily overlap with chromatin regions
associated with strong transcriptional activity. Conversely, DSBs^{high} γH2AX^{low} has a higher
overlap with heterochromatin, repressed, and weak transcription chromatin states (Figure 5G
and Figure S5J).
Our results illustrate that DSBs alone do not fully account for the variation in γH2AX across genes
(Figure 5A). Conducting GAM on genes categorized by their chromatin context revealed that
DSBs primarily account for variation in γH2AX in genes that overlap with strong enhancer regions
and least in genes that overlap with repressed or heterochromatin regions (Figure 5H). This
suggests that breaks and γH2AX are most closely associated with genes spanning strong
enhancer elements.
Collectively, these results indicate that γH2AX preferentially marks endogenous DSBs in active
chromatin compared to inactive chromatin. This differential marking may influence variations in
the repair efficiency of distinct DSBs based on their chromatin context and γH2AX signaling.
263
Genes marked with γH2AX exhibit high DSB turnover and repair efficiency
sBLISS, like other methods used to map DSBs, captures a snapshot of the DSB landscape at
the moment of fixation[41–45]. Therefore, a gene that shows high levels of DSBs could experience
frequent DSB generation and repair cycles (high DSB turnover) or infrequent DSB generation
and repair cycles (low DSB turnover). To estimate the rate of endogenous DSB repair across
the genome, we mapped DSBs in HeLa cells after disrupting the primary DSB repair pathway,
non-homologous end-joining (NHEJ)[45], by using siRNAs that target XRCC4. Regions that are

frequently repaired should accumulate excessive DSBs due to repair impairment. In contrast, regions with inherently low repair rates should show minimal DSB accumulation (Figure 6A). To objectively determine which genes accumulate DSBs following XRCC4 knockdown (Figure 6B), we employed the DESeq2 algorithm[46] on DSB count matrices, identifying 429 genes with significant DSB accumulation upon XRCC4 KD (Figure 6C). Stratifying these genes by their baseline DSB density revealed a trend in which groups with higher baseline DSB density showed progressively greater impact following XRCC4 KD (Figure 6D), supporting the notion that DSB-prone genes are repaired more effectively. Enrichment analysis indicated a strong enrichment of DSBs^{high} γH2AX^{high} genes (Figure 6E), along with the top DSB-enriched and active SE-regulated genes. In contrast, DSBs^{high} γH2AX^{low} genes and the bottom DSB-enriched and non-SE-regulated genes were not significantly enriched. Directly examining the effects of repair impairment demonstrated that NHEJ impairment leads to the most pronounced accumulation of DSBs at DSBs^{high} γH2AX^{high} and SE-regulated genes, compared to DSBs^{high} γH2AX^{low} and non-SE genes (Figure 6F). To investigate further how γH2AX patterns influence the repair and turnover of endogenous DSBs, we used the accumulation of DSBs following repair impairment as a proxy for repair efficiency (Figure 6G). Within DSBs^{high} γH2AX^{high} genes, repair efficiency markedly increases with higher DSB levels; however, within DSBs^{high} γH2AX^{low} genes, this relationship of repair efficiency is significantly weaker (Figure 6H). These findings suggest two distinct types of endogenous DSBs: γH2AX-marked DSBs, which exhibit high turnover, and poorly γH2AX-marked DSBs, which exhibit low turnover.

Efficient repair at transcriptional stress sites is crucial for SE-regulated oncogene expression.

High γH2AX marking and repair efficiency at transcription stress sites under physiological conditions suggests that these regions rely on DSB repair to sustain their elevated transcriptional levels. To test this, we treated MCF7 cells with an ATM inhibitor (ATMi), which impairs DSBs repair, and conducted sBLISS and spike-in RT qPCR. Indeed, ATM inhibition led to an accumulation of DSBs at genes, biased toward active SE-regulated genes (Figure S8A-B). Measuring mRNA levels of the top γH2AX marked SE-regulated genes indicated a significant decrease in their expression levels (Figure S8C). To determine whether this reduction in

expression was due to the inhibition of the ATM axis or the accumulation of DSBs, we knocked

down XRCC4 under the same conditions as figure 5 and performed spike-in RT qPCR.

Interestingly, the top γH2AX marked SE-regulated genes, which exhibited DSB accumulation,

significantly decreased in expression (Figure S8D). These results indicate that the failure to

repair DSBs, rather than merely inhibiting DDR signaling, impedes the expression of SE-

regulated oncogenes.

Efficiently repaired transcriptional DSBs cluster in the 3D genome.

Induced DSBs cluster in repair foci[47,48], especially when introduced within active genes[48]. We

hypothesized that endogenous transcription-associated DSBs might also cluster within the 3D

genome as a mechanism to enhance repair. To investigate this, we analyzed high-quality Hi-C

data generated from MCF7 cells[49].

To quantify the clustering of genes associated with break density, we introduced two Hi-C

metrics that measure how a group of genes with a common characteristic (e.g., break density)

is clustered (Figure 7A): intra-group interactions, which represent the sum of genomic interaction

scores between genes within the same group, and inter-group interactions, the sum of genomic

interaction scores between genes from different groups. A clustered group of genes is expected

to exhibit higher intra-group interactions and lower inter-group interactions compared to a non-

clustered group (Figure 7A). To investigate whether DSB-enriched genes exhibit clustering,

genomic bins were arranged according to the break densities of their constituent genes and

categorized into nine groups based on break density. Indeed, bins containing DSB-enriched

genes demonstrate higher intra-group interactions and lower inter-group interactions than bins

with low DSB genes (Figure 7B), indicating the clustering of high DSB genes.

The percentage of strong interactions among DSBs^{high} γH2AX^{high} genes is significantly higher

than that among DSBs^{high} γH2AX^{low} genes (Figure 7C and Figure S9A). Additionally,

transcription stress sites exhibit a substantially higher percentage of strong interactions

compared to random sites with similar chromosomal distribution (Figure 7 D and Figure S9B),

suggesting that efficiently repaired transcription stress sites are clustered together. To avoid the

confounding effects of SE regulation on 3D interactions, we analysed Hi-C data from the

MCF10A cell line, which was conducted in parallel to MCF7[49]. MCF10A is a pre-cancerous cell

line that shares many common SE-regulated genes with MCF7 cells. However, these common

genes are significantly more enriched in DSBs within MCF7 cells (Figure 7E). This enabled a
comparison of the same regions with the same SE-regulation status but differing DSB
enrichment. Indeed, bins containing common SE-regulated genes, as opposed to random
regions, showed significantly higher strong interchromosomal interactions in MCF7 cells than in
MCF10A cells (Figure 7F-G), indicating that transcription stress sites that arise during
carcinogenesis undergo increased clustering.
338
The repair of transcription stress sites is mutagenic.
Since transcription stress sites are more likely to be in close proximity (Figure 7), we suspected
that recurrent DSBs at these sites could serve as mutational hotspots. To explore this, we
downloaded and analysed mutational data from the COSMIC database[50], which contains
extensive mutational data on human cancers. While mutations within coding regions are subject
to selective pressures based on their impact on protein function and cellular fitness [51,52], non-
coding mutations—particularly those in intronic regions—are less constrained by selection and
can provide a more direct readout of a region's intrinsic susceptibility to mutation. To minimize
the confounding effects of selection, we focused on non-coding intronic mutations in breast
cancer and intersected them with our breakome data from the breast cancer cell line MCF7.
Grouping genes by their DSB or γH2AX density revealed that genes with a higher break or
γH2AX density exhibit greater intronic mutation density (Figure 8A-C). This association suggests
that the repair of transcription-associated DSBs is inherently error-prone, leading to the
accumulation of mutations over time.
353
To disentangle the effects of DSB density from repair and turnover, we modeled the relationship
between γH2AX density and break density using nonlinear least squares (NLS) regression. The
residuals from this model were then used to classify genes in each group into three subgroups:
Under-responsive genes (negative residuals), Moderately responsive genes (residuals near
zero), and Over-responsive genes (positive residuals). Notably, over-responsive genes showed
the most increase in mutation density with increased DSBs (Figure 8D-F). This suggests that
excessive γH2AX accumulation at DSB sites, which reflects recurrent DSB formation and
turnover, and heightened repair activity, is error-prone. Furthermore, DSBs^{high} γH2AX^{high} genes,
but not DSBs^{high} γH2AX^{low} genes, demonstrated intronic mutation enrichment, compared to

randomly selected genes (Figure 8G-I). Collectively, these results suggest that transcription
stress sites are prone to mutations, likely due to a combination of high DSB turnover, clustering,
and defective repair. These findings have important implications for understanding how
transcription-associated genome instability contributes to cancer evolution, as recurrent DSB
formation and error-prone repair may drive the accumulation of oncogenic mutations over time.
368
369

Discussion

Cellular processes–particularly those upregulated in cancer such as replication and
transcription–play a significant role in genomic instability[53]. Replication-dependent instability,
caused by replication stress[4,54], has been extensively studied, revealing its landscape[55,56],
causes, consequences[57], and potential for therapeutic targeting[58,59]. Transcription-replication-
dependent genomic instability[60], particularly transcription-replication conflicts (TRCs)[61], has also
been well characterized, with studies examining its genomic distribution, influencing factors, and
the impact of transcription-replication orientation[62]. Recently, attention has shifted toward
transcription-dependent, replication-independent genomic instability, namely, transcriptional
DSBs. While research has begun to uncover their causes and implications in disease[17,63],
several key aspects remain underexplored, including genomic hotspot patterns, repair
mechanisms, and their impact on cancer evolution. By utilizing multi-omic approaches, we
delineated the landscape of transcription stress and revealed that SEs shape the transcriptional
program and contribute to the formation of genomic instability hotspots in cancer.

SEs determine cell identity in normal cells[31]. In cancer cells, SEs enhance the transcription of
oncogenes to promote cell growth and proliferation [6,8,64,65]. Previous studies have established
that highly transcribed genes are prone to transcription-associated genome instability[13,15,20,30,66].
However, our data reveal that this relationship is significantly amplified in SE-regulated genes,
suggesting that transcriptional stress is not merely a function of high gene expression but is
modulated by SE activity. The enrichment of highly expressed SE-regulated genes among
transcription stress sites was over 30-fold higher than expected by chance, far exceeding that
of highly expressed TE-regulated genes or highly expressed genes lacking enhancer regulation.
This specificity suggests that SEs impose a unique transcriptional burden that predisposes their

target genes to DNA damage. The functional implications are significant, as many SE-regulated
genes encode critical oncogenes, including *CCND1*, *NEAT1*, *miR21*, and *MYC*, which play
essential roles in tumorigenesis.

The pronounced enrichment of DSBs at SE-regulated oncogenes was significantly greater in
cancer cell lines compared to their normal counterparts, indicating that SEs primarily drive
genomic instability in cancer cells. SEs have previously been recognized as hyper-mutated
regions that can alter the expression of target genes and contribute to carcinogenesis[6]. Our
findings enhance this understanding by demonstrating that SEs also dictate mutational
susceptibility at their target genes by inducing hyper-transcription.
Although sBLISS captures a snapshot of the DSB landscape at the time of fixation, we could
estimate the repair and turnover of DSBs across genes, revealing how transcription stress sites
particularly undergo high DSB turnover and repair efficiency. To accommodate this, these
regions experience constant activation of the DNA damage response (DDR) and H2AX
phosphorylation. Interestingly, not all DSB-enriched genes exhibit such efficient repair signaling,
suggesting two distinct types of DSB-bearing genes in the genome: genes that undergo frequent
breakage-repair cycles and are actively recognized by the repair machinery, and genes that,
despite being DSB-rich, experience infrequent breakage-repair cycles and remain persistent.
These persistent DSBs, found in regions with low expression, have a lower demand for repair.
Importantly, these observations pertain to physiological, endogenous DSBs under steady-state
conditions. In contrast, previous studies of induced DSBs, generated in large numbers by
exogenous agents, have reported different dynamics[40,67], likely reflecting a response to
unprogrammed and highly abundant damage. We speculate that these latter genes experience
breakage in a more programmed manner compared to genes with high turnover, which break as
a byproduct of transcription. The nature of these DSBs requires further investigation.
Additionally, we demonstrate that genes with efficient DSB repair signaling are more vulnerable
to mutations than genes with comparable DSB levels but less effective repair signaling,
emphasizing the importance of the repair process itself in mutagenesis at transcription stress
sites.
421

References

1.   Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: The next generation.
Preprint, https://doi.org/10.1016/j.cell.2011.02.013
https://doi.org/10.1016/j.cell.2011.02.013.
2.   Bradner, J.E., Hnisz, D., and Young, R.A. (2017). Transcriptional Addiction in Cancer.
Preprint at Cell Press, https://doi.org/10.1016/j.cell.2016.12.013
https://doi.org/10.1016/j.cell.2016.12.013.
3.   Lee, T.I., and Young, R.A. (2013). Transcriptional regulation and its misregulation in
disease. Preprint at Elsevier B.V., https://doi.org/10.1016/j.cell.2013.02.014
https://doi.org/10.1016/j.cell.2013.02.014.
4.   Bowry, A., Kelly, R.D.W., and Petermann, E. (2021). Hypertranscription and replication
stress in cancer. Preprint at Cell Press, https://doi.org/10.1016/j.trecan.2021.04.006
https://doi.org/10.1016/j.trecan.2021.04.006.
5.   Pott, S., and Lieb, J.D. (2015). What are super-enhancers? Preprint at Nature Publishing
Group, https://doi.org/10.1038/ng.3167 https://doi.org/10.1038/ng.3167.
6.   Bal, E., Kumar, R., Hadigol, M., Holmes, A.B., Hilton, L.K., Loh, J.W., Dreval, K., Wong,
J.C.H., Vlasevska, S., Corinaldesi, C., et al. (2022). Super-enhancer hypermutation
alters oncogene expression in B cell lymphoma. Nature *607*, 808–815.
https://doi.org/10.1038/s41586-022-04906-8.
7.   Wang, X., Cairns, M.J., and Yan, J. (2019). Super-enhancers in transcriptional regulation
and genome organization. Preprint at Oxford University Press,
https://doi.org/10.1093/nar/gkz1038 https://doi.org/10.1093/nar/gkz1038.
8.   Wang, M.T., Chen, Q.Y., Wang, S.J., Xie, H., Liu, J., Huang, R.X., Xiang, Y.F., Jiang,
Y.Y., Tian, D.S., and Bian, E.B. (2023). Super-enhancers complexes zoom in
transcription in cancer. Preprint at BioMed Central Ltd, https://doi.org/10.1186/s13046-
023-02763-5 https://doi.org/10.1186/s13046-023-02763-5.
9.   Bacabac, M., and Xu, W. (2023). Oncogenic super-enhancers in cancer: mechanisms
and therapeutic targets. Preprint at Springer, https://doi.org/10.1007/s10555-023-10103-
4 https://doi.org/10.1007/s10555-023-10103-4.
10.   Hnisz, D., Schuijers, J., Lin, C.Y., Weintraub, A.S., Abraham, B.J., Lee, T.I., Bradner,
J.E., and Young, R.A. (2015). Convergence of Developmental and Oncogenic Signaling
Pathways at Transcriptional Super-Enhancers. Mol Cell *58*, 362–370.
https://doi.org/10.1016/j.molcel.2015.02.014.
11.   Morton, A.R., Dogan-Artun, N., Faber, Z.J., MacLeod, G., Bartels, C.F., Piazza, M.S.,
Allan, K.C., Mack, S.C., Wang, X., Gimple, R.C., et al. (2019). Functional Enhancers
Shape Extrachromosomal Oncogene Amplifications. Cell *179*, 1330-1341.e13.
https://doi.org/10.1016/j.cell.2019.10.039.
12.   Hnisz, D., Abraham, B.J., Lee, T.I., Lau, A., Saint-André, V., Sigova, A.A., Hoke, H.A.,
and Young, R.A. (2013). XSuper-enhancers in the control of cell identity and disease.
Cell *155*, 934. https://doi.org/10.1016/j.cell.2013.09.053.
13.   Hidmi, O., Oster, S., Monin, J., and Aqeilan, R.I. (2024). TOP1 and R-loops facilitate
transcriptional DSBs at hypertranscribed cancer driver genes. iScience *27*, 109082.
https://doi.org/10.1016/j.isci.2024.109082.
14.   Lensing, S. V., Marsico, G., Hänsel-Hertsch, R., Lam, E.Y., Tannahill, D., and
Balasubramanian, S. (2016). DSBCapture: In situ capture and sequencing of DNA
breaks. Nat Methods *13*, 855–857. https://doi.org/10.1038/nmeth.3960.

15.   Oster, S., and Aqeilan, R.I. (2020). Programmed DNA Damage and Physiological DSBs:
469   Mapping, Biological Significance and Perturbations in Disease States. Preprint at NLM
470   (Medline), https://doi.org/10.3390/cells9081870 https://doi.org/10.3390/cells9081870.
471   16.   Girasol, M.J., Krasilnikova, M., Marques, C.A., Damasceno, J.D., Lapsley, C.,
472   Lemgruber, L., Burchmore, R., Beraldi, D., Carruthers, R., Briggs, E.M., et al. (2023).
473   RAD51-mediated R-loop formation acts to repair transcription-associated DNA breaks
474   driving antigenic variation in Trypanosoma brucei. Proc Natl Acad Sci U S A *120*.
475   https://doi.org/10.1073/pnas.2309306120.
476   17.   Stork, C.T., Bocek, M., Crossley, M.P., Sollier, J., Sanz, L.A., Dé Ric Ché Din, F.,
477   Swigut, T., and Cimprich, K.A. (2016). Co-transcriptional R-loops are the main cause of
478   estrogen-induced DNA damage. https://doi.org/10.7554/eLife.17548.001.
479   18.   Sollier, J., Stork, C.T., García-Rubio, M.L., Paulsen, R.D., Aguilera, A., and Cimprich,
480   K.A. (2014). Transcription-Coupled Nucleotide Excision Repair Factors Promote R-Loop-
481   Induced Genome Instability. Mol Cell *56*, 777–785.
482   https://doi.org/10.1016/j.molcel.2014.10.020.
483   19.   Halazonetis, T.D., Gorgoulis, V.G., and Bartek, J. An Oncogene-Induced DNA Damage
484   Model for Cancer Development.
485   20.   Cristini, A., Ricci, G., Britton, S., Salimbeni, S., Huang, S. yin N., Marinello, J., Calsou,
486   P., Pommier, Y., Favre, G., Capranico, G., et al. (2019). Dual Processing of R-Loops and
487   Topoisomerase I Induces Transcription-Dependent DNA Double-Strand Breaks. Cell
488   Rep *28*, 3167-3181.e6. https://doi.org/10.1016/j.celrep.2019.08.041.
489   21.   Milano, L., Gautam, A., and Caldecott, K.W. (2024). DNA damage and transcription
490   stress. Preprint at Cell Press, https://doi.org/10.1016/j.molcel.2023.11.014
491   https://doi.org/10.1016/j.molcel.2023.11.014.
492   22.   Min, J., Zhao, J., Zagelbaum, J., Lee, J., Takahashi, S., Cummings, P., Schooley, A.,
493   Dekker, J., Gottesman, M.E., Rabadan, R., et al. (2023). Mechanisms of insertions at a
494   DNA double-strand break. Mol Cell *83*, 2434-2448.e7.
495   https://doi.org/10.1016/j.molcel.2023.06.016.
496   23.   Reijns, M.A.M., Parry, D.A., Williams, T.C., Nadeu, F., Hindshaw, R.L., Rios Szwed,
497   D.O., Nicholson, M.D., Carroll, P., Boyle, S., Royo, R., et al. (2022). Signatures of TOP1
498   transcription-associated mutagenesis in cancer and germline. Nature *602*, 623–631.
499   https://doi.org/10.1038/s41586-022-04403-y.
500   24.   Nakazawa, Y., Hara, Y., Oka, Y., Komine, O., van den Heuvel, D., Guo, C., Daigaku, Y.,
501   Isono, M., He, Y., Shimada, M., et al. (2020). Ubiquitination of DNA Damage-Stalled
502   RNAPII Promotes Transcription-Coupled Repair. Cell *180*, 1228-1244.e24.
503   https://doi.org/10.1016/j.cell.2020.02.010.
504   25.   Gyenis, A., Chang, J., Demmers, J.J.P.G., Bruens, S.T., Barnhoorn, S., Brandt, R.M.C.,
505   Baar, M.P., Raseta, M., Derks, K.W.J., Hoeijmakers, J.H.J., et al. (2023). Genome-wide
506   RNA polymerase stalling shapes the transcriptome during aging. Nat Genet *55*, 268–
507   279. https://doi.org/10.1038/s41588-022-01279-6.
508   26.   Saponaro, M., Kantidakis, T., Mitter, R., Kelly, G.P., Heron, M., Williams, H., Söding, J.,
509   Stewart, A., and Svejstrup, J.Q. (2014). RECQL5 controls transcript elongation and
510   suppresses genome instability associated with transcription stress. Cell *157*, 1037–1049.
511   https://doi.org/10.1016/j.cell.2014.03.048.
512   27.   Min, S., Ji, J.H., Heo, Y., and Cho, H. (2022). Transcriptional regulation and chromatin
513   dynamics at DNA double-strand breaks. Preprint at Springer Nature,

https://doi.org/10.1038/s12276-022-00862-5 https://doi.org/10.1038/s12276-022-00862-
5.

28.   Machour, F.E., and Ayoub, N. (2020). Transcriptional Regulation at DSBs: Mechanisms
and Consequences. Preprint at Elsevier Ltd, https://doi.org/10.1016/j.tig.2020.01.001
https://doi.org/10.1016/j.tig.2020.01.001.

29.   Crossley, M.P., Bocek, M., and Cimprich, K.A. (2019). R-Loops as Cellular Regulators
and Genomic Threats. Preprint at Cell Press,
https://doi.org/10.1016/j.molcel.2019.01.024
https://doi.org/10.1016/j.molcel.2019.01.024.

30.   Cristini, A., Park, J.H., Capranico, G., Legube, G., Favre, G., and Sordet, O. (2016).
DNA-PK triggers histone ubiquitination and signaling in response to DNA double-strand
breaks produced during the repair of transcription-blocking topoisomerase I lesions.
Nucleic Acids Res *44*, 1161–1178. https://doi.org/10.1093/nar/gkv1196.

31.   Wang, Y., Song, C., Zhao, J., Zhang, Y., Zhao, X., Feng, C., Zhang, G., Zhu, J., Wang,
F., Qian, F., et al. (2023). SEdb 2.0: a comprehensive super-enhancer database of
human and mouse. Nucleic Acids Res *51*, D280–D290.
https://doi.org/10.1093/nar/gkac968.

32.   Shu, S., Wu, H.J., Ge, J.Y., Zeid, R., Harris, I.S., Jovanović, B., Murphy, K., Wang, B.,
Qiu, X., Endress, J.E., et al. (2020). Synthetic Lethal and Resistance Interactions with
BET Bromodomain Inhibitors in Triple-Negative Breast Cancer. Mol Cell *78*, 1096-
1113.e8. https://doi.org/10.1016/j.molcel.2020.04.027.

33.   Winter, G.E., Mayer, A., Buckley, D.L., Erb, M.A., Roderick, J.E., Vittori, S., Reyes, J.M.,
di Iulio, J., Souza, A., Ott, C.J., et al. (2017). BET Bromodomain Proteins Function as
Master Transcription Elongation Factors Independent of CDK9 Recruitment. Mol Cell *67*,
5-18.e19. https://doi.org/10.1016/j.molcel.2017.06.004.

34.   Zheng, Z.Z., Xia, L., Hu, G.S., Liu, J.Y., Hu, Y.H., Chen, Y.J., Peng, J.Y., Zhang, W.J.,
and Liu, W. (2022). Super-enhancer-controlled positive feedback loop BRD4/ERα-RET-
ERα promotes ERα-positive breast cancer. Nucleic Acids Res *50*, 10230–10248.
https://doi.org/10.1093/nar/gkac778.

35.   Geraud, M., Cristini, A., Salimbeni, S., Bery, N., Jouffret, V., Russo, M., Ajello, A.C.,
Fernandez Martinez, L., Marinello, J., Cordelier, P., et al. (2024). TDP1 mutation causing
SCAN1 neurodegenerative syndrome hampers the repair of transcriptional DNA double-
strand breaks. Cell Rep *43*. https://doi.org/10.1016/j.celrep.2024.114214.

36.   Bunch, H., Lawney, B.P., Lin, Y.F., Asaithamby, A., Murshid, A., Wang, Y.E., Chen,
B.P.C., and Calderwood, S.K. (2015). Transcriptional elongation requires DNA break-
induced signalling. Nat Commun *6*. https://doi.org/10.1038/ncomms10191.

37.   Madabhushi, R., Gao, F., Pfenning, A.R., Pan, L., Yamakawa, S., Seo, J., Rueda, R.,
Phan, T.X., Yamakawa, H., Pao, P.C., et al. (2015). Activity-Induced DNA Breaks
Govern the Expression of Neuronal Early-Response Genes. Cell *161*, 1592–1605.
https://doi.org/10.1016/j.cell.2015.05.032.

38.   Carroll, J.S., Meyer, C.A., Song, J., Li, W., Geistlinger, T.R., Eeckhoute, J., Brodsky,
A.S., Keeton, E.K., Fertuck, K.C., Hall, G.F., et al. (2006). Genome-wide analysis of
estrogen receptor binding sites. Nat Genet *38*, 1289–1297.
https://doi.org/10.1038/ng1901.

39.   γH2AX and cancer.

40. Kim, J.A., Kruhlak, M., Dotiwala, F., Nussenzweig, A., and Haber, J.E. (2007).
560   Heterochromatin is refractory to γ-H2AX modification in yeast and mammals. Journal of
561   Cell Biology *178*, 209–218. https://doi.org/10.1083/jcb.200612031.
562   41. Bouwman, B.A.M., Agostini, F., Garnerone, S., Petrosino, G., Gothe, H.J., Sayols, S.,
563   Moor, A.E., Itzkovitz, S., Bienko, M., Roukos, V., et al. (2020). Genome-wide detection of
564   DNA double-strand breaks by in-suspension BLISS. Nat Protoc *15*, 3894–3941.
565   https://doi.org/10.1038/s41596-020-0397-2.
566   42. Yan, W.X., Mirzazadeh, R., Garnerone, S., Scott, D., Schneider, M.W., Kallas, T.,
567   Custodio, J., Wernersson, E., Li, Y., Gao, L., et al. (2017). BLISS is a versatile and
568   quantitative method for genome-wide profiling of DNA double-strand breaks. Nat
569   Commun *8*. https://doi.org/10.1038/ncomms15058.
570   43. Canela, A., Sridharan, S., Sciascia, N., Tubbs, A., Meltzer, P., Sleckman, B.P., and
571   Nussenzweig, A. (2016). DNA Breaks and End Resection Measured Genome-wide by
572   End Sequencing. Mol Cell *63*, 898–911. https://doi.org/10.1016/j.molcel.2016.06.034.
573   44. Dobbs, F.M., van Eijk, P., Fellows, M.D., Loiacono, L., Nitsch, R., and Reed, S.H.
574   (2022). Precision digital mapping of endogenous and induced genomic DNA breaks by
575   INDUCE-seq. Nat Commun *13*. https://doi.org/10.1038/s41467-022-31702-9.
576   45. Wei, P.C., Chang, A.N., Kao, J., Du, Z., Meyers, R.M., Alt, F.W., and Schwer, B. (2016).
577   Long Neural Genes Harbor Recurrent DNA Break Clusters in Neural Stem/Progenitor
578   Cells. Cell *164*, 644–655. https://doi.org/10.1016/j.cell.2015.12.039.
579   46. Anders, S., and Huber, W. (2010). Differential expression analysis for sequence count
580   data. Genome Biol *11*. https://doi.org/10.1186/gb-2010-11-10-r106.
581   47. Arnould, C., Rocher, V., Saur, F., Bader, A.S., Muzzopappa, F., Collins, S., Lesage, E.,
582   Le Bozec, B., Puget, N., Clouaire, T., et al. (2023). Chromatin compartmentalization
583   regulates the response to DNA damage. Nature *623*, 183–192.
584   https://doi.org/10.1038/s41586-023-06635-y.
585   48. Aymard, F., Aguirrebengoa, M., Guillou, E., Javierre, B.M., Bugler, B., Arnould, C.,
586   Rocher, V., Iacovoni, J.S., Biernacka, A., Skrzypczak, M., et al. (2017). Genome-wide
587   mapping of long-range contacts unveils clustering of DNA double-strand breaks at
588   damaged active genes. Nat Struct Mol Biol *24*, 353–361.
589   https://doi.org/10.1038/nsmb.3387.
590   49. Barutcu, A.R., Lajoie, B.R., McCord, R.P., Tye, C.E., Hong, D., Messier, T.L., Browne,
591   G., van Wijnen, A.J., Lian, J.B., Stein, J.L., et al. (2015). Chromatin interaction analysis
592   reveals changes in small chromosome and telomere clustering between epithelial and
593   breast cancer cells. Genome Biol *16*. https://doi.org/10.1186/s13059-015-0768-0.
594   50. Sondka, Z., Dhir, N.B., Carvalho-Silva, D., Jupe, S., Madhumita, McLaren, K., Starkey,
595   M., Ward, S., Wilding, J., Ahmed, M., et al. (2024). COSMIC: a curated database of
596   somatic variants and clinical data for cancer. Nucleic Acids Res *52*, D1210–D1217.
597   https://doi.org/10.1093/nar/gkad986.
598   51. Burrell, R.A., McGranahan, N., Bartek, J., and Swanton, C. (2013). The causes and
599   consequences of genetic heterogeneity in cancer evolution. Preprint,
600   https://doi.org/10.1038/nature12625 https://doi.org/10.1038/nature12625.
601   52. Yates, L.R., and Campbell, P.J. (2012). Evolution of the cancer genome. Preprint,
602   https://doi.org/10.1038/nrg3317 https://doi.org/10.1038/nrg3317.

53.   Negrini, S., Gorgoulis, V.G., and Halazonetis, T.D. (2010). Genomic instability an
evolving hallmark of cancer. Preprint, https://doi.org/10.1038/nrm2858
https://doi.org/10.1038/nrm2858.
54.   Hills, S.A., and Diffley, J.F.X. (2014). DNA replication and oncogene-induced replicative
stress. Preprint at Cell Press, https://doi.org/10.1016/j.cub.2014.04.012
https://doi.org/10.1016/j.cub.2014.04.012.
55.   Kim, J., Sturgill, D., Sebastian, R., Khurana, S., Tran, A.D., Edwards, G.B., Kruswick, A.,
Burkett, S., Hosogane, E.K., Hannon, W.W., et al. (2018). Replication Stress Shapes a
Protective Chromatin Environment across Fragile Genomic Regions. Mol Cell *69*, 36-
47.e7. https://doi.org/10.1016/j.molcel.2017.11.021.
56.   Gaillard, H., García-Muse, T., and Aguilera, A. (2015). Replication stress and cancer.
Preprint at Nature Publishing Group, https://doi.org/10.1038/nrc3916
https://doi.org/10.1038/nrc3916.
57.   Zeman, M.K., and Cimprich, K.A. (2014). R E V I E W Causes and consequences of
replication stress.
58.   Machour, F.E., R. Abu-Zhayia, E., Kamar, J., Barisaac, A.S., Simon, I., and Ayoub, N.
(2024). Harnessing DNA replication stress to target RBM10 deficiency in lung
adenocarcinoma. Nat Commun *15*. https://doi.org/10.1038/s41467-024-50882-0.
59.   Cybulla, E., and Vindigni, A. (2023). Leveraging the replication stress response to
optimize cancer therapy. Preprint at Nature Research, https://doi.org/10.1038/s41568-
022-00518-6 https://doi.org/10.1038/s41568-022-00518-6.
60.   Corazzi, L., Ionasz, V.S., Andrejev, S., Wang, L.C., Vouzas, A., Giaisi, M., Di Muzio, G.,
Ding, B., Marx, A.J.M., Henkenjohann, J., et al. (2024). Linear interaction between
replication and transcription shapes DNA break dynamics at recurrent DNA break
Clusters. Nat Commun *15*. https://doi.org/10.1038/s41467-024-47934-w.
61.   Hamperl, S., and Cimprich, K.A. (2016). Conflict Resolution in the Genome: How
Transcription and Replication Make It Work. Preprint at Cell Press,
https://doi.org/10.1016/j.cell.2016.09.053 https://doi.org/10.1016/j.cell.2016.09.053.
62.   Hamperl, S., Bocek, M.J., Saldivar, J.C., Swigut, T., and Cimprich, K.A. (2017).
Transcription-Replication Conflict Orientation Modulates R-Loop Levels and Activates
Distinct DNA Damage Responses. Cell *170*, 774-786.e19.
https://doi.org/10.1016/j.cell.2017.07.043.
63.   Geraud, M., Cristini, A., Salimbeni, S., Bery, N., Jouffret, V., Russo, M., Ajello, A.C.,
Fernandez Martinez, L., Marinello, J., Cordelier, P., et al. (2024). TDP1 mutation causing
SCAN1 neurodegenerative syndrome hampers the repair of transcriptional DNA double-
strand breaks. Cell Rep *43*. https://doi.org/10.1016/j.celrep.2024.114214.
64.   Zhou, R.W., and Parsons, R.E. (2023). Etiology of super-enhancer reprogramming and
activation in cancer. Preprint at BioMed Central Ltd, https://doi.org/10.1186/s13072-023-
00502-w https://doi.org/10.1186/s13072-023-00502-w.
65.   Huang, H., Hu, J., Maryam, A., Huang, Q., Zhang, Y., Ramakrishnan, S., Li, J., Ma, H.,
Ma, V.W.S., Cheuk, W., et al. (2021). Defining super-enhancer landscape in triple-
negative breast cancer by multiomic profiling. Nat Commun *12*.
https://doi.org/10.1038/s41467-021-22445-0.
66.   Mosler, T., Conte, F., Longo, G.M.C., Mikicic, I., Kreim, N., Möckel, M.M., Petrosino, G.,
Flach, J., Barau, J., Luke, B., et al. (2021). R-loop proximity proteomics identifies a role

of DDX41 in transcription-associated genomic instability. Nat Commun *12*.
https://doi.org/10.1038/s41467-021-27530-y.
67. Jakob, B., Splinter, J., Conrad, S., Voss, K.O., Zink, D., Durante, M., Löbrich, M., and
Taucher-Scholz, G. (2011). DNA double-strand breaks in heterochromatin elicit fast
repair protein recruitment, histone H2AX phosphorylation and relocation to euchromatin.
Nucleic Acids Res *39*, 6489–6499. https://doi.org/10.1093/nar/gkr230.
68. Suehnholz, S.P., Nissan, M.H., Zhang, H., Kundra, R., Nandakumar, S., Lu, C., Carrero,
S., Dhaneshwar, A., Fernandez, N., Xu, B.W., et al. (2024). Quantifying the Expanding
Landscape of Clinical Actionability for Patients with Cancer. Cancer Discov *14*, 49–65.
https://doi.org/10.1158/2159-8290.CD-23-0467.
69. Hidmi, O., Oster, S., Shatleh, D., Monin, J., and Aqeilan, R.I. (2024). Protocol for
mapping physiological DSBs using in-suspension break labeling in situ and sequencing.
STAR Protoc *5*. https://doi.org/10.1016/j.xpro.2024.103059.
70. Dunham, I., Kundaje, A., Aldred, S.F., Collins, P.J., Davis, C.A., Doyle, F., Epstein, C.B.,
Frietze, S., Harrow, J., Kaul, R., et al. (2012). An integrated encyclopedia of DNA
elements in the human genome. Nature *489*, 57–74.
https://doi.org/10.1038/nature11247.
71. Luo, Y., Hitz, B.C., Gabdank, I., Hilton, J.A., Kagda, M.S., Lam, B., Myers, Z., Sud, P.,
Jou, J., Lin, K., et al. (2020). New developments on the Encyclopedia of DNA Elements
(ENCODE) data portal. Nucleic Acids Res *48*, D882–D889.
https://doi.org/10.1093/nar/gkz1062.
72. Greer, Y.E., Hernandez, L., Fennell, E.M.J., Kundu, M., Voeller, D., Chari, R., Gilbert,
S.F., Gilbert, T.S.K., Ratnayake, S., Tang, B., et al. (2022). Mitochondrial Matrix
Protease ClpP Agonists Inhibit Cancer Stem Cell Function in Breast Cancer Cells by
Disrupting Mitochondrial Homeostasis. Cancer Research Communications *2*, 1144–
1161. https://doi.org/10.1158/2767-9764.CRC-22-0142.
674
675

Method details

Cell culture and experimental conditions

MCF7 (HTB-22) and T-47D (HTB-133) cells were grown in RPMI supplemented with 10% (vol/vol) fetal bovine serum FBS (GIBCO), glutamine, and penicillin/streptomycin. HEK-293 (CRL-1573), MDA-MB-468 (HTB-132), MDA-MB-231 (HTB-26), and HeLa cells (CCL-2) were grown in DMEM supplemented with 10% (vol/vol) FBS, glutamine, and penicillin/streptomycin. A-4098 cells (HTB-44) were grown in. EMEM supplemented with 10% FBS, glutamine, and penicillin/streptomycin. SH-SY5Y (CRL-2266) were grown in DMEM/F12 supplemented with 10% FBS, glutamine, and penicillin/streptomycin. HMLE cells were grown in Promocell mammary epithelial cell basal media (C-21010) with added supplements (c-93110) whereas MCF10A were grown on DMEM/F12 supplemented with 5% Horse serum, 20 ng/ml EGF, 0.5 mg/ml Hydrocortisone, 100 ng/ml Cholera toxin, 10 mg/ml Insulin and Pen/Strep. MCF-7 cells were synchronized in the G1 cell cycle stage by culturing in serum-free media for 72 hours followed by replacing the media with 5% FBS media for 5 hours. Cells were grown at 37°C under a humidified atmosphere with 5% CO_2. Cells were routinely authenticated by STR profiling, and tested for mycoplasma, and cell aliquots from early passages were used.

Cells were treated with DMSO (Sigma-Aldrich), dBET6 (Sigma-Aldrich, SML2638), or CDK9i (NVP-2) (MCE, Cat# Y-12214A) 100 nM for 4 hours. Cells were then treated with DMSO or ATMi (20 uM, KU-55933, Sigma) for 5 hours.

Transcription stress score

Our sBLISS, DRIP-seq, TOP1cc, and TOP1 ChIP-seq data used for transcription stress score calculation are already published[13] and are available at GEO: GSE241309. For each dataset, the signal density per gene was first converted to an empirical p-value based on its rank within the distribution of all genes. Higher signal values correspond to lower p-values. These p-values were then transformed into Z-scores using the standard normal inverse cumulative distribution function:

$$Z_k = \Phi^{-1}(1 - p_k)$$

where p_k is the empirical p-value for dataset k, and Φ^{-1} denotes the inverse of the standard normal cumulative distribution function.

706 The individual Z-scores from all four datasets were combined per gene using the Liptak

707 weighted Z-test (unweighted version):

708
$$TSS_{raw} = \frac{Z_1 + Z_2 + Z_3 + Z_4}{\sqrt{4}}$$

709 For better representation, the combined score was further min-max normalized and then

710 stretched by a power transformation to emphasize differences:

712
$$TSS_{final} = TSS_{scaled}^4$$

Genes not among the top 1000 in one or more parameters were excluded.

Identification of Transcription Stress-Associated Domains using Multivariate HMM

To identify genomic regions simultaneously enriched for markers of transcription stress, we

applied a multivariate hidden Markov model (HMM) using the *mhsmm* R package. The hg38

genome was divided into 1 kb non-overlapping bins, and mean signals for TOP1, TOP1cc, R-

loops, and DSBs were calculated per bin. Signal intensities were capped at the 0.5th and 99.5th

percentiles to minimize the influence of outliers. A two-state multivariate HMM was initialized

with uniform state priors and a transition matrix favouring self-transition (0.95 probability).

Emissions were modelled using multivariate Gaussian distributions with means offset by ±0.5

standard deviations and shared covariance. The model was trained by maximum likelihood

estimation, and Viterbi decoding was used to assign each bin to a state. The state with the

highest average signal across all datasets was designated as the "co-marked" state. To refine

the identified regions, we selected bins showing high signal (above the 95th percentile) in at

least three of the four datasets, merged adjacent bins separated by ≤1 kb, and filtered to retain

regions ≥3 kb. [13]

Gene sets compilation and enrichment analysis

Gene sets tested for enrichment in this manuscript were defined as follows: SE-regulated genes

for each cell line were downloaded from SEdb2.0[31]. A list of annotated oncogenes was

downloaded from OncoKB[68]. Highly expressed genes are defined as the top 1000 expressed

genes. TE-regulated genes are defined as genes that overlap enhancer regions and are not

annotated by an SE. Non-E genes are genes that do not overlap any enhancer element.

enrichment was calculated by dividing the number of observed overlapping genes by the

theoretical number of genes predicted under a random model. One-tailed (upper-tailed for values

> 1, and lower-tailed for values < 1) hypergeometric test using the phyper function in R to

determine the statistical significance of the observed enrichment.

In-suspension break labelling *in situ* and sequencing (sBLISS)

sBLISS was conducted as previously described[41,69]. In summary, 10^6 cells were fixed in 2%

paraformaldehyde in 10% FCS/PBS for 10 minutes at room temperature. The fixation was

quenched with 125 mM glycine for 5 minutes at room temperature, followed by another 5 minutes

on ice and two washes in ice-cold PBS. Cells were lysed for 60 minutes on ice, and their nuclei

were permeabilized for 60 minutes at 37°C. Next, nuclei were rinsed twice with CutSmart Buffer

containing 0.1% Triton X-100 (CS/TX100), and double-strand break (DSB) ends were blunted in

situ using NEB's Quick Blunting Kit for 60 minutes at room temperature. The blunted nuclei were

then washed twice with 1x CS/TX100 before in situ ligation of the sBLISS adapters to the DSB

ends. Adaptor ligation was carried out with T4 DNA Ligase for 20-24 hours at 16°C, with BSA

and ATP added. Following ligation, the nuclei underwent two washes with 1x CS/TX100, and

genomic DNA was extracted using Proteinase K at 55°C for 14-18 hours while shaking at 800

rpm. Proteinase K was then heat-inactivated for 10 minutes at 95°C, followed by extraction using

Phenol:Chloroform:Isoamyl Alcohol, Chloroform, and ethanol precipitation. The purified DNA

was sonicated in 100 μL of ultra-pure water using Covaris M220 for 60 seconds. Sonicated

samples were concentrated with AMPure XP beads (Beckman Coulter), and fragment sizes

were evaluated using a BioAnalyzer 2100 (Agilent Technologies), targeting a range of 300 bp to

800 bp with a peak around 400-600 bp. The sonicated DNA was then in vitro transcribed using

the MEGAscript T7 Kit for 14 hours at 37°C. After RNA purification and ligation of the 3'-Illumina

adaptors, the RNA underwent reverse transcription. The final library indexing and amplification

step was performed with NEBNext® Ultra™ II Q5® Master Mix.

sBLISS fastq files are initially de-multiplexed using sample barcodes. Quality control is

performed with trim galore to eliminate residual adapters, trim reads to a base quality of at least

20, and filter out short reads smaller than 20 bp. Initial and final sample qualities are assessed

with fastqc. Quality-processed fastq files are aligned to the GRCh38 assembly with hisat2, then

sorted and indexed using samtools. The resulting bam files are de-duplicated with umi-tools,

utilizing genomic coordinates and Unique Molecular Identifiers (UMI). Custom Python and R

scripts are employed to identify read start positions and convert bam files to bigwig format for

downstream analysis. An additional custom R script is used to discard a blacklist of positions,

primarily within centromeres.

RNA-seq and GRO-seq data processing

Bulk RNA-seq processed data for untreated MCF7, HeLa, and T-47D cell lines were obtained

from ENCODE[70,71], while data for MDA-MB-436 and MDA-MB-468 were downloaded from GEO

(GSE212143)[72]. For MCF7, the following datasets were used: ENCFF967AOT, ENCFF328CKZ,

ENCFF921PJP, ENCFF930TQB, ENCFF030IUO, ENCFF878CJF, ENCFF057ITQ,

ENCFF317XCR, ENCFF985KJE, and ENCFF456OYZ. For HeLa, we used ENCFF750TQI,

ENCFF622TLZ, ENCFF689BFU, ENCFF151QTZ, ENCFF262EJQ, ENCFF878CJF,

ENCFF964OAC, ENCFF346JDS, and ENCFF638MLA. For T-47D, we used ENCFF234TQB,

ENCFF412DJX, ENCFF475SHD, and ENCFF846PLI. TPM values were averaged across

replicates for each cell line to determine the expression rank of each gene.

RNA-seq data for MCF7 after CDK9i treatment was downloaded from GSE129012 (4h

timepoint). And GRO-seq data for MCF7 after E2 treatment was downloaded from GSE27463

(40 min timepoint).

RT-qPCR and Spike-in RT-qPCR

Total RNA was isolated utilizing TRI reagent (Biolab), following the manufacturer's guidelines for

the phenol/chloroform extraction technique. cDNA was synthesized from 1 µg of RNA using the

QScript cDNA synthesis kit (Quantabio). The SYBR Green PCR Master Mix (Applied

Biosystems) was employed for qRT-PCR. All assays were conducted in triplicate. Experiments

were performed to assess the levels of cDNA using primers targeting: Chicken RPL4 (FW,

GAGTGACTACAACCTGCCGA; REV, TTGGCGTATGGGTTCAGCTT), SIK1 (FW,

CTCCGGGTGGGTTTTTACGAC; REV, CTGCGTTTTGGTGACTCGATG), ID3 (FW,

GAGAGGCACTCAGCTTAGCC; REV, TCCTTTTGTCGTTGGAGATGAC), XBP1 (FW,

CCCTCCAGAACATCTCCCCAT; REV, ACATGACTGGGTCCAAGTTGT), CCND1 (FW,

GCTGCGAAGTGGAAACCATC; REV, CCTCCTTCTGCACACATTTGAA), HES1 (FW,

CCTGTCATCCCCGTCTACAC; REV, CACATGGAGTCCGCCGTAA), FOS (FW,

TCCCATCGGTCCACTAGGTTT; REV, AGGGCTGCACTGAGTTCTTTG), JUNB (FW,

ACGACTCATACACAGCTACGG; REV, GCTCGGTTTCAGGAGTTTGTAGT), TFF1 (FW,

CCCTCCCAGTGTGCAAATAAG; REV, GAACGGTGTCGTCGAAACAG), UBC (FW,

ATTTGGGTCGCGGTTCTTG; REV, TGCCTTGACATTCTCGATGGT), HPRT (FW,

TGACACTGGCAAAACAATGCA; REV, GGTCCTTTTCACCAGCAAGCT), MYC (FW,

ATGCCCCTCAACGTGAACTTC; REV, CGCAACATAGGATGGAGAGCA), XRCC4 (FW,

ATGTTGGTGAACTGAGAAAAGC; REV, GCAATGGTGTCCAAGCAATAAC), GREB1 (FW,

TGGTCCGTAATGCACAAGGG; REV, CTGCGTTTAGTGAGGGGTGA), GATA3 (FW,

GCCTCTGCTTCATGGATCCC; REV, CACACTCCCTGCCTTCTGTG), ACTB (FW,

TTTTGGCTATACCCTACTGGCA; REV, CTGCACAGTCGTCAGCATATC).

In RT-qPCR experiments, Ct values were normalized to HPRT levels. For spike-in RT-qPCR,

chicken RNA was added in proportion to the number of cells in each sample, and Ct values were

normalized to chicken RPL4.

γH2AX ChIP-seq

MCF7 cells (~10^6) were crosslinked with 1% formaldehyde (methanol free, Thermo Scientific

28906) for 10 min at room temperature and quenched with glycine, 125 mM final concentration.

Fixed cells were washed twice in PBS and incubated in sonication buffer (0.5M NaCl, 0.5%SDS

RIPA buffer containing PMSF, protease and phosphatase inhibitors) for 30min on ice. Cells were

sonicated using Covaris M220 for 10min to produce chromatin fragments of ~200-300 bp. The

sheared chromatin was centrifuged 10min at maximum speed. From the supernatant, 50ul were

saved as input DNA and the rest was diluted in RIPA buffer without NaCl and SDS. The

chromatin was immunoprecipitated by incubation with 5 µg of gamma-H2AX (phospho S139;

Abcam, AB2893) antibody pre mixed with 100ul Dynabeads Protein G (ThermoFisher, 10004D)

and incubated over-night at 4°C with rotation. Immunoprecipitates were washed twice with RIPA

buffer + 150mM NaCl, twice with RIPA buffer + 300 mM NaCl, and twice with TE buffer. The

chromatin was eluted from the beads with 200ul of direct elution buffer (10mM Tris pH 8, 0.3M

NaCl, 5mM EDTA, 0.5% SDS) and incubated overnight at 65°C to reverse the cross-linking.

Samples were treated with RNAse for 1h at 37 °C and with proteinase K at 55 °C for 2 h. DNA

was cleaned up by QIAquick PCR purification column (Qiagen), according to the manufacturer's

instructions, and eluted in 30 μl of elution buffer. The ChIPed and the Input DNA were used for

real-time PCR or to prepare libraries by Truseq (Ilumina) and sequenced in Nextseq (Illumina).

ChIPed and input fastq files are quality controlled with trim galore to remove residual adapters,

trim reads to base quality of at least 20 and filter out short reads with size smaller than 20 bp.

Initial and final sample qualities are evaluated with fastqc. The quality processed fastq files are

aligned to GRCh38 genome with hisat2 followed by sorting and indexing with samtools. The

utility bamCompare of deepTools is applied on the bam files to create a peak bigwig file. This

operation is set to ignore duplicates, discard a blacklist of regions, normalize by RPKM, and

compare by ratio in tiles of 50 bp. γH2AX ChIP-seq data for HeLa and IMR90 were downloaded

from GEO: GSM4564685 and GEO: GSM2067927, respectively.

Statistical modelling and residual-based classification

For multivariate logistic regression, genes were categorized based on whether they overlapped

transcription stress sites identified by Hidden Markov Modeling (HMM). Logistic regression was

performed using the glm function in R (family = binomial).

Multivariate Least Absolute Shrinkage and Selection Operator (LASSO) regression was

conducted using the glmnet package in R, with 10-fold cross-validation (cv.glmnet) to select the

optimal regularization parameter (λ). The model was trained to classify genes based on

transcription stress site overlap or DSB enrichment. Odds ratios were derived from the final

model coefficients to assess the relative predictive power of each feature.

Generalized additive model (GAM) was applied to model the relationship between each variable

and γH2AX across all genes in (Figure 5A), and for genes classified into their overlapping

chromatin states in (Figure 5H). The model was constructed using the gam function in

the mgcv package in R.

To dissect the effects of absolute DSBs from DSB turnover on mutation frequency, we modelled

the relationship between break density and transcription stress by nonlinear least squares (NLS)

regression using the following equation:

854
$$gH2AX \ density = a \times break \ density^b$$

The model was fitted using the nls function in R, with initial parameter estimates

for "a" and "b" set to 0.1. After fitting the model, residuals were extracted and used to classify

genes into three response categories based on predefined thresholds: genes with residuals > 0

were classified as 'Over-responsive,' genes with residuals < 0 as 'Under-responsive' and genes

between -0.3 and 0.3 as 'Moderately responsive'.

Classifying individual DSB sites based on γH2AX

Each break site was assigned a γH2AX coverage level (γH2AX rank). To ensure a comparable

distribution of break scores (i.e., the number of DSBs at the same site) between the two groups,

break sites were classified into four categories: those with a break score of 1 (the large majority),

those with a break score between 2 and 4, those with a break score between 5 and 10, and

those with a break score between 11 and 50. Within each of the four categories, 1% of the top

and bottom γH2AX-ranked breaks are selected. All categories are then re-combined so that the

top set of breaks and the bottom set of breaks exhibit a similar break score distribution. Mapping

to chromatin states is then computed and plotted for these sets.

Chromatin states

Chromatin states bed files were downloaded from ENCODE[70,71]. For MCF7, ENCFF631HCR file

was used. For HeLa, ENCFF555HXM file was used.

Transient transfection

Transient transfection of siXRCC4 (Dharmacon; SMARTpool siRNA, Catalog ID: L-004494-00-

0005) and siSc (Dharmacon; D-00181010) was carried out using Lipofectamine (ThermoFisher).

On the day of transfection, cells were seeded to achieve a confluency of 60-70%. The

transfection solution was prepared by mixing 1.5 ml of serum and antibiotic-free RPMI with 30

µL of Lipofectamine and incubating for 5 minutes before adding a mixture of 1.5 ml RPMI and

0.6 nmoles of siRNA, followed by a 20-minute incubation at room temperature. This mixture was

then added to cells in a 10 cm plate cultured in antibiotic and serum-free media. The media was

replaced with fully supplemented media after 5-6 hours, and the cells were incubated in the

incubator for 48 hours.

HiC data analysis

Normalized Hi-C contact matrices for MCF7 and MCF10A cells were obtained from GEO
(GSE66733).

Genomic bins were intersected with all genes, and each bin was assigned to a single gene. Bins
overlapping multiple genes were excluded from the analysis.

In inter- and intra-group metrics calculation, genes were stratified into nine groups based on their
break density, excluding genes with zero breaks. Intra-group and inter-group interaction scores
were computed as follows: Intra-group interactions: For each gene, interaction scores with all
other genes within the same group were summed and normalized by the total number of genes
in that group. Inter-group interactions: For each gene, interaction scores with all genes outside
its group were summed and normalized by the total number of genes in all other groups. Genes,
along with their interaction scores, were ordered based on break density. The data was
visualized using loess smoothing.

COSMIC mutations data analysis

The file Cosmic_NonCodingVariants_Tsv_v101_GRCh37.tar, containing all noncoding variants,
was downloaded from COSMIC. This file was first filtered for breast cancer samples and
separated based on the type of mutation into SNVs, deletions, and insertions mutation files.
These files were then converted into genomic ranges objects and filtered to contain only
mutations within introns. To calculate mutation density, mutations per gene were counted and
normalized by the sum of intron length.

Acknowledgments

We sincerely thank the members of the Aqeilan lab for their valuable discussions and insights.
We are deeply grateful to Dr. Sheera Adar and Dr. Yotam Drier for their thoughtful guidance and
support as members of the PhD committee of O.H. We also appreciate the invaluable assistance
of Dr. Abed Nasereddin and Dr. Idit Shiff from the Core Research Facility of the Hebrew
University-Hadassah Medical School. Additionally, we extend our sincere gratitude to Prof.
Yinon Ben Neriah for generously providing dBET6 and CDK9i. This study was supported by
grants from the Israel Science Foundation (ISF) [No. 1056/21]. We also acknowledge the
support of the Carole and Andrew Harper Diversity Scholarship Program to O.H. and D.S,

Author contributions

922   O.H. and R.I.A. conceived and designed the study. O.H. and J.M. performed computational

923   analysis. S.O.F performed sBLISS on HMLE, MCF10A, MCF7, T47D, MDA-MB-436 and MDA-

924   MB-468 cell lines. D.S. performed sBLISS on MCF7 cells synchronized in G1, SH-SY5Y cells,

925   and NSCs. O.H. performed all other experiments. O.H. and R.I.A wrote the manuscript.

928
929
Figure 1: Super-enhancer-regulated oncogenes are major transcription stress sites.

A, Transcription stress sites ranked by their scores based on four transcription stress
parameters: TOP1, R-loops, TOP1cc, and DSBs in MCF7 cells. **B,** The ratio of observed to
expected numbers of genes in transcription stress sites. P-values were calculated using
hypergeometric testing (upper-tailed for enrichment in the top score and lower-tailed for the
bottom score). **C,** Genome browser snapshots depict DSBs, TOP1cc, and R-loops at selected
transcription stress sites. **D,** A boxplot illustrates DSBs enrichment at the SE-regulated genes
over other genes in each cell line (downloaded from SEdb[2.0 31]). Enrichment was calculated by
dividing the break density of each SE-regulated gene by the median break density of non-SE-
regulated genes and the average values for replicates were plotted. In the boxplot, the center
line represents the median; the box limits indicate the upper and lower quartiles; the whiskers
denote 1.5X the interquartile range, while the dots represent outliers. P-values were computed
using the Mann-Whitney U test, ****p < 0.0001.

Figure 2: Super-enhancer landscape shapes transcription stress sites in cancer.

A, Heatmap showing the normalized break density row Z-score for each cell line's unique SE-regulated genes. For each cell line, SE-regulated genes were identified, common SE-regulated

genes from other cell lines were excluded, and low to moderately expressed genes as well as
genes with low break counts were filtered out. Each column corresponds to a biological replicate.
Columns were normalized by median scaling. **B**, Boxplot illustrating the normalized break
density of cells treated with DMSO (control), dBET6, or CDK9i (100nM for 4 hours each) for SE-
regulated genes in each cell line or for randomly selected genes. values are averaged for three
biological replicates. **C**, Heatmap depicting the normalized break density row Z-score for each
cell line's unique SE-regulated genes following treatment with DMSO (control), dBET6, or
CDK9i. Each column corresponds to a biological replicate. Each sample was normalized by
downsizing to 270000 random breaks. **D**, Boxplot demonstrating the normalized break density
of cells treated with vehicle (control) or estradiol (E2, 100nM for 1 hour) for significantly affected
($p < 0.05$) estrogen receptor-occupied SE (ERSE) target genes or for randomly selected genes.
E, Heatmap representing the normalized break density row Z-score of significantly affected
ERSE target genes. Boxplot features include: center line representing median; box limits
denoting upper and lower quartiles; whiskers extending to 1.5x the interquartile range; and dots
indicating outliers. P-values were computed using the Mann-Whitney U test, with n.s. indicating
$p > 0.05$, ** indicating $p < 0.01$, and **** indicating $p < 0.0001$.

Figure 3. Specificity of transcription stress at super-enhancer (SE)–regulated genes.
A, Multivariate LASSO regression identifying predictors of transcription stress site locations in
100kb genomic bins. Odds ratios are shown. **B,** Boxplot showing normalized expression (TPM)

of SE-regulated, highly expressed, and TE-regulated genes. Despite lower expression. **C,**
LASSO regression for prediction of DSB-rich genomic sites. the genome was binned in 10kb
bins, and the top 5% bind in break density were assigned as DSB-rich. **D,** Boxplot showing the
break density in estrogen-responsive genes upon E2 treatment, grouped by degree of
transcriptional upregulation (fold change, FC). Non-SE genes were randomly downsampled to
match SE gene numbers for comparable P-values. **E,** Boxplot showing the break density in
estrogen-responsive genes following CDK9 inhibition (CDK9i), grouped by degree of
transcriptional downregulation (log2 fold change). Non-SE genes were randomly downsampled
to match SE gene numbers for comparable P-values. Boxplot features include: center line
representing the median; box limits denoting upper and lower quartiles; whiskers extending to
1.5x the interquartile range; and dots indicating outliers. P-values in boxplots were calculated
using the Mann-Whitney U test in B, and Wilcoxon paired test in D and E. n.s. $p > 0.05$, *$p < 0.05$,
**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure 3: γH2AX marks transcription stress sites.

A, Ranked plot of genes in MCF7 cells based on γH2AX density. **B**, Boxplot showing γH2AX
density for transcription stress sites and randomly selected genes. **C**, The ratio of the observed
to the expected number of transcription stress sites or top 1000 break-prone genes in MCF7
among the top γH2AX enriched genes. **D-F**, The ratio of observed to expected numbers of SE-
regulated genes in the top and bottom 300 genes enriched in γH2AX for each cell line. P-values
were determined using the hypergeometric test: n.s. $p > 0.05$, *$p < 0.05$, ***$p < 0.001$, ****$p <$
0.0001. **G,** Genome browser snapshots of representative common γH2AX targets across MCF7,
HeLa, and IMR90 cell lines. **H**, Genome browser snapshots of representative unique γH2AX
targets in MCF7, HeLa, and IMR90 cell lines. **I**, Heatmap displaying the γH2AX density column
Z-score for each cell line's unique SE-regulated genes. For each cell line, SE-regulated genes
were identified, common SE-regulated genes with other cell lines, and genes with low γH2AX
counts were filtered out. **J**, Scatter plot showing the correlation between break density and
γH2AX density for each gene. **K,** Coefficient of variation (CV) of γH2AX density as a function of
mean DNA double-strand break (DSB) signal. Genes were binned into deciles based on their
DSB signal intensity (log-transformed), and for each bin, the mean γH2AX density, standard
deviation, and CV (standard deviation divided by mean) were calculated. Boxplot features
include: center line representing the median; box limits denoting upper and lower quartiles;
whiskers extending to 1.5x the interquartile range; and dots indicating outliers. P-values in
boxplots were calculated using the Mann-Whitney U test, ****$p < 0.0001$.

Figure 4: γH2AX preferentially marks endogenous DSBs of active chromatin.
A, Barplot showing R² values for the different variables tested regarding their effect on γH2AX
variance across all genes. **B**-**C**. The ratio of observed to expected number of transcription stress
sites (**B**) and active SE-regulated genes (**C**). **D**, γH2AX signal at the two defined sets of DSBs.
E. The percentage of DSBs enriched with γH2AX (γH2AX^{high} DSBs) or lacking γH2AX (γH2AX^{low}
DSBs) in each chromatin state. Chromatin states were ordered based on what is typically known
about their transcriptional activity. **F**, The ratio of observed to expected numbers of high DSB
genes overlapping heterochromatin and repressed regions in both gene groups in MCF7 cells.
G. The percentage of overlaps between each gene group and chromatin states in MCF7. For
each group, the number of overlaps was counted for each chromatin state and then divided by
the total number of overlaps. **H**. A barplot showing R² values for the effect of DSBs on γH2AX
variance for genes overlapping different chromatin states. P-values were determined using the
hypergeometric test: ****p < 0.0001.

1015
1016 **Figure 5: Genes marked with γH2AX show high DSB turnover and repair efficiency. A**.
1017 Schematic representation of the rationale behind the experiment to estimate the rate of
1018 endogenous DSB repair and turnover. **B**. Validation of XRCC4 knockdown via RT qPCR, each
1019 dot represents the mean of three technical replicate of each biological replicate. **C**. Volcano plot
1020 depicting the results of the differential expression analysis of DSBs following XRCC4 KD,

assessed using the DESeq2 algorithm. The experiment was performed on two biological replicates per condition. **D**. Boxplot illustrating the log2FC of break density upon XRCC4 KD for significantly affected genes stratified according to their baseline break density into ten groups. **E**. The ratio of observed to expected number of genes in our defined gene sets. P-values were calculated using Fischer exact test. **F**. Boxplot of break density fold change. P-values were calculated using the Mann-Whitney U test. **G**. Genome browser snapshots of DSBs^{high} γH2AX^{high} and DSBs^{high} γH2AX^{low} gene examples, showing DSBs, γH2AX, and repair efficiency. Repair efficiency was estimated by pooling replicates, tiling the genome into 100bp tiles, subtracting the break score of each tile in the siXRCC4 sample from siSc (control), and smoothing the segments through sliding window averaging. **H**. Line plot for repair efficiency (log2FC after XRCC4 KD) for DSBs^{high} γH2AX^{high} versus DSBs^{high} γH2AX^{low} genes ordered by their break density. Lines were generated using LOESS smoothing. Boxplot: center line indicates median; box limits represent upper and lower quartiles; whiskers extend to 1.5x interquartile range; dots indicate outliers. *p < 0.05, ****p < 0.0001.

1041
Figure 7: Efficiently repaired transcriptional DSBs cluster in the 3D genome.
A. Schematic illustrating inter- and intra-group interactions and their use to determine the
clustering of genes with a common characteristic (high DSB density in this instance). **B**. Line
plot depicting inter-group (dashed line) and intra-group (solid line) genomic bins arranged
according to the break density of their overlapping genes. Lines were created using LOESS
smoothing, with shaded areas representing the 95% confidence intervals. **C**. Percentage of
strong interactions within genomic bins containing DSBs^{high} γH2AX^{high} compared to bins with
DSBs^{high} γH2AX^{low} genes. Strong interactions are defined as a log2 interaction score > 2. The
percentage is calculated by dividing the number of strong interactions by the total number of
interactions. P-values were calculated by Chi-squared test. **D**. Percentage of strong interactions
calculated as in (**C**). **E**. Boxplot reflecting break density of common SE-regulated genes between
MCF7 and MCF10A cell lines. P-values were calculated by Mann-Whitney U test. **F**. Heatmap
showing Log2 interaction scores for genomic bins containing common SE-regulated genes from
both MCF7 and MCF10A cell lines. Arrows indicate distinct sites of variation in interactions. **G**.
Count of interchromosomal strong interactions shown in the heatmap from (**F**). Random bins
serve as a reference. Boxplot: center line indicates median; box limits represent upper and lower
quartiles; whiskers extend to 1.5x interquartile range; dots indicate outliers. P-values were

calculated using chi-square test in (**c-d**) and (**g**), and Wilcoxon paired test in (e). ***p < 0.001,
****p < 0.0001.
1061

1062

Figure 7: transcriptional DSBs with high turnover are especially susceptible to mutations.
A-**C**. Intronic deletion density (**A**), insertion density (**B**), and SNV density (**C**) for genes categorized by their break or γH2AX density (excluding zero break genes). Random genes are included for reference. **D**-**F**, Rate of intronic mutations across genes categorized by their break density and further divided into three groups based on their γH2AX response to DSBs. **G-I**. Boxplot of intronic deletion density (**G**), insertion density (**H**), and SNV density (**I**) for DSBs[high] γH2AX[high] vs. DSBs[high] γH2AX[low] vs. randomly selected genes.

1070

Figure S1. Characterization of transcription stress sites defined by HMM analysis. **A,** Distribution of transcription stress sites identified by HMM across genic and intergenic regions. Genic regions were defined as gene coordinates extended by ±10 kb flanking regions. **B,** Proportion of intergenic transcription stress sites identified by HMM overlapping key chromatin states and super-enhancers. **C,** Enrichment analysis showing observed over expected overlap of intergenic transcription stress sites with chromatin states and super-enhancers. **D,** Enrichment analysis of transcription stress sites within different gene categories. **E,** Boxplot displaying expression levels of noted genes. **F,** Logistic regression analysis testing the predictive power of SE status, gene expression, GC content, and gene length for the presence of transcription stress sites within genes.

1081
1082 **Figure S2**
1083 **A**, Heatmap illustrating break density of SE-regulated genes following RNase H overexpression
1084 and TOP1 KD. **B**, Boxplot showing break density of random genes or SE-regulated genes in
1085 MCF7 cells synchronized in G1 phase or cycling cells. **C–D,** Flow cytometry analysis of DNA
1086 content in MCF7 cells. (C) DNA content profile of asynchronously cycling MCF7 cells. (D) DNA

content profile of MCF7 cells synchronized in G1 phase, demonstrating enrichment of the population in G1. Cells were stained with propidium iodide (PI) and analyzed by flow cytometry. **E**-**G**, Scatter plot demonstrating the correlation between expression and break density for all genes (**E**), non-SE-regulated genes (**F**), and SE-regulated genes (**G**). Boxplot: center line indicates median; box limits represent upper and lower quartiles; whiskers extend to 1.5x interquartile range; dots indicate outliers. P-value was calculated using Mann Whitney U test. n.s. indicating p > 0.05, * indicating p < 0.05.

Figure S3

A-**E**, Boxplot showing expression levels (**A**), R-loop density (**B**), TOP1 density (**C**), break density
(**D**), and TOP1cc density (**E**) for high expression high breaks genes versus high expression low
breaks genes. **F**, observed to expected ratio of SE genes in high expression high breaks versus
high expression low breaks genes. **g-h**, boxplot illustrating mRNA levels of sample SE-regulated
genes in MCF7 (**G**) and HeLa (**H**) cells after dBET6 or CDK9i (100nM for 4 hours). Boxplot:
center line indicates median; box limits represent upper and lower quartiles; whiskers extend to
1.5x interquartile range; dots indicate outliers. P-value was calculated using Mann Whitney U
test. n.s. indicating $p > 0.05$, **** indicating $p < 0.0001$.

1112

Figure S4. Distribution and enrichment of γH2AX signal at genic regions across different
cell lines. A, Percentage of total γH2AX peak coverage occurring within genic and intergenic
regions in MCF7, HeLa, and IMR90 cells. Genic regions were defined as gene coordinates
extended by ±10 kb flanking regions. (**B**‑**D**) γH2AX signal over input across the gene body and
flanking regions (±10 kb) for the top enriched genes in (**B**) MCF7, (**C**) HeLa, and (**D**) IMR90
cells. TSS: transcription start site; TES: transcription end site.

1119
Figure S5
A, Scatter plot showing the correlation between break density and γH2AX density for each gene
in HeLa cells. **B-E**, boxplots showing DSBs (B and D) and γH2AX density (C and E) for the two
groups of genes defined in (a) and Figure 3F. **D-E,** Genome browser snapshots of representative

examples of DSBs[high] γH2AX[high] and DSBs[high] γH2AX[low] genes in MCF7 (D), and HeLa (E) cell lines. Boxplot: center line indicates median; box limits represent upper and lower quartiles; whiskers extend to 1.5x interquartile range; dots indicate outliers. P-values were calculated using Mann Whitney U test. n.s. indicating p > 0.05, **** indicating p < 0.0001.

Figure S6
Smooth terms from Generalized Additive Models (GAMs) showing the effect of transcription-related predictors on γH2AX density. The y-axis shows the partial effect of each predictor (centered smooth function, s(x)) on γH2AX levels after accounting for non-linear relationships. The red line represents the fitted smooth function, and the shaded gray band indicates the 95% confidence interval.

1141
Figure S7
A-D, Boxplot showing expression levels (**A**), R-loop density (**B**), TOP1 density (**C**), TOP1cc
density (**D**) for DSBs^{high} γH2AX^{high} and DSBs^{high} γH2AX^{low} genes in MCF7 cells. **E**, boxplot
showing expression levels for DSBs^{high} γH2AX^{high} and DSBs^{high} γH2AX^{low} genes in HeLa cells.
F, observed to expected ratio of active SE genes in DSBs^{high} γH2AX^{high} in HeLa cell line. **G**,
heatmap showing the enrichment of γH2AX signal at the two defined sets of DSBs. **H**, The
percentage of DSBs enriched with γH2AX (γH2AX^{high} DSBs) or lacking γH2AX (γH2AX^{low} DSBs)
in each chromatin state of HeLa cell line. Chromatin states were ordered based on what is
typically known about their transcriptional activity. **I**, The ratio of observed to expected numbers
of high DSB genes overlapping heterochromatin and repressed regions in both gene groups in
HeLa cells. **J**, The percentage of overlaps between each gene group and chromatin states in
HeLa. For each group, the number of overlaps was counted for each chromatin state and then
divided by the total number of overlaps.

1155
Figure S8: Efficient repair at transcription stress sites is required for SE-regulated
oncogene expression
A, Ratio of observed to expected active SE-regulated genes and non-SE-regulated genes in
genes with significantly increased vs decreased DSBs upon ATMi treatment. P-values were
calculated using chi-square test. **B**, break density of significantly affected SE-regulated genes.
C, heatmap showing the change of mRNA levels as measured by qPCR upon ATMi treatment
of top γH2AX-enriched SE-regulated genes in MCF7 cells. P-values were calculated by Wilcoxon
paired test. **D**, heatmap showing the change of mRNA levels as measured by qPCR upon
XRCC4 KD of top γH2AX-enriched SE-regulated genes in HeLa cells. n.s. indicating p > 0.05,
** indicating p < 0.01.

1166
1167 **Figure S9**
1168 **A**, Heatmap showing Log2 interaction scores for genomic bins containing DSBs[high] γH2AX[high]
1169 (top) and DSBs[high] γH2AX[low]. (bottom). **B**, Heatmap showing Log2 interaction scores for genomic
1170 bins containing transcription stress sites or randomly selected bind with similar chromosomal
1171 distribution.