Improving Cellular Senescence Detection with Weak Gene Markers in Single-Cell RNA-seq vi Iterative Imputation

Peng Xu[1,#,*], Hantao Zhang[1,#], Siyao Zhu[2], and Yimeng Kong[1,*]

[1] Center of Clinical Laboratory Medicine, Zhongda Hospital, School of Medicine, Advanced Institute for Life and Health, Southeast University, Nanjing, 210096, China

[2] T.H. Chan School of Public Health, Harvard University, Boston, MA 02115

[#] Authors contributed equally

[*] *Corresponding author and person to whom reprint requests should be addressed*:
Peng Xu, Ph.D. Email: pxu@seu.edu.cn
Yimeng Kong, Ph.D. Email: kongyimeng@seu.edu.cn

Abstract

Cellular senescence represents a stable cell cycle arrest state that plays critical roles in tissue aging and age-related pathologies. While single-cell RNA sequencing (scRNA-seq) has enabled comprehensive profiling of millions of cells in aged tissues, reliable identification of senescent cells remains challenging due to the weak and non-specific expression of marker genes. Although existing computational methods could be used to detect active cell states through gene set scoring approaches, their performance with senescence genes has not been rigorously assessed. In this study, we showed that senescence genes display weak, non-specific expression patterns across different tissues and are susceptible to dropout events in scRNA-seq. Current scoring approaches suffered intrinsic limitations in single-cell detection using weakly expressed markers. Simply expanding the gene set of weak markers failed to improve detection accuracy. To overcome these limitations, we developed ICE (Iterative-imputation-based Cell Enrichment), a computational framework that combines expression imputation with iterative marker refinement. ICE achieved a remarkable improvement in detection precision when analyzing pancreatic α cells with weakly expressed markers, by increasing accuracy from 56% to 98%. Applied to senescence markers (*CDKN1A/p21*, *CDKN2A/p16*, *ATF3*, and *MX1*), ICE successfully identified marker-specific cell populations, including stressed β cells in aging and type-I interferon-responsive microglia in Alzheimer's disease (AD). Together, our study offered a rigorous analytical framework for single cell detection using markers with weak expressions, which will facilitate in-depth exploration of senescence heterogeneity and temporal dynamics in human tissue and disease states.

Background

Aging is a complex multifactorial process, characterized by the progressive decline of physiological function across organ systems[1]. At the cellular level, cellular senescence—a state of irreversible cell cycle arrest—plays a central role in organismal aging[2]. This process is triggered by diverse stressors, including DNA damage, telomere attrition, oncogenic activation, and oxidative stress[3]. Key regulators of senescence include cell cycle inhibitors, DNA damage response mechanisms, and metabolic reprogramming. A hallmark of senescent cells is the senescence-associated secretory phenotype (SASP), which involves the secretion of inflammatory cytokines, chemokines, and proteases that disrupt tissue microenvironments and contribute to age-related dysfunction[4,5].

Although significant, studying senescence in human aging remains challenging due to the lack of universal biomarkers[6,7]. Conventional markers such as β-galactosidase activity, *CDKN1A/p21* and

CDKN2A/p16 are context-dependent and often overlap with non-senescent processes[8]. For example, *p21*

functions in both senescence and oligodendrocyte development, limiting its specificity as a definitive marker[9].

Recent advances have identified novel senescence-associated pathway and gene markers. For instance,

multiple studies reported the derepression of LINE-1 retrotransposons and senescence-associated endogenous

retroviruses (SA-ERVs), which activate the IFN-I signaling pathway[10-12]. The transcription factor ATF3 drives

SA-ERV expression, producing double-stranded RNAs that amplify IFN-I signaling and stabilize the

senescent state[13]. ATF3 also remodels chromatin accessibility to promote senescence in endothelial cells and

aging tissues, underscoring its pleiotropic role in aging[14,15]. In the aged mouse brain, microglial IFN-I

signaling exacerbates neuroinflammation and neuronal decline, while its ablation mitigates these effects[16].

These findings highlight other non-universal biomarkers, such as *ATF3* and IFN-I signaling, acting as

promising new markers of senescence in human tissues and diseases.

scRNA-seq has transformed aging research by enabling high-resolution analysis of cellular heterogeneity.

Studies in model organisms and humans have uncovered senescence-associated transcriptional changes in

adipose, muscle, and brain tissues[17-19]. However, detecting senescent cells remains difficult due to the weak

and non-specific expression of canonical markers (e.g., *p16*, *p21*, SASP factors) and inherent technical noise

in scRNA-seq. Gene Set Enrichment Analysis (GSEA), originally developed for bulk RNA-seq data, has been

adapted to detect aging-associated cells[20,21]. More recently, methods such as AUCell and UCell have been

developed specifically for scRNA-seq for active cell detection with predefined gene markers[22,23]. However,

systematic benchmarking of these methods for weakly expressed senescence markers is lacking.

In this study, we first explored expressions of senescence genes across different tissues and cell types,

illustrating their weak and non-specific expression patterns. We then showed that current gene set scoring

methods, both methods designed for bulk and for single-cell data, are intrinsically limited in analyzing weakly

expressed markers. To address the limitation, we introduced ICE, an iterative-imputation-based analytical

framework that enhances detection performance specifically for weak markers. Applying ICE to diverse

senescence markers, we identified distinct cell populations in pancreatic β cells and Alzheimer's disease (AD)

microglia. Our method provides a robust platform for investigating senescence heterogeneity and its systemic

impacts in aging and age-related diseases at single-cell level.

Results

Senescence genes exhibit weak expressions in human Tissues and cell types

To characterize the expression patterns of senescence-associated genes, we systematically examined

senescence marker genes across multiple human tissues and cell types. Using the CellAge database (version

3)[24], we identified 274 senescence-inducing genes, including 126 oncogene-induced senescence genes, 65

replicative senescence genes, and 83 stress-induced senescence genes. To assess whether these markers exhibit

tissue specificity, we compared their expression levels with tissue-specific genes (tissue-markers) in bulk

RNA-seq data from 38 GTEx tissues. We found that senescence markers displayed weak and non-specific

expression patterns compared to tissue-markers (**Fig. 1a**). The median expression of senescence genes ranged

between 8.39–8.98 log2(CPM), significantly lower than the median expression of tissue-markers (10.93

log2(CPM); Wilcoxon test, $p < 0.01$) (**Fig. 1b**). These results suggest that senescence markers are broadly

expressed at low levels across tissues, lacking strong tissue specificity.

We next performed single-cell resolution analysis of senescence gene expression using pancreatic

scRNA-seq data[25]. To account for technical limitations of scRNA-seq (e.g., dropout events), we calculated

detection ratios (1 - dropout rate) for senescence markers. All three senescence marker categories showed low

median detection ratios (oncogene-induced: 12.5%; replicative: 10.2%; stress-induced: 8.3%) compared to

canonical cell-type markers (71.2%) (**Fig. 1c**). This pattern was consistent across all pancreatic cell types,

indicating that senescence markers are weakly expressed and distinct from conventional cell-type markers.

Fig. 1 | Expression patterns of senescence markers in human tissues and cell types.

a, Heatmap of senescence marker expression across 38 GTEx (v8) tissues. Rows represent tissues; columns show genes grouped by senescence category sourced from CellAge database. Expression values are standardized (z-score), with colors ranging from low (blue) to high (red).

b, Boxplots of expression distributions for senescence markers (grouped by type) and tissue-specific markers. Boxes denote the interquartile ranges (IQR) (25th–75th percentiles), center lines indicate medians, and whiskers extend to 1.5×IQR. Individual genes are shown as points.

c, Violin plots depicting detection rates in pancreatic scRNA-seq data. Senescence markers are colored by type.

Benchmarking gene set scoring methods with weak markers

Given the weak expression patterns of senescence genes, we systematic evaluate gene set scoring method for senescence detection using weak markers. We selected six widely used methods for active cell detection with predefined gene set, including four bulk-based scoring methods (GSEA[21], GSVA[26], zscore[27], and PLAGE[28]) and two single-cell-based methods (UCell[23] and AUCell[22]). For benchmarking, we selected scRNA-seq datasets from 3 endocrine cell types in the pancreas and 6 major cell types in the brain, all well-characterized and associated with aging-related disorders[25,29]. Since these datasets lack ground-truth senescent cells, we classified cell-type markers into strong (top 50) and weak (rankings 51–100) based on expression specificity, using the weak markers for evaluation. To further assess robustness, we randomly selected 10–50 marker genes to simulate varying gene set sizes. This benchmarking strategy allowed us to comprehensively evaluate method performance across different levels of marker weakness and gene set sizes.

In addition to the six methods above, we present ICE (Iterative-imputation-based Cell Enrichment), a novel approach designed to handle weak markers and dropout effects in scRNA-seq data (**Fig. 2a**). To address technical noise and sparsity, ICE uses a diffusion-based model to smooth cell profiles and impute zero-inflated gene expression[30]. For enrichment score (ES) calculation, ICE embeds GSEA, which performs a weighted random walk along a ranked gene list and measures the overrepresentation of a gene set at the extremes[21]. Unlike conventional methods like GSEA, ICE identifies active cells by detecting peaks in the derivative of the ES trend, providing a binary classification of enriched cells, instead of just numerical scores. To improve robustness for weak markers, ICE iteratively refines gene sets or selects top-correlated genes. This combined imputation and refinement framework ensures reliable enrichment analysis, particularly for low-signal gene sets.

Fig. 2 | **Benchmark of gene set scoring methods using weakly expressed gene markers**.

a, Schematic of the ICE pipeline: (1) denoising and imputation of zero-inflated single-cell data, (2) enrichment score (ES) calculation, (3) significance threshold determination through variation trend analysis, and (4) iterative marker gene refinement to update the ES matrix.

b-e, Pancreatic α cell detection using 10 weak markers: **b**, UMAP visualization comparing bulk-based GSEA (left) and ICE (right) detection; top 2,319 ES cells (red) versus Louvain-clustered α cells (dashed outline). **c**, ES distribution (ICE: colored; bulk-GSEA: gray) with peak-detection threshold (dotted line). Peaks were detected by derivative analysis of

the ES trend. **d**, Precision analysis across ES-ranked cell intervals (10% bins). **e**, ROC curves with AUC values for all methods.

f-i, Pancreatic β cell evaluation (format consistent with above panels).

j-m, Pancreatic γ cell assessment (format consistent with above panels).

ICE outperforms other methods in detecting pancreatic α cells with weak gene markers

Among 8,569 pancreatic cells, we first benchmarked gene set scoring methods for detecting α cells with 10 weak marker genes. Using Louvain clustering as our reference standard, we established a ground truth of 2,319 α cells in the dataset (**Methods**). To evaluate the advantage of ICE's iterative-imputation strategy, we first compared ICE directly with conventional bulk-based GSEA for α cell detection with 10 weak α markers. When analyzing the top 2,319 cells ranked by ES, the bulk-based GSEA approach correctly identified only 1,290 α cells (55.6% precision) (**Fig. 2b**). In contrast, ICE correctly classified 2,276 cells (98.1% precision), demonstrating superior performance by reducing the error rate from 44.4% to 1.9%.

Further examination of the enrichment score distributions revealed critical differences between ICE and conventional GSEA method. ICE's peak detection algorithm precisely identified the biological cutoff, with ES values showing an abrupt decline after the top 2,362 cells — closely matching the reference α cell count (**Fig. 2c**). In contrast, traditional GSEA failed to produce a clear separation between α cells and other cell populations in the ES distribution. These results suggest that ICE framework effectively recovers true expression signals that are obscured by dropout events, enabling more accurate enrichment score calculations.

Next, we systematically evaluated seven gene set scoring methods in α cells. To assess marker expression across different abundance levels, we analyzed the ES-ranked cells in cumulative 10% intervals (from top 10% to 100%). ICE demonstrated exceptional robustness, maintaining >98% α-cell identification precision across all expression thresholds (**Fig. 2d**). In contrast, the remaining six methods exhibited expression-level dependence: while achieving reasonable performance (average precision: 84.6%) for the top 10% of cells, their precision markedly declined to 55.9% when analyzing the full cell population. Receiver operating characteristic (ROC) analysis confirmed these findings, showing that ICE achieved near-perfect classification with Area Under the ROC Curve (AUC) value > 0.99 compared to other methods (AUC = 0.73-0.80; **Fig. 2e**). The results demonstrate ICE's ability to reliably detect cells regardless of marker expression strength.

ICE demonstrates superior performance in pancreatic β and γ cells

Beyond α cells, ICE demonstrated superior performance in identifying pancreatic β cells with 10 weakly

expressed markers. Among 2,505 β cells, traditional GSEA detected only 1,554 (62.0%) with 10 weak markers,

whereas ICE correctly identified 2,442 (97.5%) (**Fig. 2f**). The enrichment score (ES) distribution revealed that

ICE's peak detection effectively distinguished β cells from others, while traditional GSEA failed (**Fig. 2g**).

ICE maintained high accuracy across varying marker expression levels, whereas other methods declined to

60 – 70% as ES decreased (**Fig. 2h**). Consistently, ICE achieved an AUC value of 0.99 for β cell detection,

outperforming other methods (0.80–0.84) (**Fig. 2i**).

For γ cells, which were rare in the dataset (3.0%, n=264), cell detection proved even more challenging

with 10 weak markers. Traditional GSEA identified only 37 (14.0%) γ cells, while ICE detected 227 (86.0%)

(**Fig. 2j**). Despite the small population, ICE's ES distribution allowed peak-based cutoff determination (**Fig.**

2k). ICE exhibited 100% precision in the top 80% of ES-ranked cells, dropping to 86.9% when all cells were

included (**Fig. 2l**). In contrast, other methods performed poorly (11.5–16.2%) across all thresholds. ICE's AUC

value (0.98) far surpassed competing methods (0.69–0.74) (**Fig. 2m**).

ICE surpasses existing methods in multiple brain cell types

We further validated ICE on 16,929 brain cells, starting with inhibitory neurons (InNs, n=3,724) using 10

weak markers. Traditional GSEA correctly classified only 65.3% (2,430/3,724) of InNs, while ICE achieved

97.8% (3,641/3,724) (**Fig. 3a**). ICE's ES distribution enabled precise separation of InNs from other cell types

(**Fig. 3b**). Regardless of marker strength, ICE maintained >97.7% precision, whereas other methods showed

gradual performance decline at low ES (**Fig. 3c**). ICE also attained a near-perfect AUC score (0.99) compared

to others (0.88–0.95) (**Fig. 3d**). The advantage of ICE extended to other brain cell types, including astrocytes,

endothelial cells, excitatory neurons, oligodendrocytes, and oligodendrocyte precursor cells (**Fig. 3e-n**).

Fig. 3 | Comparative performance of cell detection methods across brain cell types.

a-d, Evaluation of inhibitory neuron (InN) detection using 10 weak markers: (**a**) UMAP visualization, (**b**) enrichment score (ES) distribution, (**c**) precision across ES-ranked cells, and (**d**) ROC curves comparing ICE with alternative methods (format consistent with **Fig. 2**).

e-i, Precision analysis for ES-ranked cells in: (**e**) astrocytes (Astro), (**f**) endothelial cells (Endo), (**g**) excitatory neurons (ExN), (**h**) oligodendrocytes (Oligo), and (**i**) oligodendrocyte precursor cells (OPC).

j-n, ROC curves with AUC values for each method in corresponding brain cell types (e-i).

Increased marker set size fails to overcome weak expression limitations by tradtional methods

Since traditional methods face inherent constraints in detecting cells using small (10-gene) weak marker sets, we investigated whether increasing the marker panel size could enhance detection performance. We systematically evaluated pancreatic α cell identification across expanding gene sets (20, 30, 40, and 50 markers). ICE maintained consistently superior performance, achieving sustained high precision (~98.2%) across all marker set sizes (**Fig. 4a-d**). In contrast, other tools showed only modest improvements, with mean precision increasing marginally from 64.5% to 74.9% with marker panels expanding. Importantly, all methods exhibited performance degradation as marker expression strength (ES) decreased. The superior capability of ICE was further corroborated by AUC analysis, where it maintained near-perfect discrimination (~0.99) regardless of marker set size (**Fig. 4e-h**). Other methods showed substantially lower performance, with the mean AUC value increasing from 0.85 to 0.92, despite the fivefold increase in marker gene number. The

results reveal that simply expanding marker set size cannot overcome the fundamental challenge of weak

marker strength in cell detection by traditional methods, while ICE can overcome this limitation by enhancing

signal from low-expression markers.

Fig. 4 | Performance assessment of cell detection methods across varying marker set sizes.

a–d, Precision of pancreatic α cell detection using weak marker sets of increasing size: (**a**) 20 markers, (**b**) 30 markers,
(**c**) 40 markers, and (**d**) 50 markers.

e-h, Corresponding ROC curves and AUC values for each method: (**e**) 20 markers, (**f**) 30 markers, (**g**) 40 markers, and
(**h**) 50 markers.

ICE reveals aging-associated *ATF3* activity in β cells

Given ICE's superior performance, we employed it to detect senescence in human pancreatic β cells (n = 7,528)

across different ages[31](**Fig. 5a**). Our evaluation focused on four senescence markers: the canonical markers

CDKN1A and *CDKN2A*, as well as two novel candidates *ATF3*, and *MX1* (an interferon-I pathway gene[32]).

For each marker, ICE identified the top 20 correlated genes to define marker-specific signatures. Using these

signatures, ICE identified 496 (*CDKN1A*), 632 (*CDKN2A*), 154 (*ATF3*), and 379 (*MX1*) marker-associated

cells (**Fig. 5b**).

While *CDKN1A*, *CDKN2A* and *MX1* displayed diffuse expression patterns, *ATF3*-associated cells were

specifically enriched in subclusters 3 and 6 (**Fig. 5b,c**). This suggests *ATF3*-associated cells formed a distinct

subpopulation of β cells. Pseudobulk analysis in subcluster 3 showed that *ATF3* and *CDKN1A* expression

increased with age, while *MX1* and *CDKN2A* decreased (**Fig. 5d**). In all *ATF3*-associated cells, expression of

ATF3 showed a positive correlation with the donor age (**Fig. 5e**). Interestngly, differential gene expression analysis revealed that *ATF3*-associated cells exhibited upregulation of pro-apoptotic unfolded protein response (UPR)-related genes (**Fig. 5f**). Specifically, *DDIT3* (*CHOP*) functions as a key mediator of ER stress-induced apoptosis[33] and *HSPA5* (*GRP78/BiP*) encodes a master regulator of UPR[34]. These findings indicate that *ATF3*-associated β cells become activated during aging and participate in chronic ER stress, UPR activation, and increased apoptosis susceptibility.

Fig. 5 | Identification of senescence marker-associated β cell subpopulations using ICE.

a, UMAP visualization of pancreatic β cell heterogeneity, revealing distinct subclusters.

b, ICE-based detection of senescence marker-associated cells (red) versus unassociated populations (gray). Gene sets for ICE analysis were constructed by selecting the top 20 correlated genes for each senescence marker.

c, Cluster-specific enrichment patterns of senescence-associated cells. Circle color: significance level from Fisher's exact test. Circle size: proportion of enriched cells per subcluster.

d, Age-related expression dynamics of senescence markers across β cell subclusters.

e, Pseudobulk analysis in *ATF3*-associated cells confirming positive correlation between *ATF3* expression and donor age.

f, Volcano plot of differentially expressed genes in *ATF3*-associated β cells. Upregulated genes associated with UPR pathway are labeled. Pink: up-regulated genes; Blue: down-regulated genes.

ICE uncovers IFN-I-active microglia linked to AD

IFN-I responsive microglia play a role in neurodegenerative disorders and contribute to neuronal engulfment during development and disease[16,35]. Therefore, we applied ICE to investigate the link between IFN-I-active microglia and AD[36]. Unsupervised clustering of 76,346 microglia from human AD brains revealed 13 distinct subclusters (**Fig. 6a**). ICE identified 3,212 (*CDKN1A*), 779 (*CDKN2A*), 1,635 (*ATF3*), and 927 (*MX1*) marker-associated cells, each displaying unique expression patterns (**Fig. 6b**). Notably, *MX1*-associated microglia localized specifically to subcluster 5, while *CDKN1A* and *CDKN2A* were restricted to subclusters 2 and 4 (**Fig. 6c**).

In subcluster 5, *MX1* and *CDKN1A* expression correlated positively with AD progression (Braak stage), whereas *ATF3* exhibited a mild negative correlation (**Fig. 6d**). Further analysis confirmed a significant positive association between *MX1* expression and Braak stage across all *MX1*-associated cells (**Fig. 6e**). These cells also showed upregulation of key regulators in IFN-I signaling genes, including *STAT1*, *STAT2*, *OAS1*, and *OAS2* (**Fig. 6f**). These findings demonstrate ICE's ability to precisely detect cell populations enriched for senescence markers, highlighting a link between IFN-I-active microglia and AD pathogenesis.

Fig. 6 | ICE reveals senescence marker-associated microglia subpopulations in AD.

a, UMAP projection of microglia heterogeneity in AD brains, showing distinct transcriptional subclusters.

b, Detection of senescence marker-associated microglia (red) versus unassociated populations (gray). Gene sets for ICE
analysis were constructed by selecting the top 20 correlated genes for each senescence marker.

c, Subcluster enrichment patterns of senescence-associated cells. Circle color: significance level from Fisher's exact test.
Circle size: proportion of enriched cells per subcluster.

d, Expression correlation of senescence markers with AD Braak stage across different subclusters.

e, Pseudobulk analysis demonstrating positive correlation between *MX1* expression and AD neuropathological
progression (Braak stage).

f, Volcano plot of differentially expressed genes in *MX1*-associated microglia. Upregulated genes associated with IFN-
I signaling pathway are labeled. Pink: up-regulated genes; Blue: down-regulated genes.

Discussion

The emergence of single-cell profiling technologies has revolutionized aging research and provided
unprecedented insights into senescent cell states. However, the reliable identification of senescent cells
remains a significant challenge due to the weak expression of senescence biomarkers. This limitation
underscores the urgent need for novel computational approaches to accurately detecting senescent cells with
subtle molecular signatures.

In this study, we first established a benchmark to evaluate the performance of existing tools in detecting
cells based on weakly expressed gene sets. Our analysis revealed critical shortcomings in current gene set
scoring methods: as marker expression strength decreased, detection accuracy declined across multiple cell
types in the pancreas and brain. For example, in pancreatic γ-cells, accuracy dropped to as low as 11.5% when
markers were particularly faint. Notably, increasing the number of marker genes did not substantially improve
performance, suggesting that the primary limitation lies in the inherent weakness of the markers rather than
the size of the gene set.

To overcome these challenges, we developed ICE, a computational framework that integrates two key
strategies: (1) imputation to recover missing or weak expression signals and (2) iterative refinement of gene
sets to enhance marker specificity. Extensive benchmarking demonstrated that ICE significantly outperformed
existing methods. In pancreatic α cells, ICE achieved over 98% precision and near-perfect AUC even with
weak markers, whereas conventional gene set scoring methods performed poorly (50-60% precision). ICE
showed similar superior performance in identifying diverse cell types in pancreas and brain tissues. These
results highlight the importance of expression data imputation and dynamic gene set optimization for reliable
cell detection in scRNA-seq data.

Applying ICE to key senescence markers (*CDKN1A*, *CDKN2A*, *ATF3*, and *MX1*), we uncovered distinct

cell populations enriched for these markers and explored their functional implications. In pancreatic β-cells, ICE identified a subpopulation with elevated *ATF3* expression, which exhibited heightened unfolded protein response (UPR) pathway activity, marked by *DDIT3* upregulation. Prior studies have shown that *DDIT3* deletion in β cells mitigates ER stress and prevents age-associated metabolic dysfunction in mice[33]. Thus, the *ATF3*-positive β cells detected by ICE may represent a senescent population under chronic ER stress during aging.

Similarly, in the context of neurodegenerative disease, ICE revealed a subset of interferon-responsive microglia enriched in late-stage Alzheimer's disease (AD). These cells, characterized by high *MX1* expression, displayed robust IFN-I signaling activity, which has been linked to AD progression. Given that IFN-I signaling can be triggered by damage-associated nucleic acids, these microglia may be activated by amyloid fibrils containing such molecules[37]. Supporting this notion, blocking IFN-I signaling has been shown to improve cognitive and synaptic function in AD models[38]. These findings suggest that *MX1*-expressing microglia represent a chronically inflamed state associated with AD pathology.

Intriguingly, we observed minimal overlap between different senescence markers, with each marking distinct cell populations. While *CDKN1A* and *CDKN2A*—classic senescence markers—were broadly expressed in pancreatic β cells, *ATF3* marked a more consistent subpopulation exhibiting age-related UPR activation, a phenotype more directly tied to stress-induced senescence. Similarly, in microglia, *CDKN1A* and *CDKN2A* exhibited variable expression across subgroups, underscoring the heterogeneity of senescence markers in human tissues. Moreover, *ATF3* and *MX1* were rarely co-expressed in the cell types examined, suggesting that distinct senescence-associated pathways may operate in different cellular contexts.

In summary, our study uncovered the weak expression patterns of senescence-associated genes across human tissues and cell types. We further demonstrated the limitations of current gene set scoring methods in detecting such subtle markers and introduced ICE, a computational framework that enhances cell-state identification through imputation-coupled iterative marker refinement. By enabling high-resolution dissection of senescence heterogeneity, ICE offers novel insights into the spatiotemporal dynamics of aging across tissues. Moreover, the ICE framework is broadly applicable to diseased cell detection with weakly expressed markers, opening new avenues for studying cellular dysfunction in diverse pathological contexts.

Methods

Overview of ICE

ICE is a single-cell gene set enrichment analysis method designed to identify cells significantly enriched

for a predefined gene set. The method integrates imputation, iterative refinement, and derivative-based cutoff

estimation to enhance the accuracy and robustness of enrichment detection, particularly in the presence of

sparse or weak marker genes. ICE requires two main inputs: a gene expression profile of single cells and a

predefined set of genes (e.g., a gene set of interest). Below, we outline the key steps of the ICE algorithm:

1. **Imputation of Single-Cell Data**

The first step in ICE involves rectifying the entire single-cell gene expression data using the MAGIC

algorithm[30], which was ranked as one of the most accurate imputation method from extensive

benchmark[39]. The imputation begins by constructing a k-Nearest Neighbor (kNN) graph, where each

cell is connected to its k-nearest neighbors based on Euclidean distance to capture cell-to-cell

similarities. Next, an affinity matrix is built using a Gaussian kernel to transform these distances into

similarity scores, weighted by a parameter to control the spread of the similarity function. Finally, a

diffusion process is applied across the kNN graph, propagating expression values based on cell

affinities to smooth the data, impute missing values, and reduce noise. The resulting imputed matrix

provides a more complete and accurate representation of cellular gene expression profiles for

downstream analyses.

2. **Calculation of Enrichment Scores (ES)**

For each cell, ICE computes an ES that reflects the expression activity of the predefined gene set. In

this step, single-sample Gene Set Enrichment Analysis is applied to calculate enrichment scores for

predefined gene sets on a per-cell basis [26]. The process begins by ranking genes within each cell based

on their expression levels, creating an ordered list from highest to lowest expression. For each gene

set, an enrichment score is then computed using a running-sum approach, which assesses the positions

of genes within the ranked list to capture their relative contribution to the gene set's activity. This

results in a matrix of enrichment scores, where each entry represents the activity level of a specific

gene set in a given cell, providing a comprehensive view of gene set expression across the cell

population and enabling insights into cellular states and pathways.

3. **Derivative-Based Cutoff Estimation**

The ES values are arranged in descending order, and forward differences are computed to approximate

the derivatives of the ES trend. The pracma package (v2.4.4) in R was used to calculate the cutoff

value from the initial peak in the derivative trend. This cutoff distinguishes cells significantly enriched

for the gene set from those that are not.

4. **Iterative Refinement via Differential Expression Analysis**

Based on the initial cell classification, ICE performs differential expression analysis using the Seurat

package[40]. The resulting positive markers are used to refine the discovery of enriched cells. The top n

(a variable can be changed depending on users) positive marker genes are chosen to replace the

predefined marker list. The enrichment scores and cutoff values are then recalculated iteratively to

optimize the identification of cells active in the predefined gene set.

The output of ICE not only includes the enrichment scores derived from denoised gene expressions but

also provides a clear cutoff for identifying significantly enriched cells.

Benchmarking different gene set scoring methods

The performance of ICE was systematically benchmarked against four bulk-based scoring methods

(GSEA, GSVA, zscore, and PLAGE) and two single-cell-based scoring methods (UCell and AUCell). GSEA

evaluates the distribution of predefined gene sets within a ranked list of genes to identify significant biological

pathways[21]. GSVA transforms gene expression data into gene set enrichment scores, enabling pathway-level

analysis across samples[26]. The z-score method calculates the standardized difference between the mean

expression of a gene set and the background, providing a simple yet effective enrichment metric[27]. PLAGE

uses singular value decomposition to model gene set activity as latent variables[28]. On the single-cell level,

UCell computes a rank-based enrichment score by assessing the distribution of gene set members within the

ranked expression profile of each cell[23]. AUCell calculates the Area Under the Curve (AUC) for the recovery

curve of gene set members, quantifying their enrichment in individual cells[22].

Two single-cell RNA-seq datasets, derived from human pancreas and brain tissues, were used for

benchmarking analysis. The pancreas dataset (GSE84133) was downloaded from GEO, while the brain dataset

was obtained from the PsychENCODE Consortium[25,29]. Both datasets were preprocessed using the Seurat

pipeline, including normalization with the SCTransform method, dimensionality reduction via PCA,

integration using Harmony, and clustering with the Louvain algorithm at a resolution of 0.2. UMAP was

employed for visualization. Cell type annotations for the pancreas dataset were predicted using the Azimuth

reference, while annotations for the brain dataset were refined by selecting the most frequent cell-type label

per cluster. Differential expression analysis was performed using the Wilcoxon test to identify marker genes

for each cell type.

To evaluate the performance of the methods, three endocrine cell types (α, β, and γ) were analyzed in the

pancreas dataset, while six cell types (inhibitory neurons, astrocytes, endothelial cells, excitatory neurons,

oligodendrocytes, and oligodendrocyte precursor cells) were chosen from the brain dataset. To assess the

impact of marker strength and sparsity, two categories of marker gene sets were defined for each cell type:

strong markers comprised the top 50 most differentially expressed genes, and weak markers included the

subsequent 50 genes. These differentially expressed genes were identified using the Wilcoxon test during cell-

type marker analysis, ranked by their adjusted p-values. Additionally, random subsets of 10, 20, 30, 40, and

50 markers were selected from each category to evaluate the methods under varying marker sparsity conditions.

For each cell type, the clustering labels were used as ground truth annotations. Precision was calculated

as the proportion of correctly classified cells relative to the total number of identified cells based on the ranked

ESs. To further evaluate performance, Receiver Operating Characteristic (ROC) analysis was conducted. The

ROC curve was constructed using the "roc" function from the "pROC" package in R, with the true positive

rate (TPR) and false positive rate (FPR) calculated based on the normalized ES scores. The Area Under the

Curve (AUC) was computed to quantify the performance of each method.

RNA-seq analysis of GTEx database

Raw RNA-seq count data were obtained from the GTEx database (v8). Tissues with fewer than 50 samples

were excluded, resulting in 14,956 samples from 38 tissues derived from 948 non-diseased donors aged 20–

80 years. For each tissue, genes with expression levels > 1 log2CPM (counts per million) in at least 25% of

samples were retained for downstream analysis. Read counts were normalized using the trimmed mean of M-

values (TMM) method to correct for library size differences[41]. Normalized expression values were log2-

transformed. Tissue-specific differentially expressed genes (DEGs) were identified using the limma package

(moderated t-test) [42]. Genes with |log2FC| > 1 and adjusted P-value < 0.05 were classified as significant DEGs

and tissue-specific markers.

Enrichment analysis with single gene markers

To evaluate ICE using single gene markers, we performed correlation analysis based on imputed gene

expression data. For each target gene, the top n most correlated genes (including the target gene itself) were

selected to construct a marker gene set for ICE. In this study, we set n = 20 as part of our exploratory analysis.

Two single-cell RNA-seq datasets related to aging and AD were analyzed: stressed β cells and type-I interferon

(IFN-I)-responsive microglia. The β cell dataset (GSE114297) was obtained from GEO and consisted of islet

samples from 12 donors without diabetes[31]. After preprocessing using the Seurat pipeline (normalization,

clustering, and cell type annotation), the β cell population was selected for ICE analysis. The microglia dataset

was obtained from the Synapse database (syn52293417) and included nuclei sequencing data from the

prefrontal cortex of human individuals with varying degrees of AD pathology and cognitive impairment[36].

The immune cell count matrix was directly downloaded from the database, and the microglia population was

isolated for downstream ICE analysis.

Statistics and Reproducibility

The performance of different methods was evaluated using precision and ROC analyses. The AUC was

computed by the "pROC" package in R to quantify the performance of each method. Enrichment of target

cells within cell subclusters was assessed using Fisher's exact test via the "ClusterProfiler" package.

Correlations between gene expression and donor age or AD pathology (Braak stage) were calculated using

pseudobulked counts for each donor, generated with the "AggregateExpression" function in Seurat. Pearson

correlation was used for these analyses. Differential expression analysis to identify cell-type markers was

performed using the Wilcoxon test from the Seurat pipeline.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Data availability

All datasets used for benchmarking are publicly available. The pancreas dataset was accessed from GEO under

accession number GSE84133. The brain dataset was obtained from the PsychENCODE Consortium. The

stressed β cell dataset was retrieved from GEO under accession GSE114297, and the AD microglia dataset

was obtained from the Synapse database (syn52293417). The R implementation of the ICE package is

available on GitHub (https://github.com/pxulab/ICE). Source data supporting the findings of this study are

provided with the manuscript.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PX and YK perceived the concept and designed the study; PX and HZ performed the data analyses, prepared the figures, and wrote the paper; SZ participated in the data analysis. All the authors participated in the discussion and interpretation of the results, reviewed and revised the paper.

Acknowledgements

This work was primarily supported by the National Natural Science Foundation of China (32470666), the Interdisciplinary Research Program for Young Scholars (2024FGC1004) and start-up fund (RF1028624054 to PX and RF1028623368 to YK) from Southeast University.

References

Cai, Y. *et al.* The landscape of aging. *Sci China Life Sci* (2022). https://doi.org/10.1007/s11427-022-2161-3

Varela-Eirin, M. & Demaria, M. Cellular senescence. *Curr Biol* **32**, R448-R452 (2022). https://doi.org/10.1016/j.cub.2022.04.003

Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks of aging: An expanding universe. *Cell* **186**, 243-278 (2023). https://doi.org/10.1016/j.cell.2022.11.001

He, S. & Sharpless, N. E. Senescence in Health and Disease. *Cell* **169**, 1000-1011 (2017). https://doi.org/10.1016/j.cell.2017.05.015

Coppe, J. P. *et al.* Senescence-associated secretory phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor. *PLoS Biol* **6**, 2853-2868 (2008). https://doi.org/10.1371/journal.pbio.0060301

Ogrodnik, M. *et al.* Guidelines for minimal information on cellular senescence experimentation in vivo. *Cell* **187**, 4150-4175 (2024). https://doi.org/10.1016/j.cell.2024.05.059

Suryadevara, V. *et al.* SenNet recommendations for detecting senescent cells in different tissues. *Nature Reviews Molecular Cell Biology* **25**, 1001-1023 (2024). https://doi.org/10.1038/s41580-024-00738-8

de Magalhaes, J. P. Cellular senescence in normal physiology. *Science* **384**, 1300-1301 (2024). https://doi.org/10.1126/science.adj7050

Zezula, J. *et al.* p21cip1 is required for the differentiation of oligodendrocytes independently of cell cycle withdrawal. *EMBO Rep* **2**, 27-34 (2001). https://doi.org/10.1093/embo-reports/kve008

10   Di Giorgio, E. *et al.* Transcription of endogenous retroviruses in senescent cells contributes to the
accumulation of double-stranded RNAs that trigger an anti-viral response that reinforces senescence.
Cell Death & Disease **15** (2024). https://doi.org/10.1038/s41419-024-06548-2
11   Liu, X. *et al.* Resurrection of endogenous retroviruses during aging reinforces senescence. *Cell* (2023).
https://doi.org/10.1016/j.cell.2022.12.017
12   De Cecco, M. *et al.* L1 drives IFN in senescent cells and promotes age-associated inflammation. *Nature*
566, 73-78 (2019). https://doi.org/10.1038/s41586-018-0784-9
13   Mao, J. *et al.* Reactivation of senescence-associated endogenous retroviruses by ATF3 drives
interferon signaling in aging. *Nature Aging* **4**, 1794-1812 (2024). https://doi.org/10.1038/s43587-024-
00745-6
14   Li, L. *et al.* Human skin rejuvenation via mRNA. *bioRxiv*, 2024.2011.2012.623261 (2024).
https://doi.org/10.1101/2024.11.12.623261
15   Zhang, C. *et al.* ATF3 drives senescence by reconstructing accessible chromatin profiles. *Aging Cell*
20 (2021). https://doi.org/10.1111/acel.13315
16   Roy, E. R., Li, S., Saroukhani, S., Wang, Y. & Cao, W. Fate-mapping and functional dissection reveal
perilous influence of type I interferon signaling in mouse brain aging. *Molecular Neurodegeneration*
19 (2024). https://doi.org/10.1186/s13024-024-00736-6
17   Lai, Y. *et al.* Multimodal cell atlas of the ageing human skeletal muscle. *Nature* **629**, 154-164 (2024).
https://doi.org/10.1038/s41586-024-07348-6
18   Whytock, K. L. *et al.* Aging human abdominal subcutaneous white adipose tissue at single cell
resolution. *Aging Cell*, e14287 (2024). https://doi.org/10.1111/acel.14287
19   Green, G. S. *et al.* Cellular communities reveal trajectories of brain ageing and Alzheimer's disease.
Nature (2024). https://doi.org/10.1038/s41586-024-07871-6
20   Xu, P. *et al.* The landscape of human tissue and cell type specific expression and co-regulation of
senescence genes. *Mol Neurodegener* **17**, 5 (2022). https://doi.org/10.1186/s13024-021-00507-7
21   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for interpreting
genome-wide expression profiles. *Proceedings of the National Academy of Sciences* **102**, 15545-15550
(2005). https://doi.org/10.1073/pnas.0506580102
22   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat Methods* **14**,
1083-1086 (2017). https://doi.org/10.1038/nmeth.4463
23   Andreatta, M. & Carmona, S. J. UCell: Robust and scalable single-cell gene signature scoring. *Comput*
Struct Biotechnol J **19**, 3796-3798 (2021). https://doi.org/10.1016/j.csbj.2021.06.043
24   Avelar, R. A. *et al.* A multidimensional systems biology analysis of cellular senescence in aging and
disease. *Genome Biol* **21**, 91 (2020). https://doi.org/10.1186/s13059-020-01990-9
25   Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas Reveals Inter-
and   Intra-cell   Population   Structure.   *Cell   Syst*   **3**,   346-360   e344   (2016).
https://doi.org/10.1016/j.cels.2016.08.011
26   Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray and RNA-
seq data. *BMC Bioinformatics* **14**, 7 (2013). https://doi.org/10.1186/1471-2105-14-7
27   Lee, E., Chuang, H. Y., Kim, J. W., Ideker, T. & Lee, D. Inferring pathway activity toward precise
disease   classification.   *PLoS   Comput   Biol*   **4**,   e1000217   (2008).
https://doi.org/10.1371/journal.pcbi.1000217
28   Tomfohr, J., Lu, J. & Kepler, T. B. Pathway level analysis of gene expression using singular value
decomposition. *BMC Bioinformatics* **6**, 225 (2005). https://doi.org/10.1186/1471-2105-6-225

29   Li, M. *et al.* Integrative functional genomic analysis of human brain development and neuropsychiatric
risks. *Science* **362** (2018). https://doi.org/10.1126/science.aat7615

30   van Dijk, D. *et al.* Recovering Gene Interactions from Single-Cell Data Using Data Diffusion. *Cell*
174, 716-729 e727 (2018). https://doi.org/10.1016/j.cell.2018.05.061

31   Xin, Y. *et al.* Pseudotime Ordering of Single Human beta-Cells Reveals States of Insulin Production
and Unfolded Protein Response. *Diabetes* **67**, 1783-1794 (2018). https://doi.org/10.2337/db18-0365

32   Sadler, A. J. & Williams, B. R. G. Interferon-inducible antiviral effectors. *Nature Reviews Immunology*
8, 559-568 (2008). https://doi.org/10.1038/nri2314

33   Yong, J. *et al.* Chop/Ddit3 depletion in beta cells alleviates ER stress and corrects hepatic steatosis in
mice. *Sci Transl Med* **13** (2021). https://doi.org/10.1126/scitranslmed.aba9796

34   Kopp, M. C., Larburu, N., Durairaj, V., Adams, C. J. & Ali, M. M. U. UPR proteins IRE1 and PERK
switch BiP from chaperone to ER stress sensor. *Nature Structural & Molecular Biology* **26**, 1053-1062
(2019). https://doi.org/10.1038/s41594-019-0324-9

35   Escoubas, C. C. *et al.* Type-I-interferon-responsive microglia shape cortical development and behavior.
Cell **187**, 1936-1954 e1924 (2024). https://doi.org/10.1016/j.cell.2024.02.020

36   Mathys, H. *et al.* Single-cell atlas reveals correlates of high cognitive function, dementia, and resilience
to   Alzheimer's   disease   pathology.   *Cell*   **186**,   4365-4385.e4327   (2023).
https://doi.org/10.1016/j.cell.2023.08.039

37   Roy, E. R. *et al.* Type I interferon response drives neuroinflammation and synapse loss in Alzheimer
disease. *J Clin Invest* **130**, 1912-1930 (2020). https://doi.org/10.1172/JCI133737

38   Roy, E. R. *et al.* Concerted type I interferon signaling in microglia and neural cells promotes memory
impairment   associated   with   amyloid   beta   plaques.   *Immunity*   **55**,   879-894   e876   (2022).
https://doi.org/10.1016/j.immuni.2022.03.018

39   Hou, W., Ji, Z., Ji, H. & Hicks, S. C. A systematic evaluation of single-cell RNA-sequencing imputation
methods. *Genome Biol* **21**, 218 (2020). https://doi.org/10.1186/s13059-020-02132-x

40   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529 (2021).
https://doi.org/10.1016/j.cell.2021.04.048

41   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
expression   analysis   of   digital   gene   expression   data.   *Bioinformatics*   **26**,   139-140   (2010).
https://doi.org/10.1093/bioinformatics/btp616

42   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* **43**, e47 (2015). https://doi.org/10.1093/nar/gkv007

565