Biomechanics of static roll posture control by body flexion in adult zebrafish

Ryota Nagaoka[1,2,3], Taisei Katayama[1,2,4], Shin-ichi Higashijima[1,2,3], Masashi

Tanimoto[1,2,3,5,*]

[1]Division of Behavioral Neurobiology, National Institute for Basic Biology, Okazaki,

Aichi 444-8787, Japan

[2]Neuronal Networks Research Group, Exploratory Research Center on Life and Living

Systems, Okazaki, Aichi 444-8787, Japan

[3]The Graduate University for Advanced Studies, SOKENDAI, Okazaki, Aichi 444-8787,

Japan

[4]Division of Biological Science, Graduate School of Science, Nagoya University, Nagoya,

Aichi 464-8602, Japan

[5]Lead contact

[*]Correspondence: tanimoto@nibb.ac.jp

Summary

Posture control is crucial for animals. Static posture control in fish remains poorly explored. Recent studies have shown that larval zebrafish perform body flexion during slight roll tilts: the body flexion displaces a gas-filled swim bladder, generating counter-rotation torque for postural recovery through misalignment of gravity and buoyancy forces. Swim bladder deflation impairs this postural recovery, suggesting its critical role. This static posture control strategy may be utilized by many fish species. However, adult fish differ from larvae in morphology and behavior, raising questions about the generality of this mechanism. Our behavioral analysis showed that adult zebrafish also flexed their body during roll tilts, with flexion persisting until recovery to an upright posture. Similarly to larvae, swim bladder deflation impaired postural recovery. These results demonstrate that adult fish employ static roll posture control through body flexion and suggest the generality of this mechanism in fish species.

Introduction

For many animals, posture control is crucial for survival, regardless of whether they live on the land or in the water. Terrestrial animals use both dynamic and static postural control mechanisms.[1–3] When posture is significantly disrupted, animals dynamically move their limb and axial muscles to recover their desired posture. Even during seemingly motionless periods, static postural control through anti-gravity muscle tone continuously maintains posture.

How about postural control in fish? Previous studies have been predominantly centered on dynamic movements involving swimming and fin dynamics.[4–8] In contrast, our understanding of static postural control mechanisms remains limited. This is particularly intriguing because most fish species are inherently unstable[7] - anesthetized or dead fish typically roll to a belly-up position due to their mass distribution.[6,9–11] Nevertheless, many fish species maintain an upright posture even while appearing motionless.[9,10] These common observations suggest the existence of static postural control mechanisms through neuromuscular activity. However, until recently, the underlying mechanisms of this static postural control remained poorly understood.

Recent studies on millimeter-sized larval zebrafish have provided valuable insights into this longstanding question. Larval zebrafish perform body flexion during both slight roll tilt and artificial vestibular stimulation mimicking a head roll tilt.[12–15] This body flexion behavior is driven by neural circuits, from a vestibular nucleus via reticulospinal neurons projecting to the spinal cord, and the posterior hypaxial muscles (PHM) near the swim bladder.[12,16] This posture control through body flexion relies on the presence of the swim bladder, a gas-filled chamber in the body: postural recovery is impaired in swim bladder-deflated larvae.[12] Biomechanics underlying this postural recovery has been proposed.[12]

The body flexion during roll tilt displaces the swim bladder to one side while placing the

head and tail to the other side. Since the density of the swim bladder is orders of

magnitude lower than the rest of the body, this asymmetric positioning of the swim

bladder results in the misalignment of gravity and buoyancy forces, which generates a

corrective moment of force for postural recovery. This fine and static posture control

strategy may be utilized by many fish species possessing a swim bladder or low-density

tissue. However, adult fish differ from larvae in musculoskeletal structure, body size, and

behavioral dynamics,[17–23] raising questions about the generality of this mechanism.

In the present study, behavioral analyses reveal that adult zebrafish utilize body flexion

to correct roll-tilted posture similarly to larvae. When sustained roll tilt stimulus is applied,

the body flexion persists throughout the stimulus. Crucially, swim bladder deflation

disturbs postural recovery, suggesting swim bladder's critical role in maintaining static

posture. These results suggest that zebrafish maintain an upright posture via the fine and

static roll posture control mechanism through body flexion throughout their life.

Results

Adult zebrafish utilize body flexion to correct roll-tilted posture

To investigate how adult zebrafish maintain an upright posture, we developed a fish

rotation apparatus (Figure 1A). A fish was placed in a cylindrical chamber in a dark room

and subjected to postural disturbance in the roll direction. Viscous water containing

methylcellulose was used to efficiently deliver roll stimulus to the fish and to slow down

its behavior for easier observation. Fish's posture was simultaneously recorded by two

cameras positioned at the front and dorsal sides of the chamber. Pigment-less *casper*[24]

fish were used as a standard fish for easier deflation of the swim bladder in a later

experiment (see below).

Rotation of the chamber induced roll tilt in the fish. As previously reported in

larvae,[12,13,15] adult fish occasionally performed dynamic postural control behaviors

accompanied by swim-like vigorous body movement and fin beating. Since the present

study focused on static postural control behaviors, we collected trials in which fish

recovered posture without these dynamic movements. When a roll-tilt stimulus was

applied towards fish's left-down side, fish was slightly tilted to the left-down direction.

Simultaneously, the fish flexed its body to the right (ear-up) side (Figure 1B middle

column, Video S1). As the tilt stimulus progressed, head roll angle and body flexion angle

increased (Figure 1C). After the roll tilt stimulus reached its final angle, fish gradually

recovered the upright posture, while the flexed body relaxed to its original state (Figure

1B right column, Figure 1C, Video S1). Similar results were obtained in wild-type fish

(Figure S1A and S1B). These results suggest that upon postural disturbance in the roll

axis, adult zebrafish recover an upright posture by the body flexion to the ear-up side.

Body flexion continues throughout sustained head roll tilt

To further clarify the relationship between the head roll tilt and body flexion, fish's

anterior body was rigidly restrained to a rectangular chamber using an acrylic holder in

fish tank water without methylcellulose, and a roll tilt stimulus was applied (Figure 2A).

With this setup, fish's head roll angle was directly coupled with the chamber's tilt angle

throughout the experiment. When a roll tilt stimulus was applied towards the fish's left-

down or right-down side and the tilt was kept for 3 s, fish continued the body flexion to

the ear-up side throughout the sustained stimulation (Figure 2B and 2C). In some trials,

fish exhibited swim-like lateral body undulation and tail flick to either left or right side

during the sustained tilt (Figure S1C). As the sustained tilt stimulation was released, the

flexed body relaxed. These results further demonstrate the tight correlation between the

head roll tilt and body flexion.

Biomechanical model of postural recovery through body flexion

Mechanistically, how does the body flexion contribute to the roll postural recovery? A

biomechanical model of the posture control has been proposed in a larval fish study.[12]

The body flexion during roll tilt causes asymmetric positioning of the low-density swim

bladder relative to the whole body, resulting in the misalignment of gravity and buoyancy

forces, which generates a corrective moment of force for postural recovery.[12] In adult

zebrafish, the center of mass (COM) and center of volume (COV) are located near the

constriction region between the anterior and posterior swim bladder chambers,[11] and both

are located on the midline when a fish is at rest: gravity acting on COM and buoyancy

acting on COV are balanced (Figure 3 left column). In a simplified model of adult

zebrafish, when tilted in the roll direction, the fish flexes its body towards the ear-up side

(Figure 3 middle column). The swim bladder shifts toward the ear-down side relative to

the whole body, while the head and tail reposition toward the ear-up side. Because the

swim bladder contains gas with a density orders of magnitude lower than the rest of the

body tissues, the ear-down side positioning of the swim bladder shifts relative positions

of COM and COV, which misaligns the gravity and buoyancy forces. This misalignment

generates a moment of force that counteracts the roll tilt, thereby restoring an upright

posture. According to this model, postural recovery is expected to be disturbed in a swim

bladder-deflated fish. Specifically, in the swim bladder-deflated fish, the density of the

body is nearly homogeneous. As a result, positions of the COM and COV hardly separate,

even when the fish performs body flexion during roll tilt (Figure 3 right column).

Consequently, little moment of force is generated, resulting in significantly impaired

postural recovery. Additionally, the swim bladder-deflated fish is expected to be prone to

roll tilt, since it is not able to efficiently counteract roll tilt through body flexion.

Swim bladder plays a pivotal role in roll posture recovery

To test whether the model holds true in adult fish, as previously demonstrated in larvae,

we performed behavioral analysis in swim bladder-deflated fish (Figure S1D). To fairly

evaluate the impact of swim bladder deflation, control experiments (sham deflation) were

also conducted: this sham treatment involved all the steps performed on the deflated fish

with the exception of rupturing the swim bladder sac and applying negative pressure to

remove the bladder gases.

In response to roll tilt stimulus, sham fish flexed their body and recovered an upright

posture similarly to intact fish. Figure 4A (top photographs) and Video S2 show an

example. When a roll-tilt stimulus was applied towards the fish's right-down side, a sham

fish was slightly tilted to the right-down direction. Simultaneously, the fish flexed its

body to the left (ear-up) side. After the stimulus reached its final angle, the head roll angle

decreased and the body flexion relaxed to the original state (Figure 4A, right column in

the top photographs).

In swim bladder-deflated fish experiments, when a roll-tilt stimulus was applied towards

the fish's left-down side, a fish was slightly tilted to the left-down direction (Figure 4B,

top photographs and Video S2). Simultaneously, the fish flexed its body to the right (ear-

up) side. In contrast to sham fish, the body flexion did not relax to the original state and

the head roll angle only slightly and gradually decreased, after the stimulus reached its

final angle (Figure 4B, right column in the top photographs).

To evaluate the effects of swim bladder deflation on behavioral kinetics, angles and

angular velocities of the head roll and body flexion were compared (Figure 4A and 4B,

bottom traces). In both sham and deflated fish, head roll angle increased as the roll tilt

stimulus angle increased. As predicted based on a biomechanical model described above,

deflated fish are expected to be prone to roll tilt, since they are not able to counteract roll

tilt efficiently through body flexion. Consistent with this model prediction, the maximum

head roll angle was larger in deflated fish (Figure 4C, $p = 0.026$, Mann-Whitney U test).

During postural recovery after head roll angles reached their peaks, the head roll angle

gradually decreased in both sham and deflated fish. Importantly, however, the speed of

head angle change was different: there was a trough in the head angular velocity traces of

sham fish (asterisk in Figure 4A), while troughs were not clearly observed in the traces

of swim bladder-deflated fish (Figure 4B). The minimum head roll angular velocity

showed less negative (i.e., smaller amplitude) values in the deflated fish group compared

to the sham group (Figure 4D, $p = 0.015$, Mann-Whitney U test). Since angular velocity

in the negative direction represents the recovery movement, these less negative (smaller

amplitude) values indicate a slower recovery speed in the swim bladder-deflated fish.

As head roll stimulus angle increased, body flexion angle also increased. In sham fish,

after the stimulus reached the final angle, the body flexion angle decreased toward the

baseline. In deflated fish, in contrast, the body flexion angle remained high and did not

immediately decrease. While there was no significant difference in the maximum body

flexion angle (Figure 4E, $p = 0.699$, Mann-Whitney U test), the recovery speed of body

flexion was slower in the deflated fish group (Figure 4F, $p = 0.002$, Mann-Whitney U

test). These results demonstrated that deflating the swim bladder disturbs posture

recovery despite fish performing sustained body flexion during roll tilt. Collectively,

these results suggest that adult zebrafish, similar to larvae, employ body flexion for static

roll posture control, where the swim bladder plays a crucial role in effective posture

recovery.

Body flexion with intact swim bladder recovers upright roll posture

To further examine whether the moment of force generated by body flexion with an intact

swim bladder is sufficient to recover upright posture, we analyzed passive movement of

chemically fixed fish specimens before and after swim bladder deflation. Fish were

euthanized and chemically fixed in a body-flexed posture (Figure 5A, top). The body

flexion angles ($\theta 1$ in Figure 5A) were between 14 and 24 degrees in the 4 specimens

examined. The fixed fish specimen was held by a pair of forceps in a chamber filled with

fish tank water. To mimic the body orientation of live fish during roll tilt (Figure 1 and

4), the fixed specimen was held in a roll-tilted orientation with its body flexed to the ear-

up side (Figure 5A, bottom). Care was taken to apply equivalent roll tilt angles ($\theta 2$ in

Figure 5A) across before-deflation and after-deflation conditions (before deflation: 24.0

± 4.6 °; after deflation: 24.0 ± 5.9 °). The tilted specimen was gently released and passive

movement was recorded by a front camera.

Once the constraint by the forceps was removed, the specimen before swim bladder

deflation started passive rolling toward the ear-up side (Figure 5B, top photographs and

Video S3). This movement was consistent with the biomechanical model (Figure 3) and

was similar to postural recovery of intact and sham fish (Figures 1 and 4A). A key

difference between intact fish and chemically fixed specimens was the relaxation of body

flexion during postural recovery. In intact fish, the body relaxes to a straight position as

the head roll angle reaches zero (Figure 1B and 1C), which reduces recovery moment of

force and settles into an upright roll posture. In chemically fixed specimens, the body

flexion angle is fixed regardless of head roll angle, which presumably generates a constant

moment of force that keeps the specimen rolling. As expected, the fish specimens kept

rolling even after they reached an upright orientation and head roll angle continuously

decreased (asterisk in Figure 5C, left panel). Similar movements were observed in all 4

fish specimens (asterisk in Figure 5C, right panel).

To examine whether swim bladder deflation disturbs passive roll recovery, after the

passive movement experiments described above, the swim bladder was deflated and the

specimens were again subjected to the same passive movement experiments. In contrast

to the specimens before swim bladder deflation, deflated specimens did not consistently

roll toward the ear-up direction (Figure 5B, bottom photographs and Video S3). They

instead did not roll or slightly rolled in a random direction (double asterisks in Figure 5C,

left panel). Similar movements were observed in all 4 fish specimens (double asterisks in

Figure 5C, right panel). Head roll angular velocities were calculated from the period from

0.2 to 0.5 s after the release. Angular velocity was significantly reduced after swim

bladder deflation (before deflation: -289.4 ± 47.9 °/s; after deflation: 29.4 ± 107.4 °/s; p

= 3.5e-23, Welch's t-test). These results further demonstrated that body flexion with an

intact swim bladder is sufficient to produce passive recovery moment of force for roll

postural recovery.

Discussion

The present study has established an experimental paradigm for analyzing roll postural

control behavior in adult zebrafish and demonstrated that when slightly roll tilted, adult

zebrafish exhibit body flexion to recover an upright posture (Figure 1). Both the tight

correlation between body flexion and head roll angle (Figure 2) and the dependence on

swim bladder for postural recovery (Figure 4) suggest that the mechanisms underlying

roll postural maintenance through body flexion are similar to those in larvae (Figure 3).[12]

The passive tilt experiment using chemically fixed fish specimens demonstrated that the

flexed body with intact swim bladder is sufficient to produce recovery moment of force

(Figure 5). The conservation of these mechanisms between larvae and adults suggests that

body flexion serves as a fundamental strategy for static roll postural maintenance

throughout their life.

Previous researches on larvae suggested that the otolith organs in the vestibular system

sense roll tilt and induce the body flexion. [12,13,15] Not only the otolith system, but fish may

also use sensory information from eyes, semicircular canals, and lateral line, to perform

the body flexion during roll tilt. In our experiments, however, the body flexion was

observed in dark conditions. Therefore, visual inputs are not necessary for the body

flexion. Moreover, in restrained behavioral experiments, fish maintained the body flexion

throughout the sustained tilt stimulus, during which angular velocity and water flow were

minimal, while gravitational input from otolith organs persisted. This observation

suggests that the gravity-sensing otolith organs play a primary role in inducing this

behavior.

Neural circuits and muscles that induces the body flexion behavior during roll tilt were

identified in zebrafish larvae.[12,13,16,25,26] A vestibular nucleus (tangential nucleus) receives

head roll tilt signals from vestibular primary afferents and send neural signals to a group

of reticulospinal neurons in the nucleus of the medial longitudinal fasciculus that projects

to the spinal cord.[12] These neuronal nuclei also exist in adult fish.[27–29] This conservation

of the neuronal nuclei and similarity of the body flexion behavior between larvae and

adults suggest that the same neural circuits likely play a principal role in inducing the

body flexion in adult fish. Regarding the muscles responsible for the body flexion during

roll tilt, adult zebrafish appeared to flex their body near the middle of the body where the

caudal chamber of the swim bladder is located (Figure 1B and 2B). In larvae, muscle

contraction during roll tilt also appears to occur only near the swim bladder.[12] Consistent

with this, the PHM muscles lateral to the swim bladder generate body flexion, with slow-

type PHM primarily responsible for this movement.[12,16] These data suggest that the slow-

type PHMs also generate the body flexion in adults. Muscular organization in adult fish,

however, is more complex and extensively developed compared to larvae.[17,19,30,31] Thus,

other muscles, such as trunk slow muscles that are recruited during weak and sustained

body movements, may also contribute to the body flexion.[32] Experimentally examining

the causal relationship between the body flexion and neuromuscular cell populations in

adult fish is technically challenging due to their size, opaque tissues, ossified skull, and

lack of gene promoters for selectively targeting a specific cell population. However,

future advancements in technology, including optical, genetic, and chemical approaches,

as well as combination of these, may help overcome these limitations and clarify neural

and muscular mechanisms.

In a biomechanical model, the body flexion toward the ear-up side displaces the low-

density swim bladder toward the ear-down side, resulting in separation of the COM and

COV (Figure 3). The magnitude of the recovery moment of force is proportional to the

distance between the COM and COV, which is maximized under a condition where the

swim bladder, the least dense part of the body, is located near the COM, and the body

flexes at the position near the swim bladder. The body organization and the body flexion

in larval zebrafish meet this condition, indicating that the body flexion is a reasonable,

energy-efficient behavior for postural correction.[12] Similarly in adult zebrafish, the

COM—positioned at approximately one third of the total body from the rostral end—is

located around a constriction region of the swim bladder separating the rostral and caudal

chambers.[11] Regarding the body flexion region, as discussed above, the body flexed

around the middle of the body (Figures 1B, and 2B, 4A and 4B), the location of which

was slightly caudal to the swim bladder. Thus, the flexion region seems to slightly

caudally deviate from the most biomechanically efficient location. This subtle deviation

may be attributed to the musculoskeletal organization in adult fish. In the abdominal

region in the rostral body, bony structures increase the body stiffness: lateral angular

stiffness of the vertebral column is higher in the abdominal region than the caudal body

region,[33] and the rib cage bones covering internal organs presumably increase the stability

in the abdominal region. These morphological characteristics make the rostral body

flexion challenging and constrain the flexion region primarily to the middle and caudal

body.[34,35] Considering these morphological constraints, the flexion at the middle of the

body is presumably an adaptive compromise between biomechanical efficiency and

structural limitations. While the results from the swim bladder deflation experiments in

live fish (Figure 4) and fixed fish specimens (Figure 5) supported the biomechanical

model (Figure 3), the precise locations of the COM and COV during body flexion have

not been provided. Further investigation through morphometric analyses would help

address these questions.

Swim bladder-deflated fish is prone to roll tilt and the maximum head roll angle was

larger than sham control (Figure 4C). According to the biomechanical model, the deflated

fish are able to counteract roll tilt by increasing their body flexion angle. However, the

maximum body flexion angle was comparable between control and deflated fish (Figure

4E). This may also be attributed to the morphological constraints described above.

Previous studies showed that during tilt in the roll axis, fish exhibit body undulation with

asymmetric motor pattern that recovers their posture.[4,5,8] Consistently, in the present

study, swim-like vigorous body movement and fin beating were observed when adult

zebrafish was roll tilted. Intriguingly, in experiments where the fish's rostral body was

restrained and sustained tilt stimulation was delivered, fish occasionally exhibited a

swim-like body undulation and tail flick behaviors in addition to their body flexion as

larva (Figure S1C).[13,15] This indicates that fish maintain roll posture primarily through

static control via body flexion, and recruit dynamic control through vigorous body

movements when static control alone is insufficient.

What are the advantages of static posture control through body flexion? Compared to

swimming and fin movements, maintaining posture through slight body flexion is less

visually conspicuous and therefore likely reduces the risk of being detected by

predators.[36,37] Additionally, body flexion presumably requires less energy compared to

vigorous axial undulation during swimming.[38] This is particularly beneficial during rest

periods when maintaining postural stability while conserving energy is crucial.

The mechanism of posture control by body flexion is likely a widely used strategy

employed by many aquatic organisms with low-density organs. While our study focused

on the swim bladder, not only the swim bladder in many fish species, but also lungs in

lungfish and lipid-rich liver in sharks may play a crucial role when these animals flex

their body upon roll tilt.[39–41] Further investigation across a broader range of species will

elucidate the generality of the body flexion-mediated, static roll posture control

mechanism.

Limitations of the study

Swim bladder-deflated fish showed slow but noticeable posture recovery (Figure 4). Technical limitations in the deflation procedure might explain this slow recovery: the swim bladder may not have been fully deflated. While the posterior chamber of the swim bladder was deflated, the anterior chamber was difficult to observe and therefore its deflation was not confirmed. Small amount of residual gas might have generated a small recovery moment of force during body flexion.

Although the present study has demonstrated that body flexion plays a crucial role in roll posture control, this mechanism is likely not the only strategy fish employ to maintain an upright posture. Previous studies on fish postural control have primarily focused on dynamic control mechanisms. Fin movements vary considerably among fish species. Some species skillfully control their fins to maintain posture even while they do not locomote.[42–44] In these species, fin movements, particularly the pectoral fins, may play a primary role in maintaining posture. Further research is needed to understand how fish selectively employ different postural control strategies - static control through body flexion versus dynamic control through fin movements - and how these strategies and body structure may be adapted to their habitat.

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Masashi Tanimoto (tanimoto@nibb.ac.jp).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All data reported in this paper will be shared by the lead contact upon request. All original code has been deposited at Zenodo at https://zenodo.org/records/14551001 and is publicly available as of the date of publication. Any additional information is available from the lead contact upon request.

Acknowledgements

The authors are grateful to Higashijima lab members for their help with fish care and discussion. This work is supported by Japan Society for the Promotion of Science KAKENHI Grant Numbers JP20K06866 and JP23K05983.

Author contributions

R.N., T.K., S.H., and M.T. designed the research; T.K. built a prototype of the experimental apparatus under the supervision of M.T.; R.N. established the apparatus, performed experiments, and analyzed data under the supervision of M.T.; R.N. and M.T. wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

STAR Methods

Key resources table

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Fish

Adult wild-type and *casper*[24] zebrafish (38-45 mm in total length) were maintained under

light-dark cycles (10:14 or 12:12 hours) under standard conditions.[45] For all experiments,

both sexes of fish that displayed normal body morphology and maintained a dorsal-up

posture while at rest were used. For unrestrained behavioral experiments of intact fish, 2

males and 2 females of *casper* fish and 2 males and 2 females of wild-type fish were used.

For restrained behavioral experiments, 2 males and 1 female of wild-type fish and 1 male

and 2 females of *casper* fish were used. For unrestrained behavioral experiments of

casper fish with swim bladder deflation, 3 males and 3 females were used for sham

deflation, and 3 males and 3 females were used for deflation. For fixed specimen passive

movement experiments, 2 males and 2 females of wild-type fish were used. All

experiments were conducted according to animal experiment protocols approved by the

animal care and use committees of the National Institutes of Natural Sciences.

METHODS DETAILS

Unrestrained behavioral experiments

The fish chamber consisted of two acrylic cylinders of different sizes (22 mm [inner

diameter] $\times$ 50 mm [length] $\times$ 1.5 mm [thickness] and 46 mm [inner diameter] $\times$ 10 mm

[length] $\times$ 2 mm [thickness]) connected by an acrylic ring plate. The two cylinders were

separated by a transparent, fluorinated ethylene propylene sheet (NR0538-002,

Fronchemical), which had similar refractive index as water. Small holes were made on

this sheet so that water passes between the two cylinders. Both ends of the connected

chamber were covered with acrylic lids. The two cylinders served different purposes: the

smaller cylinder housed a fish, while the larger cylinder supplied oxygen and, by

manually tilting the chamber, removed small air bubbles from the smaller cylinder. Air

bubble was maintained in the larger cylinder for oxygen supply. The chamber was filled

with fish tank water containing 0.8% methylcellulose (22223-65, Nacalai Tesque, inc.)

unless otherwise noted, and a fish was placed in the smaller cylinder.

The rotation stimulus device was composed of a motorized rotation stage (DDR100/M,

ThorLabs) and supporting mechanical components. The fish chamber was mounted on

this device. The stage was driven by a stage controller (BBD301, ThorLabs) using a

software (Kinesis version 1.14.49, ThorLabs). Fish were illuminated by an infrared light-

emitting diode (IR-LED) plate (LFL-100ir2-850, CCS) from the ventral side and another

IR-LED (HLV3-22IR860, CCS) from the caudal side. Experiments were conducted in a

dark room to eliminate visual cues. The chamber was rotated 45° in the roll direction

(maximum velocity: 40°/s, acceleration and deceleration: 25°/s²) unless otherwise noted.

Fish behavior was recorded at 60 frames per s (fps) by two cameras (HC-VX992MS,

Panasonic) from the frontal and dorsal sides. The timing of rotation stimulus and camera

image frames were synchronized by a flash of an IR-LED, which was oriented towards

two cameras and distinct from the illumination IR-LEDs. In each trial, a voltage pulse

was sent from the rotation stage controller to the IR-LED through a pulse generator

(SEN7203, Nihon Kohden) at 1 s before the rotation started.

Since swim bladder-deflated fish were prone to roll tilting, they frequently exhibited

dynamic posture control through swimming and tail flick when subjected to the standard

roll stimulation described above. To better observe static posture control via slight body

flexion, a weaker rotation stimulus of 35-45° (maximum velocity: 30-40°/s, acceleration

and deceleration: 20-25°/s²) was applied. The reduced buoyancy caused by swim bladder

deflation was compensated by 5-32% (in most cases 10-25%) sucrose (196-00015,

Fujifilm Wako), depending on neutral buoyancy of individual fish, in fish tank water

containing 0.8% methylcellulose (22223-65, Nacalai Tesque, inc.) solution. For wild-type

fish, which were slightly larger than *casper* fish, a cylinder chamber with larger

dimensions (26 mm [inner diameter] × 50 mm [length] × 2 mm [thickness] and 46 mm

[inner diameter] × 10 mm [length] × 2 mm [thickness]) was used.

Swim bladder deflation

Pigment-less *casper* fish were used for easier identification of the swim bladder position.

Fish were anesthetized with 0.01% MS-222 (Ethyl 3-aminobenzoate methanesulfonate,

A5040, Sigma-Aldrich) for 10 minutes, and after confirming the absence of response to

tactile stimuli, a syringe needle (27G [NN-2719S, Terumo] or 30G [Dentronics]) was

gently inserted into the caudal chamber of the swim bladder from a lateral and slightly

dorsal and caudal direction. Negative pressure was gradually applied to the syringe to

remove the air from the swim bladder. Successful swim bladder deflation was confirmed

by the caudal chamber of the swim bladder becoming invisible. Following deflation, fish

were returned to tank water, allowed to rest for 10 minutes before behavioral experiments.

Sham deflation

Sham swim bladder deflation was performed almost as those for deflation procedure

described above. The only modification was that the swim bladder sac was not penetrated

by a syringe needle and swim bladder gas was not removed for sham deflation. Even

slight piercing of the swim bladder sac with a syringe needle resulted in air leakage, which

would have compromised the intended control condition. Therefore, sham fish underwent

identical handling, surgical preparation, and recovery procedure to those used for deflated

fish, but with the swim bladder left intact avoiding penetration of the swim bladder sac.

Restrained behavioral experiments

Fish were anesthetized with 0.01% MS-222 for 10 minutes. The fish's body between the

operculum and pectoral fin was immobilized by a separable pair of acrylic plates that

have a ditch: an elliptical hole (major axis: 6.0-6.75 mm; minor axis: 4-5 mm) was formed

when these plates were combined together. To stabilize the body position, a small amount

of flexible silicone (Macks Earplugs) was applied to the space between the hole and the

fish body. The plate holding the fish was then placed into a groove of an acrylic chamber

(60 mm [length] × 30 mm [width] × 45 mm [height]) filled with fish tank water without

methylcellulose. An open-topped rectangular box (55 mm [length] × 30 mm [width] × 20

mm [height]) was placed over the fish to prevent image distortion when tilt stimulus was

applied and the water surface tilted. Fish was illuminated by an IR-LED (LFL-100ir2-

850, CCS) placed below the chamber. The rotation stimulus device was the same as that

described in 'Unrestrained behavioral experiments' section. The chamber was mounted

on the rotation stimulus device, and 20° roll-tilt stimulation (maximum velocity: 10°/s,

acceleration and deceleration: 5°/s²) was applied and held for 3 s before returning to its

original position. Fish behavior was recorded at 25 fps by a camera (daA720-520um,

Basler) with a lens (13VM308ASIRII, Tamron) from the dorsal side. The camera tilted

together with the fish chamber. The timing of the rotation stimulus and the camera image

frames were synchronized by voltage pulses. Specifically, from 2 s before the rotation

started in each trial, image frame acquisition was triggered by voltage pulses from the

rotation stage controller through a pulse generator (SEN-7203, Nihon Kohden).

Fixed-specimen passive movement experiments

Fish were euthanized by placing them with 0.04% MS-222 for 30 minutes. The fish

specimens were restrained on a sponge plate using metal pins, with the body maintained

in a bent posture such that the body flexion angle was approximately 20 degrees. The

body bend direction was to the left in 2 fish and to the right in 2 fish. The fish specimens

were fixed with 4% paraformaldehyde (158127, Sigma-Aldrich) in 1x phosphate buffered

saline (PBS) for 30 minutes at room temperature. After brief rinse with 1x PBS, the fish

specimens were photographed from the dorsal direction by a camera (HC-VX992MS,

Panasonic). The body flexion angle of the fixed fish specimens was between 14 and 24

degrees in 4 specimens examined.

The fixed fish specimen was held by a pair of forceps in a chamber filled with fish tank

water. The held specimen was manually roll tilted from the dorsal-up orientation with its

body flexed to the ear-up side. The tilted specimen was released and passive movement

was recorded by a front camera (HC-VX992MS, Panasonic) at 60 fps. Care was taken to

minimize external forces to the specimen after the release. Swim bladder deflation was

performed as described above.

QUANTIFICATION AND STATISTICAL ANALYSIS

Image analysis of live fish experiments

Captured videos were analyzed with a custom program written in Python (version 3.12.7).

Videos from unrestrained behavioral experiments were downsampled to 30 fps, which

was sufficient to capture behavioral dynamics. Videos from restrained behavioral

experiments were analyzed at the original frame rate. For analysis, videos were converted

into series of TIFF images. The head roll angle was calculated as follows: each image

from the frontal camera was binarized to isolate pixels containing the eyes, and the xy

coordinates of each eye's center were estimated. The head roll angle was then measured

as the angle between the line connecting the centers of the two eyes and the horizontal

line. To extract slow changes in head angle, a 21-frame moving average was applied.

The body flexion angle was calculated using images from the dorsal camera. Each image

was binarized to extract pixels containing the fish while excluding the fins. The fish's

midline, from snout to tail, was divided into 23 segments. The body flexion angle was

determined by calculating the angle between two lines: (1) the line connecting the second

and twelfth points from the tail, and (2) the line connecting the twelfth and twenty-second

points. These points were selected since the two lines best represent the body flexion in

most cases. In image frames where the fish's snout or tail overlapped with the contour of

the cylinder in the images, the point closest to the body flexion point (either the thirteenth,

fourteenth, or fifteenth point) was chosen as the middle point. The angle was calculated

between (1) the line connecting the second and middle points, and (2) the line connecting

the middle and twenty-second points. The dorsal camera images were distorted by optical

refraction through the curved surface of the acrylic cylinder, resulting in slight expansion

of image objects in the cylinder's short-axis direction. The magnitude of distortion

increased from the top to the bottom of the cylinder along the depth axis. It also increased

from the center to the edge along the cylinder's short-axis direction in horizontal planes,

except in the middle horizontal planes where the image slightly shrank near the cylinder's

edge. However, since the body flexion angle was measured from the body midline, this

image distortion would minimally affect the angle measurements. Also, since the fish's

location was randomly distributed in the cylinder across multiple trials, any potential

effects of image distortion would be averaged out, thus having minimal impact on the

angle analysis. Therefore, the body flexion angle was calculated from raw images.

For all measurements except dynamic movement analysis (Figure S1C), image frames

where fish exhibited swimming or tail flick were excluded from the dataset. All time-

series head roll and body flexion angle data were offset by subtracting the pre-stimulus

baseline that was average angle during the initial 1 s of each trial. Supplementary videos

were made from image series using a software (Clipchamp, Microsoft).

Image analysis of fixed-specimen passive movement experiments

Head roll angle was calculated from captured videos using the same methods described

above for live fish. Videos were analyzed at the original frame rate. For image frames

where the custom tracking program failed to detect eye positions, these eye positions were

manually annotated. Only trials where fish specimens showed no visible influence of

external forces from the pair of forceps after release were included in the analysis. For

each individual fish specimen, 10 trials each were randomly selected from the before-

deflation and after-deflation conditions.

Definition of Measured Values and Statistical Analysis

Data was analyzed using a custom code written in Python (version 3.12.7). A single trial

where head roll angle exceeded more than 40 degrees were excluded from the analysis.

Head roll angular velocity and body bend angular velocity were calculated from angle

data in each trial using the numpy.gradient function from the Python NumPy library. A

15-frame moving average was applied to the head roll angular velocity and a 13-frame

moving average to the body bend angular velocity. The maximum head roll angle and

body bend angle were defined as the peak values after the stimulus onset. The minimum

angular velocity was defined as the lowest value after the tilt stimulus onset. For fixed-

specimen passive movement, head roll angular velocities were calculated as the slope of

linear regression fitted to the angle data during the period from 0.2 to 0.5 s after

release.All statistical analyses were performed using the SciPy tool (version 1.13.1). The

mean value for each individual was calculated from in the recorded trials. Using the

individual means, the overall mean and standard deviation were calculated for both sham

and swim bladder-deflated fish. The effect of swim bladder deflation on posture recovery

was evaluated using the two-sided Mann-Whitney U test or two-sided Welch's t-test.

Supplemental information

Document S1. Figure S1

Video S1. Static posture control by body flexion during roll-tilt stimulus in intact adult

zebrafish, related to Figure 1

Video S2. Static posture control by body flexion during roll-tilt stimulus in sham deflation

and swim bladder-deflated adult zebrafish, related to Figure 4

Video S3. Passive movement of fixed fish specimens before and after swim bladder

deflation, related to Figure 5

Figure legends

Figure 1. Static roll posture control through body flexion in adult zebrafish.

(A) Schematic illustration of unrestrained behavioral experiments. Behavior of adult

zebrafish during roll tilt stimuli was recorded using dorsal and frontal view cameras in a

dark room. (B, C) Behavioral experiments on intact adult *casper* fish. (B) Snapshots of a

fish during a roll tilt stimulus towards its left-down side (top row: frontal view; bottom

row: dorsal view). Magenta lines connecting the center of eyes show head roll angle.

Black and magenta dashed lines denote the midlines of the rostral and caudal body,

respectively. Changes in the head roll angle and body flexion angle relative to the pre-

stimulus baselines are indicated at the upper right. Fish tank water contains

methylcellulose. Before stimulus onset, the fish maintains an upright posture (0.0 s). As

the fish is tilted in the roll direction, it flexes its body towards the ear-up side (2.5 s). The

fish recovers its upright posture as it straightens its body (6.0 s). Scale bars indicate 3 mm.

(C) Time courses of changes in head roll angle and body flexion angle in response to roll

tilt stimuli. Average traces from four individual fish are shown in different colors (7 to 9

trials per fish). See also Figures S1A and S1B.

Figure 2. Restrained behavior during static roll tilt.

(A) Schematic illustration of restrained behavioral experiments. A fish was restrained and

subjected to roll tilt stimuli in a dark room. Its behavior was recorded by a dorsal camera,

which tilted together with the fish chamber. (B) Snapshots of a restrained fish during a

roll tilt stimulus towards its left-down side. Before stimulus onset, the body is straight (-

2.0 s). While the sustained tilt stimulus is applied, the fish maintains body flexion (6.0 s).

As the tilt stimulus is released, the fish straightens its body (12.0 s). Scale bars indicate 3

mm. (C) Time courses of body flexion angle in response to roll tilt stimuli. Average traces

from six individual fish (5 to 8 trials per fish) are shown (red: *casper* fish; blue: wild-type

fish). See also Figure S1C.

Figure 3. A mechanistic model of static roll posture control through body flexion in adult

fish.

A biomechanical model of postural recovery from roll tilt by body flexion adapted from

a model for larval zebrafish in Sugioka et al.[12] Gravity and buoyancy forces act at the

center of mass (COM) and center of volume (COV), respectively. The COM and COV

are located near the constriction region between the anterior and posterior swim bladder

chambers.[11] Left column: At rest, the fish's COM and COV are located on the midline,

and the gravity and buoyancy forces are nearly balanced, enabling the fish to maintain an

upright posture. Middle column: When the fish is tilted in the roll direction, it flexes its

body towards the ear-up side, causing the swim bladder to shift towards the ear-down

side. The extremely low density of the swim bladder results in a shift in the relative

positions of the COM and COV. This misalignment of gravity and buoyancy axes

generates a moment of force that restores the upright posture. Right column: A fish with

a deflated swim bladder flexes its body when tilted in the roll direction. Due to nearly

uniform body density, the relative position of COM and COV remains almost unchanged.

Consequently, the moment of force is greatly reduced, preventing restoration of upright

posture.

Figure 4. Behavior of static roll posture control in swim bladder-deflated fish.

(A and B) Behavioral images and kinetics of sham control and swim bladder-deflated

casper fish. (Top photographs) Snapshots of a sham fish (A) and swim bladder-deflated

fish (B) during a roll tilt stimulus (top row: frontal view; bottom row: dorsal view). Fish

tank water contains methylcellulose (A) and methylcellulose and sucrose (B). A sham

fish maintains an upright posture before stimulus onset (0.0 s). As the fish is tilted in the

right-down direction, it flexes its body towards the left (ear-up) side (2.5 s). The fish

recovered upright posture, while its body flexion relaxed to the original state (6.0 s). A

deflated fish maintains an upright posture before stimulus onset (0.0 s). As the fish is

tilted in the left-down direction, it flexes its body towards the right (ear-up) side (2.5 s).

The fish remains roll tilted, while it continues body flexion (6.0 s). Scale bar indicates 3

mm. (Bottom traces) Time courses of changes in head roll angle, head roll angular

velocity, changes in body flexion angle, and body flexion angular velocity in response to

roll tilt stimuli. Average traces from six individual fish are shown in different colors (4 to

8 trials per fish). In angular velocity traces of head roll and body flexion, there are troughs

(asterisk and double asterisks) during postural recovery in sham fish group, while troughs

are not clearly observed in deflated fish group. (C-F) Comparison of behavioral

parameters between sham group and swim bladder-deflated group. (C) Maximum head

roll angle. $p = 0.026$, Mann-Whitney U test. (D) Minimum head roll angular velocity. p

$= 0.015$, Mann-Whitney U test. (E) Maximum body flexion angle. $p = 0.699$, Mann-

Whitney U test. (F) Minimum body flexion angular velocity. $p = 0.002$, Mann-Whitney

U test.

Figure 5. Fixed-specimen passive movement experiments.

(A) Schematic illustration of fixed-specimen passive movement experiments. A fish was euthanized and chemically fixed in a body-flexed posture with body flexion angle ($\theta 1$) approximately 20 degrees. The fixed fish specimen was held by a pair of forceps in a chamber filled with fish tank water. The held specimen was manually roll tilted approximately 24 degrees ($\theta 2$) with its body flexed to the ear-up side. The tilted specimen was released and passive movement was recorded by a front camera. (B) Snapshots of passive movement of a released fish specimen before (top row) and after (bottom row) swim bladder deflation. Before deflation, the specimen rolls toward ear-up side and continues rolling. After deflation, the specimen does not roll. Scale bar indicates 3 mm. (C) Time courses of head roll angle before (asterisk) and after swim bladder deflation (double asterisks). (Left) Data from an example fish. (Right) Data from 4 fish. Black traces represent mean and gray areas represent standard error of the mean.

Figure S1. Unrestrained behavioral experiments in wild-type fish, dynamic movement during roll tilt, and swim bladder-deflation.

(A, B) Unrestrained behavioral experiments in wild-type adult zebrafish performed in a dark room. (A) Snapshots of a fish during a roll tilt stimulus towards its left-down side (top row: frontal view; bottom row: dorsal view). Magenta lines connecting the center of eyes show head roll angle. Black and magenta dashed lines denote the midlines of the

rostral and caudal body, respectively. Changes in the head roll angle and body flexion

angle relative to the pre-stimulus baselines are indicated at the upper right. Before

stimulus onset, the fish maintains an upright posture (0.0 s). As the fish is tilted in the roll

direction, it flexes its body towards the ear-up side (2.5 s). The fish recovers its upright

posture as it straightens its body (5.0 s). Scale bars indicate 3 mm. (B) Time courses of

head roll angle and body flexion angle in response to roll tilt stimuli. Average traces from

four individual fish are shown in different colors (6 to 9 trials per fish). (C) Time course

of body flexion angle in response to sustained roll tilt stimulation in body-restrained

experiments. Example 5 trials where fish exhibited dynamic movement are shown in

different colors. (D) Lateral view images of pigment-less *casper* fish before and after

swim bladder deflation. Scale bar indicates 5 mm.

Video S1. Static posture control by body flexion during roll-tilt stimulus in intact adult

zebrafish.

Front and dorsal view video clips in intact *casper* fish. Video plays at real-time speed.

Video S2. Static posture control by body flexion during roll-tilt stimulus in sham deflation

and swim bladder-deflated adult zebrafish.

Front and dorsal view video clips in sham deflation (first half) and swim bladder-deflated

(second half) *casper* fish. Video plays at real-time speed.

Video S3. Passive movement of fixed fish specimens.

Front view video clips of a fixed fish specimen before (first half) and after (second half)

swim bladder deflation. Video plays at ×0.25 speed.

References

1.   Horak, F. B. Postural orientation and equilibrium: what do we need to know
about neural control of balance to prevent falls? *Age Ageing* 35, ii7–ii11 (2006).

2.   Le Mouel, C. & Brette, R. Mobility as the Purpose of Postural Control. *Front*
Comput Neurosci 11, (2017).

3.   Maki, B. E., Mcilroy, W. E. & Fernie, G. R. Change-in-support reactions for
balance recovery. *IEEE Engineering in Medicine and Biology Magazine* 22, 20–
26 (2003).

4.   Zelenin, P. V., Orlovsky, G. N. & Deliagina, T. G. Sensory-Motor
Transformation by Individual Command Neurons. *The Journal of Neuroscience*
27, 1024–1032 (2007).

5.   Bagnall, M. W. & McLean, D. L. Modular Organization of Axial Microcircuits in
Zebrafish. *Science (1979)* 343, 197–200 (2014).

6.   Webb, P. W. & Weihs, D. Hydrostatic stability of fish with swim bladders: not all
fish are unstable. *Can J Zool* 72, 1149–1154 (1994).

7.   Ehrlich, D. E. & Schoppik, D. Control of Movement Initiation Underlies the
Development of Balance. *Current Biology* 27, 334–344 (2017).

8.   Zelenin, P. V., Grillner, S., Orlovsky, G. N. & Deliagina, T. G. The pattern of
motor coordination underlying the roll in the lamprey. *Journal of Experimental*
Biology 206, 2557–2566 (2003).

9.   Fath, M. & Tytell, E. D. Day and night posture of the bluegill sunfish (*Lepomis*
macrochirus). Preprint at https://doi.org/10.1101/2023.07.13.548884 (2023).

10.   Webb, P. W. & Weihs, D. Stability versus Maneuvering: Challenges for Stability
during Swimming by Fishes. *Integr Comp Biol* 55, 753–764 (2015).

11.   Robertson, G. N., Lindsey, B. W., Dumbarton, T. C., Croll, R. P. & Smith, F. M.

The contribution of the swimbladder to buoyancy in the adult zebrafish (*Danio*
rerio): A morphometric analysis. *J Morphol* 269, 666–673 (2008).

12.   Sugioka, T., Tanimoto, M. & Higashijima, S. Biomechanics and neural circuits
for vestibular-induced fine postural control in larval zebrafish. *Nat Commun* 14,
1217 (2023).

13.   Beiza-Canelo, N. *et al.* Magnetic actuation of otoliths allows behavioral and
brain-wide neuronal exploration of vestibulo-motor processing in larval zebrafish.
Current Biology 33, 2438-2448.e6 (2023).

14.   Migault, G. *et al.* Whole-Brain Calcium Imaging during Physiological Vestibular
Stimulation in Larval Zebrafish. *Current Biology* 28, 3723-3735.e6 (2018).

15.   Favre-Bulle, I. A., Stilgoe, A. B., Rubinsztein-Dunlop, H. & Scott, E. K. Optical
trapping of otoliths drives vestibular behaviours in larval zebrafish. *Nat Commun*
8, 630 (2017).

16.   Thiele, T. R., Donovan, J. C. & Baier, H. Descending Control of Swim Posture by
a Midbrain Nucleus in Zebrafish. *Neuron* 83, 679–691 (2014).

17.   Thorsen, D. H. & Hale, M. E. Development of zebrafish (Danio rerio) pectoral
fin musculature. *J Morphol* 266, 241–255 (2005).

18.   Ma, X. & Xu, X. A Swimming-based Assay to Determine the Exercise Capacity of
Adult Zebrafish Cardiomyopathy Models. *Bio Protoc* 11, (2021).

19.   McMenamin, S. K. & Parichy, D. M. Metamorphosis in Teleosts. in *Current*
Topics in Developmental Biology vol. 103 127–165 (Academic Press Inc., 2013).

20.   McHenry, M. J. & Lauder, G. V. Ontogeny of form and function: Locomotor
morphology and drag in zebrafish (Danio rerio). *J Morphol* 267, 1099–1109
(2006).

21.   Burris, B., Jensen, N. & Mokalled, M. H. Assessment of Swim Endurance and
Swim Behavior in Adult Zebrafish. *Journal of Visualized Experiments* (2021)
doi:10.3791/63240.

22.   Mauguit, Q., Olivier, D., Vandewalle, N. & Vandewalle, P. Ontogeny of

swimming movements in bronze corydoras (Corydoras aeneus). *Can J Zool* 88,
378–389 (2010).

23.   Gibb, A. C., Swanson, B. O., Wesp, H., Landels, C. & Liu, C. Development of the
755   Escape Response in Teleost Fishes: Do Ontogenetic Changes Enable Improved
756   Performance? *Physiological and Biochemical Zoology* 79, 7–19 (2006).

757   24.   White, R. M. *et al. Transparent Adult Zebrafish as a Tool for in Vivo*
758   *Transplantation Analysis.*

759   25.   Favre-Bulle, I. A., Vanwalleghem, G., Taylor, M. A., Rubinsztein-Dunlop, H. &
760   Scott, E. K. Cellular-Resolution Imaging of Vestibular Processing across the
761   Larval Zebrafish Brain. *Current Biology* 28, 3711-3722.e3 (2018).

762   26.   Bianco, I. H. *et al.* The tangential nucleus controls a gravito-inertial vestibulo-
763   ocular reflex. *Current Biology* 22, 1285–1295 (2012).

764   27.   Straka, H. & Baker, R. Vestibular blueprint in early vertebrates. *Front Neural*
765   *Circuits* 7, (2013).

766   28.   SUWA, H., GILLAND, E. & BAKER, R. Otolith Ocular Reflex Function of the
767   Tangential Nucleus in Teleost Fish. *Ann N Y Acad Sci* 871, 1–14 (1999).

768   29.   K. Lee, R. K. & Eaton, R. C. Identifiable reticulospinal neurons of the adult
769   zebrafish, *Brachydanio rerio. Journal of Comparative Neurology* 304, 34–52
770   (1991).

771   30.   Stickland, N. C. Growth and development of muscle fibres in the rainbow trout
772   (Salmo gairdneri). *J Anat* 137 (Pt 2), 323–33 (1983).

773   31.   van Meer, N. M. M. E., van Leeuwen, J. L., Schipper, H. & Lankheet, M. J. Axial
774   muscle‐fibre orientations in larval zebrafish. *J Anat* 10, 261–267 (2024).

775   32.   Jayne, B. C. & Lauder, G. V. How swimming fish use slow and fast muscle fibers:
776   implications for models of vertebrate muscle recruitment. *Journal of Comparative*
777   *Physiology A* 175, (1994).

778   33.   Nowroozi, B. N. & Brainerd, E. L. Regional variation in the mechanical
779   properties of the vertebral column during lateral bending in Morone saxatilis. *J R*

780   *Soc Interface* 9, 2667–2679 (2012).

34.   TSUKAMOTO, K. Swimming motion of fishes : Their adaptation to water.
Hikaku seiri seikagaku(Comparative Physiology and Biochemistry) 10, 249–262
(1993).

35.   Hang, H., Heydari, S., Costello, J. H. & Kanso, E. Active tail flexion in concert
with passive hydrodynamic forces improves swimming speed and efficiency. *J*
Fluid Mech 932, (2022).

36.   Martel, G. & Dill, L. M. Influence of Movement by Coho Salmon
(*Oncorhynchus kisutch)* Parr on Their Detection by Common Mergansers
(*Mergus merganser). Ethology* 99, 139–149 (1995).

37.   Tang, Z.-H., Huang, Q., Wu, H., Kuang, L. & Fu, S.-J. The behavioral response
of prey fish to predators: the role of predator size. *PeerJ* 5, e3222 (2017).

38.   SP, G. & DJ, E. Resolving shifting patterns of muscle energy use in swimming
fish. *PLoS One* 9, (2014).

39.   McCune, A. R. & Carlson, R. L. Twenty ways to lose your bladder: common
natural mutants in zebrafish and widespread convergence of swim bladder loss
among teleost fishes. *Evol Dev* 6, 246–259 (2004).

40.   Maina, J. N. & Maloiy, G. M. O. The morphometry of the lung of the African
lungfish (*Protopterus aethiopicus*) : its structural-functional correlations. *Proc R*
Soc Lond B Biol Sci 224, 399–420 (1985).

41.   Baldridge, H. D. Sinking Factors and Average Densities of Florida Sharks as
Functions of Liver Buoyancy. *Copeia* 1970, 744 (1970).

42.   Drucker, E. G. & Lauder, G. V. Function of pectoral fins in rainbow trout:
behavioral repertoire and hydrodynamic forces. *Journal of Experimental Biology*
206, 813–826 (2003).

43.   Hove, J. R., O'Bryan, L. M., Gordon, M. S., Webb, P. W. & Weihs, D. Boxfishes
(Teleostei: Ostraciidae) as a model system for fishes swimming with many fins:
kinematics. *Journal of Experimental Biology* 204, 1459–1471 (2001).

44.   Hale, M. E. Developmental Change in the Function of Movement Systems:
809   Transition of the Pectoral Fins between Respiratory and Locomotor Roles in
810   Zebrafish. *Integr Comp Biol* 54, 238–249 (2014).

811   45.   Westerfield, M. The Zebrafish Book. A Guide for the Laboratory Use of
812   Zebrafish (Danio rerio), 5th Edition. *University of Oregon Press, Eugene (Book)*
813   (2007).

Figure 1

A

Dorsal side camera

Roll-tilt stimulus

Frontal side camera

B

Intact fish

Frontal view

0.0 s Δ1.1° 2.5 s Δ10.4° 6.0 s Δ-2.5°

Dorsal view

Δ0.3° Δ21.6° Δ4.5°

C

ΔHead roll angle (°)

ΔBody flexion angle (°)

Roll stimulus (°)

Time (s)

Figure 2

A

Roll-tilt stimulus

Dorsal side camera

Fish holder

B

Rostral body-restrained

-2.0 s   6.0 s   12.0 s

Δ0.1°   Δ15.1°   Δ-3.3°

C

ΔBody flexion angle (°)

-10

Roll stimulus (°)

-2 0 2 4 6 8 10 12

Time (s)

Figure 3

	Upright	Roll tilted	Swim bladder-deflated
Frontal view			
	Buoyancy / Gravity / Swim bladder	Moment of force	Moment of force
Dorsal view	Swim bladder / COV / COM	Body flexion	Body flexion

Figure 4

Figure 5

A

Chemically fixed fish specimen

Dorsal view

θ1

Flexed body shape

Released in water

Forceps

Horizontal line

θ2

Front camera

B

	-0.1 s	0.0 s	0.1 s	0.2 s	0.3 s
Before swim bladder-deflation	21.6°	20.7°	11.1°	-20.7°	-53.6°
After swim bladder-deflation	22.1°	23.2°	19.3°	23.6°	29.9°

C

Figure S1

A

Wild type

0.0 s   2.5 s   5.0 s

Δ0.0°   Δ8.9°   Δ2.1°

Δ-0.1°   Δ27.9°   Δ5.7°

B

ΔHead roll angle (°)

ΔBody flexion angle (°)

Roll stimulus (°)

Time (s)

C

ΔBody flexion angle (°)

Roll stimulus (°)

Time (s)

D

Swim bladder deflation

Before

Syringe needle

Swim bladder

After