Start Small: A Model for Tissue-wide Planar Cell Polarity without Morphogens

Abhisha Thayambath[1,2] and Julio M Belmonte[1,2]

[1]Department of Physics, North Carolina State University, Raleigh, North Carolina, USA
[2] Quantitative and Computational Developmental Biology Cluster, North Carolina State University, Raleigh, North Carolina, USA

Corresponding author: jbelmon2@ncsu.edu

Abstract

Planar cell polarity (PCP) is an important patterning mechanism in both vertebrates and invertebrates by which cells coherently polarize along the apical surface of the epithelium. This patterning mechanism acts upstream of many developmental processes, such as oriented growth, division, cell movements and orientation of body hairs. While various models have been proposed to explain PCP patterning, all rely on persistent global cues/gradients to obtain global orientation of large tissues. However, recent experimental work has shown that this process can happen independently of such global cues, challenging the current paradigm. In this work, we developed a new model for PCP using the Cellular Potts modelling framework to investigate the conditions under which global tissue orientation can be achieved without a tissue spanning morphogen. We found that a combination of a local boundary signal, a small initial tissue size and uniform proliferation can effectively establish long-range polarity without the need for global cues. We also investigated the impact of cell division planes and growth rates on final patterning. Finally, we compared the cell-autonomous and cell non-autonomous versions of our PCP model, as found in flies and mice, and found that the latter offers more robust patterning outcomes in the absence of gradients.

Author Summary

Cells must coordinate the spatial organization of certain proteins inside them to produce aligned structures like body hairs and cilia across large distances in various organisms. This polarity of cells is also important to provide directionality for several key events during animal development. It was widely assumed that a graded signal acting across the tissue provides the global cue for the cells to attain this long-range alignment. However, the identity and existence of this global cue have recently been challenged by new experimental findings. In this work we developed a new computational model showing how coordinated cell polarity can emerge from simple, local rules in the absence of a global tissue-spanning signal. Our model proposes a mechanism where local signals polarize a small initial group of cells, and their subsequent proliferation extend this initial alignment across large scales. We did extensive test of our model under different conditions and show that this proposed mechanism provides a reliable recipe for the generation of large-scale tissue polarity without the need for global signals.

Introduction

Cell polarity is the intrinsic asymmetry in the shape, structure, or spatial organization of cellular components that enables cell and tissues to perform specialized functions. **Planar Cell Polarity (PCP)** refers to the coordinated orientation of cells within the plane of an epithelial sheet and is a key mechanism for establishing long-range tissue organization along the apical surface. The establishment of

a coherent proximal-distal axis is crucial for the proper function of many cellular and developmental processes downstream of PCP such as the generation of body hairs (sensory hairs in mammals [1][2], wing hair in *Drosophila* [3] etc.) and the orientation of tissue processes such as convergent-extension [4], gut elongation [5], neural tube closure [6][7], as well as in processes in mesenchymal tissues such as gastrulation, migration and cell differentiation [8] [9].

While the mechanisms of local planar cell polarity through direct cell-cell communication are well understood experimentally and computationally, the long-range order and direction of tissue orientation remains an open question. Multiple computational models suggest the need of some global cue for the establishment of tissue wide PCP alignment in the presence of noise. Yet, experimentally, the list of possible candidates for such a global cue continues to diminish. Early in 2000's, Peter Lawrence and colleagues showed that PCP is independent of the DPP, Notch, EGF, and FGF signalling pathways [10]. The Ds and Fj proteins, which are part of the Fat–Dachsous PCP pathway, are expressed in opposing gradients and could act together as the global cue. However, uniform expression of those components does not alter the long-range PCP alignment [11]. And more recent studies show that long-range PCP polarity in the *Drosophila* wing and notum can be established independent of Wnt gradients [12] [13]. All these negative results suggest that an alternative mechanism, independent of morphogen gradients, may be required to establish global PCP orientation.

PCP pathways

The mechanisms underlying planar cell polarity have been extensively studied in *Drosophila* where PCP controls the orientation of hairs on the wing and abdomen and the organization of ommatidia in the eye [14]. Two major pathways establish PCP in *Drosophila* wing- the **core PCP pathway** and the **Fat–Dachsous pathway**. Both pathways rely on the asymmetric localization of their associated proteins on the opposing sides of each cell. This asymmetric localization is guided in part by direct communication between neighbouring cells which determines local alignment, and by a putative global cue which determines the global direction of tissue polarization.

The core pathway is composed of the cytoplasmic proteins Dishevelled(Dsh), Diego(Dg) and Prickle(Pk), transmembrane proteins Frizzled(Fz) and Van Gogh/Strabismus(Van/Stbm) and an atypical cadherin Flamingo(Fmi). Fmi is found on both opposing ends of each cell, and facilitates homophilic cell adhesion between adjacent cells [15] [16]. The communication between neighbouring cells occurs through the interaction of Fz–Fmi in one cell with Vang–Fmi in the adjacent cell. Inside a cell, Dsh binds with Fz [17], whereas Pk interacts with Van [18]. Pk also binds to Dsh preventing it from localizing to the proximal side [19] by destabilizing Fz binding [20]. Diego competes with Pk for Dsh binding favouring Dsh accumulation on the distal side and thereby enhancing Fz localization at the distal membrane [21]. As a result of this competitive inhibition, the complexes Van-Pk and Fz-Dsh-Dg localize to the opposing ends of each cell, forming the cells' proximal and distal domains, respectively [22][23][24] (see [25] and [26] for review).

The Ft-Ds pathway is composed of the protocadherins Ft and Ds which form heterophilic complexes between cells [27]. This is regulated by the transmembrane kinase Four-Jointed (Fj). In the developing wing, the asymmetric localization of Ft to the proximal side and Ds to the distal sides of the cells is driven by the complementary gradients of Ds and Fj [28][29]. It is still unclear whether the Ft-Ds pathway operates independently or together with the core pathway to coordinate polarity across the tissue [30][31].

Morphogens and global alignment

Experiments and mathematical models suggest that the spatial segregation of polarity complexes inside each cell and their preferential affinity between neighbouring cells is effective in coordinating polarity across small groups of cells. An initial uniform distribution of these core proteins spatially segregates over time, resulting in a polarized distribution (Fig1A). While the reinforcement of the initial asymmetry by feedback interactions is sufficient for the local coordination of cell polarities, the identity of the biasing cue that could determine the global alignment direction and ensure large-scale coordination remains poorly understood. It has been hypothesized that the global alignment of core PCP proteins arises from a spatial gradient of a biasing cue acting across the entire tissue. Lawrence et al. have shown that if the spatial gradient of a secreted molecule acts as the global cue, then it is not part of the DPP, Notch, EGF, or FGF signalling pathways [10]. The Ft-Ds-Fj system is thought to be a potential candidate that could act as an upstream global cue for the core proteins, given that Ds and Fj are expressed in

complementary gradients in opposing directions. However, even in conditions of uniform expression of these components, polarity can still be effectively established, suggesting that the gradients of Ft-Ds-Fj are not strictly necessary [11] (see [32] for a comprehensive review).

Figure 1: **Sub-cellular Potts Model for Planar Cell Polarity: (A)** Schematic of an epithelial cell arrangement with PCP protein distributions. Initially, the PCP proteins are uniformly distributed within each cell. Over time, Fz, Dsh, and Dgo (magenta) localize to the proximal/anterior edge, while Vang and Pk (green) segregate to the distal/posterior side. **(B)** Schematic of two epithelial cells with cluster ids 1 (orange) and 2 (blue) respectively, along with the respective domain ids. The cluster id indicates the association of each domain with a specific biological cell. **(C)** Model Overview: Each epithelial cell is subdivided into four compartments—proximal, distal, lateral, and cytoplasm. **Seed** cells are placed randomly within the lattice, which grow and rearrange their shapes to efficiently pack in the space. PCP domains are introduced with random initial distributions. The proximal and distal domains spatially segregate into opposing ends of each cell over time. Blue box highlights the evolution of a single cell during this process. **(D)** The vector field representation of PCP orientations shows the transition from a homogeneous distribution ($\phi \approx 0$) of global polarization to complete alignment ($\phi \approx 1$). (Panel A adapted from [25])

Wnt ligands, which bind to Fz receptors, are expressed as gradients in the developing Drosophila wing [33][34][35]. Together, they recruit and activate Dsh, which is also a highly conserved component of the Wnt signalling pathway, which in turn propagates the signal internally to the cell [36]. Since Fz and Dsh belong to the core pathway, Wnt was long considered to be the global cue that sets the global alignment and long-range orientation of PCP in *Drosophila* wing and other PCP systems. Although some studies have suggested the role of Wnt ligands (Wg and Wnt4) in global PCP establishment [37],

two independent recent studies show that Wnt ligands are not required to align core PCP in *Drosophila* [12] [13]. Ewen-Campen et al. conducted loss of function studies targeting different combinations of Wg and other Wnt genes (wnt2, wnt4, wnt5, wnt6, wntD, wnt10) in *Drosophila* wing and notum and found that loss of individual or combinations of Wnt ligands that they examined do not disrupt PCP alignment [12]. In another study published in the same year, Yu et al. knocked out the diffusible Wnts DWnt2, Dwnt4, DWnt6, and DWnt10 and found no disruptions in wing hair orientation [13]. Together, these results show that Wnt ligands are not required for global PCP establishment, and other mechanisms must be responsible for the long-range orientation of PCP.

Computational models of PCP

There are several models that explain the mechanisms behind planar cell polarity- ranging from explicit molecular signalling models to phenomenological/reduced models. The *explicit signalling models* consider the detailed interactions between the proteins in core and Ft-Ds pathways. For instance, both [38] and [39] describe cell polarity based on the asymmetric distribution of intercellular protein complexes with cells and their associated PCP components modelled by reaction-diffusion equations. Both models use a global cue (which could arise from a directional signal within each cell or from a tissue wide ligand gradient) to introduce a small bias in each cell which are enhanced by local feedback interactions between the core proteins. Fisher and colleagues [40] have shown that both these models require a persistent global cue to generate a robust polarity that is insensitive to noise.

The *phenomenological models* simplify the PCP mechanism by reducing the complexity of the spatial protein interactions in order to extract the fundamental interactions leading to planar polarization. Burak et al. [41] proposed a semi-phenomenological model incorporating mutual exclusion of Fz and Vang within cells and intercellular interactions, showing that a weak early global cue can establish long-range polarity. Wang et al. [42] used a cellular automaton model where cells align their polarity with the average of their neighbours, finding that stronger global cues accelerate alignment. These models used cells with a fixed shape in a fixed lattice. The work by Zhu and colleagues [43] was the first to use cells with varying shapes and dynamic neighbourhoods, revealing that directional cues are critical for PCP, while hexagonal packing has a minimal effect on patterning. Schamberg et al. [44] developed a feedback and diffusion-based model that generates polarization in a row of cells, either through a weak initial global polarization across all cells or via a travelling wave originating from a single cell or boundary. Abley et al. [45] show that individual cells can polarize through random fluctuations and spread polarity via cell-cell coupling and boundary organizers. This model also used gradients to improve the tissue-level polarity. Hazelwood and Hancock [46] developed a functional model using global cues to study the molecular functions of core proteins Pk and Dsh. Fisher et al. [47] explored how intercellular protein complexes interpret tissue-wide gradients when establishing polarity. More recently, Singh et al. [48] developed a minimal model centered on Ft-Ds heterodimer formation, demonstrating that while PCP can arise without a global gradient, such a gradient is essential for robustness to noise and stabilization of polarity patterns.

All these models, explicit and phenomenological, used a global cue - whether as a persistent directional signal within each cell, a tissue-wide ligand gradient, or a weak orienting signal - to establish and maintain polarity across the tissue. However, given the recent experimental results by [12] [13] which found no evidence for the necessity of a Wnt gradient for the establishment and maintenance of PCP in the *Drosophila* wing (the primary model system for PCP), the question arises as whether there are morphogen-independent mechanisms that can explain robust establishment of long-range tissue orientation.

A model for long-range PCP without morphogens

To address the question of how a tissue achieves large-scale PCP alignment without morphogens, we developed a phenomenological model for PCP based on the Cellular Potts (Glazier-Graner-Hogeweg) modelling framework in which we represented the asymmetrically distributed PCP proteins as distinct compartments inside cells [49] [50]. We varied the contact/interface properties of these compartments to model the mutual exclusion of these proteins inside cells and differential adhesion across cells. As discussed by Lavalou and Lecuit, adhesion motifs in the transmembrane PCP proteins play an essential role in forming the inter-cellular protein complexes in such a way that when one protein is expressed at a given cell boundary, its corresponding counterpart is also recruited to the interfacial boundary of the adjacent cell [26]. In our model, the compartments representing the proximal and distal PCP proteins at

adjacent cell boundaries interact with a higher adhesion strength, leading to their preferential binding. Within each cell, we can choose the properties of the cell compartments to model either cell autonomous polarization, where a single cell can segregate its PCP domains by itself, as seen in flies [51]; or cell non-autonomous polarization[52], where PCP domains require interaction with neighbouring cells to polarize, as seen in mice. Through these interactions alone, we show that cells can locally align their polarity in the absence of a directional signal. However, this mechanism alone is not sufficient to establish a long-range direction of orientation and swirling patterns arise as we increase tissue size, even in the presence of a boundary-orienting signal that polarizes the first row of cells. To explore how we can establish and maintain global alignment along a particular direction across large numbers of cells, we examine and test a mechanism first suggested by Sagner et al. [53], where a small-scale alignment generated by the boundary signal can propagate as the tissue grows. Our results show that uniform cell proliferation coupled with a boundary signal is a robust mechanism for large-scale polarity establishment in the absence of a tissue-wide morphogen gradient.

Model Overview

We developed our model within the Cellular Potts Modelling framework, using the CompuCell3D (CC3D) simulation software [54]. This modelling framework represents cells as extended objects formed by a collection of lattice sites on a square grid in 2D or 3D. Each cell consists of either a single domain of lattice sites (defined by a domain ID and a domain type) or multiple domains representing specific sub-cellular compartments (e.g., nucleus and cytoplasm) or abstract regions inside cells (e.g., apical and basal regions). In our model, we represent each biological cell as a 2D object subdivided into four sub-cellular compartments/domains: (i) proximal, (ii) distal, (iii) lateral, and (iv) cytoplasm. Together they represent the spatial heterogeneity in the distribution of various PCP proteins as well as the cytosolic constituents of the cell as viewed from their apical side (Fig1B'). Since each biological cell in our model is made up of the four domains, we assign a cluster ID to each domain to clearly indicate its association with a specific biological cell (Fig1B). Proximal and distal domains represent the proteins that localize to the proximal (Fz, Dsh, Dgo) and distal (Van/Stbm, Pk) regions of the cells whereas the lateral domain represents the regions of the cell membrane not populated by the core PCP proteins.

An effective energy equation H_{total} defines the properties of each of these domains, such as size, aspect ratios, adhesion preferences, and interactions with other domains (Eq. 1). Each of these properties is governed by an individual term in the overall effective energy equation: a volume constraint H_V helps to maintain the size and compressibility of cell domains and a contact/adhesion energy (H_C) specifies the relative strength of contact/adhesion between domains from the same and/or different cells.

$$H_{total} = H_V + H_C \tag{1}$$

The system evolves by minimizing the effective energy H_{total} by a series of random lattice-site copy attempts where we randomly select a lattice site i and a neighbouring lattice site j within the 3rd neighbour order of i. If the two lattice sites belong to different cells or cell domains, we evaluate the difference in energy (ΔH) if the ID of site i is copied over site j. Lattice site copy attempts that decrease the effective energy of the system is accepted, whereas those that increase the effective energy of the system are accepted with a probability:

$$P = \exp\left(-\frac{\Delta H}{T}\right) \tag{2}$$

where T is a parameter that specifies the level of membrane fluctuations in the system. The unit of time of the simulations is the Monte-Carlo step (MCS), defined as N lattice-site copy attempts, where N is the number of lattice sites in the grid.

The volume constraint (H_V) penalizes the deviations of the cell domain's volume (V) - defined as the number of lattice sites occupied by the cell domain - from its target volume (V_t):

$$H_V = \lambda_V (V - V_t)^2 \tag{3}$$

where λ_V represents the strength of the volume constraint and can be interpreted as the inverse compressibility of the cell domain. The parameters used for the implementation of all these constraints are given in Supplementary Table 1.

The contact energy (H_C) is modelled as:

$$H_C = \sum_i \sum_{j \neq i} J(\sigma_i, \sigma_j) \tag{4}$$

where the second sum is over the 4th order neighbours of site i; σ_i and σ_j are the types of cell compartments occupying sites i and j, respectively; and J is a matrix containing the contact/adhesion energy between compartment types belonging to the same cell (denoted as j_{xy}) and different cells (denoted as J_{xy}). For the sake of simplicity, we assume no variability in cells' contact energies between cells or over time.

The value of the J matrix is different depending on whether the domains belong to the same cell (same cluster ID) or to different cells (different cluster IDs). When belonging to the same cell, all contact energies of pairwise combinations of domains are set at low values, while those between the proximal and distal domains are set at high values to facilitate the sorting of the two domains inside the cells (Table 1). Even in the absence of a neighbour or an interacting environment, the proximal and distal domains segregate to opposing ends inside a cell in relatively faster time scales (Fig S1A), and hence individual cells are capable of establishing PCP asymmetry autonomously in our model. This is consistent with the findings of [51] where individual cells lacking the core protein Fmi - and thus are unable to communicate with neighbouring cells - still generate PCP asymmetry.

Table 1: *Internal* contact energies (j) between PCP domains

Cell Type	Proximal	Distal	Lateral	Cyto
Proximal	-	30	1	1
Distal	-	-	1	1
Lateral	-	-	-	1
Cyto	-	-	-	-

When belonging to different cells, all contact energies of pair-wise combinations of domains were set to higher values. We defined the external contact energies in such a way that proximal and distal domains of neighbouring cells would preferentially bind together to incorporate the observed inter-cellular coupling (Table 2). We determined the exact values for each contact energy term by extensive parameter scans aimed at optimizing cell autonomous polarization for small systems (Fig S2A, S2B and S2C). The parameter scans show that our results are valid for a wide range of parameter values.

Table 2: *External* contact energies (J) between PCP domains and Medium

Cell Type	Proximal	Distal	Lateral	Cyto	Medium
Proximal	12	7	12	30	10
Distal	-	12	12	30	10
Lateral	-	-	14	30	14
Cyto	-	-	-	50	30
Medium					0

We also used a temporary cell type, called **seed** cells, to initialize the simulations by randomly placing them in the lattice and allowing them to grow into arbitrary shapes until they reach a volume that completely fills the space. During this growth phase, cells remodel their shapes and rearrange their neighbours to achieve efficient packing within the lattice. This mechanism is consistent with the observation that epithelial cells are usually irregularly packed, as in the case of the *Drosophila* wing during larval and pre-pupal stages, where repacking into a hexagonal array occurs after polarization and tissue shear [55]. Once the cells reach their target sizes and relax their boundaries, we establish PCP domains within each cell by assigning them random spatial distributions within each cell (50% of cell volume is composed of cytoplasmic domains, 15% is composed of proximal domains and distal domains each and the remaining 20% is composed of the lateral domains). This random assignment is done to avoid imposing any pre-pattern on the tissue. The total volume of each domain inside each cell remains conserved throughout the simulation, unless there is cell growth.

Starting from a homogeneous distribution, differential contact preferences within individual cell domains and between neighbouring cells lead to the spatial segregation of PCP proteins inside each cell (Fig1C, 1D). We observe that this intercellular coupling through stronger adhesion between proximal

and distal domains across cell boundaries is sufficient for local coordination of polarity in the absence of a directional cue. The proximal and distal domains segregate to opposite ends of each cell, with lateral compartment symmetrically localizing between them. This rearrangement occurs naturally, as the internal contact energies are set to prevent the direct contact between proximal and distal domains with the lateral compartment acting as a buffer between them. The system globally aligns along a random direction on a relatively short timescale of $30,000$ MCS for a system of 4×4 cells (Fig S1B).

When implementing cell growth and division, we update the target volumes of all cell domains such that the proportions of Proximal, Distal and Lateral domains remain the same with respect to the perimeter of the cell. The rate of cell growth is a varied parameter in our model. Once the cell doubles volume, a division plane is chosen to be either along the major or minor axis of the cell. After division, the two daughter cells inherit equal amounts of all domains. These domains, however, are randomly mixed inside the new cells. Spatial segregation of the internal domains will happen according to the internal and external contact energies and the local neighbourhood of the new cells. This is done to avoid any bias and to reflect the biology of dividing cells, that loose their polarity information upon division [56]. After cells divide they experience a refractory time t_r before restarting their growth.

In order to quantify the degree of tissue polarization, we adapted the order parameter used by the flocking model community to describe the collective movement of self-propelled particles [57][58]. Here, we describe every cell in the lattice using a polarity vector ($\vec{\nu}(\sigma)$), defined as the normalized distance between the center of mass of **proximal** and **distal** domains:

$$\vec{\nu}(\sigma) = \frac{\vec{r}(\sigma_{prox}) - \vec{r}(\sigma_{dist})}{|\vec{r}(\sigma_{prox}) - \vec{r}(\sigma_{dist})|} \tag{5}$$

where $\vec{r}(\sigma_{\text{prox}})$ and $\vec{r}(\sigma_{\text{dist}})$ are the positions of the center of mass of the **proximal** and **distal** domains of cell σ, respectively. From this vector, we define the scalar order parameter ϕ that describes the degree of global tissue polarization as

$$\phi = \frac{1}{N} \Big| \sum_{\sigma} \vec{\nu}(\sigma) \Big| \tag{6}$$

where N is the total number of cells in the lattice and $\phi \in [0, 1]$, where 0 means no alignment, and 1 means perfect alignment of PCP vectors (Figure 1D).

We also quantified the local polarity order $\phi(r)$ by measuring the degree of polarization as a function of either the distance r between cells or as a function of the distance r from a signaling boundary. The first version of the metric is calculated as:

$$\phi_R(r) = \frac{1}{N} \sum_{\sigma} \frac{1}{N_{d(\sigma,\sigma')<r}} \Big| \sum_{\sigma'}^{d(\sigma,\sigma')<r} \vec{\nu}(\sigma') \Big| \tag{7}$$

Here, $\vec{\nu}(\sigma')$ is the polarity vector of cell σ', and $d(\sigma, \sigma')$ denotes the distance between cells σ and σ'. For each reference cell σ, the local average alignment is computed over all neighbouring cells σ' within a distance r from it and normalized by their number $N_{d(\sigma,\sigma')<r}$. The outer average over all cells yields $\phi_B(r)$, which describes how local polarity decay with distance.

To quantify the local polarity order as a function of distance from the left boundary, we divided the tissue along the x-axis into bins of width equal to one cell diameter. For each bin centered at distance r from the boundary, we calculated the mean polarity of all cells whose centers are less than r. The local polarity order as a function of the distance from the boundary is then calculated as

$$\phi_B(r) = \frac{1}{N_{d(b,\sigma)<r}} \Big| \sum_{\sigma}^{d(b,\sigma)<r} \vec{\nu}(\sigma) \Big| \tag{8}$$

where $d(b, \sigma)$ is the distance between boundary and cell σ and the sum is done for all cells that lie within a distance r from the left boundary.

Results

Based on the modelling framework described in the previous section, we began by studying the behaviour of the model under various scenarios and checking if we can reproduce the previously known experimental

results. We first examined the time evolution of the degree of global polarization across different boundary conditions. Then, we tried to reproduce the dominering non-autonomy effects in polarity alignment observed in *Vang* and *Fz* loss of function mutants. After this, we studied the impact of increasing system size under different boundary conditions and investigated the mechanisms by which cells can establish long-range polarity. All results presented are averages over 100 independent simulations.

Time evolution of global polarization under different boundary conditions

We first characterized our model by investigating how it behaves under different conditions. We examined the evolution of global polarization over time under different boundary conditions for a system of 8×8 cells (Fig 2A, 2B). We initialized the system by randomly placing seed cells in the simulation space and letting them grow and relax their boundaries for 10^4 MCS. After that we added the PCP domains to the cells in a random manner (Fig 1C) and studied the evolution of global polarization defined by the scalar order parameter ϕ (Fig 1D). As time increases from 10^4 MCS, global polarization increases and then stabilizes after 5×10^5 MCS for both the boundary conditions. (Fig 2D).

To circumvent the effects that an external boundary might have on tissue orientation, we first simulated our model with periodic boundary conditions to mimic a continuous and unbounded tissue (Fig2A, first panel). These boundary conditions help eliminate the edge effects present in large biological tissues by treating the tissue as if it wraps around on itself. Starting from an initial state where cell polarization vectors are randomly oriented (Fig1D, first panel and Fig2C, first panel, green), the system self-organizes towards a globally aligned state based only on the local contact preferences between the domains within the cells and between neighbouring cells without any imposed external cues. Over time, the global polarization increases and stabilizes at approximately 85% (Fig2D, red). The final tissue angles of polarization are predominantly distributed along four distinct directions —right, left, up, and down—corresponding to the symmetry axes of the lattice (Fig2A, second panel). Fig2C (second panel) illustrates one such outcome in a small 8×8 system, where the initially

random orientations of cell polarity vectors (Fig2C, first panel) evolves toward the left, representing one of the possible final states under periodic conditions.

We also investigated the effect of boundary-orienting signals on tissue polarization. We added a fixed column of **distal** domain type at the left boundary of the system (Fig2B), which acts as a directional cue that orients cells directly in contact with it towards the right (Fig2C third panel, blue). The open boundary on the front does not have a preference for proximal or distal domains. The addition of this boundary signal biases the global alignment direction to be predominantly directed along the proximal-distal direction, as it can be seen in the radial histogram of individual cell angle distributions (Fig2C, third panel, blue). We also found that the presence of this boundary slightly speeds and enhances the global polarization of the tissue, as the evolution of the order parameter ϕ is higher than the periodic boundary configuration for nearly all time steps in the simulation (Fig2D). From these results, we conclude that the presence of the boundary signal in our model successfully biases small systems to polarize along a preferred direction, as observed in other models for PCP [41][42].

Loss of function mutants produce domineering non-autonomy

Clones of cells lacking *Fz* function disrupt polarity in the wild-type cells distal to the mutant clones, whereas clones lacking *Vang* function disrupt polarity in wild type cells on the proximal side of the mutant clone [38]. This effect is called "domineering non-autonomy" because the mutant cells affect the polarity of the neighbouring wild-type cells. To test whether our model reproduces this result, we created a 8×12 simulation space with a left boundary signal to induce global polarization to the right, and introduced a patch of cells that contain only proximal or distal surface domains, mimicking the loss of function for *Fz* and *Vang* activity, respectively. We found that the polarity vectors of cells distal to the *Fz*-like mutant patch reoriented facing the mutant clone (Fig2E). Conversely, we found that the polarity vectors in wild-type cells proximal to the *Vang*-like mutant patch pointed away from the clone (Fig2E'). Hence, our model successfully reproduces the domineering non-autonomous effects of *Fz* and *Vang* mutants, as expected given that PCP alignment mechanism in our model is determined solely by local effects.

Effect of system size on global and local polarization

After establishing the basic characteristics of the model and verifying that it is able to reproduce key results from experimental observations and other published models, we investigated the effect of system

Figure 2: **Time Evolution of Global Polarization under different Boundary Conditions and Domineering Non-autonomy:** **(A)** Polarization outcomes under periodic boundary conditions applied along both the x and y directions. The system exhibits four distinct final states, with the mean tissue polarization vector aligning along one of the four directions: right, left, up, or down. **(B)** Polarization under a directional cue applied at the left boundary, with periodic boundary conditions maintained along the y axis. This configuration introduces a bias to the right and sets a definitive proximal-distal axis to the right. **(C)** Distribution of individual cell polarization angles in a system comprising 8×8 cells. The first panel (green) shows the initial distribution of cell polarization vectors. The second panel (red) depicts the final distribution under periodic boundary conditions for a system that polarizes towards the left direction, similar to the output highlighted in red in panel **(A)**. The third panel (blue) shows the final distribution under the left-boundary orienting signal, with polarization vectors highly aligned along the proximal-to-distal axis. **(D)** Time evolution of the global polarization, quantified by the scalar order parameter ϕ, under periodic (red circles) and left-boundary (blue triangles) conditions. In both periodic and left boundary configurations, global polarization increases with increasing time in a system of 8×8 cells examined for 10^7 MCS. **(E)** Domineering non-autonomous behaviour of cells in loss-of-function mutants for *Fz*. In *Fz* loss-of-function mutants (magenta), polarity vectors distal to the patch point towards the mutant clone. **(E')** Domineering non-autonomous behaviour of cells in loss-of-function mutants for *Vang*. In *Vang* loss-of-function mutants (green), polarity vectors proximal to the patch point away from the mutant clone.

size on global and local polarization. We increased the number of cells across different lattice sizes and ran simulations for 10^6 Monte Carlo Steps to allow sufficient time for the system to polarize. As in the previous section, we considered two boundary conditions: periodic boundaries and left boundary signal (Fig2A, B). We found that as the number of cells increases, global polarization decreases in both configurations (Fig3A, see Fig S1C for time series data). In the periodic boundary configuration with 30×30 cells, the distribution of the mean angle of tissue polarization is nearly uniform (Fig3B, first panel). However, for the 30×30 system with a left boundary signal (Fig3B, second panel) the mean angle does not exhibit the strong preference along the proximal-distal axis as observed in the 8×8 system (Fig2C, third panel). Instead, the alignment direction is more dispersed around proximal-distal axis (Fig3B, second panel). This suggests that the effect of the boundary orienting signal weakens with the distance from the boundary.

For the periodic boundary configuration, the representative simulation output for an 8×8 system shows that the final cell polarity vectors align along a specific direction (Fig3C, top panel) while the 30×30 system shows that cells align within small local domains and swirling patterns emerge (Fig3C, bottom panel). This clearly suggests that when system size increases, there is no long-range order across the entire system in the absence of a directional cue. For the configuration with the left boundary signal, 8×8 system shows cell polarity vectors aligned along proximal-distal axis (Fig3D, top panel). However, when system size is increased to 30×30, although the first few columns of cells remain well aligned along proximal-distal axis, global alignment is lost as we move further away from the left boundary (Fig3D, bottom panel).

We also examined how polarization changes as a function of distance using the two versions of the local polarity order parameter $\phi(r)$. Local order is high (over 80%) over small distances (radius < 5 cell diameters), but decreases steeply as the radius (measured in units of cell diameters) increases for the periodic boundary configuration independently of tissue size (Fig3E, top panel). This suggests that the distance over which the polarity alignment is maintained, the correlation length, is very small. In the left boundary signal configuration, cells close to the boundary are highly polarized with $\phi > 90\%$ for distances up to 5 cell diameters. The degree of polarization still declines with distance from the boundary, but the decrease is less steep compared to the periodic case (Fig3E, bottom panel). This indicates that the boundary signal helps maintain the cell polarity correlation higher and for longer distances as compared to periodic boundary configuration. However, it does not propagate tissue polarity effectively across large distances from the boundary. This reinforces the idea that additional mechanisms might be required for the coordination of polarity across a large tissue.

Cell proliferation is sufficient to establish long-range polarity

Our findings from the previous section suggest that establishing polarity across a large tissue requires additional mechanisms. A key question is whether this can be achieved without a graded biasing signal acting across the entire tissue. We hypothesize that the cells may first establish polarity in a small domain with the help of a margin/boundary-oriented signal and then expand this domain while conserving alignment. To test this, we start with a small patch of tissue that is already polarized with the help of the left boundary-orienting signal. We then allow the cells to proliferate by gradually increasing their volume. We let these cells grow slowly while maintaining the relative proportions of the proximal, distal, lateral, and cytoplasmic domains so that the cells maintain their polarity during this process. When a cell exceeds twice its volume, it divides along its minor axis. Following division, daughter cells inherit equal fractions of the cell domains from the parent cell but lose internal polarity as we redistribute these domains uniformly within each daughter cell (Fig4B). These cells re-polarize over time by spatially re-organizing their domains based on the polarity of the neighbouring cells. This mechanism is consistent with the findings from Devenport et al., where they describe how PCP components are equally inherited by daughter cells and how they re-establish their polarity in accordance with their neighbours in basal epidermal stem cells in mouse embryo [56].

We tested two different growth scenarios: growth at the distal end of the tissue (opposite to the boundary signal) and uniform tissue growth. We begin our simulations with a 10×4 system of cells that are already polarized. There is a periodic boundary condition along the vertical direction, a fixed boundary wall of **distal** domain type at the left side of the tissue, and an open space to the right that the tissue occupies as the cells grow and divide. The open boundary on the front has no preference between proximal and distal domains. We compare the results of the two scenarios to the polarization of tissues with fixed/pre-established sizes under the same boundary conditions that polarizes all at once. To allow for the comparison of growing tissues at different times, we varied the number of cell columns

Figure 3: **Effect of System Size on Polarization for Periodic Boundary and Left Boundary Signal Configurations (A)** Final average global polarization (ϕ) for different system sizes under periodic and left boundary signal configurations (average over 100 simulations). As system size increases, global polarization decreases for both boundary conditions. **(B)** Final tissue angle distribution of polarization for the 30×30 system under periodic (red) and left boundary (blue) conditions. For periodic boundaries, the polarization angles become nearly uniform, while for the left boundary signal, the alignment direction is more dispersed around the proximal-distal axis. **(C)** Final cell polarity vectors for the 8×8 (top panel) and 30×30 (bottom panel) system with periodic boundary conditions. In the smaller 8×8 system, all cells align in a single direction (upward in this case). For the 30×30 system, local alignment occurs within small domains, and swirling patterns emerge. **(D)** Final cell polarity vectors for the 8×8 (top panel) and 30×30 (bottom panel) system with left boundary signal configuration. For the smaller 8×8 system, all cells align along proximal-distal direction. For the 30×30 system, alignment is maintained in the first few columns but dissipates as cells move farther from the boundary. **(E)** Local polarization as a function of radial distance for periodic boundary configuration (top panel) and as a function of distance from the boundary for left boundary signal configuration (bottom). Local polarization decreases sharply with distance in the periodic boundary case, whereas the decrease in polarization for the left boundary signal configuration is less steep.

in the non-growing tissue.

As expected, for the simulations with fixed-sized tissues, the more cell columns we added, the smaller the global polarization, reaching $\phi = 0.3$ for the largest system size examined (Fig4E, red line). The decrease in ϕ is steeper with increasing number of cells when compared to the previous case with square shaped tissues (Fig2A first panel, 2B). In this case, each additional column of cells are farther away from the boundary signal, as the tissue has more cells along x-direction. This causes a steeper decline in tissue polarization as a function of distance from boundary than the square tissue case, where cells are added

on both directions (Fig4E, S3).

Figure 4: **Cell Proliferation-Based Mechanism for Maintaining Polarity.** **(A)** Schematic of the system used to study the effect of cell proliferation. 10 rows of cells are arranged in a box with periodic conditions along the vertical direction, a boundary signal on the left, and an open space on the right. **(B)** Cells grow to double their volume and then divide. After each division, daughter cells inherit the PCP domains equally, lose polarity, and redistribute these domains based on neighbouring cells' polarity. **(C)** Simulation showing cell growth and division at the distal/front boundary of a system. **(D)** Simulation showing uniform cell proliferation. Cells about to divide are indicated by an 'X' mark. **(E)** Comparison of global polarization ϕ for systems with and without cell proliferation as a function of the number of cells. For fixed tissue sizes, global polarization decreases with system size (red). Systems with cell proliferation on front/distal boundary stabilize at a higher ϕ, approximately 0.89 (blue). Uniform cell proliferation further enhances global polarization at a higher value 0.92 (green). **(F)** Distribution of mean angle of polarization for systems with proliferating cells, showing a shift towards $\pm 60°$ for front/distal boundary proliferation and proximal-distal alignment for uniform cell proliferation. **(G)** Comparison of final global polarization when number of cells in the system is 900 for varying growth rates for front/distal proliferation (blue) and uniform cell proliferation (green). For uniform cell proliferation, the refractory time between cell divisions is set to 10^4 MCS. **(H)** Simulation output showing polarity vector alignment for the configuration without cell proliferation. Domains of local alignment can be observed. **(I)** Simulation output showing polarity vector alignment for the proliferation on front configuration. **(J)** Simulation output showing preference for alignment along the proximal-distal axis for the uniform cell proliferation configuration.

Next, we tested whether pre-polarizing a small (10×4) tissue and allowing cells at the distal end to

grow and divide would improve polarization. As the number of cells increases, we observed a decrease in global polarization ϕ until the system size reaches 200 cells (Fig4E, S4A). After this, ϕ gradually recovers and stabilizes at approximately 0.9 as system size reaches 900 cells (Fig4E). This suggests that cell proliferation on the boundary may help maintain the initially established polarity. However, when we examine the mean angle distribution for these systems, we observe that the alignment is not strictly along the x-direction but instead shifts towards $\pm 60°$ (Fig4F). Simulation snapshots confirms that after a few rows away from the left boundary, the polarity vectors tend to point either upward or downward, explaining the initial decrease observed in global polarization ϕ (Fig4I, S5). This trend can be observed in the plot of local polarization as a function of distance from the boundary, where ϕ decreases with distance and stabilizes after approximately 20 columns of cells (Fig S3). The results presented are for growth rates of 0.001 lattice sites per MCS (or a 1.44×10^5 MCS volume doubling time). Results remain the same for slower growth rates, but polarization is gradually lost for faster growth rates, suggesting that cells need some time to adjust their polarity before the next round of division to maintain local tissue alignment (Fig S6). We found that slower growth rates help maintain the initially established polarity as the cells grow (Fig4G, S6).

Instead of restricting cell growth to just the front boundary, we allow all cells to grow, but at randomly assigned times. The time at which each cell begins to grow is decided by selecting a random number between 0 and refractory time t_r. Once a cell reaches twice its initial volume, it divides. The newly divided cells would start to grow again only after a random time t_{random} (selected between 0 and t_r) has passed. The growth rate is set to 0.001 lattice sites per MCS (or a 1.44×10^5 MCS volume doubling time). As a result, the time scale for polarity adjustment is much shorter than the growth and division time scale, giving the cell ample time to adjust its polarity between divisions. We find that the global polarization, ϕ, remains roughly constant at values greater than 0.9 as the number of cells in the system increases. Although there is a slight progressive drop in ϕ as the system size expands, ϕ stays above 0.9 up to a system size of 900 cells (Fig4E). A similar trend is observed for the local order, which also remains above 0.9 (Fig S3). Unlike the previous growth scenario, the mean angle distribution plot shows that the system stays preferentially aligned along the proximal-distal axis (Fig4G). The simulation output also shows that the polarity vectors tend to point along the proximal-distal direction (Fig4J). Thus, uniform cell proliferation appears to be an effective mechanism that robustly maintains the initially established polarity.

Cell autonomous vs cell non-autonomous polarization

The question of whether cells require contact with neighbours to establish polarization has been investigated in different systems- mouse and *Drosophila*. The studies conducted in the epidermis of chimeric mouse embryos (with dual PCP reporter cells with *Fz* and *Vang* markers and unlabelled Celsr1 mutant cells) by Basta et. al show that intercellular complexes formed by the atypical cadherin Celsr1 at cell-cell junctions are essential for cells to sort the PCP proteins across cell junctions [52]. As a consequence, individual cells that don't have contact with neighbours can't establish PCP asymmetry. However, Weiner et. al conducted experiments in *Drosophila* wing that block the intercellular homodimer formation by atypical cadherin *Fmi* (by replacing *Fmi* with *Fmi*Δcad which cannot form homodimers with neighboring cells but can help the assembly of PCP proteins inside cells), which show that individual cells can polarize autonomously [51].

To investigate how the model behaves under these two scenarios- cell autonomous versus cell non-autonomous polarization- we created a second version of our PCP model to simulate the cell non-autonomous case. While in the cell-autonomous case the internal contact energy were all positive ($j_{pd} = 30$ and all other internal energies were set to 1), for the cell non-autonomous case, all internal contact energies were set to be negative (Figure 5A). To check whether the model versions lead to consistent domain segregation or mixing, we tracked the distance between proximal–distal compartments and their common interface over time (Fig5B). In the cell-autonomous case, the distance remained approximately 0.8 times the cell diameter, indicating that compartments remained spatially separated at all times. However in the cell non-autonomous case, the distance fluctuated between 0 and 0.8 times cell diameter, showing that the compartments failed to spatially segregate (Fig5B left panel). Conversely, the interface between proximal-distal compartments remained near zero in the cell-autonomous case, whereas it fluctuated at non-zero values in the cell non-autonomous case (Fig5B right panel).

Next, we investigated the effect of system size on cell non-autonomous polarization under different boundary conditions. We found that global polarization decreases with increasing system size (Fig5C). However, for both boundary conditions this decrease is less pronounced compared to the decrease ob-

Figure 5: **Cell non-autonomous polarization (A)** Simulation snapshots showing cell-autonomous (as observed in *Drosophila* wing) and cell non-autonomous polarization (as observed in mouse epidermis), where the PCP proteins can either self-polarize or not in single cells. **(B)** Comparison of the distance (left panel) and interface (right panel) between proximal and distal compartments over time for both model versions. In the cell-autonomous case, the compartments have no common interface and remain spatially separated, while in the cell non-autonomous case the interface area is high and their distance fluctuates at values lower than 0.8, indicating mixing. **(C)** Global polarization of cell non-autonomous PCP model as a function of system size for periodic and left boundary signal configurations. As system size increases, global polarization decreases under both boundary conditions. **(D)** Final mean angle of tissue polarization for a 30×30 cell system: under periodic boundary conditions (left, red), the angles are uniformly distributed between $0°$ and $360°$; under the left boundary condition (right, blue), the angles are more dispersed along the proximal–distal axis. **(E)** Final cell polarity vectors for a 30×30 system with periodic boundary conditions (left): local alignment occurs within small domains. For the left boundary signal configuration (right), alignment is maintained in the first few columns but dissipates with distance from the boundary. **(F)** Comparison of global polarization for systems with and without cell proliferation as a function of the number of cells. For fixed tissue sizes, global polarization decreases with system size (red). Uniform cell proliferation enhances global polarization at a higher value 0.93 (green). **(G)** Simulation output showing preference for alignment along the proximal-distal axis for the uniform cell proliferation configuration.

served for cell autonomous polarization (Fig3A) and (Fig S7A and S7B). The final mean angle of tissue polarization for a 30×30 cell system under periodic boundary conditions were uniformly distributed in $[0, 360°]$ whereas under the left boundary condition, the angles were more dispersed along the proximal–distal axis (Fig5D). Simulation snapshots show that periodic boundary conditions produced small,

locally aligned domains (Fig5E, left), whereas left-boundary conditions maintained alignment along the proximal–distal axis near the boundary but lost alignment farther away (Fig5E, right).

We also examined the time taken by the system to reach 50% global polarization ($\phi = 0.5$) for these boundary conditions for cell autonomous and non-autonomous polarization scenarios. Systems with periodic boundaries polarized more slowly, while those with left-boundary signals polarized faster (Fig S8A). Across 100 simulations, most systems reached 50% polarization even with increasing size, but compared to the non-autonomous case, cell-autonomous polarization required nearly an order of magnitude more time, and the fraction of simulations achieving 50% polarization dropped sharply with size (Fig S8A, left). When we increased system size along the x-axis while keeping 10 cells along the y-axis, the time to reach 50% polarization grew with size in both cases, but much more sharply in the autonomous case. The fraction of simulations reaching 50% polarization decreased only slightly for non-autonomous cells but fell to 20% for autonomous cells (Fig S8B).

As in the case of cell autonomous polarization (Fig4E), uniform cell proliferation rescued the decline in global polarization observed for fixed tissues with increasing system size (Fig 5F). Uniform proliferation enhanced and stabilized global polarization at a higher mean value (0.93) compared to the cell autonomous scenario (Fig S7C). Simulation outputs show a preference for alignment along the proximal-distal axis for the uniform cell proliferation configuration (Fig 5G).

Discussion

Planar Cell Polarity signalling is a very common process used by multiple organisms to pattern both mesenchymal and epithelial tissues. Given its relevance, much effort has been put by both experimentalists and theoreticians to understand how this process takes place. All models of PCP to date require some sort of global cue to ensure robust tissue alignment at large scales, a result that found support on the existing assumption within the field that morphogenetic gradients such as Wnts were responsible for the observed coherent cell polarization at tissue scale. Given that two recent works put that assumption in check, new modelling efforts were needed to check whether long-range PCP alignment is possible without global cues and what is required to achieve such result.

To investigate how tissues can achieve long-range polarity without the use of global morphogens, we created a stochastic phenomenological model of PCP dynamics that relies only on the contact affinities between cell compartments within and across cells. By carefully choosing our model parameters, these local interactions lead to the segregation of proximal and distal compartments within cells and alignment of proximal-distal orientation among neighbouring cells (Fig 1C). Like all other models to date, we assume that the PCP proteins that make the distal and proximal domains have the ability to reorganize their spatial orientation using the local information provided by their immediate neighbours. This assumption was recently demonstrated experimentally by the work of the Devenport and Axelrod groups for both mice and flies, respectively [52] [51]. We systematically increased the size of our simulated tissues and found that global ordering is quickly lost with the number of simulated cells, confirming that local interactions alone cannot guarantee global ordering, a result shared with many other PCP models [41] [42]. These simulations were done with squared geometries and with periodic boundary conditions along both directions to avoid unwanted boundary effects that could interfere with the alignment of cells.

Next we tested whether the presence of a local signalling boundary would be sufficient to provide a bias that would orient the whole tissue in the same direction. Many tissues where PCP is present exhibit such local orienting signals. For example, in the *Drosophila* wing imaginal disc, PCP is aligned with respect to multiple signalling centers such as Wingless (Wg) and Notch at the dorso-ventral boundary, Hedgehog (Hh) and Decapentaplegic (Dpp) at the anterior-posterior compartment boundary, and Dachsous (Ds) at the hinge fold [53]. Planarians also have 3 signalling centers, two at the anterior and posterior poles of the animal, and one extending along the body margins [59]. In our simulations, we added a rigid wall on the left side of the tissue with adhesion properties of the distal compartment, which should induce the first row of cells to polarize proximal-distally, causing the subsequent rows to do the same. On the right side the tissue sees an open space which has no contact preference to proximal or distal cell compartments. This was done to prevent a double boundary with possible competing bias. Although the presence of a boundary signal does bias the overall tissue direction, global polarization still decreases with system size at the same rate as in the previous simulations without a boundary signal. Addition of a second boundary on the right side with proximal properties does reinforce the directional bias and increases the overall range of patterning, but does not prevent the loss of global polarization with increasing number of cells (Fig S9A,B). We also tested whether a wall with adhesion preference to lateral cell compartments

(i.e. neither proximal nor distal identities) could induce any polarity, but we found that this leads to independent rows of cells that can polarize either up or down (Fig S10A,B and C).

Given that for small systems a local boundary signal successfully leads to full tissue polarization in the direction specified by the boundary, we tested whether the combination of an initially small tissue, a local boundary and subsequent cell proliferation would suffice to generate large scale tissue orientation - a hypothesis first proposed by Sanger et al. in 2012 [53]. In our model for cell growth and division, we incorporated two important observations made in mammalian skin cells: that PCP components are inherited by daughter cells; and that cells lose their polarity information during division due to the temporary internalization of PCP proteins [56]. Therefore, cells in our model must reorient their proximal and distal compartments after division using only the local information from their immediate neighbours. Our results show that cell proliferation is sufficient to sustain and propagate the initially established polarity in an expanding tissue (Fig 4J). The ability of the tissue to sustain global polarization while growing is dependent on cell growth rates, with faster division cycles leading to an increasing loss in tissue polarization (Fig4G, Fig S6, Fig S11). We attribute this result to the need of the cells to properly re-establish their polarity and alignment with the neighbours before the next division and not disrupting the polarity of neighbours while growing. We note that as long as a sufficiently slow growth rate is chosen, the patterning mechanism is very robust. Figure 4E shows that the global polarization metric ϕ remains high throughout the whole course of the simulations while the tissue expands, with little variation in global orientation (Fig4F).

Our results also indicate a difference between uniform cell proliferation within the tissue, and restricted proliferation at the front, as seen in wound healing, regeneration and PCP establishment in the retina growth cone [60]. For proliferation on front, we observed a shift in the direction of polarity at either ±60 degrees from the proximal-distal axis that remained coherent for the rest of the simulations. We believe that the shift happens due to the lack of sufficient polarity cues for daughter cells at the front. We noticed a great impact of cell division planes on the onset of this shift, which occurs sooner when cell divisions are restricted along the minor axis compared to divisions restricted along the major axis, as the former division plane leaves the distal-most daughter cell with fewer polarity cues (Fig S12, Fig S4A and B). This is further supported by the simulations where cells undergo a single division cycle either inside the tissue or at the front, where we found that cells dividing inside the tissue re-polarize and align their directions ten times faster than cells dividing at the front (Fig S13). The observed ±60° deviation from the proximal-distal axis is surprising as it is not what would be expected due to pinning effects of the lattice. We believe that this angle preference is due to the geometry of our simulation space and the hexagonal packing that arises from front-restricted proliferations, where cells form rows along the direction orthogonal to the proximal-distal axis.

Whether single cells have the ability to polarize cell autonomously, however, seems to be organism specific: in mice, cells are not able to polarize autonomously [52], while in *Drosophila* they can [51]. In our CPM model, we can choose our parameters to model both cell autonomous and cell non-autonomous polarization and our simulations reveal clear differences between both these polarization mechanisms. Although the key results remain qualitatively the same in both autonomous and non-autonomous systems, the latter is better at propagating polarity signals across the entire tissue and establishing long range alignment. This is evident from the results where we observed consistently higher levels of global polarization for non-autonomous systems across all system sizes (Fig S7). These systems reached 50% global polarization threshold in nearly all simulation runs, and required significantly lesser time to reach that threshold (Fig S8). Even in a proliferating tissue, the polarization is slightly higher than that observed for autonomous systems (< 1%). Constraints imposed on the affinities between internal cell compartments in the cell autonomous case could be reducing the cell's ability to readjust its polarity direction and possibly letting the system to get trapped in a local minima state. In contrast, the dynamic internal compartments in the non-autonomous systems provide cells with greater flexibility to adapt their polarity to PCP signals from neighbours leading to an optimized tissue alignment. A similar effect was also observed in migratory cells and analysed in more detail by Nandan and Koseska [61][62].

All our results and analyses focus on the question of which mechanisms best create order from an initial unordered state. However, an equally interesting question is what is the maximum amount of ordering that different systems can maintain. Simulations of tissues that already start from a perfectly polarized initial condition shows a linear decrease in their polarization over time, with higher rates of decrease for larger systems (Fig S14). When compared to equivalent simulations where cells start from disordered states we noticed that the loss of polarization happens at a slower rate than the gain of polarization for smaller systems. We also noticed that global polarization is preserved when cells start with a perfectly hexagonal configuration (Fig S15), suggesting a strong influence of cell packing in the

maintenance of PCP polarization [55][63].

The mechanism we proposed here relies on cell proliferation from an initially small tissue where the low number of cells ensures an initial coherent pattern that can be expanded. However, this raises the question of how global alignment can be established in biological systems where tissues may already be large before PCP activation. A few alternate mechanisms may play a role in such cases. One possible mechanism is the use of two signalling boundaries (at the proximal and distal sides) that increases the maximum sizes that pre-growth tissues could polarize coherently (see Fig S9). Another mechanism would be a sequence of PCP cell activation from a boundary signal, where the next row of cells only polarizes after the preceding row has already polarized and aligned with the rest of the tissue (Fig S16). A similar mechanism was implemented by Salbreux et al. [60] to model the zebrafish retina development, where tissue scale mechanical stresses also played a role in elongating cells and biasing orientations in a specific direction. Other models also made use of mechanical stresses to aid global PCP alignment. In 2010, Aigouy et al. [64] proposed a vertex model where anisotropic tensions oriented cell elongations and rearrangements, leading to global realignment. A similar mechanism was proposed by Sugimura et al. [65], where anisotropy from cell elongation together with asymmetric cell-cell coupling along a particular axis can act as a global cue for polarity alignment. However, these models are based on a pre-established PCP order from which the applied tension realigns the orientation [64] [65], so it is not clear if these mechanisms could by themselves promote alignment from a highly disordered initial state.

There are multiple ways to test whether the mechanism proposed here is at play in any specific tissue. The most direct test is by manipulating the growth rates of cells as the model predicts better(worst) PCP alignments for slower(faster) growth rates. Another model prediction is that the presence of a signalling boundary is necessary to establish the proximal-distal direction in a growing tissue. If this boundary is removed or made unable to signal/influence the cells, the growing tissue is likely to orient in a different direction. Our model also predicts that the ability of a large tissue to keep its orientation after the removal/blockage of a signalling boundary will depend on cell packing: irregularly packed tissues will experience a slow decrease in their global polarizations proportional to the tissue size, while tissues that are packed in a highly ordered hexagonal tiling would keep their polarization. In the absence of a global cue, our model predicts that a boundary signal alone cannot ensure global polarization of static and large tissues. If PCP is activated in an already large tissue, or blocked until a tissue has grown large enough, the cells would only align locally and global polarization will not be achieved. Our model also suggests that cell non-autonomous PCP polarization provides a more robust polarization mechanism than cell autonomous PCP polarization. This is a species-specific difference, but may be tested by comparing the degree or rate of polarization between tissues of comparable size under similar conditions. Any perturbation that can switch the cell polarization between autonomous and non-autonomous would open the way to test our model predictions about the higher robustness of the cell non-autonomous model.

This work presents the first description of planar cell polarity using the Cellular Potts Model, which offers some advantages and disadvantages compared to the previous models of PCP. Because we are using a phenomenological model, the work here cannot account for the details in molecular pathways nor account for the effects of different PCP components concentrations as done in [38] [39] [40]. This shortcoming can be amended by associating each PCP domain with ODE models that capture molecular pathways and signalling, as was done in other CP models [54][66]. Off-grid models, such as cell center and vertex models avoid the undesired effects of lattice anisotropy and allow for an easy implementation of shear stresses on tissues, which would be challenging to do in CP models.

The use of the CPM framework, however, offers some unique advantages by allowing for a detailed representation of cell behaviours in a dynamically evolving tissue environment. Most previous models have considered non-motile cells with fixed shapes [38][39][42][41], with some exceptions incorporating variations in cell shape [43] or allowing cells to divide, change shape, and rearrange their neighbours [64]. In our model, cells are highly dynamic — they can change shape, exchange neighbours, and actively pack themselves through growth and rearrangement. Since PCP is also crucial for providing directional information and thereby guiding other developmental processes, our model based on Cellular Potts formalism provides a versatile framework for studying PCP-driven morphogenesis in other contexts for which Cellular Potts models have been used in the past, such as convergent-extension [67][68], limb-bud growth [69] and primitive streak formation [70]. The mechanisms of long-range polarity without morphogens that we explored here, however, are general and do not rely on our specific choice of modelling framework. We believe that the results of our work should be easy to replicate in other models where cells are able to proliferate and will be interesting to check how other processes such as stress and details on PCP signalling at the molecular level could impact the results.

Author contributions

JMB conceptualized the study. AT and JMB developed the computational model. AT conducted the investigation, formal analysis and visualization. AT and JMB wrote the manuscript.

Competing interests

The authors declare no conflicts of interest.

Materials and Methods

The model was developed using the CompuCell3D (CC3D) simulation software [54], with simulations executed on a computer cluster using a set of custom-developed Python scripts. The simulation code and the analysis scripts are publicly available at https://github.com/abhisha-ramesh/PCP_Subcellular_Potts_Model.

Acknowledgments

We would like to thank Dr. Eric Brooks for his valuable feedback and comments on this work. We also thank Dr. François Graner and Dr. James Glazier for their feedback with the implementation of certain aspects of the model. We also thank Dr. Juan Manuel Gomez, Dr. Lutz Brusch, Dr. Benedikt Best and Pedro Cencil Dal Castel for their feedback and comments on this work. We acknowledge the computing resources provided by North Carolina State University High Performance Computing Services Core Facility (RRID:SCR_022168). This work was supported by the North Carolina State University.

References

[1]   Suzanne Eaton. "Planar polarization of Drosophila and vertebrate epithelia". en. In: *Current Opinion in Cell Biology* 9.6 (Dec. 1997), pp. 860–866. ISSN: 09550674. DOI: 10.1016/S0955-0674(97)80089-0.

[2]   John B. Wallingford. "Planar Cell Polarity and the Developmental Control of Cell Behavior in Vertebrate Embryos". en. In: *Annual Review of Cell and Developmental Biology* 28.1 (Nov. 2012), pp. 627–653. ISSN: 1081-0706, 1530-8995. DOI: 10.1146/annurev-cellbio-092910-154208.

[3]   D. Gubb and A. García-Bellido. "A genetic analysis of the determination of cuticular polarity during development in *Drosophila melanogaster*". en. In: *Development* 68.1 (Apr. 1982), pp. 37–57. ISSN: 0950-1991, 1477-9129. DOI: 10.1242/dev.68.1.37.

[4]   Ray Keller. "Shaping the Vertebrate Body Plan by Polarized Embryonic Cell Movements". en. In: *Science* 298.5600 (Dec. 2002), pp. 1950–1954. ISSN: 0036-8075, 1095-9203. DOI: 10.1126/science.1079478.

[5]   Michael K. Dush and Nanette M. Nascone-Yoder. "Vangl2 coordinates cell rearrangements during gut elongation". eng. In: *Developmental Dynamics: An Official Publication of the American Association of Anatomists* 248.7 (July 2019), pp. 569–582. ISSN: 1097-0177. DOI: 10.1002/dvdy.61.

[6]   Zoha Kibar et al. "Ltap, a mammalian homolog of Drosophila Strabismus/Van Gogh, is altered in the mouse neural tube mutant Loop-tail". en. In: *Nature Genetics* 28.3 (July 2001), pp. 251–255. ISSN: 1546-1718. DOI: 10.1038/90081.

[7]   Amber Huffine Bogart and Eric R. Brooks. "Canonical Wnt pathway modulation is required to correctly execute multiple independent cellular dynamic programs during cranial neural tube closure". en. In: *Developmental Biology* 523 (July 2025), pp. 115–131. ISSN: 00121606. DOI: 10.1016/j.ydbio.2025.04.013.

[8]   Panpan Yao et al. "Wnt/PCP pathway regulates the migration and neural differentiation of mesenchymal stem cells in vitro". eng. In: *Folia Histochemica Et Cytobiologica* 60.1 (2022), pp. 44–54. ISSN: 1897-5631. DOI: 10.5603/FHC.a2022.0006.

[9]   John B. Wallingford et al. "Dishevelled controls cell polarity during Xenopus gastrulation". en. In: *Nature* 405.6782 (May 2000), pp. 81–85. ISSN: 1476-4687. DOI: 10.1038/35011077.

[10] Peter A. Lawrence, José Casal, and Gary Struhl. "Towards a model of the organisation of planar polarity and pattern in the *Drosophila* abdomen". en. In: *Development* 129.11 (June 2002), pp. 2749–2760. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.129.11.2749.

[11] Hitoshi Matakatsu and Seth S. Blair. "Interactions between Fat and Dachsous and the regulation of planar cell polarity in the *Drosophila wing*". en. In: *Development* 131.15 (Aug. 2004), pp. 3785–3794. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.01254.

[12] Ben Ewen-Campen et al. "No Evidence that Wnt Ligands Are Required for Planar Cell Polarity in Drosophila". en. In: *Cell Reports* 32.10 (Sept. 2020), p. 108121. ISSN: 22111247. DOI: 10.1016/j.celrep.2020.108121.

[13] Joyce J.S. Yu et al. "Frizzled-Dependent Planar Cell Polarity without Secreted Wnt Ligands". en. In: *Developmental Cell* 54.5 (Sept. 2020), 583–592.e5. ISSN: 15345807. DOI: 10.1016/j.devcel.2020.08.004.

[14] Lisa V. Goodrich and David Strutt. "Principles of planar polarity in animal development". en. In: *Development* 138.10 (May 2011), pp. 1877–1892. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.054080.

[15] Tadao Usui et al. "Flamingo, a Seven-Pass Transmembrane Cadherin, Regulates Planar Cell Polarity under the Control of Frizzled". en. In: *Cell* 98.5 (Sept. 1999), pp. 585–595. ISSN: 00928674. DOI: 10.1016/S0092-8674(00)80046-X.

[16] Yuko Shimada et al. "Asymmetric colocalization of Flamingo, a seven-pass transmembrane cadherin, and Dishevelled in planar cell polarization". en. In: *Current Biology* 11.11 (June 2001), pp. 859–863. ISSN: 09609822. DOI: 10.1016/S0960-9822(01)00233-0.

[17] Hing-C. Wong et al. "Direct Binding of the PDZ Domain of Dishevelled to a Conserved Internal Sequence in the C-Terminal Region of Frizzled". In: *Molecular Cell* 12.5 (Nov. 2003), pp. 1251–1260. ISSN: 1097-2765. DOI: 10.1016/S1097-2765(03)00427-1.

[18] Rebecca Bastock, Helen Strutt, and David Strutt. "Strabismus is asymmetrically localised and binds to Prickle and Dishevelled during *Drosophila* planar polarity patterning". en. In: *Development* 130.13 (July 2003), pp. 3007–3014. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.00526.

[19] David R.P. Tree et al. "Prickle Mediates Feedback Amplification to Generate Asymmetric Planar Cell Polarity Signaling". en. In: *Cell* 109.3 (May 2002), pp. 371–381. ISSN: 00928674. DOI: 10.1016/S0092-8674(02)00715-8.

[20] Samantha J. Warrington et al. "A Dual Function for Prickle in Regulating Frizzled Stability during Feedback-Dependent Amplification of Planar Polarity". en. In: *Current Biology* 27.18 (Sept. 2017), 2784–2797.e3. ISSN: 09609822. DOI: 10.1016/j.cub.2017.08.016.

[21] Andreas Jenny et al. "Diego and Prickle regulate Frizzled planar cell polarity signalling by competing for Dishevelled binding". en. In: *Nature Cell Biology* 7.7 (July 2005), pp. 691–697. ISSN: 1476-4679. DOI: 10.1038/ncb1271.

[22] D. I. Strutt. "Asymmetric localization of frizzled and the establishment of cell polarity in the Drosophila wing". eng. In: *Molecular Cell* 7.2 (Feb. 2001), pp. 367–375. ISSN: 1097-2765. DOI: 10.1016/s1097-2765(01)00184-8.

[23] J. D. Axelrod. "Unipolar membrane association of Dishevelled mediates Frizzled planar cell polarity signaling". eng. In: *Genes & Development* 15.10 (May 2001), pp. 1182–1187. ISSN: 0890-9369. DOI: 10.1101/gad.890501.

[24] Gishnu Das et al. "Diego interacts with Prickle and Strabismus/Van Gogh to localize planar cell polarity complexes". en. In: *Development* 131.18 (Sept. 2004), pp. 4467–4476. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.01317.

[25] Danelle Devenport. "The cell biology of planar cell polarity". en. In: *Journal of Cell Biology* 207.2 (Oct. 2014), pp. 171–179. ISSN: 1540-8140, 0021-9525. DOI: 10.1083/jcb.201408039.

[26] Jules Lavalou and Thomas Lecuit. "In search of conserved principles of planar cell polarization". en. In: *Current Opinion in Genetics & Development* 72 (Feb. 2022), pp. 69–81. ISSN: 0959437X. DOI: 10.1016/j.gde.2021.11.001.

[27] Hitoshi Matakatsu and Seth S. Blair. "Separating the adhesive and signaling functions of the Fat and Dachsous protocadherins". en. In: *Development* 133.12 (June 2006), pp. 2315–2324. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.02401.

[28] Abhijit A. Ambegaonkar et al. "Propagation of Dachsous-Fat Planar Cell Polarity". en. In: *Current Biology* 22.14 (July 2012), pp. 1302–1308. ISSN: 09609822. DOI: 10.1016/j.cub.2012.05.049.

[29] Amy Brittle, Chloe Thomas, and David Strutt. "Planar Polarity Specification through Asymmetric Subcellular Localization of Fat and Dachsous". en. In: *Current Biology* 22.10 (May 2012), pp. 907–914. ISSN: 09609822. DOI: 10.1016/j.cub.2012.03.053.

[30] José Casal, Peter A. Lawrence, and Gary Struhl. "Two separate molecular systems, Dachsous/Fat and Starry night/Frizzled,act independently to confer planar cell polarity". en. In: *Development* 133.22 (Nov. 2006), pp. 4561–4572. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.02641.

[31] Helen Strutt and David Strutt. "How do the Fat–Dachsous and core planar polarity pathways act together and independently to coordinate polarized cell behaviours?" en. In: *Open Biology* 11.2 (Feb. 2021), p. 200356. ISSN: 2046-2441. DOI: 10.1098/rsob.200356.

[32] Wen Yih Aw and Danelle Devenport. "Planar cell polarity: global inputs establishing cellular asymmetry". en. In: *Current Opinion in Cell Biology* 44 (Feb. 2017), pp. 110–116. ISSN: 09550674. DOI: 10.1016/j.ceb.2016.08.002.

[33] H.-Arno J. Müller, Reba Samanta, and Eric Wieschaus. "Wingless signaling in the *Drosophila* embryo: zygotic requirements and the role of the *frizzled* genes". en. In: *Development* 126.3 (Feb. 1999), pp. 577–586. ISSN: 0950-1991, 1477-9129. DOI: 10.1242/dev.126.3.577.

[34] Chiann-mun Chen and Gary Struhl. "Wingless transduction by the Frizzled and Frizzled2 proteins of *Drosophila*". en. In: *Development* 126.23 (Dec. 1999), pp. 5441–5452. ISSN: 0950-1991, 1477-9129. DOI: 10.1242/dev.126.23.5441.

[35] Purnima Bhanot et al. "Frizzled and DFrizzled-2 function as redundant receptors for Wingless during *Drosophila* embryonic development". en. In: *Development* 126.18 (Sept. 1999), pp. 4175–4186. ISSN: 0950-1991, 1477-9129. DOI: 10.1242/dev.126.18.4175.

[36] Monica Sharma et al. "Dishevelled: A masterful conductor of complex Wnt signals". en. In: *Cellular Signalling* 47 (July 2018), pp. 52–64. ISSN: 08986568. DOI: 10.1016/j.cellsig.2018.03.004.

[37] Jun Wu et al. "Wg and Wnt4 provide long-range directional input to planar cell polarity orientation in Drosophila". en. In: *Nature Cell Biology* 15.9 (Sept. 2013), pp. 1045–1055. ISSN: 1465-7392, 1476-4679. DOI: 10.1038/ncb2806.

[38] Keith Amonlirdviman et al. "Mathematical Modeling of Planar Cell Polarity to Understand Domineering Nonautonomy". en. In: *Science* 307.5708 (Jan. 2005), pp. 423–426. ISSN: 0036-8075, 1095-9203. DOI: 10.1126/science.1105471.

[39] Jean-François Le Garrec, Philippe Lopez, and Michel Kerszberg. "Establishment and maintenance of planar epithelial cell polarity by asymmetric cadherin bridges: A computer model". en. In: *Developmental Dynamics* 235.1 (Jan. 2006), pp. 235–246. ISSN: 1058-8388, 1097-0177. DOI: 10.1002/dvdy.20617.

[40] Sabine Fischer et al. "Is a persistent global bias necessary for the establishment of planar cell polarity?" eng. In: *PloS One* 8.4 (2013), e60064. ISSN: 1932-6203. DOI: 10.1371/journal.pone.0060064.

[41] Yoram Burak and Boris I. Shraiman. "Order and Stochastic Dynamics in Drosophila Planar Cell Polarity". en. In: *PLoS Computational Biology* 5.12 (Dec. 2009). Ed. by Anand R. Asthagiri, e1000628. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1000628.

[42] Yanshu Wang, Tudor Badea, and Jeremy Nathans. "Order from disorder: Self-organization in mammalian hair patterning". en. In: *Proceedings of the National Academy of Sciences* 103.52 (Dec. 2006), pp. 19800–19805. ISSN: 0027-8424, 1091-6490. DOI: 10.1073/pnas.0609712104.

[43] Hao Zhu. "Is anisotropic propagation of polarized molecular distribution the common mechanism of swirling patterns of planar cell polarization?" en. In: *Journal of Theoretical Biology* 256.3 (Feb. 2009), pp. 315–325. ISSN: 00225193. DOI: 10.1016/j.jtbi.2008.08.029.

[44] S. Schamberg et al. "Modelling and Analysis of Planar Cell Polarity". en. In: *Bulletin of Mathematical Biology* 72.3 (Apr. 2010), pp. 645–680. ISSN: 0092-8240, 1522-9602. DOI: 10.1007/s11538-009-9464-0.

[45] Katie Abley et al. "An intracellular partitioning-based framework for tissue cell polarity in plants and animals". en. In: *Development* 140.10 (May 2013), pp. 2061–2074. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.062984.

[46] Lee D Hazelwood and John M Hancock. "Functional modelling of planar cell polarity: an approach for identifying molecular function". en. In: *BMC Developmental Biology* 13.1 (May 2013), p. 20. ISSN: 1471-213X. DOI: 10.1186/1471-213X-13-20.

[47] Katherine H. Fisher and David Strutt. "A theoretical framework for planar polarity establishment through interpretation of graded cues by molecular bridges". en. In: *Development* 146.3 (Feb. 2019), dev168955. ISSN: 1477-9129, 0950-1991. DOI: 10.1242/dev.168955.

[48] Divyoj Singh et al. "Emergence of planar cell polarity from the interplay of local interactions and global gradients". In: *eLife* 13 (Oct. 2024). Ed. by Karsten Kruse and Aleksandra M Walczak, e84053. ISSN: 2050-084X. DOI: 10.7554/eLife.84053.

[49] François Graner and James A. Glazier. "Simulation of biological cell sorting using a two-dimensional extended Potts model". en. In: *Physical Review Letters* 69.13 (Sept. 1992), pp. 2013–2016. ISSN: 0031-9007. DOI: 10.1103/PhysRevLett.69.2013.

[50] J. A. Glazier and F. Graner. "Simulation of the differential adhesion driven rearrangement of biological cells". eng. In: *Physical Review. E, Statistical Physics, Plasmas, Fluids, and Related Interdisciplinary Topics* 47.3 (Mar. 1993), pp. 2128–2154. ISSN: 1063-651X. DOI: 10.1103/physreve.47.2128.

[51] Alexis T Weiner et al. *Cell autonomous polarization by the planar cell polarity signaling pathway.* en. Sept. 2023. DOI: 10.1101/2023.09.26.559449.

[52] Lena P. Basta et al. "Epithelial polarization by the planar cell polarity complex is exclusively non–cell autonomous". en. In: *Science* 387.6740 (Mar. 2025), eads5704. ISSN: 0036-8075, 1095-9203. DOI: 10.1126/science.ads5704.

[53] Andreas Sagner et al. "Establishment of Global Patterns of Planar Polarity during Growth of the Drosophila Wing Epithelium". en. In: *Current Biology* 22.14 (July 2012), pp. 1296–1301. ISSN: 09609822. DOI: 10.1016/j.cub.2012.04.066.

[54] Maciej H. Swat et al. "Multi-Scale Modeling of Tissues Using CompuCell3D". en. In: *Methods in Cell Biology*. Vol. 110. Elsevier, 2012, pp. 325–366. ISBN: 9780123884039. DOI: 10.1016/B978-0-12-388403-9.00013-8.

[55] Anne-Kathrin Classen et al. "Hexagonal Packing of Drosophila Wing Epithelial Cells by the Planar Cell Polarity Pathway". en. In: *Developmental Cell* 9.6 (Dec. 2005), pp. 805–817. ISSN: 15345807. DOI: 10.1016/j.devcel.2005.10.016.

[56] Danelle Devenport et al. "Mitotic internalization of planar cell polarity proteins preserves tissue polarity". en. In: *Nature Cell Biology* 13.8 (Aug. 2011), pp. 893–902. ISSN: 1465-7392, 1476-4679. DOI: 10.1038/ncb2284.

[57] Tamás Vicsek et al. "Novel Type of Phase Transition in a System of Self-Driven Particles". en. In: *Physical Review Letters* 75.6 (Aug. 1995), pp. 1226–1229. ISSN: 0031-9007, 1079-7114. DOI: 10.1103/PhysRevLett.75.1226.

[58] Tamás Vicsek and Anna Zafeiris. "Collective motion". en. In: *Physics Reports* 517.3-4 (Aug. 2012), pp. 71–140. ISSN: 03701573. DOI: 10.1016/j.physrep.2012.03.004.

[59] Hanh Thi-Kim Vu et al. "Dynamic Polarization of the Multiciliated Planarian Epidermis between Body Plan Landmarks". en. In: *Developmental Cell* 51.4 (Nov. 2019), 526–542.e6. ISSN: 15345807. DOI: 10.1016/j.devcel.2019.10.022.

[60] Guillaume Salbreux et al. "Coupling Mechanical Deformations and Planar Cell Polarity to Create Regular Patterns in the Zebrafish Retina". en. In: *PLoS Computational Biology* 8.8 (Aug. 2012). Ed. by Stanislav Shvartsman, e1002618. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1002618.

[61] Akhilesh Nandan and Aneta Koseska. "Non-asymptotic transients away from steady states determine cellular responsiveness to dynamic spatial-temporal signals". en. In: *PLOS Computational Biology* 19.8 (Aug. 2023). Ed. by Attila Csikász-Nagy, e1011388. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1011388.

[62] Akhilesh Nandan et al. "Cells use molecular working memory to navigate in changing chemoattractant fields". en. In: *eLife* 11 (June 2022), e76825. ISSN: 2050-084X. DOI: 10.7554/eLife.76825.

[63] Dali Ma et al. "Cell packing influences planar cell polarity signaling". en. In: *Proceedings of the National Academy of Sciences* 105.48 (Dec. 2008), pp. 18800–18805. ISSN: 0027-8424, 1091-6490. DOI: 10.1073/pnas.0808868105.

[64]   Benoît Aigouy et al. "Cell Flow Reorients the Axis of Planar Polarity in the Wing Epithelium of Drosophila". en. In: *Cell* 142.5 (Sept. 2010), pp. 773–786. ISSN: 00928674. DOI: 10.1016/j.cell.2010.07.042.

[65]   Kaori Sugimura and Hiroshi Kori. "A reduced cell-based phase model for tissue polarity alignment through global anisotropic cues". en. In: *Scientific Reports* 7.1 (Dec. 2017), p. 17466. ISSN: 2045-2322. DOI: 10.1038/s41598-017-17611-8.

[66]   Susan D. Hester et al. "A Multi-cell, Multi-scale Model of Vertebrate Segmentation and Somite Formation". en. In: *PLoS Computational Biology* 7.10 (Oct. 2011). Ed. by Edmund J. Crampin, e1002155. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1002155.

[67]   Mark Zajac, Gerald L. Jones, and James A. Glazier. "Simulating convergent extension by way of anisotropic differential adhesion". In: *Journal of Theoretical Biology* 222.2 (May 2003), pp. 247–259. ISSN: 0022-5193. DOI: 10.1016/S0022-5193(03)00033-X.

[68]   Julio M. Belmonte, Maciej H. Swat, and James A. Glazier. "Filopodial-Tension Model of Convergent-Extension of Tissues". en. In: *PLOS Computational Biology* 12.6 (June 2016), e1004952. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1004952.

[69]   Gaja Lesnicar-Pucko et al. *Cellular mechanisms of chick limb bud morphogenesis*. en. Sept. 2020. DOI: 10.1101/2020.09.10.292359.

[70]   Bakhtier Vasiev et al. "Modeling Gastrulation in the Chick Embryo: Formation of the Primitive Streak". en. In: *PLOS ONE* 5.5 (May 2010), e10571. ISSN: 1932-6203. DOI: 10.1371/journal.pone.0010571.