1 **Amino acid sensing by the α-cell mitochondrial phosphoenolpyruvate cycle regulates**
2 **intracellular Ca^{2+} levels independently of glucagon secretion**

3 Erli Jin[1], Hannah R. Foster[1], Evgeniy Potapenko[1], Shih Ming Huang[1], Xinhang Dong[2], Jing W.
4 Hughes[2], and Matthew J. Merrins[1,3]

5
[1]Department of Medicine, Division of Endocrinology, Diabetes, and Metabolism, University of
Wisconsin-Madison, Madison, WI 53705, USA
[2]Department of Cell Biology and Physiology, Washington University School of Medicine, St.
Louis, MO 63110, USA
[3]William S. Middleton Memorial Veterans Hospital, Madison, WI 53705, USA
*Correspondence: mmerrins@medicine.wisc.edu (M.J.M)

Keywords: Glucagon secretion, pancreatic α-cells, amino acid metabolism, pyruvate kinase,
mitochondrial phosphoenolpyruvate carboxykinase.

Abstract

Pancreatic islet α-cells are increasingly recognized as amino acid sensors for the organism.
Building on our prior work in β-cells, we sought to determine whether the mitochondrial
phosphoenolpyruvate (PEP) cycle is involved in α-cell amino acid sensing. Three different
methods were used to probe the PEP cycle, including pyruvate kinase activators (TEPP-46), and
mice with α-cell specific deletion of pyruvate kinase (PKM1/2-αKO) or mitochondrial PEP
carboxykinase (PCK2-αKO). The mitochondrial fuels glutamine/leucine antagonized
alanine/arginine-stimulated Ca^{2+} influx and glucagon secretion under hypoglycemic conditions.
Both PKM1/2 and PCK2 were required for glutamine/leucine to close K$_{ATP}$ channels and limit
amino acid-stimulated membrane depolarization. The Ca^{2+} response to amino acids was
suppressed by pyruvate kinase activation with TEPP-46, and enhanced by α-cell deletion of
pyruvate kinase or PCK2 – all without changing glucagon secretion. Finally, using
diazoxide/KCl to probe the pathways downstream of membrane depolarization, we identified an
essential role of the PEP cycle in homeostatically restoring intracellular Ca^{2+} levels. In sum, the
α-cell mitochondrial PEP cycle senses glutamine/leucine and inhibits K$_{ATP}$ channels similarly to
β-cells, while restricting amino acid-stimulated membrane depolarization and Ca^{2+} influx.
However, defying expectations, none of the amino acids tested, including alanine/arginine,
regulate glucagon secretion by modulating membrane depolarization or intracellular Ca^{2+}.

Graphical abstract

α-cells as amino acids sensors. Arginine, alanine, and glutamine potentiate glucagon secretion,
while leucine has a suppressive effect (left). Independently of glucagon secretion, glutamine and
leucine suppress alanine and arginine-stimulated Ca^{2+} influx via the phosphoenolpyruvate (PEP)
cycle, which closes K_{ATP} channels and suppresses V_m depolarization (right).

Highlights

• Our studies identify a role for the α-cell PEP cycle in sensing amino acids under
hypoglycemic conditions.
• Pyruvate kinase and PCK2 are required for glutamine/leucine to close α-cell K_{ATP} channels
and limit membrane depolarization and Ca^{2+} influx.
• Glutamine/leucine oppose alanine/arginine-stimulated Ca^{2+} influx and glucagon secretion.
• All of the amino acids tested regulate glucagon secretion, but none do so by modulating
membrane depolarization or intracellular Ca^{2+} levels.

Introduction

Glucagon release from pancreatic islet α-cells is elevated under fasting and hypoglycemic
conditions. Under these conditions, when inhibitory paracrine signals from neighboring islet β/δ-
cells are reduced, α-cells respond strongly to non-glucose fuels including fatty acids and amino
acids, which modulate Ca^{2+} and glucagon secretion (1–5). Glucagon secretion is dependent on
voltage-dependent Ca^{2+} channels (VDCC) (6–8), which trigger downstream endoplasmic
reticulum Ca^{2+} release (9). The activity of these channels is tightly regulated by membrane
potential (V_m). Among the many ion channels that regulate α-cell V_m, the activity of ATP-
sensitive K^+ (K_{ATP}) channels is thought to be an important determinant of α-cell V_m, Ca^{2+}, and
glucagon release (10–15). When glucose is elevated, glycolysis inhibits α-cell K_{ATP} channels and
restricts Ca^{2+} entry (10,16–20). However, the metabolic pathways that control α-cell function in
response to amino acid stimulation remain unknown.

How exactly do individual amino acids impact glucagon secretion? The mechanisms are diverse. Some amino acids, like glutamate and glycine, stimulate glucagon secretion through α-cell membrane receptors (21–23). Charged amino acids like arginine may stimulate Ca^{2+} influx by directly depolarizing V_m (24,25). Alanine, which is uncharged, may also regulate V_m due to co-transport with Na^+ (26,27). In contrast, the branched chain amino acids are inhibitory to glucagon release (28–30). Leucine, for example, has been shown to suppress glucagon release by lowering cAMP, both via a direct effect on the α-cell and by stimulating inhibitory paracrine signaling via β/δ-cells (31). Beyond secretion, glutamine is required for amino acid stimulated α-cell proliferation (32), a mechanism that links glucagon release to amino acid catabolism in the liver-α-cell axis (3,33). Each of these findings indicate that amino acids are crucial to α-cell viability and function, but little is known about their intracellular metabolism.

In this paper, we focused on the mitochondrial fuels glutamine and leucine, and their control of α-cell V_m and Ca^{2+} by the glucagonotropic amino acids alanine and arginine. Due to the paucity of α-cell data, our working model was based on prior work in β-cells (18,34,35). At low glucose, glutamine and leucine evoke insulin secretion by stimulating mitochondrial phosphoenolpyruvate (PEP) production by the PCK2 enzyme. Mitochondria PEP is then used by plasma-membrane associated pyruvate kinase (PK), which raises the ATP/ADP ratio to close K_{ATP} channels, depolarize V_m, and open VDCC (18,34–36). In the PEP cycle, pyruvate generated by the PK reaction is sequentially metabolized by mitochondrial pyruvate carboxylase and PCK2, and returned to cytosol as PEP to augment flux through PK. Similar to β-cells, α-cell K_{ATP} channels are locally controlled by plasma membrane-associated enzymes including PK (18). However, these experiments were performed using excised patch-clamp, leaving it unclear whether PK and the PEP cycle are active in α-cells and if they regulate V_m, Ca^{2+}, or glucagon secretion.

Research Design and methods

Mice

Wild-type C57BL/6J mice were obtained from The Jackson Laboratory. *GcgCre^ERT* mice (Jax 030346) (46) were crossed with *GCaMP6s* mice (Jax 028866), a Cre-dependent Ca^{2+} indicator strain. *Pkm^{f/f}* mice were generated using CRISPR and sequence-verified by the University of Wisconsin-Madison Genome Editing and Animal Model Core, as done previously to generate *Pck2^{f/f}* mice (35). *Pkm^{f/f}* and *Pck2^{f/f}* mice were crossed with *GcgCre^ERT:GCaMP6s* mice to generate *GcgCre^ERT:GCaMP6s:Pkm^{f/f}* (PKM1/2-αKO) and *GcgCre^ERT:GCaMP6s:Pck2^{f/f}* (PCK2-αKO) and their littermate *GcgCre^ERT:GCaMP6s* controls. As in our prior study, *GcgCre^ERT* was present in the dams to facilitate *loxP* excision in the study animals without the need for tamoxifen (31). Mice were used between 11-24 weeks of age. Although 4-hydroxytamoxifen (100 nM) was only strictly necessary for adenoviral expression of Cre-dependent JEDI-2P biosensors, it was included in the culture media in all islet studies for consistency, after verifying that hormone secretion was unaffected (Supplemental Fig. 6). All procedures involving animals were approved by the Institutional Animal Care and Use Committees of the William S. Middleton Memorial Veterans Hospital and the University of Wisconsin-Madison and followed the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

In vivo **assays**

For meal tolerance tests, mice were fasted for 5 hours prior to oral gavage of liquid Ensure (10
mL/kg). For insulin tolerance tests, animals were fasted for 5 hours, followed by IP-injection of
0.75 mg/kg Humilin-R (Eli Lilly). Blood glucose was measured from the tail using a glucometer
(Contour). Glucagon was assayed from blood plasma using a mouse glucagon ELISA kit (Crystal
Chem 81518) assayed with a TECAN Spark plate reader.

Islet Perifusion Assays

Islets from 5-6 mice of each genotype were pooled and then divided in equal numbers (80-100
islets/chamber) and placed into a 12-channel perifusion system (BioRep Peri-5 or homemade
perifusion system) containing KRPH buffer (in mM: 140 NaCl, 4.7 KCl, 2.5 $CaCl_2$, 1 NaH_2PO_4,
1 $MgSO_4$, 2 $NaHCO_3$, 5 HEPES, 2 glucose, 0.1% fatty acid-free BSA, pH 7.4) using 100 µL Bio-
Gel P-4 Media (Bio-Rad #1504124) to stabilize the islets. Islets were equilibrated at 2 mM
glucose for 48 min prior to perfusion with AA at 37°C. Glucagon secretion was assayed using
Promega Lumit Glucagon Immunoassay (W8020) and measured using a TECAN Spark plate
reader. The Quant-IT PicoGreen dsDNA Assay Kit (Invitrogen P7589) was used to determine
DNA content after lysis in 0.1% Triton X-100.

Immunofluorescence staining and quantification

Isolated islets were fixed in 4% paraformaldehyde, permeabilized in CAS-Block (Invitrogen)
containing 0.3% Triton X-100 for 1 hour at room temperature, and subsequently incubated
overnight at 4°C with primary antibodies diluted in CAS-Block/Triton: rabbit-anti-PKM1/2
(CST3190T, 1:200); rabbit-anti-PCK2 (CST6924S, 1:200); mouse-anti-glucagon (ab10988,
1:800). Following five 10-minute washes in PBST (PBS with 0.1% Triton X-100), islets were
incubated with secondary antibodies (Abberior STAR Orange anti-mouse and STAR Red anti-
rabbit, 1:200) for 4 hours at room temperature, followed by five 10-minute PBST washes.
Stained islets were mounted on glass slides using ProLong Gold Antifade (Invitrogen) and
imaged on a Nikon AXR NSPARC confocal microscope. Z-stacks were acquired at 1 µm step
size, covering 10-30 planes per islet. For quantification, glucagon signal was used to define α-
cell vs. non-α-cell regions of interest (ROI). Whole-cell PKM1/2 and PCK2 staining intensities
were measured within randomly selected ROIs across z-planes and islets. Statistical comparisons
were performed using ordinary one-way ANOVA.

FACS sorting and qPCR

Islets isolated from 6mice/group were pooled and dispersed with trypsin (TrypLE, Thermo
fisher). Cells were sorted using FACSAriaII at University of Wisconsin Carbone Cancer Center
Flow Lab. α and non α-cells were sorted based on GCaMP6s signal. RNA was isolated using
QIAGEN RNeasy mini kit according to the manufacturer's instructions. cDNA was prepared
using monkey Moloney virus reverse transcriptase (Life Technologies) then analyzed by
quantitative PCR (qPCR) using Power SYBR Master Mix (Life Technologies)(Forward primer:
TGGCCGTGCAATCCAGAAAA; Reverse primer: TTGGTGATGCCCAAAATCAGCAT). All
transcripts were normalized to β-actin. Fold changes were calculated relative to vehicle only
controls.

Electrophysiology

Mouse islets were dispersed with Accutase (Sigma-Aldrich A6964100ML) and plated on
sterilized uncoated glass shards and cultured for up to 3 days. Recording electrodes made of
microfilament borosilicate glass (Harvard Apparatus) were used to pull ~3 MΩ patch pipettes
using a Flaming/Brown micropipette puller (P-1000, Sutter Instruments), and polished
(Narishige MF-830) to a final tip resistance of 4-5 MΩ. Cell-attached recording started after
formation of a stable gigaseal (>3 GΩ). Recordings were made by using HEKA Instruments
EPC10 patch-clamp amplifier at room temperature. The pipette solution contained (in mM): 10
sucrose, 130 KCl, 2 $CaCl_2$, 0.9 $MgCl_2$, 10 EGTA, 20 HEPES, pH 7.2, adjusted with KOH. The
extracellular bath solution contained (in mM): 140 NaCl, 5 KCl, 1.2 $MgCl_2$, 2.5 $CaCl_2$, 0.5
glucose, 10 HEPES, pH 7.4, adjusted with NaOH. Holding membrane potential was set at −50
mV during recording. Data was filtered online at 1 kHz with a Bessel filter and analyzed offline
using Clampfit 10.7 software (Molecular Devices). During each condition, recordings are held
for several minutes for the activity to stabilize, such that no significant further changes in
channel conductance were observed. A 120-s window at the end of each condition was used to
analyze K_{ATP} activity in terms of power, frequency, and duration and open time as previously
described (18), all of which were normalized to control conditions. Membrane potential
measurement was performed as in(47). Briefly, Intact islets were perfused with a standard
external solution containing (135 mM NaCl, 4.8 mM KCl, 5 mM $CaCl_2$, 1.2 mM $MgCl_2$, 20 mM
HEPES, 10 mM glucose; pH 7.35) with 2mM glucose and amino acids as indicated. Pipette tips
were filled with an internal solution (28.4 mM K_2SO_4, 63.7 mM KCl, 11.8 mM NaCl, 1 mM
$MgCl_2$, 20.8 mM HEPES, 0.5 mM EGTA; 40 mM sucrose; pH 7.2) containing 0.36 mg/mL
amphotericin B. Islet α and β-cells were identified by their characteristic I-V curves
(Supplemental Fig. 3)

Islet Ca^{2+} and V_m imaging

Reagents were obtained from Sigma-Aldrich unless indicated otherwise. For imaging of α-cell
Ca^{2+}, islets isolated from *GcgCre^{ERT}:GCaMP6s* mice were cultured in the presence of 100 nM 4-
hydroxytamoxifen and imaged 3 days post isolation. Islets with different genotypes were
barcoded
391 by preincubation with 2 µM DiR (Thermo Fisher Scientific D12731) in 2ml islet media for
15 min at 37 °C. Islet barcode was then imaged using Cy7 (for DiR) from Chroma.For imaging
of α-cell membrane potential, islets from *GcgCre^{ERT}* mice were incubated with adenovirus
expressing Cre-dependent JEDI-2P biosensors (Ad-CMV-FlexOn-JEDI-2P, VectorBuilder) for 2
hours, transferred to fresh media containing 100 nM 4-hydroxytamoxifen, and imaged 3 days
post isolation. Islets were imaged on an RC-41LP chamber (Warner Instruments) mounted on a
Nikon Ti2 microscope equipped with a 10×/0.5 NA SuperFluor objective (Nikon). The imaging
solution contained (in mM): 137 NaCl, 5.6 KCl, 1.2 $MgCl_2$, 0.5 NaH_2PO_4, 4.2 $NaHCO_3$, 10
HEPES, and 2.6 $CaCl_2$ (pH 7.4). Metabolites, mouse GIP (Pheonix Pharmaceuticals), and
pyruvate kinase activator (TEPP-46, Calbiochem) were added as indicated. Islets were imaged at
a constant flow rate of 250 µL/min using a feedback-controlled flow cell (Fluigent) and the
temperature was maintained at 37°C with chamber and solution heaters (Warner). Filters for
GCAMP6s and JEDI-2P imaging were as follows: exciter, ET500/20x; dichroic, ET-Fura2-GFP;
emitter, ET535/30m (Chroma). Images were captured with a Hamamatsu ORCA-Flash4.0
CMOS camera every 6 s. A single region of interest was used to quantify the average response of
each islet using NIS-Elements software (Nikon Instruments). Ca^{2+} and membrane potential were

reported as the fluorescence intensity normalized to the initial condition (F/F$_o$). Single α-cell
quantification was performed with the Imaris analysis software (Oxford Instruments).
198
199

Results

Glutamine and leucine suppress α-cell Ca^{2+} influx stimulated by alanine and arginine

Mixed amino acids, provided at their concentrations measured in mouse portal blood
(QLRA; 0.6 mM glutamine, 0.5 mM leucine, 0.2 mM arginine, 2 mM alanine) (26), stimulated
biphasic glucagon secretion from mouse islets under low glucose conditions (Fig. 1A). To
determine how amino acids regulate α-cell Ca^{2+}, we imaged islets isolated from mice expressing
GCaMP6s biosensors in α-cells (*GcgCre^{ERT}:GCaMP6s*). Although single α-cells exhibited
heterogeneity in their Ca^{2+} response, the aggregate islet response to mixed amino acids was a
sustained rise in Ca^{2+} (Fig. 1B). Opening K$_{ATP}$ channels with 200 μM diazoxide (Fig. 1C) or
blocking L-type VDCCs with 1 μM isradipine (Fig. 1D) suppressed the Ca^{2+} rise stimulated by
amino acids, confirming the requirement for membrane depolarization and L-type VDCCs
(6,9,15).

We applied various combinations of amino acids at low glucose to identify their
individual effects on α-cell Ca^{2+}. Arginine induced only a transient Ca^{2+} rise, while alanine
provoked a sustained increase (Fig. 1E,F). The effects of arginine and alanine were additive. On
their own, the combination of glutamine and leucine only slightly reduced average Ca^{2+},
however, they impaired the intracellular Ca^{2+} response to alanine and arginine by 42% (cf. Figs.
1G and H). In a perifusion assay, glutamine and leucine suppressed glucagon secretion
stimulated by arginine and alanine (Fig. 1I). Glucagon release fell in response to glutamine
withdrawal, and the glucagonostatic effect of leucine was reduced (Fig. 1J). Taken together,
alanine and arginine stimulate Ca^{2+} influx and glucagon secretion, while the combination of
glutamine and leucine has a suppressive effect. Since amino acids were applied to the entire islet
in theses assays, the inhibitory effects of paracrine signaling cannot be completely ignored (*e.g.*,
glutamine/leucine stimulate insulin release, Supplemental Fig. 1). We therefore sought additional
approaches to isolate the effects of amino acids on α-cell metabolism.

    Figure 1

227
Glutamine/leucine suppress α-cell Ca^{2+} influx stimulated by alanine/arginine. *A*: Perifusion assay
was used to measure glucagon secretion from mouse islets in the presence of 2 mM glucose (2G)
and amino acids (QLRA, in mM: 0.6 mM glutamine, 0.5 mM leucine, 0.2 mM arginine, 2 mM
and alanine). *B*: Heatmap (left) and example traces (right) of Ca^{2+} responses in intact mouse
islets expressing GCaMP6s biosensors. Average Ca^{2+} is shown in the white trace (left panel). *C-*
D Amino acid stimulated Ca^{2+} influx is blocked by 200μM diazoxide (K$_{ATP}$ channel opener) (*C*)
and 10 μM isradipine (L-type VDCC blocker) (*D*). *E-J*: α-cell Ca^{2+} (*E-H*) and glucagon secretion
(*I-J*) stimulated by QLRA alone or in combination. Imaging data (*B-H*) represent averaged
normalized GCaMP6s intensity from *n* = 40-56 islets from 3 mice imaged per protocol.

Perifusion data (*A*, *I*, and *J*) represent the average of $n = 3$ chambers from $n = 3$ mice per
condition with 80–100 mouse islets per chamber. Data are shown as mean ± SEM with AUC for
each condition compared with control by Student's *t* test. ***P* < 0.05, ****P* < 0.01, *****P* <
0.0001.

Mice lacking α-cell PKM1/2 or PCK2 respond normally to changing glucose levels

To test the function of the α-cell PEP cycle, we generated mice with α-cell specific
knockout of PKM1/2 or PCK2 by crossing *GcgCre^{ERT}:GCaMP6s* mice with *Pkm^{f/f}* mice
(PKM1/2-αKO,   *GcgCre^{ERT}:GCaMP6s:Pkm^{f/f})*   or   *Pck2^{f/f}*   mice   (PCK2-αKO,
GcgCre^{ERT}:GCaMP6s:Pck2^{f/f}). All studies were performed with littermate *GcgCre^{ERT}:GCaMP6s*
controls. α-Cell specific knockdown of PKM1/2 and PCK2 was evident by immunostaining of
isolated islets (Fig. 2A,B), however high background fluorescence made absolute quantification
difficult. After pooling FACS-sorted GCaMP6s$^+$ α-cells from 6 mice per genotype, we confirmed
by Western blot that PKM1 and PKM2 were virtually absent in α-cells from PKM1/2-αKO islets
(Fig. 2C). Although PCK2 was undetectable by Western blot, we confirmed the loss of *Pck2*
mRNA using QPCR of FACS-sorted α-cells; *Pck2* mRNA was reduced by 88% in PCK2-αKO α-
cells relative to controls.

To investigate glycemic control in the PKM1/2-αKO and PCK2-αKO mice, we
performed meal tolerance tests (MTT) by gavage of protein shake after a 5 hour fast. No
differences in blood glucose clearance were observed in the PCK2-αKO mice relative to controls
(Fig. 2D). A small improvement in glucose tolerance was observed in the PCK2-αKO mice
relative to controls, however this effect appears to be driven by elevated glucose in the control
mice at a single timepoint (Fig. 2E). To test the function of PKM1/2-αKO and PCK2-αKO mice
under low glucose conditions, fasted mice were subjected to an insulin tolerance test. No
difference in fasting glucagon was observed in either strain, and insulin lowered blood glucose
and stimulated glucagon secretion to a similar extent in PKM1/2-αKO and PCK2-αKO mice as
controls (Fig. 2F,G). Thus, both strains of knockout mice responded normally to raising or
lowering blood glucose.

 Figure 2

Mice lacking α-cell PKM1/2 or PCK2 respond normally to changing glucose levels. *A-B*:
Representative images of stained islets showing targeted deletion of PCK2 in Glucagon[+] α-cells.
Insets highlight α-cells (asterisks) with intact (control) or reduced (PCK2-αKO) enzyme
expression. Scale bar, 50 µm. Quantification represents *n* = 100 cells from 10 islets per genotype.
C: Western blot showing PKM1/2 knockdown in FACS sorted α-cells (GCaMP6s[+]) and non-α-
cells (GCaMP6s[-]) from PKM1/2-αKO and controls (pooled samples from 6 mice per group). D-
E: Meal tolerance test (MTT) with PKM1/2 and PCK2 α-KO. *n* = 10 mice per group. *F-G*:
Plasma glucose and glucagon levels in PKM1/2-αKO, PCK2α-KO, and littermate controls in
response to an insulin tolerance test (ITT). Data are shown as mean ± SEM with AUC for each
condition compared with control by Student's *t* test (*C-D*) or one-way ANOVA (*B*, *E-F*). *P <
0.05, ***P < 0.001, ****P < 0.0001, ns, not significant.

The α-cell PEP cycle closes K$_{ATP}$ channels and restricts V$_m$ depolarization in response to
amino acid stimulation

Previous studies in β-cells show that, at low glucose, glutamine and leucine stimulate
insulin secretion by fueling the mitochondrial PEP cycle to close K$_{ATP}$ channels, depolarize V$_m$,
and open VDCCs (35). Given that α-cell K$_{ATP}$ channels are also regulated by PK (18), we tested
whether the PEP cycle impacts electrical activity similarly in α-cells. K$_{ATP}$ channel activity was
directly measured by patch-clamp using the fluorescent GCaMP6s reporter to positively identify
α-cells. K$_{ATP}$ channel activity was increased by glutamine and reduced by leucine in the control

group, effects that were blunted or blocked by α-cell deletion of PKM1/2 and PCK2 (Figs. 3A
and 3B). Notably, leucine retained some ability to close K_{ATP} channels in the PKM1/2-αKO,
while failing to close K_{ATP} channels in the PCK2-αKO cells. These data suggest that the
remaining α-cell PKL isoform can partially compensate for the loss of PKM1/PKM2, while
cutting off the mitochondrial PEP supply for all three PK isoforms (i.e., by deleting PCK2) is
more severe.

We next tested how PEP cycle affects α-cell V_m using perforated patch-clamp recordings
of surface α-cells in the intact islet. α-Cells were identified by their characteristic I-V curves
(Supplemental Fig. 2). The application of mixed amino acids (QLRA) depolarized α-cell V_m by
4.7 mV on average in control islets (Fig. 3C). Relative to controls, the amino acid-stimulated
increase in ΔV_m trended upward in PKM1/2-αKO α-cells, but only reached significance in the
PCK2-αKO (Fig. 3C,D). Given the high degree of α-cell variability, and the low throughput
inherent to the patch-clamp approach, we performed orthogonal measurements using JEDI-2P
voltage sensors (37) to record ΔV_m from hundreds of α-cells. In these assays, we isolated islets
from PKM1/2-αKO and PCK2-αKO mice without GCaMP6s. Voltage-sensitive JEDI-2P
biosensors (37) were expressed in α-cells using Cre-dependent adenovirus. Islets from PKM1/2-
αKO and PCK2-αKO mice were measured simultaneously with their controls using barcoding
with near-IR dyes. Confirming the patch-clamp studies, the α-cell ΔV_m response to amino acids
was augmented by deletion of PCK2, while α-cell deletion of PKM1/2 had no significant effect
(Fig. 3E,F).

    Figure 3

310
The α-cell PEP cycle closes K_{ATP} channels and restricts α-cell membrane potential in response to
amino acid stimulation. *A-B*: Example traces and power analysis of K_{ATP} channel activity in
response to 0.5 mM glutamine (0.5Q) and 1.5 mM leucine (1.5L) in dispersed α-cells from
PKM1/2-αKO (A) and PCK2-αKO (B) and controls. *C-D*: Example traces and membrane
potential (ΔV_m) measured by patch-clamp in response 2 mM glucose and amino acids (QLRA;
0.6 mM glutamine, 0.5 mM leucine, 0.2 mM arginine, 2 mM and alanine) in intact islets from
PKM1/2-αKO (*C*) and PCK2-αKO mice (*D*) as indicated. Data represent 12-18 cells from 3 mice
per group. *E-F*: Average normalized membrane potential (ΔV_m) measured by JEDI-2P voltage
sensors in response to 2 mM glucose and QLRA in islets from PKM1/2-αKO (*E*) and PCK2-
αKO mice (*F*) as indicated. Data represent 64-92 islets from 3 mice imaged per protocol. Data
are shown as mean ± SEM for each condition compared with control by either by one way
ANOVA (*A*, *B*) or Student's *t* test (*C-F*). Controls were averaged in A-B and C-D. *$P < 0.05$, ns,
not significant.

The PEP cycle suppresses α-cell Ca^{2+} in response to amino acid stimulation without
impacting glucagon release

Our electrophysiological analyses of K_{ATP} channels and V_m suggested a mechanism by
which PKM1/2 and PCK2 might regulate α-cell Ca^{2+}. Consistently, the application of small
molecule PK activators (10 μM TEPP-46), which allosterically activate PKM2 and PKL to drive
the PEP cycle (34,36,38), nearly abolished the Ca^{2+} response to an amino acid ramp (Fig. 4A). To
confirm that this effect was due to a direct effect on the α-cell, we applied the same amino acid
ramp to islets isolated from PKM1/2-αKO and PCK2-αKO mice. Control islets were again
measured simultaneously using islet barcoding. Elevated α-cell Ca^{2+} was observed in response to
a range of amino acid levels in both PKM1/2-αKO and PCK2-αKO islets relative to controls
(Fig. 4B,C). As in the electrophysiological assays, α-cell deletion of PCK2 had a stronger effect
on Ca^{2+} levels than deletion of PKM1/2. Taking advantage of this phenotype, we used PCK2-
αKO islets to test effects of individual amino acids on Ca^{2+} levels (Supplemental Fig. 3). Alanine
and arginine were able to stimulate Ca^{2+} influx regardless of whether glutamine and leucine are
present, as expected for their direct role in controlling V_m. However, the Ca^{2+} phenotype of the
PCK2-αKO islets was lost in the absence of glutamine, consistent with its role in providing
anaplerotic glutamate carbons that drive oxaloacetate formation for PCK2. These data indicate
that the α-cell PEP cycle cannot function in experiments lacking glutamine.

We next investigated whether Ca^{2+} regulation by the PEP cycle impacts glucagon
secretion using islet perfusion assays. Amino acids and GIP were sequentially applied under low
glucose conditions to maximally stimulate glucagon release. However, we observed no
difference in glucagon release upon PK activation with TEPP-46 (Fig. 4D), or upon α-cell
deletion of PKM1/2 or PCK2 (Fig. 4E,F). Similar results were obtained in PCK2-αKO islets and
controls were stimulated with alanine and GIP in the absence of leucine (Supplemental Fig. 4).
The enhanced intracellular Ca^{2+} levels observed in α-cells lacking PKM1/2 and PCK2 is
consistent with the permissive role of Ca^{2+} in controlling glucagon release. The ability of PK
activation to suppress amino acid stimulated Ca^{2+} influx without impacting secretion not only
indicates that Ca^{2+} influx at 2 mM glucose is sufficient to trigger glucagon secretion, but that
amino acid stimulation of Ca^{2+} influx has no bearing on secretion.

Changes in cytosolic Ca^{2+} can also be modulated by Ca^{2+} extrusion. To test for a role of
the PEP cycle downstream of V_m depolarization, we performed experiments in the presence of
the K_{ATP} channel opener diazoxide (200 μM) and then triggered Ca^{2+} influx with 30 mM KCl.
Even with V_m clamped, Ca^{2+} reached a higher level in α-cells lacking PKM1/2 or PCK2 (Fig.
4G,H). These findings indicate that the α-cell PEP cycle, in addition to controlling V_m,
homeostatically suppresses intracellular Ca^{2+} levels after glucagon is released.

Figure 4

Pyruvate kinase and PCK2 suppress α-cell Ca^{2+} without impacting glucagon secretion. *A-C*:
Average α-cell Ca^{2+} in response to amino acid ramp (1x QLRA; 0.6 mM glutamine, 0.5 mM
leucine, 0.2 mM arginine, 2 mM and alanine) in islets from α-cell GCaMP6s mice treated with
pyruvate kinase activator (PKa, 10μM TEPP-46), islets from control and PKM1/2-αKO mice
(*B*), or control and PCK2-αKO mice (*C*). Data represent 57-105 islets from 3 mice per group. *D-*
E: Average α-cell Ca^{2+} in response to 200 μM diazoxide followed by 30 mM KCl in intact islets
from control and PKM1/2-αKO (*D*) and PCK2-αKO mice (*E*). Data represent 904-1194 α-cells
from 4 mice. *F-H*: Glucagon secretion in response to QLRA, GIP (10 nM) in wild-type mice
treated with PKa (*F*), control and PKM1/2-αKO mice (*G*), or control and PCK2-αKO mice (*H*).
Data represent 6 chambers from 6 mice per condition with 80–100 mouse islets per chamber.
Data are shown as mean ± SEM with AUC for each condition compared with control by
Student's *t* test. **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns, not significant.

Discussion

Our data provide strong support for the concept of α-cells as amino acids sensors (3). We
confirmed that alanine and arginine stimulate Ca^{2+} influx and glucagon secretion at physiological
concentrations. Glutamine and leucine, while having only a small effect on Ca^{2+} at low glucose,
strongly antagonized alanine/arginine-stimulated V_m depolarization and Ca^{2+} influx. We used
three different models to show that glutamine and leucine suppress Ca^{2+} via α-cell PEP cycle,
including pharmacological PK activators and α-cell specific deletion of PKM1/2 and PCK2.
Strikingly, all three of these manipulations modulated Ca^{2+} without changing glucagon secretion.
Thus, the glucagon released in response to amino acids, including electrogenic alanine and
arginine, is not dependent on their ability to stimulate Ca^{2+} influx.

At the mechanistic level, we found that both PCK2 and PK are required for
glutamine/leucine to close K_{ATP} channels, mirroring findings in β-cells (18,34–36), but the
functional outcomes are different. While β-cells use the PEP cycle to initiate secretion, α-cells
use it primarily to modulate Ca^{2+} dynamics. Specifically, we provided genetic evidence that PK is
involved in suppressing intracellular Ca^{2+} levels in response to mitochondrial fuels. Part of this
effect may arise from the ability of PK to close K_{ATP} channels and suppress membrane
depolarization. These findings are consistent with our prior report indicating that a glycolytic
metabolon is present on the plasma membrane of islet cells, and regulates α-cell K_{ATP} channels
via the ATP/ADP ratio (18). Similar to glycolysis, mitochondrial PCK2 can generate PEP in
response to glutamine/leucine. Deletion of PCK2 or PKM1/2 in α-cells blocked K_{ATP} closure in
response to glutamine/leucine, indicating a conserved pathway in both α and β-cells (35). Of
course, K_{ATP} closure has different effects on electrical excitability in α- and β-cells (11). In α-
cells, we observed that both membrane potential and Ca^{2+} levels increased in response to amino
acid stimulation in the PCK2 and PKM1/2 deletion models. In addition to K_{ATP} channels, there
are other potential mechanisms of V_m regulation that we did not explore, such as metabolic
control of α-cell Na^+/K^+ ATPases (39), which facilitate V_m hyperpolarization in β-cells (40).

Several factors may explain why elevated Ca^{2+} in our knockout models failed to enhance
secretion. First is that Ca^{2+} is only permissive for hormone secretion in islet cells (6,41). From
this perspective, it makes sense that elevated Ca^{2+} in the PCK2-αKO and PKM1/2-αKO has little
impact on glucagon secretion, particularly in the presence of arginine and alanine that strongly
stimulate Ca^{2+}. Previous findings show that glucagon release is largely controlled by mouse α-
cell P/Q-type Ca^{2+} channels, although they only account for ~20% of Ca^{2+} current (10). Second,
since glucagon release is not changed by PK or the PEP cycle, it is possible that K_{ATP} channels
primarily impact L-type VDCC, as in β-cells. In any case, the field must look beyond Ca^{2+} to
understand amino acid regulation of glucagon secretion. The glucagonostatic effect of leucine is
both Ca^{2+}- and PEP cycle-independent, and may be primarily mediated by its ability to reduce
cAMP (31). Of note, gain-of-function mutations in glutamate dehydrogenase cause
hyperinsulinemic hypoglycemia in response to protein rich meals (42). We believe this effect is
due to glutamine carbons funneling through PCK2 and PK to close K_{ATP} channels (43). However,
in α-cells, a change in glucagon secretion is not observed even when flux through PEP cycle is
increased, potentially explaining why glutamate dehydrogenase mutations do not impact
glucagon secretion.

Importantly, the elevated levels of Ca^{2+} in the knockout models were also observed in the
presence of diazoxide/KCl, which stimulates Ca^{2+} influx independently of K_{ATP} closure. These
findings indicate that the PEP cycle restores intracellular Ca^{2+} levels after depolarization. In cell

lines, PK has been previously shown to play a role in ER Ca^{2+} storage (44). At low glucose, and in the absence of fatty acids (17), glutamine/leucine are key mitochondrial substrates. Defects in metabolism induced by α-cell PKM1/2 and PCK2 deletion could therefore impair Ca^{2+} extrusion. Mitochondria are also important for Ca^{2+} sequestration (45). In β-cells, the PEP cycle has been shown to hyperpolarize the mitochondrial membrane as PK lowers ADP (i.e., ADP privation) (34), and could affect mitochondrial Ca^{2+} handling. Further studies are needed to understand how amino acids affect organellar Ca^{2+} handling. In the meantime, our work shows that α-cells use the PEP cycle not as a direct regulator of secretion, but rather to limit Ca^{2+} rise during amino acid stimulation.

Article Information

Acknowledgements. The authors thank the University of Wisconsin Carbone Cancer Center Flow Lab for assistance with cell sorting. The authors also thank Matthew Flowers from Dawn Davis lab from University of Wisconsin-Madison for assistance with qPCR. Graphics were created using BioRender.com.

Funding. The Merrins laboratory gratefully acknowledges support from the NIH/NIDDK (R01DK113103 and R01DK127637 to M.J.M.) and the United States Department of Veterans Affairs Biomedical Laboratory Research and Development Service (I01BX005113 to M.J.M.). The Hughes laboratory gratefully acknowledges support from the NIH/NIDDK (R01DK138974 and R01DK140365 to J.W.H.).

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Authors contributions. E.J., H.R.F., E.P., S.M.H., X.D, and J.W.H. designed and performed experimental work and statistical analysis. E.J. and M.J.M. wrote and edited the manuscript. M.J.M. supervised the work.

Supplementary data

Supplemental Fig. 1. Leucine stimulates insulin secretion at low glucose. Insulin secretion from mouse islets measured as in Fig. 1I-J. Data are shown as mean ± SEM with AUC for each condition compared with control by Student's *t* test. **$P < 0.01$.

Supplemental Figure 2. I–V curves used for α-cell identification in intact islets. Example traces and I-V curves measured from surface α-cells (*A*) and β-cells (*B*) in bath solution containing 2 mM glucose and 0.5 mM glutamine. Cells were depolarized from a holding potential of -70 mV to voltages between −60 and +40 mV in 10 mV increments. Data represent 12-18 cells from 3 mice per group as in Figure 3*C-D*.

Supplemental Figure 3. The α-cell Ca^{2+} PEP cycle requires glutamine. *A-D*: Averaged Ca^{2+} from control and PCK2-αKO mice in response to 2 mM glucose and combinations of amino acids as indicated. Data represent 42-55 islets from 3 mice per group. Data are shown as mean ± SEM with AUC for each condition compared with control by Student's *t* test. **P < 0.01, ****P < 0.0001, ns, not significant.

Supplemental Figure 4. The α-cell Ca^{2+} PEP cycle is not required for glucagon secretion stimulated by alanine and GIP. Glucagon secretion control and PCK2-αKO islets in response to

alanine (2 mM) and GIP (10 nM GIP). Data represent 3-6 chambers from 3-5 mice per condition with 80–100 mouse islets per chamber. Data are shown as mean ± SEM with AUC for each condition compared with control by Student's *t* test. ns, not significant.

Supplemental Figure 5. Culturing mouse islets 4-hydroxytamoxifen does not impact glucagon and insulin secretion. Hormone secretion measured from mouse islets after 3-day treatment with 1, 10, and 100 nM 4-hydroxytamoxifen (4-HT) in the culture media. Islets were measured in the presence of glucose (2G: 2mM glucose; 10G: 10mM glucose) and amino acids as indicated (QLRA, in mM: 0.6 mM glutamine, 0.5 mM leucine, 0.2 mM arginine, 2 mM and alanine). Data represent the average of 3 chambers from 3 mice per condition with 80 mouse islets per chamber. Data are shown as mean ± SEM with AUC for each condition compared with control by 2-way ANOVA. ns, not significant.

Reference

1. Capozzi ME, D'Alessio DA, Campbell JE. The past, present, and future physiology and pharmacology of glucagon. Cell Metabolism. 2022 Nov 1;34(11):1654–74.

2. Armour SL, Stanley JE, Cantley J, Dean ED, Knudsen JG. Metabolic regulation of glucagon secretion. J Endocrinol. 2023 Jul 1;JOE-23-0081.

3. Dean ED. A Primary Role for α-Cells as Amino Acid Sensors. Diabetes. 2020 Apr;69(4):542–9.

4. Hamilton A, Eliasson L, Knudsen JG. Amino acids and the changing face of the α-cell. Peptides. 2023 Aug;166:171039.

5. Gilon P. The Role of α-Cells in Islet Function and Glucose Homeostasis in Health and Type 2 Diabetes. J Mol Biol. 2020 06;432(5):1367–94.

6. Le Marchand SJ, Piston DW. Glucose Decouples Intracellular Ca2+ Activity from Glucagon Secretion in Mouse Pancreatic Islet Alpha-Cells. Kulkarni R, editor. PLoS ONE. 2012 Oct 15;7(10):e47084.

7. Onyango AN. Mechanisms of the Regulation and Dysregulation of Glucagon Secretion. Oxid Med Cell Longev. 2020 Jul 21;2020:3089139.

8. Dai XQ, Camunas-Soler J, Briant LJB, Santos T, Spigelman AF, Walker EM, et al. Heterogenous impairment of α cell function in type 2 diabetes is linked to cell maturation state. Cell Metabolism. 2022 Feb;34(2):256-268.e5.

9. Acreman S, Ma J, Denwood G, Gao R, Tarasov A, Rorsman P, et al. The endoplasmic
reticulum plays a key role in α-cell intracellular Ca2+ dynamics and glucose-regulated
glucagon secretion in mouse islets. iScience. 2024 Apr 5;27(5):109665.

10. Zhang Q, Ramracheya R, Lahmann C, Tarasov A, Bengtsson M, Braha O, et al. Role of
KATP Channels in Glucose-Regulated Glucagon Secretion and Impaired Counterregulation
in Type 2 Diabetes. Cell Metabolism. 2013 Dec;18(6):871–82.

11. Zhang Q, Dou H, Rorsman P. 'Resistance is futile?' – paradoxical inhibitory effects of K $_{ATP}$
channel closure in glucagon☐secreting α☐cells. The Journal of Physiology. 2020
Nov;598(21):4765–80.

12. Gromada J, Ma X, Høy M, Bokvist K, Salehi A, Berggren PO, et al. ATP-Sensitive K+
Channel–Dependent Regulation of Glucagon Release and Electrical Activity by Glucose in
Wild-Type and *SUR1* –/– Mouse α-Cells. Diabetes. 2004 Dec 1;53(suppl_3):S181–9.

13. Briant LJB, Dodd MS, Chibalina MV, Rorsman NJG, Johnson PRV, Carmeliet P, et al.
CPT1a-Dependent Long-Chain Fatty Acid Oxidation Contributes to Maintaining Glucagon
Secretion from Pancreatic Islets. Cell Reports. 2018 Jun;23(11):3300–11.

14. Armour SL, Frueh A, Chibalina MV, Dou H, Argemi-Muntadas L, Hamiltion A, et al.
Glucose controls glucagon secretion by regulating fatty acid oxidation in pancreatic alpha
cells. Diabetes. 2023 Jul 26;db230056.

15. Ramracheya R, Ward C, Shigeto M, Walker JN, Amisten S, Zhang Q, et al. Membrane
potential-dependent inactivation of voltage-gated ion channels in alpha-cells inhibits
glucagon secretion from human islets. Diabetes. 2010 Sep;59(9):2198–208.

16. MacDonald PE, Marinis YZD, Ramracheya R, Salehi A, Ma X, Johnson PRV, et al. A KATP
Channel-Dependent Pathway within α Cells Regulates Glucagon Release from Both Rodent
and Human Islets of Langerhans. Vidal-Puig AJ, editor. PLoS Biol. 2007 May 15;5(6):e143.

17. Armour SL, Frueh A, Chibalina MV, Dou H, Argemi-Muntadas L, Hamilton A, et al.
Glucose Controls Glucagon Secretion by Regulating Fatty Acid Oxidation in Pancreatic α-
Cells. Diabetes. 2023 Oct 1;72(10):1446–59.

18. Ho T, Potapenko E, Davis DB, Merrins MJ. A plasma membrane-associated glycolytic
metabolon is functionally coupled to KATP channels in pancreatic α and β cells from humans
and mice. Cell Rep. 2023 Apr 13;42(4):112394.

19. Bahl V, Rifkind R, Waite E, Hamdan Z, May CL, Manduchi E, et al. G6PC2 controls
glucagon secretion by defining the set point for glucose in pancreatic α cells. Sci Transl Med.
2025 Jan;17(779):eadi6148.

20. Bahl V, Lee May C, Perez A, Glaser B, Kaestner KH. Genetic activation of α-cell
glucokinase in mice causes enhanced glucose-suppression of glucagon secretion during
normal and diabetic states. Mol Metab. 2021 Jul;49:101193.

21. Cabrera O, Jacques-Silva MC, Speier S, Yang SN, Köhler M, Fachado A, et al. Glutamate Is
a Positive Autocrine Signal for Glucagon Release. Cell Metabolism. 2008 Jun;7(6):545–54.

22. Li C, Liu C, Nissim I, Chen J, Chen P, Doliba N, et al. Regulation of glucagon secretion in
normal and diabetic human islets by γ-hydroxybutyrate and glycine. J Biol Chem. 2013 Feb
8;288(6):3938–51.

23. Zhang Z, Ye WW, Piro AL, Wang DS, Untereiner A, Lyons SA, et al. Glycine receptor
activation promotes pancreatic islet cell proliferation via the PI3K/mTORC1/p70S6K
pathway. JCI Insight. 10(8):e178754.

24. Malaisse WJ, Blachier F, Mourtada A, Camara J, Albor A, Valverde I, et al. Stimulus-
secretion coupling of arginine-induced insulin release. Metabolism of l-arginine and l-
ornithine in pancreatic islets. Biochimica et Biophysica Acta (BBA) - Molecular Cell
Research. 1989 Sep;1013(2):133–43.

25. Spears E, Stanley JE, Shou M, Yin L, Li X, Dai C, et al. Pancreatic islet α cell function and
proliferation requires the arginine transporter SLC7A2 [Internet]. bioRxiv; 2023 [cited 2023
Aug 24]. p. 2023.08.10.552656. Available from:
https://www.biorxiv.org/content/10.1101/2023.08.10.552656v1

26. El K, Gray SM, Capozzi ME, Knuth ER, Jin E, Svendsen B, et al. GIP mediates the incretin
effect and glucose tolerance by dual actions on α cells and β cells. Sci Adv. 2021 Mar
10;7(11):eabf1948.

27. Simmons TW, Moseley RH, Boyer JL, Ballatori N. Modulation of Na+/alanine cotransport in
liver sinusoidal membrane vesicles by internal divalent cations. Biochimica et Biophysica
Acta (BBA) - Biomembranes. 1990 Apr 30;1023(3):462–8.

28. Rocha DM, Faloona GR, Unger RH. Glucagon-stimulating activity of 20 amino acids in
dogs. J Clin Invest. 1972 Sep 1;51(9):2346–51.

29. Li C, Najafi H, Daikhin Y, Nissim IB, Collins HW, Yudkoff M, et al. Regulation of leucine-
stimulated insulin secretion and glutamine metabolism in isolated rat islets. J Biol Chem.
2003 Jan 31;278(5):2853–8.

30. Sener A, Malaisse WJ. L-leucine and a nonmetabolized analogue activate pancreatic islet
glutamate dehydrogenase. Nature. 1980 Nov;288(5787):187–9.

31. Knuth ER, Foster HR, Jin E, Ekstrand MH, Knudsen JG, Merrins MJ. Leucine Suppresses α-
Cell cAMP and Glucagon Secretion via a Combination of Cell-Intrinsic and Islet Paracrine
Signaling. Diabetes. 2024 Jun 13;73(9):1426–39.

32. Dean ED, Li M, Prasad N, Wisniewski SN, Von Deylen A, Spaeth J, et al. Interrupted
Glucagon Signaling Reveals Hepatic α Cell Axis and Role for L-Glutamine in α Cell
Proliferation. Cell Metabolism. 2017 Jun;25(6):1362-1373.e5.

33. Armour SL, Stanley JE, Cantley J, Dean ED, Knudsen JG. Metabolic regulation of glucagon
secretion. J Endocrinol. 2023 Sep 1;259(1):e230081.

34. Lewandowski SL, Cardone RL, Foster HR, Ho T, Potapenko E, Poudel C, et al. Pyruvate
Kinase Controls Signal Strength in the Insulin Secretory Pathway. Cell Metabolism. 2020
Nov;32(5):736-750.e5.

35. Foster HR, Ho T, Potapenko E, Sdao SM, Huang SM, Lewandowski SL, et al. β-cell deletion
of the PKm1 and PKm2 isoforms of pyruvate kinase in mice reveals their essential role as
nutrient sensors for the KATP channel. Elife. 2022 Aug 23;11:e79422.

36. Abulizi A, Cardone RL, Stark R, Lewandowski SL, Zhao X, Hillion J, et al. Multi-Tissue
Acceleration of the Mitochondrial Phosphoenolpyruvate Cycle Improves Whole-Body
Metabolic Health. Cell Metab. 2020 Nov 3;32(5):751-766.e11.

37. Liu Z, Lu X, Villette V, Gou Y, Colbert KL, Lai S, et al. Sustained deep-tissue voltage
recording using a fast indicator evolved for two-photon microscopy. Cell. 2022 Sep
1;185(18):3408-3425.e29.

38. Anastasiou D, Yu Y, Israelsen WJ, Jiang JK, Boxer MB, Hong BS, et al. Pyruvate kinase M2
activators promote tetramer formation and suppress tumorigenesis. Nat Chem Biol. 2012
Oct;8(10):839–47.

39. Barg S, Galvanovskis J, Göpel SO, Rorsman P, Eliasson L. Tight coupling between electrical
activity and exocytosis in mouse glucagon-secreting alpha-cells. Diabetes. 2000 Sep
1;49(9):1500–10.

40. Dickerson MT, Dadi PK, Zaborska KE, Nakhe AY, Schaub CM, Dobson JR, et al. Gi/o
protein-coupled receptor inhibition of beta-cell electrical excitability and insulin secretion
depends on Na+/K+ ATPase activation. Nat Commun. 2022 Oct 29;13(1):6461.

41. Hughes JW, Ustione A, Lavagnino Z, Piston DW. Regulation of islet glucagon secretion:
Beyond calcium. Diabetes Obes Metab. 2018;20 Suppl 2:127–36.

42. Luczkowska K, Stekelenburg C, Sloan-Béna F, Ranza E, Gastaldi G, Schwitzgebel V, et al.
Hyperinsulinism associated with GLUD1 mutation: allosteric regulation and functional
characterization of p.G446V glutamate dehydrogenase. Hum Genomics. 2020 Mar 6;14:9.

43. Merrins MJ, Kibbey RG. Glucose Regulation of β-Cell KATP Channels: It Is Time for a New
Model! Diabetes. 2024 Jun 1;73(6):856–63.

44. Ho PC, Bihuniak JD, Macintyre AN, Staron M, Liu X, Amezquita R, et al.
Phosphoenolpyruvate Is a Metabolic Checkpoint of Anti-tumor T Cell Responses. Cell. 2015
Sep 10;162(6):1217–28.

45. Griesche N, Sanchez G, Hermans C, Idevall-Hagren O. Cortical mitochondria regulate
insulin secretion by local Ca2+ buffering in rodent beta cells. J Cell Sci. 2019 Apr
30;132(9):jcs228544.

46. Ackermann AM, Zhang J, Heller A, Briker A, Kaestner KH. High-fidelity Glucagon-CreER
mouse line generated by CRISPR-Cas9 assisted gene targeting. Mol Metab. 2017;6(3):236–
44.

47. Merrins MJ, Poudel C, McKenna JP, Ha J, Sherman A, Bertram R, et al. Phase Analysis of
Metabolic Oscillations and Membrane Potential in Pancreatic Islet β-Cells. Biophys J. 2016
Feb 2;110(3):691–9.

611