Cross-platform Hi-C meta-analysis identifies functional insulators that actively block

enhancer-promoter interactions

Jian Cui[1,*], Wanying Xu[1,2,*], Xiuyuan Lang[1,2,*], Shanshan Zhang[1,2], Leina Lu[1,2], Xiaoxiao Liu[1],

Yan Li[1,#], Fulai Jin[1,3,4,5,#]

[1]Department of Genetics and Genome Sciences, School of Medicine, Case Western Reserve

University, Cleveland, OH 44106, USA

[2]The Biomedical Sciences Training Program (BSTP), School of Medicine, Case Western

Reserve University, Cleveland, Ohio 44106, USA

[3]Department of Computer and Data Sciences, School of Engineering, Case Western Reserve

University, Cleveland, OH 44106, USA

[4]Department of Population and Quantitative Health Sciences, School of Medicine, Case

Western Reserve University, Cleveland, Ohio 44106, USA

[5]Case Comprehensive Cancer Center, Case Western Reserve University, Cleveland, OH

44106, USA

[*]Equal contribution. [#]Co-senior authors.

Correspond with: yxl1379@case.edu (Y.L.), fxj45@case.edu (F.J.)

Summary

CTCF forms the boundaries of topologically associating domains (TADs) but their insulating

function in transcription is controversial. Here we report that functional insulators (FINs) shall be

determined not only by their ability to form static loops or boundaries, but also by the dynamic

context if their flanking sequences form new loops upon boundary loss. We performed a meta-

analysis of nine independent Hi-C and micro-C datasets upon acute CTCF- or cohesin protein-

depletion and found that only at several hundred loci, newly gained enhancers-promoters (E-P)

loops can be reproducibly observed upon CTCF depletion. The new E-P loops are cohesin-

dependent and associated with recurrent gene activation. These findings allow us to map FINs

and define their direct target genes genome-wide. FINs are mostly present in euchromatin, do

not overlap with TAD boundaries, and can be perturbed by *Wapl*-depletion or CTCF-

displacement. We therefore conclude that FINs, but not TAD boundaries, are *bona fide*

insulators.

Introduction

It is generally accepted that TADs confine the communication between genes and enhancers for transcriptional control[1, 2, 3, 4, 5, 6]. Most of the TADs are maintained by CTCF and cohesin, which are also known as insulator proteins. Cohesin is a multimeric complex consisting of *SMC1A*, *SMC3*, *RAD21* and one SA subunit (*STAG1* or *STAG2*). TAD forms when a DNA loop extrudes out of the cohesin complex ring before blocked by CTCF binding sites in convergent orientation, and cohesin serves as an enzymatic engine for this process. Deletion of CTCF or cohesin core subunit RAD21 leads to near complete loss of TADs[7, 8]. Cohesin function is also regulated by cohesin loading complex MAU2/NIPBL[9] and cohesin release factor WAPL[10, 11, 12]: TADs disappear upon loss of NIPBL[13, 14], while loss of WAPL causes bigger loops because cohesin can pass through CTCF blocks due to prolonged loop extrusion[14, 15].

Insulators are canonically defined as the DNA elements that inhibit gene expression when put between enhancers and promoters[16, 17, 18]. The discovery of thousands of CTCF-demarcated TADs immediately suggests a model that CTCF sites at TAD boundaries are enhancer-blocking insulators, which is supported by genetic experiments focusing on selected loci[19, 20, 21, 22, 23]. However, this model is challenged by a genome-wide Hi-C study in mESCs (Nora *et al.*, **Fig. 1a**) which only observed mild transcription changes despite near complete loss of TADs upon CTCF-depletion[7]. This conclusion is reproduced in two other genome-wide studies done with *in situ* Hi-C and micro-C[24, 25] (**Fig. 1a**). Notably, Kubo *et al*. reported a role of CTCF in mediating E-P interactions[24], and Hsieh *et al.* reported short-range cohesin-independent cross-TAD E-P interactions[25]. Taken together, although all these studies questioned the insulating function of TAD boundaries, they did not address the fundamental question about the whereabouts of the real enhancer-blocking insulators.

In this study we demonstrated that "functional insulators" (FINs) can be identified by examining if their flanking regions gain E-P interactions upon CTCF depletion. Challenges exist to map FINs with Hi-C-like genome-wide technologies: (i) most of the public Hi-C datasets do not have billion-scale read depth usually required for loop-level analysis (**Fig. 1a**); (ii) E-P interactions are generally weaker than CTCF loops in Hi-C and therefore harder to detect; (iii) the gain of E-P loop signal can be subtle, requiring robust quantitation of loop strength; (iv) it is difficult to rule out the possibility that some loop changes are secondary effects after prolonged CTCF or cohesin depletion[7, 25]. These challenges can be addressed by comparing independent datasets, but existing CTCF-depletion data are from different Hi-C protocols, which perform very

differently in loop-level analyses with conventional pipelines[26]. We previously developed *HiCorr*

and *DeepLoop* pipelines to allow robust comparison of loop profiles from orthogonal Hi-C and

micro-C platforms, even when sequencing depth is low[27, 28]. Here we use *DeepLoop* to perform

a meta-analysis of orthogonal Hi-C datasets to map recurrent DNA rewiring events upon

depletion of CTCF and other cohesin proteins (**Fig. 1a**), thus generating a reliable genome-wide

map of enhancer-blocking insulators.

Results

DeepLoop **allows cross-platform comparison of independent CTCF- and cohesin-degron**

Hi-C datasets at loop-resolution

We collected nine pairs of independent Hi-C or micro-C datasets from six publications in which

CTCF[7, 24, 25], RAD21[25, 29, 30], or WAPL[25, 29, 31] were acutely depleted in three different mESC lines.

The number of contacts in these Hi-C/micro-C datasets are highly variable ranging from 43

million to ~1 billion. For every target protein, the choice of degron methods (AID or dTAG) and

the timespan of protein degradation (range from 3hr to 3 days) are also variable (**Fig. 1a**).

To assess how these variations complicate cross-platform comparisons. We first performed low

resolution compartment- and TAD-level analysis on the three CTCF-degron datasets with

conventional pipelines. At compartment level (500kb resolution), all datasets display rather

consistent compartment correlation heatmaps, and the compartmental organization is largely

preserved after CTCF depletion (**Supplementary Fig. 1a-c**). When plotting ICE-normalized

contact matrices[32] at 40kb resolution followed by DI (directionality index)-based TAD analyses,

all three studies consistently show similar TAD structures and a massive loss of TADs upon

CTCF depletion despite a trend that micro-C detected more bins with stronger directionality

biases (**Supplementary Fig. 1d-f**). These observations confirmed the findings from the original

reports and showed that conventional pipelines are overall robust across different Hi-C

platforms at low-resolution.

However, Hi-C analysis at high resolution is severely affected by data sparsity and protocol

biases[33, 34], thus the robustness of cross-platform comparison is highly dependent on the choice

of analytical pipelines. We generated 5kb resolution contact heatmaps (each pixel in a heatmap

represents a pair of 5kb bins) using conventional *ICE*[32] (with or without distance correction),

HiCorr[34], and *DeepLoop*[28], then used scatter plots to compare control cell heatmaps from

independent experiments to assess the consistency of different pipelines in analyzing

orthogonal Hi-C platforms (**Supplementary Fig. 2**). Here we only compare control cell

heatmaps because before protein depletion, the control cells from all studies are expected to be

at similar biological states. Protein-depletion cells from independent studies, even if they

depleted the same protein, can be still biologically diverse due to many variations such as time-

span of protein depletion, incomplete protein depletion, loss of stemness, *etc.* We found that

although conventional ICE pipelines show poor cross-platform consistency with R-square values

often close to zero, *HiCorr* and *DeepLoop* clearly improved the consistency with much higher

cross-platform R-square values (**Supplementary Fig. 2**, summarized in **Fig. 1b**).

We also use scatter plots to compare the heatmaps before and after protein depletion at pixel-

by-pixel level and interestingly, found that only HiCorr and DeepLoop heatmaps show

recognizable populations of loop pixels that retain their strength upon CTCF- or Rad21-

depletion (**Supplementary Fig. 2**), suggesting their robustness in revealing the unchanged loop

pixels that are masked by noises in the other pipelines. To further evaluate the cross-platform

consistency when applying different pipelines to analyze dynamic loops, we generated

differential heatmaps by subtracting each pair of 5kb-resolution contact heatmaps (depletion –

control), then computed the correlations between the differential heatmaps from all nine pairs of

degron experiments (**Fig. 1c**, **Supplementary Fig. 3**). Only the matrices from *HiCorr* and

DeepLoop clustered CTCF- and Rad21-degron datasets together and *Wapl*-degron datasets

are separated out (**Fig. 1c**, **Supplementary Fig. 3**). This is consistent with the knowledge that

depletion of *CTCF* or Rad21, but not Wapl, causes massive loss of TADs and CTCF loops.

Taking together, these results support the robustness of *DeepLoop* in cross-platform analysis

regardless of read depth[28].

Hi-C meta-analysis reveals recurrent E-P loop-gaining events upon CTCF depletion

We first focused on the CTCF-degron Hi-C datasets. In the example of **Fig. 1d** (top two rows),

with conventional ICE-normalization, we observed dampened TAD structure at 20kb resolution

as the original study reported[7], but at 5kb resolution, loop signal from ICE-normalized heatmaps

are weak and often invisible. From the *DeepLoop* enhanced heatmaps, we can consistently

observe multiple CTCF loops that appear in all control cells heatmaps but are substantially

weakened upon CTCF depletion. A notable change in this region is the gain of long-range loops

(**Fig. 1d**, second row, loop pixels in dashed circle) connecting *Notch1* gene to a distal enhancer

region with multiple H3K27ac peaks (**Fig. 1e**). Consistent with gain of E-P loop, *Notch1* is

upregulated upon CTCF-depletion (**Fig. 1e**). The gain of loop signal at this locus is also

observable in *Wapl*-depleted cells (**Fig. 1d**, bottom two rows) and two other CTCF-degron

studies (**Supp. Fig. 4**), but not in the *Rad21*-depleted cells (**Fig. 1d**, 3rd and 4th rows, **Supp. Fig.**

4g). These observations suggest that the two CTCF loop anchors at the TAD boundary region

(**Fig. 1d**, highlighted in pink) are blocking the *Notch1*-enhancer interaction. This example also

demonstrated that *DeepLoop* may detect recurrent insulating events from CTCF-degron Hi-C

data without the need for ultra-high sequencing depth.

To perform a genome-wide analysis of insulating events, we call 300K~500K loop pixels from

each CTCF-degron dataset and use the pixel intensity in *DeepLoop* heatmaps to measure the

loop strength (**Fig. 2a, Supp. Fig. 5**). Notably, *DeepLoop* outputs are enhanced from ratio

(obs/exp) heatmaps instead of normalized contact frequency used by ICE/KR. We previously

showed that this approach promises cross-platform convergence between orthogonal Hi-C and

micro-C data[28].

In all CTCF-degron studies, the scatter plots are clearly down-skewed reflecting a massive loss

of chromatin loops upon CTCF deletion (**Fig. 2a, Supp. Fig. 5a, d**, left panels). Using a simple

fold-change cutoff, we called loop pixels that lose strength ("lost"), retain strength ("retained"), or

gain strength ("gained") from each dataset (**Fig. 2a, Supp. Fig. 5a, d**, middle panels). We

further classified the loop pixels based on their epigenetic marks (**Methods**, examples of loop

classification in **Supp. Fig. 6**). "CTCF-only" loop pixels, defined as those with at least one

anchor overlapping a CTCF peak, were predominantly found among the "lost" pixels. In

comparison, "retained" and "gained" loop pixels are enriched with "CTCF-EP" or "EP-only" loops

characterized by promoter (H3K4me3) or enhancer (H3K27ac) marks at one or both anchors

(**Fig. 2b, Supp Fig. 5b, e**). Interestingly, in all three studies the "gained" loops are much larger

than the "retained" loops (**Fig. 2a, Supp. Fig. 5a, d**, right panels); we hypothesized that this is

because the "gained" E-P loops form through cohesin loop extrusion (more discussions below).

We also examine the robustness of the "lost", "retained", and "gained" pixels across

independent Hi-C/micro-C studies. As expected, the "lost" pixels from any study generally also

lose strength in the other two studies (**Fig. 2c, Supp Fig. 5c,f**, first row). However, the

consistency of "retained" loop pixels is study specific: (i) although the "retained" loop pixels from

Nora *et al.* are mostly also "retained" in Kubo *et al.*, some of them lose strength upon CTCF

degradation in Hsieh *et al.* (**Fig. 2c**, second row); (ii) most of the "retained" loop pixels from

Kubo *et al.* lose strength in the other two studies (**Supp. Fig. 5c**, second row); (iii) the "retained"

loops from Hsieh *et al.* remain "retained" in both Nora *et al.* and Kubo *et al.* (**Supp. Fig. 5f**,

second row). Since incomplete CTCF depletion causes the leftover CTCF loops[7], these

observations suggest that CTCF depletion in Kubo *et al.* may be less complete than the other

two studies. On the other hand, Hsieh *et al.* appear to have achieved better CTCF depletion,

perhaps due to its use of an N-terminal degron system (**Fig. 1a**), although platform-specific

biases, differences in assay sensitivity and sequencing depth may also explain some of the

discrepancies between studies.

Interestingly, the "gained" pixels from any one of the three studies generally also gain strength

in the other two studies (**Fig. 2c, Supp Fig. 5c,f,** third row). The only exception is that some

"gained" loop pixels from Kubo *et al.* lose strength in Hsieh *et al.* (**Supp Fig. 5c**). Again, this is

likely because Kubo *et al.* retains more CTCF loops due to incomplete CTCF degradation.

Nevertheless, most of the "gained" pixels from Nora *et al.* and Hsieh *et al.* do gain strength in

Kubo *et al.* (**Fig. 2c, Supp Fig. 5f**), supporting the consistency between the three studies.

Notably, E-P loop gaining events are largely neglected in the original Hi-C/micro-C studies

because the changes are often too subtle to distinguish from background with conventional

analytical tool (*e.g.*, **Fig. 1d**). Our meta-analyses prove that although there are only a small

number of newly gained E-P loops, they are reliable consequences of CTCF-depletion; affected

genes such as *Notch1* in **Fig. 1** are putative target genes of CTCF-mediated insulation.

Hi-C Meta-analyses of *Rad21*- and *Wapl*-depletion data distinguish consensus loop

regulatory mechanisms from system variations

We also performed meta-analyses of the *Rad21*- and *Wapl*-depletion Hi-C datasets. Ablation of

Rad21 also causes massive loss of CTCF loops, while the "retained" and "gained" loops are

also enriched with E-P loops (**Supp. Fig. 7a-i**). However, there are important differences

between *Rad21*- and *CTCF*-depleted cells. (i) In *Rad21*-depleted cells, the "retained" and

"gained" loop pixels (except the "gained" loop pixels in Rhodes *et al.*) are much smaller (mostly

within 100kb) than those in *CTCF*-depleted cells, indicating that cohesin-independent loops are

short-range (**Supp. Fig. 7a, d, g**, right panels). One alternative explanation is that the greatly

reduced pool of cohesin only allows rare and short-lived extrusion events, resulting in smaller

loops. (ii) Unlike in CTCF-depleted cells, the "gained" loop pixels are NOT recurrent between

the three *Rad21*-degon datasets (**Supp. Fig. 7c, f, i**, last row). (iii) The "retained" and "gained"

loop pixels in *Rad21*-depleted cells (except the "gained" loop pixels in Rhodes *et al.*) are more

enriched with H3K36me3 mark, suggesting transcription associated chromatin interactions

(**Supp. Fig. 7b, e, h**).

The "gained" loops in *Rad21*-depleted cells from Rhodes *et al*. are enriched with H3K27me3

and tends to be larger in size (**Supp. Fig. 7a, b**, right panels). These observations are

consistent with the conclusion from the original study that cohesin removal enhances long-range

interactions between a subset of polycomb-occupied regions[30]. However, these features (the

larger loop size and the enrichment with H3K27me3) are specific to "gained" loops in Rhodes *et*

al. but not observed in the other two *Rad21*-degron studies (**Supp. Fig. 7d, e, g, h**, right

panels). We manually examined the "gained" H3K27me3 loops in Rhodes *et al.* and found that

in almost all cases, the long-range H3K27me3 loops are already present in the control cells

from Kriz *et al.* and Hsieh *et al.* before *Rad21* depletion (two examples are shown in **Supp. Fig.**

8). Therefore, at least in the control mESCs from Kriz *et al.* and Hsieh *et al.*, cohesin does not

prevent the formation of long-range H3K27me3 loops.

In contrast, ablation of cohesin unloading factor *Wapl* generates up-skewed scatterplots in all

three studies (**Supp. Fig. 7j, m, p**, left panels). The skewness of Hsieh *et al.* is less apparent,

probably because Hsieh *et al.* performed *Wapl* depletion for much shorter time (3hrs) than Liu *et*

al. (24hrs) and Kriz *et al*. (72hrs) (**Fig. 1a**). In all *Wapl*-degron studies, the "lost", "retained", and

"gained" loops are almost equally enriched with CTCF and E-P loops (**Supp. Fig. 7k, n, q**).

Consistent with the model that *Wapl*-depletion stabilizes chromatin-bound cohesin and promote

loop extrusion[14, 15], the "gained" loops are significantly bigger than "lost" and "retained" loops in

all three studies (**Supp. Fig. 7j, m, p**, right panels). The "lost", "retained", and "gained" loop

pixels are generally consistent between studies (**Supp. Fig. 7l, o, r**) despite some discrepancy

for "lost" loop pixels from Hsieh *et al.* (**Supp. Fig. 7r**), again likely because Hsieh *et al.* only

depleted *Wapl* for a short period of time.

Newly gained E-P loops in CTCF-depleted cells are cohesin dependent.

We next take the "lost", "retained", and "gained" loop pixels from *CTCF*-degron studies and

examine how they change in *Rad21*- and *Wapl*-degron data. Firstly, nearly all "lost" loops in

CTCF-depleted cells also disappear in *Rad21*-depleted cells (**Fig. 2d**, first row, **Supp Fig. 9a,b**,

first row), suggesting that most of the CTCF loops are cohesin-dependence. Secondly, a subset

of the "retained" loops in CTCF-depleted cells are also "retained" upon *Rad21*-depletion (**Fig.**

2d, second row, **Supp Fig. 9a, b**, second row), suggesting cohesin-independence. The other

"retained" loops in CTCF-depleted cells lose strength upon *Rad21*-depletion likely because they

are: (i) leftover CTCF loops due to incomplete CTCF-depletion, or (ii) non-CTCF loops formed

via cohesin-dependent mechanisms. Interestingly, most of the "retained" loops in CTCF-

depleted cells from Kubo *et al.* are lost upon *Rad21*-depletion (**Supp Fig. 9a**, second row), but

most of the "retained" loops from Hsieh *et al.* are also retained upon *Rad21*-depletion (**Supp**

Fig. 9b, second row), again suggesting that Hsieh *et al.* may have achieved more complete

CTCF-depletion than Kubo *et al.*

Finally, the "gained" loops in CTCF-degron studies, which are weak in both CTCF-degron and

Rad21-degron control mESCs, do not gain strength upon *Rad21*-depletion cells (**Fig. 2d, Supp**

Fig. 9a, b, third row), suggesting that the formation of these "gained" loops requires cohesin. In

contrast, we observed that these "gained" loops in CTCF-depleted cells showed an apparent

upward bias in the scatter plots of *Wapl*-degron studies (**Fig. 2d, Supp Fig. 9a, b**, third row).

This observation is consistent with the example in **Fig. 1d** and further supports a role of cohesin

in forming those "gained" loops. *Wapl*-depletion may perturb the function of insulators by

releasing short-range CTCF loops and allow insulated E-P interactions to skip some insulators.

Define, map, and characterize functional insulators (FINs) in mESC genome

Among the total 8,141 "gained" loop pixels from all three CTCF-degron Hi-C/micro-C datasets,

4,944 pixels are recurrent in at least two studies with a median size ~250kb (**Fig. 3a**). In each

dataset, the "gained" loop pixels can be merged into ~300 "loop-gaining loci" (**Methods**) and

there are substantial overlaps between independent studies: 174 loop-gaining loci are shared by

all three studies and 304 are shared by at least two studies (**Fig. 3b**). We hypothesize that

these recurrent E-P loop-gaining loci are caused by loss of CTCF insulation; indeed, 4,665 out

of the 4,944 loop-gaining pixels (94%) are insulated by at least one CTCF loop in wildtype cells

(**Fig. 3c**). We further curated the reproducible loop-gaining loci based on the types and numbers

of CREs involved (**Fig. 3d**). Nearly all the curated loci (283 of 299) gain E-P loops (2 loci gain

CTCF contacts and the others cannot be clearly classified). We further classified the E-P loop-

gaining loci into three categories (**Fig. 3d**). In *Type I*, one of the two E/P sites is insulated in a

CTCF loop while the other site is not (exampled in **Supp. Fig. 10a**); *Type II* involves at least two

CTCF loops insulating two E/P-sites (exampled in **Fig. 1e, Supp. Fig. 10b**). In contrast, the

gained E-P loops in *Type III* do not have insulating CTCF loops (examled in **Supp. Fig. 10c**).

Type I and *II* represents most of the E-P loop gaining loci (263 out of 283), while only 20 loci

belong to *Type III*, again suggesting that most of the loop gaining loci are due to the removal of

CTCF insulation.

We therefore propose to define "functional insulators" (FINs) as the CTCF binding sites within

the "lost" loop anchors that when bound by CTCF, can actively block the interactions between

flanking *cis*-regulatory elements (CREs) (model *iii* in **Fig. 3e, Methods**). Therefore, FINs can be

found from "loop-gaining loci" and in fact, most "loop-gaining loci" contain at least one FIN, and

we therefore also call them "FIN loci" (**Methods**). The definition of FIN is based on the gain-of-

loop upon loss of insulator function, which is conceptually analogous to the conventional model

to define insulators based on the gain-of-expression upon loss of insulator function (model *i* in

Fig. 3e) but distinct from the TAD-based definition of insulators (model *ii* in **Fig. 3e**), which is a

static definition that does not need the knowledge about the outcomes after loss of insulators.

In **Fig. 3f**, we show a FIN (highlighted in pink) that is not at TAD boundary. In this example we

also included the browser tracks for "insulation score" (InsScore)[35] and "directionality index"

(DI)[1]. TAD boundaries can be defined as bins where InsScore reaches local minimum or

positions where DI changes its sign (**Supp. Fig. 11a**), but the FIN in this example does not fit

into either criterion. There are much fewer FINs than TAD boundaries (**Fig. 3g**) and only a small

percentage of the FINs overlap with TAD boundaries (**Fig. 3h, Supp. Fig. 11b**). These

observations indicate that most FINs are inside TADs and conversely, loss of TAD boundaries

usually does not cause loop formation across their flanking regions. Change of InsScore also

does not predict FINs because although we observed an overall trend of increased InsScore at

FINs upon CTCF depletion, TAD boundaries also show the same trend and cannot be

distinguished from FINs (**Fig. 3i, Supp. Fig. 11c**). Interestingly, most (96%) of the FINs are

within compartment A (**Fig. 3j**), indicating that active E-P blocking mainly happens within

euchromatin region. Regardless, it is important to note that low-resolution 3D genome features,

including compartment and TAD boundaries, lack the precision to reveal the dynamic changes

of loop profiles as we used to define FINs.

Cohesin degron Hi-C data further supports a cohesin-dependent mechanism of FINs. We found

that almost none of the curated loop-gaining events appear in the *Rad21*-degron cells (**Fig. 3d**),

indicating cohesin dependency. Interestingly, a significant fraction of the *Type I* and *II*, but not

the *Type III* "gain of E-P loop" loci, can be also observed in *Wapl*-degron datasets (**Fig. 3d**).

This phenomenon can be explained by previous findings that *Wapl* deletion allows loop

extrusion to skip CTCF blocks[14, 15]. As shown in the carton in **Fig. 3k**, because *Wapl*-depletion

stabilizes chromatin-bound cohesin and leads to longer loop extrusion, previously insulated

enhancers and promoters can interact inside the bigger CTCF loop due to "insulator skipping",

creating the same E-P loop as observed in CTCF-depleted cells. All these results support FINs

as *bona fide* insulators of cohesin-mediated loop extrusion.

To explore which types of enhancers or promoters would gain chromatin loops upon CTCF

depletion, we picked the FIN-blocked ATAC-peaks in mESCs (n = 995) and scan for enriched or

depleted motifs compared to all mESC ATAC-peaks (n = 59,871). Interestingly, the loop-gaining

ATAC-peaks are enriched with many G-rich motifs but depleted with A/T-rich motifs (**Fig. 3l**).

This result is reminiscent of previous reports that G-quadruplexes (G4) structure may regulate

chromatin looping[36, 37]. Notably, many of the putative transcription factors binding to these G-rich

motifs have been reported to regulate loop formation in various cell types, such as Kruppel-like

factors[38, 39, 40], SP1[41], MAZ and several zinc-finger proteins[42, 43, 44]. Therefore, our results

suggest that G-rich *cis*-regulatory elements are more likely to act as new loop anchors upon

CTCF-depletion.

FINs are associated with consistent early transcription responses upon CTCF depletion.

We found that in all CTCF-degron studies FIN proximal regions (+/- 100kb) are 2~3 fold more

enriched with DEGs than TAD-boundary proximal regions, and the "gained" loop anchors are 10

times more enriched with DEGs (**Fig. 4a**). Furthermore, early response DEGs are clearly more

enriched at gained loop anchors (**Fig. 4b**), suggesting that FIN-blocked enhancers and

promoters are the direct targets of insulators. In 16.8% of FIN loci (49 out of 299), we observed

the blocking of 43 super-enhancers (there are 231 super-enhancers in total) (**Fig. 4c**). As

expected, genes gaining super-enhancer interactions are more likely to be upregulated than

genes gaining interactions with regular enhancers (**Fig. 4d**).

Using common cutoffs (2-fold for RNA-seq and 1.5-fold for NET-seq, p<0.05), each of the

CTCF-degron studies identified a few hundred early response DEGs, and the reproducibility

between independent studies is limited due to complex reasons (**Fig. 4e**). Both Nora *et al.* and

Kubo *et al.* performed RNA-seq and the earliest timepoint was 24hr CTCF-depletion, which may

still include late-response DEGs caused by indirect mechanisms (*e.g.*, loss of pluripotency).

Among the two RNA-seq studies, Kubo *et al.* detected fewer DEGs than Nora *et al.*, but 70% of

Kubo *et al.*'s DEGs are also observed in Nora *et al.* (47% of the upregulated DEGs and 81% of

the downregulated DEGs, **Fig. 4e**). This observation is consistent with our discussion above

that CTCF depletion in Kubo *et al.* was less complete. Hsieh *et al.* performed both NET-seq and

RNA-seq after only 3hr CTCF-depletion, which should include fewer late-response genes, but

the short treatment time may also reduce the sensitivity of DEG detection. We use NET-seq

data because Hsieh *et al.* reported that NET-seq has higher sensitivity in identifying DEGs[25] as

it maps newly synthesized RNA, while RNA-seq measures the amount of steady state RNA in

the cells.

We decided to focus on the comparison between Hsieh *et al.* and Nora *et al.* as they represent

a more complete set of CTCF direct targets. We next hypothesize that DEGs near FINs are

more likely to be the direct targets of CTCF-mediated enhancer-promoter insulation. If this

hypothesis is true, DEGs near FINs should be more reproducible between independent studies.

Indeed, the upregulated DEGs near FINs (*i.e.*, at "gained" loop anchors upon CTCF depletion)

are more reproducible, but downregulated DEGs near FINs are less reproducible than controls

(**Fig. 4f**, red bars). In contrast, the "lost" loop anchors does not predict the reproducibility of

DEGs. This observation suggests that transcription activation is the primary consequence of

insulator loss, consistent with the convention concept that insulators block enhancer-mediated

gene activation.

In **Fig. 4g** we plotted 49 upregulated DEGs at FIN loci as putative "direct targets" of insulators

(from Hsieh *et al.* or Nora *et al.*), among them 34 are considered as "high-confidence" targets

supported by both studies. One representative example is shown in **Fig. 4h** and **Supp. Fig.**

12a, in which we observed gain of loop between two putative enhancer clusters upon CTCF

depletion, leading to the upregulation of *Cdiptos* gene. Similarly, we also plotted 37 reproducibly

down-regulated DEGs near FINs supported by both studies (**Fig. 4i**). Strikingly, 18 of the

recurrent down-regulated DEGs are from protocadherin-γ, a well-known insulator loci (**Fig. 4i,j**).

Protocadherin proteins play key roles in the formation of synaptic connections between neurons

and the *Pcdh*-γ genes are distinguished by their variable promoters (exon1): γ*a*'s and γ*b*'s are

stochastically expressed in individual neurons and the γ*c*'s are constitutively expressed[45, 46, 47,]

[48]. In mESCs, the promoters of γ*a*'s and γ*b*'s are occupied by CTCF, H3K4me3, and the

repressive H3K27me3 mark (**Fig. 4j**). A cluster of putative enhancers marked by H3K27ac is

present at the 3' of γ*c* promoters (**Fig. 4j**, highlighted in green) forming loop interactions with the

promoters of γ*a*'s and γ*b*'s (**Fig. 4j**, blue arches), which are restricted by a FIN with multiple

CTCF peaks (**Fig. 4j**, highlighted in pink). Upon CTCF depletion, the H3K27ac-interactions are

weakened, and the *Pcdh-γ* promoter loops extend into a distal H3K27ac-poor region marked by

H3K27me3 (**Fig. 4j**), consistent with the downregulation of γ*a* and γ*b* genes. To verify the

function of the putative FIN, we designed a pair of sgRNAs to delete the four CTCF peaks in the

FIN and confirm the down-regulation of all γ*a* and γ*b* genes (**Fig. 4k-l**). Taking together, these

results support the transcription consequences of FINs and nominate "direct targets" of FINs in

mESCs.

Validate FINs with multiplexed CTCF displacement assay

To validate the *cis*-regulatory functions of FINs, we need to verify if the gained loops are truly

due to the loss of insulation. Because a FIN often involves multiple CTCF binding sites that

contribute to the insulating function collectively, we need to simultaneously perturb multiple

CTCF sites in a relatively big region. To achieve this, we combine a multi-sgRNA delivering

system name CARGO[49] with a dCas9-based CTCF-displacement method[47, 50] to perturb

multiple CTCF binding sites in the same cells. In this approach, 4-10 sgRNAs exactly

overlapping the cognate motifs of multiple CTCF binding sites are expressed from the same

CARGO plasmid. When co-transfected with a dCas9 construct, the sgRNAs can displace CTCF

from genome at multiple target sites simultaneously (**Fig. 5a**).

We used this approach to validate the gain-of-loop events at two FIN loci. The first FIN is close

to *Cdiptos*, a recurrent upregulated DEG (**Fig. 5b**). This example belongs to a simpler type of

FIN in which the newly gained loop (**Fig. 5b**, red arches) is blocked by only one FIN (**Fig. 5c**).

We used one CARGO construct targeting 4 CTCF motifs and confirmed with CUT&RUN that we

successfully weakened the target CTCF peaks but did not affect other CTCF peaks in this

region (**Fig. 5b**, highlighted in pink). We performed 4C-seq and confirmed the gain of long-range

chromatin interactions consistent with Hi-C data (**Fig. 5b**, highlighted in green). We also

validated with RT-qPCR that *Cdiptos* is upregulated upon displacement of target CTCF binding

sites (**Fig. 5d**). The second example is more complex as it may involve more than two FINs

(**Fig. 5e,f**). In this example *Mesp1* lost its loop interaction to a nearby FIN (**Fig. 5e**, blue arches)

and upon CTCF depletion, gained promiscuous distal interactions to a large region that is

~200kb in size (**Fig. 5e**, red arches). For this locus we used one CARGO construct to target 9

CTCF motifs and successfully weakened the target CTCF peaks (**Fig. 5e**, highlighted in pink).

Again, using 4C-seq we confirmed the gain of interactions to the large distal region containing

several clusters of putative enhancers (**Fig. 5e**, highlighted in green), and confirmed the

upregulation of *Mesp1* with RT-qPCR (**Fig. 5g**). Taking together, these experiments verified the

insulating functions of FINs identified from CTCF-degron Hi-C data.

Discussion

In this study we propose to define functional insulators, or FINs, as CTCF binding sites that

actively block the loop formation between flanking *cis*-regulatory elements (**Fig. 3e**). Mapping

FINs requires robust detection of newly gained E-P loops, which is challenging for Hi-C analysis

because E-P loops are usually weak and hard to detect. With a meta-analysis of three

independent CTCF-degron Hi-C datasets, we demonstrate that the subtle gain of loop signal

detected by *DeepLoop* are quantitatively robust across different Hi-C or micro-C datasets,

therefore proves the feasibility to map FINs reliably with Hi-C at intermediate read depth.

The concept of FIN is distinct from the common assumption that TAD boundaries are insulators.

Most TADs disappear upon CTCF depletion, and we typically do not observe gain of new loops

across TAD boundaries. In contrast, most of the FINs are anchors of CTCF loops inside TADs

in control cells, and we call an CTCF loop anchor "FIN" only if its flanking sequences gain new

loops (**Fig. 3e**). Importantly, there are only a few hundred FINs in mESCs, which readily

explains why CTCF-depletion only causes mild changes of transcriptome. It should be also

noted that the gained E-P loops do not always predict significant gene activation due to complex

reasons[51]. Furthermore, target gene expression can be quantitative tuned by multiple enhancers

in an additively fashion, and each single enhancer may only have a small effect size[52].

Nevertheless, we showed that FINs better explain DEGs, especially the recurrent upregulated

DEGs in CTCF-depleted cells. This is consistent with the classic definition of insulators as DNA

elements that repress enhancer-mediated gene activation.

It remains unclear why only a small subset of CTCF loop anchors are FINs, *i.e.*, why is gain-of-

loop only observed at some loci but not others? Our analysis provided some hints. Firstly, the

fact that almost all FINs are in compartment A (euchromatin) suggests their dependency on

surrounding active chromatin. Notably, promoters and enhancers may play key roles in loop

initiation and stalling and are known to co-localize with cohesin and its loading factors[53]. Since

genome compartmentalization is a highly dynamic 3D genome feature, we speculate that FINs

are also likely to be cell type- or cell state-specific. Secondly, meta-analysis with *Rad21*- and

Wapl-degron Hi-C data shows that the newly gained E-P loops in CTCF-depleted cells do not

form in cohesin-depleted cells and are also likely to be governed by cohesin regulators. It is

therefore plausible that local cohesin activity is a key determinant. Finally, *cis*-regulatory

elements at the anchors of "gained loops" are enriched with G-rich TF motifs, suggesting that

some CREs are particularly capable of forming stable long-range regulatory loops but such

function may undergo tight control by insulators. Taking together, all these observations suggest

context-specific roles of FINs, and it will be interesting to further explore how FINs exert their

specific functions in health and diseases.

Methods

Experiments

Cell culture

Mouse embryonic stem cells (E14Tg2a) were cultured in DMEM+Glutamax (ThermoFisher ,

#10566-016) with the addition of 15% Fetal Bovine Serum (ThermoFisher, # SH30071.03),

550μM β-mercaptoethanol (ThermoFisher, #21985-023), 1mM Sodium Pyruvate

(ThermoFisher, #11360-070), 1X non-essential amino acids (ThermoFisher, #11140-50), and

10^4 U of Leukemia inhibitory factor (Millipore, #ESG1107). The cells were maintained at a

density of $0.2-1.5x10^5$ cells/cm^2 and were passaged every 24-48 hours using TrypLE

(ThermoFisher, #12563011) on dishes coated with 0.1% gelatin (Millipore, # ES-006-B) at 37°C

and 5% CO2. The medium was refreshed daily when cells were not being passaged. Regular

checks for mycoplasma infection were performed every 3-4 months, with negative results.

4C-seq

4C-seq was conducted following a published protocol[54]. Briefly, 5 million cells fixed with 2%

formaldehyde, then quenched with 125 mM glycine. The fixed cells were lysed using a cell lysis

buffer containing 50 mM Tris-Cl pH 7.5, 150 mM NaCl, 5 mM EDTA, 0.5% NP-40, 1% Triton X-

100, and 1x protease inhibitor cocktail (Roche, Cat#11873580001) for 20–30 minutes on ice.

Following lysis, nuclei were collected by centrifugation at 2,500g for 5 minutes at 4°C and

washed once with 1x restriction enzyme buffer. The nuclei pellets were resuspended in 1x

restriction enzyme buffer and treated with 0.3% SDS for 1 hour at 37°C with shaking, followed

by an additional hour with 2.5% Triton X-100. Chromatin was then digested overnight with the

appropriate restriction enzyme at the recommended temperature while rotating in an air bath.

The specific restriction enzymes used for each locus are detailed in **Supplemental Table 6**.

After digestion, the enzymes were inactivated by heating at 65°C, and nuclei were ligated with

50 μl of T4 DNA ligase (Invitrogen, Cat# 15224–090) in a 7 ml ligation solution at 16°C

overnight. Reverse cross-linking was performed by treating the samples with proteinase K to

isolate the proximity-ligated DNA. The purified DNA was quantified and subjected to secondary

restriction enzyme digestion using roughly 1 unit of restriction enzyme per 1 μg of DNA at the

recommended temperature overnight. After enzyme inactivation, the samples were self-ligated

with T4 DNA ligase. The ligated DNA was recovered using sodium acetate and ethanol and

quantified using the Qubit dsDNA HS assay kit (Thermo Fisher, Cat# Q32851). The 4C

templates were amplified using designed primers to generate sequencing libraries. The primer
system was modified to be compatible with the Illumina Nextera system, using two sequential
PCRs. Locus-specific inverse PCR primers are listed in **Supplemental Table 6**. For each locus,
the 4C templates were amplified with locus-specific primers using 200 ng of template per
reaction, and products from five parallel amplifications were pooled to generate the final 4C
library. A 50 µl PCR product aliquot was purified with homemade Sera-Mag beads. One-fifth of
the purified DNA was used for a second PCR with primers N7xx and N5xx, identical to Illumina
Nextera sample preparation primers. The final products were purified and sequenced. Reads
from the first cutting site were used for data analysis.

Plasmid cloning

The dCas9 expression vectors used in this study were created using the Cas9 expression
vector backbone from the pX330 plasmid (Addgene; plasmid 42230) through the In-Fusion
cloning technique. The dCas9 genes were amplified from the pHAGE EF1α dCas9-KRAB
plasmid (Addgene; plasmid 50919) via PCR and independently cloned into the AgeI and EcoRI
sites of the pX330 plasmid, substituting the Cas9 ORF. Detailed information regarding the
primers can be found in **Supplementary Table 5**. All sgRNAs in this study were designed using
the CCTop-CRISPR/Cas9 target online predictor.

The Cas9 and *Pcdhg* loop anchor target gRNA expression vector (pDonor_Rosa26_Cas9) was
constructed using the backbone of the plasmid pDonor MCS Rosa26 (Addgene, plasmid
#37200). An EF-1α-Cas9-P2A fragment, amplified from LentiV_Cas9_Puro (Addgene, plasmid
#108100), and a Puromycin-SV40 poly(A) signal fragment, derived from pR26 (Addgene,
plasmid #127372), were inserted into the backbone. A paired *Pcdhg* loop anchor target gRNA
expression cassette (sequences listed in **Supplementary Table 5**) was then introduced into the
construct.

CARGO assembly

The CARGO array assembly follows a published protocol[49]. The protocol consists of two main
steps: constant region preparation and array assembly. In the constant region preparation step,
the constant region with proper sticky ends is released by digesting the pGEMT-hU6-spSL
(provided by Dr. Joanna Wysocka) plasmid with BsmBI. The released constant region is then
separated from the pGEMT backbone on a 1% agarose gel and gel purified using a commercial
kit (QIAGEN, Cat#28704). Oligonucleotides (50-60 bp) with variable regions were ordered

through IDT (**Supplementary Table 5**). The forward and reverse oligos are annealed and
phosphorylated and then stored at -20°C.
In the array assembly step, the annealed chimeric oligo is mixed with the digested constant
region, and a minimum of 4-hour ligation with high-concentration T4 ligase (Thermofisher
Scientific, Cat#EL0013) is performed. The ligation step yields ligation products, referred to as
minicircles. After the stoichiometry check, individual minicircles are treated with plasmid-safe
exonuclease to clean up the un-ligated products. Then, the plasmid-safe exonuclease treated
individual minicircles are pooled together and purified. The pooled circle mixes are assembled
directly into the digested destination vector pX332-mCherry (provided by Dr. Joanna Wysocka)
through an optimized golden gate assembly reaction. The product of the golden-gate reaction is
then subjected to plasmid-safe exonuclease (Epicentre, Cat#E3101K) treatment. After that, the
exonuclease-treated reaction is then transformed into competent cells (Bioline, Cat#BIO-
85027). For each plate, individual colonies are picked for plasmid purification using a miniprep
kit (QIAGEN, Cat#27204) and test digestion with Acc65I, KpnI, and XbaI. Plasmids with the
correct digestion pattern are then verified by Sanger sequencing

ChIPmentation

We employed ChIPmentation[55] to map CTCF in various samples. Briefly, cells and tissues were
fixed in 1% formaldehyde at room temperature for 15 minutes, followed by glycine quenching.
To isolate nuclei, we lysed cell with lysis buffer 1 (50 mM HEPES; pH 7.5, 140 mM NaCl, 1 mM
EDTA, 10% glycerol, 0.5% Nonidet-40, 0.25% Triton X-100, and a protease inhibitor cocktail) for
10 minutes at 4°C. The nuclei were pelleted at 1,800 g for 10 minutes at 4°C and then
resuspended in nuclei lysis buffer (10 mM Tris-Cl pH 8.0, 100 mM NaCl, 1 mM EDTA, 0.5 mM
EGTA, 0.1% Na-Deoxycholate, 0.5% N-lauroylsarcosine, and a protease inhibitor cocktail) for
sonication. For CTCF pulldown, 100-150 µg of chromatin was used. Dynabeads M-280 (Life
Technologies, Sheep Anti-Rabbit IgG, #11204D) were washed three times with 0.5 mg/mL of
BSA/PBS on ice, then incubated with the CTCF antibody (Abcam, #ab70303) for at least 2
hours at 4°C. The beads/antibody complexes were then washed with BSA/PBS. Pulldown was
performed in binding buffer (1% Triton-X 100, 0.1% Sodium Deoxycholate, and a protease
inhibitor cocktail in 1X TE) by mixing the beads/antibody complexes with chromatin. After
overnight incubation, the beads/antibody/chromatin complexes were washed with RIPA buffer
(50 mM HEPES pH 8.0, 1% NP-40, 0.7% Sodium Deoxycholate, 0.5 M LiCl, 1 mM EDTA, and a
protease inhibitor cocktail). The beads complexes were then incubated with homemade Tn5
transposase in tagmentation reaction buffer (10 mM Tris-Cl pH 8.0 and 5 mM MgCl2) for 10

minutes at 37°C. To remove free DNA, the beads were washed twice with 1X TE on ice. The

pulldown DNA was recovered by reversing crosslinking overnight, followed by purification with

PCR Clean DX beads (Aline Biosciences, #C-1003-50). ChIP-seq libraries were generated by

amplifying the pulldown DNA with Illumina Nextera primers. Size selection was then performed

with PCR Clean DX beads to select fragments ranging from 100 bp to 1000 bp.

Pcdhg loop anchor deletion

Mouse embryonic stem cells (E14Tg2a) were transfected with the plasmid pX330-sgR26

(Addgene, plasmid #127376), which expresses mammalian codon-optimized Cas9 and the

Rosa26 sgRNA, together with the pDonor_Rosa26_Cas9 plasmid, using Lipofectamine™ 3000

reagent (Invitrogen) according to the manufacturer's instructions. Two days after transfection,

the cells were treated with puromycin (2 µg/mL) for three days. Drug-resistant clones were then

dissociated into single-cell suspensions, mixed thoroughly, and replated at low density to allow

single-colony formation, while maintaining puromycin selection (2 µg/mL). After four days,

individual colonies were picked for genotyping using primers specific for the *Pcdhg* loop anchor

deletion (**Supplementary Table 5**). Knockout cell clones were identified and selected for

subsequent experiments.

Data analysis

Hi-C, micro-C, and 4C-seq

Hi-C and micro-C sequencing reads were mapped to mm10 reference genome separately using

bowtie[56] (v.1.1.2) as previously described[28]. After removing non-unique reads and PCR

duplication reads, the bam files are used as input into *HiCorr* in '*Bam-process-HindIII*', '*Bam-*

process-DpnII', and '*Bam-process-microC*' modes for bias correction with conventional Hi-C, *in-*

situ Hi-C, and micro-C, respectively. After bias correction, we enhance the Hi-C/micro-C data

with *DeepLoop*. The *DeepLoop* package provides multiple models optimized for Hi-C data with

different read depths[28]. (Note that we use the number of *cis*-contact within 2Mbp as the

measurement of read depth.) To choose the appropriate model, we usually selected the model

with the closest read depth. The models chosen for each dataset are listed in **Supplementary**

Table 2. Because each dataset has a pair of Hi-C experiments (control and KO conditions), if

the read depths of the two Hi-C experiments are significantly different, we will down-sample the

high-depth dataset to lower depth to reduce potential biases (**Supplementary Table 2**).

4C-seq data were analyzed using pipe4C[54] (v.1.1.3) to generate bam and bigwig files for data
visualization on UCSC genome browser.

ChIP–seq and CUT&RUN

ChIP-seq and CUT&RUN data were mapped to mm10 (mouse) using bowtie. After mapping
step, we remove PCR duplication reads and use Macs2[57] to call peak and generated bigwig to
for data visualization on UCSC genome browser.

Motif analysis

Motif enrichment analysis was performed by comparing motif occurrences at ATAC peaks
located on gained loop anchors against all ATAC peaks. Background motif frequencies were
obtained by scanning all ATAC peak sequences using the mouse HOCOMOCO v11[58] motif
database with FIMO (MEME suite)[59]. Motif occurrences in ATAC peaks on gained loop anchors
were similarly identified by FIMO scanning. Odds ratios were calculated using Fisher's exact
test to quantify relative motif enrichment. To determine whether motifs were significantly
enriched or depleted, a binomial test was applied based on the observed verse expected motif
frequencies, and the resulting *p*-values were adjusted to q-values for motif testing correction.

RNA-seq and mNET-seq

RNA-seq data are aligned to mm10 using HISAT2[60] and read counts for each gene are called
with featureCounts[61]. DESeq2[62] is used to identify differential expressed genes (DEGs). The
DEGs are identified using Wald test with the p-value <0.01 and log2FC>1. For mNET-seq
analysis, we followed the customized pipeline from the Hsieh *et al.*[25] RNA-seq data are aligned
to mm10 using HISAT2[60] and read counts for each gene are called with featureCounts[61].
DESeq2[62] is used to identify differential expressed genes (DEGs). The DEGs are identified
using Wald test with the p-value <0.01 and log2FC>1. For NET-seq analysis, we followed the
customized pipeline from Hsieh *et al.* with the following steps: (i) Trim the adapter using
TrimGalore (v.0.6.7) (https://github.com/FelixKrueger/TrimGalore) to remove adapter sequences
'AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC' and
'GATCGTCGGACTGTAGAACTCTGAAC' at each side of the reads; (ii) Map trimmed read to
mm10 using STAR; (iii) Identify the last nucleotide incorporated by Pol II by using mNET_snr
(https://github.com/tomasgomes/mNET_snr) to locate the 3′-nucleotide of the second read and
the strand sign of the first read. (iv) Annotate and count the reads of each gene using

FeatureCounts[61]; (v) Use DEseq2[62] to quantify the changes of the mNET-seq signal at the gene

body between control and KO cells.

Pairwise loop strength comparison

For each pair of CTCF/cohesin-degron study, we took top 500k loop pixels from the *DeepLoop*

enhanced contact matrices of control and IAA/dTAG-treated samples separately, then union

them together. The loop strength of each pixel in the two experiments are normalized linearly:

$$Y_i \sim k * X_i + \varepsilon$$

$$i: A \ given \ loop \ pixel$$

$$X_i: i \ loop \ strength \ in \ WT \ condition$$

$$Y_i: i \ loop \ strength \ in \ KO \ condition$$

However, because CTCF- and *Rad21*-depletion leads to massive loss of loop signal, much

fewer loop pixels are expected in the IAA/dTAG-treated cells than in control cells. To adjust for

this, we have to manually pick a regression slope *k* based on the shape of the scatter plots.

The slopes we chose are listed in **Supplementary Table 2**. We then use the minimum strength

of the control cell loop pixels to adjust the minimum cutoff to call loop pixels in KO samples:

$Y_{min} = k * \min(X_i)$. In another word, a pixel is called as loop pixel in KO sample only if its

strength is bigger than Y_{min}. This step leads to much fewer loop pixels in CTCF- and *Rad21*-

depleted cells. We typically obtain ~600k chromatin loop for each study. We used a simple fold

change cutoff (FC > 4) to define gained or lost loop pixels. Finally, we used the following criteria

to call retained loops: (i) the strength of the pixel in both WT and KO conditions must be higher

than their minimum cutoffs ($X_i > X_{min}$ & $Y_i > Y_{min}$); (ii) the absolute value of residual must be

higher than the adjusted minimum of loop pixels in KO samples ($|\varepsilon| < Y_{min}$).

Classify chromatin loops based on epigenome data

For the purpose of being consistent, we intersected loop anchors with the same set of mESC

CTCF, H3K27ac, H3K4me3, H3K27me3 and H3K36me3 ChIP-seq peaks from ENCODE[63] to

classify the loop pixels. We classification loop pixels in the following order:

(1) EP-only: both anchors have either H3K4me3 or H3K27ac mark, but they do not have

CTCF peaks.

(2) CTCF-EP: after excluding EP-only, look for pixels which only one anchor has CTCF, and

the other anchor has either H3K4me3 or H3K27ac marker.

(3) CTCF-only: after excluding EP-only and CTCF-EP, look for pixels with at least one
anchor bound by CTCF.
(4) H3K27me3: after excluding (1) - (3) above, look for pixels with at least one anchor bound
by H3K27me3.
(5) H3K36me3: after excluding (1) - (4) above, look for pixels with at least one anchor bound
by H3K36me3.
(6) The rest pixels are classified as "other".

Define Loop-gaining loci, FINs, FIN loci, and FIN-blocked regular enhancers and super
enhancers (SE)

Loop-gaining loci are defined from "gained" loop pixels in CTCF-depleted cells. We identified
thousands of "gained" loop pixels and each pixel represents a pair of 5kb-bins. However, these
pixels often occupy overlapping regions. We therefore merge all overlapping pixels into one
region and define them as loop-gaining loci. This step allows us to count the number of regions
in the genome containing "gained" loops. In our study, we have ~300 loop-gaining loci after
merging ~5,000 recurrent "gained" loop pixels.

We define FINs as a subset of CTCF binding sites that satisfy the following criteria: (i) they
located at the anchors of CTCF loops that are lost upon CTCF depletion; (ii) the FIN should be
inside a newly gained loop in CTCF-depleted cells; (iii) the other anchor of the CTCF loop
should be outside of the newly gained loop; (iv) CTCF sites satisfying (i-iii) within 20kb of each
other will be merged into one FIN. Note that we do not have limits on the number of CTCF sites
to define FINs. Therefore, although a FIN can be just one single CTCF site, most FINs have
multiple CTCF sites.

As described above, all FINs must be inside loop-gaining loci. "FIN loci" are "loop-gaining loci"
that containing FINs. In fact, almost all the "loop-gaining loci" are also "FIN loci". We therefore
often use the two terms interchangeably.

We call enhancers as H3K27ac peaks and use a published list of super enhancers in mESCs[64]
(**Supplementary Tables 11 & 12**). This allows us to determine the types of enhancers blocked
by FINs.

Recurrent loop-gaining pixels and loop-gaining loci in CTCF depleted cells

A recurrent loop-gaining pixel is called if it satisfies FC > 4 in at least two studies. We define the
location of each loop pixel as the entire region between the left and right anchors (including the

anchors themselves). To define loop-gaining loci, we merge all gained loop pixels from the

CTCF-degron studies using the "merge" option in *bedtools*; each loop-gaining locus usually has

multiple loop pixels. Loop loci with gap size < 20kb will be merge into one locus. In this study,

we merged all the recurrent loop-gaining pixels to obtain 299 recurrent loop events. To test the

consistency of loop-gaining loci between independent CTCF-degron study, we call loop-gaining

loci for each individual CTCF-degron study and the share loop-gaining loci from two different

studies are called if they have at least one base overlap.

Test the recurrence of loop-gaining loci from CTCF-degron in Rad21- and Wapl- degron data

Nearly all loop-gaining loci observed in Hi-C degron studies have at least one E-P loop pixels.

We further classify the loop-gaining loci depending on the number of CTCF loops separating the

two anchors of the gained E-P loops. For all loci, we also manually examined all loci to confirm

the classification. Several loci cannot be clearly classified into the categories shown in **Fig. 3d**

because they have different marks or have multiple events. To test if a loop gaining locus is

recurrent in one of the CTCF/cohesin-degron study in **Fig. 3d**, we examine all loop-gaining

pixels in the locus and if one of pixels also gain signal it will be considered recurrent.

Compartment and TADs in mESCs

Hi-C compartment analysis is performed in WT mESCs following previous publication[65]. After QC

step, we constructed 500kb resolution contact matrices for each chromosome and then

normalized them using the genome distance. We next generated correlation matrices for each

chromosome and performed the principal component analysis on the correlation matrix and then

assigned the genome into two compartments by using PC1. We used the density of transcription

starting sites (TSS) of each 500-kb bin to determine compartment A and compartment B.

TAD boundary calling with Insulation score or Directionality index (DI) methods

Insulation scores were computed for contact matrices using 40kb bins and 2Mb sliding window

using the *cooltools*[66] package. Insulating loci were identified through a local minima detection

procedure based on peak prominence. Highly insulating regions corresponding to strong TAD

boundaries were defined using the Li thresholding method implemented in the *cooltools*

insulation score module. In this method, TAD boundaries are a list of 40kb bins.

We also called TAD domains with control mESC Hi-C data using the DI method described by

Dixon *et al.*[1], which quantifies the degree of upstream or downstream interaction bias for each

genome bin. Directionality indices were calculated also with 40kb bins using a 2Mb sliding

window. A hidden Markov model (HMM) was applied to infer hidden bias states across the

genome, classifying each bin as upstream-biased, downstream-biased, or neutral. Consecutive

bins sharing the same bias state were merged into domains. The start and end positions of

each domain are defined as TAD boundaries. In this way, the TAD boundaries will be also a list

of single bases. The TAD boundaries from the WT samples of the three datasets are listed in

Supplementary Table 9.

To compare the TAD boundaries from two methods, we consider two TAD boundaries

overlapping if their distance gap is smaller than 40kb. The same distance cutoff (40kb) is also

used to decide if a FIN overlaps TAD boundary.

Ideogram plot

We generate ideogram plots with Rideogram[67] to visualized the locations of recurrent gaining-

loop pixels, compartment, and TADs.

Density and enrichment of DEGs upon CTCF-depletion

For any region in the genome, the density of DEGs is defined as the average number of DEGs

per Mb-region. We computed DEG density for TAD boundary +/- 100kb, FIN +/- 100kb, gained

loop anchors, lost loop anchors, and retained loop anchors. We also compute the enrichment of

DEGs at regions of interest as their DEG density divided by the DEG density in the whole

genome.

Hi-C data visualization

Heatmap was used to visualize the raw, ratio, and *DeepLoop*-enhanced heatmaps. We use the

following strategies as described before[28] to determine the color scales these heatmaps. (i) Raw

or ICE-normalized heatmaps represent raw read counts; the brightest red color indicates the

98th percentile of the contact matrix. Color is proportionally scaled down to one read (white). (ii)

HiCorr heatmap represents the ratio value after Hi-C bias correction; the brightest red color

indicates at least twofold enrichment of a given pixel comparing to its background. Color is

proportionally scaled down to onefold (no enrichment). (iii) DeepLoop heatmaps outputs

'transformed fold change' that represents only relative levels of signal enrichment (for example,

a value of one may no longer represents no enrichment). We therefore set the brightest red

color as the lower limit of the top 300k pixels genome wide. Color is proportionally scaled down

to half of that lower limit or onefold, whichever is higher. All loop arches shown in figures were

generated with UCSC Genome Browser[68].

Data availability

Accession numbers for third-party data used in this study can be found in **Supplementary**

Table 1. The raw data of 4C–seq and ChIP-seq generated in this study, and reanalyzed

published data, can be found at accession number GSE243728. The 18 Hi-C and micro-C

datasets analyzed by *DeepLoop* can be found at https://hiview.case.edu/public/DeepLoop/.

To review GEO accession GSE243728:

Go to https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE243728

Enter token uvixkcaajpajpih into the box

Code availability

707   The code is available is available at github (https://github.com/JinLabBioinfo/FIN_project).

References

1. Dixon JR*, et al.* Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**, 376-380 (2012).

2. Nora EP*, et al.* Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* **485**, 381-385 (2012).

3. Bickmore WA, van Steensel B. Genome architecture: domain organization of interphase chromosomes. *Cell* **152**, 1270-1284 (2013).

4. Dekker J, Mirny L. The 3D Genome as Moderator of Chromosomal Communication. *Cell* **164**, 1110-1121 (2016).

5. McCord RP, Kaplan N, Giorgetti L. Chromosome Conformation Capture and Beyond: Toward an Integrative View of Chromosome Structure and Function. *Mol Cell* **77**, 688-708 (2020).

6. Merkenschlager M, Nora EP. CTCF and Cohesin in Genome Folding and Transcriptional Gene Regulation. *Annual review of genomics and human genetics* **17**, 17-43 (2016).

7. Nora EP*, et al.* Targeted Degradation of CTCF Decouples Local Insulation of Chromosome Domains from Genomic Compartmentalization. *Cell* **169**, 930-944 e922 (2017).

8. Rao SSP*, et al.* Cohesin Loss Eliminates All Loop Domains. *Cell* **171**, 305-320.e324 (2017).

9. Nasmyth K, Haering CH. Cohesin: its roles and mechanisms. *Annu Rev Genet* **43**, 525-558 (2009).

10. Chan KL, Roig MB, Hu B, Beckouet F, Metson J, Nasmyth K. Cohesin's DNA exit gate is distinct from its entrance gate and is regulated by acetylation. *Cell* **150**, 961-974 (2012).

11. Huis in 't Veld PJ*, et al.* Characterization of a DNA exit gate in the human cohesin ring. *Science* **346**, 968-972 (2014).

12. Kueng S*, et al.* Wapl controls the dynamic association of cohesin with chromatin. *Cell* **127**, 955-967 (2006).

13. Schwarzer W*, et al.* Two independent modes of chromatin organization revealed by cohesin removal. *Nature* **551**, 51-56 (2017).

14. Haarhuis JHI*, et al.* The Cohesin Release Factor WAPL Restricts Chromatin Loop Extension. *Cell* **169**, 693-707 e614 (2017).

15. Wutz G*, et al.* Topologically associating domains and chromatin loops depend on cohesin and are regulated by CTCF, WAPL, and PDS5 proteins. *The EMBO journal* **36**, 3573-3599 (2017).

16. Wallace JA, Felsenfeld G. We gather together: insulators and genome organization. *Curr Opin Genet Dev* **17**, 400-407 (2007).

17. Wendt KS*, et al.* Cohesin mediates transcriptional insulation by CCCTC-binding factor. *Nature* **451**, 796-801 (2008).

18. Parelho V*, et al.* Cohesins functionally associate with CTCF on mammalian chromosome arms. *Cell* **132**, 422-433 (2008).

19. Franke M*, et al.* Formation of new chromatin domains determines pathogenicity of genomic duplications. *Nature* **538**, 265-269 (2016).

20. Weischenfeldt J*, et al.* Pan-cancer analysis of somatic copy-number alterations implicates IRS4 and IGF2 in enhancer hijacking. *Nat Genet* **49**, 65-74 (2017).

21. Lupianez DG*, et al.* Disruptions of topological chromatin domains cause pathogenic rewiring of gene-enhancer interactions. *Cell* **161**, 1012-1025 (2015).

22. Anania C*, et al.* In vivo dissection of a clustered-CTCF domain boundary reveals developmental principles of regulatory insulation. *Nat Genet* **54**, 1026-1036 (2022).

23. Zuin J*, et al.* Nonlinear control of transcription through enhancer-promoter interactions. *Nature* **604**, 571-577 (2022).

24. Kubo N*, et al.* Promoter-proximal CTCF binding promotes distal enhancer-dependent gene activation. *Nat Struct Mol Biol* **28**, 152-161 (2021).

25. Hsieh TS, Cattoglio C, Slobodyanyuk E, Hansen AS, Darzacq X, Tjian R. Enhancer-promoter interactions and transcription are largely maintained upon acute loss of CTCF, cohesin, WAPL or YY1. *Nat Genet* **54**, 1919-1932 (2022).

26. Akgol Oksuz B*, et al.* Systematic evaluation of chromosome conformation capture assays. *Nature methods*, (2021).

27. Lu L*, et al.* Robust Hi-C Maps of Enhancer-Promoter Interactions Reveal the Function of Non-coding Genome in Neural Development and Diseases. *Mol Cell* **79**, 521-534.e515 (2020).

28. Zhang S*, et al.* DeepLoop robustly maps chromatin interactions from sparse allele-resolved or single-cell Hi-C data at kilobase resolution. *Nat Genet* **54**, 1013-1025 (2022).

29. Kriz AJ, Colognori D, Sunwoo H, Nabet B, Lee JT. Balancing cohesin eviction and retention prevents aberrant chromosomal interactions, Polycomb-mediated repression, and X-inactivation. *Mol Cell* **81**, 1970-1987 e1979 (2021).

30. Rhodes JDP*, et al.* Cohesin Disrupts Polycomb-Dependent Chromosome Interactions in Embryonic Stem Cells. *Cell reports* **30**, 820-835 e810 (2020).

31. Liu NQ*, et al.* WAPL maintains a cohesin loading cycle to preserve cell-type-specific distal gene regulation. *Nat Genet* **53**, 100-109 (2021).

32.   Imakaev M, *et al.* Iterative correction of Hi-C data reveals hallmarks of chromosome
810   organization. *Nature methods* **9**, 999-1003 (2012).
811

812   33.   Yaffe E, Tanay A. Probabilistic modeling of Hi-C contact maps eliminates systematic
813   biases to characterize global chromosomal architecture. *Nat Genet* **43**, 1059-1065
814   (2011).
815

816   34.   Lu L, *et al.* Robust Hi-C Maps of Enhancer-Promoter Interactions Reveal the Function of
817   Non-coding Genome in Neural Development and Diseases. *Mol Cell* **79**, 521-534 e515
818   (2020).
819

820   35.   Crane E, *et al.* Condensin-driven remodelling of X chromosome topology during dosage
821   compensation. *Nature* **523**, 240-244 (2015).
822

823   36.   Li L, *et al.* YY1 interacts with guanine quadruplexes to regulate DNA looping and gene
824   expression. *Nat Chem Biol* **17**, 161-168 (2021).
825

826   37.   Wulfridge P, *et al.* G-quadruplexes associated with R-loops promote CTCF binding. *Mol
827   Cell* **83**, 3064-3079 e3065 (2023).
828

829   38.   Schoenfelder S, *et al.* Preferential associations between co-regulated genes reveal a
830   transcriptional interactome in erythroid cells. *Nat Genet* **42**, 53-61 (2010).
831

832   39.   Di Giammartino DC, *et al.* KLF4 is involved in the organization and regulation of
833   pluripotency-associated three-dimensional enhancer networks. *Nat Cell Biol* **21**, 1179-
834   1190 (2019).
835

836   40.   Wei Z, *et al.* Klf4 organizes long-range chromosomal interactions with the oct4 locus in
837   reprogramming and pluripotency. *Cell stem cell* **13**, 36-47 (2013).
838

839   41.   Bryan AF, Justice M, Stutzman AV, McKay DJ, Dowen JM. Cohesin stabilization at
840   promoters and enhancers by common transcription factors and chromatin regulators.
841   *Epigenetics Chromatin* **18**, 33 (2025).
842

843   42.   Ortabozkoyun H, *et al.* CRISPR and biochemical screens identify MAZ as a cofactor in
844   CTCF-mediated insulation at Hox clusters. *Nat Genet* **54**, 202-212 (2022).
845

846   43.   Xiao T, Li X, Felsenfeld G. The Myc-associated zinc finger protein (MAZ) works together
847   with CTCF to control cohesin positioning and genome organization. *Proceedings of the
848   National Academy of Sciences of the United States of America* **118**, (2021).
849

850   44.   Ortabozkoyun H, *et al.* Members of an array of zinc-finger proteins specify distinct Hox
851   chromatin boundaries. *Mol Cell* **84**, 3406-3422 e3406 (2024).
852

853   45.   Hirayama T, Yagi T. Regulation of clustered protocadherin genes in individual neurons.
854   *Seminars in cell & developmental biology* **69**, 122-130 (2017).
855

856   46.   Flaherty E, Maniatis T. The role of clustered protocadherins in neurodevelopment and
857   neuropsychiatric diseases. *Curr Opin Genet Dev* **65**, 144-150 (2020).
858

47. Jia Z, Li J, Ge X, Wu Y, Guo Y, Wu Q. Tandem CTCF sites function as insulators to balance spatial chromatin contacts and topological enhancer-promoter selection. *Genome Biol* **21**, 75 (2020).

48. Hirayama T, Tarusawa E, Yoshimura Y, Galjart N, Yagi T. CTCF is required for neural development and stochastic expression of clustered Pcdh genes in neurons. *Cell reports* **2**, 345-357 (2012).

49. Gu B*, et al.* Transcription-coupled changes in nuclear mobility of mammalian cis-regulatory elements. *Science* **359**, 1050-1055 (2018).

50. Tarjan DR, Flavahan WA, Bernstein BE. Epigenome editing strategies for the functional annotation of CTCF insulators. *Nature communications* **10**, 4258 (2019).

51. Pollex T*, et al.* Enhancer-promoter interactions become more instructive in the transition from cell-fate specification to tissue differentiation. *Nat Genet* **56**, 686-696 (2024).

52. Fulco CP*, et al.* Activity-by-contact model of enhancer-promoter regulation from thousands of CRISPR perturbations. *Nat Genet* **51**, 1664-1669 (2019).

53. Kagey MH*, et al.* Mediator and cohesin connect gene expression and chromatin architecture. *Nature* **467**, 430-435 (2010).

54. Krijger PHL, Geeven G, Bianchi V, Hilvering CRE, de Laat W. 4C-seq from beginning to end: A detailed protocol for sample preparation and data analysis. *Methods* **170**, 17-32 (2020).

55. Schmidl C, Rendeiro AF, Sheffield NC, Bock C. ChIPmentation: fast, robust, low-input ChIP-seq for histones and transcription factors. *Nature methods* **12**, 963-965 (2015).

56. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).

57. Zhang Y*, et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

58. Vorontsov IE*, et al.* HOCOMOCO in 2024: a rebuild of the curated collection of binding models for human and mouse transcription factors. *Nucleic acids research* **52**, D154-D163 (2024).

59. Grant CE, Bailey TL, Noble WS. FIMO: scanning for occurrences of a given motif. *Bioinformatics* **27**, 1017-1018 (2011).

60. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. *Nature methods* **12**, 357-360 (2015).

61. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

62. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

910
911 63.   Yue F, *et al.* A comparative encyclopedia of DNA elements in the mouse genome.
912   *Nature* **515**, 355-364 (2014).
913
914 64.   Hnisz D, *et al.* Super-enhancers in the control of cell identity and disease. *Cell* **155**, 934-
915   947 (2013).
916
917 65.   Lieberman-Aiden E, *et al.* Comprehensive mapping of long-range interactions reveals
918   folding principles of the human genome. *Science* **326**, 289-293 (2009).
919
920 66.   Abdennur N, Mirny LA. Cooler: scalable storage for Hi-C data and other genomically
921   labeled arrays. *Bioinformatics* **36**, 311-316 (2020).
922
923 67.   Hao Z, *et al.* RIdeogram: drawing SVG graphics to visualize and map genome-wide data
924   on the idiograms. *PeerJ Comput Sci* **6**, e251 (2020).
925
926 68.   Navarro Gonzalez J, *et al.* The UCSC Genome Browser database: 2021 update. *Nucleic*
927   *acids research* **49**, D1046-D1057 (2021).
928
929

Acknowledgements

This work was supported by grants from NIH R01HG009658 and R01CA267872 (to F.J.),
R01DK113185 (to Y.L.), R01DK131437, R01HG012384, and UG3NS132061 (to Y.L. and F.J.).
F.J. is also supported by a subaward from NIH project (U01AG072579) through University of
Miami and partially by several other NIH grants R01CA252224, R01GM148662, and
P01CA272161. Y.L. is also partially supported by NIH grants R01CA264320. This work made use
of the High-Performance Computing Resource in the Core Facility for Advanced Research
Computing at Case Western Reserve University.
938

Author Contributions Statement

940 F.J. and Y.L conceived the project and secured funding for this project. J.C. designed and
941 performed the FIN validation experiments, W.X. and X.L. performed the data analysis. L.L. help
942 with the experiments and S.Z. and X.L. helped with the data analysis. F.J., W.X., J.C.,X.L., and
943 Y.L. wrote the manuscript. F.J. and Y.L. also provided critical feedback and interpretation of
944 results as well as helping to revise the manuscript.
945

946 **Competing Interests Statement**

947 The authors declare no competing interests
948

 Figure Legend

Figure 1. *DeepLoop* allows cross-platform comparison between independent CTCF- and
cohesin-degron Hi-C datasets at loop-resolution.

a, A list of published Hi-C and micro-C studies with acute depletion of *Ctcf*, *Rad21*, or *Wapl* in
mESCs. **b,** Matrices summarizing R-square values when comparing contact heatmaps from
different pipelines and orthogonal Hi-C/micro-C datasets in control mESCs. **c,** Correlation
matrices comparing the differential matrices of all nine degron experiments. For each degron
experiment, the differential matrix is the difference between two Hi-C contact matrices (depletion
– control) at 5kb resolution. Left, Hi-C contact matrices from *ICE* are used to compute
differential matrices; right, Hi-C contact matrices from *DeepLoop* are used. **d,** Contact heatmaps
of the CTCF-, Rad21-, and Wapl-degron Hi-C data at *Notch1* locus with *ICE*, *HiCorr* and
DeepLoop. Red dashed circle: gained loops. Numbers in brackets indicate the color scale of
heatmaps: for *ICE* normalization and *HiCorr* heatmaps, the maximum intensity corresponds to
the 98th percentile of the corresponding contact matrix; for *DeepLoop* heatmaps, the maximum
intensity corresponds to the lower cutoff of top 500,000 pixels genome-wide. **e**, Genome
browser tracks showing ChIP-seq data and Hi-C loops from Nora *et al.* at *Notch1* locus. Blue
highlighted region showing Notch1 gene which is at the anchor of a gained loop. Pink region
highlights the potential insulator region that blocks the formation of *Notch1* E-P loop. RNA-seq
results are also shown for *Notch1* gene in WT (black) and CTCF-depleted cells (red). Bar plot:
quantitation of *Notch1* expression from RNA-seq. P-value: one-side Wald test provided by DE-
seq2.

Figure 2. Identify and characterize recurrent gain of E-P loop events in CTCF-depleted
cells through cross-platform Hi-C meta-analyses.

a-c, Meta-analysis of independent CTCF-degron Hi-C datasets. **a**, Left: scatterplot comparing
the strength of loop pixels between WT and CTCF-depleted mESCs (every dots represent one
pixel in the contact heatmap); middle: the same scatterplot as left but highlight the lost (blue),
retained (black), and gained (red) loop pixels; right: violin plot comparing the size of loop pixels
and the value below each plot indicates the median loop size for the corresponding category. **b,**
Pie charts showing the components of lost, retained and gained loop pixels based on epigenetic
marks. **c**, Scatter plots showing the consistency of lost, retained and gained loop pixels between
three CTCF degron studies. Plots in the left column highlight the lost, retained, and gained loop

pixels from Nora *et al.*, the right two columns show the quantitative changes of the

corresponding pixels in *Kubo et al.* or *Hsieh et al.* **d,** The lost, retained, and gained loop pixels

from one CTCF-degron Hi-C data (Nora *et al.*) are plotted onto Hi-C scatter plots from *Rad21-*

or *Wapl*-degron studies to examine their quantitative changes.

Figure 3. Characterization of functional insulators (FINs) in mESC genome.

a, Heatmap showing the union of all gained loop pixels from three CTCF-degron Hi-C datasets.

Every row represents one CTCF-degron study, and every column represents one loop pixel.

Loop pixels are organized into clusters based on their reproducibility. The number of loop pixels

in each cluster and their median size are also shown. Color scale indicates the increase of loop

signal in log scale. **b**, Venn diagram showing the consistence of loop-gaining loci identified from

three datasets. **c,** Pie chart showing that most of the gained loops in CTCF-depleted cells have

at least one CTCF loop anchor in between in wildtype cells. **d,** A table summarizing the

classification and recurrence of loop gaining loci between independent CTCF- or cohesin-

degron studies. **e,** Cartoons summarizing three possible ways to define insulators. **f,** An

example of non-TAD boundary FIN. The FIN is highlighted in pink. The tracks of InsScore and

DI are also included. TAD boundaries determined from InsScore and DI are highlighted in

purple and indicated with black arrows, respectively. **g,** Ideogram showing the locations of loop-

gaining loci from three studies, TAD boundaries, and compartments on chromosome 1. FINs

from three different CTCF-degron studies are shown in different colors, and the color scale

shows the fold increase of gained loop in log scales. **h,** Pie chart showing the percentage of

FINs that overlap TAD boundary. **i,** Boxplot comparing the insulation scores at FINs and TAD

boundaries before and after CTCF depletion computed with HI-C data from Nora *et al*. **h,** Stack

bar plot comparing the proportion of FINs and TAD boundaries in compartment A or B. **i,** A

model to explain why some of the newly gained E-P loops in CTCF-depleted cells are also

observed in Wapl-depleted cells. **j,** A list of TF motifs enriched or depleted in the mESC ATAC-

seq peaks that will gain chromatin loops upon CTCF depletion. Enrichment or depletion are

relative to all mESC ATAC-seq peaks.

Figure 4. FINs are associated with consistent early transcriptional consequences in

independent CTCF-degron studies.

a, Bar plots comparing the density of DEGs upon CTCF-depletion across different genomic

regions. The y-axis indicates the number of DEGs per Mb. The number of DEGs were

calculated from the following datasets: 3-hour NET-seq for Hsieh *et al.*, 24-hour RNA-seq for

Nora *et al*., and 24-hour RNA-seq for Kubo *et al.* **b,** Enrichment of DEGs at different types of

genomic regions compared to whole genome as background. X-axis: time of IAA treatment (hrs)

in each study; y-axis: fold enrichment. Each panel represents one CTCF-degron study. **c,** Pie

chart showing the percentage of FINs that are blocking super enhancers; **d,** Boxplot showing

the expression changes of genes gaining enhancer loop. Red: genes gaining super-enhancer

loop; Blue: genes gaining regular enhancer loop. *, p < 0.05, Wilcoxon signed-rank test. **e,** Venn

diagrams showing the overlap of all up- (left panel) and down-regulated (right panel) DEGs

between three CTCF-degron studies. **f,** Bar plot comparing the reproducibility scores (measured

by Jaccard Index) of up- or down-regulated DEGs between two CTCF-degron studies (Hsieh *et*

al. and Nora *et al.*). Gray bars: all DEGs; blue bars: DEGs at the anchors of lost loops; red bars:

DEGs at the anchors of gained loops. P-values are from binomial test. **g,** Heatmap showing the

union of upregulated DEGs at the anchors of gained loops from *Hsieh et al* and *Nora et al.*;

DEGs are clustered based on consistency. **h,** Left panel shows the browser tracks near *Cdiptos*

locus and right panel shows the Hi-C heatmaps in the same region from *Hsieh et al* and *Nora et*

al. The direction of CTCF motifs are indicated by triangles. Blue arches (left) and blue dashed

circles (right): lost loop pixels; red arches and red dashed circles: gained loop. Regions

highlighted in green: enhancers that gained loops; the region highlighted in pink: a CTCF loop

anchor is defined as *FIN*. **i,** Similar as **g**, a heatmap showing the union of down-regulated DEGs

at the anchors of gained loops from *Hsieh et al.* and *Nora et al*. Among these genes 18 genes

are from the Pcdh-γ locus (shown as a separate heatmap). **j,** Left panel shows the browser

tracks near Pcdh-γ locus and right panel shows the Hi-C heatmaps from two CTCF-degron

studies. **k,** Design of sgRNAs for insulator deletion. **l,** RT-qPCR shows the down-regulation of

Pcdh-γ genes in mESCs after insulator deletion.

Figure 5. Validate FINs with multiplexed CTCF displacement assay.

a, Scheme of the CARGO vector that can express multiple sgRNAs. In this system, the

sequences of *n* sgRNAs are broken into left and right halves and inserted into (*n*+1) minicircles.

The *i*-th minicircle contains the right half of the (*i-1*)-th sgRNA and the left half of the *i*-th sgRNA

sequences divided by a *BpiI* restriction site. Final CARGO vector can be assembled by ligating

the re-linearized minicircles and vector backbone; each sgRNA is driven by a U6 promoter. Co-

transfect CARGO vector and a dCas9-expressing plasmid can displace multiple CTCF binding

sites in the same cell. **b,** Validation of a FIN at the *Cdiptos* locus. From the top, 1st and 2nd

tracks are H3K4me3 and H3K27ac ChIP-seq data showing the promoter and enhancers in

mESC. 3rd and 4th tracks show CTCF ChIP-seq data before and after targeting 4 CTCF motifs in

FIN (highlighted in pink) with CARGO. 5th and 6th are Hi-C arches (*DeepLoop* pixels with data

from Nora *et al.*) before and after CTCF-depletion (blue: lost loops; red: gained loops). The

bottom 3 tracks show 4C-seq data (bait highlighted in yellow) before and after targeting CTCF

sites; regions highlighted in green gain 4C signal. The raw and ICE, HiCorr, or DeepLoop

processed contact heatmaps are also included. **c,** A model cartoon showing that the gained

loop at this FIN locus only involves one CTCF loop. **d,** RT-qPCR results showing that *Cdiptos* is

upregulated after CTCF displacement. ** p<0.01; one-side t-test. **e-g,** Validation of a FIN at

Mesp1 locus similar as shown in **b-d**. Note the gained loop at this FIN locus involves at least

two CTCF loops (**f**).

1054

Supplementary figure legend

Supplementary Figure 1. Compare the compartments and TADs between orthogonal

CTCF-degron datasets from Hi-C, in situ Hi-C, and micro-C.

a-c, Compartment analysis of CTCF-degron Hi-C datasets at 500kb resolution. **a,** Correlation

heatmaps of the three CTCF-degron Hi-C datasets at 500kb resolution showing the

compartments in control and CTCF-depleted cells. PC1 values are shown on top of each

heatmap indicating compartment A (positive PC1) and B (negative PC1). **b,** Scatter plots

comparing the PC1 values for each 500kb bin before and after CTCF depletion. **c,** Top row:

scatter plots comparing the PC1 values of Hi-C data in control cells from three independent

studies; bottom row: scatter plots comparing the PC1 values of Hi-C data in CTCF-depleted

cells from three studies. **d-f,** TAD analysis of CTCF-degron Hi-C datasets at 40kb resolution. **d,**

ICE normalized contact heatmaps of the three CTCF-degron Hi-C datasets at 40kb resolution.

TADs are recognizable as squares along the diagonals. Note the loss of TADs in CTCF-

depleted cells. **e,** Scatter plots compare the directionality index (DI) values of all 40kb bins

before and after CTCF-depletion in all three studies. Note that the DI values of many bins

become close to zero in CTCF-depleted cells, indicating that these bins lost directional biases,

consistent with massive loss-of-TAD upon CTCF-depletion. **f,** Top row: scatter plots comparing

the DI values of all 40kb bins in control cells from three independent studies; bottom row: scatter

plots comparing the DI values of all 40kb bins in CTCF-depleted cells from three studies. Note

that although the R-squares between Nora et al. and Kubo et al. are rather high (R-square

0.718 and 0.612 for control and △CTCF cells), the DI values from micro-C data (Hsieh et al.)

deviate significantly from the other two studies (R-square 0.295~0.458). In micro-C data, more

bins show polarized DI values (*i.e.*, stronger directionality biases), consistent with the general

belief that with ICE-based pipeline, micro-C identifies more loops than Hi-C or in situ Hi-C.

Supplementary Figure 2. Compare the loop-level analyses of orthogonal Hi-C datasets

with ICE, HiCorr, and DeepLoop, related to Fig. 1b.

Part I. Cross-platform comparison of CTCF-degron datasets. **a-c,** Loop analysis of three CTCF-

degron Hi-C datasets with ICE at 5kb resolution. **a,** ICE heatmaps of the three CTCF-degron Hi-

C datasets at 5kb resolution. Note that there are more noises in Hi-C (Nora) and *in situ* Hi-C

(Kubo) heatmaps than in micro-C (Hsieh) heatmaps. **b,** Scatter plots of ICE heatmap pixels

comparing the control cells contact heatmaps from three independent studies. Note that the

consistency between orthogonal Hi-C protocols is low (R-square values close to zero). **c,**

Scatter plots comparing the ICE heatmaps of control and ΔCTCF cells in three different studies

(5kb resolution). Every dot in the scatter plot represents one pixel in the contact matrix and the

x-axis and y-axis indicate the intensity of the pixel in two experiments. Note that △CTCF cells

have many pixels stronger than WT cells because the read-depth of micro-C data in △CTCF

cells is higher than in control cells. **d-f,** Similar as **a-c** but the comparisons are between ICE

heatmaps after correcting distance biases, which are observed/expected ratio heatmaps. Note

that in **f**, the control vs. △CTCF scatter plot for Hsieh *et al.* correctly show stronger loop pixels

in control cells than △CTCF cells. However, the scatter plots in **e** show that the consistency

between independent studies remains poor with R-square values still close to zero, indicating

that the pixel-level correlations are dominated by random noise pixels. **g-i,** the same

comparisons between HiCorr-corrected heatmaps, which are also distance-corrected ratio

heatmaps. Note that the scatter plots in **i** show that in each of the control vs. △CTCF

comparisons, we start to observe retained pixels with similar strength before and after CTCF

depletion. Furthermore, the scatter plots in **h** also show that HiCorr achieves higher R-square

values across different Hi-C platforms. **j-l,** comparisons between *DeepLoop* heatmaps. Note

that DeepLoop highlights loop pixels and suppresses noise pixels (**j**). The retained pixels and

weakened pixels are more clearly recognizable in each of the control vs. △CTCF scatter plots

(**l**). DeepLoop also shows the best cross-platform consistency at pixel level (**k**).

Part II. Cross-platform comparison of Rad21-degron datasets. Same as Part I and all

conclusions are the same.

Part III. Cross-platform comparison of Wapl-degron datasets. Same as Part I and all
conclusions are the same.

Supplementary Figure 3. Correlation matrices comparing the profiles of dynamic loops
from all nine pairs of degron experiments, related to Fig. 1c.

For each degron experiment, we first obtain a pair of 5kb resolution Hi-C contact matrices in
control and depletion cells. Next, we computed the delta matrices (depletion – control) for all
nine experiments. A correlation matrix is then generated between the nine delta matrices. **a-d**
are correlation matrices when contact matrices are generated with different pipelines.

Supplementary Figure 4. Recurrent gain of E-P loop at *Notch1* locus between
independent CTCF-degron Hi-C studies, related to Fig. 1d,e.

a, Contact heatmaps of the CTCF-degron Hi-C data by Kubo *et al.* at *Notch1* locus with *ICE*,
HiCorr and *DeepLoop*. Red dashed circle highlights the same locations where we observed a
gained E-P loop in the contact heatmap from Nora *et al.* (shown in **Fig. 1d**). **b**, Genome browser
tracks showing ChIP-seq data and Hi-C loops from Kubo *et al.* Blue highlighted region indicates
Notch1 gene. Pink region highlights the potential insulator region that blocks the formation of
Notch1 E-P loop. **c,** RNA-seq results from Kubo *et al.* are also shown as bar plot. p-value: one-
side Wald test provided by DE-seq2. **d-f,** the same analysis as **a-c** at *Notch1* locus using data
from Hsieh *et al.* Note that Hsieh *et al.* used mNET-seq for transcriptome profiling. **g,** Contact
heatmaps of *Notch1* locus from two *Rad21*-degron datasets (Kriz *et al.* and Hsieh *et al.*) in
addition to the one included in **Fig. 1d** (Rhodes *et al.*). **h,** Contact heatmaps of *Notch1* locus
from two *Wapl*-degron datasets (Kriz *et al.* and Hsieh *et al.*) in addition to the one included in
Fig. 1d (Liu *et al.*).

Supplementary Figure 5. Meta-analysis of independent CTCF-degron Hi-C datasets to
examine the consistency and epigenetic features of dynamic loops.

a, Using the *in situ* Hi-C data from the CTCF-degron study by Kubo *et al.*, left: scatterplot
comparing the strength of loop pixels between WT and CTCF-depleted mESCs; middle: the
same scatterplot as left but highlight the lost (blue), retained (black), and gained (red) loop
pixels; right: violin plot comparing the size of loop pixels. **b,** Pie charts showing the components
of lost, retained and gained loop pixels based on epigenetic marks. **c**, Scatter plots in the left

column highlight the lost, retained, and gained loop pixels from Kubo *et al.*, the right two

columns show the quantitative changes of the corresponding pixels in Nora *et al.* or Hsieh *et al.*

d-f, The same analyses as **a-c** are performed with CTCF-degron micro-C data by Hsieh *et al.*

Supplementary Figure 6. Examples of chromatin loop classification.

1140 Each panel shows the genome browser tracks with loop arches. These examples include **a**, EP

1141 loop; **b**, CTCF-EP loop; **c**, CTCF loop; **d**, H3K27me3 loop; **e**, H3K36me3 loop.

Supplementary Figure 7. Meta-analysis of independent *Rad21*- and *Wapl*- degron Hi-C

datasets to examine the consistency and epigenetic features of dynamic loops.

Part I: Meta-analysis of *Rad21*-degron datasets. **a,** Left and middle panels define the lost,

retained, and gained loop pixels upon *Rad21*-depletion from the study by Rhodes *et al.*, and the

right panel compares their loop size distribution. **b,** Pie charts showing the components of lost,

retained and gained loops based on epigenetic marks. **c,** Scatter plots testing if the lost,

retained and gained loop pixels from Rhodes *et al.* are quantitatively consistent in the other two

Rad21-degron studies. **d-f,** the same analyses as **a-c** are performed with *Rad21*-degron Hi-C

data by Kriz *et al.* **g-i,** the same analyses as **a-c** are performed with *Rad21*-degron micro-C data

by Hsieh *et al.*

Part II: Meta-analysis of *Wapl*-degron datasets. **j-r,** Same as Part I except that the analyses

were done with Wapl-degron datasets. Note there is an overall trend that *Wapl*-depletion causes

more "gained" loops.

Supplementary Figure 8. "Gained" long-range H3K27me3 loops in Rad21-depleted cells

from Rhodes *et al.* exist before Rad21-depletion in two other studies.

1157 Two examples are shown in **a** and **b**. In each example, Hi-C data are shown as both contact

1158 heatmaps (DeepLoop enhanced) and loop arches. Browser tracks of CTCF, H3K27ac,

1159 H3K4me3, and H3K27me3 are also included. Note that the gained H3K27me3 loop in Rhodes

1160 et al. already exist before Rad21-depletion in two other studies.

Supplementary Figure 9. Meta-analysis reveals the fate of dynamic loops from CTCF-

degron studies upon depletion of cohesin factors.

a, The lost, retained, and gained loop pixels from one CTCF-degron Hi-C data (Kubo *et al.*) are

plotted onto Hi-C scatter plots from *Rad21*- or *Wapl*-degron studies to examine their quantitative

changes. **b,** Same analyses are done with the lost, retained, and gained loop pixels from the

CTCF-degron Hi-C data by Hsieh *et al.*

Supplementary Figure 10. Examples of loop gaining loci, related to Fig. 3d.

a, An example of Type I EP loop gaining loci, in which one of the two E/P sites is insulated in a

CTCF loop while the other site is outside the CTCF loop; **b,** An example of Type II EP-loop

gaining loci, in which two E/P sites are insulated in two CTCF loops; **c,** An example of Type III

EP loop gaining loci, in which neither of the two E/P sites are insulated by CTCF loops; **d,** An

example of CTCF loop gaining loci, in which the newly gained loop are between incompletely

degraded CTCF sites.

Supplementary Figure 11. Limited overlap between FINs and TAD boundaries, related to

Fig. 3f,h,i.

a, Venn diagrams comparing the TAD boundaries called by two methods from three studies. **b,**

Pie chart showing the percentage of FINs that overlap TAD boundaries called from different

datasets and algorithms. **c,** Compare insulation scores at FINs and TAD boundaries in different

studies.

Supplementary Figure 12. Example contact heatmaps at putative insulator target genes,

related to Fig. 4h,j.

a, Raw, *ICE*, *HiCorr*, and *DeepLoop* contact heatmaps at *Cdiptos* locus in control and CTCF-

depleted cells from Hsieh *et al.* and Nora *et al.* **b,** Similar as **a** but the heatmaps are at the

Pcdh-γ locus.

Figure 1

a Published Hi-C data in mESCs upon acute depletion of architectural proteins

Publication	mESC line	KO protocol	C-technique	# of contacts	
				Ctrl	IAA/dTAG
Nora et al.	E14TG2a	C-Ctcf AID (48hrs)	HindIII Hi-C	244M	242M
Kubo et al.	F123	C-Ctcf AID (48hrs)	in-situ Hi-C	185M	1 16M
Hsieh et al.	JM8.N4	N-Ctcf AID (3hrs)	micro-C	291M	830M
Rhodes et al.	E14TG2a	C-Rad21 AID (6hrs)	in-situ Hi-C	63M	43M
Kriz et al.	F123	C-Rad21 dTAG (8hrs)	In-situ Hi-C	136M	100M
Hsieh et al.	JM8.N4	C-Rad21 AID (3hrs)	micro-C	283M	1.00B
Liu et al.	E14TG2a	C-Wapl AID (24hrs)	in-situ Hi-C	77M	88M
Kriz et al.	F123	C-Wapl dTAG (72hrs)	in-situ Hi-C	193M	147M
Hsieh et al.	JM8.N4	N-Wapl AID (3hrs)	micro-C	298M	670M

b Pixel-by-pixel correlation between contact heatmaps from orthogonal studies

c Correlation between differential matrices (depletion - control)

d ICE normalization | HiCorr | DeepLoop

e chr2:26,176,653-27,145,048

Figure 2

Figure 3

a Non-recurrent N=3,197 | Recurrent N=4,944

Nora, Kubo, Hsieh

log2(ΔCTCF/Ctrl): <1, [1,2], [2,3], >=3

N=1,969; 170kb / N=493; 455kb / N=735; 305kb / N=354; 235kb / N=2,621; 200kb / N=803; 435kb / N=1,166; 340kb

b Nora N=332, Kubo N=271: 42, 48, 68, 174, 35, 37, 14; Hsieh N=293

c Gained loops: 279; 4,665. Insulated by CTCF loops? Yes (grey) / No (orange)

d Classification and recurrence of loop gaining loci between independent CTCF/cohesin degron studies

Type of loci	Classification of loci	#of loci	Ctcf degron			Rad21 degron			Wapl degron		
			Nora	Kubo	Hsieh	Rhodes	Kriz	Hsieh	Liu	Kriz	Hsieh
EP N=283	I.	107	105	83	87	1	0	1	21	26	26
	II.	156	150	122	135	0	0	0	47	51	48
	III.	20	20	14	17	0	0	0	0	0	1
CTCF N=2		2	2	2	2	0	0	0	2	2	1
Undef N=14	Undetermined	14	12	10	13	1	1	2	11	8	10

⬡ CTCF ▶◀ CTCF motif ⬭ cohesin ▬ Enhancer or Promoter

e i. Define enhancer-blocking insulator by its transcription consequences

Enhancer — Insulator — Promoter → OFF
Enhancer — Insulator (X) — Promoter → ON

ii. Define insulator as TAD boundary

CTCF Insulator — Enhancer — CTCF Insulator — Promoter — CTCF Insulator

iii. Define insulator by loop dynamics

Anchor — Enhancer — CTCF Insulator — Promoter → OFF
Anchor — Enhancer — Insulator — Promoter → ON

Functional insulator (FIN)

f chr7:126,656,644-127,322,620 FIN

CTCF motif; CTCF; K4me3; K27ac

TAD boundary ... TAD boundary

Nora Control: InsScore; Direction index; TAD boundary

Cdiptos

Nora Control / ΔCTCF

g Chr1

FIN loci-Nora / FIN loci-Kubo / FIN loci-Hsieh / TAD boundary / Compartment

Log2(FC) Ctcf-AID/Ctrl: 0–2
Compartment A / B

h FIN & TAD boundary Nora et al.: 20.4% on-TAD boundary; 79.6% non-TAD boundary

i Nora et al.

log2(Insulation Score)

**** ****

FIN N=785 / TAD-bdry N=3,336

Control / ΔCTCF

j FIN & compartment

percentage

FIN N=785 / TAD-bdry N=3,336

Compartment A / B

k Both ΔWapl and ΔCtcf may cause FIN-skipping

- Ctcf / - Wapl

l Loop-gaining ATAC peaks in ΔCTCF cells vs. all ATAC peaks

	Enriched	Depleted
SP5	7.53	0.13
SP4	6.17	0.16
SP3	6.05	0.17
KLF3	5.93	0.17
ZBT17	5.74	0.17
KLF15	5.45	0.18
SP1	5.34	0.19
MAZ	5.10	0.20
KLF6	4.97	0.20
SP2	4.82	0.21
ZN281	4.82	0.21
SALL1	4.75	0.21
ZN148	4.60	0.22
WT1	4.42	0.23
EGR2	3.10	0.32
ARI5B	0.53	1.88
MSX3	0.54	1.84
GATA5	0.59	1.71
PROP1	0.59	1.69
HXA7	0.59	1.69
PAX6	0.61	1.65
FOXQ1	0.62	1.62
FOXD1	0.62	1.62
TGIF1	0.62	1.61
HXA10	0.62	1.60
CEBPB	0.63	1.59
SOX13	0.64	1.57
CDX1	0.65	1.53
FOXO4	0.66	1.52

−log10(qvalue): 0, 100, 200

Figure 4

Figure 5