Map functional insulators with cross-platform Hi-C meta-analysis

Wanying Xu[1,2,*], Jian Cui[1,2,*], Shanshan Zhang[1,2], Leina Lu[1,2], Xiaoxiao Liu[1], Yan Li[1,#], Fulai

Jin[1,3,4,#]

[1]Department of Genetics and Genome Sciences, School of Medicine, Case Western Reserve

University, Cleveland, OH 44106, USA

[2]The Biomedical Sciences Training Program (BSTP), School of Medicine, Case Western

Reserve University, Cleveland, Ohio 44106, USA

[3]Department of Computer and Data Sciences, Case Western Reserve University, Cleveland,

OH 44106, USA

[4]Department of Population and Quantitative Health Sciences, Case Comprehensive Cancer

Center, Case Western Reserve University, Cleveland, OH 44106, USA

[*] Equal contribution. [#] Co-senior authors.

Correspond with: yxl1379@case.edu (Y.L.), fxj45@case.edu (F.J.)

Summary

CTCF forms the boundaries of topologically associating domains (TADs) but their insulating

function in transcription is controversial. Here we report that functional insulators (FINs) shall be

determined not only by their ability to form static loops or boundaries, but also by the dynamic

context if their flanking sequences form new loops upon boundary loss. We performed a meta-

analysis of nine independent Hi-C and micro-C datasets upon acute CTCF- or cohesin protein-

depletion and found that only at several hundred loci, newly gained enhancers-promoters (E-P)

loops can be reproducibly observed upon CTCF depletion. The new E-P loops are cohesin-

dependent and associated with recurrent gene activation. These findings allow us to map FINs

and define their direct target genes genome-wide. FINs are mostly present in euchromatin, do

not overlap with TAD boundaries, and can be perturbed by *Wapl*-depletion or CTCF-

displacement. We therefore conclude that FINs, but not TAD boundaries, are *bona fide*

insulators.

Introduction

It is generally accepted that TADs confine the communication between genes and enhancers for transcriptional control(*1-6*). Most of the TADs are maintained by CTCF and cohesin, which are also known as insulator proteins. Cohesin is a multimeric complex consisting of *SMC1A*, *SMC3*, *RAD21* and one SA subunit (*STAG1* or *STAG2*). TAD forms when a DNA loop extrudes out of the cohesin complex ring before blocked by CTCF binding sites in convergent orientation, and cohesin serves as an enzymatic engine for this process(*7-15*). Deletion of CTCF or cohesin core subunit RAD21 leads to near complete loss of TADs(*16, 17*). Cohesin function is also regulated by cohesin loading complex MAU2/NIPBL(*18*) and cohesin release factor WAPL(*19-21*): TADs disappear upon loss of NIPBL(*22, 23*), while loss of WAPL causes bigger loops because cohesin can pass through CTCF blocks due to prolonged loop extrusion(*23, 24*).

Insulators are canonically defined as the DNA elements that inhibit gene expression when put between enhancers and promoters(*25-27*). The discovery of thousands of CTCF-demarcated TADs immediately suggests a model that CTCF sites at TAD boundaries are enhancer-blocking insulators, which is supported by genetic experiments focusing on selected loci(*28-32*). However, this model is challenged by a genome-wide Hi-C study in mESCs (Nora *et al.*, **Fig. 1a**) which only observed mild transcription changes despite near complete loss of TADs upon CTCF-depletion(*16*). This conclusion is reproduced in two other genome-wide studies done with *in situ* Hi-C and micro-C(*33, 34*) (**Fig. 1a**). Notably, Kubo *et al*. reported a role of CTCF in mediating E-P interactions(*33*), and Hsieh *et al.* reported short-range cohesin-independent cross-TAD E-P interactions(*34*). Taken together, although all these studies questioned the insulating function of TAD boundaries, they did not address the fundamental question about the whereabouts of the real enhancer-blocking insulators.

In this study we demonstrated that "functional insulators" (FINs) can be identified by examining if their flanking regions gain E-P interactions upon CTCF depletion. Challenges exist to map FINs with Hi-C-like genome-wide technologies: (i) most of the public Hi-C datasets do not have billion-scale read depth usually required for loop-level analysis (**Fig. 1a**); (ii) E-P interactions are generally weaker than CTCF loops in Hi-C and therefore harder to detect; (iii) the gain of E-P loop signal can be subtle, requiring robust quantitation of loop strength; (iv) it is difficult to rule out the possibility that some loop changes are secondary effects after prolonged CTCF or cohesin depletion(*16, 34*). These challenges can be addressed by comparing independent datasets, but existing CTCF-depletion data are from different Hi-C protocols, which perform very

differently in loop-level analyses with conventional pipelines(*35*). We previously developed

HiCorr and *DeepLoop* pipelines to allow robust comparison of loop profiles from orthogonal Hi-C

and micro-C platforms, even when sequencing depth is low(*36, 37*). Here we use *DeepLoop* to

perform a meta-analysis of orthogonal Hi-C datasets to map recurrent DNA rewiring events

upon depletion of CTCF and other cohesin proteins (**Fig. 1a**), thus generate a reliable genome-

wide map of enhancer-blocking insulators.

Results

***DeepLoop* reveals insulated E-P loops from CTCF-degron Hi-C data**

We collected nine pairs of independent Hi-C or micro-C datasets from six publications in which

CTCF(*16, 33, 34*), RAD21(*34, 38, 39*), or WAPL(*34, 38, 40*) were acutely depleted in three

different mESC lines. The number of contacts in these Hi-C/micro-C datasets are highly variable

ranging from 43 million to ~1 billion. For every target protein, the choice of degron methods (AID

or dTAG) and the timespan of protein degradation (range from 3hr to 3 days) are also variable

(**Fig. 1a**). As shown in the examples in **Fig. 1b**, for Hi-C and *in situ* Hi-C studies, conventional

ICE-normalization only support compartment- or TAD-level analysis at 20kb or lower resolution

due to the lack of read depth. In contrast, *HiCorr*-based bias correction(*36*) followed by

DeepLoop enhancement(*37*) reveals detailed loop profiles at 5kb resolution (**Fig. 1b**). From this

example, we observed that the enhanced loop profiles in wildtype mESCs from independent

studies are rather similar (**Fig. 1b**, **Supplementary Fig. 1**), supporting the robustness of

DeepLoop in cross-platform analysis regardless of read depth(*37*). On the other hand, loss of

CTCF, *Rad21*, and *Wapl* leads to distinct loop profiles (**Fig. 1b**).

We firstly focused on the CTCF-degron Hi-C data from Nora *et al*. In the example of **Fig. 1b** (top

two rows), with conventional ICE-normalization, we observed dampened TAD structure at 20kb

resolution as the original study reported(*16*), but at 5kb resolution, loop signal from ICE-

normalized heatmaps are weak. From the *DeepLoop* enhanced heatmaps, we can clearly

observe that multiple CTCF loops are substantially weakened upon CTCF depletion. However,

the most notable change in this region is the gain of a long-range loop (**Fig. 1b**, second row,

loop pixels in dashed circle) connecting *Notch1* gene to a distal enhancer region with multiple

H3K27ac peaks (**Fig. 1c**). Consistent with gain of E-P loop, *Notch1* is upregulated upon CTCF-

depletion (**Fig. 1c**). The gain of loop signal at this locus is also observable in *Wapl*-depleted

cells (**Fig. 1b**, bottom two rows) and two other CTCF-degron studies (**Supp. Fig. 1**), but not in

the *Rad21*-depleted cells (**Fig. 1b**, 3rd and 4th rows). These observations suggest that the two

CTCF loop anchors at the TAD boundary region (**Fig. 1c**, highlighted in pink) are blocking the

Notch1-enhancer interaction. This example also demonstrated that *DeepLoop* may detect

recurrent insulating events from CTCF-degron Hi-C data without the need for ultra-high

sequencing depth.

Hi-C meta-analysis reveals recurrent E-P loop-gaining events upon CTCF depletion

To perform a genome-wide analysis of insulating events, we call 300K~500K loop pixels from

each CTCF-degron dataset and use the pixel intensity in *DeepLoop* heatmaps to measure the

loop strength (**Fig. 1d, Supp. Fig. 2**). Notably, *DeepLoop* outputs are enhanced from ratio

(obs/exp) heatmaps instead of normalized contact frequency used by ICE/KR. We previously

showed that this approach promises cross-platform convergence between orthogonal Hi-C and

micro-C data(*37*).

In all CTCF-degron studies, the scatter plots are clearly down-skewed reflecting a massive loss

of chromatin loops upon CTCF deletion (**Fig. 1d, Supp. Fig. 2a, d**, left panels). Using a simple

fold-change cutoff, we called loop pixels that lose strength ("lost"), retain strength ("retained"), or

gain strength ("gained") from each dataset (**Fig. 1d, Supp. Fig. 2a, d**, middle panels). We

further classified the loop pixels based on their epigenetic marks and found that CTCF loops

dominate the "lost" pixels, while the "retained" and "gained" pixels are enriched with enhancer or

promoter marks (**Fig. 1e, Supp Fig. 2b, e**). Interestingly, in all three studies the "gained" loops

are much bigger than the "retained" loops (**Fig. 1d, Supp. Fig. 2a, d**, middle panels),

suggesting that these "gained" E-P loops form through loop extrusion (more discussions below).

We also examine the robustness of the "lost", "retained", and "gained" pixels across

independent Hi-C/micro-C studies. As expected, the "lost" pixels from any study generally also

lose strength in the other two studies (**Fig. 1f, Supp Fig. 2c,f**, first row). However, the

consistency of "retained" loop pixels is study specific: (i) although the "retained" loop pixels from

Nora *et al.* are mostly also "retained" in Kubo *et al.*, some of them lose strength in Hsieh *et al.*

(**Fig. 1f**, second row); (ii) most of the "retained" loop pixels from Kubo *et al.* lose strength in the

other two studies (**Supp. Fig. 2c**, second row); (iii) the "retained" loops from Hsieh *et al.* remain

"retained" in both Nora *et al.* and Kubo *et al.* (**Supp. Fig. 2f**, second row). Since incomplete

CTCF depletion causes the leftover CTCF loops(*16*), these observations indicate that CTCF

depletion in Kubo *et al.* is less complete than the other two studies; Hsieh *et al.* achieves the

best CTCF depletion, perhaps because it uses N-terminal degron system (**Fig. 1a**).

Interestingly, the "gained" pixels from any one of the three studies generally also gain strength

in the other two studies (**Fig. 1f, Supp Fig. 2c,f,** third row). The only exception is that some

"gained" loop pixels from Kubo *et al.* lose strength in Hsieh *et al.* (**Supp Fig. 2c**). Again, this is

likely because Kubo *et al.* retains more CTCF loops due to incomplete CTCF degradation.

Nevertheless, the "gained" pixels from Nora *et al.* and Hsieh *et al.* do gain strength in Kubo *et al.*

(**Fig. 1f, Supp Fig. 2f**), supporting the consistency between the three studies. Notably, E-P loop

gaining events are largely neglected in the original Hi-C/micro-C studies because the changes

are often too subtle to distinguish from background with conventional analytical tool (*e.g.*, **Fig.**

1b). Our meta-analyses prove that although there are only a small number of newly gained E-P

loops, they are reliable consequences of CTCF-depletion; affected genes such as *Notch1* in

Fig. 1 are likely to be the direct targets of insulators.

Newly gained E-P loops in CTCF-depleted cells are cohesin dependent.

We also performed meta-analyses of the *Rad21*- and *Wapl*-depletion Hi-C datasets. Ablation of

Rad21 also causes massive loss of CTCF loops, while the "retained" and "gained" loops are

also enriched with E-P loops (**Fig. 2a-b, Supp. Fig. 3**). However, there are important

differences between *Rad21*- and *CTCF*-depleted cells. (i) The "retained" and "gained" loop

pixels in *Rad21*-depleted cells are much smaller (mostly within 100kb) than those in *CTCF*-

depleted cells, indicating that cohesin-independent loops are short-range (**Fig. 2a, Supp. Fig.**

3a, d, right panels). (ii) Unlike in CTCF-depleted cells, the "gained" loop pixels are NOT

recurrent between the three *Rad21*-degon datasets (**Fig. 2c, Supp. Fig. 3c, f**, last row). (iii) The

"retained" and "gained" loop pixels in *Rad21*-depleted cells are more enriched with H3K36me3

mark, suggesting transcription associated chromatin interactions (**Fig. 2b, Supp. Fig. 3b, e, h**).

(iv) The "gained" loops from Rhodes *et al.* are enriched with H3K27me3 mark and tends to be

longer (**Supp. Fig. 3a, b**, right panels), which is consistent with the conclusion from the original

study(*39*). However, this phenomenon is specific to Rhodes *et al.* and it is not observed in the

other two *Rad21*-degron studies.

Conversely, ablation of cohesin unloading factor *Wapl* generates up-skewed scatterplots in all

three studies (**Fig. 2d, Supp. Fig. 4a, d**). The skewness of Hsieh *et al.* is less apparent,

probably because Hsieh *et al.* performed *Wapl* depletion for much shorter time (3hrs) than Liu *et*

al. (24hrs) and Kriz *et al*. (72hrs) (**Fig. 1a**). In all *Wapl*-degron studies, the "lost", "retained", and

"gained" loops are almost equally enriched with CTCF and E-P loops (**Fig. 2e, Supp. Fig. 4b,**

e). Consistent with the model that *Wapl*-depletion stabilizes chromatin-bound cohesin and

promote loop extrusion(*23, 24*), the "gained" loops are significantly bigger than "lost" and

"retained" loops in all three studies (**Fig. 2d, Supp. Fig. 4a, d**, right panels). The "lost",

"retained", and "gained" loop pixels are generally consistent between studies (**Fig. 2f, Supp.**

Fig. 4c,f) despite some discrepancy for "lost" loop pixels from Hsieh *et al.* (**Supp. Fig. 4f**), again

likely because Hsieh *et al.* only depleted *Wapl* for a short period of time.

We next take the "lost", "retained", and "gained" loop pixels from *CTCF*-degron studies and

examine how they change in *Rad21*- and *Wapl*-degron data. Firstly, nearly all *CTCF*-dependent

loops ("lost" in *CTCF*-depleted cells) are also *Rad21*-dependent (**Fig. 2g, Supp Fig. 5a,b**, first

row). Secondly, the *CTCF*-independent loops ("retained" in CTCF-depleted cells) include both

cohesin-dependent and cohesin-independent fractions, since some of them lose strength in

Rad21-degron data but some do not (**Fig. 2g, Supp Fig. 5a, b**, second row). More of the CTCF-

independent loops from Hsieh *et al.* are also cohesin independent (**Supp Fig. 5b**, second row),

again suggesting that Hsieh *et al.* achieves more complete CTCF-depletion. Finally, the

"gained" loops in CTCF-depleted cells do not gain strength in *Rad21*-depleted cells (**Fig. 2g,**

Supp Fig. 5a, b, third row). These results strongly indicate that the recurrently gained E-P

interactions upon CTCF depletion are created by loop extrusion.

Interestingly, the "gained" loops in CTCF-depleted cells have apparent upward biases in the

scatter plots from *Wapl*-degron data (**Fig. 2h, Supp Fig. 5b, d**, third row). These results are

consistent with the example in **Fig. 1b** and suggest that *Wapl*-depletion can perturb insulator

function by releasing short-range CTCF loops and allows insulated E-P interactions to skip

insulators (more analysis below).

Define, map, and characterize functional insulators (FINs) in mESC genome

Among the total 8,141 "gained" loop pixels from all three CTCF-degron Hi-C/micro-C datasets,

4,944 pixels are recurrent in at least two studies with a median size ~250kb (**Fig. 3a**). In each

dataset, the "gained" loop pixels are found in ~300 loci and there are substantial overlaps

between independent studies: 174 loop-gaining loci are shared by all three studies and 304 are

shared by at least two studies (**Fig. 3b**). We hypothesize that these recurrent E-P loop-gaining

loci are caused by loss of CTCF insulation; indeed, 4,665 out of the 4,944 loop-gaining pixels

(94%) are insulated by at least one CTCF loop in wildtype cells (**Fig. 3c**). We further curated the

reproducible loop-gaining loci based on the types and numbers of CREs involved (**Fig. 3d**).

Nearly all the curated loci (283 of 299) gain E-P loops (2 loci gain CTCF contacts and the others

cannot be clearly classified). We further classified the E-P loop-gaining loci into three categories

(**Fig. 3d**). In *Type I*, one of the two E/P sites is insulated in a CTCF loop while the other site is

not; *Type II* involves at least two CTCF loops insulating two E/P-sites. In contrast, the gained E-

P loops in *Type III* do not have insulating CTCF loops. *Type I* and *II* represents most of the E-P

loop gaining loci (263 out of 283), while only 20 loci belong to *Type III*, again suggesting that

most of the loop gaining loci are due to the removal of CTCF insulation.

We therefore propose to define "functional insulators" (FINs) as the CTCF loop anchors that are

actively blocking the interactions between flanking *cis*-regulatory elements (CREs) (model *iii* in

Fig. 3e). The definition of FIN is conceptually analogous to the conventional method to define

enhancer-blocking insulators by transcriptional effects (model *i* in **Fig. 3e**), but distinct from the

TAD-based definition of insulators (model *ii* in **Fig. 3e**). There are clearly much fewer FINs than

TAD boundaries (**Fig. 3f**): only 13% of the FINs overlap TAD boundaries (**Fig. 3g**). These

observations indicate that most FINs are inside TADs and conversely, loss of TAD boundaries

usually does not cause loop formation across their flanking regions. We also found that most

(96%) of the FIN loci are within compartment A (**Fig. 3h**), indicating that active E-P blocking

mainly happens within euchromatin region.

Cohesin degron Hi-C data further support a cohesin-dependent mechanism of FINs. We found

that almost none of the curated loop-gaining events appears in the *Rad21*-degron cells (**Fig.**

3d), indicating cohesin dependency. Interestingly, a significant fraction of the *Type I* and *II*, but

not the *Type III* "gain of E-P loop" loci, can be also observed in *Wapl*-degron datasets (**Fig. 3d**).

This phenomenon can be explained by previous findings that *Wapl* deletion allows loop

extrusion to skip CTCF blocks(*23, 24*). As shown in the carton in **Fig. 3i**, because *Wapl*-

depletion stabilizes chromatin-bound cohesin and leads to longer loop extrusion, previously

insulated enhancers and promoters can interact inside the bigger CTCF loop due to "insulator

skipping", creating the same E-P loop as observed in CTCF-depleted cells. Taken together,

these results support FINs as *bona fide* insulators of cohesin-mediated loop extrusion.

FINs are associated with consistent early transcription responses upon CTCF depletion.

We found that in all CTCF-degron studies FIN proximal regions (+/- 100kb) are 2~3 fold more
enriched with DEGs than TAD-boundary proximal regions, and the "gained" loop anchors are 10
times more enriched with DEGs (**Fig. 4a**). Furthermore, early response DEGs are clearly more
enriched at gained loop anchors (**Fig. 4b**), suggesting that FIN-blocked enhancers and
promoters are the direct targets of insulators. 16.8% of FINs (49 out of 299) are blocking 43
super-enhancers (out of 231, or 19%) (**Fig. 4c**). As expected, genes gaining super-enhancer
interactions are more likely to be upregulated than genes gaining interactions with regular
enhancers (**Fig. 4d**).

Using common cutoffs (2-fold for RNA-seq and 1.5-fold for NET-seq, $p < 0.05$), each of the
CTCF-degron studies identified a few hundred early response DEGs, and the reproducibility
between independent studies is limited due to complex reasons (**Fig. 4e**). Using NET-seq,
Hsieh *et al*. identified more DEGs than RNA-seq despite shorter time of CTCF depletion (3hr),
but it remains possible that some DEGs are caused by indirect mechanisms. Among the two
RNA-seq studies, Kubo *et al.* found fewer DEGs than Nora *et al.*, and most of the DEGs from
Kubo et al. are also observed in Nora *et al*. (**Fig. 4e**), consistent our conclusion above that the
CTCF-depletion in Kubo *et al.* is less complete. We therefore focused on the reproducible DEGs
between Hsieh *et al.* and Nora *et al.* as they represent a more reliable set of CTCF direct
targets. Interestingly, although downregulated DEGs are more reproducible than upregulated
DEGs between the two studies (**Fig. 4f**, grey bars), near FINs (at the anchors of gained E-P
loops) the upregulated DEGs are significantly more reproducible than downregulated DEGs
(**Fig. 4f**, red bars). This observation supports transcription activation as the primary
consequence of insulator loss, consistent with the convention concept that insulators block
enhancer-mediated gene activation.

In **Fig. 4g** we plotted 49 upregulated DEGs at FIN loci as direct targets of insulators (from Hsieh
et al. and Nora *et al.*), among them 34 are considered as "high-confidence" targets supported by
both studies. **Fig. 4h** shows one representative example, in which we observed gain of loop
between two enhancer clusters upon CTCF depletion, leading to the upregulation of *Cdiptos*
gene. Similarly, we also plotted 37 reproducibly down-regulated DEGs near FINs supported by
both studies (**Fig. 4i**). Strikingly, 18 of the recurrent down-regulated DEGs are from
protocadherin-γ, a well-known insulator loci (**Fig. 4i,j**). Protocadherin proteins play key roles in
the formation of synaptic connections between neurons and the *Pcdh-*γ genes are distinguished
by their variable promoters (exon1): γ*a*'s and γ*b*'s are stochastically expressed in individual

neurons and the γc's are constitutively expressed(*41-44*). In mESCs, the promoters of γa's and

γb's are occupied by CTCF, H3K4me3, and the repressive H3K27me3 mark (**Fig. 4j**). An

enhancer cluster marked by H3K27ac is present at the 3' of γc promoters (**Fig. 4j**, highlighted in

green) forming loop interactions with the promoters of γa's and γb's, which are restricted by a

FIN with multiple CTCF peaks (**Fig. 4j**, blue arches). Upon CTCF depletion, the enhancer

interactions are weakened, and the *Pcdh-γ* promoter loops extend into a distal enhancer-poor

region marked by H3K27me3, which explains the downregulation of γa and γb genes. Taken

together, these results demonstrated the transcription consequences of FINs and mapped their

direct targets in mESCs.

Validate FINs with multiplexed CTCF displacement assay

To validate the *cis*-regulatory functions of FINs, we need to verify if the gained loops are truly

due to the loss of insulation. Because a FIN often involves multiple CTCF binding sites that

contribute to the insulating function collectively, we need to simultaneous perturb multiple CTCF

sites in a relatively big region. To achieve this, we combine a multi-sgRNA delivering system

name CARGO(*45*) with a dCas9-based CTCF-displacement method(*43, 46*) to perturb multiple

CTCF binding sites in the same cells. In this approach, 4-10 sgRNAs exactly overlapping the

cognate motifs of multiple CTCF binding sites are expressed from the same CARGO plasmid.

When co-transfected with a dCas9 construct, the sgRNAs can displace CTCF from genome at

multiple target sites simultaneously (**Fig. 5a-b**).

We used this approach to validate the gain-of-loop events at two FIN loci. The first FIN is close

to *Cdiptos*, a recurrent upregulated DEG (**Fig. 5c**). This example belongs to a simpler type of

FIN in which the newly gained loop (**Fig. 5c**, red arches) is blocked by only one FIN (**Fig. 5d**).

We used one CARGO construct targeting 4 CTCF motifs and confirmed with CUT&RUN that we

successfully weakened the target CTCF peaks but did not affect other CTCF peaks in this

region (**Fig. 5c**, highlighted in pink). We performed 4C-seq and confirmed the gain of long-range

chromatin interactions consistent with Hi-C data (**Fig. 5c**, highlighted in green). We also

validated with RT-qPCR that *Cdiptos* is upregulated upon displacement of target CTCF binding

sites (**Fig. 5e**). The second locus is more complex as the newly gained loop is blocked by more

than two FINs (**Fig. 5f,g**). For this locus we used one CARGO construct to target 9 CTCF motifs

and successfully weakened the target CTCF peaks (**Fig. 5f**, highlighted in pink). Again, we

confirmed the gain of loop interactions with 4C-seq (**Fig. 5f**, highlighted in green), and the

upregulation of *Mesp1* gene since the new loops bring distal enhancers to its proximity. Taken

together, these experiments verified that FINs execute their insulating functions in *cis*.

Discussion

Discussion

In this study we propose to define functional insulators, or FINs, as CTCF binding sites that

actively block the loop formation between flanking *cis*-regulatory elements (**Fig. 3e**). Mapping

FINs requires robust detection of newly gained E-P loops, which is challenging for Hi-C analysis

because E-P loops are usually weak and hard to detect. With a meta-analysis of three

independent CTCF-degron Hi-C datasets, we demonstrate that the subtle gain of loop signal

detected by *DeepLoop* are quantitatively robust across different Hi-C or micro-C datasets,

therefore proves the feasibility to map FINs reliably with Hi-C at intermediate read depth.

The concept of FIN is distinct from the common assumption that TAD boundaries are insulators.

In fact, FINs are usually inside the TAD, and most TAD boundaries do not gain cross-boundary

loops upon CTCF depletion. Importantly, there are only a few hundred FINs in mESCs, which

readily explains why CTCF-depletion only cause mild changes of transcriptome. It should be

also noted that the gained E-P loops do not always predict significant gene activation due to

complex reasons(*47*). Furthermore, target gene expression can be quantitative tuned by

multiple enhancers in an additively fashion, and each single enhancer may only have a small

effect size(*48*). Nevertheless, we showed that FINs better explain DEGs, especially the

recurrent upregulated DEGs in CTCF-depleted cells. This is consistent with the classic definition

of insulators as DNA elements that repress enhancer-mediated gene activation.

It remains unclear why only a small subset of CTCF loop anchors are FINs, *i.e.*, why is gain-of-

loop only observed at some loci but not others? Our analysis provided some hints. Firstly, meta-

analysis with *Rad21*- and *Wapl*-degron Hi-C data shows that the newly gained E-P loops in

CTCF-depleted cells are cohesin-dependent and governed by cohesin regulators. It is therefore

plausible that local cohesin activity is a key determinant. Secondly, the fact that almost all FINs

are in compartment A (euchromatin) is also consistent with high cohesin activity because

cohesin and its loading factors are known to co-localize with promoters and enhancers(*49*).

Finally, since genome compartmentalization is a highly dynamic 3D genome feature, we

speculate that FINs are also likely to be cell type- or cell state-specific. It will be interesting to

further explore the context-specific roles of FINs in health and diseases.

Figure Legend

Figure 1. *DeepLoop* reveals recurrent gain of E-P loops from independent CTCF-degron Hi-C datasets.

a, A list of published Hi-C and micro-C studies with acute depletion of *Ctcf*, *Rad21*, or *Wapl* in mESCs. **b,** Contact heatmaps of the CTCF-, Rad21-, and Wapl-degron Hi-C data at *Notch1* locus with *ICE*, *HiCorr* and *DeepLoop*. Red dashed circle: gained loops. **c**, Genome browser tracks showing ChIP-seq data and Hi-C loops from Nora *et al.* at *Notch1* locus. Blue highlighted region showing Notch1 gene which is at the anchor of a gained loop. Pink region highlights the potential insulator region that blocks the formation of *Notch1* E-P loop. RNA-seq results are also shown for *Notch1* gene in WT (black) and CTCF-depleted cells (red). Bar plot: quantitation of *Notch1* expression from RNA-seq. P-value: one-side Wald test provided by DE-seq2. **d-f,** Meta-analysis of independent CTCF-degron Hi-C datasets. **d**, Left: scatterplot comparing the strength of loop pixels between WT and CTCF-depleted mESCs; middle: the same scatterplot as left but highlight the lost (blue), retained (black), and gained (red) loop pixels; right: violin plot comparing the size of loop pixels. **e,** Pie charts showing the components of lost, retained and gained loop pixels based on epigenetic marks. **f**, Scatter plots showing the consistency of lost, retained and gained loop pixels between three CTCF degron studies. Plots in the left column highlight the lost, retained, and gained loop pixels from Nora *et al*., the right two columns show the quantitative changes of the corresponding pixels in *Kubo et al.* or *Hsieh et al*.

Figure 2. Meta-analysis of CTCF- and cohesin-degron Hi-C datasets reveals the fate and mechanisms of dynamic loops.

a-c, Meta-analysis of independent *Rad21* degron Hi-C datasets. **a,** Left and middle panels define the lost, retained, and gained loop pixels upon *Rad21*-depletion from the study by Kriz *et al.*, and the right panel compares their loop size distribution. **b,** Pie charts showing the components of lost, retained and gained loops based on epigenetic marks. **c,** Scatter plots testing if the lost, retained and gained loop pixels from Kriz et al. are consistent in the other two *Rad21* degron studies. **d-e,** The same meta-analysis of three independent *Wapl*-degron Hi-C datasets. **g,** The lost, retained, and gained loop pixels from one CTCF-degron Hi-C data (Nora

et al.) are plotted onto Hi-C scatter plots from *Rad21-* or *Wapl*-degron studies to examine their

quantitative changes.

Figure 3. Characterization of functional insulators (FINs) in mESC genome.

a, Heatmap showing the union of all gained loop pixels from three CTCF-degron Hi-C datasets.

Every row represents one CTCF-degron study and every column represents one loop pixel.

Loop pixels are organized into clusters based on their reproducibility. The number of loop pixels

in each cluster and their median size are also shown. Color scale indicates the increase of loop

signal in log scale. **b,** Venndiagram showing the consistence of *FIN* loci identified from three

datasets. **c,** Pie chart showing that most of the gained loops in CTCF-depleted cells have at

least one CTCF loop anchor in between in wildtype cells. **d,** A table summarizing the

classification and recurrence of loop gaining loci between independent CTCF- or cohesin-

degron studies. **e,** Cartoons summarizing three possible ways to define insulators. **f,** Ideogram

showing the locations of FIN loci from three studies, TAD boundaries, and compartments on

chromosome 1. FIN from three different CTCF-degron studies are shown in different color, and

the color scale shows the fold increase of gained loop in log scale. **g,** Pie chart showing the

percentage of FINs that overlap TAD boundary. **h,** Stack bar plot comparing the proportion of

FINs and TAD boundaries in compartment A or B. **i,** A model to explain why some of the newly

gained E-P loops in CTCF-depleted cells are also observed in Wapl-depleted cells.

Figure 4. FINs are associated with consistent early transcriptional consequences in

independent CTCF-degron studies.

a, Bar plots comparing the density of DEGs upon CTCF-depletion at different type of genomic

regions. y-axis indicates the number of DEGs per Mb. **b,** Enrichment of DEGs at different types

of genomic regions compared to whole genome as background. X-axis: time of IAA treatment

(hrs) in each study; y-axis: fold enrichment. Each panel represent one CTCF-degron study. **c,**

Left: pie chart showing the percentage of FINs that are blocking super enhancers; Right: pie

chart showing the percentage of super enhancers that are blocked by *FIN*. **d,** Boxplot showing

the expression changes of genes gaining enhancer loop. Red: genes gaining super-enhancer

loop; Blue: genes gaining regular enhancer loop. *, $p < 0.05$, Wilcoxon signed-rank test. **e,** Venn

diagrams showing the overlap of all up- (left panel) and down-regulated (right panel) DEGs

between three CTCF-degron studies. **f,** Bar plot comparing the reproducibility (measured by

Jaccard Index) of up- or down-regulated DEGs between two CTCF-degron studies (Hsieh *et al.*

and Nora *et al.*). Gray bars: all DEGs; blue bars: DEGs at the anchors of lost loops; red bars:

DEGs at the anchors of gained loops. P-values are from binomial test. **g,** Heatmap showing the

union of upregulated DEGs at the anchors of gained loops from *Hsieh et al* and *Nora et al.*;

DEGs are clustered based on consistency. **h,** Left panel shows the browser tracks near *Cdiptos*

locus and right panel shows the Hi-C heatmaps in the same region from *Hsieh et al* and *Nora et*

al. The direction of CTCF motifs are indicated by triangles. Blue arches (left) and blue dashed

circles (right): lost loop pixels; red arches and red dashed circles: gained loop. Regions

highlighted in green: enhancers that gained loops; the region highlighted in pink: a CTCF loop

anchor is defined as *FIN*. **i,** Similar as **g**, a heatmap showing the union of down-regulated DEGs

at the anchors of gained loops from *Hsieh et al.* and *Nora et al*. Among these genes 18 genes

are from the Pcdh-γ locus (shown as a separate heatmap). **j,** Left panel shows the browser

tracks near Pcdh-γ locus and right panel shows the Hi-C heatmaps from two CTCF-degron

studies.

Figure 5. Validate FINs with multiplexed CTCF displacement assay.

a, Scheme of the CARGO vector that can express multiple sgRNAs. In this system, the

sequences of *n* sgRNAs are broken into left and right halves and inserted into (n+1) minicircles.

The *i*-th minicircle contains the right half of the (*i-1*)-th sgRNA and the left half of the *i*-th sgRNA

sequences divided by a *BpiI* restriction site. Final CARGO vector can be assembled by ligating

the re-linearized minicircles and vector backbone; each sgRNA is driven by a U6 promoter. **b,**

Co-transfect CARGO vector and a dCas9-expressing plasmid can displace multiple CTCF

binding sites in the same cell. **c,** Validation of a FIN at the *Cdiptos* locus. From the top, 1st and

2nd tracks are H3K4me3 and H3K27ac ChIP-seq data showing the promoter and enhancers in

mESC. 3rd and 4th tracks show CTCF ChIP-seq data before and after targeting 4 CTCF motifs in

FIN (highlighted in pink) with CARGO. 5th and 6th are Hi-C arches (*DeepLoop* pixels with data

from Nora *et al.*) before and after CTCF-depletion (blue: lost loops; red: gained loops). The

bottom 3 tracks show 4C-seq data (bait highlighted in yellow) before and after targeting CTCF

sites; regions highlighted in green gain 4C signal. **d,** A model cartoon showing that the gained

loop at this FIN locus only involves one CTCF loop. **e,** RT-qPCR results showing that *Cdiptos* is

upregulated after CTCF displacement. ** p<0.01; one-side t-test. **f-h,** Validation of a FIN at

Mesp1 locus similar as shown **c-e**. Note the gained loop at this FIN locus involves at least two

CTCF loops (**g**).

Supplementary figure legend

Supplementary Figure 1. Recurrent gain of E-P loop at *Notch1* locus between
independent CTCF-degron Hi-C studies.

a, Contact heatmaps of the CTCF-degron Hi-C data by Kubo *et al.* at *Notch1* locus with *ICE*,
HiCorr and *DeepLoop*. Red dashed circle highlights the same locations where we observed a
gained E-P loop in the contact heatmap from Nora *et al.* (shown in **Fig. 1b**). **b**, Genome browser
tracks showing ChIP-seq data and Hi-C loops from Kubo *et al.* Blue highlighted region indicates
Notch1 gene. Pink region highlights the potential insulator region that blocks the formation of
Notch1 E-P loop. **c,** RNA-seq results from Kubo *et al.* are also shown as bar plot. p-value: one-
side Wald test provided by DE-seq2. **d-f,** The same analysis as **a-c** at *Notch1* locus using data
from Hsieh *et al.* Note that Hsieh *et al.* used mNET-seq for transcriptome profiling.

Supplementary Figure 2. Meta-analysis of independent CTCF-degron Hi-C datasets to
test the consistency and reveal the epigenetic features of dynamic loops.

a, Using the *in situ* Hi-C data from the CTCF-degron study by Kubo *et al.*, left: scatterplot
comparing the strength of loop pixels between WT and CTCF-depleted mESCs; middle: the
same scatterplot as left but highlight the lost (blue), retained (black), and gained (red) loop
pixels; right: violin plot comparing the size of loop pixels. **b,** Pie charts showing the components
of lost, retained and gained loop pixels based on epigenetic marks. **c**, Scatter plots in the left
column highlight the lost, retained, and gained loop pixels from Kubo *et al.*, the right two
columns show the quantitative changes of the corresponding pixels in Nora *et al.* or Hsieh *et al.*
d-f, The same analyses as **a-c** are performed with CTCF-degron micro-C data by Hsieh *et al.*

Supplementary Figure 3. Meta-analysis of independent *Rad21*-degron Hi-C datasets to
test the consistency and reveal the epigenetic features of dynamic loops.

a, Left and middle panels define the lost, retained, and gained loop pixels upon *Rad21*-depletion
from the study by Rhodes *et al.*, and the right panel compares their loop size distribution. **b,** Pie
charts showing the components of lost, retained and gained loops based on epigenetic marks.
c, Scatter plots testing if the lost, retained and gained loop pixels from Rhodes *et al.* are
quantitatively consistent in the other two *Rad21*-degron studies. **d-f,** The same analyses as **a-c**
are performed with *Rad21*-degron micro-C data by Hsieh *et al.*

Supplementary Figure 4. Meta-analysis of independent *Wapl*-degron Hi-C datasets to test the consistency and reveal the epigenetic features of dynamic loops.

a, Left and middle panels define the lost, retained, and gained loop pixels upon *Wapl*-depletion from the study by Kriz *et al.*, and the right panel compares their loop size distribution. **b,** Pie charts showing the components of lost, retained and gained loops based on epigenetic marks. **c,** Scatter plots testing if the lost, retained and gained loop pixels from Kriz *et al.* are quantitatively consistent in the other two *Wapl*-degron studies. **d-f,** The same analyses as **a-c** are performed with *Wapl*-degron micro-C data by Hsieh *et al.*

Supplementary Figure 5. Meta-analysis reveals the fate of dynamic loops from CTCF-degron studies upon depletion of cohesin factors.

a, The lost, retained, and gained loop pixels from one CTCF-degron Hi-C data (Kubo *et al.*) are plotted onto Hi-C scatter plots from *Rad21*- or *Wapl*-degron studies to examine their quantitative changes. **b,** Same analyses are done with the lost, retained, and gained loop pixels from the CTCF-degron Hi-C data by Hsieh *et al.*

Method

Experiments

Cell culture

Mouse embryonic stem cells (E14Tg2a) were cultured in DMEM+Glutamax (ThermoFisher , #10566-016) with the addition of 15% Fetal Bovine Serum (ThermoFisher, # SH30071.03), 550µM β-mercaptoethanol (ThermoFisher, #21985-023), 1mM Sodium Pyruvate (ThermoFisher, #11360-070), 1X non-essential amino acids (ThermoFisher, #11140-50), and 10^4 U of Leukemia inhibitory factor (Millipore, #ESG1107). The cells were maintained at a density of $0.2\text{-}1.5\times10^5$ cells/cm^2 and were passaged every 24-48 hours using TrypLE (ThermoFisher, #12563011) on dishes coated with 0.1% gelatin (Millipore, # ES-006-B) at 37°C and 5% CO2. The medium was refreshed daily when cells were not being passaged. Regular checks for mycoplasma infection were performed every 3-4 months, with negative results.

4C-seq

4C-seq was conducted following a published protocol(*50*). Briefly, 5 million cells fixed with 2%

formaldehyde, then quenched with 125 mM glycine. The fixed cells were lysed using a cell lysis

buffer containing 50 mM Tris-Cl pH 7.5, 150 mM NaCl, 5 mM EDTA, 0.5% NP-40, 1% Triton X-

100, and 1x protease inhibitor cocktail (Roche, Cat#11873580001) for 20–30 minutes on ice.

Following lysis, nuclei were collected by centrifugation at 2,500g for 5 minutes at 4°C and

washed once with 1x restriction enzyme buffer. The nuclei pellets were resuspended in 1x

restriction enzyme buffer and treated with 0.3% SDS for 1 hour at 37°C with shaking, followed

by an additional hour with 2.5% Triton X-100. Chromatin was then digested overnight with the

appropriate restriction enzyme at the recommended temperature while rotating in an air bath.

The specific restriction enzymes used for each locus are detailed in **Supplemental Table 6**.

After digestion, the enzymes were inactivated by heating at 65°C, and nuclei were ligated with

50 µl of T4 DNA ligase (Invitrogen, Cat# 15224–090) in a 7 ml ligation solution at 16°C

overnight. Reverse cross-linking was performed by treating the samples with proteinase K to

isolate the proximity-ligated DNA. The purified DNA was quantified and subjected to secondary

restriction enzyme digestion using roughly 1 unit of restriction enzyme per 1 µg of DNA at the

recommended temperature overnight. After enzyme inactivation, the samples were self-ligated

with T4 DNA ligase. The ligated DNA was recovered using sodium acetate and ethanol and

quantified using the Qubit dsDNA HS assay kit (Thermo Fisher, Cat# Q32851). The 4C

templates were amplified using designed primers to generate sequencing libraries. The primer

system was modified to be compatible with the Illumina Nextera system, using two sequential

PCRs. Locus-specific inverse PCR primers are listed in **Supplemental Table 6**. For each locus,

the 4C templates were amplified with locus-specific primers using 200 ng of template per

reaction, and products from five parallel amplifications were pooled to generate the final 4C

library. A 50 µl PCR product aliquot was purified with homemade Sera-Mag beads. One-fifth of

the purified DNA was used for a second PCR with primers N7xx and N5xx, identical to Illumina

Nextera sample preparation primers. The final products were purified and sequenced. Reads

from the first cutting site were used for data analysis.

Plasmid cloning

The dCas9 expression vectors used in this study were created using the Cas9 expression

vector backbone from the pX330 plasmid (Addgene; plasmid 42230) through the In-Fusion

cloning technique. The dCas9 genes were amplified from the pHAGE EF1α dCas9-KRAB

plasmid (Addgene; plasmid 50919) via PCR and independently cloned into the AgeI and EcoRI

sites of the pX330 plasmid, substituting the Cas9 ORF. Detailed information regarding the

primers can be found in **Supplementary Table 5**. All sgRNAs in this study were designed using

the CCTop-CRISPR/Cas9 target online predictor.

CARGO assembly

The CARGO array assembly follows a published protocol(*45*). The protocol consists of two main

steps: constant region preparation and array assembly. In the constant region preparation step,

the constant region with proper sticky ends is released by digesting the pGEMT-hU6-spSL

(provided by Dr. Joanna Wysocka) plasmid with BsmBI. The released constant region is then

separated from the pGEMT backbone on a 1% agarose gel and gel purified using a commercial

kit (QIAGEN, Cat#28704). Oligonucleotides (50-60 bp) with variable regions were ordered

through IDT (**Supplementary Table 5**). The forward and reverse oligos are annealed and

phosphorylated and then stored at -20°C.

In the array assembly step, the annealed chimeric oligo is mixed with the digested constant

region, and a minimum of 4-hour ligation with high-concentration T4 ligase (Thermofisher

Scientific, Cat#EL0013) is performed. The ligation step yields ligation products, referred to as

minicircles. After the stoichiometry check, individual minicircles are treated with plasmid-safe

exonuclease to clean up the un-ligated products. Then, the plasmid-safe exonuclease treated

individual minicircles are pooled together and purified. The pooled circle mixes are assembled

directly into the digested destination vector pX332-mCherry (provided by Dr. Joanna Wysocka)

through an optimized golden gate assembly reaction. The product of the golden-gate reaction is

then subjected to plasmid-safe exonuclease (Epicentre, Cat#E3101K) treatment. After that, the

exonuclease-treated reaction is then transformed into competent cells (Bioline, Cat#BIO-

85027). For each plate, individual colonies are picked for plasmid purification using a miniprep

kit (QIAGEN, Cat#27204) and test digestion with Acc65I, KpnI, and XbaI. Plasmids with the

correct digestion pattern are then verified by Sanger sequencing

ChIPmentation

We employed ChIPmentation(*51*) to map CTCF in various samples. Briefly, cells and tissues

were fixed in 1% formaldehyde at room temperature for 15 minutes, followed by glycine

quenching. To isolate nuclei, we lysed cell with lysis buffer 1 (50 mM HEPES; pH 7.5, 140 mM

NaCl, 1 mM EDTA, 10% glycerol, 0.5% Nonidet-40, 0.25% Triton X-100, and a protease

inhibitor cocktail) for 10 minutes at 4°C. The nuclei were pelleted at 1,800 g for 10 minutes at

4°C and then resuspended in nuclei lysis buffer (10 mM Tris-Cl pH 8.0, 100 mM NaCl, 1 mM

EDTA, 0.5 mM EGTA, 0.1% Na-Deoxycholate, 0.5% N-lauroylsarcosine, and a protease

inhibitor cocktail) for sonication. For CTCF pulldown, 100-150 μg of chromatin was used.

Dynabeads M-280 (Life Technologies, Sheep Anti-Rabbit IgG, #11204D) were washed three

times with 0.5 mg/mL of BSA/PBS on ice, then incubated with the CTCF antibody (Abcam,

#ab70303) for at least 2 hours at 4°C. The beads/antibody complexes were then washed with

BSA/PBS. Pulldown was performed in binding buffer (1% Triton-X 100, 0.1% Sodium

Deoxycholate, and a protease inhibitor cocktail in 1X TE) by mixing the beads/antibody

complexes with chromatin. After overnight incubation, the beads/antibody/chromatin complexes

were washed with RIPA buffer (50 mM HEPES pH 8.0, 1% NP-40, 0.7% Sodium Deoxycholate,

0.5 M LiCl, 1 mM EDTA, and a protease inhibitor cocktail). The beads complexes were then

incubated with homemade Tn5 transposase in tagmentation reaction buffer (10 mM Tris-Cl pH

8.0 and 5 mM $MgCl_2$) for 10 minutes at 37°C. To remove free DNA, the beads were washed

twice with 1X TE on ice. The pulldown DNA was recovered by reversing crosslinking overnight,

followed by purification with PCR Clean DX beads (Aline Biosciences, #C-1003-50). ChIP-seq

libraries were generated by amplifying the pulldown DNA with Illumina Nextera primers. Size

selection was then performed with PCR Clean DX beads to select fragments ranging from 100

bp to 1000 bp.

Data analysis

Hi-C, micro-C, and 4C-seq

Hi-C and micro-C sequencing reads were mapped to mm10 reference genome separately using

bowtie(_52_) (v.1.1.2) as previously described(_37_). After removing non-unique reads and PCR

duplication reads, the bam files are used as input into _HiCorr_ in '_Bam-process-HindIII_', '_Bam-_

process-DpnII', and '_Bam-process-microC_' modes for bias correction with conventional Hi-C, _in-_

situ Hi-C, and micro-C, respectively. After bias correction, we enhance the Hi-C/micro-C data

with _DeepLoop_. Note that _DeepLoop_ has multiple models to enhance Hi-C data with different

read depth. The enhance models we chose for each dataset are listed in **Supplementary Table**

2. Because each datasets have a pair of Hi-C experiments (control and KO conditions), if the

read depths of the two Hi-C experiments are significantly different, we will down-sample the

high-depth dataset to lower depth to ensure fair comparison (**Supplementary Table 2**). 4C-seq

data were analyzed using pipe4C(*50*) (v.1.1.3) to generate bam and bigwig files for data

visualization on UCSC genome browser.

<u>*ChIP–seq and CUT&RUN*</u>

ChIP-seq and CUT&RUN data were mapped to mm10 (mouse) using bowtie. After mapping

step, we remove PCR duplication reads and use Macs2(*53*) to call peak and generated bigwig

to for data visualization on UCSC genome browser.

<u>*RNA-seq and mNET-seq*</u>

RNA-seq data are aligned to mm10 using HISAT2(*54*) and read counts for each gene are

called with featureCounts(*55*). DESeq2(*56*) is used to identify differential expressed genes

(DEGs). The DEGs are identified using Wald test with the p-value <0.01 and log2FC>1. For

mNET-seq analysis, we followed the customized pipeline from the Hsieh *et al.*(*34*) with the

following steps: (i) Trim the adapter using TrimGalore (v.0.6.7)

(https://github.com/FelixKrueger/TrimGalore) to remove adapter sequences

'AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC' and

'GATCGTCGGACTGTAGAACTCTGAAC' at each side of the reads; (ii) Map trimmed read to

mm10 using STAR; (iii) Identify the last nucleotide incorporated by Pol II by using mNET_snr

(https://github.com/tomasgomes/mNET_snr) to locate the 3′-nucleotide of the second read and

the strand sign of the first read. (iv) Annotate and count the reads of each gene using

FeatureCounts(*55*); (v) Use DEseq2(*56*) to quantify the changes of the mNET-seq signal at the

gene body between control and KO cells.

<u>*Pairwise loop strength comparison*</u>

For each pair of CTCF/cohesin-degron study, we took top 500k loop pixels from the *DeepLoop*

enhanced contact matrices of control and IAA/dTAG-treated samples separately, then union

them together. The loop strength of each pixel in the two experiments are normalized linearly:

$$Y_i \sim k * X_i + \varepsilon$$

$$i: A\ given\ loop\ pixel$$

$$X_i: i\ loop\ strength\ in\ WT\ condition$$

$$Y_i: i\ loop\ strength\ in\ KO\ condition$$

However, because CTCF- and *Rad21*-depletion leads to massive loss of loop signal, much

fewer loop pixels are expected in the IAA/dTAG-treated cells than in control cells. To adjust for

this, we have to manually pick a regression slope *k* based on the shape of the scatter plots.

The slopes we chose are listed in **Supplementary Table 2**. We then use the minimum strength

of the control cell loop pixels to adjust the minimum cutoff to call loop pixels in KO samples:

$Y_{min} = k * \min(X_i)$. In another word, a pixel is called as loop pixel in KO sample only if its

strength is bigger than Y_{min}. This step leads to much fewer loop pixels in CTCF- and *Rad21*-

depleted cells. We typically obtain ~600k chromatin loop for each study. We used a simple fold

change cutoff (FC > 4) to define gained or lost loop pixels. Finally, we used the following criteria

to call retained loops: (i) the strength of the pixel in both WT and KO conditions must be higher

than their minimum cutoffs ($X_i > X_{min}$ & $Y_i > Y_{min}$); (ii) the absolute value of residual must be

higher than the adjusted minimum of loop pixels in KO samples ($|\varepsilon| < Y_{min}$).

Classify chromatin loops based on epigenome data

For the purpose of being consistent, we intersected loop anchors with the same set of mESC

CTCF, H3K27ac, H3K4me3, H3K27me3 and H3K36me3 ChIP-seq peaks from ENCODE(*57*) to

classify the loop pixels. We classification loop pixels in the following order:

(1) EP-only: both anchors have either H3K4me3 or H3K27ac marker, but they do not have

600   CTCF peaks.

601   (2) CTCF-EP: after excluding EP-only, look for pixels which only one anchor has CTCF, and

602   the other anchor has either H3K4me3 or H3K27ac marker.

603   (3) CTCF-only: after excluding EP-only and CTCF-EP, look for pixels with at least one

604   anchor bound by CTCF.

605   (4) H3K27me3: after excluding (1) - (3) above, look for pixels with at least one anchor bound

606   by H3K27me3.

607   (5) H3K36me3: after excluding (1) - (4) above, look for pixels with at least one anchor bound

608   by H3K36me3.

(6) The rest pixels are classified as "other".

Define FINs and FIN-blocked regular enhancers and super enhancers (SE)

We define FINs as a subset of CTCF binding sites that satisfy the following criteria: (i) they

located at the anchors of CTCF loops that are weakened upon CTCF depletion; (ii) the FIN

should be inside a newly gained loop in CTCF-depleted cells; (iii) the other anchor of the CTCF

loop should be outside of the newly gained loop. We call enhancers as H3K27ac peaks and use

a published list of super enhancers in mESCs(*58*) (**Supplementary Tables 11 & 12**). This

allows us to determine the types of enhancers blocked by FINs.

Recurrent loop-gaining pixels and loop-gaining loci in CTCF depleted cells

A recurrent loop-gaining pixel is called if it satisfies FC > 4 in at least two studies. We define the

location of each loop pixel as the entire region between the left and right anchors (including the

anchors themselves). To define loop-gaining loci, we merge all gained loop pixels from the

CTCF-degron studies using the "merge" option in *bedtools*; each loop-gaining locus usually has

multiple loop pixels. Loop loci with gap size < 20kb will be merge into one locus. In this study,

we merged all the recurrent loop-gaining pixels to obtain 299 recurrent loop events. To test the

consistency of loop-gaining loci between independent CTCF-degron study, we call loop-gaining

loci for each individual CTCF-degron study and the share loop-gaining loci from two different

studies are called if they have at least one base overlap.

Test the recurrence of loop-gaining loci from CTCF-degron in Rad21- and Wapl- degron data

Nearly all loop-gaining loci observed in Hi-C degron studies have at least one E-P loop pixels.

We further classify the loop-gaining loci depending on the number of CTCF loops separating the

two anchors of the gained E-P loops. For all loci, we also manually examined all loci to confirm

the classification. Several loci cannot be clearly classified into the categories shown in **Fig. 3d**

because they have different marks or have multiple events. To test if a loop gaining locus is

recurrent in one of the CTCF/cohesin-degron study in **Fig. 3d**, we examine all loop-gaining

pixels in the locus and if one of pixel also gain signal it will be consider recurrent.

Compartment and TADs in mESCs

Hi-C compartment analysis is performed in WT mESCs following previous publication(*59*). After

QC step, we constructed 500kb resolution contact matrices for each chromosome and then

normalized them using the genome distance. We next generated correlation matrices for each

chromosome and performed the principal component analysis on the correlation matrix and then

assigns the genome into two compartments by using PC1. We used the density of transcription

starting sites (TSS) of each 500-kb bin to determine compartment A and compartment B. For

TADs and their boundaries, we directly use the TADs in mESCs defined by Dixon *et al.*

(**Supplementary Table 9**).

Ideogram plot

We generate ideogram plots with Rideogram(*60*) to visualized the locations of recurrent gaining-
loop pixels, compartment, and TADs.

Density and enrichment of DEGs upon CTCF-depletion

For any region in the genome, the density of DEGs is defined as the average number of DEGs
per Mb-region. We computed DEG density for TAD boundary +/- 100kb, FIN +/- 100kb, gained
loop anchors, lost loop anchors, and retained loop anchors. We also compute the enrichment of
DEGs at regions of interest as their DEG density divided by the DEG density in the whole
genome.

Hi-C data visualization

Heatmap was used to visualize the raw, ratio, and *DeepLoop*-enhanced heatmaps. We use the
following strategies as described before(*37*) to determine the color scales these heatmaps. (i)
Raw or ICE-normalized heatmaps represent raw read counts; the brightest red color indicates
the 98th percentile of the contact matrix. Color is proportionally scaled down to one read (white).
(ii) HiCorr heatmap represents the ratio value after Hi-C bias correction; the brightest red color
indicates at least twofold enrichment of a given pixel comparing to its background. Color is
proportionally scaled down to onefold (no enrichment). (iii) DeepLoop heatmaps outputs
'transformed fold change' that represents only relative levels of signal enrichment (for example,
a value of one may no longer represents no enrichment). We therefore set the brightest red
color as the lower limit of the top 300k pixels genome wide. Color is proportionally scaled down
to half of that lower limit or onefold, whichever is higher. All loop arches shown in figures were
generated with UCSC Genome Browser(*61*).

Data availability

Accession numbers for third-party data used in this study can be found in **Supplementary**
Table 1. The raw data of 4C–seq and ChIP-seq generated in this study, and reanalyzed
published data, can be found at accession number GSE243728. The 18 Hi-C and micro-C
datasets analyzed by *DeepLoop* can be found at https://hiview.case.edu/public/DeepLoop/.

To review GEO accession GSE243728:

Go to https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE243728

Enter token uvixkcaajpajpih into the box

Code availability

The code is available is available at github (https://github.com/JinLabBioinfo/FIN_project).

Acknowledgements

This work was supported by grants from NIH R01HG009658 and R01CA267872 (to F.J.),
R01DK113185 (to Y.L.), R01DK131437, R01HG012384, and UG3NS132061 (to Y.L. and F.J.).
F.J. is also supported by a subaward from NIH project (U01AG072579) through University of
Miami and partially by several other NIH grants R01CA252224, R01GM148662, and
P01CA272161. Y.L. is also partially supported by NIH grants R01CA264320. This work made use
of the High-Performance Computing Resource in the Core Facility for Advanced Research
Computing at Case Western Reserve University.

References

1. J. R. Dixon, S. Selvaraj, F. Yue, A. Kim, Y. Li, Y. Shen, M. Hu, J. S. Liu, B. Ren,
Topological domains in mammalian genomes identified by analysis of chromatin
interactions. *Nature* **485**, 376-380 (2012).
2. E. P. Nora, B. R. Lajoie, E. G. Schulz, L. Giorgetti, I. Okamoto, N. Servant, T. Piolot, N.
L. van Berkum, J. Meisig, J. Sedat, J. Gribnau, E. Barillot, N. Bluthgen, J. Dekker, E.
Heard, Spatial partitioning of the regulatory landscape of the X-inactivation centre.
Nature **485**, 381-385 (2012).
3. W. A. Bickmore, B. van Steensel, Genome architecture: domain organization of
interphase chromosomes. *Cell* **152**, 1270-1284 (2013).
4. J. Dekker, L. Mirny, The 3D Genome as Moderator of Chromosomal Communication.
Cell **164**, 1110-1121 (2016).
5. R. P. McCord, N. Kaplan, L. Giorgetti, Chromosome Conformation Capture and Beyond:
Toward an Integrative View of Chromosome Structure and Function. *Mol Cell* **77**, 688-
708 (2020).
6. M. Merkenschlager, E. P. Nora, CTCF and Cohesin in Genome Folding and
Transcriptional Gene Regulation. *Annual review of genomics and human genetics* **17**,
17-43 (2016).

7. Z. Tang, O. J. Luo, X. Li, M. Zheng, J. J. Zhu, P. Szalaj, P. Trzaskoma, A. Magalska, J.
Wlodarczyk, B. Ruszczycki, P. Michalski, E. Piecuch, P. Wang, D. Wang, S. Z. Tian, M.
Penrad-Mobayed, L. M. Sachs, X. Ruan, C. L. Wei, E. T. Liu, G. M. Wilczynski, D.
Plewczynski, G. Li, Y. Ruan, CTCF-Mediated Human 3D Genome Architecture Reveals
Chromatin Topology for Transcription. *Cell* **163**, 1611-1627 (2015).
8. A. L. Sanborn, S. S. Rao, S. C. Huang, N. C. Durand, M. H. Huntley, A. I. Jewett, I. D.
Bochkov, D. Chinnappan, A. Cutkosky, J. Li, K. P. Geeting, A. Gnirke, A. Melnikov, D.
McKenna, E. K. Stamenova, E. S. Lander, E. L. Aiden, Chromatin extrusion explains key
features of loop and domain formation in wild-type and engineered genomes.
Proceedings of the National Academy of Sciences of the United States of America **112**,
E6456-6465 (2015).
9. J. Nuebler, G. Fudenberg, M. Imakaev, N. Abdennur, L. A. Mirny, Chromatin
organization by an interplay of loop extrusion and compartmental segregation.
Proceedings of the National Academy of Sciences of the United States of America **115**,
E6697-E6706 (2018).
10. K. Nasmyth, Disseminating the genome: joining, resolving, and separating sister
chromatids during mitosis and meiosis. *Annu Rev Genet* **35**, 673-745 (2001).
11. G. Fudenberg, M. Imakaev, C. Lu, A. Goloborodko, N. Abdennur, L. A. Mirny, Formation
of Chromosomal Domains by Loop Extrusion. *Cell reports* **15**, 2038-2049 (2016).
12. I. F. Davidson, B. Bauer, D. Goetz, W. Tang, G. Wutz, J. M. Peters, DNA loop extrusion
by human cohesin. *Science* **366**, 1338-1345 (2019).
13. Y. Kim, Z. Shi, H. Zhang, I. J. Finkelstein, H. Yu, Human cohesin compacts DNA by loop
extrusion. *Science* **366**, 1345-1349 (2019).
14. M. J. Rowley, V. G. Corces, Organizational principles of 3D genome architecture. *Nat
Rev Genet* **19**, 789-800 (2018).
15. L. L. P. Hanssen, M. T. Kassouf, A. M. Oudelaar, D. Biggs, C. Preece, D. J. Downes, M.
Gosden, J. A. Sharpe, J. A. Sloane-Stanley, J. R. Hughes, B. Davies, D. R. Higgs,
Tissue-specific CTCF-cohesin-mediated chromatin architecture delimits enhancer
interactions and function in vivo. *Nat Cell Biol* **19**, 952-961 (2017).
16. E. P. Nora, A. Goloborodko, A. L. Valton, J. H. Gibcus, A. Uebersohn, N. Abdennur, J.
Dekker, L. A. Mirny, B. G. Bruneau, Targeted Degradation of CTCF Decouples Local
Insulation of Chromosome Domains from Genomic Compartmentalization. *Cell* **169**, 930-
944 e922 (2017).
17. S. S. P. Rao, S. C. Huang, B. Glenn St Hilaire, J. M. Engreitz, E. M. Perez, K. R. Kieffer-
Kwon, A. L. Sanborn, S. E. Johnstone, G. D. Bascom, I. D. Bochkov, X. Huang, M. S.
Shamim, J. Shin, D. Turner, Z. Ye, A. D. Omer, J. T. Robinson, T. Schlick, B. E.
Bernstein, R. Casellas, E. S. Lander, E. L. Aiden, Cohesin Loss Eliminates All Loop
Domains. *Cell* **171**, 305-320 e324 (2017).
18. K. Nasmyth, C. H. Haering, Cohesin: its roles and mechanisms. *Annu Rev Genet* **43**,
525-558 (2009).
19. K. L. Chan, M. B. Roig, B. Hu, F. Beckouet, J. Metson, K. Nasmyth, Cohesin's DNA exit
gate is distinct from its entrance gate and is regulated by acetylation. *Cell* **150**, 961-974
(2012).
20. P. J. Huis in 't Veld, F. Herzog, R. Ladurner, I. F. Davidson, S. Piric, E. Kreidl, V.
Bhaskara, R. Aebersold, J. M. Peters, Characterization of a DNA exit gate in the human
cohesin ring. *Science* **346**, 968-972 (2014).
21. S. Kueng, B. Hegemann, B. H. Peters, J. J. Lipp, A. Schleiffer, K. Mechtler, J. M. Peters,
Wapl controls the dynamic association of cohesin with chromatin. *Cell* **127**, 955-967
(2006).

22.   W. Schwarzer, N. Abdennur, A. Goloborodko, A. Pekowska, G. Fudenberg, Y. Loe-Mie,
N. A. Fonseca, W. Huber, H. H. C, L. Mirny, F. Spitz, Two independent modes of
chromatin organization revealed by cohesin removal. *Nature* **551**, 51-56 (2017).
23.   J. H. I. Haarhuis, R. H. van der Weide, V. A. Blomen, J. O. Yanez-Cuna, M. Amendola,
M. S. van Ruiten, P. H. L. Krijger, H. Teunissen, R. H. Medema, B. van Steensel, T. R.
Brummelkamp, E. de Wit, B. D. Rowland, The Cohesin Release Factor WAPL Restricts
Chromatin Loop Extension. *Cell* **169**, 693-707 e614 (2017).
24.   G. Wutz, C. Varnai, K. Nagasaka, D. A. Cisneros, R. R. Stocsits, W. Tang, S.
Schoenfelder, G. Jessberger, M. Muhar, M. J. Hossain, N. Walther, B. Koch, M.
Kueblbeck, J. Ellenberg, J. Zuber, P. Fraser, J. M. Peters, Topologically associating
domains and chromatin loops depend on cohesin and are regulated by CTCF, WAPL,
and PDS5 proteins. *The EMBO journal* **36**, 3573-3599 (2017).
25.   J. A. Wallace, G. Felsenfeld, We gather together: insulators and genome organization.
Curr Opin Genet Dev **17**, 400-407 (2007).
26.   K. S. Wendt, K. Yoshida, T. Itoh, M. Bando, B. Koch, E. Schirghuber, S. Tsutsumi, G.
Nagae, K. Ishihara, T. Mishiro, K. Yahata, F. Imamoto, H. Aburatani, M. Nakao, N.
Imamoto, K. Maeshima, K. Shirahige, J. M. Peters, Cohesin mediates transcriptional
insulation by CCCTC-binding factor. *Nature* **451**, 796-801 (2008).
27.   V. Parelho, S. Hadjur, M. Spivakov, M. Leleu, S. Sauer, H. C. Gregson, A. Jarmuz, C.
Canzonetta, Z. Webster, T. Nesterova, B. S. Cobb, K. Yokomori, N. Dillon, L. Aragon, A.
G. Fisher, M. Merkenschlager, Cohesins functionally associate with CTCF on
mammalian chromosome arms. *Cell* **132**, 422-433 (2008).
28.   M. Franke, D. M. Ibrahim, G. Andrey, W. Schwarzer, V. Heinrich, R. Schopflin, K. Kraft,
R. Kempfer, I. Jerkovic, W. L. Chan, M. Spielmann, B. Timmermann, L. Wittler, I. Kurth,
P. Cambiaso, O. Zuffardi, G. Houge, L. Lambie, F. Brancati, A. Pombo, M. Vingron, F.
Spitz, S. Mundlos, Formation of new chromatin domains determines pathogenicity of
genomic duplications. *Nature* **538**, 265-269 (2016).
29.   J. Weischenfeldt, T. Dubash, A. P. Drainas, B. R. Mardin, Y. Chen, A. M. Stutz, S. M.
Waszak, G. Bosco, A. R. Halvorsen, B. Raeder, T. Efthymiopoulos, S. Erkek, C. Siegl,
H. Brenner, O. T. Brustugun, S. M. Dieter, P. A. Northcott, I. Petersen, S. M. Pfister, M.
Schneider, S. K. Solberg, E. Thunissen, W. Weichert, T. Zichner, R. Thomas, M. Peifer,
A. Helland, C. R. Ball, M. Jechlinger, R. Sotillo, H. Glimm, J. O. Korbel, Pan-cancer
analysis of somatic copy-number alterations implicates IRS4 and IGF2 in enhancer
hijacking. *Nat Genet* **49**, 65-74 (2017).
30.   D. G. Lupianez, K. Kraft, V. Heinrich, P. Krawitz, F. Brancati, E. Klopocki, D. Horn, H.
Kayserili, J. M. Opitz, R. Laxova, F. Santos-Simarro, B. Gilbert-Dussardier, L. Wittler, M.
Borschiwer, S. A. Haas, M. Osterwalder, M. Franke, B. Timmermann, J. Hecht, M.
Spielmann, A. Visel, S. Mundlos, Disruptions of topological chromatin domains cause
pathogenic rewiring of gene-enhancer interactions. *Cell* **161**, 1012-1025 (2015).
31.   C. Anania, R. D. Acemel, J. Jedamzick, A. Bolondi, G. Cova, N. Brieske, R. Kuhn, L.
Wittler, F. M. Real, D. G. Lupianez, In vivo dissection of a clustered-CTCF domain
boundary reveals developmental principles of regulatory insulation. *Nat Genet* **54**, 1026-
1036 (2022).
32.   J. Zuin, G. Roth, Y. Zhan, J. Cramard, J. Redolfi, E. Piskadlo, P. Mach, M.
Kryzhanovska, G. Tihanyi, H. Kohler, M. Eder, C. Leemans, B. van Steensel, P. Meister,
S. Smallwood, L. Giorgetti, Nonlinear control of transcription through enhancer-promoter
interactions. *Nature* **604**, 571-577 (2022).
33.   N. Kubo, H. Ishii, X. Xiong, S. Bianco, F. Meitinger, R. Hu, J. D. Hocker, M. Conte, D.
Gorkin, M. Yu, B. Li, J. R. Dixon, M. Hu, M. Nicodemi, H. Zhao, B. Ren, Promoter-
proximal CTCF binding promotes distal enhancer-dependent gene activation. *Nat Struct*
Mol Biol **28**, 152-161 (2021).

34.   T. S. Hsieh, C. Cattoglio, E. Slobodyanyuk, A. S. Hansen, X. Darzacq, R. Tjian,
806   Enhancer-promoter interactions and transcription are largely maintained upon acute loss
807   of CTCF, cohesin, WAPL or YY1. *Nat Genet* **54**, 1919-1932 (2022).
808   35.   B. Akgol Oksuz, L. Yang, S. Abraham, S. V. Venev, N. Krietenstein, K. M. Parsi, H.
809   Ozadam, M. E. Oomen, A. Nand, H. Mao, R. M. J. Genga, R. Maehr, O. J. Rando, L. A.
810   Mirny, J. H. Gibcus, J. Dekker, Systematic evaluation of chromosome conformation
811   capture assays. *Nature methods*, (2021).
812   36.   L. Lu, X. Liu, W. K. Huang, P. Giusti-Rodriguez, J. Cui, S. Zhang, W. Xu, Z. Wen, S. Ma,
813   J. D. Rosen, Z. Xu, C. F. Bartels, R. Kawaguchi, M. Hu, P. C. Scacheri, Z. Rong, Y. Li,
814   P. F. Sullivan, H. Song, G. L. Ming, Y. Li, F. Jin, Robust Hi-C Maps of Enhancer-
815   Promoter Interactions Reveal the Function of Non-coding Genome in Neural
816   Development and Diseases. *Mol Cell* **79**, 521-534 e515 (2020).
817   37.   S. Zhang, D. Plummer, L. Lu, J. Cui, W. Xu, M. Wang, X. Liu, N. Prabhakar, J. Shrinet,
818   D. Srinivasan, P. Fraser, Y. Li, J. Li, F. Jin, DeepLoop robustly maps chromatin
819   interactions from sparse allele-resolved or single-cell Hi-C data at kilobase resolution.
820   *Nat Genet* **54**, 1013-1025 (2022).
821   38.   A. J. Kriz, D. Colognori, H. Sunwoo, B. Nabet, J. T. Lee, Balancing cohesin eviction and
822   retention prevents aberrant chromosomal interactions, Polycomb-mediated repression,
823   and X-inactivation. *Mol Cell* **81**, 1970-1987 e1979 (2021).
824   39.   J. D. P. Rhodes, A. Feldmann, B. Hernandez-Rodriguez, N. Diaz, J. M. Brown, N. A.
825   Fursova, N. P. Blackledge, P. Prathapan, P. Dobrinic, M. K. Huseyin, A. Szczurek, K.
826   Kruse, K. A. Nasmyth, V. J. Buckle, J. M. Vaquerizas, R. J. Klose, Cohesin Disrupts
827   Polycomb-Dependent Chromosome Interactions in Embryonic Stem Cells. *Cell reports*
828   **30**, 820-835 e810 (2020).
829   40.   N. Q. Liu, M. Maresca, T. van den Brand, L. Braccioli, M. Schijns, H. Teunissen, B. G.
830   Bruneau, E. P. Nora, E. de Wit, WAPL maintains a cohesin loading cycle to preserve
831   cell-type-specific distal gene regulation. *Nat Genet* **53**, 100-109 (2021).
832   41.   T. Hirayama, T. Yagi, Regulation of clustered protocadherin genes in individual neurons.
833   *Seminars in cell & developmental biology* **69**, 122-130 (2017).
834   42.   E. Flaherty, T. Maniatis, The role of clustered protocadherins in neurodevelopment and
835   neuropsychiatric diseases. *Curr Opin Genet Dev* **65**, 144-150 (2020).
836   43.   Z. Jia, J. Li, X. Ge, Y. Wu, Y. Guo, Q. Wu, Tandem CTCF sites function as insulators to
837   balance spatial chromatin contacts and topological enhancer-promoter selection.
838   *Genome Biol* **21**, 75 (2020).
839   44.   T. Hirayama, E. Tarusawa, Y. Yoshimura, N. Galjart, T. Yagi, CTCF is required for
840   neural development and stochastic expression of clustered Pcdh genes in neurons. *Cell*
841   *reports* **2**, 345-357 (2012).
842   45.   B. Gu, T. Swigut, A. Spencley, M. R. Bauer, M. Chung, T. Meyer, J. Wysocka,
843   Transcription-coupled changes in nuclear mobility of mammalian cis-regulatory
844   elements. *Science* **359**, 1050-1055 (2018).
845   46.   D. R. Tarjan, W. A. Flavahan, B. E. Bernstein, Epigenome editing strategies for the
846   functional annotation of CTCF insulators. *Nature communications* **10**, 4258 (2019).
847   47.   T. Pollex, A. Rabinowitz, M. C. Gambetta, R. Marco-Ferreres, R. R. Viales, A.
848   Jankowski, C. Schaub, E. E. M. Furlong, Enhancer-promoter interactions become more
849   instructive in the transition from cell-fate specification to tissue differentiation. *Nat Genet*
850   **56**, 686-696 (2024).
851   48.   C. P. Fulco, J. Nasser, T. R. Jones, G. Munson, D. T. Bergman, V. Subramanian, S. R.
852   Grossman, R. Anyoha, B. R. Doughty, T. A. Patwardhan, T. H. Nguyen, M. Kane, E. M.
853   Perez, N. C. Durand, C. A. Lareau, E. K. Stamenova, E. L. Aiden, E. S. Lander, J. M.
854   Engreitz, Activity-by-contact model of enhancer-promoter regulation from thousands of
855   CRISPR perturbations. *Nat Genet* **51**, 1664-1669 (2019).

49. M. H. Kagey, J. J. Newman, S. Bilodeau, Y. Zhan, D. A. Orlando, N. L. van Berkum, C.
C. Ebmeier, J. Goossens, P. B. Rahl, S. S. Levine, D. J. Taatjes, J. Dekker, R. A.
Young, Mediator and cohesin connect gene expression and chromatin architecture.
Nature **467**, 430-435 (2010).
50. P. H. L. Krijger, G. Geeven, V. Bianchi, C. R. E. Hilvering, W. de Laat, 4C-seq from
beginning to end: A detailed protocol for sample preparation and data analysis. *Methods*
170, 17-32 (2020).
51. C. Schmidl, A. F. Rendeiro, N. C. Sheffield, C. Bock, ChIPmentation: fast, robust, low-
input ChIP-seq for histones and transcription factors. *Nature methods* **12**, 963-965
(2015).
52. B. Langmead, C. Trapnell, M. Pop, S. L. Salzberg, Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).
53. Y. Zhang, T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson, B. E. Bernstein, C.
Nusbaum, R. M. Myers, M. Brown, W. Li, X. S. Liu, Model-based analysis of ChIP-Seq
(MACS). *Genome Biol* **9**, R137 (2008).
54. D. Kim, B. Langmead, S. L. Salzberg, HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods* **12**, 357-360 (2015).
55. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
56. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
57. F. Yue, Y. Cheng, A. Breschi, J. Vierstra, W. Wu, T. Ryba, R. Sandstrom, Z. Ma, C.
Davis, B. D. Pope, Y. Shen, D. D. Pervouchine, S. Djebali, R. E. Thurman, R. Kaul, E.
Rynes, A. Kirilusha, G. K. Marinov, B. A. Williams, D. Trout, H. Amrhein, K. Fisher-Aylor,
I. Antoshechkin, G. DeSalvo, L. H. See, M. Fastuca, J. Drenkow, C. Zaleski, A. Dobin, P.
Prieto, J. Lagarde, G. Bussotti, A. Tanzer, O. Denas, K. Li, M. A. Bender, M. Zhang, R.
Byron, M. T. Groudine, D. McCleary, L. Pham, Z. Ye, S. Kuan, L. Edsall, Y. C. Wu, M. D.
Rasmussen, M. S. Bansal, M. Kellis, C. A. Keller, C. S. Morrissey, T. Mishra, D. Jain, N.
Dogan, R. S. Harris, P. Cayting, T. Kawli, A. P. Boyle, G. Euskirchen, A. Kundaje, S. Lin,
Y. Lin, C. Jansen, V. S. Malladi, M. S. Cline, D. T. Erickson, V. M. Kirkup, K. Learned, C.
A. Sloan, K. R. Rosenbloom, B. Lacerda de Sousa, K. Beal, M. Pignatelli, P. Flicek, J.
Lian, T. Kahveci, D. Lee, W. J. Kent, M. Ramalho Santos, J. Herrero, C. Notredame, A.
Johnson, S. Vong, K. Lee, D. Bates, F. Neri, M. Diegel, T. Canfield, P. J. Sabo, M. S.
Wilken, T. A. Reh, E. Giste, A. Shafer, T. Kutyavin, E. Haugen, D. Dunn, A. P. Reynolds,
S. Neph, R. Humbert, R. S. Hansen, M. De Bruijn, L. Selleri, A. Rudensky, S.
Josefowicz, R. Samstein, E. E. Eichler, S. H. Orkin, D. Levasseur, T. Papayannopoulou,
K. H. Chang, A. Skoultchi, S. Gosh, C. Disteche, P. Treuting, Y. Wang, M. J. Weiss, G.
A. Blobel, X. Cao, S. Zhong, T. Wang, P. J. Good, R. F. Lowdon, L. B. Adams, X. Q.
Zhou, M. J. Pazin, E. A. Feingold, B. Wold, J. Taylor, A. Mortazavi, S. M. Weissman, J.
A. Stamatoyannopoulos, M. P. Snyder, R. Guigo, T. R. Gingeras, D. M. Gilbert, R. C.
Hardison, M. A. Beer, B. Ren, E. C. Mouse, A comparative encyclopedia of DNA
elements in the mouse genome. *Nature* **515**, 355-364 (2014).
58. D. Hnisz, B. J. Abraham, T. I. Lee, A. Lau, V. Saint-Andre, A. A. Sigova, H. A. Hoke, R.
A. Young, Super-enhancers in the control of cell identity and disease. *Cell* **155**, 934-947
(2013).
59. E. Lieberman-Aiden, N. L. van Berkum, L. Williams, M. Imakaev, T. Ragoczy, A. Telling,
I. Amit, B. R. Lajoie, P. J. Sabo, M. O. Dorschner, R. Sandstrom, B. Bernstein, M. A.
Bender, M. Groudine, A. Gnirke, J. Stamatoyannopoulos, L. A. Mirny, E. S. Lander, J.
Dekker, Comprehensive mapping of long-range interactions reveals folding principles of
the human genome. *Science* **326**, 289-293 (2009).

60.   Z. Hao, D. Lv, Y. Ge, J. Shi, D. Weijers, G. Yu, J. Chen, RIdeogram: drawing SVG
907   graphics to visualize and map genome-wide data on the idiograms. *PeerJ Comput Sci* **6**,
908   e251 (2020).
909   61.   J. Navarro Gonzalez, A. S. Zweig, M. L. Speir, D. Schmelter, K. R. Rosenbloom, B. J.
910   Raney, C. C. Powell, L. R. Nassar, N. D. Maulding, C. M. Lee, B. T. Lee, A. S. Hinrichs,
911   A. C. Fyfe, J. D. Fernandes, M. Diekhans, H. Clawson, J. Casper, A. Benet-Pages, G. P.
912   Barber, D. Haussler, R. M. Kuhn, M. Haeussler, W. J. Kent, The UCSC Genome
913   Browser database: 2021 update. *Nucleic Acids Res* **49**, D1046-D1057 (2021).
914

Figure 1

a

Published Hi-C data in mESCs upon acute depletion of architectural proteins

Publication	mESC line	KO protocol	C-technique	# of contacts	
				Ctrl	IAA/dTAG
Nora et al.	E14TG2a	C-Ctcf AID (48hrs)	HindIII Hi-C	244M	242M
Kubo et al.	F123	C-Ctcf AID (48hrs)	in-situ Hi-C	185M	116M
Hsieh et al.	JM8.N4	N-Ctcf AID (3hrs)	micro-C	291M	830M
Rhodes et al.	E14TG2a	C-Rad21 AID (6hrs)	in-situ Hi-C	63M	43M
Kriz et al.	F123	C-Rad21 dTAG (8hrs)	In-situ Hi-C	136M	100M
Hsieh et al.	JM8.N4	C-Rad21 AID (3hrs)	micro-C	283M	1.00B
Liu et al.	E14TG2a	C-Wapl AID (24hrs)	in-situ Hi-C	77M	88M
Kriz et al.	F123	C-Wapl dTAG (72hrs)	in-situ Hi-C	193M	147M
Hsieh et al.	JM8.N4	N-Wapl AID (3hrs)	micro-C	298M	670M

b

ICE normalization | HiCorr | DeepLoop

c

chr2:26,176,653-27,145,048

d

CTCF-degron; Nora et al.

e

f

CTCF-degron

Figure 2

Figure 3

a

Non-recurrent N=3,197   Recurrent N=4,944

Nora
Kubo
Hsieh

log2(DCTCF/Ctrl)
<1
[1,2)
[2,3)
>=3

N=1,969, 170kb
N=493, 455kb
N=735, 305kb
N=354, 235kb
N=2,621, 200kb
N=803, 435kb
N=1,166, 340kb

b

Nora N=332   Kubo N=271
42   48
68   174   35
   14
37
Hsieh N=293

c

Gained loops
279
4,665
Insulated by CTCF loops?
Yes   No

d

Classification and recurrence of loop gaining loci between independent CTCF/cohesin degron studies

Type of loci	Classification of loci	#of loci	Ctcf degron			Rad21 degron			Wapl degron		
			Nora	Kubo	Hsieh	Rhodes	Kriz	Hsieh	Liu	Kriz	Hsieh
EP N=283	I. — CTCF	107	105	83	87	1	0	1	21	26	26
	II. — CTCF	156	150	122	135	0	0	0	47	51	48
	III. - CTCF	20	20	14	17	0	0	0	0	0	1
CTCF N=2	- CTCF	2	2	2	2	0	0	0	2	2	1
Undef N=14	Undetermined	14	12	10	13	1	1	2	11	8	10

⬡ CTCF   ▶◀ CTCF motif   ⬭ cohesin   ▬ Enhancer or Promoter

e

i. Define enhancer-blocking insulator by its transcription consequences

Enhancer   Insulator   Promoter   OFF
Enhancer   Insulator✗   Promoter   ON

ii. Define insulator as TAD boundary

CTCF Insulator   Enhancer   CTCF Insulator   Promoter   CTCF Insulator

iii. Define insulator by loop dynamics

Anchor   Enhancer   CTCF Insulator   Promoter   OFF
Anchor   Enhancer   Insulator   Promoter   ON

Functional insulator (FIN)

f

Chr1

FIN loci-Nora
FIN loci-Kubo
FIN loci-Hsieh
TAD boundary
Compartment

Log2(FC) Ctcf-AID/Ctrl
0 — 2
0 — 2
0 — 2
A B

g

FIN & TAD boundary
13%
87%
non-TAD boundary
on-TAD boundary

h

FIN & compartment
percentage
FIN N=785   TAD-bdry N=5,556
Compartment A B

i

Wapl-KO and Ctcf-KO may both cause FIN-skipping

- Ctcf   - Wapl

Figure 4

Figure 5

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

■ CTCF-loop ■ CTCF_EP ■ EP_only ■ H3K27me3 ■ H3K36me3 ■ Other

Supplementary Figure 4

a

Wapl-degron; Kriz *et al.*

dTAG *vs.* Control — R²=0.842

dTAG *vs.* Control — R²=0.842

Loop size

Loop Size (kb)

Lost 241kb   Retain 343kb   Gain 545kb

b

Lost loop N=1,903 — 54%, 30%, 1%, 11%, 4%

Retain loop N=310,005 — 56%, 27%, 2%, 11%, 4%

Gain loop N=32,814 — 63%, 15%, 2%, 12%, 8%

c

Wapl-degron

Kriz *et al.*   Liu *et al.*   Hsieh *et al.*

Lost loop   Retain loop   Gain loop

d

Wapl-degron; Hsieh *et al.*

IAA *vs.* Control — R²=0.854

IAA *vs.* Control — R²=0.854

Loop size

Loop Size (kb)

Lost 170kb   Retain 185kb   Gain 270kb

e

Lost loop N=1,837 — 62%, 10%, 1%, 1%, 13%, 13%

Retain loop N=373,532 — 53%, 25%, 3%, 13%, 6%

Gain loop N=90,311 — 65%, 13%, 2%, 12%, 8%

CTCF-loop   CTCF_EP   EP_only   H3K27me3   H3K36me3   Other

f

Wapl-degron

Hsieh *et al.*   Liu *et al.*   Kriz *et al.*

Lost loop   Retain loop   Gain loop

Supplementary Figure 5