Title: The influence of nucleus accumbens shell D1 and D2 neurons
on outcome-specific Pavlovian instrumental transfer

Octavia Soegyono, Elise Pepin, Beatrice K. Leung, Billy C. Chieng, Bernard W.
Balleine & Vincent Laurent

Decision Neuroscience Laboratory, School of Psychology, UNSW Sydney, NSW
2052, Australia

Corresponding author: Vincent Laurent; Email: v.laurent@unsw.edu.au

Abstract

The nucleus accumbens shell (NAc-S) and its projections to the ventral pallidum (VP) are thought to be critical for stimulus-based decisions. The NAc-S is predominantly composed of spiny projection neurons (SPNs) that express either the dopamine D1 (D1-SPNs) or the dopamine D2 receptor (D2-SPNs). Yet, the role of these two neuronal subpopulations and their inputs to the VP in stimulus-based decisions remains unknown. Here, we used optogenetics in female and male knock-in rats to selectively silence D1- or D2-SPNs and their projections to the VP at a time when the rats were required to use predictive stimuli to choose between two instrumental actions. Silencing either population of NAc-S SPNs disrupted choice. Silencing NAc-S D1-SPNs terminals in the VP also disrupted choice. However, choice was left intact by silencing NAc-S D2-SPNs terminals in the VP. Together, these findings provide novel insights into the cellular mechanisms and circuitry underlying stimulus-based decisions. We discuss how these insights are consistent with a recent model proposing that these decisions are controlled by an opioid-based memory system residing in the NAc-S.

Introduction

Our choices and decisions are often influenced by predictive signals available in the environment (Hollis, 1984). This influence is studied in the laboratory using the outcome-specific Pavlovian-instrumental transfer (PIT) task (Trapold & Overmier, 1972; Colwill & Rescorla, 1988; Holmes et al., 2010; Cartoni et al., 2016; Laurent & Balleine, 2021; Leung et al., 2024a), which comprises three stages. The first two stages are Pavlovian and instrumental conditioning, which can be administered in any order. In Pavlovian conditioning, two stimuli (S1 and S2; e.g., sounds or lights) are paired with two distinct and motivationally significant outcomes (O1 and O2; e.g., food outcomes). In instrumental conditioning, subjects are trained to perform two actions (A1 and A2), with each earning one of the two outcomes (i.e., A1→O1 and A2→O2). The final stage is the PIT test, which evaluates choice between the two actions in the presence (and absence) of the stimuli. Typically, each of the Pavlovian stimuli bias choice towards the action with which it shares an outcome: i.e., S1: A1>A2 and S2: A2>A1.

At a neural level, PIT has been found to rely on interactions between a number of structures, including cortical (Keistler et al., 2015; Bradfield et al., 2015; Bradfield et al., 2018; Tensaouti et al., 2025; Ostlund & Balleine, 2007; Lichtenberg et al., 2021; Sias et al., 2021; Lichtenberg et al., 2017), amygdala (Corbit & Balleine, 2005; Morse et al., 2020; Sias et al., 2024; Lichtenberg et al., 2017; Shiflett & Balleine, 2010; Derman et al., 2020; Prévost et al., 2012), basal ganglia (Corbit & Balleine, 2011; Corbit et al., 2001; Corbit & Janak, 2010; Morse et al., 2020; Laurent et al., 2012; Bertran-Gonzalez et al., 2013; Laurent et al., 2014; Corbit et al., 2016; Leung et al., 2024b; Leung & Balleine, 2015; Leung & Balleine, 2013; Laurent et al., 2015; Prévost et al., 2012), midbrain (Corbit et al., 2007; Sias et al., 2024; Seitz et al., 2022) and thalamic (Ostlund & Balleine, 2008; Leung et al., 2024b) territories. Among these, the nucleus accumbens shell (NAc-S) stands out because, unlike the others, it does not itself contribute to learning the stimulus-outcome (S-O) or the action-outcome (A-O) associations formed during the Pavlovian and instrumental stages of the PIT task and appears only to mediate their interaction (Corbit & Balleine, 2011; Corbit et al., 2001; Morse et al., 2020; Laurent et al., 2012; Laurent et al., 2014). Therefore, it has been proposed that the NAc-S acts as a central hub integrating S-O and A-O information at the time of PIT to guide choice between

actions in an outcome-selective manner (Shiflett & Balleine, 2010; Bertran-Gonzalez
& Laurent, 2018; Laurent & Balleine, 2021). Accordingly, PIT is disrupted by
manipulations undermining NAc-S function during the PIT test (Morse et al., 2020;
Corbit et al., 2016; Laurent et al., 2012; Laurent et al., 2014; Laurent et al., 2015).

The NAc-S is predominantly composed of spiny projection neurons (SPNs), which
can be classified into two distinct subpopulations depending on the dopamine
receptor they express (Gerfen & Surmeier, 2011). One population harbors the
dopamine D1 receptors (D1Rs; D1-SPNs) whereas the other population express the
dopamine D2 receptors (D2Rs; D2-SPNs). To date, only one study has examined
whether activity in both populations is recruited during PIT (Laurent et al., 2014),
finding that PIT is associated with an increase in ERK1/2 phosphorylation in D1-
SPNs but not D2-SPNs. Further, blockade of D1Rs in NAc-S during choice between
actions disrupted PIT whereas D2Rs blockade had no effect. Although these findings
indicate a major involvement of NAc-S D1-SPNs in PIT, caution must be exerted
when drawing conclusions on the role of D2-SPNs. While D1Rs are exclusively
expressed on NAc-S D1-SPNs, D2Rs are present on cholinergic interneurons (CINs)
and local presynaptic dopamine terminals in addition to D2-SPNs (Gerfen &
Surmeier, 2011). Additionally, research shows that ERK1/2 phosphorylation may not
capture all transcriptional events that occur in D2-SPNs (Bertran-Gonzalez et al.,
2008; Matamales et al., 2020). Lastly, pharmacological D2R blockade has been
found to be quite ineffective at influencing D2-SPNs function (Tozzi et al., 2007).
Thus, the relative contribution of NAc-S D1- and D2-SPNs in driving choice between
actions during PIT remains unclear.

The present experiments therefore aimed to provide an unambiguous assessment of
the roles played by the two main populations of NAc-S SPNs during choice between
actions in an outcome-specific PIT task. This assessment was achieved through
optogenetic silencing in two knock-in rat lines that express Cre recombinase in either
D1- or D2-SPNs (Pettibone et al., 2019). We first used tract-tracing and ex-vivo
electrophysiology to confirm our capacity to selectively silence activity in NAc-S D1-
or D2-SPNs. We then conducted two experiments that assessed outcome-specific
task while D1- or D2-SPNs in the NAc-S were silenced during presentations of the
predictive stimuli in the choice test. Finally, we assessed the degree to which the
influence of the two SPNs populations on outcome-specific PIT depends on their

projections to the ventral pallidum (VP). We focused on the VP for two main reasons:
First, both NAc-S D1- and D2-SPNs send dense projections to this region (Lu et al.,
1997; Kupchik et al., 2015); and, second, communication between the NAc-S and
VP has previously been found to be critical for outcome-specific PIT (Leung &
Balleine, 2013).

Results

Anterograde tracing and ex-vivo cell recordings in D1-Cre and A2a-Cre rats

We first examined the connectivity of striatal D1- and D2-SPNs in transgenic rats
expressing Cre recombinase in neurons harboring the dopamine D1 receptor (D1-
Cre rats) or the adenosine A2a receptor (A2a-Cre rats) (Pettibone et al., 2019). A2a
receptors are predominantly expressed on striatal D2-SPNs (Schiffmann et al.,
2007), thereby enabling selective targeting of these SPNs in the A2a-Cre line. D1-
Cre (n = 1 female and 1 male) and A2a-Cre rats (n = 1 female and 1 male) were
unilaterally infused with a Cre-dependent eYFP virus in the NAc-S (Figure 1 A). To
validate expression in these rat lines, other D1-Cre (n = 4 males) and A2a-Cre rats (n
= 3 males) were given this unilaterally infusion in the dorsal striatum (DS) with the
same virus. In both transgenic strains, viral expression was largely restricted to local
SPNs in the two targeted regions: ~95% of eYFP-positive neurons expressed
DARPP-32 (D32 | eYFP; Figures 1 C-F). Viral expression was consistent with D1-
and D2-SPNs representing two neuronal populations similar in size: in both
transgenic strains and targeted regions ~40% of DARPP-32-postive neurons
expressed eYFP (eYFP | D32; Figures 1 C-F). DS infusion in D1-Cre rats resulted in
eYFP-positive terminals (Figures 1 B-C) in the substantia nigra pars reticulata (SNr)
and the globus pallidus externus (GPe). By contrast, the same infusion in A2a-Cre
rats only produced eYFP-positive terminals in the GPe (Figures 1 B and D). NAc-S
infusion in both D1-Cre and A2a-Cre rats resulted in eYFP-positive terminals in the
VP (Figures 1 E-F). By contrast, eYFP-positive terminals in the lateral hypothalamus
(Figures 1 E-F) were only observed in D1-Cre rats. Overall, these results are in line
with current knowledge about the cellular composition of the striatum (Gerfen &
Surmeier, 2011; Bertran-Gonzalez et al., 2010) and well-established projection
patterns of striatal D1- and D2-SPNs (Gerfen & Surmeier, 2011; Lu et al., 1997;
Kupchik et al., 2015; O'Connor et al., 2015). We were therefore confident in the

capacity of the two rat transgenic lines to reveal the function of NAc-S D1- and D2-SPNs.

Next, we used ex-vivo electrophysiological recordings to assess our capacity to silence activity of NAc-S D1- and D2-SPNs. D1-Cre and A2a-Cre rats were bilaterally infused in the NAc-S with either a null Cre-dependent eYFP virus (D1-Cre: 2 females; A2a-Cre: 2 females) or a Cre-dependent inhibitory halorhodospin (eNpHR3.0; D1-Cre: 2 females; A2a-Cre: 2 females) virus (Figures 2 A-B). LED light illumination (625 nm) of either NAc-S D1-SPNs (Figure 2 A) or NAc-S D2-SPNs (Figure 2 B) transfected with the eYFP virus had no effect on action potential firing. By contrast, the same illumination onto NAc-S D1- (Figure 2 A) or D2-SPNs (Figure 2 B) transfected with eNpHR3.0 suppressed action potential firing. Importantly, there was no difference in firing prior to or following LED light illumination, indicating that the optical manipulation did not alter D1- or D2-SPNs physiology. Together, these results validate the efficacy of eNpHR3.0 to specifically silence D1- and D2-SPNs activity at the time of LED light delivery.

NAc-S D1-SPNs mediate outcome-specific PIT

This experiment examined the effects of silencing NAc-S D1-SPNs on outcome-specific PIT. D1-Cre rats were bilaterally infused in the NAc-S with either the null Cre-dependent eYFP virus (eYFP: 5 females and 5 males) or the Cre-dependent halorhodopsin (eNpHR3.0: 5 females and 5 males) virus and were implanted with fiber-optic cannulas above the NAc-S (Figure 3 A and Supplementary Figures S1 A-B). Using the outcome-specific PIT protocol depicted in Figure 3 B, the rats first received Pavlovian conditioning during which two stimuli (S1 and S2; white noise or clicker) were paired with two food outcomes (O1 and O2; grain pellets or sucrose solution). Next, the rats underwent instrumental conditioning and learned that pressing one lever earned one of the food outcomes whereas pressing a second lever delivered the other outcome. Finally, an outcome-specific PIT test was administered and assessed choice between the two lever press actions in the presence or absence of each stimulus. Critically, half of the trials for each stimulus took place under LED light stimulation (625 nm, ON trials) whereas the other half of the trials was conducted without the stimulation (OFF trials). Thus, NAc-S D1-SPNs were silenced on half of the trials in rats infused with eNpHR3.0.

Pavlovian and instrumental conditioning proceeded as expected (Supplementary
Figures S1 C-D). The data of most interest are those from the outcome-specific PIT
test and are presented in Figure 3 C (see also Supplementary Figure S1 E). They
are plotted as the mean number of lever presses per minute when the stimulus
predicted the same outcome as the action (Same) and when the stimulus predicted a
different outcome to the action (Different). Thus, A1 was identified as "Same" and A2
as "Different" in the presence of S1. Conversely, A2 was identified as "Same" and A1
as "Different" in the presence of S2. Further, baseline responding (lever presses per
minute during the two minutes preceding each stimulus presentation) was subtracted
from the rates of responding during the stimuli since it did not differ between groups
(Group – eYFP vs. eNpHR3.0: $p = 0.42$). This approach allowed us to focus on the
net effect of the stimuli on choice performance.

Silencing NAc-S D1 SPNs eliminated outcome-specific PIT. Overall lever press rates
were similar between groups (Group: $p = 0.48$), LED light condition (LED – ON vs
OFF: $p = 0.14$), and the two factors did not interact (Group x Light: $p = 0.07$).
However, lever press rates were higher on the action earning the same outcome as
the stimuli compared to the action earning the different outcome (Lever – Same vs.
Different: $F_{(1,18)} = 50.06$; $\eta^2 = 0.74$; $p < 0.001$), regardless of group (Group x Lever: p
$= 0.77$). There was no Lever by LED light condition interaction (Lever x LED: $p =$
0.72) but critically, there was an interaction between Group, LED light condition, and
Lever during the presentation of the predictive stimuli (Group x LED x Lever: $F_{(1,18)} =$
7.19; $\eta^2 = 0.29$; $p = 0.02$). Follow-up analyses revealed that rats in the eYFP group
displayed intact outcome-specific PIT whether the LED light was OFF ($F_{(1,9)} = 6.14$;
$\eta^2 = 0.67$; $p < 0.05$) or ON ($F_{(1,9)} = 39.46$; $\eta^2 = 0.81$; $p < 0.001$). By contrast, rats in
the eNpHR3.0 group displayed outcome-specific PIT when the light was OFF ($F_{(1,9)} =$
12.38; $\eta^2 = 0.58$; $p < 0.01$) but not when it was ON ($p = 0.67$). Together, these results
indicate that activity in NAc-S D1-SPNs is necessary for the outcome-specific
influence exerted by predictive stimuli on choice between actions. This is consistent
with previous pharmacological work showing that NAc-S D1Rs blockade removes
specific PIT (Laurent et al., 2014). Importantly, the impairment produced by silencing
NAc-S D1-SPNs was restricted to the influence of predictive stimuli on choice
between actions. In the same rats, we found that this silencing had no effect on
valued-based choice (Figure S1 F) using an outcome devaluation procedure: all rats

performed the action that previously earned a valuable outcome more than the
action that earned a devalued outcome. This agrees with studies reporting that NAc-
S lesion or inactivation spares value-based choice (Corbit & Balleine, 2011; Corbit et
al., 2001; Morse et al., 2020; Laurent et al., 2012; Laurent et al., 2014).

NAc-S D2-SPNs mediate outcome-specific PIT

The removal of outcome-specific PIT under silencing of NAc-S D1-SPNs reproduced
the impairment previously observed with blockade of NAc-S D1Rs (Laurent et al.,
2014). Interestingly, the latter pharmacological study did not find any effect following
NAc-S infusion of a D2R antagonist. To evaluate this finding, the present experiment
examined the effects of silencing NAc-S D2-SPNs on outcome-specific PIT. A2a-Cre
rats were bilaterally infused in the NAc-S with either the null Cre-dependent eYFP
virus (eYFP: 4 females and 4 males) or the Cre-dependent halorhodospin
(eNpHR3.0: 4 females and 2 males) virus and were implanted with fiber-optic
cannulas above the NAc-S (Figure 4 A and Supplementary Figures S2 A-B). The rats
then received the outcome-specific PIT protocol previously described (Figure 3 B).

Pavlovian and instrumental conditioning went as expected (Supplementary Figures
S2 C-D). The data from the outcome-specific PIT test are presented in Figure 4B,
again with baseline subtracted since it did not differ between groups (Group: $p =$
0.89; see also Supplemental Figure 2 E). Silencing NAc-S D2-SPNs activity
eliminated outcome-specific PIT. Overall lever press rates during the test were
similar between groups (Group: $p = 0.86$), LED light condition (LED: $p = 0.21$), and
the two factors did not interact (Group x LED: $p = 0.70$). Lever press rates were
higher on the action earning the same outcome as the stimuli compared to the action
earning the different outcome (Lever: $F_{(1,12)} = 22.77$; $\eta^2 = 0.66$; $p < 0.001$), regardless
of group (Group x Lever: $p = 0.46$). There was no Lever by LED light condition
interaction (Lever x LED: $p = 0.15$) but critically, there was an interaction between
group, LED light condition, and Lever during the presentation of the predictive stimuli
(Group x LED x Lever: $F_{(1,12)} = 8.73$; $\eta^2 = 0.42$; $p < 0.01$). Follow-up analyses
revealed that control eYFP rats expressed outcome-specific PIT whether the LED
light was OFF ($F_{(1,7)} = 16.25$; $\eta^2 = 0.70$; $p < 0.01$) or ON ($F_{(1,7)} = 20.58$; $\eta^2 = 0.75$; $p <$
0.01). By contrast, rats in the eNpHR3.0 group displayed outcome-specific PIT when
the light was OFF ($F_{(1,5)} = 6.94$; $\eta^2 = 0.58$; $p < 0.05$) but not ON ($p = 0.34$). Thus,

these results show for the first time that activity in NAc-S D2-SPNs is necessary for
the outcome-specific influence exerted by predictive stimuli on choice between
actions.

The present finding contrasts with our previous observation that outcome-specific
PIT remains unaffected by NAc-S infusion of a D2R antagonist (Laurent et al., 2014).
However, the capacity of such pharmacological manipulation to reveal the function of
D2-SPNs is unclear. For instance, a D2R antagonist could be expected to enhance
D2-SPN activity, since D2Rs are Gi coupled and so their activation should reduce
adenylyl cyclase activity. Further, D2R are not exclusively expressed on this cell
population (Gerfen & Surmeier, 2011), as they can be found on local CINs and
presynaptic dopamine terminals. Finally, previous work found that pharmacological
treatment targeting D2R can leave D2-SPN activity unaffected (Tozzi et al., 2007).
These issues are overcome by the present optical manipulation in A2a-Cre rats, and
we are therefore confident that NAc-S D2-SPNs activity is in fact critical for outcome-
specific PIT. Importantly, we also found that the impairment produced by silencing
NAc-S D2-SPNs was restricted to the influence of predictive stimuli on choice
between actions. In the same rats, this silencing had no effect on valued-based
choice (Figure S2 F).

NAc-S D1-SPNs projections to the VP mediate outcome-specific PIT

Previous work indicates that communication between the NAc-S and VP is critical for
outcome-specific PIT (Leung & Balleine, 2013). Consistent with the literature (Lu et
al., 1997; Kupchik et al., 2015), we observed that both NAc-S D1- and D2-SPNs
densely innervate the VP (Figure 1 E-F). Since we found that activity in both neuron
populations mediates the outcome-specific PIT effect, we investigated whether
projections from each population contribute to the effect. We first focused on those
originating from NAc-S D1-SPNs. D1-Cre rats were bilaterally infused in the NAc-S
with either the null Cre-dependent eYFP virus (eYFP: 3 females and 5 males) or the
Cre-dependent halorhodopsin (eNpHR3.0: 6 females and 6 males) virus and were
implanted with fiber-optic cannulas above the VP (Figure 5 A and Supplementary
Figures S3 A-B). The rats then received the behavioral protocol previously described
(Figure 3 B).

Pavlovian and instrumental conditioning went as expected (Supplementary Figures
S3 C-D). The data from the outcome-specific PIT test are presented in Figure 5 B in
the manner described previously since baseline responding did not differ between
groups (Group: $p = 0.34$; see also Supplemental Figure 3 E). Silencing NAc-S D1-
SPNs projections to the VP eliminated outcome-specific PIT. Overall lever press
rates during the test were similar between groups (Group: $p = 0.11$). LED activation
reduced these rates (LED: $F_{(1,18)} = 7.58$; $\eta^2 = 0.30$; $p < 0.05$) but this reduction
depended on the group considered (Group x LED: $F_{(1,18)} = 9.78$; $\eta^2 = 0.35$; $p < 0.01$).
Lever press rates were higher on the action earning the same outcome as the stimuli
compared to the action earning the different outcome (Lever: $F_{(1,18)} = 35.73$; $\eta^2 =$
0.67; $p < 0.001$), regardless of group (Group x Lever: $p = 0.19$). There was a Lever
by LED light condition interaction (Lever x LED: $F_{(1,18)} = 7.56$; $\eta^2 = 0.29$; $p < 0.05$) and
critically, there was an interaction between group, LED light condition, and Lever
during the presentation of the predictive stimuli (Group x LED x Lever: $F_{(1,18)} = 5.01$;
$\eta^2 = 0.22$; $p < 0.05$). Follow-up analyses revealed that control eYFP rats expressed
outcome-specific PIT whether the LED light was OFF ($F_{(1,7)} = 26.63$; $\eta^2 = 0.79$; $p <$
0.001) or ON ($F_{(1,7)} = 12.31$; $\eta^2 = 0.64$; $p < 0.01$). By contrast, rats in the eNpHR3.0
group displayed outcome-specific PIT when the light was OFF ($F_{(1,11)} = 26.53$; $\eta^2 =$
0.71; $p < 0.001$) but not ON ($p = 0.48$). Thus, these results show for the first time that
NAc-S D1-SPNs mediate outcome-specific PIT via their projections to the VP.
Importantly, we also found that the impairment produced by silencing NAc-S D1-
SPNs terminals in the VP was restricted to the influence of predictive stimuli on
choice between actions. In the same rats, this silencing had no effect on valued-
based choice (Figure S3 F).

NAc-S D2-SPN projections to the VP do not mediate outcome-specific PIT

Since activity in NAc-S D2-SPNs is required for outcome-specific PIT (Figure 3) and
these neurons innervate the VP (Figure 1 F), we assessed whether D2-SPNs
projections to the VP are involved in outcome-specific PIT. A2-Cre rats were
bilaterally infused in the NAc-S with either the null Cre-dependent eYFP virus (eYFP:
4 females and 5 males) or the Cre-dependent halorhodopsin (eNpHR3.0: 5 females
and 6 males) virus and were implanted with fiber-optic cannulas above the VP
(Figure 6 A and Supplementary Figures S4 A-B). The rats then received the
behavioral protocol previously described (Figure 3 B).

Pavlovian and instrumental conditioning went as expected (Supplementary Figures
S4 C-D). The data from the outcome-specific PIT test are presented in Figure 6 B as
previously since baseline responding did not differ between groups (Group: $p = 0.90$;
see also Supplemental Figure 4 E). Silencing NAc-S D2-SPNs projections to the VP
had no effect on outcome-specific PIT. Overall lever press rates were similar
between groups (Group: $p = 0.56$), LED light condition (LED: $p = 0.75$), and the two
factors did not interact (Group x LED: $p = 0.73$). The rates were higher on the action
earning the same outcome as the stimuli relative to the action earning the different
outcome (Lever: $F_{(1,13)} = 22.71$; $\eta^2 = 0.64$; $p < 0.001$), irrespective of group (Group x
Lever: $p = 0.55$). There was no interaction between groups, LED light condition, and
Lever (Group x Light x Lever: $p = 0.59$). Thus, NAc-S D2-SPNs do not appear to
mediate outcome-specific PIT via their projections to the VP. Likewise, we found no
evidence that these projections influence valued-based choice (Figure S4 F).

Discussion

The present experiments investigated the role of NAc-S D1- and D2-SPNs in choice
between actions using an outcome-specific PIT task. First, they combined
anatomical tract-tracing and ex-vivo electrophysiology to demonstrate that two
recently developed knock-in rat lines (Pettibone et al., 2019) enable selective
silencing of activity in either population of NAc-S SPNs. Consistent with previous
findings (Laurent et al., 2014), they found that NAc-S D1-SPNs are necessary for
PIT since their silencing eliminated outcome-specific choice. Additionally, they
showed that this choice is also abolished by silencing NAc-S D2-SPNs, providing the
first evidence that both SPNs populations contribute to PIT expression. Finally, the
last two experiments revealed for the first time that outcome-specific choice is likely
to involve downstream regulation of VP function by NAc-S D1-SPNs but not NAc-S
D2-SPNs. Together, these findings offer novel insights into the cellular mechanisms
that govern the outcome-specific influence of predictive stimuli on choice between
actions.

Convincing evidence had been provided for a significant role of NAc-S D1-SPNs in
PIT (Laurent et al., 2014). The evidence was based on the observations that PIT is
associated with an increase in ERK1/2 phosphorylation within these SPNs and that
pharmacological blockade of NAc-S D1Rs eliminates outcome-specific choice

between actions. These findings align with the present study, showing that PIT is
eliminated when NAc-S D1-SPNs are silenced during presentations of the predictive
stimuli at the time of choice. Importantly, the effect of NAc-S D1-SPNs silencing was
specific to the assessment of choice between actions in the presence of predictive
stimuli. For instance, the ability of the stimuli to elicit approaches towards the
magazine where food used to be delivered remained intact despite NAc-S D1-SPNs
silencing. Furthermore, the same silencing preserved the capacity to select between
the two actions based on the value attributed to their respective outcomes. This
selectivity in the impairment produced by NAc-S D1-SPNs is consistent with previous
findings demonstrating that the NAc-S does not contribute to learning and retrieving
the S-O and A-O associations produced by Pavlovian and instrumental conditioning
(Corbit & Balleine, 2011; Corbit et al., 2001; Morse et al., 2020; Laurent et al., 2012;
Laurent et al., 2014).

Previous research failed to provide any evidence supporting the involvement of NAc-
S D2-SPNs in outcome-specific choice (Laurent et al., 2014). Specifically, PIT was
found to produce no change in ERK1/2 phosphorylation in these neurons and was
left intact by NAc-S D2Rs blockade. However, these assessments have significant
limitations, including the widespread distribution of striatal D2Rs (Gerfen & Surmeier,
2011) and the inability of ERK1/2 phosphorylation or D2Rs antagonism to capture or
distort activity in NAc-S D2-SPNs (Bertran-Gonzalez et al., 2008; Tozzi et al., 2007).
The present study addressed these limitations by implementing optogenetic silencing
in a knock-in rat line giving access to D2-SPNs through the targeting of the A2a
receptor that is predominantly expressed in these neurons in the NAc-S (Schiffmann
et al., 2007). We found that PIT is eliminated when NAc-S D2-SPNs are silenced
during presentations of the predictive stimuli at the time of choice. Moreover, the
same silencing left intact the ability of the predictive stimuli to elicit magazine
approaches and preserved the capacity to choose between actions based on the
value of their outcomes. Thus, we conclude that both NAc-S D1- and D2-SPNs are
indispensable to outcome-selective PIT.

Consistent with the literature (Lu et al., 1997; Kupchik et al., 2015), we found that
both NAc-S D1- and D2-SPNs send dense projections to the VP, which is often
described as the major efferent of the nucleus accumbens (Heimer et al., 1991;
Zahm & Heimer, 1990). Importantly, the VP has been shown to play a crucial role in

outcome-specific choice (Leung & Balleine, 2013; Leung & Balleine, 2015; Leung et al., 2024b). VP neurons exhibit enhanced cFos activation following an outcome-specific PIT test, and their levels of activation correlate with PIT performance. Moreover, pharmacological inactivation of the VP eliminates outcome-specific choice, and the same elimination is observed when the NAc-S is disconnected from the VP. We therefore tested for a role of NAc-S D1- and D2-SPNs projections to the VP during choice between actions in our outcome-specific PIT task. We found that PIT was abolished by silencing NAc-S D1-SPNs terminals but not by silencing NAc-S D2-SPNs terminals. Neither silencing affected magazine approaches elicited by the predictive stimuli or choice between actions based on the value of their outcomes. It is important to note that our study does not provide any evidence about the efficacy of NAc-S D2-SPNs terminals silencing in the VP, and future experiments should aim to provide such evidence. Yet, the same silencing for NAc-S D1-SPNs terminals resulted in PIT elimination. Therefore, it seems reasonable to conclude that NAc-S D1-SPNs, but not NAc-S D2 SPNs, projections to the VP are indispensable to observe outcome-specific choice in a PIT task. Since NAc-S D2-SPNs appear to exclusively project to the VP (Humphries & Prescott, 2010; Kupchik et al., 2015), our findings suggest that these SPNs mediate outcome-specific PIT by locally regulating NAc-S function. By contrast, it remains to be determined whether NAc-S D1-SPNs also coordinate outcome-specific PIT via their projections to the lateral hypothalamus (O'Connor et al., 2015) and/or the ventral tegmental area (Humphries & Prescott, 2010), in addition to the VP.

The present findings are consistent with a recent model proposing that outcome-specific choice in PIT relies on an opioid-based memory residing in the NAc-S (Leung et al., 2024a; Laurent & Balleine, 2021). In this model, as the basolateral amygdala encodes and stores outcome-specific S-O associations across Pavlovian conditioning, it also drives the formation of the NAc-S memory, which involves the durable accumulation of delta-opioid receptors (DOPRs) on the somatic membrane of local cholinergic interneurons (CINs) (Bertran-Gonzalez et al., 2013; Morse et al., 2020). Although this memory is not necessary for Pavlovian conditioning per se, its expression is later required to enable the predictive stimuli to guide outcome-specific choice during the PIT test (Morse et al., 2020). The model specifically proposes that memory expression is controlled by fluctuations in glutamatergic and dopamine

release from cortical and midbrain inputs to the NAc-S. One consequence is to activate NAc-S D2-SPNs and enkephalin discharge by these neurons. Enkephalin then binds onto DOPRs that had accumulated on local CINs and dampens acetylcholine secretion from the interneurons. The sudden drop in NAc-S acetylcholine frees D1-SPNs from the inhibitory tone imposed by acetylcholine occupancy of muscarinic M4 receptors that are exclusively found on these SPNs (Tayebati et al., 2004; Lobo et al., 2006; Guo et al., 2010; Jeon et al., 2010). The ultimate consequence is to promote NAc-S D1-SPNs function, including the coordination of outcome-specific choice in PIT by regulating activity in downstream brain regions such as the VP. The model predicts all the findings presented here. NAc-S D2-SPNs silencing eliminates PIT at the time of choice by preventing enkephalin release and thereby, expression of the DOPR-based memory. The main consequence of this expression is precluded by NAc-S D1-SPNs silencing at the time of choice, while silencing the terminals of these SPNs in the VP squash their ability to regulate the activity of this brain region to coordinate outcome-specific choice in PIT. Thus, one fundamental implication of the present findings is to strengthen the proposal that an opioid-based memory system in the NAc-S enables outcome-specific choice between actions.

In summary, the present experiments found that the two main populations of SPNs in the NAc-S are indispensable for outcome-specific choice between actions in a PIT task. They also revealed that NAc-S D1-SPNs coordinate this choice by downstream regulation of VP activity. By contrast, NAc-S D2-SPNs function in PIT appears to be restricted to modulating local activity. These findings provide novel insights in the cellular mechanisms and circuitry underlying the outcome-specific influence of predictive stimuli on choice between actions, and are consistent with a recent model proposing that this influence is mediated by an opioid-based memory system in the NAc-S. Beyond these mechanistic considerations, the findings offer an opportunity to gain novel insights about various disorders in which the outcome-specific influence of predictive stimuli over our choices and decisions is dysfunctional. These disorders include depression (Geurts et al., 2013; Nord et al., 2018), anxiety disorders (Quail et al., 2017; Krypotos & Engelhard, 2020), substance-use disorders (Heinz et al., 2019; Hogarth et al., 2018; Hogarth et al., 2018; Steins-Loeber et al., 2020; Garbusow et al., 2016; Garbusow et al., 2019; Hogarth et al., 2018), gambling

418   disorders (Genauck et al., 2019), anorexia nervosa (Vogel et al., 2019) and obesity
419   (Watson et al., 2014; Lehner et al., 2017; Meemken & Horstmann, 2019).

Materials and methods

Subjects

Two genetically modified knock-in rat lines were used and obtained from the
breeding facility at the University of New South Wales (Sydney, Australia). D1-Cre
rats (Rat Research & Resource Center, Columbia, MO, USA; LE-Drd1$^{em1(iCre)Berke}$,
RRRC#: 00856) expressed Cre recombinase in neurons expressing the dopamine
D1 receptors (D1R). A2a-Cre rats (Rat Research & Resource Center, Columbia, MO,
USA; LE-*Adora2a$^{em1(iCre)Berke}$*, RRRC#: 00857) expressed Cre recombinase in
neurons expressing the adenosine A2A receptors (A2R). The rats were at least 8
weeks old at the beginning of each experiment. Efforts were made to allocate an
equal number of female and male rats in each group. However, these efforts were
hampered by exclusions following post-mortem assessments of viral spread and
fiber-optic cannulas placement. The final number of female and male rats did not
provide sufficient power to analyse an effect of sex. Therefore, all test data present
individual performance for female and male rats. Rats were housed in transparent
plastic boxes with their littermates (up to four rats per box) in a climate-controlled
colony room maintained on a 12 h light-dark cycle (lights on at 7:00am). Behavioural
procedures were conducted during the light phase (8:00am to 6:00pm). Water and
standard lab chow were available *ad libitum* prior to the start of each experiment.
Rats were food restricted to maintain them at ~90% of their *ad libitum* body weight
during behavioural protocols. The Animal Care and Ethics Committee at the
University of New South Wales approved all procedures.

Viruses

443   The following viruses were used: AAV5-EF1a-DIO-eYFP (eYFP; AddGene;
444   Watertown, MA, USA) and AAV5-EF1a-DIO-eNpHR3.0-eYFP (eNpHR3.0; UNC
445   Vector Core; Chapel Hill, NC, USA).

Surgeries

447   Stereotaxic surgery was conducted under isoflurane gas anaesthesia (0.8L/min;
448   induction at 4-5%, maintenance 2-2.5%). Animals were placed in a stereotaxic frame

(Kopf Instruments, Tujunga, CA, USA), the surgical area was shaved, and betadine iodine antiseptic solution was applied. At the incision site, bupivacaine hydrochloride (0.5%; 0.1-0.2mL) was injected subcutaneously. In the lower flank, Metacam (1mg/kg) was injected subcutaneously. An incision was made along the midline to expose the skull, the membrane on top of the skull was cleared, and the skull was adjusted to align bregma and lambda on the same horizontal plane.

For viral infusion, small holes were drilled in the skull above the NAc-S using the following coordinates (indicated in mm from bregma): +1.8 or +1.9 anteroposterior (A/P); ±0.85 or ±0.95 mediolateral (M/L); -7.7 or 7.85 dorsoventral (D/V). An Infuse/Withdraw Pump (Standard Infuse/Withdraw Pump 11 Elite; Holliston, Harvard Apparatus, MA, USA) in combination with glass 1µL syringes (86200; Hamilton Company, Reno, NV, USA) were used to perform infusions. 0.5µL of the virus was infused in each hemisphere at a rate of 0.1µL/min. Following each infusion, the needle was left in position for 5-10 min to allow for diffusion. The incision site was closed and secured with surgical sutures (684G; Ethicon, North Ryde, NSW, Australia) or surgical staples (59027; Stoelting, Wood Dale, IL, USA), and betadine antiseptic ointment was applied to the area. Animals then received subcutaneous injections of the antibiotic Duplocillin (0.15ml/kg) and sodium chloride (0.9%, 2mL) and placed on a heat mat for recovery. Once alert and responsive, animals were returned to their home cages and monitored daily as they recovered for a minimum of one week following surgery with *ad libitum* access to food and water.

Rats received a second surgery to bilaterally implant fibre-optic cannulas (10mm length, 0.48mm diameter; Doric Lenses, Québec, QC, Canada) after completion of the initial 16 days of training (Pavlovian and Instrumental). The surgical area was prepared using the same method as described for viral surgeries. Three jewellers' screws were inserted into the skull, distributed anterior and posterior to the implantation site and secured in place with dental cement. Fibre-optic cannulas were implanted into the targeted brain region. The following coordinates (indicated in mm from bregma) were used for the NAc-S: +1.8 or +1.9 A/P; -0.85 or -0.95 M/L for left hemisphere; +2.2 M/L with -10° angle for right hemisphere; -6.75 D/V for left hemisphere; -6.8 or -6.9 D/V for right hemisphere. The following coordinates (indicated in mm from bregma) were used for the VP: +0.24 A/P; ±1.8 M/L; -7.5 or -

481 8.0 D/V. Post-implantation, rats received the same care as described for viral
482 surgeries.

Behavioral apparatus

Training and testing were conducted in 12 identical operant chambers (ENV-007-VP;
L 29.53 x W 23.5 x H 27.31 cm; MED Associates, Fairfax, VT, USA) enclosed within
light- and sound-resistant cabinets (ENV-018MD; L 59.69 x W 40.64 x H 55.88 cm;
MED Associates, Fairfax, VT, USA). The chambers consisted of two stainless-steel
side walls, three clear polycarbonate walls and ceiling, and a stainless-steel grid
floor. Each chamber was equipped with a recessed magazine connected to a 45mg
grain pellet (F0165; Bio-Serv Biotechnologies, Flemington, NJ, USA) dispenser and
a pump fitted with a 60mL syringe that delivered a 0.2mL sucrose solution (20% w/v;
white sugar; Winc, Erskine Park, NSW, Australia) into the left side of the magazine.
Head entries into the magazine were detected via an infrared beam that crossed the
magazine opening. One retractable lever was located on either side of the magazine.
A house light (3W, 24V) situated at the top of the wall opposite to the food magazine
was used to illuminate the chamber during the training and test sessions. Each
chamber contained a Sonalert white noise generator (80dB), and a 28V DC
mechanical relay that delivered a 2Hz clicker stimulus. Training and testing sessions
were programmed and controlled by computers external to the testing rooms using
MED-PC V software (MED Associates, Fairfax, VT, USA) which also recorded
experimental data from each session. All cabinets were fitted with cameras (IPC-
K35AP; Dahua, Artarmon, NSW, Australia) to record and view real-time chamber
activity using D-ViewCam DCS-100 software (D-Link, North Ryde, NSW, Australia).
The ceiling of these chambers contained a hole which enabled the connection of the
patch cable between the LED and each freely moving rat.

Outcome devaluation was conducted in a separate room equipped with 16 individual,
open-top plastic boxes with stainless steel wire mesh lids. During devaluation, the
lights were dimmed, and each individual chamber was fitted with either a ceramic
dish for grain pellet devaluation, or a plastic sipper bottle for sucrose solution
devaluation.

Optogenetic equipment

Fibre-optic cannulas (10mm; Doric Lenses, Québec, QC, Canada) consisted of a
zircona ferrule contained within a stainless-steel receptacle from which a
silica/polymer flat-tip fibre extended from. Fibre-optic patch cords (custom; Doric
Lenses, Québec, QC, Canada) were enclosed in a flexible cladding that consisted of
one transparent fibre that split into two smaller fibres for attachment to the implanted
fibre-optic cannulas. Delivery of the LED light was controlled through a TTL adapter
which converted 28V DC output to a TTL transition (MED Associates, Fairfax, VT,
USA). The adaptor was connected to an LED driver (Doric Lenses, Québec, QC,
Canada) and Connectorised LED light source (Doric Lenses, Québec, QC, Canada)
with a fibre-optic rotary joint for attachment to the fibre-optic patch cords. LED light
was delivered as orange light (625nm) and measured at least 10mW at the cannula
tip using a photometer (PM200; ThorLabs, Newton, NJ, USA) when the LED was
switched on.

Behavioral procedures

Pavlovian conditioning

Pavlovian conditioning involved 8 daily 1hr training sessions. Two auditory
conditioned stimuli (S1 and S2; white noise or clicker) were paired with two food
outcomes (O1 and O2; grain pellets or sucrose solution). Each stimulus was
presented for 2min in duration, 4 times each in a pseudorandom order within each
training session. Stimulus presentations were separated by pseudorandom intertrial
intervals (ITIs) ranging between 3 to 5min in length (4min average). The stimulus
order and ITIs were changed daily. Stimulus-outcome (S-O) pairings were
counterbalanced within- and between- groups. Food outcomes were delivered on a
random time 30s schedule throughout the stimulus presentation. Magazine entries
were recorded throughout the session, and conditioned responding was analysed by
comparing magazine entries in the 2min prior to stimulus presentation and during the
2min stimulus presentation period.

Instrumental conditioning

During instrumental conditioning, two actions (A1 and A2; left or right lever presses)
were trained with the two different outcomes (O1 and O2) in 2 separate daily
sessions for 8 days. The order of the sessions was counterbalanced, as were the A-
O relationships, which were further counterbalanced with the S-O relationships. Each

session ended when 20 outcomes were earned or when 30min had elapsed. For the first 2 days, actions were continuously reinforced (i.e. one action earned one outcome delivery). Across the next three sessions, actions were rewarded on a random ratio 5 (RR5) schedule (i.e. each action delivered an outcome with a probability of 0.2). For the final 3 days, animals were rewarded on an RR10 schedule (i.e. each response delivered an outcome with a probability of 0.1). In the last experiment (NAc-S D1-SPN→VP projections), rats only received 2 days on RR10. Lever press actions were measured throughout the duration of the session, and performance during instrumental training was assessed using the mean number of lever presses per min.

Pavlovian Instrumental transfer test

Across the three days preceding this test, rats were given one session of Pavlovian retraining and two separate sessions of instrumental retraining. Immediately before the test, patch cables were attached to the fibre-optic cannulas and the LED. During the test, both levers were inserted into the chamber simultaneously and remained in the chamber for the duration of the test. The test was conducted in extinction (i.e. no outcomes were delivered). To reduce baseline (pre S) lever presses, responding on both levers was extinguished for 8min prior to any stimuli being presented. Each stimulus was presented for 2min, four times each during the test in the following order: noise-clicker-clicker-noise-clicker-noise-noise-clicker. Each stimulus presentation was separated by a fixed 3min interval. Critically, the LED was activated during half of the trials (ON) but left inactive during the other half (OFF). The order for ON and OFF trials was counterbalanced. Responses were recorded prior to and during the S presentation. These were then collated into 'pre' (lever press rates in the two min prior to S presentation), 'same' (when S shares the *same* outcome with the action) and 'different' (when S has a *different* outcome with the action) following data collection.

Choice tests following outcome devaluation

Across the two consecutive days preceding these tests, rats were given 2 daily instrumental conditioning sessions (RR10). In the last experiment (NAc-S D1-SPNs→VP projections), rats first received one day of Pavlovian conditioning followed by 3 days of instrumental conditioning (RR10). On the last two days of instrumental

conditioning, following the second session, rats were habituated to the devaluation
chambers for 30min. During these habituation sessions, animals had access to one
outcome (counterbalanced). On the first test day, rats were placed in the devaluation
chambers for 1hr where they had *ad libitum* access to one of the previously trained
outcomes (O1 or O2). Once removed from the devaluation boxes, patch cables were
attached to the fibre-optic cannulas and the LED and the first choice test was
conducted. Both levers were made available for 2min each, four times during the
session with an intertrial interval of 2min where the levers were retracted. The LED
was activated during 2 of the 2min trials where the levers were available (ON) and
was left inactive during the other 2 trials (OFF). The order of the ON and OFF trials
was counterbalanced. A similar procedure was implemented the following day except
that the other outcome was devalued. Lever presses and magazine entries were
recorded while the levers were available.

Tissue preparation and histology

Transcardial fixation and brain sectioning

Rats were euthanised with an intraperitoneal injection of sodium pentobarbital (1mL)
and transcardially perfused with 4% paraformaldehyde (PFA) in 0.1M sodium
phosphate buffer (PBS, pH 7.5; 400mL for rats). Brains were immediately removed
and placed in individual specimen jars containing PFA and stored at 4°C overnight
for post-fixation. Brains were sliced into 40µm coronal or sagittal sections in 0.1M
Tris-buffered saline (TBS, 0.25M Tris, 0.5M NaCl, pH 7.5) using a vibratome
(VT1200; Leica Microsystems, North Ryde, NSW, Australia) and stored at -20°C in
cryoprotective solution containing 30% ethylene glycol, 30% glycerol and 0.1M PBS.

Immunofluorescence

Free-floating sections were selected in TBS to contain the region of interest and
washed three times for 10 min each in TBS solution. Sections were then incubated in
0.5% Triton-X, 10% NHS in TBS blocking solution for two hours. Sections were
washed again in TBS solution three times for 10 min each. Sections were then
incubated between 12-48h (dependent on antibody) in a solution containing the
primary antibodies in 0.2% Triton-X, 2% NHS in TBS at 4°C. The primary antibodies
were the following: purified mouse anti DARPP-32 (1:300; BD Biosciences,
Macquarie Park, Australia; #AB398980) and polyclonal rabbit anti-GFP (1:1000;

Invitrogen, Mulgrave, Australia; #AB221569). After the specified incubation period,
the sections were washed for three times in TBS for 10min each and then
transferred into a solution containing the secondary antibodies in TBS solution at
room temperature for 1h. The secondary antibodies were the following: donkey anti-
mouse Cy3 (1:400; Jackson ImmunoResearch Laboratories, West Grove, USA;
#AB2340813) and donkey anti-rabbit Alexa-488 (1:500; Jackson ImmunoResearch
Laboratories, West Grove, USA; #AB2313584). Next, sections were washed three
times in TBS for 10min and then mounted onto glass slides using Vectashield
hardset antifade mounting medium (H-1400; Vector Laboratories, Newark, CA,
USA). Sections that did not require immunofluorescence staining were washed three
times for 10min each in TBS and mounted using Vectashield hardset antifade
mounting medium.

Microscopy and imaging

Imaging of the mounted brain sections was performed on a spinning disk confocal
microscope (Zyla 4.2 sCMOS; Andor Technology, Belfast, UK) using the Diskovery
multi-modal imaging platform (Oxford Instruments, Belfast, UK). The Diskovery
platform was mounted on an inverted microscope body with a motorised stage and
Perfect Focus System (Eclipse Ti; Nikon, Melville, NY, USA). Image capture was
controlled using the NIS-Elements Software (Nikon, Melville, NY, USA). Single high-
magnification field-of-view images were generated by automatic stitching of multiple
adjacent frames from within a defined area using the motorised stage (optical
magnification: 20X; pixel depth: 16-bit). All images were processed using Fiji ImageJ
open-source software (Rueden et al., 2017). Additional images were taken with a
confocal microscope (BX61W1; Olympus, Shinjuku, Tokyo, Japan) using Fluoview
software (FV1000; Olympus, Shinjuku, Tokyo, Japan) for further placement analysis.
The images were used to assess neuronal pathways, as well as the spread of viral
infusions and placement of fibre-optic cannulas. Based on the latter, the following
numbers of rats were excluded from the final statistical analyses due to misplaced
viral or fibre-optics misplacements: NAc-S D1-SPNs silencing (n=18), NAc-S D2-
SPNs silencing (n=9), NAc-S D1-SPNs→VP silencing (n=X), NAc-S D2-SPNs→VP
silencing (n=5).

Electrophysiological recordings

Brain slice preparation

Rats were euthanized under isoflurane gas anesthesia (4% in air). Brains were immediately removed and sliced using a vibratome (VT1200; Leica Microsystems, North Ryde, NSW, Australia) in ice-cold oxygenated sucrose buffer solution (consisting of [in mM]: 241 sucrose, 28 $NaHCO_3$, 11 glucose, 1.4 NaH_2PO_4, 3.3 KCl, 0.2 $CaCl_2$, 7 $MgCl_2$). Coronal brain slices (300µm thickness) containing the NAc were sampled and maintained at 33°C in a submerged chamber containing physiological saline (consisting of [in mM]: 124 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 1 $MgCl_2$, 1 $CaCl_2$, 10 glucose and 26 $NaHCO_3$) and equilibrated with 95% O_2 and 5% CO_2.

Recordings

Following 1h equilibration, slices were transferred to a recording chamber. An upright microscope (BX50WI; Olympus, Shinjuku, Tokyo, Japan) was used to visualise neurons using differential interference contrast (DIC) Dodt tube optics and eYFP fluorescence, and superfused continuously (1.5mL/min) with oxygenated physiological saline at 33°C. Whole-cell patch-clamp recordings were performed using glass pipettes (tip size 2–5 MΩ) containing internal solution (consisting of [in mM]: 115 K gluconate, 20 NaCl, 1 $MgCl_2$, 10 HEPES, 11 EGTA, 5 Mg-ATP, 0.33 Na-GTP, and 5 phosphocreatine di(tris) salt, pH 7.3, osmolarity 285-290 mOsm/L). Neurons sampled during recording were marked with Biocytin (0.1%) added to the internal solution. A Multiclamp 700B amplifier (Molecular Devices, San Jose, CA, USA) connected to a Macintosh computer and interface ITC-18 (InstruTECH, Longmont, CO, USA) was used for data acquisition. Liquid junction potentials of -10mV were not corrected. In current-clamp mode, membrane potentials were sampled at 5kHz (low pass filter 2 kHz; Axograph X, Axograph, CA, USA). Stock solution of drug was diluted to working concentration in the extracellular solution immediately before use and applied by continuous superfusion. Data from whole-cell recordings were included in analyses if the neurons appeared healthy under DIC on the monitor screen.

LED stimulation

Transfected neurons with eNpHR3.0 or eYFP in the NAc-S were illuminated by a continuous wave of LED light (625nm, 2 minutes, 2mW; ThorLabs, Newton, NJ,

USA) onto the brain slice under the 40X water-immersion objective. Since SPNs in

ex-vivo brain slices are not spontaneously active, brief injection of positive current

step pulses were applied to make them fire action potentials.

Post-hoc histological analysis

Immediately after physiological recording, brain slices were fixed overnight using 4%

PFA in 0.16M phosphate buffer (PB) solution, washed and then transferred into 0.5%

Triton-X in PB solution for 3 days to permeabilise cells. Slices were washed,

mounted onto glass slides, and then coverslipped with Fluoromount-G mounting

medium (Southern Biotech, Birmingham, AL, USA). A 2D projection image was then

obtained from a collated image stack using the confocal microscope (X61W1;

Olympus, Shinjuku, Tokyo, Japan).

Statistical analyses

The data presented met the assumptions of the statistical test used. The main

measures were the number of magazine entries and lever presses per minute. The

difference between groups were analyzed using a planned orthogonal contrast

procedure controlling the per contrast error rate (Hays, 1963). A first contrast (Group)

compared rats that had been infused with the control eYFP virus and rats that had

been infused with the inhibitory eNpHR3.0 virus. Changes across days (Day) during

Pavlovian and instrumental conditioning were assessed by planned linear trend

analyses. Pavlovian conditioning and the PIT tests included a within-subject factor

(Period) comparing magazine entry rates during the 2min stimuli against rates during

a 2min baseline period immediately preceding the stimuli. During the PIT tests, a

within-subject factor (Lever) examined the net (i.e., minus baseline) rates of lever

presses on the response (Same) earning the same outcome as predicted by the

stimulus against those on the response (Different) earning the different outcome. A

second within-subject factor (LED) examined these rats when the LED was activated

(ON) or not activated (OFF). The choice tests following outcome devaluation used a

within-subject factor of response identity (Lever) comparing rates of lever presses on

the response (Valued) earning the valued outcome against those on the response

(Devalued) earning the devalued outcome. It also used the LED within-subject factor

described above. All analyses were carried out using the PSY statistical software

(School of Psychology, the University of New South Wales, Australia). The Type I

error rate was controlled at alpha = 0.05 for each contrast tested. If interactions were
detected, follow-up simple effects analyses were calculated to determine the source
of the interactions. For each significant statistical comparison, we report measures of
effect size partial eta-squared ($\eta2$; $\eta2 = 0.01$ is a small effect, $\eta2 = 0.06$ a medium
effect and $\eta2 = 0.14$ a large effect). Data will be made available upon request.

Acknowledgements

710 The research reported in this manuscript was supported by a Future Fellowship from
711 the Australian Research Council (FT220100474) to V.L.

References

Bertran-Gonzalez, J., Hervé, D., Girault, J. A., & Valjent, E. (2010). What is the
Degree of Segregation between Striatonigral and Striatopallidal Projections.
Front Neuroanat, *4*, 136. https://doi.org/10.3389/fnana.2010.00136

Bertran-Gonzalez, J., & Laurent, V. (2018). Studying Integrative Processing and
Prospected Plasticity in Cholinergic Interneurons: The Importance of Pavlovian–
Instrumental Transfer. *Goal-Directed Decision Making*.

Bertran-Gonzalez, J., Bosch, C., Maroteaux, M., Matamales, M., Herve, D., Valjent,
E., & Girault, J.-A. (2008). Opposing Patterns of Signaling Activation in
Dopamine D1 and D2 Receptor-Expressing Striatal Neurons in Response to
Cocaine and Haloperidol. *Journal of Neuroscience*, *28*(22), 5671-5685.
https://doi.org/10.1523/jneurosci.1039-08.2008

Bertran-Gonzalez, J., Laurent, V., Chieng, B. C., Christie, M. J., & Balleine, B. W.
(2013). Learning-related translocation of δ-opioid receptors on ventral striatal
cholinergic interneurons mediates choice between goal-directed actions. *J
Neurosci*, *33*(41), 16060-16071. https://doi.org/10.1523/JNEUROSCI.1927-
13.2013

Bradfield, L. A., Dezfouli, A., van Holstein, M., Chieng, B., & Balleine, B. W. (2015).
Medial Orbitofrontal Cortex Mediates Outcome Retrieval in Partially Observable
Task Situations. *Neuron*, *88*(6), 1268-1280.
https://doi.org/10.1016/j.neuron.2015.10.044

Bradfield, L. A., Hart, G., & Balleine, B. W. (2018). Inferring action-dependent
outcome representations depends on anterior but not posterior medial
orbitofrontal cortex. *Neurobiol Learn Mem*, *155*, 463-473.
https://doi.org/10.1016/j.nlm.2018.09.008

Cartoni, E., Balleine, B., & Baldassarre, G. (2016). Appetitive Pavlovian-instrumental
Transfer: A review. *Neurosci Biobehav Rev*, *71*, 829-848.
https://doi.org/10.1016/j.neubiorev.2016.09.020

Colwill, R. M., & Rescorla, R. A. (1988). Associations between the discriminative
stimulus and the reinforcer in instrumental learning. *Journal of Experimental*
Psychology: Animal Behavior Processes, *14*(2), 155-164.

Corbit, L. H., Muir, J. L., Balleine, B. W., & Balleine, B. W. (2001). The role of the
nucleus accumbens in instrumental conditioning: Evidence of a functional
dissociation between accumbens core and shell. *J Neurosci*, *21*(9), 3251-3260.
http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=11312
310&retmode=ref&cmd=prlinks

Corbit, L. H., & Balleine, B. W. (2005). Double dissociation of basolateral and central
amygdala lesions on the general and outcome-specific forms of pavlovian-
instrumental transfer. *J Neurosci*, *25*(4), 962-970.
https://doi.org/10.1523/JNEUROSCI.4507-04.2005

Corbit, L. H., & Balleine, B. W. (2011). The general and outcome-specific forms of
Pavlovian-instrumental transfer are differentially mediated by the nucleus
accumbens core and shell. *J Neurosci*, *31*(33), 11786-11794.
https://doi.org/10.1523/JNEUROSCI.2711-11.2011

Corbit, L. H., Janak, P. H., & Balleine, B. W. (2007). General and outcome-specific
forms of Pavlovian-instrumental transfer: the effect of shifts in motivational state
and inactivation of the ventral tegmental area. *Eur J Neurosci*, *26*(11), 3141-
3149. https://doi.org/10.1111/j.1460-9568.2007.05934.x

Corbit, L. H., Fischbach, S. C., & Janak, P. H. (2016). Nucleus accumbens core and
shell are differentially involved in general and outcome-specific forms of
Pavlovian-instrumental transfer with alcohol and sucrose rewards. *Eur J*
Neurosci, *43*(9), 1229-1236. https://doi.org/10.1111/ejn.13235

Corbit, L. H., & Janak, P. H. (2010). Posterior dorsomedial striatum is critical for both
selective instrumental and Pavlovian reward learning. *Eur J Neurosci*, *31*(7),
1312-1321. https://doi.org/10.1111/j.1460-9568.2010.07153.x

Derman, R. C., Bass, C. E., & Ferrario, C. R. (2020). Effects of hM4Di activation in
CamKII basolateral amygdala neurons and CNO treatment on sensory-specific
vs. general PIT: refining PIT circuits and considerations for using CNO.
Psychopharmacology (Berl), *237*(5), 1249-1266.
https://doi.org/10.1007/s00213-020-05453-8

Garbusow, M., Nebe, S., Sommer, C., Kuitunen-Paul, S., Sebold, M., Schad, D. J.,
Friedel, E., Veer, I. M., Wittchen, H. U., Rapp, M. A., Ripke, S., Walter, H.,
Huys, Q. J. M., Schlagenhauf, F., Smolka, M. N., & Heinz, A. (2019). Pavlovian-
To-Instrumental Transfer and Alcohol Consumption in Young Male Social
Drinkers: Behavioral, Neural and Polygenic Correlates. *J Clin Med*, *8*.
https://doi.org/10.3390/jcm8081188

Garbusow, M., Schad, D. J., Sebold, M., Friedel, E., Bernhardt, N., Koch, S. P.,
Steinacher, B., Kathmann, N., Geurts, D. E., Sommer, C., Müller, D. K., Nebe,
S., Paul, S., Wittchen, H. U., Zimmermann, U. S., Walter, H., Smolka, M. N.,
Sterzer, P., Rapp, M. A., Huys, Q. J., Schlagenhauf, F., & Heinz, A. (2016).
Pavlovian-to-instrumental transfer effects in the nucleus accumbens relate to
relapse in alcohol dependence. *Addict Biol*, *21*(3), 719-731.
https://doi.org/10.1111/adb.12243

Genauck, A., Andrejevic, M., Brehm, K., Matthis, C., Heinz, A., Weinreich, A.,
Kathmann, N., & Romanczuk-Seiferth, N. (2019). Cue-induced effects on
decision-making distinguish subjects with gambling disorder from healthy
controls. *Addict Biol*, e12841. https://doi.org/10.1111/adb.12841

Gerfen, C. R., & Surmeier, D. J. (2011). Modulation of striatal projection systems by
dopamine. *Annu Rev Neurosci*, *34*, 441-466. https://doi.org/10.1146/annurev-
neuro-061010-113641

Geurts, D. E. M., Huys, Q. J. M., den Ouden, H. E. M., & Cools, R. (2013). Serotonin
and aversive pavlovian control of instrumental behavior in humans. *J Neurosci*,
33(48), 18932-18939. https://doi.org/10.1523/JNEUROSCI.2749-13.2013

Guo, M. L., Fibuch, E. E., Liu, X. Y., Choe, E. S., Buch, S., Mao, L. M., & Wang, J.
Q. (2010). CaMKIIalpha interacts with M4 muscarinic receptors to control
receptor and psychomotor function. *EMBO J, 29*(12), 2070-2081.
https://doi.org/10.1038/emboj.2010.93

Hays, W. L. (1963). *Statistics for psychologists*. Holt, Rinehart and Winston.

Heimer, L., Zahm, D. S., Churchill, L., & Kalivas…, P. W. (1991). Specificity in the
projection patterns of accumbal core and shell in the rat. *Neuroscience*.
https://www.sciencedirect.com/science/article/pii/030645229190202Y

Heinz, A., Beck, A., Halil, M. G., Pilhatsch, M., Smolka, M. N., & Liu, S. (2019).
Addiction as Learned Behavior Patterns. *J Clin Med, 8*.
https://doi.org/10.3390/jcm8081086

Hogarth, L., Lam-Cassettari, C., Pacitti, H., Currah, T., Mahlberg, J., Hartley, L., &
Moustafa, A. (2018). Intact goal-directed control in treatment-seeking drug
users indexed by outcome-devaluation and Pavlovian to instrumental transfer:
critique of habit theory. *European Journal of Neuroscience*.
https://doi.org/10.1111/ejn.13961

Hollis, K. L. (1984). The biological function of Pavlovian conditioning: the best
defense is a good offense. *Journal of Experimental Psychology: Animal
Behavior Processes, 10*(4), 413. https://psycnet.apa.org/record/1985-19329-
001

Holmes, N. M., Marchand, A. R., & Coutureau, E. (2010). Pavlovian to instrumental
transfer: a neurobehavioural perspective. *Neurosci Biobehav Rev, 34*(8), 1277-
1295. https://doi.org/10.1016/j.neubiorev.2010.03.007

Humphries, M. D., & Prescott, T. J. (2010). The ventral basal ganglia, a selection
mechanism at the crossroads of space, strategy, and reward. *Prog Neurobiol,
90*(4), 385-417. https://doi.org/10.1016/j.pneurobio.2009.11.003

Jeon, J., Dencker, D., Wörtwein, G., Woldbye, D. P., Cui, Y., Davis, A. A., Levey, A.
I., Schütz, G., Sager, T. N., Mørk, A., Li, C., Deng, C. X., Fink-Jensen, A., &
Wess, J. (2010). A subpopulation of neuronal M4 muscarinic acetylcholine

receptors plays a critical role in modulating dopamine-dependent behaviors. *J Neurosci*, *30*(6), 2396-2405. https://doi.org/10.1523/JNEUROSCI.3843-09.2010

Keistler, C., Barker, J. M., & Taylor, J. R. (2015). Infralimbic prefrontal cortex interacts with nucleus accumbens shell to unmask expression of outcome-selective Pavlovian-to-instrumental transfer. *Learn Mem*, *22*(10), 509-513. https://doi.org/10.1101/lm.038810.115

Krypotos, A.-M., & Engelhard, I. M. (2020). Pavlovian-to-instrumental transfer in subclinical obsessive–compulsive disorder. *Journal of Experimental Psychopathology*, *11*(3), 2043808720925244. https://journals.sagepub.com/doi/pdf/10.1177/2043808720925244

Kupchik, Y. M., Brown, R. M., Heinsbroek, J. A., & Lobo…, M. K. (2015). Coding the direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections. *Nature …*. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4551610

Laurent, V., & Balleine, B. W. (2021). How predictive learning influences choice: Evidence for a GPCR-based memory process necessary for Pavlovian-instrumental transfer. *Journal of Neurochemistry*, *157*(5), 1436-1449. https://doi.org/10.1111/jnc.15339

Laurent, V., Bertran-Gonzalez, J., Chieng, B. C., & Balleine, B. W. (2014). δ-Opioid and Dopaminergic Processes in Accumbens Shell Modulate the Cholinergic Control of Predictive Learning and Choice. *J Neurosci*, *34*(4), 1358-1369. https://doi.org/10.1523/JNEUROSCI.4592-13.2014

Laurent, V., Leung, B., Maidment, N., & Balleine, B. W. (2012). μ- and δ-opioid-related processes in the accumbens core and shell differentially mediate the influence of reward-guided and stimulus-guided decisions on choice. *J Neurosci*, *32*(5), 1875-1883. https://doi.org/10.1523/JNEUROSCI.4688-11.2012

Laurent, V., Wong, F. L., & Balleine, B. W. (2015). δ-Opioid receptors in the accumbens shell mediate the influence of both excitatory and inhibitory predictions on choice. *Br J Pharmacol*, *172*(2), 562-570. https://doi.org/10.1111/bph.12731

Lehner, R., Balsters, J. H., Bürgler, A., Hare, T. A., & Wenderoth, N. (2017). Food-
Predicting stimuli Differentially influence eye Movements and goal-Directed
Behavior in normal-Weight, Overweight, and Obese individuals. *Frontiers in*
psychiatry, *8*, 230. https://doi.org/10.3389/fpsyt.2017.00230

Leung, B. K., & Balleine, B. W. (2013). The ventral striato-pallidal pathway mediates
the effect of predictive learning on choice between goal-directed actions. *J*
Neurosci, *33*(34), 13848-13860. https://doi.org/10.1523/JNEUROSCI.1697-
13.2013

Leung, B. K., & Balleine, B. W. (2015). Ventral pallidal projections to mediodorsal
thalamus and ventral tegmental area play distinct roles in outcome-specific
Pavlovian-instrumental transfer. *J Neurosci*, *35*(12), 4953-4964.
https://doi.org/10.1523/JNEUROSCI.4837-14.2015

Leung, B. K., Laurent, V., & Balleine, B. W. (2024a). The influence of predictive
learning on choice and decision-making and its neural bases. *Elsevier*.
https://doi.org/10.1016/B978-0-443-15754-7.00067-5

Leung, B. K., Chieng, B., Becchi, S., & Balleine, B. W. (2024b). A ventral pallidal-
thalamocortical circuit mediates the cognitive control of instrumental action.
Curr Biol, S0960-9822(24)00763. https://doi.org/10.1016/j.cub.2024.06.011

Lichtenberg, N. T., Pennington, Z. T., Holley, S. M., Greenfield, V. Y., Cepeda, C.,
Levine, M. S., & Wassum, K. M. (2017). Basolateral Amygdala to Orbitofrontal
Cortex Projections Enable Cue-Triggered Reward Expectations. *J Neurosci*,
37(35), 8374-8384. https://doi.org/10.1523/JNEUROSCI.0486-17.2017

Lichtenberg, N. T., Sepe-Forrest, L., Pennington, Z. T., Lamparelli, A. C., Greenfield,
V. Y., & Wassum, K. M. (2021). The Medial Orbitofrontal Cortex-Basolateral
Amygdala Circuit Regulates the Influence of Reward Cues on Adaptive
Behavior and Choice. *J Neurosci*, *41*(34), 7267-7277.
https://doi.org/10.1523/JNEUROSCI.0901-21.2021

Lobo, M. K., Karsten, S. L., Gray, M., Geschwind, D. H., & Yang, X. W. (2006).
FACS-array profiling of striatal projection neuron subtypes in juvenile and adult
mouse brains. *Nat Neurosci*, *9*(3), 443-452. https://doi.org/10.1038/nn1654

Lu, X. Y., Ghasemzadeh, M. B., & Kalivas, P. W. (1997). Expression of D1 receptor,
D2 receptor, substance P and enkephalin messenger RNAs in the neurons
projecting from the nucleus accumbens. *Neuroscience*.
https://www.sciencedirect.com/science/article/pii/S0306452297003278

Matamales, M., McGovern, A. E., Mi, J. D., Mazzone, S. B., Balleine, B. W., &
Bertran-Gonzalez, J. (2020). Local D2- to D1-neuron transmodulation updates
goal-directed learning in the striatum. *Science*, *367*(6477), 549-555.
https://doi.org/10.1126/science.aaz5751

Meemken, M.-T., & Horstmann, A. (2019). Appetitive Pavlovian-to-Instrumental
Transfer in Participants with Normal-Weight and Obesity. *Nutrients*, *11*(5),
1037. https://doi.org/10.3390/nu11051037

Morse, A. K., Leung, B. K., Heath, E., Bertran-Gonzalez, J., Pepin, E., Chieng, B. C.,
Balleine, B. W., & Laurent, V. (2020). Basolateral Amygdala Drives a GPCR-
Mediated Striatal Memory Necessary for Predictive Learning to Influence
Choice. *Neuron*, *106*(5), 855-869.e8.
https://doi.org/10.1016/j.neuron.2020.03.007

Nord, C. L., Lawson, R. P., Huys, Q. J. M., Pilling, S., & Roiser, J. P. (2018).
Depression is associated with enhanced aversive Pavlovian control over
instrumental behaviour. *Sci Rep*, *8*(1), 12582. https://doi.org/10.1038/s41598-
018-30828-5

O'Connor, E. C., Kremer, Y., Lefort, S., Harada, M., Pascoli, V., Rohner, C., &
Lüscher, C. (2015). Accumbal D1R Neurons Projecting to Lateral
Hypothalamus Authorize Feeding. *Neuron*, *88*(3), 553-564.
https://doi.org/10.1016/j.neuron.2015.09.038

Ostlund, S. B., & Balleine, B. W. (2007). Orbitofrontal cortex mediates outcome
encoding in Pavlovian but not instrumental conditioning. *J Neurosci*, *27*(18),
4819-4825. https://doi.org/10.1523/JNEUROSCI.5443-06.2007

Ostlund, S. B., & Balleine, B. W. (2008). Differential involvement of the basolateral
amygdala and mediodorsal thalamus in instrumental action selection. *J
Neurosci*, *28*(17), 4398-4405. https://doi.org/10.1523/JNEUROSCI.5472-
07.2008

Pettibone, J. R., Yu, J. Y., Derman, R. C., Faust, T. W., Hughes, E. D., Filipiak, W.
E., Saunders, T. L., Ferrario, C. R., & Berke, J. D. (2019). Knock-In Rat Lines
with Cre Recombinase at the Dopamine D1 and Adenosine 2a Receptor Loci.
eNeuro, 6(5). https://doi.org/10.1523/ENEURO.0163-19.2019

Prévost, C., Liljeholm, M., Tyszka, J. M., & O'Doherty, J. P. (2012). Neural correlates
of specific and general Pavlovian-to-Instrumental Transfer within human
amygdalar subregions: a high-resolution fMRI study. *J Neurosci, 32*(24), 8383-
8390. https://doi.org/10.1523/JNEUROSCI.6237-11.2012

Quail, S. L., Morris, R. W., & Balleine, B. W. (2017). Stress associated changes in
Pavlovian-instrumental transfer in humans. *Q J Exp Psychol (Hove), 70*(4), 675-
685. https://doi.org/10.1080/17470218.2016.1149198

Schiffmann, S. N., Fisone, G., Moresco, R., Cunha, R. A., & Ferré, S. (2007).
Adenosine A2A receptors and basal ganglia physiology. *Prog Neurobiol, 83*(5),
277-292. https://doi.org/10.1016/j.pneurobio.2007.05.001

Seitz, B. M., Hoang, I. B., DiFazio, L. E., Blaisdell, A. P., & Sharpe, M. J. (2022).
Dopamine errors drive excitatory and inhibitory components of backward
conditioning in an outcome-specific manner. *Current Biology*.
https://doi.org/10.1016/j.cub.2022.06.035

Shiflett, M. W., & Balleine, B. W. (2010). At the limbic-motor interface: disconnection
of basolateral amygdala from nucleus accumbens core and shell reveals
dissociable components of incentive motivation. *Eur J Neurosci, 32*(10), 1735-
1743. https://doi.org/10.1111/j.1460-9568.2010.07439.x

Sias, A. C., Jafar, Y., Goodpaster, C. M., Ramírez-Armenta, K., Wrenn, T. M., Griffin,
N. K., Patel, K., Lamparelli, A. C., Sharpe, M. J., & Wassum, K. M. (2024).
Dopamine projections to the basolateral amygdala drive the encoding of
identity-specific reward memories. *Nat Neurosci*.
https://doi.org/10.1038/s41593-024-01586-7

Sias, A. C., Morse, A. K., Wang, S., Greenfield, V. Y., Goodpaster, C. M., Wrenn, T.
M., Wikenheiser, A., Holley, S. M., Cepeda, C., Levine, M. S., & Wassum, K. M.
(2021). A bidirectional corticoamygdala circuit for the encoding and retrieval of

detailed reward memories. *Elife*, *10*, e68617.
https://doi.org/10.7554/eLife.68617

Steins-Loeber, S., Lörsch, F., van der Velde, C., Müller, A., Brand, M., Duka, T., &
Wolf, O. T. (2020). Does acute stress influence the Pavlovian-to-instrumental
transfer effect? Implications for substance use disorders. *Psychopharmacology*
(Berl). https://doi.org/10.1007/s00213-020-05534-8

Tayebati, S. K., Di Tullio, M. A., & Amenta, F. (2004). Age-related changes of
muscarinic cholinergic receptor subtypes in the striatum of Fisher 344 rats. *Exp*
Gerontol, *39*(2), 217-223. https://doi.org/10.1016/j.exger.2003.10.016

Tensaouti, Y., Morel, L., & Parkes, S. L. (2025). Contribution of rat insular cortex to
stimulus-guided action. *J Neurosci*, e1923242025.
https://doi.org/10.1523/JNEUROSCI.1923-24.2025

Tozzi, A., Tscherter, A., Belcastro, V., Tantucci, M., Costa, C., Picconi, B., Centonze,
D., Calabresi, P., & Borsini, F. (2007). Interaction of A2A adenosine and D2
dopamine receptors modulates corticostriatal glutamatergic transmission.
Neuropharmacology, *53*(6), 783-789.
https://doi.org/10.1016/j.neuropharm.2007.08.006

Trapold, M.A., Overmier. J. B. (1972). The second learning process in instrumental
conditioning. *Current research and theory*, *2*, 427-452.
https://cir.nii.ac.jp/crid/1571417125145479296

Vogel, V., Dittrich, M., Horndasch, S., Kratz, O., Moll, G. H., Erim, Y., Paslakis, G.,
Rauh, E., & Steins-Loeber, S. (2019). Pavlovian-to-instrumental-transfer in
Anorexia Nervosa: A pilot-study on conditioned learning and instrumental
responding to low- and high-calorie food stimuli. *European Journal of*
Neuroscience. https://doi.org/10.1111/ejn.14592

Watson, P., Wiers, R. W., Hommel, B., & de Wit, S. (2014). Working for food you
don't desire. Cues interfere with goal-directed food-seeking. *Appetite*, *79*, 139-
148. https://doi.org/10.1016/j.appet.2014.04.005

Zahm, D. S., & Heimer, L. (1990). Two transpallidal pathways originating in the rat
nucleus   accumbens.   *Journal*   *of*   *comparative*   *Neurology*.
https://doi.org/10.1002/cne.903020302

Figures and corresponding legends

Figure 1 – Anterograde tracing in D1-Cre and A2a-Cre rats

A. D1-Cre or A2a-Cre rats were unilaterally infused in the dorsal striatum (DS) or
nucleus accumbens shell (NAc-S) with DIO-eYFP. B. Sagittal micrographs obtained

in D1-Cre (top) and A2A-Cre (bottom) rats following viral infusion in the DS. C. DIO-
eYFP infusion in the DS of D1-Cre rats. Micrographs show eYFP expression in the
DS, D32 staining and co-labelling (eYFP + D32) in the DS. They also show that DS
D1-SPNs project to the substantia nigra pars reticulata (SNr) and the globus pallidus
externus (GPe). Viral expression was restricted to putative SPNs (D32 | eYFP), with
~40% of SPNs expressing eYFP (eYFP | D32). D. DIO-eYFP infusion in the DS of
A2a-Cre rats. Micrographs show eYFP expression in the DS, D32 staining and co-
labelling (eYFP + D32) in the DS. They also show that DS D2-SPNs project to the
SNr but not the GPe. Viral expression was restricted to putative SPNs (D32 | eYFP),
with ~40% of SPNs expressing eYFP (eYFP | D32). E. DIO-eYFP infusion in the
NAc-S of D1-Cre rats. Micrographs show eYFP expression in the NAc-S, D32
staining and co-labelling (eYFP + D32) in the DS. They also show that NAc-S D1-
SPNs project to the ventral pallidum (VP) and the lateral hypothalamus (LH). Viral
expression was restricted to putative SPNs (D32 | eYFP), with ~41% of SPNs
expressing eYFP (eYFP | D32). F. DIO-eYFP infusion in the NAc-S of A2a-Cre rats.
Micrographs show eYFP expression in the NAc-S, D32 staining and co-labelling
(eYFP + D32) in the DS. They also show that NAc-S D2-SPNs project to the VP but
not the LH. Viral expression was restricted to putative SPNs (D32 | eYFP), with
~40% of SPNs expressing eYFP (eYFP | D32).

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

Figure 2 – Ex-vivo cell recordings in D1-Cre and A2a-Cre rats

A. D1-Cre were bilaterally infused in the NAc-S with DIO-eYFP (black) or DIO-eNpHR3.0 (blue). The representative raw traces of cell attached recordings are those of transfected neurons that were depolarized to elicit action potentials by injecting a brief positive current step (+150pA, 200ms duration, 0.5Hz). 625nm LED illumination (orange bar, continuous wave, 2mW) had no effect in eYFP transfected neurons (black) but it inhibited action potential in eNpHR3.0 transfected neurons (blue). The grouped data for recordings include 5 animals (overlapping data points).

B. A2a-Cre were bilaterally infused in the NAc-S with DIO-eYFP (black) or DIO-eNpHR3.0 (red). The representative raw traces of cell attached recordings are those

1023 of transfected neurons that were depolarized to elicit action potentials by injecting a

1024 brief positive current step (+150pA, 200ms duration, 0.5Hz). 625nm LED illumination

1025 (orange bar, continuous wave, 2mW) had no effect in eYFP transfected neurons

1026 (black) but it inhibited action potential in eNpHR3.0 transfected neurons (red). The

1027 grouped data for recordings include 7 animals (overlapping data points).

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1050

Figure 3 – NAc-S D1-SPNs mediate outcome-specific PIT

A. D1-Cre rats were bilaterally infused in the NAc-S with DIO-eYFP (black) or DIO-eNpHR3.0 (blue). Fiber-optic cannulas were implanted above the NAc-S to provide 625nm LED illumination B. Schematic representation of the behavioral design; S1 and S2: noise and clicker stimuli (counterbalanced); O1 and O2: grain pellets and sucrose solution (counterbalanced); A1 and A2: left and right lever press (counterbalanced). At test, S1 and S2 were presented 4 times each, in a pseudorandom order. Half of the trials for each stimulus was conducted under 625nm LED illumination (ON; continuous wave; ~10mW) whereas the LED remained inactivated during the other half of the trials (OFF). ON/OFF trials were counterbalanced. C. Outcome-specific PIT test: net lever presses when the stimuli predicted the same outcome as the action (Same) or when the stimuli predicted the different outcome (Different). Lever presses are shown for each group in trials conducted under 625nm LED illumination (ON) and in trials without illumination (OFF). Data are shown as mean ± SEM. Panel C includes individual data points for female (filled circle) and male (open circle) rats. Asterisks denote significant effect (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; *n.s.*, nonsignificant).

1068

Figure 4 – NAc-S D2-SPNs mediate for outcome-specific PIT

A. A2a-Cre rats were bilaterally infused in the NAc-S with DIO-eYFP (black) or DIO-
eNpHR3.0 (red). Fiber-optic cannulas were implanted above the NAc-S to provide
625nm LED illumination. B. Outcome-specific PIT test: net lever presses when the
stimuli predicted the same outcome as the action (Same) or when the stimuli
predicted the different outcome (Different). Lever presses are shown for each group
in trials conducted under 625nm LED illumination (ON) and in trials without
illumination (OFF). Data are shown as mean ± SEM. Panel B includes individual data
points for female (filled circle) and male (open circle) rats. Asterisks denote
significant effect (* $p < 0.05$; ** $p < 0.01$; *n.s.*, nonsignificant).

1081

1082

1083

1084

1085

1086

1087

1088

1089

Figure 5 – NAc-S D1-SPNs projections to the VP mediate outcome-specific PIT

A. D1-Cre rats were bilaterally infused in the NAc-S with DIO-eYFP (black) or DIO-eNpHR3.0 (blue). Fiber-optic cannulas were implanted above the VP to provide 625nm LED illumination B. Outcome-specific PIT test: net lever presses when the stimuli predicted the same outcome as the action (Same) or when the stimuli predicted the different outcome (Different). Lever presses are shown for each group in trials conducted under 625nm LED illumination (ON) and in trials without illumination (OFF). Data are shown as mean ± SEM. Panel B includes individual data points for female (filled circle) and male (open circle) rats. Asterisks denote significant effect (** *p* < 0.01; *** *p* < 0.01; *n.s.*, nonsignificant).

1110

Figure 6 – NAc-S D2-SPNs projections to the VP do not mediate outcome-specific PIT

A. A2a-Cre rats were bilaterally infused in the NAc-S with DIO-eYFP (black) or DIO-eNpHR3.0 (red). Fiber-optic cannulas were implanted above the VP to provide 625nm LED illumination. B. Outcome-specific PIT test: net lever presses when the stimuli predicted the same outcome as the action (Same) or when the stimuli predicted the different outcome (Different). Lever presses are shown for each group in trials conducted under 625nm LED illumination (ON) and in trials without illumination (OFF). Data are shown as mean ± SEM. Panel B includes individual data points for female (filled circle) and male (open circle) rats.

1121

1122

1123

1124

1125

1126

1127

1128