Local adaptation in climate tolerance at a small geographic scale contrasts with broad latitudinal
patterns

Greg M. Walter[1,2]*, Avishikta Chakraborty[1,3], Fiona E. Cockerell[1], Vanessa Kellermann[1,4], Lesley A.
Alton[1], Craig R. White[1], Matthew D. Hall[1], Carla M. Sgrò[1]

[1] *School of Biological Sciences, Monash University, Melbourne 3800.*
[2] *School of Natural Sciences, University of Tasmania, Hobart 7001.*
[3] *Department of Genetics, Evolution & Environment, University College London, London WC1E 6BT.*
[4] *Department of Environment and Genetics, La Trobe University, Melbourne 3086.*

* Corresponding Author: Greg M. Walter
10   Email: greg.walter@utas.edu.au; Phone: +614 15 246 846
11   Address: Life Sciences Building, University of Tasmania,
12   Churchill Ave, Sandy Bay, Hobart 7005 Australia

Abstract

If environmental heterogeneity at small geographic scales has a similar capacity to generate adaptive
genetic variation to that found across broad latitudinal gradients, then we currently underestimate
adaptive capacity to rapid global change. To contrast adaptive genetic variation across geography, we
quantified population variation in tolerance traits and their thermal plasticity for five populations of
Drosophila melanogaster from a 3000-km latitudinal gradient across east coast Australia, with eight
populations from an environmentally heterogeneous 600×300km area within Victoria. Surprisingly,
population variation in stress tolerance within Victoria was comparable to that across latitude. Consistent
with local adaptation, heat and desiccation tolerance were greater for populations from warmer and drier
environments, and populations from more predictable environments showed greater thermal plasticity.
Adaptation to environmental heterogeneity at smaller geographic scales can therefore contribute more to
adaptive capacity than originally thought. However, patterns of adaptation often changed across
geography and treatments, which makes predicting responses to global change challenging.

Keywords: desiccation, *Drosophila*, geographic variation, heat tolerance, heterogeneous landscape,
intraspecific variation, latitudinal cline, local adaptation, plasticity, thermal tolerance

Introduction

Ongoing global change is creating increasingly stressful and unpredictable environments, which is fragmenting available habitat for mobile species, and increasing the vulnerability of long-lived species and those with low dispersal (Parmesan & Yohe 2003; Parmesan 2006). It is therefore likely that many organisms will need to adapt *in situ* if they are to persist. Rapid adaptation to global change will be more likely if gene flow readily moves adaptive genetic variation across an environmentally heterogeneous landscape (Sexton *et al.* 2011; Sgrò *et al.* 2011; Willi *et al.* 2022; Sexton *et al.* 2024). Predicting the vulnerability of species then relies on understanding how much adaptive genetic variation is available and how it is distributed among populations across landscapes (Razgour *et al.* 2019; Urban *et al.* 2024). However, we still know little about the scale at which environmental heterogeneity generates and maintains adaptive genetic variation, which limits our ability to predict the adaptive capacity of species facing rapid global change (Byers 2005; Whitlock 2015; Alvarado *et al.* 2022; Meek *et al.* 2023).

The geographic scale at which adaptation arises across a landscape is rarely considered when assessing vulnerability to global change (Bennett *et al.* 2019). Understanding how environmental heterogeneity generates adaptation across geography reveals how gene flow and local selection pressures interact to determine the extent of adaptive genetic differentiation (Slatkin 1987; García-Ramos & Kirkpatrick 1997; Richardson *et al.* 2014; DeMarche *et al.* 2019; Booker 2024). Such knowledge is crucial for predicting how species will respond to ongoing global change. Local adaptation at smaller geographical scales can facilitate more rapid movement of adaptive alleles across a metapopulation, which increases the adaptive capacity of populations (Meek *et al.* 2023). While patterns of adaptation across broad environmental gradients are common (e.g., Huey *et al.* 2000; Etterson 2004; Ren *et al.* 2020; Aspinwall *et al.* 2023), we critically lack tests of whether these same patterns of adaptation emerge at small geographic scales.

Studies of climatic adaptation typically sample broadly across latitude covering thousands of kilometres. Latitudinal clines in stress tolerance traits reflect adaptive genetic variation because genetic differences are shaped by local climatic selection that changes along the latitudinal gradient (Huey *et al.* 2000; Hoffmann *et al.* 2002; Etterson 2004; Trotta *et al.* 2006; Sgrò *et al.* 2010; Lasne *et al.* 2018; Lush *et al.* 2024). However, if local adaptation occurs at smaller geographic scales (e.g., tens or hundreds of kilometres), then adaptive genetic variation could be generated and maintained at scales where gene flow is more likely to move adaptive alleles among populations (Stotz *et al.* 2021; Alvarado *et al.* 2022). While the idea that environmental heterogeneity can generate adaptive genetic variation at different geographic scales has been around for some time (e.g., Cowling & Pressey 2001; Pressey *et al.* 2007; Sgrò *et al.* 2011), few studies have empirically contrasted the extent of adaptation across geographic scales.

Empirical tests of adaptation at multiple geographic scales across a heterogenous landscape are scarce. Hoffmann *et al.* (2001) found little evidence of variation among populations of *Drosophila* c.20–100km apart compared to latitudinal extremes (>2000km). By contrast, subtle differentiation in genes across

distinct tidal environments at a local scale (<100km) were associated with temperature in a sea urchin despite little population structure across latitude (Rumberger *et al.* 2025). Moreover, local adaptation across elevation reveals the potential for adaptation across an ecological gradient (Gonzalo-Turpin & Hazard 2009; de Villemereuil *et al.* 2018; Halbritter *et al.* 2018; Tregenza *et al.* 2021). While these studies show that local environmental differences can generate adaptive differentiation, we still do not understand how adaptive capacity differs for populations distributed across environments at a local versus broad geographic scale. We therefore need studies that directly contrast climate adaptation along a broad gradient with populations across a small geographic scale where local adaptation could still arise.

In addition to adaptation, plastic changes in climate tolerance traits are also expected to play an important role in helping populations buffer global change (Reed *et al.* 2011; Gunderson & Stillman 2015; Magozzi & Calosi 2015). Environments that are variable and predictable should favour the evolution of greater plasticity (Scheiner 1993; Leung *et al.* 2020), which can lead to populations that differ in plasticity when they adapt to different environments (Lind & Johansson 2007; Murren *et al.* 2014; Matesanz & Ramírez-Valiente 2019; Walter *et al.* 2022). While populations sampled across latitude vary in plasticity for thermal traits that reflect climatic selection for plasticity (Sgrò *et al.* 2010; van Heerwaarden & Sgrò 2017), we still lack the ability to connect environmental predictability with the evolution of adaptive plasticity across a heterogeneous landscape.

To contrast the extent of adaptive genetic variation in key traits across geographic scales, we focus on geographic variation in populations of the vinegar fly, *Drosophila melanogaster,* in Australia. Since its introduction to the east coast of Australia, c.100 years ago, *D. melanogaster* has developed clines across c.3000km for numerous traits, including thermal tolerance, desiccation and starvation tolerance and body size (reviewed in Hoffmann & Weeks 2007; Lasne *et al.* 2018). Molecular data suggests that population structure is weak along latitude due to some level of gene flow at distances up to 1500km (Gockel *et al.* 2001; Kennington *et al.* 2003). The presence of latitudinal clines in ecologically important traits provides strong evidence that adaptive genetic variation is generated across latitude (Huey *et al.* 2000; Hoffmann *et al.* 2002). Such broad patterns of adaptation and the potential for high gene flow means that *D. melanogaster* provides a powerful conservative test of whether small-scale geographic variation can generate and maintain adaptive genetic variation.

We sampled five populations of *D. melanogaster* across 3000-km in latitude along east coast Australia (18°S–43°S), which we compared to eight populations sampled from a variety of climates within a 600km×300km area in Victoria (**Fig. 1a**). Compared to the latitudinal populations, locations sampled in Victoria showed greater variation in mean temperature, both across days (**Fig. 1b**) and within days (**Fig. 1c**). To ensure our estimates of among-population variation would be conservative, we sampled flies in Autumn, which is after the summer peak population size that has the greatest potential for homogenization of populations via dispersal (Shpak *et al.* 2010; Behrman *et al.* 2015). Using a

manipulative laboratory experiment, we reared larvae from all 13 populations under three temperature

treatments and measured wing size and tolerance traits (heat, cold and desiccation) on emergent adults.

The data for the latitudinal populations were used previously to show that latitudinal clines changed more

across temperature treatments than in response to a low-calorie diet (Walter *et al.* 2025). Here, we test

three hypotheses to contrast the extent of adaptive differentiation in climate tolerance across geography:

(I) We would detect variation among populations in Victoria for climate tolerance traits, but populations

from across latitude would vary to a greater extent. (II) If the among-population variation is adaptive, we

would observe associations between climate of origin and tolerance traits for both latitudinal and

Victorian populations. (III) If adaptation generates differences in plasticity, then population variation in

plasticity would be associated with environmental predictability.

Methods

Collection and maintenance of experimental populations

In March–May 2021, we collected 20–50 field-inseminated female *D. melanogaster* from thirteen

locations in Australia using sweep nets over fruit waste. For the five latitudinal populations, we sampled

flies at fruit farms in Tasmania, New South Wales, Victoria and Northern Queensland (**Fig. 1a**; **Table**

S1). In Victoria, we sampled flies from discarded grape skins at eight wineries from a variety of climates

(**Fig. 1a**; **Table S1**). We chose wineries due to the abundance of breeding habitat at the same time of year,

allowing us to sample broadly across the state within a short period.

In the lab, we generated isofemale lines by allowing inseminated females to individually lay for 4 days in

40mL vials containing 7mL of standard yeast-potato-dextrose medium (Brewer's yeast 36.4g/L; potato

flakes 18.2g/L; dextrose 27.3g/L; agar 6.4g/L, nipagin [10% w/v in ethanol] 10.9mL/L; and propionic

acid 4.6mL/L), and leaving the larvae to develop at 25°C on a 12:12 hour light:dark cycle. In the

following generation, we checked species identification to prevent contamination from the

morphologically similar *D. simulans*. In the third generation, we treated each isofemale line with

0.3mg/mL of tetracycline in the food to remove differences among populations in their endosymbionts,

particularly *Wolbachia*, which is more prevalent in tropical populations. After three generations, for each

population we combined five males with five virgin females from each isofemale line (19–60 lines per

population, mean=44.6; **Table S1**) into a mass bred population that we maintained in five 300mL bottles

containing 62.5mL of standard food. Each generation, we collected adults and kept them in bottles until

they reached peak fertility (c.4 days), after which they were allowed to lay 250–300 eggs in each of the

five bottles. After laying, we removed the adults, added cards for pupation and left the larvae to develop.

We maintained the 13 populations at 25°C for 12 generations (c.9 months) to ensure that genotypes were

sufficiently mixed, and to avoid conducting experiments during COVID-19 lockdowns in Melbourne.

Manipulative experiment

In January 2022, we conducted a manipulative laboratory experiment with the 13 mass-bred populations

to test for variation among populations in cold, hot and desiccation tolerance in response to temperature

treatments. We picked eggs from all 13 populations into three temperatures (13, 25 and 29°C). 25°C

represents control conditions at which the populations were maintained, and 13 and 29°C represent the

cold and hot margins that larvae in natural populations are often exposed to in winter and summer,

respectively. For each population, we placed c.200 adult flies in each of two laying pots with a thin

(c.3mm) layer of food with double agar and let them lay for 6 hours. We then transferred 20 eggs into

each vial (n=36–42 vials/population/treatment; **Table S2**) containing 7mL of standard food and left larvae

to develop at their treatment temperatures. To manage the replication, we split the experiment into two

blocks so that adults emerged 21–28 January (block 1) and 1–8 March (block 2).

For each block, we conducted 2–3 rounds of egg-picks for each treatment split over 4–5 days to maximise

the overlap in eclosion across treatments (n=14–21 vials/population/treatment/block). We timed the egg

picks for the different treatments according to pilot experiments so that adult flies would emerge

simultaneously. Overall, we used 1456 vials in total (n=455–546 vials per treatment).

During eclosion, adults from each treatment were collected in separate vials each day so that flies would

be the same age across treatments. Flies were then kept at their experimental treatments until they reached

maturity and had mated. Pilot experiments suggested flies kept at 13°C would not mature, and so they

were kept at 16°C to mature in time for the experiment.

Stress tolerance assays and body size measurements

We measured females as their body size and stress tolerance are more sensitive to variation in nutrition

and temperature (Kutz *et al.* 2019; Chakraborty *et al.* 2020), and because sex-specific adaptation was not

our focus. We collected females under light CO_2 anaesthesia, let them recover for 48 hours and then used

established methods to measure heat, cold, or desiccation tolerance, or body size (Hoffmann *et al.* 2002;

Lasne *et al.* 2018). We measured 30 females per assay. To assay heat tolerance, we measured heat

knockdown by putting individual females into sealed 5mL glass vials, submerging them in a 39°C water

bath and measuring the time for each female to cease moving. To assay cold tolerance, we measured

chill-coma recovery by placing individual females into sealed 1.5mL plastic tubes that we submerged in a

0°C water bath for 4 hours and measured the time they took to regain movement. To measure desiccation

tolerance, we put individual females in unsealed 5mL vials placed in sealed glass tanks containing silica

gel that maintained relative humidity at <5%. We then recorded the time it took for each fly to cease

moving. To measure wing size, we removed the right wing of 30 females, mounted them on a microscope

slide and photographed them using a Leica M60 stereo microscope (Leica, Heerbrugg, Switzerland). We

then used *imageJ* to quantify 10 vein intersection landmarks (**Fig. S1**), which we used to estimate

centroid size.

Statistical analyses

175 We used *R* (v.4.3.2) statistical software for all analyses (R Core Team 2024). With *glmmTMB* (Brooks *et*

176 *al.* 2017), we used linear mixed models to test how each trait varied across the 13 populations, and how

177 such variation was associated with the environment of origin. We checked model assumptions using

178 *DHARMa* (Hartig 2022).

Hypothesis I: Testing for among-population variation in tolerance traits and body size

To test for significant variation among populations in response to treatment, we included population and

temperature as fixed effects, and tested for a significant population×temperature interaction using type-III

ANOVA. We performed the analysis on latitudinal and Victorian populations separately. Replicate run

nested within block was the only random effect.

To compare differences in plasticity, we extracted the population marginal means from the models above

using *emmeans* (Lenth 2019), and then calculated plasticity using: $plasticity =$

$\frac{(treatment\ mean - control\ mean)}{control\ mean}$ (Valladares *et al.* 2006), which represents plasticity as the proportional

change in phenotype from control conditions (25°C).

Hypotheses II: Testing whether among-population variation is adaptive

We tested for associations between climate of origin and stress tolerance traits. For all 13 sampling

locations, we first extracted fine-scale rainfall (daily) and temperature (hourly) data using *Nichemapr*

(Kearney & Porter 2020) for the four months preceding population sampling. We calculated daily mean

temperature and then quantified predictability of temperature by calculating autocorrelation across days

using the *acf* function. We then tested whether climate tolerance traits and wing size were associated with

environment of origin using the linear mixed model

$$y_{ijklm} = O \times G_i \times T_j + p_{k(ij)} + r_l + e_{m(ijkl)} \qquad (1)$$

where the three-way interaction of the fixed effects ($O \times G_i \times T_j$) includes environment of origin (O)

measured at the population locations (continuous variable), the geographical comparison (G_i; Victoria vs

latitude) and treatment temperature (T_j). Random effects included the *k*th population nested within

treatment temperature and geography ($p_{k(ij)}$) to avoid pseudoreplication, and the *l*th replicate run within

experimental block (r_l). $e_{m(ijkl)}$ represents the residual, and y_{ijklm} the response variable. We applied

equation 1 to each tolerance trait and body size separately. We included mean temperature as the

environmental covariate (O) for analysing thermal tolerance and body size, and mean daily rainfall when

analysing desiccation tolerance.

A significant slope for O (effect of environment) provides evidence that environment of origin is

associated with that particular trait. Where the two-way interaction between environment of origin and

geography is significant suggests that latitudinal and Victorian populations differ in their association

between tolerance traits (or body size) and environment of origin, and therefore show different patterns of

adaptation. A significant three-way interaction then provides evidence that associations between

environment of origin and the tolerance trait (or body size) is different for latitudinal and Victorian

populations, and this also depends on the treatment temperature.

Finally, to test whether environmental predictability was associated with population variation in thermal

plasticity, we applied equation 1 but used estimates of plasticity in heat tolerance as the response variable,

and autocorrelation of temperature across days (capturing predictability of temperature) as the

environmental covariate.

Results

Hypothesis I: High variation among Victorian populations for tolerance traits

Temperature created significant changes in stress tolerance and body size (**Fig. 2**, **Table 1**). As expected,

flies exposed to colder temperatures were larger, more cold tolerant and less heat tolerant. We found

significant variation among populations in all tolerance traits and body size (**Table 1**). Body size was the

only trait where variation among populations was much greater across latitude compared to Victoria (**Fig.**

S2), which was consistent across all treatments (**Fig. 2**). Somewhat surprisingly, this meant that Victorian

populations showed similar levels of variation among populations compared to the latitudinal populations

for all tolerance traits (**Fig. 2**, **Fig. S2**). These results suggest that environmental heterogeneity at smaller

geographic scales can generate substantial among-population variation in tolerance traits.

Statistically significant interactions between population and temperature quantifies how populations vary

in response to treatments, and therefore reflects population variation in plasticity. For cold and heat

tolerance, both Victorian and latitudinal populations showed significant temperature×population

interactions (**Table 1a-b**). Only desiccation tolerance showed no significant population×temperature

interactions (**Table 1c**). For wing size, only latitudinal populations showed a significant

population×temperature interaction (**Table 1d**). Therefore, Victorian and latitudinal populations both

displayed significant variation in thermal plasticity (**Fig. 3**).

Hypothesis II: Population variation is associated with climate of origin

Along the east coast, temperatures were higher and more consistent at lower latitudes. By contrast,

Victorian sampling locations had greater variation both in temperature across days (**Fig. 1b**) and within

each day (**Fig. 1c**). The Victorian locations therefore experience greater thermal variation compared to

latitudinal locations. Rainfall was lower at high latitudes, ranging 193–984mm south–north, while

Victoria varied between 130–428mm (**Table S1**). To test whether among-population variation in

tolerance traits and body size was adaptive, we quantified associations between climate of origin with the

traits measured in the laboratory. Below we summarise the results for each trait.

<u>Cold tolerance:</u> We found a significant three-way interaction between temperature of origin (O),

geography (G; latitude vs Victoria) and treatment temperature (T; **Fig. 4a**; **Table S3**). While there was no

association between temperature of origin and cold tolerance at 13°C, at the warmer temperatures (25°C

and 29°C), latitudinal populations from warmer temperatures showed slower cold recovery (reduced cold

tolerance), as expected (**Fig. 4a**). By contrast, Victorian populations showed the opposite trend where

populations from warmer environments showed higher cold tolerance (**Fig. 4a**), suggesting that they were

better at coping with a cold shock after acclimating to warmer conditions.

<u>Heat tolerance:</u> We found no significant effect of geography, suggesting that associations between

temperature and heat tolerance were consistent for Victorian and latitudinal populations (**Fig. 4b; Table**

S3). Associations between temperature of origin and heat tolerance changed depending on the treatment

temperature, which was supported by a significant temperature of origin × treatment temperature

interaction (**Fig. 4b; Table S3**). While we found no strong associations at 13°C, at the warmer (25°C and

29°C) treatments, populations from warmer locations showed higher heat tolerance, creating a positive

association between temperature of origin and heat tolerance (**Fig. 4b**). We therefore found evidence of

local adaptation in warmer treatments, which was consistent for Victorian and latitudinal populations.

<u>Wing size:</u> Consistent with latitudinal studies, populations from lower latitudes had smaller wings (**Fig.**

4c). However, Victorian populations showed a contrasting trend where populations from warmer

environments had slightly larger wings (**Fig. 4c**). This result was supported by a significant temperature

of origin × geography (latitudinal vs Victoria) interaction (**Fig. 4c**; **Table S3**). Wing size therefore

showed contrasting associations with temperature of origin for Victorian versus latitudinal populations,

and this was consistent across treatment temperatures.

<u>Desiccation tolerance:</u> We found a significant interaction between rainfall of origin × geography (**Fig. 4d**;

Table S3). This meant that latitudinal populations showed no association between rainfall and desiccation

tolerance, whereas Victorian populations from drier environments showed greater desiccation tolerance

(**Fig. 4d**). This result was consistent across treatment temperatures. Victorian populations from drier

environments therefore showed greater desiccation tolerance, suggesting local adaptation to water

availability.

To ensure that body size did not affect the association between climate of origin and each tolerance trait,

we conducted the same analyses, but we used population means and included wing size as a covariate.

Including wing size in the analysis did not change the strength of the association between climate of

origin and tolerance traits, suggesting the patterns observed were not due mainly to variation in body size.

Hypothesis III: Heat tolerance plasticity is linked to environmental predictability

To test for an association between environmental predictability and plasticity, we focussed on heat

tolerance as it showed larger variation in plasticity among populations compared to other traits (**Fig. 3**)

and the most consistent association with climate (**Fig. 4**). Greater autocorrelation (i.e., predictability) of

temperature across days was associated with greater plasticity for the Victorian populations, but less

plasticity along latitude (**Fig. 5**). This was supported by a significant interaction between autocorrelation

of mean temperature and geography (latitudinal vs Victorian populations) (**Table S4**). Victorian

populations from more predictable thermal environments therefore showed greater increases in heat

tolerance when they were exposed to the warmest treatment temperature (**Fig. 5**). We found the same

positive trend at 13°C, but because plasticity reduced heat tolerance at the cooler treatment, populations

from more predictable environments showed a smaller reduction in heat tolerance. These results suggest

that Victorian populations (but not latitudinal populations) from more predictable environments showed

adaptive thermal plasticity.

Discussion

Using populations of *Drosophila melanogaster*, we contrasted population variation in tolerance traits and

body size across latitude along east-coast Australia with population variation at a smaller geographical

scale within Victoria. Not only was among-population variation in Victoria significant for all traits, but

the variation was similar in extent to that seen across latitude (*Hypothesis I*: **Figs. 2–3**). This among-

population variation in tolerance traits was associated with the climate of origin, which provides evidence

for local adaptation across Victorian populations (*Hypothesis II*: **Fig. 4**). Plasticity in heat tolerance was

associated with temperature predictability, but only for Victorian populations (*Hypothesis III*: **Fig. 5**).

Together, our results suggest that environmental heterogeneity at a relatively small geographic scale can

generate local adaptation that mirrors the extent of adaptive genetic variation across 3000-km in latitude.

Local-scale environmental heterogeneity can therefore generate adaptive genetic variation that increases

the adaptive capacity of the metapopulation. However, we also showed that latitudinal and Victorian

populations often showed contrasting patterns of adaptation, and that patterns of adaptation changed with

treatment temperature (**Figs. 4–5**). Accurately predicting population responses to global change therefore

needs to consider how adaptation arises across geographic scales.

Contrasting patterns of adaptation across geographic scales

While the presence of latitudinal clines suggest that extensive gene flow could transfer beneficial alleles

to declining populations at the broader geographical context (Meek *et al.* 2023), our results indicate that this process can occur on a much smaller geographic scale. We show that beneficial alleles are likely to segregate among local (and potentially fragmented) populations at a smaller scale than previous anticipated, which increases the adaptive capacity of the metapopulation to global change. Given that gene flow increases at smaller spatial scales (Sexton *et al.* 2024), populations could rely on locally segregating adaptive genetic variation rather than the movement of adaptive alleles across broad geographic scales (Meek *et al.* 2023; Sexton *et al.* 2024). However, our results raise several concerns about predicting population responses to global change based on patterns of adaptation across geography. First, patterns of climate adaptation often disappeared under different temperatures, which highlights the need to understand adaptive potential in response to environmental variation that better represent the breadth of environments created by global change. Second, associations between climate of origin and all traits (except heat tolerance), as well as plasticity in heat tolerance, differed for Victoria populations compared to latitudinal populations. Inferences for natural populations facing climate change are therefore likely to change with geography. While heat tolerance was consistently greater in populations from warmer environments, predicting population responses based on other traits and environmental variables will depend on how adaptation has been shaped across geography.

Segregation of adaptive genetic variation across a heterogeneous landscape
Few studies have directly contrasted adaptation across geographic scales. In lodgepole pines, regional variation accounted for 20% of total genetic variance in key ecological traits, suggesting that environmental heterogeneity maintains genetic variation (Yeaman & Jarvis 2006). At a molecular level, subtle genetic differences in a sea urchin emerged at a local scale (<100km) and were associated with sea temperature, which provides evidence of local adaptive genetic differentiation (Rumberger *et al.* 2025). While Hoffmann *et al.* (2001) found evidence of variation among populations for cold tolerance in *D. melanogaster* at local scales (c.20-100km), they did not test heat tolerance or connect among population variation to the environment of origin. Nadeau and Urban (2024) demonstrated variation in thermal tolerance among *Daphnia* from nearby pools with different thermal conditions, but could not link this variation with environmental differences, suggesting that microgeographic variation did not generate patterns of adaptation. While these studies show that small-scale environmental heterogeneity can generate among-population variation, they struggled to connect such variation with environmental heterogeneity at the appropriate scale, or link adaptive genetic variation to key ecological traits. By contrast, our study focused on a variety of climatic environments from a 300×600km area, which maximised our chance of connecting patterns of local adaptation with environmental heterogeneity despite the homogenising effects of gene flow that occurs regularly within 500km (Kennington *et al.* 2003).

Predictions about adaptation change across scales

Our key result is that our understanding of climatic adaptation changes depending on geographic scale (O'Brien *et al.* 2017). While Victorian populations mimicked broader heat tolerance trends (populations from warmer areas show higher heat tolerance), this was not the case for any other tolerance trait or body size. Therefore, our predictions are consistent for adapting to heat, but become inaccurate when considering plasticity in heat tolerance, or any other trait. This is because Victorian populations showed the opposite patterns to the latitudinal populations for cold and desiccation tolerance, and body size. Therefore, while latitudinal clines are a powerful tool for understanding adaptation to broad climatic gradients, if we scale down the same predictions, they can become wildly inaccurate and misrepresent the relationship between temperature and climatic tolerance traits. Accurately predicting the adaptive potential of natural population facing global change therefore requires understanding how adaptation occurs across geographical scales (O'Brien *et al.* 2017; Sexton *et al.* 2024; Urban *et al.* 2024).

Our results suggest that predicting adaptation to climate change based on latitudinal patterns are likely to be inaccurate at other geographical scales. Along latitude, populations from lower (warmer) latitudes showed higher heat tolerance, lower cold tolerance and smaller body size (**Fig. 4a,b,d**), suggesting that thermal tolerance and body size are selected together across latitude. While Victorian populations from warmer locations had higher heat tolerance, they contrasted with latitudinal populations by showing higher cold tolerance and larger body sizes (**Fig. 4a,d**). Selection for body size across the cline could be due to autocorrelation of environmental variables (e.g., humidity) or genetic factors (e.g., inversions; Rako *et al.* 2006) driving correlated trait responses to latitude. Compared to latitudinal populations, Victorian populations experienced lower humidity and greater temperature fluctuations (**Fig. 1b-c**), which could increase heat tolerance while creating different changes in body size and cold tolerance (Hangartner *et al.* 2020). In support of this, Victorian populations showed greater adaptive genetic variation in desiccation tolerance than latitudinal populations (**Fig. 2d, 4d**). Selection could therefore favour different combinations of traits away from the coast compared to across latitude. Accurately predicting species' responses to climate change then requires understanding how populations from across a species' range, rather than from a subset of environments, respond to environmental variation that includes climate extremes (Matesanz & Ramírez-Valiente 2019; Walter *et al.* 2020; Meek *et al.* 2023; Rumberger *et al.* 2025). Our results also suggest that understanding the drivers of body size, particularly latitudinal trends (i.e., Bergmann's rule), requires further consideration of correlated responses in other traits (e.g., desiccation tolerance) and environmental factors other than temperature.

Temperature-dependent patterns of adaptation

Compared to latitudinal populations, patterns of adaptation changed across treatment temperatures to a greater extent for Victorian populations (**Fig. 4**). The cold treatment tended to reduce variation among populations, whereas the hot temperature increased variation among populations. However, greater

variation at the high temperature often weakened patterns of adaptation in desiccation and heat tolerance,

and created opposing patterns to what was predicted for cold tolerance (populations from warmer

environments were more cold-tolerant; **Fig. 4a**). These results suggest that across a species range, some

populations are likely to be more sensitive to temperature changes than others. This is consistent with

studies showing that patterns of adaptation can weaken under stressful or variable conditions (Booker

2024; Walter *et al.* 2025), which increases vulnerability to climate change (Anderson & Wadgymar

2020). Accurately predicting the response of natural populations to global change therefore requires

considering how locally adapted populations will vary in their response to the particular conditions

created by environmental change.

Selection for plasticity across Victoria but not across latitude

Victorian populations from locations where temperature was more predictable across days showed greater

thermal plasticity (**Fig. 5**). These results support theory and empirical observations that environmental

predictability generates adaptive plasticity (Reed *et al.* 2010; Diaz *et al.* 2020; Leung *et al.* 2020;

Gallegos *et al.* 2024), which helps populations adapt to thermal variation. Local adaptation across a

heterogeneous landscape can therefore generate variation in plasticity, which could determine population

responses to environmental change (Vinton *et al.* 2022). While such adaptive plasticity can help

populations to buffer familiar thermal variation, it could also increase their vulnerability to unpredictable

changes in temperature created by global change (Ashander *et al.* 2016). Furthermore, latitudinal

populations showed no association between thermal predictability and plasticity, suggesting that not all

predictable environments will favour the evolution of adaptive plasticity. Understanding how adaptive

plasticity evolves across a heterogeneous landscape, and then determining the consequences for persisting

under environmental change remains a substantial challenge for predicting population responses to global

change (Meek *et al.* 2023).

Conclusions

We show that adaptive genetic variation that emerges at a relatively small-scale is comparable to that

found across a broad latitudinal gradient. Such local adaptive genetic variation improves the adaptive

potential of species responding to rapid environmental change, but changes in environment, such as

exposure to temperature extremes, can obscure the detection of such adaptive genetic variation. More

broadly, our results support the idea that local adaptation at smaller geographic scales can generate

adaptive genetic variation, which means that adaptive alleles can be harnessed more efficiently than

previously anticipated. Decisions on managing landscapes should then aim to maintain the processes that

generate adaptative variation at multiple geographic scales. By focusing on conserving environmental

heterogeneity, including at local scales, adaptive potential to environmental change can be enhanced.

Acknowledgements

We thank Ary Hoffmann for comments on a previous version. We are grateful to Carmen da Silva, Julian

Beaman and Belinda van Heerwaarden for helping to sample the latitudinal populations. We thank

Andrew Murch for his insights into climatic variation across Victorian wine regions. We also thank Emily

Lombardi, Mia Wansbrough and Sandra Hangartner for their help with some of the assays. GMW was

supported by Australian Research Council fellowships DE200101019 and FT240100466.

References

1.
Alvarado, A.H., Bossu, C.M., Harrigan, R.J., Bay, R.A., Nelson, A.R.P., Smith, T.B. *et al.* (2022). Genotype-environment associations across spatial scales reveal the importance of putative adaptive genetic variation in divergence. *Evolutionary Applications*, 15, 1390-1407.

2.
Anderson, J.T. & Wadgymar, S.M. (2020). Climate change disrupts local adaptation and favours upslope migration. *Ecol. Lett.*, 23, 181-192.

3.
Ashander, J., Chevin, L.M. & Baskett, M.L. (2016). Predicting evolutionary rescue via evolving plasticity in stochastic environments. *Proceedings of the Royal Society B-Biological Sciences*, 283.

4.
Aspinwall, M.J., Blackman, C.J., Maier, C., Tjoelker, M.G., Rymer, P.D., Creek, D. *et al.* (2023). Aridity drives clinal patterns in leaf traits and responsiveness to precipitation in a broadly distributed Australian tree species. *Plant-Environment Interactions*, 4, 70-85.

5.
Behrman, E.L., Watson, S.S., O'Brien, K.R., Heschel, M.S. & Schmidt, P.S. (2015). Seasonal variation in life history traits in two *Drosophila* species. *J. Evol. Biol.*, 28, 1691-1704.

6.
Bennett, S., Duarte, C.M., Marbà, N. & Wernberg, T. (2019). Integrating within-species variation in thermal physiology into climate change ecology. *Philos. Trans. R. Soc. Lond. B.*, 374, 20180550.

7.
Booker, T.R. (2024). The structure of the environment influences the patterns and genetics of local adaptation. *Evolution Letters*, 8, 787-798.

8.
Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C., Nielsen, A. *et al.* (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, 9, 378-400.

9.

Byers, D.L. (2005). Evolution in heterogeneous environments and the potential of maintenance of genetic variation in traits of adaptive significance. *Genetica*, 123, 107-124.

10.

Chakraborty, A., Sgrò, C.M. & Mirth, C.K. (2020). Does local adaptation along a latitudinal cline shape plastic responses to combined thermal and nutritional stress? *Evolution*, 74, 2073-2087.

11.

Cowling, R.M. & Pressey, R.L. (2001). Rapid plant diversification: planning for an evolutionary future. *Proc. Natl. Acad. Sci. USA*, 98, 5452-5457.

12.

de Villemereuil, P., Mouterde, M., Gaggiotti, O.E. & Till-Bottraud, I. (2018). Patterns of phenotypic plasticity and local adaptation in the wide elevation range of the alpine plant *Arabis alpina*. *J. Ecol.*, 106, 1952-1971.

13.

DeMarche, M.L., Doak, D.F. & Morris, W.F. (2019). Incorporating local adaptation into forecasts of species' distribution and abundance under climate change. *Global Change Biol.*, 25, 775-793.

14.

Diaz, F., Kuijper, B., Hoyle, R.B., Talamantes, N., Coleman, J.M. & Matzkin, L.M. (2020). Environmental predictability drives adaptive within- and transgenerational plasticity of heat tolerance across life stages and climatic regions. *Funct. Ecol.*, 35, 153-166.

15.

Etterson, J.R. (2004). Evolutionary potential of *Chamaecrista fasciculata* in relation to climate change. I. Clinal patterns of selection along an environmental gradient in the great plains. *Evolution*, 58, 1446-1458.

16.

Gallegos, C., Chevin, L.M., Hodgins, K.A. & Monro, K. (2024). Predicting adaptation and evolution of plasticity from temporal environmental change. *Methods in Ecology and Evolution*, 16, 84-96.

17.

García-Ramos, G. & Kirkpatrick, M. (1997). Genetic models of adaptation and gene flow in peripheral populations. *Evolution*, 51, 21-28.

18.

Gockel, J., Kennington, W.J., Hoffmann, A., Goldstein, D.B. & Partridge, L. (2001). Nonclinality of molecular variation implicates selection in maintaining a morphological cline of *Drosophila melanogaster*. *Genetics*, 158, 319-323.

19.

Gonzalo-Turpin, H. & Hazard, L. (2009). Local adaptation occurs along altitudinal gradient despite the existence of gene flow in the alpine plant species *Festuca eskia*. *J. Ecol.*, 97, 742-751.

20.

Gunderson, A.R. & Stillman, J.H. (2015). Plasticity in thermal tolerance has limited potential to buffer ectotherms from global warming. *Proceedings of the Royal Society B-Biological Sciences*, 282, 20150401.

21.

Halbritter, A.H., Fior, S., Keller, I., Billeter, R., Edwards, P.J., Holderegger, R. *et al.* (2018). Trait differentiation and adaptation of plants along elevation gradients. *J. Evol. Biol.*, 31, 784-800.

22.
Hangartner, S., Lasne, C., Sgrò, C.M., Connallon, T. & Monro, K. (2020). Genetic covariances promote
climatic adaptation in Australian *Drosophila*. *Evolution*, 74, 326-337.

23.
Hartig, F. (2022). DHARMa: Residual diagnostics for hierarchical (multi-Level / mixed) regression
models. R package.

24.
Hoffmann, A.A., Anderson, A. & Hallas, R. (2002). Opposing clines for high and low temperature
resistance in *Drosophila melanogaster*. *Ecol. Lett.*, 5, 614-618.

25.
Hoffmann, A.A., Hallas, R., Sinclair, C. & Mitrovski, P. (2001). Levels of variation in stress resistance in
Drosophila among strains, local populations, and geographic regions: patterns for desiccation,
starvation, cold resistance, and associated traits. *Evolution*, 55, 1621-1630.

26.
Hoffmann, A.A. & Weeks, A.R. (2007). Climatic selection on genes and traits after a 100 year-old
invasion: a critical look at the temperate-tropical clines in *Drosophila melanogaster* from eastern
Australia. *Genetica*, 129, 133-147.

27.
Huey, R.B., Gilchrist, G.W., Carlson, M.L., Berrigan, D. & Serra, L. (2000). Rapid evolution of a
geographic cline in size in an introduced fly. *Science*, 287, 308-309.

28.
Kearney, M. & Porter, W.P. (2020). NicheMapR – an R package for biophysical modelling: the
ectotherm and Dynamic Energy Budget models. *Ecography*, 43, 85-96.

29.
Kennington, W.J., Gockel, J. & Partridge, L. (2003). Testing for asymmetrical gene flow in a *Drosophila*
melanogaster body-size cline. *Genetics*, 165, 667-673.

30.
Kutz, T.C., Sgrò, C.M. & Mirth, C.K. (2019). Interacting with change: Diet mediates how larvae respond
to their thermal environment. *Funct. Ecol.*, 33, 1940-1951.

31.
Lasne, C., Hangartner, S.B., Connallon, T. & Sgrò, C.M. (2018). Cross-sex genetic correlations and the
evolution of sex-specific local adaptation: Insights from classical trait clines in *Drosophila*
melanogaster. *Evolution*, 72, 1317-1327.

32.
Lenth, R.V. (2019). emmeans: Estimated Marginal Means, aka Least-Squares Means. In: *R Package*
https://CRAN.R-project.org/package=emmeans.

33.
Leung, C., Rescan, M., Grulois, D. & Chevin, L.M. (2020). Reduced phenotypic plasticity evolves in less
predictable environments. *Ecol. Lett.*, 23, 1664-1672.

34.
Lind, M.I. & Johansson, F. (2007). The degree of adaptive phenotypic plasticity is correlated with the
spatial environmental heterogeneity experienced by island populations of *Rana temporaria*. *J.*
Evol. Biol., 20, 1288-1297.

35.
Lush, J., Sgrò, C.M. & Hall, M.D. (2024). Anticipating change: the impact of simulated seasonal
heterogeneity on heat tolerances along a latitudinal cline. *Ecology*.

36.
Magozzi, S. & Calosi, P. (2015). Integrating metabolic performance, thermal tolerance, and plasticity
enables for more accurate predictions on species vulnerability to acute and chronic effects of
global warming. *Global Change Biol.*, 21, 181-194.

37.
Matesanz, S. & Ramírez-Valiente, J.A. (2019). A review and meta‐analysis of intraspecific differences
in phenotypic plasticity: Implications to forecast plant responses to climate change. *Global Ecol.*
Biogeogr., 28, 1682-1694.

38.
Meek, M.H., Beever, E.A., Barbosa, S., Fitzpatrick, S.W., Fletcher, N.K., Mittan-Moreau, C.S. *et al.*
(2023). Understanding local adaptation to prepare populations for climate change. *Bioscience*, 73,
36-47.

39.
Murren, C.J., Maclean, H.J., Diamond, S.E., Steiner, U.K., Heskel, M.A., Handelsman, C.A. *et al.* (2014).
Evolutionary change in continuous reaction norms. *Am. Nat.*, 183, 453-467.

40.
Nadeau, C.P. & Urban, M.C. (2024). Macroecological predictors of evolutionary and plastic potential do
not apply at microgeographic scales for a freshwater cladoceran under climate change. *Evolution*
Letters, 8, 43-55.

41.
O'Brien, E.K., Higgie, M., Reynolds, A., Hoffmann, A.A. & Bridle, J.R. (2017). Testing for local
adaptation and evolutionary potential along altitudinal gradients in rainforest *Drosophila*: beyond
laboratory estimates. *Global Change Biol.*, 23, 1847-1860.

42.
Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Ann. Rev. Ecol.*
Evol. Syst., 37, 637-669.

43.
Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across
natural systems. *Nature*, 421, 37-42.

44.
Pressey, R.L., Cabeza, M., Watts, M.E., Cowling, R.M. & Wilson, K.A. (2007). Conservation planning in
a changing world. *Trends Ecol. Evol.*, 22, 583-592.

45.
R Core Team (2024). R: A language and environment for statistical computing. R Foundation for
Statistical Computing Vienna, Austria.

46.
Rako, L., Anderson, A.R., Sgrò, C.M., Stocker, A.J. & Hoffmann, A.A. (2006). The association between
inversion *In(3R)Payne* and clinally varying traits in *Drosophila melanogaster*. *Genetica*, 128,
373-384.

47.

Razgour, O., Forester, B., Taggart, J.B., Bekaert, M., Juste, J., Ibáñez, C. *et al.* (2019). Considering adaptive genetic variation in climate change vulnerability assessment reduces species range loss projections. *Proc. Natl. Acad. Sci. USA*, 116, 10418-10423.

48.
Reed, T.E., Schindler, D.E. & Waples, R.S. (2011). Interacting effects of phenotypic plasticity and evolution on population persistence in a changing climate. *Conserv. Biol.*, 25, 56-63.

49.
Reed, T.E., Waples, R.S., Schindler, D.E., Hard, J.J. & Kinnison, M.T. (2010). Phenotypic plasticity and population viability: the importance of environmental predictability. *Proc. R. Soc. Lond. B.*, 277, 3391-3400.

50.
Ren, L., Guo, X., Liu, S., Yu, T., Guo, W., Wang, R. *et al.* (2020). Intraspecific variation in *Phragmites australis*: Clinal adaption of functional traits and phenotypic plasticity vary with latitude of origin. *J. Ecol.*, 108, 2531-2543.

51.
Richardson, J.L., Urban, M.C., Bolnick, D.I. & Skelly, D.K. (2014). Microgeographic adaptation and the spatial scale of evolution. *Trends Ecol. Evol.*, 29, 165-176.

52.
Rumberger, C., Armstrong, M., Kim, M., Ponce, R., Melendez, J., DeBiasse, M. *et al.* (2025). Selection Over Small and Large Spatial Scales in the Face of High Gene Flow. *Mol. Ecol.*, 34, e17700.

53.
Scheiner, S.M. (1993). Genetics and Evolution of Phenotypic Plasticity. *Annu. Rev. Ecol. Syst.*, 24, 35-68.

54.
Sexton, J.P., Clemens, M., Bell, N., Hall, J., Fyfe, V. & Hoffmann, A.A. (2024). Patterns and effects of gene flow on adaptation across spatial scales: implications for management. *J. Evol. Biol.*, 37, 732-745.

55.
Sexton, J.P., Strauss, S.Y. & Rice, K.J. (2011). Gene flow increases fitness at the warm edge of a species' range. *Proc. Natl. Acad. Sci. USA*, 108, 11704-11709.

56.
Sgrò, C.M., Lowe, A.J. & Hoffmann, A.A. (2011). Building evolutionary resilience for conserving biodiversity under climate change. *Evolutionary Applications*, 4, 326-337.

57.
Sgrò, C.M., Overgaard, J., Kristensen, T.N., Mitchell, K.A., Cockerell, F.E. & Hoffmann, A.A. (2010). A comprehensive assessment of geographic variation in heat tolerance and hardening capacity in populations of *Drosophila melanogaster* from eastern Australia. *J. Evol. Biol.*, 23, 2484-2493.

58.
Shpak, M., Wakeley, J., Garrigan, D. & Lewontin, R.C. (2010). A structured coalescent process for seasonally fluctuating populations. *Evolution*, 64, 1395-1409.

59.
Slatkin, M. (1987). Gene flow and the geographic structure of natural populations. *Science*, 236, 787-792.

60.

Stotz, G.C., Salgado-Luarte, C., Escobedo, V.M., Valladares, F. & Gianoli, E. (2021). Global trends in phenotypic plasticity of plants. *Ecol. Lett.*, 24, 2267-2281.

61.

Tregenza, T., Rodriguez-Muñoz, R., Boonekamp, J.J., Hopwood, P.E., Sørensen, J.G., Bechsgaard, J. *et al.* (2021). Evidence for genetic isolation and local adaptation in the field cricket *Gryllus campestris*. *J. Evol. Biol.*, 34, 1624-1636.

62.

Trotta, V., Calboli, F.C., Ziosi, M., Guerra, D., Pezzoli, M.C., David, J.R. *et al.* (2006). Thermal plasticity in *Drosophila melanogaster*: a comparison of geographic populations. *BMC Evol. Biol.*, 6, 67.

63.

Urban, M.C., Swaegers, J., Stoks, R., Snook, R.R., Otto, S.P., Noble, D.W.A. *et al.* (2024). When and how can we predict adaptive responses to climate change? *Evolution Letters*, 8, 172-187.

64.

Valladares, F., Sanchez-Gomez, D. & Zavala, M.A. (2006). Quantitative estimation of phenotypic plasticity: bridging the gap between the evolutionary concept and its ecological applications. *J. Ecol.*, 94, 1103-1116.

65.

van Heerwaarden, B. & Sgrò, C.M. (2017). The quantitative genetic basis of clinal divergence in phenotypic plasticity. *Evolution*, 71, 2618-2633.

66.

Vinton, A.C., Gascoigne, S.J.L., Sepil, I. & Salguero-Gómez, R. (2022). Plasticity's role in adaptive evolution depends on environmental change components. *Trends Ecol. Evol.*, 37, 1067-1078.

67.

Walter, G.M., Catara, S., Bridle, J.R. & Cristaudo, A. (2020). Population variation in early development can determine ecological resilience in response to environmental change. *New Phytol.*, 226, 1312-1324.

68.

Walter, G.M., Chakraborty, A., Cockerell, F.E., Kellermann, V., Hall, M.D., White, C.R. *et al.* (2025). Variation in temperature but not diet determines the stability of latitudinal clines in tolerance traits and their plasticity. *bioRxiv*.

69.

Walter, G.M., Clark, J., Cristaudo, A., Nevado, B., Catara, S., Paunov, M. *et al.* (2022). Adaptive divergence generates distinct plastic responses in two closely related *Senecio* species. *Evolution*, 76, 1229-1245.

70.

Whitlock, M.C. (2015). Modern Approaches to Local Adaptation. *Am. Nat.*, 186, S1-4.

71.

Willi, Y., Kristensen, T.N., Sgrò, C.M., Weeks, A.R., Orsted, M. & Hoffmann, A.A. (2022). Conservation genetics as a management tool: The five best-supported paradigms to assist the management of threatened species. *Proc. Natl. Acad. Sci. USA*, 119, e2105076119.

72.

Yeaman, S. & Jarvis, A. (2006). Regional heterogeneity and gene flow maintain variance in a quantitative trait within populations of lodgepole pine. *Proc. R. Soc. Lond. B.*, 273, 1587-1593.

Tables & Figures

Fig. 1 (a) Map of sampling locations along east coast Australia (circles; n=5) compared to locations
within Victoria (triangles; n=8). Shading represents the average mean annual temperature, where lighter
shading (warmer colours) depict warmer areas. **(b)** Mean daily temperature for the sampled locations. **(c)**
Daily coefficient of variation in temperature for all sampled locations. Data produced using *NicheMapR*
(Kearney & Porter 2020).

Table 1 ANOVA summaries for latitudinal and Victorian populations for the four traits (**a-d**). Parameters
in bold are significant at p<0.05. Temp = treatment temperature, and Pop = population.

Geography	Parameter	df	(a) Cold χ^2	P	(b) Heat χ^2	P	(c) Desiccation χ^2	P	(d) Wing size χ^2	P
Latitude	Intercept	1	582.7	**<0.001**	1531.6	**<0.001**	7600.0	**<0.001**	627800.2	**<0.001**
	Temp.	2	498.1	**<0.001**	1004.9	**<0.001**	13.9	**<0.001**	2777.4	**<0.001**
	Pop.	4	33.4	**<0.001**	69.2	**<0.001**	12.4	**0.014**	482.6	**<0.001**
	Temp×Pop	8	10.5	0.234	30.4	**<0.001**	6.8	0.558	39.6	**<0.001**
Victoria	Intercept	1	868.9	**<0.001**	1723.5	**<0.001**	10869.8	**<0.001**	940758.3	**<0.001**
	Temp.	2	589.5	**<0.001**	1789.7	**<0.001**	26.1	**<0.001**	4142.1	**<0.001**
	Pop.	7	24.0	**0.001**	73.4	**<0.001**	42.6	**<0.001**	28.0	**<0.001**
	Temp×Pop	14	32.0	**0.004**	45.8	**<0.001**	13.0	0.527	20.1	0.126

676 **Fig. 2** The effect of treatment temperature on stress tolerance and body size for populations along latitude

677 (open circles) and within Victoria (closed circles). We measured **(a)** cold tolerance as chill coma recovery

678 time (mins), **(b)** heat tolerance as heat knockdown time (mins), **(c)** desiccation tolerance (hours), and **(d)**

679 body size as wing centroid size. Points and error bars represent the marginal means (±1SE). Data points

680 are offset for clarity.

Fig. 3 Population variation in thermal plasticity for each trait (**a-d**). Triangles represent Victorian

populations, and circles the latitudinal populations. Horizontal grey lines represent the control conditions

(25°C), and the points represent the plasticity for each population as the change in phenotype from the

control condition. Data points are offset for clarity.

Fig. 4 Association between climate of origin and each trait (**a–d**) after exposure to three temperatures

(different colours). Panels **a–c** compare mean daily temperature to cold tolerance, heat tolerance and wing

size, while panel **d** compares average daily rainfall to desiccation tolerance. Points and error bars

represent the marginal means (±1SE). Significant interactions are presented at the top of each panel,

where T=treatment temperature, O=environment of origin and G=geographical comparison (latitude vs

Victoria). **Table S3** contains the full ANOVA summaries. **Fig. S3** separates out the treatment

temperatures for desiccation tolerance to observe the patterns more easily.

Fig. 5 Association between autocorrelation of mean temperature across days (for the sampled locations),

with plasticity in heat tolerance. Plasticity is calculated as heat tolerance relative to the 25°C control

temperature. Triangles represent Victorian populations, and open circles the latitudinal populations. The

significant O×G interaction is presented, where O=environment of origin and G=geographical

comparison (latitude vs Victoria). This provides evidence that Victorian and latitudinal populations differ

in their relationship between autocorrelation in temperature and thermal plasticity. **Table S4** contains the

full ANOVA summary.