Infanticide in a colonial cooperative bird is not associated with direct reproductive

benefits

Rita Covas [1,2,3], Liliana R. Silva.[1,2], André C. Ferreira[1,2,4,5], Pietro B. D'Amelio[3,5], Rita

Fortuna[1,2,6], Delphine Duval [7], Matthieu Paquet[8], Claire Doutrelant[3,5]

[1]CIBIO-InBio, Centro de Investigação em Biodiversidade e Recursos Genéticos, Laboratório

Associado, University of Porto, Campus Agrário de Vairão, 4485-661 Vairão, Portugal

[2]BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de

Vairão, 4485-661 Vairão, Portugal

[3]FitzPatrick Institute of African Ornithology, University of Cape Town, Rondebosch 7701,

South Africa

[4]Department of Evolutionary Biology and Environmental Science, University of Zurich, Zurich

8057, Switzerland

[5]CEFE CNRS, University of Montpellier, EPHE, IRD, Montpellier, France

[6]Department of Biology, Norwegian University of Science and Technology, Norway

[7]Département de Biologie, Ecole Normale Supérieure de Lyon, Lyon, France

[8]Centre National de la Recherche Scientifique (CNRS), Station d'Écologie Théorique et
Expérimentale, Moulis, France

ABSTRACT

Infanticide, the killing of dependent young by adult conspecifics, is an extreme form of

conflict that remains poorly understood. Previous studies suggest that it is associated with

reproductive benefits, either by making breeders newly available for reproduction or by

increasing offspring care and reducing competition for resources. In addition, infanticide may

result from a reproductive trade-off, where parents kill their offspring if environmental

conditions compromise their reproductive value. Here, we combine video, genetic and

individual-level reproductive data collected over 12 years to describe infanticide in a highly

social, cooperative bird, the sociable weaver *Philetairus socius* and to establish whether

direct reproductive benefits or life-history trade-offs are possible motivators for this

behaviour. We documented 50 cases of infanticide and identified 27 aggressors with known

histories. These were mostly born outside the colonies where they committed infanticide

(63%) and there was some bias towards females and younger individuals. Aggressors were

usually not part of the victim's breeding group (74%), but in two cases infanticide was

committed by the breeding female, and in five by helpers. In most cases, aggressors were not

reproducing at the time of the infanticide, were never recorded mating with any of the

breeders from the attacked nest, nor occupied the nest from where the chicks were evicted.

Only eight of the known aggressors were confirmed as breeders after the attacks, and these

mostly reproduced at the colony where the attacks were committed. Overall, we found no

evidence for direct reproductive benefits of infanticide in sociable weavers. However, the

younger age of aggressors, lack of genetic relatedness to the victims and future breeding at

the colony suggest that competition between breeding groups, and hence indirect

reproductive benefits, could provide motivation for infanticide in this population. Additional

work over longer periods is needed to identify these possible indirect fitness benefits of

infanticide.

Keywords

Aggression; Cooperative breeding; Conflict; Reproductive competition; Sociality

INTRODUCTION

Infanticide is an extreme form of conflict where dependent young are killed by adult conspecifics. It has been described in a wide range of taxa such as insects (Ratnieks, Foster and Wenseleers, 2006), fish (Jindal *et al.*, 2017), mammals (Lukas and Huchard, 2014, 2019), reptiles (O'Connor and Shine, 2004) and birds (Heinsohn *et al.*, 2011; Riehl, 2016; Cheng *et al.*, 2019). From an evolutionary perspective, infanticide is intriguing because, in addition to the extreme cost for the victims, the costs for the aggressors can also be important. For example, studies in mammals found that infanticide was associated with an increase in aggressive behaviours between adults (both as a way of preventing infanticide and retaliation) and social exclusion of the individuals that could potentially commit infanticide in the future (see reviews in Agrell et al., 1998; Ebensperger, 1998). Hence, for infanticide to evolve and be maintained in any population, the fitness benefits to the actors should be more important than the costs incurred.

The benefits of infanticide appear to be most often associated with direct reproductive benefits, either through access to a mate or competition for resources for developing offspring. In mammals, comparative work has shown a link between infanticide and reproductive benefits that differ among the sexes (Lukas and Huchard, 2014, 2019). For males, killing young brings their mothers into earlier oestrous, which can make them available to reproduce with that male (the sexual selection hypothesis; (Hrdy, 1979) see also (Ebensperger, 1998; Lukas and Huchard, 2014). For females, infanticide appears to be

associated with competition for the resources available for reproduction and the developing

offspring (resource competition hypothesis; Hrdy, 1979; Lukas and Huchard, 2019), but the

evolutionary determinants seem to be more complex than for males (Lukas and Huchard,

2019). The spatial proximity of other females can increase competition for food or the

parental care provided to the offspring by other group members, but also for space (nest or

dens; (Lukas and Huchard, 2019; Brown *et al.*, 2021). Evidence for these different forms of

resource competition has also been found in other taxonomic groups. For example, in the

greater ani *Crotophaga major*, more than one female lay eggs in one nest and females can

toss eggs or kill hatchlings of other females, leading to reduced starvation of their own chicks

(Riehl, 2016). In green-rumped parrotlet *Forpus passerines*, where cavities to breed are in

short supply, non-breeding pairs (or individuals) can commit infanticide in an attempt to gain

access to the cavity where the young were developing (or to mate with one of the parents;

Beissinger and Berg, 2024). Reproductive benefits can also arise indirectly, through kin

selection or through competition among matrilines or clans for access to future reproduction,

to higher social status (which is necessary for achieving reproductive status) or reduced

competition for future offspring (Rubenstein, 2012; Lukas and Huchard, 2019; Brown *et al.*,

2021). Finally, in social insects, workers caring for the developing larvae can identify those

that are not their sisters and selectively kill them (Ratnieks, Foster and Wenseleers, 2006),

providing an example where infanticide is favoured by indirect fitness benefits through

spiteful behaviour (see also West and Gardner, 2010).

Life history trade-offs might also explain cases where infanticide is committed by parents.

For example, in grey-capped social weavers *Pseudonigritta arnaudi,* there is evidence of

infanticide when parents suffered partial predation in the nest (Cheng *et al.*, 2019). In these

cases, the parents ejected the whole clutch and started a new clutch instead of raising the

remaining nestlings, as the cost of raising a partial clutch was unlikely to overcome the

benefit (Cheng *et al.*, 2019). A similar life-history trade-off was suggested in eclectus parrots

Eclectus roratus, where killing male offspring appeared advantageous for parents in poorer-

quality nest hollows, as broods with a single female offspring were found to have higher

reproductive value than broods in which the female had a younger brother (Heinsohn *et al.*,

2011). Further (albeit anecdotal) support for this hypothesis comes from a mammal, the

golden takin *Budorcas taxicolor bedfordi*, where mothers were recorded to kill their

newborns when they failed to stand up within the usual time (Zhao *et al.*, 2020).

Hence, the benefits of infanticide appear to vary according to sex, reproductive status of the

aggressors and reproductive costs and benefits. However, infanticide events are infrequent

and can take place rapidly, which means that they are difficult to document. Therefore,

infanticide and its evolutionary drivers remain poorly understood, in particular in non-

mammalian animals.

Here, we use video recordings collected over 12 years of breeding monitoring to identify

infanticide in a highly social, colonial cooperative bird, the sociable weaver *Philetairus*

socius. Using our individual-based long-term data, we describe the identity, individual and

social attributes of the aggressors: sex, age, colony of origin and resident/immigrant status,

and breeding and helping activity. Based on these data, we examine two types of hypotheses

that could explain the adaptive significance of infanticide in this species, i) reproductive

benefits arising either from sexual selection (access to a mate) or competition for resources

(breeding chamber or food for developing offspring) and ii) life-history trade-offs.

If infanticide is motivated by reproductive benefits, it should be associated with increased

access to reproduction, *i.e.* it should lead to mating with one of the parents of the young killed

to access to a breeding chamber, or both. If the reproductive benefits are associated with

competition for food for the developing offspring, we expect that the aggressors will be

reproducing at the time of committing infanticide or soon after. Alternatively, if infanticide in

sociable weavers is associated with a life-history trade-off, we predict that it should be

committed by the parents after suffering partial predation, or when conditions for raising

nestlings are poor. Partial nest predation is relatively rare in sociable weavers (the main nest

predators are snakes that typically take entire clutches; Covas, Du Plessis, et al., 2008;

D'Amelio et al., 2022). However, climatic conditions are extremely variable and associated

with offspring mortality (D'Amelio *et al.*, 2022), which could lead to situations where raising

the brood is no longer viable for parents.

METHODS

The sociable weaver is a small (ca. 27.5g) sexually monomorphic passerine endemic to the

semi-arid Kalahari savannahs of southern Africa (Maclean, 1973a). Both sexes display a

black plumage patch that is typically fully developed between 4-5 months of age (Maclean,

1973b). They are obligate colonial, living in colony structures that vary in size from less than

10 to several hundred individuals (Maclean, 1973a). Sociable weavers are facultative

cooperative breeders and within each colony there are social clusters comprising breeding

pairs and their helpers. The number of pairs with helpers varies between breeding seasons,

ranging 30-80% (Covas et al., 2006). Helpers are usually the offspring from previous

breeding attempts (from one or both breeders, and can be male and female), but up to 14% of

helpers are more distant relatives (*e.g.*, cousins) or unrelated (Covas *et al.*, 2006; Ferreira,

2021).

The breeding season usually starts in September or October and continues for as long as the

conditions remain suitable, often until April or longer, depending to a large extent on the

amount and pattern of rainfall (Mares *et al.*, 2017). Females can therefore lay several clutches

in the same breeding season (D'Amelio *et al.*, 2024). These are hereafter referred to as

"breeding attempts". Clutch size in this species varies between 2-5 eggs (exceptionally 6;

Covas et al., 2008; Fortuna et al., 2021; Maclean, 1973b). Incubation lasts 15 days and

typically starts on the first or second day of laying, leading to asynchronous hatching. The

nestling period usually lasts 21-24 days (Maclean, 1973b).

This study was conducted at Benfontein Nature Reserve, ca. 6 km south-east of Kimberley

(Northern Cape Province, South Africa, 28°520S, 24°500E, 1190 m.). Most of the colonies in

our study area have been captured and ringed since 1993. Since 2008, all birds in the study

colonies have consistently been marked with a unique colour combination during annual

captures that usually take place between August-September (before the onset of the breeding

season; Covas et al., 2002; Paquet et al., 2015). These captures therefore allow us to identify

new (unringed) individuals that have moved into our study population and track movement

between colonies. Upon capture, a blood sample is taken from the birds' brachial vein to

genetically determine sex (Griffith *et al.*, 1998), parentage and relatedness among individuals

(using 16 polymorphic markers; Van Dijk et al., 2015), which allows to differentiate helpers

from breeders (see below).

Breeding monitoring and individual identification

In this study, we use data from the detailed monitoring of breeding activity that was conducted between September 2010 – March 2023 at 13-15 study colonies per year, and videos recorded at these same colonies from 2014-2023. All nest chambers in these colonies are numbered and were inspected every 1-3 days to detect initiation of new clutches and monitor the fate of eggs and nestlings. The day when the first nestling hatches is considered day 1 of the nestling period and these nests are visited again on day 9 (to collect a blood sample weigh and mark each nestling with a metal leg ring) and on day 17 (to weigh, measure and mark nestlings with an individual colour combination). As part of a long-term experiment (Fortuna *et al.*, 2021; D'Amelio *et al.*, 2022), during this period we have artificially reduced nest predation at 3 to 8 colonies per year by wrapping tree trunks with plastic film. This generally prevented snakes from climbing up trunks and reaching the colonies (see details in D'Amelio et al., 2022). Both protected and natural colonies were included in the present study.

To identify the individuals feeding at each nest, to study nest attendance and determine group size, we collected video-recordings (from 2014 onwards) by placing a camera directly below the nest entrances. Details on the criteria for determining group size and identification of breeders and helpers are given below (see also Fortuna et al., 2021; D'amelio et al., 2024). The video-recordings were conducted in the morning (typically starting 1-3h after sunrise) and recorded both incubation (when mainly the breeding pair visits the nest, as helpers rarely incubate) and the nestling period, when both parents and helpers feed the nestlings. Incubation videos recorded ca. 1.5h (range 1-15 incubation days, 1.94 ± 0.74, mean $\pm$ SD hours) and we analysed 1332 videos for 1208 breeding attempts. Nestling provisioning videos recorded ca. 4-8h with a stronger effort around ages 9 and 17 days (range 1-23 days old, number of days: 2.42 ± 0.87, mean $\pm$ SD hours) and we analysed 5423 videos for 2109 breeding attempts.

Infanticide recordings

The videos obtained during this period allowed us to record several infanticide attempts and, in some cases, to identify the individuals performing them – hereafter called 'aggressors'. We considered that there was an infanticide attempt whenever we saw clear aggression of nestlings that were alive (e.g. repeated pecking or unambiguous bloody pecking marks) and/or we saw an attempt to expel live nestlings from inside the nest. If the nestling was not seen moving at any point during the ejection event, we considered that it had died for other reasons and was being removed from the nest for sanitation. Removal of chicks that were already dead inside the nest occurred in 15 cases, which were excluded from our analyses (Fig. ESM1). We also excluded seven cases where nestlings fell off the nest when no adult was present (Fig. ESM1). This could result from pushing by other nestlings or from an accidental fall. Additionally, we recorded ejections of 12 eggs (from seven broods; Fig. ESM1) but these were not considered in this analysis given that it was impossible to know whether the eggs were fertile, and hence we could not apply the same criteria used for the nestling ejections. In addition, there is greater uncertainty about group membership (i.e. future helpers) at the egg stage (but see Fig. ESM1 suggesting that both parents and intruders can remove eggs).

For each infanticide attempt observed, we recorded whether it was successful (i.e. the chick was seen being pushed out of the nest chamber) or non-successful (i.e. the aggressor was interrupted by other birds or the nestling was able to physically resist the ejection).

Age, sex, and relatedness to the nestlings

Age corresponds to the exact age for all individuals born in the study colonies (where all nestlings have been ringed since 2010). Immigrants were attributed an age of 690 days for immigrant males and 727 days for immigrant females on the date of first capture, which is

based on the average known age of first dispersal from our long-term captures data (n= 4441

birds of known age; 9.84 % of these were immigrants that moved within the study area;

unpublished data). Sociable weavers present a fully developed adult plumage after ca. 6 months

old and from this time their age can no longer be determined by morphological features (R.

Covas pers. obs.).

Sex was determined by genetic analyses and the genetic relatedness to nestlings was obtained

from a pedigree dataset that resulted from a combination of genetic, breeding history and video

analyses (for more details see Fortuna et al., 2021; Silva et al., 2018). These data allowed us to

differentiate three categories of individuals: (1) breeders, (2) helpers or (3) intruders. Breeders

were individuals that were seen feeding the nestlings and that were assigned as parents through

the genetic analyses (or combination of genetic analyses and incubation videos in case of

ambiguity). Helpers were individuals that were seen feeding the nestlings at least three times,

but were not assigned as parents through the criteria described above. All remaining individuals

seen entering the nest are defined as 'intruders'. In addition, we use the term 'aggressors' to

refer to any birds that conducted an attack, regardless of whether they were members of the

breeding group or intruders. Two individuals (n=4 nestling ejections) were inconclusive in their

relationship to the brood (i.e. genetic data was missing and life-history and video data were

insufficient to conclude on relatedness to the nestlings) and thus were not considered in these

relatedness calculations (but were included in the sex-ratio and age calculations, when these

data were available).

Movement and reproductive history of the aggressors

Colony of origin and residency of the aggressors were the criteria used to determine

resident/immigrant status. We considered as resident the birds born at the colony where they

committed infanticide.

We used the aggressors' reproductive history to assign them to one of three breeding categories.

1) Currently breeding, if the individual was known to have a nest with eggs or nestlings at the

time of the infanticide attempt; 2) previous breeder, if the bird was known to have been

previously reproducing at that same colony and 3) non-breeder, if the bird was never recorded

breeding until the date when the infanticide attempt took place.

Future reproduction and nest occupancy

To understand whether aggressors could increase their access to reproduction because of

infanticide, we assessed if they reproduced after the attack, and further scrutinised their future

partner acquisition and nest occupancy. For partner acquisition, we determined if aggressors

subsequently reproduced over the next 365 days, and if they did, whether they paired with one

of the breeders of the attacked nest. For nest occupancy, we determined if aggressors

subsequently reproduced in the chamber that was attacked or a nearby chamber (since sociable

weavers breeding groups can occupy up to 3 nearby chambers; Covas R., Doutrelant C., Silva

L.R., Ferreira A.C. and Fortuna R. unpublished data). We only considered future couples or

chamber occupation involving ringed birds.

All statistical analyses and calculations were conducted in R and the respective code is

available as an electronic appendix.

RESULTS

We recorded on video a total of 50 infanticide attempts of nestlings (out of a total of 70377.15 hours of video over the course of 9 years, but see Fig. ESM2 and below for the distribution of video effort in relation to nestlings' age). These attempts were conducted at 34 nest-chambers in 11 colonies. Of the 50 infanticide attempts, 64.7% (33 nestlings) ended in a successful infanticide (i.e. the aggressors successfully expelled the nestling out of the chamber). In the remainder, the victim's breeding group members chased the aggressors, or the nestling resisted eviction. The latter was often the case with older nestlings (>9 days old). Specifically, we only recorded five cases of nestlings $\geq$ 9 days old being evicted even though >61% of our videos were recorded between days 9-17 of the nestling period (Fig ESM2). The average duration of the successful events was 2.91±4.01 minutes (min 0.08, max 17.43, from the moment an individual enters the nest until it concluded the ejection, i.e. one or more nestlings were ejected from the nest; ESM video).

Sex and age of the aggressors

Of the 27 aggressors with known sex, 59.2% were females (n=16) and 40.7% were males (n=11), including one instance where two birds jointly killed the chicks of a nest (Fig. 1). However, the sex of the aggressors was unknown in 32.5% of cases, as they were not ringed (n=13 assumed different aggressors). For these cases, we estimate that ca. 71% were females, since that corresponds to the percentage of immigrant females that are captured every year in our population (n= 416 immigrant individuals captured from 2014-2023 in the same colonies, and which were subsequently sexed). Hence, after assessing our population sex ratio (47.1 % for females, 52.8 % for males, calculated using 4441 captured individuals with known histories for the seasons and colonies used in this work), we find a tendency for female-biased infanticide attempts (Prop.Test: χ^2= 4.28, df = 1, p-value = 0.04; n= 27 female aggressors as the sum of 16 known females and 11 predicted females; n=15 male aggressors as the sum of 11 known males

and 4 predicted males). However, without such assumption, and using only the average overall

sex ratio obtained from our yearly captures of 47% females ($n=$ 441l), we see no significant

association between sex and infanticide (Prop.Test: $\chi^2 = 1.14$, df $= 1$, p-value $= 0.29$; $n=$ 16

known female aggressors; $n=$ 11 known male aggressors, assuming the same sex ratio as

previous).

Figure 1: Number of aggressors that attempted infanticide ($n=$ 40) categorized by sex and status

(breeder, helper, intruder). Aggressor sex (females on the left, males on the right) also indicates

certainty: 'known' individuals (in dark grey) correspond to the individuals with confirmed sex

information, whereas 'predicted' (in light grey) correspond to the unringed and unsexed

individuals to which we applied the observed immigration sex ratio (71% of adult first captured

as adults were sexed as females; we assumed that the same sex ratio was observed throughout

the year).

All identified aggressors were adult birds. Based on a smaller sample of aggressors and

breeders of know age, 79.1% of aggressors were younger than the same sex breeder of the

brood they had attacked (*n*= 24 aggressor-breeder dyads; Fig 2).

Figure 2: Age difference in days between corresponding nest-chamber pairs of known age

aggressors and breeders (i.e., same-sex breeder of the attacked brood). Negative values mean

that the aggressor is younger than the same sex-breeder of the focal nest. The aggressors' sex

is represented in red and blue for females and males, respectively.

Relationship to the nestlings

Aggressors were usually individuals external to the breeding group. We found that 82.5% (of

the 40 aggressors identified) were not members of the breeding group (*i.e.* breeders nor

helpers). There was, however, evidence of two breeding females ejecting a live nestling from

their own nests (5%) and five cases of helpers ejecting nestlings of the group that they were

helping (12.5%). Four of these were male helpers, with various degrees of relatedness to the

attacked nestling: brother, uncle and great-uncle, and a "social half-brother" that was born as

part of a cross-fostering experiment and helped its social parents (*i.e.* it was not genetically

related to the nestling ejected). The only female helper that committed infanticide was not

related to the attacked nestling but was born in that colony.

Movement and reproductive history of aggressors

The majority of aggressors were born outside the colony where they committed infanticide

(62.5%, *n*= 25 aggressors; Fig. 3). The remainder 37.5% were born at the colony (*n*= 15

aggressors), including the two breeders and the five helpers (Fig. 3). Those that were not part

of the breeding group (*i.e.* those considered "intruders") were mostly born outside of the colony

(75.6%, *n*= 25 aggressors). All non-ringed individuals were considered as not born at the

colonies of the attacks since all the nestlings born in the studied colonies and all adults captured

there were ringed.

Figure 3: Number of aggressors in relation to their migration status (N=40 aggressors) and their relationship to the nestlings attacked (N= 50 nestlings; aggressors could attack more than one nestling). Aggressor status was defined as breeder, helper and intruder. Migration status was defined as being born (in dark grey) or not (in light grey) at the colony where the attack took place.

The reproductive status of the aggressors at the moment of the attack was unknown in 32.5% of cases, as the birds were not ringed (*n*= 13 assumed different aggressors, which were all intruders). For the birds with known histories, we found that 88.9% (*n*= 24) of the aggressors were not breeding at the time of the attack (Fig. 4), although five were helpers at other nests (or had recent helping activity in the 15 days before or after the attack; one of these birds was a helper at another colony). Among these non-breeding intruders, 54.1% (*n*= 13) were never

seen reproducing before committing infanticide, while the rest had previously attempted to breed (45.8%, *n*= 11, all at the same colony of the attack).

Figure 4: Number of birds that attempted infanticide across different aggressor status categories
(breeder, helper and intruder; *n*=27 aggressors). Reproductive status categories are: 1)
"currently breeding", if the individual was known to have a nest with eggs or nestlings at the
time of the infanticide attempt (light grey); 2) "previous breeder", if the bird was known to
have been previously reproducing at the same colony (intermediate grey) and 3) "non-breeder",
if the bird was never recorded breeding until the date when the infanticide attempt took place
(dark grey).

Future reproductive activity

With the exception of the two breeders that committed infanticide, we could only identify the

subsequent breeding activity of 57.9% (*n*= 22) of aggressors (i.e. birds of known identity and

known to be present in monitored colonies after the infanticide). Of these, only 36.4% (*n*= 8; 5

females and 3 males, 3 helpers and 5 intruders) were observed reproducing after the infanticide

(i.e. 121-774 days after the infanticide attempt), and for all of these, that reproductive attempt

corresponded to their first breeding attempt ever recorded. Finally, the majority of these birds

reproduced in the same colony where they conducted the attack (75%; $n = 6$).

None of these aggressors was recorded reproducing with one of the breeders (female or male)

from the victims' nest up to 365 days after the attack (i.e. encompassing a full breeding season

after). One female was seen reproducing with the male of a nest-chamber that she had attacked,

but only 435 days later. Similarly, none of the 22 known aggressors was found to gain access

to the chamber attacked or to a nearby chamber over the following breeding season.

DISCUSSION

Infanticide is challenging to study as it is unpredictable in occurrence and can be very brief

(average time was less than 3 min in this study). Our long-term study has enabled us to gather

multiple observations across the years, providing unambiguous evidence that infanticide

behaviour occurs in sociable weavers at the nestling stage (and also at the egg stage; Fig.

ESM1). Over 80% of the infanticide attempts considered here, were conducted by birds that

were not part of the breeding group, and over 60% of these birds were born in a different

colony. Our results suggest that the aggressors tended to be more often females than males, but

this result requires confirmation, as a large number of aggressors were not individually marked.

In addition, aggressors were usually younger than the parents of the victims. Aggressors were

usually not reproducing when they committed the attacks. No evidence was found that the

aggressors subsequently paired to one of the breeders from the nest attacked, or that they

occupied the chamber from where the chicks were evicted or a nearby chamber. Only a

minority (n=6) of the identified aggressors were found as breeders after the attack, but most of

these reproduced at the same colony, and for the first time.

These findings allow us to exclude several mechanisms as possible benefits of infanticide

behaviour in this species. First, given that the large majority of birds committing infanticide

were not the parents or other members of the breeding group, there is no convincing evidence

that infanticide of nestlings in sociable weavers is generally motivated by a life-history trade-

off. Convincing data of life-history trade-offs motivating infanticide in birds was found in

Eclectus parrots, where sons in poor-quality nest hollows were killed (Heinsohn *et al.*, 2011),

and in grey-capped social weavers where infanticide was associated with partial nest predation,

making it more advantageous to eject the whole clutch and start again, than to raise a reduced

clutch (Cheng *et al.*, 2019). In our study, we had no information about the condition of the

chicks that were ejected by their own mother, and they could have been sick or in poor

condition. In one of the cases, the other nestlings from the brood were all ejected; in the other

case, the three other siblings in the nest fledged. Hence, it is possible that a life-history trade-

off explains some cases of infanticide in sociable weavers, but it is not possible to draw any

conclusions based on these two cases, and this mechanism cannot account for the large majority

of cases of infanticide behaviour in this species.

Second, our results do not provide any evidence of increased access to a mate or nesting

chamber arising from infanticide behaviour. Here, none of the birds committing infanticide

was observed to later reproduce with a parent of one of the victims (with one exception, but

that subsequent pairing took place more than one year later). We also found no evidence that

the aggressors gained access to a nesting chamber, which is an important asset in this species

since it is used for reproduction, roosting and shelter throughout the year (Maclean, 1973a).

Access to a chamber could therefore have improved the probability of breeding for these

aggressors. Finally, with the exception of the two breeding females, only one aggressor (a male)

was reproducing when committing infanticide, suggesting that direct competition for feeding

offspring was not a main factor motivating the majority of the attacks. Hence, our results contrast with studies conducted in mammals, which suggest that infanticide can lead to increased chances of reproduction or to a larger share of resources for the offspring of individuals that are breeding simultaneously (Lukas and Huchard, 2014, 2019).

Of the identified aggressors that could be tracked after the attacks, the majority were born in another colony, and hence had no relatives in the colony where they committed the attacks, but became breeders there at a later stage. It might be possible that reducing the reproductive success of other colony members could contribute to reduce competition for resources for future offspring, and directly for the aggressors, favouring the aggressor's genetic lineage over others in the colony. Although we are not aware of empirical studies demonstrating such a mechanism (but see Hrdy, 1979), we speculate that this could take place in two ways. First, given that sociable weavers are cooperative breeders and the helpers are most commonly the offspring of one or both breeders (Covas et al., 2006; Ferreira, 2021), current reproductive success will influence the future probability of having helpers at the nest. A large group with more helpers is likely to be a stronger competitor within a colony than a single pair or a small group. Helpers contribute to groups tasks such as nest maintenance and defence (Ferreira, 2015; Silva, Covas and Doutrelant pers obs) and larger groups occupy more chambers within the colony (Silva, Covas and Doutrelant pers obs) and have higher reproductive success (D'Amelio et al., 2022b). Hence, infanticide could contribute indirectly to increase the aggressors' future reproductive output or to benefit close relatives through improved access to resources and a more favourable social environment for their future offspring, i.e. it could be a form of spite (West and Gardner, 2010). Studies in some group-living species showing that groups can kidnap the offspring of others (Ridley et al., 2022)) suggest that competition for larger group sizes can be important in animal societies (Heinsohn, 1991; Ridley et al., 2022).

We also found a tendency for the attackers to be younger than the breeders, which is compatible

with the hypothesis of competition among family groups within the colonies, as newcomers

could attempt to increase their status or (future) share of resources in relation to other groups.

This, however, is to some extent contradicted by the five cases of helpers that committed

infanticide at the nest that they were helping. Four of these were males that were related to the

nestlings attacked, although the kinship ranged from brother to great uncle (two of which

reproduced at the colony after the attacks). Increased infanticide by younger group members

was also found in free-ranging house mice *Mus musculus*, where virgin females were more

likely to commit infanticide than pre-pubertal ones (Mccarthy and Saal, 1985). However, in

the house mouse, it is not known whether this age-effect on infanticide behaviour is adaptive.

Future work will need to examine the trajectories of the pairs and families that have committed

and suffered infanticide in sociable weavers to better evaluate whether competition among

groups or clans is a possible explanation for infanticide in this species.

Another non-exclusive explanation is that infanticide could arise from direct competition for

food around the colony, as a way of reducing the numbers of competitors (see also Hrdy, 1979).

We have found previous evidence that in sociable weavers, the number of nest ejections

increases under hot and dry conditions (D'Amelio *et al.*, 2022a), when food resources are

presumably scarce. In that study, it was not possible to determine which proportion of these

ejections were victims of infanticide or were already dead in the nest. In addition, an increase

in infanticide under hot and dry conditions could also be a direct consequence of parents and

helpers spending more time away from the nests in order to find food, thereby leaving the

nestlings alone for longer. Examining the link between hot and dry conditions and infanticide

is particularly relevant, in this and other species, given that the frequency of hot and dry periods

is increasing as a result of global climate change.

Finally, our data suggest that infanticide behaviour is also likely to be associated with costs.

We found that successfully committing infanticide takes effort, with an average of nearly three

minutes per successful infanticide attempt, and our videos show that the aggressors physically

wrest with the nestlings in order to evict them (ESM video). In addition, both the breeding pair

and their helpers fight intruders that attempt to enter the chamber with physical attacks and

displacements. Hence, committing infanticide appears to be physically demanding, aggressors

undertake the risk of being pursued, and even possibly injured. Other potential costs are likely

to involve social retaliation, or a decrease in colony size (since sociable weavers' survival tends

to be higher in larger colonies; (Brown *et al.*, 2003). A better understanding of the costs of

infanticide is important in order to establish whether infanticide is an adaptive behaviour, since

costly behaviours are only expected to evolve if the benefits outweigh the costs.

In conclusion, this study provided clear evidence of infanticide in a highly social and

cooperative bird, and provided results that are not compatible with prevalent hypotheses

suggesting that it provides reproductive benefits, such as increased access to breeding partners,

nest-chambers or a greater share of resources for offspring. Life-history trade-offs might

motivate infanticide in this species, but on rare occasions. Competition among breeding groups

or clans and indirect reproductive benefits are potential mechanisms that are compatible with

the majority of the cases presented here. However, future work following individuals and

groups over extended periods and examining possible interaction with climatic conditions are

required to establish the benefits – and costs – of infanticide in this species, and whether it is

promoted by specific environmental conditions.

Acknowledgements

We are very grateful to the field managers, assistants and student involved in the long-term data collection, especially for help with the yearly captures and breeding monitoring. We thank Franck Theron for his work in the field, training and coordination of the field team and management of the sociable weaver long-term database. De Beers Mining Corporation provided access to Benfontein Nature Reserve and logistical assistance. The sexing and genotyping analyses were conducted at the CTM lab, CIBIO, and we thank Susana Lopes and all the technical staff that conduct the work.

Data, scripts, code, and supplementary information availability

Datasets, scripts, supplementary figures and texts are available on BiorXiv.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article. M. Paquet is a recommender and data editor at PCI.

Funding

The study was funded by FCT (Portugal, grants IF/01411/2014/CP1256/CT0007 and PTDC/BIA-EVF/5249/2014), ERC (EU, Consolidator grant 866489) and the DST-NRF Centre of Excellence at the Fitzpatrick Institute of African Ornithology University of Cape Town (South Africa) to R.C and additional funding from ANR (France, grants 15-CE32-0012-02 and 19-CE02-0014-01) to C.D.. A.C.F was funded by University of Zurich Forschungskredit postdoc grant (K-74312-01-01 University of Zurich), Swiss Federal Commission for Scholarships and by ERC grants 866489 to R.C and 850859 awarded to Damien Farine, R.C.

was funded by FCT (IF/01411/2014/CP1256/CT0007 and CEECIND/03451/2018). PBDA

was supported by a South African Claude Leon Postdoctoral Fellowship 2019-20, a Marie

Curie Individual Fellowship "SOCIAL MATCH" N. 896475 and by ANR 19CE02-0014-01.

R.F. was funded by FCT (SFRH/BD/130134/2017 and COVID/BD/151899/2022). The

Sociable Weaver Project is supported by the French OSU-OREME, and the CNRS programme

on Long-Term Studies in Ecology and Evolution (SEE-Life).

References

Agrell, J., Wolff, J.O. and Ylönnen, H. (1998) Counter-Strategies to Infanticide in Mammals :
Costs and Consequences in mammals, *Oikos*, 83, pp. 507–517.

Beissinger, S.R. and Berg, K.S. (2024) Eviction-driven infanticide and sexually selected
adoption and infanticide in a neotropical parrot, *Proceedings of the National Academy of*
Sciences of the United States of America, 121(20). https://doi.org/10.1073/pnas.2317305121.

Brown, A.K. *et al.* (2021) Infanticide by females is a leading source of juvenile mortality in a
large social carnivore, *American Naturalist*, 198(5), pp. 642–652.
https://doi.org/10.1086/716636.

Brown, C.R. *et al.* (2003) Multistate estimates of survival and movement in relation to colony
size in the sociable weaver, *Behavioral Ecology*, 14(4).
https://doi.org/10.1093/beheco/arg034.

Cheng, Y.R., Rubenstein, D.R. and Shen, S.F. (2019) Nest predation predicts infanticide in a
cooperatively breeding bird, *Biology Letters*, 15(8). https://doi.org/10.1098/rsbl.2019.0314.

Covas, R. *et al.* (2002) Stabilizing selection on body mass in the sociable weaver Philetairus
socius, *Proceedings of the Royal Society of London B*, 269, pp. 1905–1909.

Covas, R. *et al.* (2006) Kin associations and direct vs indirect fitness benefits in colonial
cooperatively breeding sociable weavers Philetairus socius, *Behavioral Ecology and*
Sociobiology, 60, pp. 323–331. https://doi.org/10.1007/s00265-006-0168-2.

Covas, R., Du Plessis, M.A. and Doutrelant, C. (2008) Helpers in colonial cooperatively
breeding sociable weavers Philetairus socius contribute to buffer the effects of adverse
breeding conditions, *Behavioral Ecology and Sociobiology*, 63(1) pp. 103-112.
https://doi.org/10.1007/s00265-008-0640-2.

D'Amelio, P.B. *et al.* (2022) Disentangling climatic and nest predator impact on reproductive
output reveals adverse high-temperature effects regardless of helper number in an arid-region
cooperative bird, *Ecology Letters*, 25(1), pp. 151–162. https://doi.org/10.1111/ele.13913.

D'Amelio, P.B. *et al.* (2024) Benefits of Pair-Bond Duration on Reproduction in a Lifelong
Monogamous Cooperative Passerine, *American Naturalist*, 203(5), pp. 576–589.
https://doi.org/10.1086/729436.

D'Amelio, P.B. *et al.* (2024) Highly variable breeding phenology in an arid region colonial
cooperative breeder, the Sociable Weaver Philetairus socius 95 *Ostrich*.
https://doi.org/10.2989/00306525.2024.2328654.

Van Dijk, R.E. *et al.* (2015) Fine-scale genetic structure reflects sex-specific dispersal
strategies in a population of sociable weavers (Philetairus socius), *Molecular Ecology*,
24(16). Available at: https://doi.org/10.1111/mec.13308.

Ebensperger, L.A. (1998) Strategies and counterstrategies to infanticide in mammals,
Biological Reviews, 73, pp. 321–346.

Ferreira, A.C. (2021) *Social preferences, kinship and helping behaviour in a colonial
cooperatively breeding bird*. PhD Thesis, University of Montpellier.

Ferreira, A.M.C. (2015) *Benefits and costs of helpers: investigating the underlying
mechanisms*. MSc Thesis, Universidade do Porto.

Fortuna, R. *et al.* (2021) Maternal allocation in relation to weather, predation and social
factors in a colonial cooperative bird, *Journal of Animal Ecology*, 90(5), pp. 1122–1133.
https://doi.org/10.1111/1365-2656.13438.

Griffith, R. *et al.* (1998) A DNA test to sex most birds, *Molecular Ecology*, 7, pp. 1071–
1075.

Heinsohn, R. *et al.* (2011) Adaptive Secondary Sex Ratio Adjustments via Sex-Specific
Infanticide in a Bird, *Current Biology*, 21, pp. 1–4. https://doi.org/10.1016/j.cub.2011.08.064.

Heinsohn, R.G. (1991) Kidnapping and reciprocity in cooperatively breeding white-winged
choughs, *Animal Behaviour*, 41, pp. 1097–1100.

Hrdy, S.B. (1979) Infanticide Among Animals: A Review, Classification, and Examination of
the Implications for the Reproductive Strategies of Females, Ethology and Sociobiology 1,
pp. 13–40.

Jindal, S. *et al.* (2017) A test of male infanticide as a reproductive tactic in a cichlid fish,
R.Soc. Open Sci. 4: 160891. http://dx.doi.org/10.1098/rsos.160891

Lukas, D. and Huchard, E. (2014a) The evolution of infanticide by males in mammalian
societies, *Science*, 346, pp. 841–844. https://doi.org/10.1126/science.1257226.

Lukas, D. and Huchard, E. (2019) The evolution of infanticide by females in mammals,
Philosophical Transactions of the Royal Society B: Biological Sciences, 374(1780).
https://doi.org/10.1098/rstb.2018.0075.

Maclean, G.L. (1973a) The sociable weaver, part 2: Nest architecture and social organisation,
Ostrich, 44, pp. 191–218.

Maclean, G.L. (1973b) The sociable weaver, part 3: Breeding biology and moult, *Ostrich*, 44,
pp. 219–240.

Mares, R. *et al.* (2017) Breeding decisions and output are correlated with both temperature
and rainfall in an arid-region passerine, the sociable weaver, *Royal Society Open Science*,
4(9), p. 170835. https://doi.org/10.1098/rsos.170835.

Mccarthy, M.M. and Saal, F.S. Vom (1985) The Influence of Reproductive State On
Infanticide by Wild Female House Mice (*Mus musculus*), *Physiology & Behavior*, 35, pp.
843–849.

O'Connor, D.E. and Shine, R. (2004) Parental care protects against infanticide in the lizard
Egernia saxatilis (Scincidae), *Animal Behavior*, 68, pp. 1361–1369.
https://doi.org/10.1016/j.anbehav.2004.02.014.

Paquet, M. *et al.* (2015) Antagonistic effect of helpers on breeding male and female survival
in a cooperatively breeding bird, *Journal Animal Ecology*, 84, pp. 1354–1362.
https://doi.org/10.1111/1365-2656.12377.

Ratnieks, F.L.W., Foster, K.R. and Wenseleers, T. (2006) Conflict resolution in insect
societies, *Annual Review of Ethomology*, 51, pp. 581–608.
https://doi.org/10.1146/annurev.ento.51.110104.151003.

Ridley, A.R. *et al.* (2022) Kidnapping intergroup young: An alternative strategy to maintain
group size in the group-living pied babbler (Turdoides bicolor), *Philosophical Transactions*
of the Royal Society B: Biological Sciences, 377(1851).
https://doi.org/10.1098/rstb.2021.0153.

Riehl, C. (2016) Infanticide and within-clutch competition select for reproductive synchrony
in a cooperative bird, pp. 1760–1769. https://doi.org/10.1111/evo.12993.

Rubenstein, D.R. (2012) Family feuds: Social competition and sexual conflict in complex
societies, *Philosophical Transactions of the Royal Society B: Biological Sciences*. Royal
Society, pp. 2304–2313. https://doi.org/10.1098/rstb.2011.0283.

Silva, L.R. *et al.* (2018) Females pay the oxidative cost of dominance in a highly social bird,
Animal Behaviour, 144, pp. 135–146. https://doi.org/10.1016/j.anbehav.2018.08.006.

West, S.A. and Gardner, A. (2010) Altruism, Spite, and Greenbeards, *Science*, 327, pp. 1341–
1344.

Zhao, H.T. *et al.* (2020) 'Observation of mother-perpetrated infanticide in golden takins
(Budorcas taxicolor bedfordi)', *Zoological Research*. Science China Press, pp. 455–457.
Available at: https://doi.org/10.24272/j.issn.2095-8137.2020.041.