PINK1 SUMOylation regulates basal mitophagy

Nitheyaa Shree Ramesh[1], Richard Seager[1], Kevin A. Wilkinson[2*],

and Jeremy M. Henley[1*]

[1]Centre for Synaptic Plasticity, School of Biochemistry, Biomedical Sciences Building, University of Bristol, University Walk, Bristol, BS8 1TD, UK.

[2]Centre for Synaptic Plasticity, School of Physiology, Pharmacology and Neuroscience, Biomedical Sciences Building, University of Bristol, University Walk, Bristol, BS8 1TD, UK.

[*]Corresponding authors: kevin.wilkinson@bristol.ac.uk and j.m.henley@bristol.ac.uk

Abstract

The detection and removal of damaged mitochondria is essential for cells to maintain a healthy mitochondrial network. PTEN-induced kinase 1 (PINK1) is a critical sensor that detects mitochondrial damage. PINK1 is subject to several post-translational modifications including phosphorylation, ubiquitination, and S-nitrosylation, each playing a key role in regulating its stability, subcellular localization, and/or function. Here, we show that PINK1 is additionally modified by non-canonical, lysine-independent SUMOylation, mediated by mitochondrial anchored protein ligase (MAPL). Levels of PINK1 SUMOylation are reduced by the mitochondrial uncoupler CCCP, whereas they are increased by the inhibitor of proteasomal degradation MG132. MAPL knockdown increases mitophagy in wild-type, but not PINK1$^{-/-}$ HEK293T cells. Moreover, selective deSUMOylation of PINK1 also increases mitophagy. These findings indicate that SUMOylation of PINK1 is a key factor in the regulation of mitophagy and mitochondrial quality control.

Introduction

Mitophagy, the removal of damaged mitochondria, is fundamental for the maintenance of cellular health and function (Youle & Narendra, 2011). This is emphasized by the observations that dysregulation of mitophagy is a prominent feature of a wide range of diseases including neurodegenerative disorders, cancer, liver disease, and cardiovascular disease (Wang et al., 2023).

PTEN-induced kinase 1 (PINK1) is an essential sensor for the initial detection and subsequent clearance of damaged mitochondria (Jin et al., 2010; Narendra et al., 2010). Moreover, mutations in PINK1 cause autosomal recessive early-onset Parkinson's disease, emphasizing the essential role of PINK1-dependent mitophagy in maintaining cellular health (Gonçalves & Morais, 2021; Valente et al., 2004).

The stress-induced PINK1/Parkin mitophagy pathway has been extensively studied and well characterized (for recent review see (Narendra & Youle, 2024)). Briefly, PINK1 accumulates on the outer mitochondrial membrane in response to membrane depolarization (Jin et al., 2010). It then autophosphorylates to activate its kinase domain, leading to the phosphorylation and subsequent activation of the cytosolic E3 ubiquitin ligase, Parkin (Okatsu et al., 2012; Zhuang et al., 2016). PINK1 also directly phosphorylates ubiquitin at serine 65, which can also activate Parkin recruitment to the mitochondria (Kane et al., 2014). Phosphorylated Parkin, in turn, then ubiquitinates other outer mitochondrial membrane (OMM) proteins. In a positive feedback loop, ubiquitinated OMM proteins recruit more Parkin to ubiquitinate more OMM proteins (Narendra et al., 2008). This accumulation of ubiquitinated OMM proteins leads to the recruitment of

adaptor proteins such as p62, which bind to LC3-II receptors on the autophagosome membrane (Narendra et al., 2010). Although stress-induced PINK1-mediated mitophagy is well characterized, the role, if any, of PINK1 in regulating basal mitophagy remains highly controversial (McWilliams et al., 2018, Y.T. Liu et al., 2021).

Under basal conditions, PINK1 is imported into the mitochondria, and subsequently cleaved by two proteases, mitochondrial protein peptidase (MPP) between residues P34 and G35 (Greene et al., 2012), and presenilin-associated rhomboid-like protein (PARL) between residues A103 and F104 (Deas et al., 2011). This double cleavage yields a mature ~52 kDa form of PINK1, which is then retrotranslocated to the cytosol and targeted for proteasomal degradation (Jin et al., 2010). Treatment of cells with the mitochondrial uncoupler, carbonyl cyanide chlorophenylhydrazone (CCCP), stabilizes PINK1 at the mitochondrial membrane, while treatment of cells with the proteasomal inhibitor, MG132, inhibits degradation of the retrotranslocated cytosolic mature PINK1 by the proteasome (Narendra et al., 2010).

PINK1 is subject to extensive post-translational modification, including phosphorylation, ubiquitination, and S-nitrosylation. PINK1 contains multiple phosphorylation sites and, while the functions of some remain unknown (Luo et al., 2024; Waddell et al., 2023), it has been established that autophosphorylation at S228 or S465 regulate the kinase activity of PINK1 (Aerts et al., 2015). In addition, S167 and C412 have also been identified recently as autophosphorylation sites affecting PINK1 activation (Luo et al., 2024; Waddell et al., 2023).

Ubiquitination of the 52 kDa form of PINK1 at K137 targets it for proteasomal degradation (Liu et al., 2017). Additionally, PINK1 can undergo degradation via the N-end ubiquitination rule, independent of K137 ubiquitination (Yamano & Youle, 2013). In response to specific stressors, PINK1 can be S-nitrosylated at C568, leading to a decrease in kinase activity and reduced mitochondrial translocation of Parkin (Oh et al., 2017).

SUMOylation is the covalent addition of a small ubiquitin like modifier (SUMO) protein to a substrate, canonically at a lysine residue within a SUMO consensus motif (Vertegaal, 2022; Wilkinson & Henley, 2010). SUMO conjugation plays critical roles in protein subcellular localization, stability, and function (Henley et al., 2018; Wilkinson & Henley, 2010). Briefly, SUMO is a ~11 kDa protein, of which there are three mammalian isoforms (SUMO1, SUMO2, and SUMO3). SUMO2 and SUMO3 differ by only three amino acid residues (~97% identity) and are generally collectively referred to as SUMO2/3 (Saitoh & Hinchey, 2000). In contrast, SUMO1 shares only ~50% identity with SUMO2/3. Both SUMO-1-ylation and SUMO-2/3-ylation are highly dynamic, reversible post-translational modifications with conjugation mediated by an E1, E2, and E3 enzyme cascade and deconjugation mediated by specific SUMO proteases, most notably SENPs (Celen & Sahin, 2020; Hay, 2005).

Here, we show that PINK1 can undergo SUMO-2/3-ylation by the SUMO E3 ligase MAPL. Remarkably, however, SUMO-2/3-ylation of PINK1 does not occur at lysine-residues. Nonetheless, our data demonstrate that SUMOylation of PINK1 mediates PINK1-dependent mitophagy. Together these findings indicate that, in a non-canonical form of

SUMO-2/3-ylation, MAPL-mediated SUMOylation of PINK1 contributes to the regulation of basal mitophagy.

Results

PINK1 is modified by SUMO2/3.

To test if PINK1 can be modified by SUMOylation, PINK1-GFP was overexpressed in HEK293T cells and lysed in 2% SDS-containing buffer to disrupt non-covalent interactions but conserve post-translational modifications. PINK1-GFP was immunoprecipitated and samples were blotted for SUMO2/3. As shown in Figure 1, a higher molecular weight ladder was detected, indicative of poly-SUMO chains (Fig 1A). We also immunoblotted PINK1-GFP IPs for ubiquitin. Comparable to SUMO2/3, we observed a ladder indicative of PINK1-GFP polyubiquitination. Co-transfecting the SBP-tagged SUMO protease, SBP-SENP1, significantly decreased the SUMO2/3 smear on PINK1-GFP (Fig 1B), but did not alter the ubiquitin smear. Interestingly, we also observed that the total amount of PINK1-GFP in cells increased significantly upon co-expression of SBP-SENP1 (Fig 1C).

To further confirm the specificity of the SUMO2/3 smear observed on PINK1, cells were lysed in the presence or absence of n-ethylmaleimide (NEM) before PINK1-GFP was immunoprecipitated (Fig 1D). NEM is a cysteine protease inhibitor that prevents activity of sentrin-specific proteases (SENPs). The PINK1-SUMO2/3 smear was significantly increased in cells lysed in NEM (Fig 1E). Taken together, these data indicate that exogenously expressed PINK1 is SUMO-2/3-ylated.

We next tested if endogenous PINK1 is SUMOylated. Mitochondrial-enriched fractions were prepared from HEK293T cells, and PINK1 was immunoprecipitated from these fractions using an anti-PINK1 antibody. A single band appears when PINK1 pulldown samples from mitochondrial fractions were blotted for SUMO2/3. This single SUMO-2/3-ylated PINK1 band was absent in PINK1$^{-/-}$ HEK293T cells (Fig 1F). In the endogenous IP, only one clear band appears and overexposure of the blot does not reveal a distinct higher molecular weight smear, differing from overexpression experiments present in Fig 1A. This could be due to the low levels of SUMOylated PINK1 in cells, especially given that levels of PINK1 in healthy cells are low (Narendra et al., 2010).

Since we observe both poly-SUMOylation and poly-ubiquitination on PINK1 (Fig 1A), we wondered if these may be mixed SUMO-ubiquitin chains (Hendriks et al., 2014). Therefore, to assess the composition of the higher molecular weight smear on PINK1-GFP, we applied constitutively active recombinant SENP1 or the deubiquitinating enzyme USP2 to immunoprecipitated PINK1-GFP from HEK293T cells (Fig 1G). Active SENP1 removed SUMO2/3 from PINK1-GFP but did not affect ubiquitin whereas active USP2 removed ubiquitin but did not affect SUMO2/3. Treatment with both active SENP1 and USP2 completely removed ubiquitin and SUMO2/3 smears on PINK1-GFP (Fig 1G). These data indicate that PINK1 can be simultaneously SUMOylated and ubiquitinated but that mixed SUMO-ubiquitin chains are not present. To further probe the enzymes that regulate PINK1 SUMOylation, we overexpressed SENP1, SENP3, SENP5, SENP6, or SENP7 with PINK1-GFP in HEK293T cells. Each of the individual SENPs decreased levels of SUMOylated PINK1 (Fig S1).

PINK1 is SUMOylated in a non-canonical lysine-independent manner.

It is very well established that SUMO conjugates to lysines residues. Therefore, to identify where PINK1 is SUMOylated, we systematically and additively mutated lysines to non-SUMOylatable arginine residues. Initially, we mutated lysines previously identified to be ubiquitinated as well as lysines located within a SUMO consensus motif (ψ-K-x-D/E, where ψ is a hydrophobic residue) (Fig 2A). Unexpectedly, none of these mutations individually or collectively altered PINK1-GFP SUMOylation. Indeed, PINK1-GFP in which all 21 lysines were mutated to arginines (21KR PINK1-GFP) remained SUMOylated. Importantly, however, SUMOylation in both wild-type PINK1-GFP and 21KR PINK1-GFP was dramatically decreased on co-transfection with SENP1 (Fig 2B).

Although SUMO is canonically conjugated to lysines, it has been reported recently that SUMO can conjugate to the amino group at the N-terminal end of a protein, analogous to N-terminal ubiquitination (Weng et al., 2023). To test if this happens for PINK1, we co-transfected cells with WT PINK1-GFP and N-α-acetyltransferase 60 (Naa60), which acetylates the N-terminal end of proteins in an irreversible manner thereby preventing any further posttranslational modifications. Overexpression of FLAG-Naa60 with either wild-type PINK1-GFP or 21KR PINK1-GFP did not reduce PINK1 SUMOylation (Fig S2).

To ensure that the SUMOylation site(s) were not in the GFP tag we exchanged the GFP for a SPOT tag (amino acid sequence: PDRVRAVSHWSS), thereby removing all lysine residues from the entire expressed protein (21KR PINK1-SPOT). Remarkably, both WT and 21KR PINK1-SPOT were still SUMOylated (Fig 2C). We interpret these data to indicate that PINK1 SUMOylation must occur in a non-canonical lysine-independent manner.

MAPL promotes PINK1 SUMOylation.

We hypothesized that the dual specificity SUMO and ubiquitin ligase mitochondrial anchored protein ligase (MAPL) was likely to mediate PINK1 SUMOylation. To test this we used a SMART pool of siRNA targeting human MAPL to determine the effects of knocking down MAPL in HEK293T cells (Fig 3A). As predicted, MAPL knock down significantly decreased levels of PINK1 SUMOylation (Fig 3B-3C). We next tested if MAPL binds to PINK1 by immunoprecipitating overexpressed PINK1-GFP and blotting for MAPL. As shown in Fig 3D, a single band of MAPL was present in the WT PINK1-GFP lane but not in the GFP control (Fig 3D). We also showed that the OMM protein VDAC did not co-IP with PINK1, consistent with a selective interaction between PINK1 and MAPL.

PINK1 SUMOylation and stability

PINK1 is constitutively degraded in healthy mitochondria with a reported half-life of ~30 minutes (Lee et al., 2015; Lin & Kang, 2008). This constant turnover is mediated by PINK1 import into the mitochondria and cleavge by MPP at P34/G35 and PARL at A103/F104 (Deas et al., 2011; Greene et al., 2012). The remaining fragment is then translocated back to the cytosol where it is degraded via the proteasome (Jin et al., 2010). Conversely, in stressed mitochondria PINK1 escapes MPP and PARL cleavage by being retained and stabilized at the OMM (Narendra et al., 2010).

To determine the effects of SUMOylation on PINK1 turnover we carried out an array of experiments with carbonyl cyanide m-chlorophenyl hydrazone (CCCP) and MG132, as well as two different point mutants of PINK1, P95A and F104A. CCCP dissipates the

mitochondrial membrane and inhibits oxidative phosphorylation, which stabilizes full-length PINK1 at the OMM (Narendra et al., 2010) whereas the proteasomal inhibitor MG132 accumulates the cleaved form of PINK1 (Lin & Kang, 2008). The P95A PINK1 mutant accumulates full-length PINK1 at the OMM, while the F104A mutant accumulates the cytosolic fragment of PINK1 (Deas et al., 2011).

CCCP significantly decreased levels of SUMO-2/3-ylated PINK1-GFP in HEK293T cells (Fig 4A-4B). In contrast, the proteasomal inhibitor MG132, induced a significant increase in PINK1 SUMOylation (Fig 4C-4D). Immunoprecipitation of the PINK1 cleavage mutants showed that SUMOylation was significantly reduced in P95A PINK1-GFP mutant but not the F104A PINK1-GFP mutant compared to WT PINK1-GFP (Fig 4E-4F). Additionally, for cells treated with CCCP, we blotted the total lysates for phospho-ubiquitin to confirm full length PINK1 activity (Fig 4A). The level of total phospho-ubiquitin was significantly higher in the P95A PINK1-GFP mutant compared to WT PINK1-GFP (Fig 4G).

In summary, CCCP treatment decreases, whereas MG132 treatment increases levels of SUMOylated PINK1. Moreover, the cleavage-resistant PINK1-P95A mutant exhibits reduced SUMOylation (Fig 4H). Taken together, these data indicate that non-SUMOylated PINK1 is more stable.

Counteracting SUMOylation of PINK1 promotes mitophagy.

PINK1 is a key mediator of mitophagy, so we next tested how SUMOylation affects PINK1-dependent mitophagy. WT and PINK1$^{-/-}$ cells were transfected with a control siRNA or human MAPL siRNA for 48 hours and treated with 10µM CCCP (or DMSO control) for one hour. Samples were lysed and immunoblotted for a range of autophagy and

mitophagy markers (Fig 5A). MAPL knockdown increased the ratio of LC3-II/LC3-I, indicative of increased autophagy (Fig 5B). As expected, this increase was also observed in cells treated with CCCP. Importantly, the MAPL-knockdown mediated increase in autophagy did not occur in PINK1$^{-/-}$ cells, confirming it occurs through a PINK1-dependent pathway. It is notable, however, that CCCP does increase the LC3-II/LC3-I ratio in PINK1$^{-/-}$ cells, consistent with both PINK1-dependent and independent pathways, as previously reported (Seabright et al., 2020).

MAPL knockdown increased levels of the ubiquitin-autophagy adaptor, p62 (SQSTM1) in WT but not in PINK1$^{-/-}$ HEK293T cells (Fig 5C-5D). Phosphorylated p62 was also increased by MAPL knockdown in WT cells, whereas levels of phosphorylated p62 were significantly lower in PINK1$^{-/-}$ cells and were not further decreased by knockdown of MAPL (Fig 5E-5F). These results suggest that MAPL knockdown, which strongly reduces SUMOylation of PINK1, leads to enhanced PINK1-dependent basal mitophagy.

We extended the autophagy and mitophagy marker blotting data by visualizing mitophagy using a mitochondria-targeted Keima-Red mitophagy probe. WT and PINK1$^{-/-}$ HEK293T cells were transfected with mtKeima (pMitophagy Keima Red mPark2) and an siRNA-targeting human MAPL (Fig 5G). Consistent with the blotting data, MAPL knockdown increased mitophagy in WT, but not in PINK1$^{-/-}$ cells (Fig 5H). However, MAPL knockdown did not cause a further increase mitophagy in either WT or PINK1$^{-/-}$ cells treated with CCCP for one hour, suggesting MAPL knockdown is specifically affecting basal mitophagy.

Also in agreement with the blotting data, CCCP caused mitophagy in both WT and PINK1[-/-] cells, indicative of a PINK1-independent mitophagy pathway (Seabright et al., 2020) (Fig 5H). Together, these results indicate that MAPL promotes basal mitophagy through its actions on PINK1.

Our data demonstrate that PINK1 is SUMOylated but that this occurs in a non-canonical lysine-independent manner. In consequence, despite extensive efforts, we were therefore unable to identify and construct a non-SUMOylatable mutant for loss of function studies. To circumvent this issue as far as possible we used an alternative inducible deSUMOylation approach. To achieve this, we expressed an anti-GFP-nanobody fused to the catalytic domain of SENP1 under a tetracycline promoter (GNb-SENP1) as described previously (Ibrahim et al., 2020; Mojsa et al., 2020). Alongside this we expressed PINK1-GFP. We reasoned that doxycycline induced GNb-SENP1 expression will target SENP1 to deSUMOylate PINK1-GFP (Fig 6A). We validated this system by transfecting HEK293T cells with PINK1-GFP in the presence of the anti-GFP-nanobody (GNb), or the SENP1-fused anti-GFP-nanobody (GNb-SENP1) (Fig 6B-6C). Doxycycline induction of GNb-SENP1 dramatically reduced PINK1 SUMOylation, with no evidence of differences in overall SUMO conjugation from blots of total lysates.

DeSUMOylating PINK1 increased the ratio of LC3-II/LC3-I (Fig 6D-6E), in agreement with our data from MAPL knockdown. Moreover, expression of PINK1-GFP in PINK1[-/-] cells increased mitophagy (Fig 6F-6G) which was enhanced when PINK1-GFP was deSUMOylated (Fig 6F-6G). These results demonstrate that reducing PINK1 SUMOylation increases mitophagy.

Discussion

Most documented SUMO substrates are nuclear proteins but an extensive and increasing number of extranuclear SUMO substrates have been identified (Henley et al., 2018). In particular, mitochondria are a SUMOylation hub with increasing numbers of integral and associated mitochondrial proteins emerging as SUMO substrates. These include dynamin-related protein 1 (Drp1), mitochondrial fission factor (MFF), mitochondrial fission 1 protein (Fis1), mitofusin-2 (Mfn2), and Fas-associated protein with Death Domain (FADD) (Braschi et al., 2009; Choi et al., 2017; Kim et al., 2021; Prudent et al., 2015; Seager et al., 2024; Waters et al., 2022).

Intriguingly, mitochondrial SUMO substrate proteins are involved in specific mitophagy pathways. For example, SUMOylation of Fis1 at K149 is critical for its mitochondrial localization, and SENP3-mediated deSUMOylation of Fis1 promotes mitophagy induced by the iron-chelator deferiprone (Waters et al., 2022). Mfn2 is SUMOylated in response to CCCP or MG132 treatment which promotes perinuclear aggregation of damaged mitochondria, and subsequent targeting for mitophagy (Kim et al., 2021).

MAPL mediates PINK1 SUMOylation.

Here, we demonstrate that MAPL-mediated SUMOylation of PINK1 plays a key role in regulating mitophagy. MAPL is an OMM localized SUMO and ubiquitin E3 ligase that can SUMOylate Drp1 and MFF (Prudent et al., 2015; Seager et al., 2024). We show that knockdown of MAPL significantly decreased SUMO-2/3-ylated PINK1 and that MAPL co-immunoprecipitates with PINK1-GFP, indicating MAPL binds to and SUMOylates PINK1.

Three previous studies have suggested a link between MAPL and PINK1 in regulating mitophagy (Igarashi et al., 2020; Rojansky et al., 2016; Yun et al., 2014). However, each study proposed a different role and none identified a role for SUMOylation. Briefly, using the anticancer drug, gemcitabine, it was reported that upstream actions of MAPL stabilize PINK1 at the OMM to promote PINK1-dependent, Parkin-independent mitophagy (Igarashi et al., 2020). In contrast, a study using both *Drosophila* PINK1/Parkin and mouse cortical neuronal cultures suggested that MAPL and PINK/Parkin act in parallel by regulating mitofusin ubiquitination and stability (Yun et al., 2014). The third study reported that MAPL and Parkin act downstream of PINK1 in parallel pathways to eliminate paternal mitochondria in mouse embryos (Rojansky et al., 2016).

Our results show that MAPL can bind to PINK1 and promotes PINK1 SUMOylation, providing new and important mechanistic detail on the relationship between these key mitochondrial proteins.

SUMOylation regulates PINK1 stability.

Levels of SUMOylated PINK1 are significantly decreased by CCCP. Furthermore, the cleavage-resistant PINK1 P95A mutant is less SUMOylated than wild-type PINK1. We interpret these data to suggest that SUMOylation prevents stabilization of PINK1 at the OMM and/or that SUMOylation promotes PINK1 import and subsequent degradation. Furthermore, our observations that both the P95A PINK1 mutant and CCCP treatment increase PINK1 activity while reducing SUMOylation support the hypothesis that non-SUMOylated PINK1 displays enhanced activity.

We also show that MG132 significantly increases PINK1 SUMOylation. These data are consistent with a previous proteomics study which identified PINK1 as a potential SUMOylation target in HEK293T cells stressed with MG132 (Hendriks et al., 2018).

In healthy mitochondria PINK1 is constitutively turned over. Under basal conditions, or data indicate that this degradative process is regulated by SUMOylation. Thus, we propose that SUMOylation enhances PINK1 cleavage and/or impedes PINK1 stabilization at the membrane (Fig 4H).

Site(s) of PINK1 SUMOylation

Despite exhaustive attempts using different approaches, we were unable to definitively determine the sites of PINK1 SUMOylation. Notably, a mutant PINK1, in which all lysines were replaced with arginines, was still SUMOylated. Moreover, the irreversible N-terminal acetylase, Naa60 ruled out N-terminal SUMOylation of PINK1. These results demonstrate PINK1 undergoes non-canonical lysine-independent SUMOylation. Although unusual, there is an analogous precedent for non-canonical ubiquitination of residues such as serines, cysteines, or threonines (Kelsall, 2022). Interestingly, the chemical reaction involving a nucleophilic acyl transfer during ubiquitination resembles SUMOylation, providing a potential molecular basis for serine, threonine, or cysteine SUMOylation. These possibilities are outside the scope of the current study but represent exciting future avenues of research.

PINK1 SUMOylation and mitophagy

In agreement with previous reports (Li et al., 2015; Yuan et al., 2019), using immunoblot approaches we show that MAPL knockdown increases the LC3-II/LC3-I ratio and p62 levels in WT cells. Crucially, these changes do not occur in PINK1$^{-/-}$ HEK293T cells. Moreover, live imaging of cells transfected with mitoKeima showed increased mitophagy in response to MAPL knockdown in WT but not PINK1$^{-/-}$ cells. These results indicate that MAPL plays a key role in induction of PINK1-dependent mitophagy.

It should be noted that, in our experiments, we used global MAPL knockdown. Therefore, we cannot fully exclude the possibility that the effects we observe on mitophagy could be accounted for by MAPL acting via proteins other than PINK1. Nonetheless, the fact that MAPL knockdown in PINK1$^{-/-}$ cells did not elicit any significant changes in mitophagy markers provides strong evidence for a PINK1-dependent pathway. To more specifically interrogate SUMOylation of PINK1, we used an inducible system in which a nanobody against GFP was fused to the catalytic domain of SENP1. In agreement with the MAPL knockdown studies, these experiments showed that reducing PINK1 SUMOylation increases mitophagy.

In conclusion, we have identified that SUMOylation of PINK1 attenuates basal mitophagy. The removal of damaged mitochondria is critical for cellular health, as highlighted by diseases with impaired mitophagy and dysfunction of mitochondria (Wang et al., 2023). Therefore, mitophagy must be stringently regulated to remove damaged, but spare healthy mitochondria. We describe a novel pathway in which PINK1 SUMOylation, mediated by MAPL, regulates basal mitophagy. These findings are important because understanding how PINK1 SUMOylation fits into the picture of basal mitophagy, and how

it affects overall mitochondrial health and function, is an essential first step towards modulating this process for future therapeutic applications.

Materials and Methods

Reagents and antibodies

HEK293T were acquired from European Collection of Cell Cultures (ECACC). PINK1[-/-] HEK293T cells (Abcam, ab266393) were a kind gift Prof Ian Collinson (University of Bristol, UK). Dulbecco's Modified Eagle's Medium (D5796), fetal bovine serum (FBS), and poly-l-lysine (PLL), N-ethylmaleimide (E3876), Doxycycline (D3447), CCCP, and MG132 (47490) were purchased from Sigma. Sodium pyruvate (11360-070), Dulbecco's Phosphate Buffered Saline (14200-059), L-glutamine (25030-024), 0.05% trypsin-EDTA and Dulbecco's Modified Eagle Medium for live imaging experiments (A14430-01) was obtained from Gibco. Lipofectamine 2000 transfection reagent (11668-019) and sheep IgG Isotype Control (31243) were purchased from Invitrogen.

Complete protease inhibitor cocktail tablets and Bovine Serum Albumin Fraction V (10735094001) were obtained from Roche. Protein G Sepharose 4 Fast Flow beads (17-0618-01) were obtained from GE Healthcare Biosciences. ChromoTek GFP-trap and Spot-trap agarose beads were obtained from Proteintech. Enzymes and buffers used for cloning were obtained from New England Biolabs (NEB): Cut-smart buffer (B7204S), BamHI (R3136S), HindIII (R3104S), Dpn1 (R0176L), and Quick CIP (M0525S). Solution I ligase was obtained from Takara (6022-1).

For western blotting, a list of primary antibodies used in this study can be found in Table 1. HRP-conjugated secondary antibodies were used as follows: donkey anti-sheep IgG HRP (Abcam, Ab6900), goat anti-mouse IgG HRP (Sigma, A3682), rabbit anti-rat IgG HRP (Sigma, A5795), and goat anti-rabbit IgG HRP (Sigma, A6154).

Table 1. List of antibodies used for immunoblotting

Protein	Supplier	Reference Number	Species
GFP	ChromoTek	pabg-1-100	Rat
LC3A/B	Cell Signalling	4108	Rabbit
MAPL	Abcam	ab155511	Rabbit
PINK1	Cell Signalling	6946	Rabbit
PINK1	Novus Biologicals	NBP2-36488	Mouse
PINK1	MRC PPU Reagents and Services	DU34557	Sheep
p-SQSTM1 (S403)	Cell Signalling	39786	Mouse
p-Ubiquitin (S65)	Cell Signalling	62802	Rabbit
SBP	Millipore	MAB10764	Mouse
Spot-Tag	ChromoTek	28A5	Mouse
SQSTM1	Abnova	M01	Mouse
SUMO2/3	Cell Signalling	4971S	Rabbit
Ubiquitin	Cell Signalling	3936	Mouse
VDAC	Cell Signalling	4661	Rabbit
β-actin	Sigma	A5441	Mouse

Plasmids and siRNA

For MAPL knockdown experiments, a SMART pool of siRNA targeting MAPL (Dharmacon, L-007062-00-0005) or a non-targeting pool (Dharmacon

(D-001810-10-05) was used. To generate PINK1-GFP and mutant plasmids, RNA was extracted from HEK293T cells using the Qiagen RNAeasy mini kit, cDNA was made from this RNA using Thermo RevertAid First Strand Kit, and human PINK1 sequence was amplified from this cDNA by PCR using the following primers: Fwd: CACAAGCTTGCCACCATGGCGGTGCGACAGGCGCTGGGC, Rev:

GTGGGATCCTCCAGGGCTGCCCTCCATGAGCAGAG. The PCR product (insert) and vector (peGFP-N1) was then digested using BamH1 and HindIII, ligated at a 3:1 ratio of plasmid:insert, and transformed into DH5α bacteria. The GFP tag was added to the C-terminus of PINK1 since it has previously been shown that N-terminally tagged PINK1 is trapped in the mitochondria, possibly due to N-terminal cleavage of PINK1 (Beilina et al., 2005). For site-directed mutagenesis and PINK1-SPOT, mutants were made using WT PINK1-GFP as the template and mutagenesis primers were comprised of 21 base pairs before and after the desired mutation. 1 µL of Dpn1 was added to this PCR product for 1 hour at 37°C before being purified using GeneJet Gel Extraction kit (#K0691) and transformed into DH5α bacteria.

The mitochondria-targeted Keima probe (pMitophagy Keima Red mPark2, from MBL (AM-V0259M)) was a kind gift from Prof Ian Collinson (University of Bristol, UK). GNb and GNb-SENP1 were modified from a GFP nanobody-NLS-catalytic SENP1 plasmid from Prof Ron Hay (University of Dundee, UK) as described in (Mojsa et al., 2020). Briefly, GNb was produced by introducing a stop codon after the GNb coding sequence, and the GNb-SENP1 construct was modified to remove the nuclear localization sequence and favor cytosolic expression of the SENP1-nanobody fusion protein.

SBP-tagged SENP1, 3, 5, 6 and 7 constructs were produced by amplifying the corresponding coding regions from rat neuronal cDNA and cloning into a variant of the plasmid pcDNA3.1 containing an N-terminal SBP tag.

Cell culture and transfection

HEK293T cells were grown in complete media (Dulbecco's Modified Eagle Medium (DMEM) with 10% fetal bovine serum (FBS), penicillin (1000U)), and streptomycin (0.1mg). Cells were passaged regularly for maintenance in a T75 flask and cells were used for experiments anywhere from P2 till P20. For immunoprecipitation experiments, approximately 1.75 million HEK293T cells were plated into 60 mm dishes and transfected when ~60-70% confluent. In plain DMEM (without FBS, penicillin, or streptomycin), 2.5 µg of DNA (for a double transfection) or 5 µg of DNA (for a single transfection of DNA) was added along with lipofectamine 2000 (Thermo Fisher, 1.5 µL/µg of DNA). For siRNA-mediated knockdown, JetPrime transfection reagent (Polyplus) was used. Similarly, siRNA was added to JetPrime buffer, along with JetPrime transfection reagent (2 µL/µg). This transfection mix was kept at room temperature for 30 minutes and then added dropwise onto cells.

Immunoblotting

Total cell lysates or immunoprecipitation samples were run on 10% or 15% SDS-PAGE gels and for 90 minutes at 120V. Proteins were transferred onto PVDF membranes at 400mA for 90-120 minutes. PVDF membranes were blocked with 5% milk (low-fat milk powder in PBST) or 5% BSA (bovine serum albumin in PBST) for one hour. Membranes were incubated in primary antibody for (Table 1) one hour at room temperature or

overnight at 4°C, and HRP-conjugated secondary antibody for one hour at room temperature. Membranes were placed directly in 1 mL of HRP substrates and developed on the Odyssey Fc LiCor system. Bands were quantified using Image Studio.

Immunoprecipitation and Co-immunoprecipitation

48-72 hours post transfection, cells were lysed in lysis buffer (150 mM NaCl, 50 mM Tris, 1% Triton, protease inhibitors, 20 mM NEM, and 0.1%-2% SDS), sonicated, and spun down for 20 minutes at 16,100g. The soluble fraction was then added to GFP-Trap or Spot-Trap beads (ChromoTek) and gently rotated for 1 hour at 4°C to immunoprecipitate GFP-tagged proteins. Beads were washed 3X with wash buffer (lysis buffer without NEM, protease inhibitors, or SDS). Sample buffer (125 mM Tris pH 6.4, 4% SDS, 10% glycerol, 0.004% bromophenol blue, 10% β-mercaptoethanol) was added to the beads and boiled for 10 minutes at 95°C. Before immunoprecipitation, 6% of total cell lysate was taken for input, added to sample buffer, and boiled for 10 minutes at 95°C. For co-immunoprecipitation experiments, cells were lysed in (20 mM Tris, 137 mM NaCl, 2 mM $Na_4P_2O_7$, 2 mM EDTA, 1% Triton, 25 mM β-glycerophosphate, 10% glycerol, protease inhibitors, and 20 mM NEM) and samples were not sonicated.

Mitochondrial Isolation and Endogenous Immunoprecipitation of PINK1

Confluent 150 mm dishes of HEK293T cells were detached using trypsin, centrifuged 1500 rpm 3 mins, washed once with 1XPBS, pelleted, and frozen at -80° overnight. Cells were then thawed on ice, and mitochondria were isolated as per the manufacturer's protocol (Mitochondrial Isolation Kit for Cultured Cells, Abcam, ab110170). (Note: 20 mM NEM was added to each of the reagents (A, B, and C) provided in the kit to inhibit sentrin-

specific protease activity). Mitochondrial pellet retrieved from isolation protocol was then resuspended in 1XTGS buffer (0.1M Tris-HCl, 0.15M NaCl, 10% glycerol, 0.5% glyco-diosgenin, 20 mM NEM, and protease inhibitors), incubated on ice for 15 minutes, and centrifuged at 13,200 rpm for 10 minutes at 4°C. Lysates were pre-cleared with 30 µL of protein G-sepharose beads for 2 hours. 10% of the pre-cleared samples were taken for total lysate, added to sample buffer, and boiled 95°C for 10 minutes. Sheep anti-PINK1 (S085D) or control sheep-IgG (Thermo, #31243) was incubated with pre-cleared lysate overnight. Samples were then incubated with 20 µL of protein G beads for 1 hour. Beads were washed 3X, added to sample buffer, and boiled 95°C for 10 minutes.

In-vitro deSUMOylation and deubiquitination assay

PINK1-GFP was transfected in HEK293T cells and immunoprecipitated using GFP-trap beads as described above. Beads were washed three times, and then subsequently treated with constitutively active SENP1 (100 nM, purified in-house as described in (Rocca et al., 2017)) or active USP2 (500 nM, Biotechne, E-504) for 2 hours at 37°C. 2X laemmli buffer was added to the samples, boiled 95°C for 10 minutes, run on SDS-page, and immunoblotted for SUMO2/3 and ubiquitin.

mKeima-Red mitophagy assay

The mitochondrial-targeted mKeima probe (pMitophagy Keima-Red mPark2) has a bimodal excitation spectrum that is sensitive to pH, with an excitation wavelength of 405 nm around a pH of 8, and 561 nm around a pH of 4. The ratio of these two fluorescence values can be used as measure of mitophagy, with greater fluorescence in 561 nm excitation channel denoting more mitophagy. 35 mm glass bottom dishes (Greiner) were

coated with poly-L-lysine for at least one hour in 37°C to increase cell adhesion. 250,000 cells were seeded in each dish and transfected the following day. 36-48 hours after transfection, cells were imaged live using the SpinSR microscope. At least one hour before imaging, cell media was changed to "extra-buffering" media without phenol red (DMEM (A14430-01), 10% FBS, 2mM L-glutamine, 4.5 g/L glucose, 1mM sodium pyruvate, 40 mM HEPES pH 7.4). Cells were imaged live on the Olympus IXplore SpinSR system (Yokogawa CSU-W1 SoRa spinning disk). Images were acquired using a 60X oil immersion lens (1.5 numerical aperture, 0.11 mm working distance). During image acquisition, glass dishes were kept in a 37°C chamber with constant CO_2 supply. All acquisition settings (laser power, gain, exposure time, etc.) were kept constant across all repeats and conditions within each experiment. Cells were excited using 405 nm, 488nm, and 561 nm lasers and z-stacks of 10 µm range with 0.2 µm step size was obtained. To quantify percent of mitophagy ("mitophagy index"), the fluorescence in the 561 nm channel was divided by the total fluorescence (a sum of both 405 and 561 nm).

Statistical analysis

A one sample t-test (where the control was set to 100%) or as appropriate, a one-way or two-way ANOVA followed by Tukey's post-hoc test was conducted to determine significance. For comparisons with more than two groups, all data was normalized to the average of control condition values across all repeats. For imaging experiments, within one biological repeat, all values were normalized to the average of the control condition. Outliers were removed using the ROUT method (Q= 1%). D'Agostino & Pearson test was used to test normality of data, and Kruskal-Wallis test (followed by Dunn's post-hoc test)

was conducted to test significance. All data are shown as mean ± SEM. Graphs and statistical tests were done on GraphPad Prism.

Acknowledgements

The authors thank Prof Ian Collinson (University of Bristol, UK) for providing the mitochondria-targeted Keima probe and PINK1$^{-/-}$ HEK293T cells. We are grateful to Prof Ron Hay (University of Dundee, UK) for the GFP nanobody-NLS-catalytic SENP1 plasmid. The authors also gratefully acknowledge the Wolfson Bioimaging Facility (University of Bristol, UK) for their support and assistance in this work.

Funding: This work was supported by the BBSRC (BB/R00787X/1) and the Wellcome Trust (105384/Z/14/A). NSR is funded by a Marshall Scholarship.

Author contributions:

Conceptualisation: NSR, RS, KAW, JMH

Methodology: NSR, RS, KAW, JMH

Experimentation: NSR

Data analysis: NSR, RS

Supervision: KAW, JMH

Writing – original draft: NSR, JMH

Writing – review and editing: NSR, RS, KAW, JMH

Competing interests: Authors declare they have no competing interests.

References

Aerts, L., Craessaerts, K., De Strooper, B., & Morais, V. A. (2015). PINK1 Kinase Catalytic Activity Is Regulated by Phosphorylation on Serines 228 and 402. *Journal of Biological Chemistry*, *290*(5), 2798–2811. https://doi.org/10.1074/JBC.M114.620906

Beilina, A., Van Der Brug, M., Ahmad, R., Kesavapany, S., Miller, D. W., Petsko, G. A., & Cookson, M. R. (2005). Mutations in PTEN-induced putative kinase 1 associated with recessive parkinsonism have differential effects on protein stability. *Proceedings of the National Academy of Sciences*, *102*(16), 5703–5708. https://doi.org/10.1073/pnas.0500617102

Braschi, E., Zunino, R., & McBride, H. M. (2009). MAPL is a new mitochondrial SUMO E3 ligase that regulates mitochondrial fission. *EMBO Reports*, *10*(7), 748-754–754. https://doi.org/https://doi.org/10.1038/embor.2009.86

Celen, A. B., & Sahin, U. (2020). Sumoylation on its 25th anniversary: mechanisms, pathology, and emerging concepts. *The FEBS Journal*, *287*(15), 3110–3140. https://doi.org/https://doi.org/10.1111/febs.15319

Choi, S.-G., Kim, H., Jeong, E. Il, Lee, H.-J., Park, S., Lee, S.-Y., Lee, H.-J., Lee, S. W., Chung, C. H., & Jung, Y.-K. (2017). SUMO-modified FADD recruits cytosolic Drp1 and caspase-10 to mitochondria for regulated necrosis. *Molecular and Cellular Biology*, *37*(2), e00254-16.

Deas, E., Plun-Favreau, H., Gandhi, S., Desmond, H., Kjaer, S., Loh, S. H. Y., Renton, A. E. M., Harvey, R. J., Whitworth, A. J., Martins, L. M., Abramov, A. Y., & Wood, N. W. (2011). PINK1 cleavage at position A103 by the mitochondrial protease PARL. *Human Molecular Genetics*, *20*(5), 867–879. https://doi.org/10.1093/hmg/ddq526

Gonçalves, F. B., & Morais, V. A. (2021). PINK1: A Bridge between Mitochondria and Parkinson's Disease. *Life*, *11*(5). https://doi.org/10.3390/life11050371

Greene, A. W., Grenier, K., Aguileta, M. A., Muise, S., Farazifard, R., Haque, M. E., McBride, H. M., Park, D. S., & Fon, E. A. (2012). Mitochondrial processing peptidase regulates PINK1 processing, import and Parkin recruitment. *EMBO Reports*, *13*(4), 378–385. https://doi.org/https://doi.org/10.1038/embor.2012.14

Hay, R. T. (2005). SUMO: A History of Modification. *Molecular Cell*, *18*(1), 1–12. https://doi.org/https://doi.org/10.1016/j.molcel.2005.03.012

Hendriks, I. A., D'souza, R. C. J., Yang, B., Verlaan-de Vries, M., Mann, M., & Vertegaal, A. C. O. (2014). Uncovering global SUMOylation signaling networks in a site-specific manner. *Nature Structural & Molecular Biology*, *21*(10), 927–936.

Hendriks, I. A., Lyon, D., Su, D., Skotte, N. H., Daniel, J. A., Jensen, L. J., & Nielsen, M. L. (2018). Site-specific characterization of endogenous SUMOylation across species and organs. *Nature Communications*, *9*(1), 2456. https://doi.org/10.1038/s41467-018-04957-4

Henley, J. M., Carmichael, R. E., & Wilkinson, K. A. (2018). Extranuclear SUMOylation in Neurons. *Trends in Neurosciences*, *41*(4), 198–210. https://doi.org/https://doi.org/10.1016/j.tins.2018.02.004

Ibrahim, A. F. M., Shen, L., Tatham, M. H., Dickerson, D., Prescott, A. R., Abidi, N., Xirodimas, D. P., & Hay, R. T. (2020). Antibody RING-Mediated Destruction of Endogenous Proteins. *Molecular Cell*, *79*(1), 155-166.e9. https://doi.org/https://doi.org/10.1016/j.molcel.2020.04.032

Igarashi, R., Yamashita, S., Yamashita, T., Inoue, K., Fukuda, T., Fukuchi, T., & Kanki, T. (2020). Gemcitabine induces Parkin-independent mitophagy through mitochondrial-resident E3 ligase MUL1-mediated stabilization of PINK1. *Scientific Reports*, *10*(1), 1465. https://doi.org/10.1038/s41598-020-58315-w

Jin, S. M., Lazarou, M., Wang, C., Kane, L. A., Narendra, D. P., & Youle, R. J. (2010). Mitochondrial membrane potential regulates PINK1 import and proteolytic destabilization by PARL. *Journal of Cell Biology*, *191*(5), 933–942. https://doi.org/10.1083/jcb.201008084

Kane, L. A., Lazarou, M., Fogel, A. I., Li, Y., Yamano, K., Sarraf, S. A., Banerjee, S., & Youle, R. J. (2014). PINK1 phosphorylates ubiquitin to activate Parkin E3 ubiquitin ligase activity. *Journal of Cell Biology*, *205*(2), 143–153. https://doi.org/10.1083/jcb.201402104

Kelsall, I. R. (2022). Non-lysine ubiquitylation: Doing things differently. *Frontiers in Molecular Biosciences*, *9*. https://www.frontiersin.org/journals/molecular-biosciences/articles/10.3389/fmolb.2022.1008175

Kim, C., Juncker, M., Reed, R., Haas, A., Guidry, J., Matunis, M., Yang, W. C., Schwartzenburg, J., & Desai, S. (2021). SUMOylation of mitofusins: A potential mechanism for perinuclear mitochondrial congression in cells treated with mitochondrial stressors. *Biochimica et Biophysica Acta - Molecular Basis of Disease*, *1867*(6). https://doi.org/10.1016/j.bbadis.2021.166104

Lee, S., Zhang, C., & Liu, X. (2015). Role of Glucose Metabolism and ATP in Maintaining PINK1 Levels during Parkin-mediated Mitochondrial Damage Responses * . *Journal of Biological Chemistry*, *290*(2), 904–917. https://doi.org/10.1074/jbc.M114.606798

Li, J., Qi, W., Chen, G., Feng, D., Liu, J., Ma, B., Zhou, C., Mu, C., Zhang, W., Chen, Q., & Zhu, Y. (2015). Mitochondrial outer-membrane E3 ligase MUL1 ubiquitinates ULK1 and regulates selenite-induced mitophagy. *Autophagy*, *11*(8), 1216–1229. https://doi.org/10.1080/15548627.2015.1017180

Lin, W., & Kang, U. J. (2008). Characterization of PINK1 processing, stability, and subcellular localization. *Journal of Neurochemistry*, *106*(1), 464–474. https://doi.org/https://doi.org/10.1111/j.1471-4159.2008.05398.x

Liu, Y., Guardia-Laguarta, C., Yin, J., Erdjument-Bromage, H., Martin, B., James, M., Jiang, X., & Przedborski, S. (2017). The Ubiquitination of PINK1 Is Restricted to Its Mature 52-kDa Form. *Cell Reports*, *20*(1), 30–39. https://doi.org/https://doi.org/10.1016/j.celrep.2017.06.022

Luo, S., Wang, D., & Zhang, Z. (2024). Post-translational modification and mitochondrial function in Parkinson's disease. *Frontiers in Molecular Neuroscience*, *16*. https://www.frontiersin.org/journals/molecular-neuroscience/articles/10.3389/fnmol.2023.1329554

McWilliams, T. G., Prescott, A. R., Montava-Garriga, L., Ball, G., Singh, F., Barini, E., Muqit, M. M. K., Brooks, S. P., & Ganley, I. G. (2018). Basal Mitophagy Occurs Independently of PINK1 in Mouse Tissues of High Metabolic Demand. *Cell Metabolism*, *27*(2), 439-449.e5. https://doi.org/10.1016/j.cmet.2017.12.008

Mojsa, B., Tatham, M. H., Davidson, L., Liczmanska, M., Wright, J. E., Wiechens, N., Gierlinski, M., Owen-Hughes, T., & Hay, R. T. (2020). SUMO maintains the chromatin environment of human induced pluripotent stem cells. *BioRxiv*, 2020.12.22.423944. https://doi.org/10.1101/2020.12.22.423944

Narendra, D., Kane, L. A., Hauser, D. N., Fearnley, I. M., & Youle, R. J. (2010). p62/SQSTM1 is required for Parkin-induced mitochondrial clustering but not mitophagy; VDAC1 is dispensable for both. *Autophagy*, *6*(8), 1090–1106. https://doi.org/10.4161/auto.6.8.13426

Narendra, D. P., Jin, S. M., Tanaka, A., Suen, D.-F., Gautier, C. A., Shen, J., Cookson, M. R., & Youle, R. J. (2010). PINK1 Is Selectively Stabilized on Impaired Mitochondria to Activate Parkin. *PLOS Biology*, *8*(1), e1000298-. https://doi.org/10.1371/journal.pbio.1000298

Narendra, D. P., & Youle, R. J. (2024). The role of PINK1–Parkin in mitochondrial quality control. *Nature Cell Biology*, *26*(10), 1639–1651. https://doi.org/10.1038/s41556-024-01513-9

Narendra, D., Tanaka, A., Suen, D.-F., & Youle, R. J. (2008). Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. *Journal of Cell Biology*, *183*(5), 795–803. https://doi.org/10.1083/jcb.200809125

Oh, C.-K., Sultan, A., Platzer, J., Dolatabadi, N., Soldner, F., McClatchy, D. B., Diedrich, J. K., Yates III, J. R., Ambasudhan, R., Nakamura, T., Jaenisch, R., & Lipton, S. A. (2017). S-Nitrosylation of PINK1 Attenuates PINK1/Parkin-Dependent Mitophagy in hiPSC-Based Parkinson's Disease Models. *Cell Reports*, *21*(8), 2171–2182. https://doi.org/10.1016/j.celrep.2017.10.068

Okatsu, K., Oka, T., Iguchi, M., Imamura, K., Kosako, H., Tani, N., Kimura, M., Go, E., Koyano, F., Funayama, M., Shiba-Fukushima, K., Sato, S., Shimizu, H., Fukunaga, Y., Taniguchi, H., Komatsu, M., Hattori, N., Mihara, K., Tanaka, K., & Matsuda, N. (2012). PINK1 autophosphorylation upon membrane potential dissipation is essential for Parkin recruitment to damaged mitochondria. *Nature Communications*, *3*(1), 1016. https://doi.org/10.1038/ncomms2016

Prudent, J., Zunino, R., Sugiura, A., Mattie, S., Shore, G. C., & McBride, H. M. (2015). MAPL SUMOylation of Drp1 Stabilizes an ER/Mitochondrial Platform Required for Cell Death. *Molecular Cell*, *59*(6), 941–955. https://doi.org/10.1016/J.MOLCEL.2015.08.001

Rojansky, R., Cha, M.-Y., & Chan, D. C. (2016). Elimination of paternal mitochondria in mouse embryos occurs through autophagic degradation dependent on PARKIN and MUL1. *ELife*, *5*, e17896. https://doi.org/10.7554/eLife.17896

Saitoh, H., & Hinchey, J. (2000). Functional heterogeneity of small ubiquitin-related protein modifiers SUMO-1 versus SUMO-2/3. *Journal of Biological Chemistry*, *275*(9), 6252–6258.

Seabright, A. P., Fine, N. H. F., Barlow, J. P., Lord, S. O., Musa, I., Gray, A., Bryant, J. A., Banzhaf, M., Lavery, G. G., Hardie, D. G., Hodson, D. J., Philp, A., & Lai, Y.-C. (2020). AMPK activation induces mitophagy and promotes mitochondrial fission while activating TBK1 in a PINK1-Parkin independent manner. *The FASEB Journal*, *34*(5), 6284–6301. https://doi.org/https://doi.org/10.1096/fj.201903051R

Seager, R., Ramesh, N. S., Cross, S., Guo, C., Wilkinson, K. A., & Henley, J. M. (2024). SUMOylation of MFF coordinates fission complexes to promote stress-induced mitochondrial fragmentation. *Science Advances*, *10*(40), eadq6223. https://doi.org/10.1126/sciadv.adq6223

Valente, E. M., Abou-Sleiman, P. M., Caputo, V., Muqit, M. M. K., Harvey, K., Gispert, S., Ali, Z., Del Turco, D., Bentivoglio, A. R., Healy, D. G., Albanese, A., Nussbaum, R., González-Maldonado, R., Deller, T., Salvi, S., Cortelli, P., Gilks, W. P., Latchman, D.

S., Harvey, R. J., … Wood, N. W. (2004). Hereditary Early-Onset Parkinson's Disease Caused by Mutations in PINK1. *Science*, *304*(5674), 1158–1160. https://doi.org/10.1126/science.1096284

Vertegaal, A. C. O. (2022). Signalling mechanisms and cellular functions of SUMO. *Nature Reviews Molecular Cell Biology*, *23*(11), 715–731. https://doi.org/10.1038/s41580-022-00500-y

Waddell, A. D., Ojha, H., Agarwal, S., Clarke, C. J., Terriente-Felix, A., Zhou, H., Kakade, P., Knebel, A., Shaw, A. M., Gourlay, R., Varghese, J., Soares, R. F., Toth, R., Macartney, T., Eyers, P. A., Morrice, N., Bayliss, R., Whitworth, A. J., Eyers, C. E., & Muqit, M. M. K. (2023). Regulation of Human PINK1 ubiquitin kinase by Serine167, Serine228 and Cysteine412 phosphorylation. *BioRxiv*, 2023.03.31.534916. https://doi.org/10.1101/2023.03.31.534916

Wang, S., Long, H., Hou, L., Feng, B., Ma, Z., Wu, Y., Zeng, Y., Cai, J., Zhang, D., & Zhao, G. (2023). The mitophagy pathway and its implications in human diseases. *Signal Transduction and Targeted Therapy*, *8*(1), 304. https://doi.org/10.1038/s41392-023-01503-7

Waters, E., Wilkinson, K. A., Harding, A. L., Carmichael, R. E., Robinson, D., Colley, H. E., & Guo, C. (2022). The SUMO protease SENP3 regulates mitochondrial autophagy mediated by Fis1. *EMBO Reports*, *n/a*(n/a), e48754. https://doi.org/https://doi.org/10.15252/embr.201948754

Weng, W., Gu, X., Yang, Y., Zhang, Q., Deng, Q., Zhou, J., Cheng, J., Zhu, M. X., Feng, J., Huang, O., & Li, Y. (2023). N-terminal α-amino SUMOylation of cofilin-1 is critical for its regulation of actin depolymerization. *Nature Communications*, *14*(1), 5688. https://doi.org/10.1038/s41467-023-41520-2

Wilkinson, K. A., & Henley, J. M. (2010). Mechanisms, regulation and consequences of protein SUMOylation. *Biochemical Journal*, *428*(2), 133–145. https://doi.org/10.1042/BJ20100158

Yamano, K., & Youle, R. J. (2013). PINK1 is degraded through the N-end rule pathway. *Autophagy*, *9*(11), 1758–1769. https://doi.org/10.4161/auto.24633

Youle, R. J., & Narendra, D. P. (2011). Mechanisms of mitophagy. *Nature Reviews Molecular Cell Biology*, *12*(1), 9–14. https://doi.org/10.1038/nrm3028

Yuan, Y., Li, X., Xu, Y., Zhao, H., Su, Z., Lai, D., Yang, W., Chen, S., He, Y., Li, X., Liu, L., & Xu, G. (2019). Mitochondrial E3 ubiquitin ligase 1 promotes autophagy flux to suppress the development of clear cell renal cell carcinomas. *Cancer Science*, *110*(11), 3533–3542. https://doi.org/https://doi.org/10.1111/cas.14192

Yun, J., Puri, R., Yang, H., Lizzio, M. A., Wu, C., Sheng, Z.-H., & Guo, M. (2014). MUL1 acts in parallel to the PINK1/parkin pathway in regulating mitofusin and compensates for loss of PINK1/parkin. *ELife*, *3*, e01958. https://doi.org/10.7554/eLife.01958

Zhuang, N., Li, L., Chen, S., & Wang, T. (2016). PINK1-dependent phosphorylation of PINK1 and Parkin is essential for mitochondrial quality control. *Cell Death & Disease*, *7*(12), e2501–e2501. https://doi.org/10.1038/cddis.2016.396

Figures

A SUMO-2/3-ylation of PINK1-GFP

B

C

D NEM assay

E

F Endogenous PINK1 SUMOylation

G In-vitro deSUMOylation and deUbiquitination assay

Figure 1. PINK1 is SUMO-2/3-ylated. (A) HEK293T cells were transfected with GFP, PINK1-GFP, and pcDNA3.1 or SBP-SENP1. Representative blot of SUMO2/3 and Ub higher molecular weight smears on immunoprecipitated PINK1-GFP and corresponding input samples. **(B)** Quantification of SUMO2/3 smear on PINK1-GFP immunoprecipitation (SUMO2/3/GFP normalized to control (PINK1-GFP + pcDNA)) (n=3, one-sample t-test, ***p<0.001) **(C)** Quantification of total PINK1-GFP expressed in cells (GFP/actin normalized to control (PINK1-GFP + pcDNA)) (n=3, one-sample t-test, *p<0.05) **(D)** HEK293T cells overexpressing PINK1-GFP were lysed in the presence or absence of n-ethylmaleimide (NEM), a cysteine protease inhibitor. Representative blots of SUMO2/3 higher molecular weight smear on PINK1-GFP and corresponding inputs. **(E)** Quantification of SUMO2/3 smear on immunoprecipitated PINK1 (SUMO2/3/GFP normalized to control (+NEM)) (n=3, one-sample t-test, ****p<0.0001) **(F)** Endogenous PINK1 was immunoprecipitated from mitochondrial fractions prepared from WT and PINK1$^{-/-}$ HEK293T cells, and samples were immunoblotted for SUMO2/3 and PINK1. 10% input of both WT and PINK1$^{-/-}$ cells were immunoblotted for PINK1 and β-actin. **(G)** HEK293T cells were transfected with GFP or PINK1-GFP for 48 hours. Cells were lysed and PINK1-GFP was immunoprecipitated using GFP-trap beads. Active SUMO protease, SENP1 (100 nM) or active deubiquitinase, USP2 (500 nM) were added to IP samples for 2 hours at 37° C, samples were run on SDS-PAGE gel and immunoblotted for SUMO2/3 and ubiquitin.

Selection of KR PINK1-GFP mutants

Figure 2. PINK1 is not SUMOylated at a lysine. (A) Initial set of PINK1-GFP lysine

mutants. HEK293T cells were transfected with GFP, WT PINK1-GFP, or KR mutants of

PINK1-GFP for 48 hours. PINK1 was immunoprecipitated using GFP trap beads, and

samples were immunoblotted for SUMO2/3 and GFP. **(B)** HEK293T cells were transfected

with WT PINK1-GFP or 21KR PINK1-GFP along with SBP-SENP1. GFP-Trap performed

on lysate, and samples probed for SUMO2/3 and GFP. Lysates were probed for PINK1

and SBP. **(C)** WT and 21KR PINK1-Spot were overexpressed in HEK293T cells and

PINK1 was immunoprecipitated using ChromoTek Spot-Trap beads. Samples were then

immunoblotted for SUMO2/3, Spot, and PINK1.

Figure 3. MAPL promotes PINK1 SUMOylation. (A) HEK293T cells were transfected with PINK-GFP and a control siRNA (20 nM) or MAPL siRNA (20 nM) for 48 hours. Representative blot of SUMO2/3 higher molecular weight smears on immunoprecipitated PINK1-GFP and corresponding input samples. **(B)** Quantification of SUMO2/3 smear on PINK1-GFP immunoprecipitation (SUMO2/3/GFP normalized to control (siCtrl)) (n=3, one-sample t-test, *p<0.05)**. (C)** Quantification of MAPL knockdown in total cell lysate (MAPL/actin normalized to control (siCtrl)) (n=3, one-sample t-test, *p<0.05) **(D)** HEK293T cells were transfected with WT PINK1-GFP and cells were lysed in buffer without SDS to preserve non-covalent interactions. PINK1 was immunoprecipitated using GFP trap beads and samples were immunoblotted for MAPL, VDAC, and GFP. Representative blot of at least three independent experiments.

Figure 4. PINK1 SUMOylation and stability. (A) SUMOylation of PINK1 in response to CCCP treatment. HEK293T cells were transfected with PINK1-GFP and treated with 10 µM CCCP (or DMSO control) for one hour prior to lysis. Representative blots of higher molecular weight SUMO2/3 smear of immunoprecipitated PINK1 and corresponding input samples. **(B)** Quantification of higher molecular weight SUMO2/3 smear on PINK1 (SUMO2/3/GFP normalized to control (DMSO)) (n=4, one sample test *p<0.05) **(C)** SUMOylation of PINK1 in response to MG132 treatment. HEK293T cells were transfected with PINK1-GFP and treated with 20 µM MG132 (or DMSO control) overnight before immunoprecipitating PINK1 using GFP trap beads. Representative blots of higher molecular weight SUMO2/3 and ubiquitin smear on immunoprecipitated PINK1. **(D)** Quantification of SUMO-2/3-ylated PINK1 in DMSO and MG132 conditions (SUMO2/3/GFP normalized to control (DMSO)) (n=4, one sample t-test, *p<0.05) **(E)** HEK293T cells were transfected with WT, P95A, or F104A PINK1-GFP for 48 hours. Representative blots of higher molecular weight SUMO2/3 and ubiquitin smear on immunoprecipitated PINK1 along with corresponding inputs. **(F)** Quantification of higher molecular weight SUMO2/3 smear on PINK1 (SUMO2/3/GFP normalized to control (WT PINK1-GFP)) (n=4, one sample t-test, **p<0.01) **(G)** Quantification of total phosho-ubiquitin (S65) in lysates (pUb/GFP normalized to control (WT PINK1-GFP)) (n=4, one sample t-test, *p<0.05) **(H)** Potential model of the role of PINK1 SUMOylation in its stabilization: Both the P95A mutant of PINK1 and treatment of cells with CCCP decreased levels of SUMOylated PINK1 while treatment of cells with MG132 increased levels of SUMOylated PINK1. *Created in BioRender.*

Figure 5. Knockdown of MAPL increases basal mitophagy. (A and C) Representative

blots of WT and PINK1$^{-/-}$ cells, transfected with a control or human MAPL targeting siRNA

(20 nM). Cells were treated with 10 µM CCCP (or DMSO control) for one hour before

lysis. Quantification of **(B)** LC3-II:LC3-I ratio (two-way ANOVA with Tukey's post hoc test,

n=5, *$p<0.05$, **$p<0.01$). Quantification of **(D)** p62, **(E)** phosphorylated p62 at S403, and

(F) p-p62:p62 ratio (two-way ANOVA with Tukey's post hoc test, n=4, *$p<0.05$, **$p<0.01$)

(G) Representative images of WT and PINK1$^{-/-}$ cells transfected with mitochondrial

mtKeima (0.5 µg) and siRNA control (20 nM) or siRNA targeting human MAPL (20 nM)

for 48 hours before imaging. Cells were treated with 10 µM CCCP (or DMSO control) for

one hour before imaging. Scale bar denotes 10 µm. **(H)** Quantification of "mitophagy

index": fluorescence in 561 nm excitation channel divided by the total fluorescence in 405

nm and 561 nm excitation channels as a measure of "total mitophagy: total mitochondrial

area" (C561/(C561 + C405)) (Kruskal-Wallis test followed by Dunn's multiple comparison

test, N=3, n= 101-189 cells, ****$p<0.0001$)

Figure 6. DeSUMOylation of PINK1 increases mitophagy. (A) Schematic of GNb-SENP1 deSUMOylating PINK1-GFP. **(B)** GNb-SENP1 selectively deSUMOylates PINK1. HEK293T cells were transfected with GFP or PINK1-GFP along with GNb or GNb-SENP1 for 72 hours. Cells were treated with doxycycline (or DMSO control) for 48 hours before immunoprecipitation. Representative blots of higher molecular weight SUMO2/3 smear of immunoprecipitated PINK1 and corresponding input samples. **(C)** Quantification of higher molecular weight SUMO2/3 smear on PINK1 (SUMO2/3/GFP) (n=3, one-way ANOVA with Tukey's post-hoc test, **p<0.01) **(D)** Representative blots of total lysates from (B) immunoblotted for LC3A/B. **(E)** Quantification of LC3-II:LC3-I ratio (n=4, one-way ANOVA with Tukey's post-hoc test, *p<0.05, **p<0.01) **(F)** Representative images of PINK1$^{-/-}$ cells transfected with GFP or PINK1-GFP, mt-Keima, and GNb or GNb-SENP1. Cells were treated with 1 µg/mL doxycycline (or DMSO control) for 48 hours before imaging. Scale bar denotes 10 µm. **(G)** Quantification of "mitophagy index": fluorescence in 561 nm excitation channel divided by the total fluorescence in 405 nm and 561 nm excitation channels as a measure of "total mitophagy: total mitochondrial area" (C561/(C561 + C405)) (Kruskal-Wallis test followed by Dunn's multiple comparison test, N=3, n= 90-115 cells, ***p<0.001, ****p<0.0001)

Figure 7. Proposed model of PINK1 SUMOylation and PINK1-mediated mitophagy.

(A) When PINK1 is SUMOylated, it undergoes cleavage and is targeted for degradation, unaffecting basal mitophagy. **(B)** When PINK1 is not SUMOylated, PINK1 is stabilized, increasing levels of PINK1-mediated mitophagy. *Created in BioRender.*