Destabilization of Helix III Initiates Early Serum Amyloid A Misfolding by Exposing Its Amyloidogenic Core

Haidara Nadwa,[†] Z. Faidon Brotzakis,[*,‡,¶] Annalisa Santucci,[†] Daniela Braconi,[†] and Michele Vendruscolo[*,¶]

†Dipartimento di Biotecnologie, Chimica e Farmacia, Università degli Studi di Siena, Siena, Italy

‡Institute for Bioinnovation, Biomedical Sciences Research Center "Alexander Fleming", Vari, Greece

¶Centre for Misfolding Diseases, Department of Chemistry, University of Cambridge, Cambridge, United Kingdom

E-mail: brotzakis@fleming.gr; mv245@cam.ac.uk

Abstract

Serum amyloid A (SAA) is the principal precursor of AA amyloidosis, yet the early molecular steps that trigger its pathological misfolding remain unclear. Here, we combine high-temperature molecular dynamics simulations, harmonic linear discriminant analysis (HLDA), and parallel-tempering metadynamics (PT-MetaD) to dissect the earliest conformational transitions of the disease-relevant SAA_{1-76} fragment. By constructing an optimized one-dimensional collective variable (sHLDA) from inter-helix contacts and helical root-mean-square deviations, we perform nearly 4 μs of enhanced sampling across 79 replicas (300–450 K). Free-energy surfaces projected onto sHLDA and radius of gyration reveal a possible misfolding trajectory comprising thirteen metastable intermediates. We identify that helix III destabilizes first, preceding loss of helix II and helix I, while global compactness is retained. This stepwise mechanism underscores the key role of helix III in the establishment of the native fold of SAA. Solvent-accessible surface-area analyses pinpoint a transient surge in exposure of the aggregation-prone core (residues 42–48) within specific misfolded intermediates, implicating localized core exposure rather than wholesale unfolding as the trigger for misfolding. Temperature-dependent secondary-structure profiling further confirms that SAA_{1-76} behaves as a folded bundle with disordered loops, rather than a fully disordered protein. These insights identify helix III stabilization and masking of the amyloidogenic segment as strategic avenues for therapeutic intervention.

1 Introduction

AA amyloidosis is a severe systemic disorder and a life-threatening complication of chronic inflammation, characterized by the deposition of insoluble amyloid fibrils in various organs and tissues, leading to progressive dysfunction. This condition arises as a long-term complication of chronic inflammatory diseases, including persistent infections, rheumatoid arthritis, and certain cancers.[1,2] Recent findings also pointed out a link between alkaptonuria, an ultra-rare genetic disorder of tyrosine metabolism, persistent low-grade inflammation, and AA amyloidosis,[3–5] with a pro-aggregating effect of homogentisic acid (HGA) towards SAA highlighted in vitro.[6,7]

The principal component of AA amyloid deposits is the serum amyloid A (SAA) protein,[8] a family of acute-phase apolipoproteins primarily produced by the liver in response to inflammatory stimuli. SAA proteins are normally associated with high-density lipoprotein (HDL) in circulation and play a critical role in modulating the immune response.[9] SAA isoforms are acute-phase response proteins that are synthesized predominantly in the liver and expressed constitutively (constitutive SAA) or in response to inflammatory stimuli (acute phase SAA). SAA proteins are encoded by the SAA1, SAA2, SAA3, and SAA4 genes, located on chromosome 11p15.1 and expressed isoforms consist of 103-104 residues that are highly conserved throughout evolution.[10] SAA1 and SAA2 are the predominant acute-phase isoforms and their serum levels rise dramatically, up to 1000-fold, in response to infections, traumas, or other stimuli. The exact biological function of SAA is still partly understood with increasing evidence suggesting SAA roles in cholesterol transport, antibody regulation, inhibition of platelet aggregation, and modulation of macrophage activity.[11]

Structural studies, including the crystallographic analysis of the native human Serum Amyloid A-1 (SAA1) (PDB ID 4IP9) at 2.5 Å resolution.[9] suggest that SAA1 (104 a.a.) has a four-helix bundle structure with a cone-shaped array, in which the N termini of helices 1

and 3 and the C termini of helices 2 and 4 are packed together. The residues 1-27, 32–47, 50–69, and 73–88 form helices 1, 2, 3 and 4, respectively.[9] It has been reported that SAA helix bundle features a hydrophilic interior partially filled with water[12] and α-helices 1 and 3 feature a strong amphipathic property with a hydrophobic face.[13] The C-terminal tail forms multiple salt bridges and hydrogen bonds with the α-helices 1, 2 and 4 wrapping around the bundle plays a key role to stabilize the helix bundle structure.[9] Amphipathic helices 1 and 3 form an elongated concave hydrophobic surface with a curvature radius complementary to that of HDL suggesting the possible HDL binding site.[1] The formation of amyloid fibrils in AA amyloidosis occurs in the kidneys, spleen and liver,[14] stemming from the misfolding of circulating SAA1 after significant increment of serum SAA1 up to 1000 fold reaching $1 \, \mathrm{mg \, mL^{-1}}$ due to chronic inflammation.[15] This process begins with the dissociation from HDL, followed by truncation of C-terminal that generates an unstable $\approx$76 fragment (most commonly)[16] prone to misfolding and subsequent fibril formation.[14] However, the detailed misfolding mechanism leading to this pathological aggregation remains unclear.

Given the conflicting findings about the classification of SAA as an intrinsically disordered protein (IDP)[1] or a folded protein with only partially disordered segments[9], our work aims to clarify this debate by providing an atomistic description of the early events in the misfolding of SAA along and place them on the free energy landscape of the monomeric protein. We focused our study on the N-terminal 1–76 fragment of SAA ($\mathrm{SAA_{1-76}}$) that is the principal component of pathogenic fibrils. In AA amyloidosis, serum amyloid A undergoes a highly specific proteolytic event in its interdomain linker-between residues 76 and 77. Biochemical and mass-spectrometry analyses of ex vivo AA deposits have repeatedly shown that, although SAA is initially synthesized as a 104-residue acute-phase protein, the insoluble fibrillar core recovered from patient tissues consists almost exclusively of the first 76 amino acids. By simulating $\mathrm{SAA_{1-76}}$ rather than the full-length precursor, we therefore capture the true disease-relevant species and avoid introducing artifacts from a C-terminal tail that, in vivo, is removed prior to misfolding and aggregation. In addition, by quantifying the hydrophobic

residues and the aggregation-prone regions (APRs) exposure in intermediate conformational states, we investigate whether the aggregation propensity of SAA is driven primarily by its high concentration or by substantial exposure of APRs. To address these issues, we employ a supervised learning classification method (HLDA) to construct a collective variable for biasing simulations in PT-MetaD. This approach enables comprehensive exploration of the phase space of the system, providing free energy landscapes projected onto various variables and allowing us to characterize the structural ensembles corresponding to metastable states. Our main objective is to elucidate the early events in the misfolding of SAA, by probing the structural features of the identified metastable states and proposing a possible misfolding pathway.

2   Methods

2.1   Molecular Dynamics (MD)

The crystallographic structure of SAA with code 4IP9 in the Protein Data Bank (PDB) was used to derive the fragment SAA_{1-76} as a starting configuration after deleting residues 77-104.[9] This truncated SAA form is the most common in pathological amyloid deposits, due to the pro-amyloidogenic destabilization of the four helix bundle after removal of the C-terminal residues. .[8,9,17] All MD simulations were performed using the charmm36 force field[18] and GROMACS 2022.3 software package.[19] The protein was solvated in TIP3P water[20] and counter ions and additional salt ions (sodium and chloride) were added to neutralize the system and get a final salt concentration of 0.15 M. The system was then energy-minimized using the steepest descent method, with the simulations successfully terminating upon fulfill the energy criteria of a maximum force less than 1000 kJ/mol/nm. Followingly, an NPT equilibration at 310 K and 1 atm was performed for 100 ps, using the V-rescale thermostat[21] and Parrinello-Rahman barostat[22]. The particle-mesh Ewald method[23] was used for long-range electrostatics with a short-range cutoff of 1.0 nm. A cutoff of 1.0 nm was used for the Lennard-Jones interactions. All bonds were constrained to their equilibrium length with the LINCS algorithm[24]. A 2 fs timestep was used. Finally, two 20 ns NVT simulations were performed at 310 K and 500 K respectively.

2.2   Devising Collective Variables

According to Ref. 25, the Harmonic Linear Discriminant Analysis (HLDA) has been used to construct a 1-D collective variable (CV) that can describe the SAA folding process. This is a modification of Fisher's linear discriminant analysis (LDA) that estimates the optimal dimensional projection W to achieve maximum separation for the unbiased distributions of the folded and unfolded states within an N_d-dimensional descriptor space. In HLDA, the optimization of W is performed by maximizing the ratio of the so-called between-class scatter

(S_b) to the within-class scatter (S_w), expressed as:

$$J(W) = \frac{W^T S_b W}{W^T S_w W}, \tag{1}$$

where S_b quantifies the separation of the mean values of the two classes (folded and unfolded states) and is defined as:

$$S_b = (\mu_A - \mu_B)(\mu_A - \mu_B)^T, \tag{2}$$

with μ_A and μ_B representing the mean values of the descriptors for the folded and unfolded states, respectively.

In contrast, S_w measures the spread within each class. HLDA employs the harmonic average of the covariance matrices, defined as:

$$S_w = \left(\frac{1}{\Sigma_A} + \frac{1}{\Sigma_B}\right)^{-1}, \tag{3}$$

where Σ_A and Σ_B are the covariance matrices of the folded and unfolded states. Substituting these definitions into **Equation** 1, the optimization problem becomes the maximization of $J(W)$:

$$J(W) = \frac{W^T S_b W}{W^T S_w W}. \tag{4}$$

The optimal direction W^* is then determined by solving the eigenvalue problem:

$$W^* = S_w^{-1}(\mu_A - \mu_B). \tag{5}$$

The resulting collective variable, denoted as $\mathbf{s}_{\text{HLDA}}(\mathbf{R})$, is computed as the projection of the descriptors $d(R)$ along the optimal direction W^*, corresponding to the highest eigenvalue in **Equation** 5, formulated as:

$$s_{\text{HLDA}}(R) = (\mu_A - \mu_B)^T \left(\frac{1}{\Sigma_A} + \frac{1}{\Sigma_B} \right) d(R). \tag{6}$$

This approach is able to systematically construct a CV capable of distinguishing between folded and unfolded states, providing a one-dimensional representation of the complex folding dynamics.

Based on our intuition, we chose a set of six descriptors which can potentially describe the misfolding process of SAA. The first set probes the alpha helical content of each α-helix of SAA (d1, d2, d3) (αRMSD-1, αRMSD-2, and αRMSD-3). The second set consisted of three distances between the centers of each α-helix calculated and transformed by a switching function to a contact map (d4, d5, d6) (CM-1, CM-2, and CM-3) as shown in **Figure 1a**. The descriptors were calculated along the folded and unfolded state unbiased trajectories using the open-source, community-developed PLUMED library.[26–29]

By applying HLDA, we estimated the hyperplanes that best distinguished the unbiased distributions of the folded and unfolded states within the space defined by each set of descriptors constructing a collective variables as linear combinations of the set of descriptors.

2.3 Parallel-Tempering Metadynamics (PTMetaD) Simulations

PTMetaD simulations[30] were performed in the NPT ensemble by starting from 79 equidistant time-interval frames from the 20 ns folded state NPT MD obtained in the previous step. We ran 79 replicas in a temperature range between 300 K to 450 K for 50 ns each and cumulative simulation time of 3.9 μs. Each replica was biased on the previously defined CV. The Gaussian height and width were 3.0 KJ/mol and 0.02, respectively, and a bias factor of 8 and a Gaussian deposition rate of 100 were used in all simulations. The MD and force field parameters were the same as in the unbiased NPT MD.

3   Results and Discussion

3.1   Optimization of Collective Variables

After HLDA optimization, we selected the eigenvectors corresponding to the highest eigen-value, as shown in **Table** 1. This choice ensures the maximum separation between the folded and unfolded states. The analysis of the weight distributions, illustrated in **Figure 1b**, provides structural insights into the system, indicating that the features of the folding process were captured by the CVs. Notably, the majority of the weight is attributed to the descriptors contact map-1 (CM-1), which represents the distance between α-Helix I and α-Helix II, followed by the αRMSD values of α-Helix III and α-Helix II. The optimized HLDA CV, namely the one with higher eigenvalue was defined as:

$$s_{\text{HLDA}}(R) = (-0.026 * d1) + (-0.406 * d2) + (-0.540 * d3)$$
$$+(-0.735 * d4) + (-0.021 * d5) + (-0.043 * d6). \tag{7}$$

Table 1: Eigenvalues and their corresponding eigenvectors.

Eigenvalues	Eigenvectors					
30406.110	-0.026	-0.406	-0.540	-0.735	-0.021	-0.043
7.581e-14	0.003	0.003	-0.039	0.978	0.099	0.180
-5.120e-16	-0.001	-0.003	0.022	-0.270	-0.954	0.128
-4.380e-14	-0.002	-0.001	0.028	-0.946	0.111	0.303
-3.638e-12	-0.624	0.317	0.422	0.575	0.016	0.033
-4.490e-12	-0.312	-0.410	0.546	0.661	0.011	-0.012

(a)

(b)

Figure 1: (a) Cartoon structure of SAA_{1-76} with the three α-helices shown in red; the distances between the centers of mass of the α-helices used as descriptors are also indicated. (b) Weights assigned by HLDA to each of the descriptors.

3.2 SAA misfolding mechanism

Using the HLDA derived CV, $\mathbf{s}_{\mathrm{HLDA}}(\mathbf{R})$, we performed PT-MetaD simulations to explore the misfolding mechanism of SAA, in terms of different free energy landscape projections. The overlap of potential energy distributions across consecutive replicas, as illustrated in **Figure S1**, highlights the efficiency of the sampling and exchange rate between replicas. Additionally, the diffusion of the HLDA CV at T = 310 K over time (**Figure 2**) under the effect of the PT-MetaD potential demonstrates increased fluctuations reflecting enhanced exploration of phase space and improved sampling. The simulations convergence is confirmed through the effective diffusion of replica 06 (at a temperature of 310 K) across the temperature space (shown in **Figure S2**) along with the superposition of the time-dependent FES along various CVs (**Figures S3, S4, S5**) where the FES becomes static as a function of time.

From the MD simulations we obtained two 2D free energy landscapes projected on HLDA CV and αRMSD for the first one (**Figure 3a**), while for the second one is on αRMSD and

radius of gyration (Rg) (**Figure 3b**). The possible misfolding process is represented as a pathway connecting two basins: the initial folded structure and the final unfolded state, crossing a series of metastable minima. Using the MEPSA (Minimum Energy Path Surface Analysis) software[31], we identified the minimum energy pathways as a function of collective variables we deemed informative on the misfolding mechanism. We would like to stress these pathways are neither kinetic pathways nor necessarily describe the committor function but are rather informative pathways along FES projections. These pathways are illustrated as a black line in the first free energy surface plot (**Figure 3a**) and as three distinct colored lines in the second FES plot (**Figure 3b**), passing through a series of basins that represent configurations sampled during the simulations. Thirteen distinct metastable states along the possible misfolding pathway were identified in the first FES (**Figure 3a**), labeled 1 to 13. Representative conformations from these basins are shown in panel A (**Figure 3c**), where the conformer in minimum 1 corresponds to a fully folded structure, while metastable state 13 represents an almost fully unfolded structure forming up to 25% of helical content with respect to the folded state structure.

Figure 2: HLDA CV time dependence taken from T=310K replica of PTMetaD simulations.

Figure 3: (a) FES as a function of αRMSD and HLDA showing a possible misfolding pathway. (b) FES as a function of the αRMSD and Rg showing three possible misfolding pathways and the 14 minima crossed by them. (c) A: Structures of the 13 minima crossed along the possible misfolding pathway shown in (a). B-D: Representative structures sampled during the simulations along the first, second and third possible misfolding pathways. Common structures found in shared basins among these pathways are enclosed within dotted boxes

To gain deeper insight into the structural changes along the possible misfolding pathway, we analyzed the α-helical content of the three α-helices in each of the nine minima illustrated in **Figure 4a**. The results indicate that α-Helix III unfolds first, followed by a sudden dissolution of α-Helix II after metastable state 5. The last helix to unfold is α-Helix I, which starts to decay after metastable state 6, with half of it disappearing by metastable state 13. The order of α-helix unfolding aligns with the weights of the descriptors shown in **Figure 1b**. As shown in **Figure 4b**, the overall α-helical content gradually decreases along the misfolding pathway, accompanied by an increase in random coil structure, ultimately reaching an almost fully unfolded state in metastable state 13. Rg analysis of structures in

Figure 4c suggests that, despite the loss of secondary structure, the compactness of the fragment is not significantly affected. This observation is further supported by the contact map analysis (**Figure 4d**). Although there are fluctuations in the distance between the α helices, the distances between the centers of mass of the three α-helices remain relatively stable as we can notice in the metastable 13 with respect to the folded state structure.

Along the misfolding pathway, we observe that the α-helices gradually transform into random coil in a specific sequence while maintaining structural compactness and preserving the tertiary structure. This process results in a structure composed almost entirely of random coil with some remaining α-helical content, yet the compactness remains intact. These findings suggest that the likely misfolding mechanism of SAA is consistent with the nucleation–condensation model of folding.[32] According to this model, the folding process begins with the formation of a marginally stable nucleus, due to a few correct secondary and tertiary structure interactions, acting as a template and enabling the rapid condensation of adjacent structural elements.

Figure 4: (a) Alpha helical content of SAA as measured for each representative structure sampled during the simulation. (b) Secondary structure population for each representative structure sampled during the simulation. (c) The radius of gyration for each representative structure sampled during the simulation. (d) The distances between the centers of mass of each α helix, calculated as contact map, for each representative structure sampled during the simulation.

The solvent-accessible surface area (SASA) analysis (**Figure S6**) along the possible misfolding pathway shows that the total SASA does not increase significantly along the unfolding pathway at the misfolded intermediate states. Similarly, the SASA of the hydrophobic residues (**Figure S6**) remain relatively stable during the misfolding. However, the SASA of the predicted aggregation-prone region (APR, corresponding to residues **42-48** as predicted by FuzDrop[33–35]) (**Figure 5a**) remain relatively stable during the unfolding except the basins 9, 10, 11 and 12 in which the APR region becomes more exposed to the solvent.

These results suggest that the aggregation of SAA can be triggered by the increased solvent exposure of the APR of the aforementioned metastable states.

(a)

(b)

Figure 5: (a) Total SASA for the APR for the representative structure sampled during the simulations along aRMSD and HLDA-CV. (b) Total SASA for the APR along the first (un)folding path along aRMSD and Rg.

Figure 3b presents a 2D FES plotted as a function of α-RMSD and Rg, showing three possible misfolding pathways represented by black, red and yellow dotted lines, respectively. The first proposed pathway crosses eight basins, with the conformer in minimum 1 corresponding to a fully folded structure and basin 8 representing an almost fully unfolded structure as shown in **Figure 3c**, panel B. This pathway is aligned with the projected one along α-RMSD and HLDA-CV that have been discussed before, however, it is less tuned as we can see in **Figure 3b**. The basins 2, 3, 4 and 5 in the 2D FES along α-RMSD and HLDA-CV are gathered in basin 2 in the 2D FES along α-RMSD and Rg. Similarly, basins 7, 8 and 9 in the 2D FES along α-RMSD and HLDA-CV are gathered in basin 4 in the 2D FES along α-RMSD and Rg.

The results of the α-helicity analysis align with previous results, showing that α-Helix III unfolds first, followed by α-Helix II, and finally α-Helix I, as illustrated in **Figure 6a**. Along this pathway, the overall α-helical content decreases steadily while the random coil content increases, ending with a nearly fully unfolded structure in basin 8 as shown in **Figure 6b**. Despite the loss of secondary structure, the compactness of the protein remains stable throughout the pathway, as revealed by the Rg analysis in **Figure 6c**. The contact map analysis further supports this, demonstrating that the distances between the centers of mass of the α-helices remain relatively unchanged in basin 8 compared to basin 1. However, in

metastable states 5, 6 and 7, there is a marked increase in the distances between the α helices which align with the contact map analysis of the 2D FES along αRMSD and HLDA-CV where we can notice the increase of distances in basins 10, 11, and 12 (**Figure 4d**) which are the same metastable states in both pathways (**Figure 3c**).

Figure 6: (a) Alpha helical content of SAA as measured for each representative structure sampled during the simulation along the first suggested unfolding pathway. (b) secondary structure population for each representative structure sampled during the simulation along the first suggested unfolding pathway. (c) The radius of gyration for each representative structure sampled during the simulation along the first suggested unfolding pathway. (d) The distances between the centers of mass of each α helix, calculated as contact map, for each representative structure sampled during the simulation along the first suggested unfolding pathway.

SASA analysis reveals that the overall solvent-exposure area and the SASA attributed to hydrophobic residues remains largely unchanged across the misfolded metastable states (**Figure S7**). However, the SASA attributed to the APR increased significantly during unfolding in the basins 4, 5, 6 and 7 (**Figure 5b**) which are the same basins 9, 10, 11 and 12 in the 2D FES along αRMSD and HLDA-CV that show a significant increase of

SASA attributed to APR (**Figure 5a**). These findings support the hypothesis that SAA aggregation can be driven by an increased solvent exposure of APRs in these previously mentioned metastable states.

The second identified pathway represented by red dotted line in **Figure 3b** crosses nine metastable states but does not lead to a fully unfolded structure. The conformer in metastable state 11 remains partially unfolded, as shown in **Figure 3c** Panel C sharing the first 6 basins with the first pathway.

The α-helicity analysis reveals that α-Helix III unfolds completely first, followed by α-Helix II, and finally α-Helix I within the first six metastable states, which overlap with the initial part of the first suggested pathway. However, in the rest three basins (9, 10 and 11), α-Helix I remains stable while α-Helix-II increases slightly as we can see in **Figure 7a**. Additionally, **Figure 7b** shows a steady decrease in overall α-helical content and a corresponding increase in random coil content up to basin 5, after which a significant increase occurs in the last four basins ending with a structure forming up to 50% of helical content with respect to the folded state structure.

The Rg analysis indicates a significant increase in the last four metastable states, signifying a loss of protein compactness, as shown in **Figure 7c**. These findings are supported by the contact map analysis (**Figure 7d**), which reveals a significant increase in the distances between the α-helices, particularly between α-Helix I and II and between α-Helix II and III.

Figure 7: (a) Alpha helical content of SAA as measured for each representative structure sampled during the simulation along the second suggested unfolding pathway. (b) Secondary structure population for each representative structure sampled during the simulation along the second suggested unfolding pathway. (c) The radius of gyration for each representative structure sampled during the simulation along the second suggested unfolding pathway. (d) The distances between the centers of mass of each α helix for each representative structure sampled during the simulation along the second suggested unfolding pathway.

Similar observations are noticed in the third suggested pathway represented by yellow dotted line in **Figure 3b**, which passes from metastable state 1 to metastable state 14. In this pathway, the α-helices I and II were stable while the α-helix III content decreases slightly (**Figure 8a**) ending with a structure that has around 80% of helical content with respect to the folded state structure as shown in **Figure 8b**. However, the protein's compactness decreases, as evidenced by an increase in the Rg value (**Figure 8c**), and the distance between α-Helix I and III and between α-Helix II and III also increase as illustrated in **Figure 8d**.

Figure 8: (a) Alpha helical content of SAA as measured for each representative structure sampled during the simulation along the third suggested unfolding pathway. (b) Secondary structure population for each representative structure sampled during the simulation along the third suggested unfolding pathway. (c) The radius of gyration for each representative structure sampled during the simulation along the third suggested unfolding pathway. (d) The distances between the centers of mass of each α helix, calculated as contact map, for each representative structure sampled during the simulation along the third suggested unfolding pathway.

The final two off-pathways correspond to the second generic protein misfolding mechanism: the diffusion–collision mechanism.[36] In this model, local elements of secondary structure form first, then diffuse, collide, and adhere to establish the correct tertiary structure. However, for the analyzed SAA fragment, this mechanism appears improbable, as the pathways fail to reach a fully unfolded structure, further corroborating the hypothesis of a nucleation-condensation model. Additionally, the total SASA (**Figure S8**) and the SASA of the APR (**Figure 9a**) increase along the second pathway suggesting that the increased exposure of the APR drives the aggregation process. In contrast, pathway 3 shows no sig-

nificant change in total SASA and hydrophobic SASA (**Figure S9**), or APR SASA (**Figure 9b**). However, since these two off-pathways do not lead to complete unfolding, these results assert once again the hypothesis that SAA aggregation process is driven by the APR solvent exposure.

(a)

 (b)

Figure 9: (a) Total SASA for the APR along the second misfolding path. (b) Total SASA for the APR along the third misfolding path.

There have been controversial experimental findings about the nature of SAA as IDP[1] or folded protein with only partially disordered segments.[9] Our temperature-dependent secondary structure analysis of SAA_{1-76} structural ensembles, as shown in **Figure 10**, indicate that SAA_{1-76} is not an IDP. This conclusion is based on the characteristic behavior of IDPs, which become more ordered structures at higher temperatures. In contrast, we observed that SAA_{1-76} adopts highly unstructured conformations (predominantly random coil) at elevated temperatures, while structured conformations are more prominent at lower temperatures.

Figure 10: The secondary structure analysis of SAA over different temperatures

4   Conclusions

In this study, we explored the early misfolding steps of the pro-amyloidogenic 1-76 fragment of SAA (SAA_{1-76}), which is the principal component of pathogenic fibrils, by combining a range of molecular dynamics simulations methods.

By implementing Harmonic Linear Discriminant Analysis (HLDA) to construct collective variables (CVs) and utilizing Parallel-Tempering Metadynamics (PT-MetaD), we mapped the free energy landscapes of the monomeric state of SAA_{1-76}, identifying key metastable states along its possible misfolding pathway. Notably, by utilizing HLDA with a priori set of descriptors, we generated a linear collective variable that drives the PT-MetaD simulation toward the unfolded state providing free energy estimates. Furthermore, the weight distributions of the calculated HLDA CV reveals useful information, offering physical insight into the misfolding process. The simulations revealed a sequential misfolding process, where α-helix III unfolds first, followed by α-helix II and finally α-helix I. Despite the loss of secondary structure, the overall compactness of SAA remained largely intact. Moreover, SASA analyses show that the overall SASA, as well as that of hydrophobic residues remain relatively constant along the misfolding pathway at the misfolded intermediate states. However, the SASA of the predicted APR (residues 42–48) increases in a specific metastable states. This observation implies that SAA aggregation can be driven by the significant increase in the solvent exposure of APRs. Furthermore, the secondary structure analysis of SAA across different temperatures reveals that, contrary to the typical behavior of IDPs, which tend to become more ordered at elevated temperatures, SAA shows predominantly unstructured, random coil conformations at higher temperatures, while more structured conformations are evident at lower temperatures.

These results provide insights into the molecular basis of SAA misfolding and identify metastable states that could serve as potential targets for therapeutic intervention, such

as strategies aim at stabilizing native α-helical regions. The approach presented here enhances our understanding of SAA aggregation and offers a framework for studying the early misfolding events of other amyloidogenic proteins. Future research should focus on experimental validation of these findings and on exploring small-molecule misfolding inhibitors, known as pharmacological chaperones[37,38], that stabilize the native structure of SAA to mitigate its pathogenic aggregation.

Acknowledgement

H.N acknowledges the Italian Ministry of University and Research (MUR) for a "Bando PON 37 ciclo - REACT-EU FSE DM 1061" grant. H.N acknowledges the CINECA award under the ISCRA initiative for the availability of high-performance computing resources and support. D.B. acknowledges support from the University of Siena (grant PSR 2021 F-CUR).

Supporting Information Available

The overlap of potential energy probability distributions across consecutive replicas is illustrated in **Figure** S1. The effective diffusion of replica 06 (at a temperature of 310 K) across the temperature space is shown in **Figure** S2. The superposition of the time-dependent FES along αRMSD, HLDA CV and Rg are illustrated in **Figures** S3, S4, S5, respectively. Total SASA for the protein and for the hydrophobic residues for the representative structure sampled during the simulations along αRMSD and HLDA CV is shown in **Figures** S6. Total SASA for the protein and for the hydrophobic residues for the representative structure sampled during the simulations along the first, second and third misfolding path along αRMSD and Rg are illustrated in **Figures** S7, S8, S9, respectively.

References

(1) Frame, N. M.; Gursky, O. Structure of serum amyloid A suggests a mechanism for selective lipoprotein binding and functions: SAA as a hub in macromolecular interaction networks. *FEBS letters* **2016**, *590*, 866–879.

(2) Pettersson, T.; Konttinen, Y. T.; Maury, C. P. J. Treatment strategies for amyloid A amyloidosis. *Expert opinion on pharmacotherapy* **2008**, *9*, 2117–2128.

(3) Bernini, A.; Spiga, O.; Santucci, A. Structure-function relationship of homogentisate 1, 2-dioxygenase: understanding the genotype-phenotype correlations in the rare genetic disease alkaptonuria. *Current Protein and Peptide Science* **2023**, *24*, 380–392.

(4) Braconi, D.; Geminiani, M.; Psarelli, E. E.; Giustarini, D.; Marzocchi, B.; Rossi, R.; Bernardini, G.; Spiga, O.; Gallagher, J. A.; Le Quan Sang, K.-H.; others Effects of nitisinone on oxidative and inflammatory markers in alkaptonuria: results from SONIA1 and SONIA2 studies. *Cells* **2022**, *11*, 3668.

(5) Braconi, D.; Giustarini, D.; Marzocchi, B.; Peruzzi, L.; Margollicci, M.; Rossi, R.; Bernardini, G.; Millucci, L.; Gallagher, J. A.; Sang, K.-H. L. Q.; others Inflammatory and oxidative stress biomarkers in alkaptonuria: data from the DevelopAKUre project. *Osteoarthritis and cartilage* **2018**, *26*, 1078–1086.

(6) Braconi, D.; Millucci, L.; Bernini, A.; Spiga, O.; Lupetti, P.; Marzocchi, B.; Niccolai, N.; Bernardini, G.; Santucci, A. Homogentisic acid induces aggregation and fibrillation of amyloidogenic proteins. *Biochimica et Biophysica Acta (BBA)-General Subjects* **2017**, *1861*, 135–146.

(7) Mastroeni, P.; Trezza, A.; Geminiani, M.; Frusciante, L.; Visibelli, A.; Santucci, A. HGA Triggers SAA Aggregation and Accelerates Fibril Formation in the C20/A4 Alkaptonuria Cell Model. *Cells* **2024**, *13*, 1501.

(8) Sack Jr, G. H. Serum amyloid A–a review. *Molecular medicine* **2018**, *24*, 46.

(9) Lu, J.; Yu, Y.; Zhu, I.; Cheng, Y.; Sun, P. D. Structural mechanism of serum amyloid A-mediated inflammatory amyloidosis. *Proceedings of the National Academy of Sciences* **2014**, *111*, 5189–5194.

(10) Sun, L.; Richard, D. Y. Serum amyloid A1: structure, function and gene polymorphism. *Gene* **2016**, *583*, 48–57.

(11) Kisilevsky, R.; Manley, P. N. Acute-phase serum amyloid A: perspectives on its physiological and pathological roles. *Amyloid* **2012**, *19*, 5–14.

(12) Smole, U.; Kratzer, B.; Pickl, W. F. Soluble pattern recognition molecules: guardians and regulators of homeostasis at airway mucosal surfaces. *European journal of immunology* **2020**, *50*, 624–642.

(13) Hu, Z.; Bang, Y.-J.; Ruhn, K. A.; Hooper, L. V. Molecular basis for retinol binding by serum amyloid A during infection. *Proceedings of the National Academy of Sciences* **2019**, *116*, 19077–19082.

(14) Rennegarbe, M.; Lenter, I.; Schierhorn, A.; Sawilla, R.; Haupt, C. Influence of C-terminal truncation of murine Serum amyloid A on fibril structure. *Scientific Reports* **2017**, *7*, 6170.

(15) Sipe, J. D.; Benson, M. D.; Buxbaum, J. N.; Ikeda, S.-i.; Merlini, G.; Saraiva, M. J.; Westermark, P. Amyloid fibril proteins and amyloidosis: chemical identification and clinical classification International Society of Amyloidosis 2016 Nomenclature Guidelines. *Amyloid* **2016**, *23*, 209–213.

(16) Uhlar, C. M.; Whitehead, A. S. Serum amyloid A, the major vertebrate acute-phase reactant. *European journal of biochemistry* **1999**, *265*, 501–523.

(17) Yamada, T.; Liepnieks, J.; Kluve-Beckerman, B.; Benson, M. Cathepsin B generates the most common form of amyloid A (76 residues) as a degradation product from serum amyloid A. *Scandinavian journal of immunology* **1995**, *41*, 94–97.

(18) Best, R. B.; Zhu, X.; Shim, J.; Lopes, P. E.; Mittal, J.; Feig, M.; MacKerell Jr, A. D. Optimization of the additive CHARMM all-atom protein force field targeting improved sampling of the backbone ϕ, ψ and side-chain $\chi 1$ and $\chi 2$ dihedral angles. *Journal of chemical theory and computation* **2012**, *8*, 3257–3273.

(19) Páll, S.; Abraham, M. J.; Kutzner, C.; Hess, B.; Lindahl, E. Tackling exascale software challenges in molecular dynamics simulations with GROMACS. Solving Software Challenges for Exascale: International Conference on Exascale Applications and Software, EASC 2014, Stockholm, Sweden, April 2-3, 2014, Revised Selected Papers 2. 2015; pp 3–27.

(20) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. *The Journal of chemical physics* **1983**, *79*, 926–935.

(21) Bussi, G.; Donadio, D.; Parrinello, M. Canonical sampling through velocity rescaling. *The Journal of chemical physics* **2007**, *126*.

(22) Parrinello, M.; Rahman, A. Polymorphic transitions in single crystals: A new molecular dynamics method. *Journal of Applied physics* **1981**, *52*, 7182–7190.

(23) Darden, T.; York, D.; Pedersen, L. Particle mesh Ewald: An N log (N) method for Ewald sums in large systems. *The Journal of chemical physics* **1993**, *98*, 10089–10092.

(24) Hess, B. P-LINCS: A parallel linear constraint solver for molecular simulation. *Journal of chemical theory and computation* **2008**, *4*, 116–122.

(25) Mendels, D.; Piccini, G.; Brotzakis, Z. F.; Yang, Y. I.; Parrinello, M. Folding a small protein using harmonic linear discriminant analysis. *The Journal of chemical physics* **2018**, *149*.

(26) Bonomi, M.; Branduardi, D.; Bussi, G.; Camilloni, C.; Provasi, D.; Raiteri, P.; Donadio, D.; Marinelli, F.; Pietrucci, F.; Broglia, R. A.; others PLUMED: A portable plugin for free-energy calculations with molecular dynamics. *Computer Physics Communications* **2009**, *180*, 1961–1972.

(27) Tribello, G. A.; Bonomi, M.; Branduardi, D.; Camilloni, C.; Bussi, G. PLUMED 2: New feathers for an old bird. *Computer physics communications* **2014**, *185*, 604–613.

(28) Promoting transparency and reproducibility in enhanced molecular simulations. *Nature methods* **2019**, *16*, 670–673.

(29) Tribello, G. A.; Bonomi, M.; Bussi, G.; Camilloni, C.; Armstrong, B. I.; Arsiccio, A.; Aureli, S.; Ballabio, F.; Bernetti, M.; Bonati, L.; others PLUMED Tutorials: a collaborative, community-driven learning ecosystem. *arXiv preprint arXiv:2412.03595* **2024**,

(30) Bussi, G.; Gervasio, F. L.; Laio, A.; Parrinello, M. Free-energy landscape for β hairpin folding from combined parallel tempering and metadynamics. *Journal of the American Chemical Society* **2006**, *128*, 13435–13441.

(31) Marcos-Alcalde, I.; Setoain, J.; Mendieta-Moreno, J. I.; Mendieta, J.; Gomez-Puertas, P. MEPSA: minimum energy pathway analysis for energy landscapes. *Bioinformatics* **2015**, *31*, 3853–3855.

(32) Abkevich, V. I.; Gutin, A. M.; Shakhnovich, E. I. Specific nucleus as the transition state for protein folding: evidence from the lattice model. *Biochemistry* **1994**, *33*, 10026–10036.

(33) Hardenberg, M.; Horvath, A.; Ambrus, V.; Fuxreiter, M.; Vendruscolo, M. Widespread occurrence of the droplet state of proteins in the human proteome. *Proceedings of the National Academy of Sciences* **2020**, *117*, 33254–33262.

(34) Vendruscolo, M.; Fuxreiter, M. Sequence determinants of the aggregation of proteins within condensates generated by liquid-liquid phase separation. *Journal of Molecular Biology* **2022**, *434*, 167201.

(35) Hatos, A.; Tosatto, S. C.; Vendruscolo, M.; Fuxreiter, M. FuzDrop on AlphaFold: visualizing the sequence-dependent propensity of liquid–liquid phase separation and aggregation of proteins. *Nucleic acids research* **2022**, *50*, W337–W344.

(36) Karplus, M.; Weaver, D. L. Protein folding dynamics: The diffusion-collision model and experimental data. *Protein science* **1994**, *3*, 650–668.

(37) Cohen, F. E.; Kelly, J. W. Therapeutic approaches to protein-misfolding diseases. *Nature* **2003**, *426*, 905–909.

(38) Vendruscolo, M. The thermodynamic hypothesis of protein aggregation. *Molecular Aspects of Medicine* **2025**, *103*, 101364.