The signature of reproduction on wood anatomy in European beech reveals potential for masting reconstruction

Data Availability Statement

Our Mandates Data Policy requires data to be shared and a Data Availability Statement, so please enter one in the space below. Sample statements can be found here. Please note that this statement will be published alongside your manuscript, if it is accepted for publication.

The data that support the findings of this study are available from the corresponding author, GR, upon request.

Title

2 The signature of reproduction on wood anatomy in European beech reveals potential for

3 masting reconstruction

List of authors

5 Giulia Resente[1], Alan Crivellaro[1,2], Emilie Fleurot[1], Alma Piermattei[1,2], Francesco Maimone[1], Martin

6 Wilmking[3], Andrew Hacket-Pain[4], Renzo Motta[1], Davide Ascoli[1]

Institutional affiliation

1 Department of Agricultural, Forest and Food Sciences, University of Torino, Grugliasco (TO), Italy.

2 Forest Biometrics Laboratory, Faculty of Forestry, "Stefan cel Mare" University of Suceava, Suceava,

Romania.

3 Institute of Botany and Landscape Ecology, University Greifswald, Greifswald, Germany.

4 Department of Geography and Planning, School of Environmental Sciences, University of Liverpool,

Liverpool, UK

Contact information

15 Giulia Resente, +39 3464307457, giuliaantonia.resente@unito.it

Word count (Introduction, Materials and Methods, Results, Discussion)

17 7067

Number of figures

19 5 (Fig. 1-4 should be included with colors)

Summary

1) We investigated the potential of wood anatomical traits to improve the reconstruction of masting

events—variable and synchronized patterns of seed production —which are key to understanding

tree species' responses to current and predicted climate variability. Traditional reliance on tree-

ring width as a proxy for reproduction is limited, as growth reductions can also result from drought

and other stressors.

2) We analyzed 12 beech cores from North-East Germany, building a 52-year dataset. A wide range

of wood anatomical traits was assessed to disentangle the effects of masting and drought. We

used multivariate regression and developed a random forest model to evaluate the predictive

power of these traits compared to tree ring width alone.

3) Results suggested a complex mechanism of carbon reallocation towards reproduction, while

reflecting a compensatory strategy to maintain hydraulic function and mechanical stability under

resource-constrained conditions. Number of parenchyma cells, vessel density, and lignin content

estimates emerged as key predictors for masting, outperforming tree-ring width in capturing the

reproductive signal.

4) Our findings establish a novel link between wood anatomy and masting events, demonstrating

that quantitative wood anatomical traits offer a more accurate and ecologically relevant approach

for reconstructing past masting dynamics.

Keywords (5 – 8)

"mast seeding", "dendroecology", "climate change", "quantitative wood anatomy",

"reconstruction", "beech", "reproduction", "drought".

Introduction

Masting, or mast seeding, refers to the reproduction strategy where plant populations exhibit highly variable and synchronized patterns of seed production (Pearse et al., 2021). While masting enhances reproductive fitness by increasing reproduction efficiency via economies of scale (Kelly and Sork, 2002), the associated demands of massive reproductive effort also affect plant growth and carbon stocks. These changes in resource allocation trigger major cascading ecological effects (Bogdziewicz et al., 2021) involving several ecosystem aspects, such as soil nutrient dynamics (Brumme et al., 2021), sporocarps fruiting reduction (Michaud et al., 2024), and shifts in animal population dynamics due to fluctuating food availability (Ostfeld et al., 2000).

Climate variation influences masting patterns on an inter-annual and decadal scale (Ascoli et al., 2021). Rising temperatures due to climate change have been influencing masting occurrence and changing masting patterns (Foest et al., 2024), with strong negative effects on seed production (Bogdziewicz et al., 2023). However, our understanding of the effects of the ongoing climate change on masting patterns is limited by the length of the available time series of masting, which are often too short for a comparison of their inter-annual variability with the fluctuations and decadal trends in climate. Consequently, the limited availability of long-term observations of masting prevents the evaluation of recent trends against long-term variability in masting (i.e., baseline variability) (Prebble et al., 2022; Wion et al., 2025). Additionally, available data is strongly biased geographically and taxonomically, with long-term records restricted to regions and species valued in forestry (Hacket-Pain et al., 2022).

Reconstructing past masting patterns is essential for understanding how plant reproductive behavior responds to long-term climate variability. By identifying how masting variability shifts across climatic phases, researchers can better anticipate the ecological consequences of ongoing climate change (Mundo et al., 2021; Prebble et al., 2022). In woody species such as Araucaria (*Araucaria araucana* (Molina) K. Koch), Oak (*Quercus robur* L.), and European beech (*Fagus sylvatica* L.), most reconstruction efforts to date have relied on tree-ring width as a proxy for reproductive investment (Drobyshev et al., 2014; Hacket-Pain et al., 2024; Koenig et al., 2020; Mundo et al., 2021). European beech in particular has emerged as a relevant species in masting research, with numerous studies demonstrating reduced growth during mast years, supporting the hypothesis of a direct growth–reproduction trade-off (Bajocco et al., 2021; Bogdziewicz et al., 2021; Drobyshev et al., 2014; Hacket-Pain et al., 2015; Nussbaumer et al., 2021; Stolz et al., 2023).

Despite the established use of growth rings for reconstruction purposes, the accuracy of dendrochronology remains limited by the challenge of disentangling the effects of masting from other

radial growth-reducing factors such as drought, late frost, or insect outbreaks (Drobyshev et al., 2014;

Hacket-Pain et al., 2015; Hadad et al., 2021). In beech, for instance, drought can produce similar

reductions in tree ring width as masting (Ding et al., 2017; Piovesan et al., 2008; Scharnweber et al.,

2011). However, since masting is primarily driven by internal carbon and nutrient dynamics (Han et al.,

2011; Han and Kabeya, 2017; Pearse et al., 2016), its physiological footprint may extend beyond radial

growth alone. Quantitative wood anatomy measures anatomical traits at an intra-annual level,

providing insights into tree potential hydraulic conductance (e.g., lumen area, hydraulic diameter,

specific conductivity, vessel density), and in relation to carbon and nutrient storage (e.g., tree ring

width, earlywood and latewood width, cell wall thickness, degree of lignification, axial parenchyma

cells). For these reasons, specific anatomical traits may carry distinct signatures of reproductive

investment (Redmond et al., 2019), potentially allowing for a more reliable identification of mast years

and opening new possibilities for high-resolution, anatomically based masting reconstructions.

To evaluate the potential of quantitative wood anatomy, we tested the response of ring width and a

suite of wood anatomical traits to masting and drought aiming to disentangle the effects of both

events. Specifically, we hypothesize that quantitative wood anatomy could help in the identification of

a specific "masting signature", where masting exerts a clearer effect on the traits related to growth

and nutrient mobilization, rather than traits involved in plant hydraulics. This analysis laid the basis for

the reconstruction of past masting events, and a deeper understanding of the allocation trade-offs

associated with masting (Redmond et al., 2019). By successfully linking specific anatomical

signatures to masting, our approach represents a significant step forward in understanding long-term

reproductive strategies in trees and their interactions with environmental variability.

Material and Methods

Sampling site

101   Trees were sampled in the forest of Carpin in the North-East of Germany (Fig. **1**). The site belongs to

102   the Müritz National Park, which extends over the Mecklenburg Lake District. Specifically, the sampling

103   site (53.20 N; 13.14 E) is in proximity to the Schweingartensee lake, at an elevation of 80 meters above

104   sea level.

Figure 1. Geographic location of the study site. (A) *Map of Germany with a subset of the Mecklenburg Lake District (blue rectangle). (B) Red dots represent the sampling site located in the Serrahner Wald, close to the Schweingartensee, one of the lakes of the area.*

Sample collection and processing

In February 2022, we sampled twelve healthy, dominant, and fecund trees, collecting two cores per tree at breast height with a 5 mm increment borer. For every tree, GPS position, diameter at breast height, and height measurements were collected. Stem diameter ranged between 62 and 127 centimeters, tree height between 34 and 43 meters.

Cross-dating was performed using one core per tree producing individual tree ring width time series. In the laboratory, cores were mounted on a wooden support, sanded to highlight tree ring visibility, and then scanned with an A3 optical flat-bed scanner with a resolution of 2400 dpi. Tree ring width was then measured with CooRecorder, and crossdated with cDendro 9.3 version (Cybis Electronic 2013). Tree age, together with the collected field data, is reported in Table **S1**. The second core was used for wood anatomical analyses. Cores were cut into 3-4 cm sections and processed with a Leica RM 2245 rotary microtome to produce thin sections of 12 µm thickness. These were stained with a 1:1 Safranin and Astra Blue solution to highlight the lignin and cellulose content, and to improve contrast. Thin sections were mounted on a glass slide and sealed with Euparal mounting media and dried at 65–

123 70 °C for 48 h. Images of the slides were captured using a Zeiss Axio Scan.Z1 slide scanner (Carl Zeiss

124 AG, Germany) with a 10x lens and a resolution of 2.267574 pixel/μm.

Wood anatomical traits time series

Wood anatomical traits for each individual tree were measured for the period 1970-2021 with the

CARROT software (Resente et al., 2024, 2021) on the thin section images. CARROT contains models

for measuring tree ring width (TRW), tree ring area computation, and a pretrained model for beech

vessel identification. From these analyses, we derived the following traits: coefficient of variation of

the lumen area within each ring (CV LA), average hydraulic diameter (Dh), hydraulic diameters of the

biggest and the smallest decile of the distribution (Dh min, Dh max), specific conductivity (Ks), and

vessel density (VD). Furthermore, CARROT allowed for the creation of a specific model to detect and

analyze axial parenchyma cells through the inbuilt retraining function (available at

https://zenodo.org/uploads/15262231). This new model allowed the computation of parenchyma

cells lumen area (PC LA), number (PC N), and density (PC D). A summary of the abbreviation is

provided in Table **1**.

We accounted for the vessel widening effect from the pith to the bark due to increasing tree height and

crown size (Anfodillo et al., 2013; Lechthaler et al., 2019) as well as the age effect, detrending all the

calculated traits with a cubic smoothing spline function with a 50% frequency and a cut-off of 10

years. The optimal cut-off time window was established by selecting the highest R-bar measure

among each trait time series (Supporting Information Fig. **S1**). For these analyses we used the detrend

function of the dplR Package (Bunn, 2010) that calculates the ratio between the observed and fitted

values for each annual tree ring to obtain detrended series (Cook and Kairiukstis, 1990). The function

then averages the detrended series employing a bi-weight robust mean to build the mean anatomical

chronologies. To account for the ecological phenomenon of masting—which is driven by mechanisms

developing over multiple years—and to specifically capture the relevance of interannual temperature

variation (i.e., the temperature difference from the previous year) (Vacchiano et al., 2017), we applied

a first-order differencing method to detrend the wood anatomical series.

In addition to the wood anatomical traits, we used the methodology described in Resente et al. (in

review) to compute the Red Intensity (RI) profile of each tree. RI quantifies the red component in thin

sections resulting from the staining process with Safranin and is a proxy for the lignin content in each

growth ring averaged by year in order to obtain one data point per year of the chronology. RI was

normalized using the slide identification number as a grouping factor and dividing each tree ring RI for

the average RI value of the slide. This allowed to avoid bias created by the single thin section

thickness. Anatomical Wood Density (AWD) was calculated with the methodology proposed in

Kollmann and Côté, (1968) where the percentage of white pixels corresponding to cell lumina was

divided by 100 and multiplied by the density of the cell walls (1.53 g/cm^3).

Table 1. **Table showing traits name and abbreviation.** *Traits were analyzed within each growth ring, after applying different*

detrending methods, which resulted in dimensionless measures.

Abbreviation	Trait
TRW	Tree ring width
CV LA	Coefficient of variation of the lumen area
Dh	Hydraulic diameter
Dh $_{min, max}$	Hydraulic diameter of the smallest and the biggest decile
Ks	Specific conductivity
VD	Vessel density
PC LA	Lumen area of the parenchymal cells
PC N	Number of the parenchymal cells
PC D	Parenchyma cells density
AWD	Anatomical Wood Density
RI	Red Intensity

Masting and environmental data

A regional masting chronology was developed using seed production time series retrieved from three different sources: version 2.0 of the MASTREE+ dataset (Hacket-Pain et al., 2022), the publication from Vacchiano et al., (2017), and the German National Inventory FGRDEU (https://fgrdeu.genres.de/erntehandel/ernteaufkommen). Eighteen time series of seed production were selected within a 310 km radius of the sampling site, excluding data from Denmark and Sweden due to their low resolution and discontinuities. This radius was specifically chosen to include the centroids of all the regions adjacent to the study area. To ensure comparability, continuous seed production data were converted into ordinal data (1-5), with 1 representing low and 5 high seed production (Ascoli et al., 2017). Since masting in beech is known to exhibit strong spatial synchrony up to a 300km radius (r~0.5) (Bogdziewicz et al., 2023; Gallego Zamorano et al., 2018), we first evaluated the relationship between time series similarity and geographic distance from the sampling site using a Mantel test (r^2 = -0.154, $p < 0.05$). Recognizing that spatial synchrony tends to decline with increasing distance, we incorporated a distance-based weighting scheme to adjust the influence of each time series, applying the formula: *Correction factor = 1−(Distance/1000)* to each ordinal time series (Table **S2**). In order to obtain a binary classification of masting events (hereafter referred to as masting $_{Y/N}$), where M_Y denotes a masting event and M_N represents a non-masting event, we applied a threshold criterion to each year for each time series. A given year was classified as M_Y if at least one-third of the time series exhibited a value of either 4 or 5. This method led to the identification of 13 masting events (Table 1) from which we excluded 1973, as masting events are unlikely to occur in two successive years. Overall, this was considered as the most reliable evaluation approach since many individual series were incomplete and applying different thresholds than the 1/3 rule would result in an insufficient and non-biologically plausible number of masting events.

To quantify drought, we calculated the Standardized Precipitation Evapotranspiration Index (SPEI) from the temperature and precipitation using the SPEI package (Beguería and Vicente-Serrano, 2023). Climate data were downloaded from the DWD - National Meteorological Service of Germany that holds records of local weather stations. Climate data were sourced from five local weather stations near the sampling site (Müritz, Angermünde, Wittstock, Carpin-Serrahn, and Wesenberg) and averaged to produce complete chronologies for both temperature and precipitation. According to the classification of the SPEI (McKee et al., 1993) and the site-specific conditions, we considered drought events those years for which SPEI is classified as "severely dry" (SPEI $_{MAY-JUL}$ < -1.49). Where SPEI was calculated as the average of the months May, June, and July to account for the importance of spring moisture condition for leaf unfolding (Skvareninova et al., 2024) and for the relevance of early summer drought condition in determining growth reduction (Arnič et al., 2021). As for the masting events, this

196   classification created a binary drought indicator: D_Y for drought years and D_N for non-drought years

197   (Table **2**).

198 *Table 2. List of the masting and drought events identified from 1970 to 2021.* Data were calculated according to local

199 masting records and weather stations. A star was placed for every year in common with both events.

Masting event	Drought event
1970*	1970*
1974	1971
1976*	1976*
1983*	1983*
1990	1992*
1992*	1994
1995	1997
2000	1999
2004	2003
2011	2006
2014	2008
2019	2018

Analysis

To investigate the link between masting and wood anatomy, we first analyzed the main effect of masting on beech TRW and verified that our findings matched results from literature (Drobyshev et al., 2014; Hacket-Pain et al., 2015). Notably, masting in beech involves multiple phenological stages starting from 2 years before flowering and fruiting (t-2) when resource priming occurs, to 1 year before (t-1) when floral buds are initiated, to 1 year after (t+1) where resources remain depleted (Vacchiano et al., 2017). For this reason, we defined the variable "masting phase", where TRW for each tree and each year was associated to the relative time lag with respect to the year of masting occurrence: TRW_{t-2}, TRW_{t-1}, TRW_{t0}, TRW_{t+1}, $TRW_{t\,null}$, where "t0" is the year of masting and "t null" represents years not associated to any of the defined lag. A Generalized Additive Model (GAM) was used to identify the effect of the masting phases with the following formula (Eq.**1**)

$$Eq.\,1\; gam\,(TRW\,\sim masting\,phase, \qquad random\,=\,\sim (masting\,phase|\,individual\,tree))$$

Time interaction was excluded as the TRW time series are detrended. We employed the *estimate_means()* function from the *modelbased* package in R (Hartig et al., 2024) to visualize the mean and the confidence interval of the prediction according the category and *estimate_contrasts()* function to perform the marginal contrast analysis. P-values were adjusted for multiple testing via the Benjamini-Yekutieli method.

Next, we evaluated the joint and individual effects of masting and drought on wood anatomical traits to determine whether anatomical responses could be attributed to masting alone or confounded by drought. Binary classifications were used for both phenomena ($M_{Y/N}$ and $D_{Y/N}$). Detrended wood anatomical traits were modelled fitting a generalized linear mixed model (GLMM) using Template Model Builder (TMB) according to the following variables: $M_{Y/N}$, $D_{Y/N}$, the cumulative precipitation of the growing season (from March to September) and the mean of the temperature for the same period (Eq. **2**).

$$Eq.\,2\; glmmTMB\,\big(anatomical\,trait\,\sim M_{Y/N}+D_{Y/N}+prec_{sum}+temp_{mean}\big)$$

The glmmTMB package allowed for the use of Student's t distribution which was observed to reach a better fit than normal gaussian for all models except the one involving TRW detrended with the first order difference. Random effect of the single tree was avoided as variance associated to the single trees resulted negligible. Diagnostic tests were performed on all models, including a check on the posterior predictions, collinearity, and uniformity of the residuals using the *performance* package (Lüdecke et al., 2021), and heteroskedasticity, outliers, and overdispersion using the *DHARMa* package (Hartig et al., 2024) (Table **S3**). Differences among drought and masting categories were tested with marginal contrast analysis, and P-values were adjusted for multiple testing via Benjamini-

Yekutieli method. Concerning the analysis on the effects of masting and drought, we relied on a

confusion matrix to organize the events occurrence, where we defined $M_N D_N$ as the case when neither

masting nor drought occurred (the control group), $M_Y D_N$ and $M_N D_Y$ were masting and drought events

occurring in isolation, respectively, and $M_Y D_Y$ corresponded to both events occurring in the same year

(Table **3**). The comparison between masting and drought effects against none of the events happening

($M_Y D_N$ and $M_N D_Y$ vs $M_N D_N$) constitutes the ground level of our analysis, which was further implemented

with the comparison of the effect of $M_Y D_Y$. Results were evaluated with the computation of marginal

means from the *model based* package (Makowski et al., 2025) to visualize the mean and the

confidence interval of the prediction according to the category, while the *estimate contrast* function

was used to perform the marginal contrast analysis. P-values were adjusted for multiple testing via

the Benjamini-Yekutieli method.

Table 3. Confusion matrix to interpret drought and masting event occurrence. *M_ND_N (no masting, no drought), M_YD_N*

(masting only) and M_ND_Y (drought only), M_YD_Y (both masting and drought).

	Masting $_N$	Masting $_Y$
Drought $_N$	M_ND_N	M_YD_N
Drought $_Y$	M_ND_Y	M_YD_Y

Since the analysis revealed key insights into the effects of masting on wood anatomical traits, we first

evaluated trait collinearity using a correlation matrix, highlighting values over 0.7 (Supporting

Information Fig. **S2**). Trait selection was then performed with the goal of maximizing the detectable

influence of masting on anatomical variation. Specifically, we developed a random forest

classification model to predict masting events using three trait groups: 1. QWA: only quantitative

wood anatomical traits; 2. TRW: only tree ring width (cubic spline detrended); and 3. QWA + TRW:

combined anatomical traits and TRW. Wood anatomical traits considered for the analysis were: RI

and VD detrended with the first difference method, and AWD, Dh max, PC N detrended with the cubic

spline method. Due to collinearity with TRW, PC N was excluded from the combined model (QWA +

TRW).

To assess model performance while accounting for inter-individual structure, we implemented a

nested resampling strategy. The outer loop consisted of 12 resamples using group-wise initial splits,

with tree identification code as the grouping factor, retaining 75% of the data for training and 25% for

testing in each split.

The random forest model was then specified using the *parsnip* package with engine set to *ranger*, and

hyperparameters (mtry, min_n, and trees) tuned via grid search (100 combinations). Within each outer

resample, we performed inner resampling using 4-fold grouped cross-validation for model tuning. The

best model for each split was selected based on the best result of the ROC AUC parameter.

Model evaluation on the outer test sets included the computation of ROC AUC, precision, recall, and

F1 score. Variable importance was extracted using impurity-based measures. Additionally, we

assessed model interpretability using Individual Conditional Expectation (ICE) plots via the *iml*

package (Casalicchio et al., 2025). These plots were generated for each predictor to visualize feature

effects on the predicted probability of the positive class.

The same random forest model settings and variable selection were successively applied to predict

drought events and understand the performance of the same model on a different target event. ICE

plots for drought classification were compared to those for masting classification to understand the

extent and nature of each variable's contribution to the correct prediction.

Results

Assessing tree ring width effect

Mean contrast analysis revealed that tree-ring width (TRW) was significantly narrower during mast

years (t0) compared to years preceding or following masting events ($p < 0.001$; Fig. **2**). Among the other

time lags, most comparisons were non-significant, with two exceptions: TRW in t–1 was significantly

greater than in t–2 (p < 0.05) and t+1 (p < 0.001). (Full p-value results are provided in the

supplementary, Table **S4**).

Figure 2. Estimated marginal means on the detrended TRW according to the time lag with respect to the mast year.

Confidence intervals set to 95%. Lag categories are represented by t0 year of the masting, t-1 one year prior to the masting, t-

2 two years prior to masting, and t+1 one year after masting. Horizontal black line and grey shaded area correspond to the

marginal mean and confidence interval (95%) of t null.

Drought and masting events comparison evaluation

The marginal effects of drought and masting, and their interactions were evaluated across wood

anatomical traits. Results are presented by trait groups: 1. ring structural traits (TRW, AWD, and RI), 2.

hydraulic traits (Dh, Dh max, Dh min, Ks, VD, and CV LA), and 3. storage and nutrient mobilizations

traits (PC LA, PC N, and PC D). Here we show results on the trait detrended with the cubic spline

method (Fig. **3**). A table with p-value significance for each comparison is included in the

supplementary material (Table **S5**).

Ring Structural Traits

The group exhibited significant responses across treatment conditions. Tree ring width (TRW) was

significantly reduced in mast years compared to non-mast years, whether masting occurred in the

presence (MyDy) or absence (MyDn) of drought (p < 0.01). The lowest growth was associated with the

coincidence of masting and drought (MyDy). TRW was highest under MnDy, and significantly reduced

under MnDn (p <0.01) (Fig. **3**, TRW). Anatomical wood density (AWD) increased significantly under

masting, regardless of drought (MyDn, MyDy) (p <0.01), compared to MnDn and MyDy, for which there

was no significant difference (Fig. **3**, AWD). Red intensity (RI) did not differ between MnDn and MyDy

but was significantly higher under MnDy and significantly lower under MyDn, showing the divergence between anatomy responses when the two single events occur (Fig. **3**, RI).

Hydraulic traits

Hydraulic traits showed variable responses to drought, masting, and their combination. Hydraulic diameter (Dh) showed significant reduction in the presence of masting. Dh was lower when masting occurred with respect to non-masting, in the absence of drought (MyDn < MnDn, $p < 0.01$), and the same reduction was observed for both events occurring (MyDy, $p < 0.01$) (Fig. **3**, Dh). Dh max showed the same results, only with a lower degree of significance ($p < 0.05$). Minimum hydraulic diameter (Dh min) was the only parameter that clearly showed sensitivity only to drought (Fig. **3**, Dh max). Dh min was significantly reduced in drought years ($p < 0.01$), but there was no detected effect of masting (Fig. **3**, Dh min). Specific hydraulic conductivity (Ks) showed no significant comparison. Vessel density (VD) increased significantly under MyDn and MyDy ($p < 0.01$) with respect to both MnDn and MnDy, which were not significantly different from each other (Fig. **3**, VD). Coefficient of variation of vessel lumen area (CV LA) increased significantly under the combined effects of masting and drought (MyDy), while there was no significant difference among MyDn and MnDy. These two events differed significantly from the MnDn condition ($p < 0.05$ and $p < 0.01$, respectively) (Fig. **3**, CV LA).

Storage and Nutrient Mobilization Traits

These traits were also significantly affected by masting, and to a lesser degree by drought. Parenchyma cell lumen area (PC LA) was significantly lower in MyDn and MyDy with respect to both MnDn and MnDy ($p < 0.01$) (Fig. **3**, PC LA). Parenchyma cell number (PC N) followed the same pattern as PC LA but, while for PC LA masting and drought seemed to have the same effect that was significant only for masting, in PC N masting and drought exhibit opposite behavior with respect to the control (MnDn) (Fig. **3**, PC N). However, the confidence interval revealed that masting is the only significant event affecting this trait. Parenchyma cell density (PC D) showed no significant comparisons.

Repeating the analysis for the same traits detrended with the first order difference method produced largely similar results, with differences only for certain traits. Specifically, within the ring structural traits, comparison between MnDy and MnDn in TRW were no longer significant, while AWD MnDy versus MnDn showed significant difference, with higher values for MnDy, and even greater difference for MyDn. RI relationship remained unchanged, aside from the MnDn vs MyDy comparison which differed significantly. For the hydraulic traits, the only notable difference regarded CV LA, with a heightened difference between MyDn and MnDy, where masting showed no significant difference with MnDn, as well as drought compared to MyDy. Similarly to the other groups of traits, for the storage and nutrient mobilizations traits only PC LA exhibited difference due to the detrending method and

specifically heightened the difference between MnDy towards MnDn, and MyDn towards MyDy.

Results for the traits detrended with the first order difference method are available in the

supplementary material (Supporting Information Fig. **S3**) along with the table with p-value significance

for each comparison (Table **S5**).

Results summary

Masting, drought and their interaction significantly influenced wood anatomical traits, which were

analyzed grouped into structural, hydraulic, and storage-related categories. Analyses based on cubic

spline detrended data revealed distinct patterns:

Ring Structural Traits: Tree-ring width (TRW) was highest during drought-only years and lowest when

both drought and masting co-occurred. Anatomical wood density (AWD) increased with masting,

while red intensity (RI) responded in opposite directions under drought and masting alone.

Hydraulic Traits: Masting led to narrower vessel diameters (Dh, Dh max), while Dh min was the only

tested trait that showed responses to drought but not to masting. Vessel density (VD) rose with

masting, and lumen area variation (CV LA) increased when both events occurred. Specific hydraulic

conductivity (Ks) showed no clear response.

Storage and Nutrient Traits: Masting caused a marked reduction in parenchyma cell lumen area (PC

LA) and number (PC N), suggesting a trade-off in storage capacity. Cell density (PC D) remained

unaffected.

Figure 3. Estimated marginal means of masting events vs drought events for each parameter considered. Time series

were detrended with cubic smooth spline with 10 year cutoff. Grey color refers to MnDn (control group), yellow to MnDy

(drought), green to MyDn (masting) and orange to MyDy when both events occur.

Mast year prediction

The predictive performance analysis revealed that the combined QWA+TRW model achieved

marginally higher metrics (F1-score: 0.91±0.006; precision: 0.88±0.01; recall: 0.95±0.02; ROC AUC:

0.86±0.02) compared to the QWA-only model (F1-score: 0.91±0.01; precision: 0.87±0.01; recall:

0.92±0.02; ROC AUC: 0.85±0.03). However, the results varied across runs and showed overlapping

confidence intervals between QWA+TRW and QWA-only model. In contrast, the TRW-only model

consistently demonstrated the lowest performance across all metrics (F1-score: 0.88±0.01;

precision: 0.84±0.01; recall: 0.92±0.05; ROC AUC: 0.79±0.02). The narrow error bars observed for

most metrics (Fig. 4a) indicated stable model performance across resampling iterations.

Mean variable importance for the combined model (QWA+TRW) showed that the most influential

predictor was TRW, followed by RI, and VD. Other traits such as AWD and Dh max also contributed to

the model but with comparatively lower importance. The sample ID ranked lowest in importance,

indicating minimal influence from individual-specific variation. Beside this difference the two groups

perform similarly in the variable importance evaluation (Fig. 4b).

Figure 4. Performance evaluation and variable importance of Random Forest models in predicting masting behavior.

(A) Evaluation metrics (F-1 score, precision, recall, and ROC AUC) for three model types: QWA (quantitative wood anatomical

traits only), QWA + TRW (combined wood anatomical traits and tree-ring width), and TRW (tree-ring width only) . Standard

deviation bars indicate variability across cross-validation folds.

(B) Variable importance rankings (VIP) of the top predictors in the Random Forest models. Error bars represent variability in

importance estimates across models. Traits are labeled according to the wood anatomical traits considered; all traits were

detrended using the cubic spline method, except for RI and VD, which were detrended using the first order difference

method.

Results on the same models applied to predict drought events were also assessed and showed that

the main evaluation parameter – ROC AUC – was substantially lower for the same three categories of

traits (TRW = 0.54, QWA = 0.68, and TRW + QWA = 0.68). Recall and precision showed more divergent

results, with higher recall for all three groups, but lower precision (Supporting Information Fig. **S4**).

In this regard, ICE plots were evaluated to assess the nature of the prediction of masting and drought

according to the same QWA variables. Results showed that VD and AWD show a positive relationship

with masting probability: as values increase, so does the likelihood of a masting event. RI and PCN

exhibit a strong negative relationship: lower values of these traits are associated with higher

probabilities of masting. The same is verified for Dh max but with lower intensity. Tree identification

code (ID) shows relatively minor variation, indicating that individual-level effects are less critical than

anatomical traits for predicting masting (Fig. **5**). The same traits were evaluated for drought prediction,

where we found ICE curves to be generally flatter and more diffuse, indicating weaker predictive

relationships for drought of these traits. RI together with AWD are the only traits showing a moderate

inflection at intermediate values. Dh max, AWD, PC N, and ID show only subtle changes across their

range, supporting the earlier observation that drought events do not induce strong or consistent

anatomical changes.

Figure 5. Individual Conditional Expectation (ICE) plots for masting prediction models. Each line represents an individual

observation, illustrating how predicted probability changes across the range of each variable while all other variables are

held constant. ICE plots for masting prediction showing the influence of vessel density (VD), maximum hydraulic diameter

(Dhmax), red intensity (RI), anatomical wood density (AWD), number of parenchyma cells (PC N), and tree identification code

(ID) on the model's predicted probability of a masting year

Discussion

Our results demonstrated that wood anatomical traits contain a clear and consistent imprint signal of masting events, and that the responses are distinct from those induced by drought. Results enabled us to reliably predict the occurrence of large mast years in beech avoiding confounding signals like drought. These findings suggest a high potential for masting reconstructions based on quantitative wood anatomy (QWA) representing a significant advancement compared to methods based on tree ring width (TRW) (Hacket-Pain et al., 2019).

Effects of masting and drought on wood anatomy

The distinct anatomical signatures of masting and drought observed in our study provide mechanistic insights into how trees balance resource allocation under different stressors.

We verified that our findings are in line with literature in terms of the effect of masting on TRW, which, predominantly in beech but also in other species, was reported to cause ring width reduction (Drobyshev et al., 2014; Hacket-Pain et al., 2024, 2015; Selås et al., 2002). Our analysis further indicates that this reduction is evident not only when comparing masting to non-masting years but also when considering the physiological patterns preceding or following beech masting events, such as floral induction, resource priming the previous year, and resource depletion the year after masting (Vacchiano et al., 2017) (Fig. **2**).

While it is well established that beech masting reduces growth, the effects of masting on wood anatomical traits alone and in combination to drought remain largely unexplored, with only one study specifically tackling this research gap, investigating the effects of masting on beech hydraulic structure (Rodríguez-Ramírez et al., 2019). However, relying on macroscopical images of beech cores holds the disadvantage of poorly capturing vessel variability. Disentangling drought from masting effects was essential to isolate the true effect of reproductive investment on quantitative wood anatomy, to reveal distinct physiological trade-offs that would otherwise be obscured by overlapping stress responses. A general observation on the effect of drought and masting on wood anatomy, is that drought did not seem to have a strong impact on our samples. This is primarily supported by the fact that TRW was not reduced by the occurrence of drought, as it is otherwise widely documented in beech (Leuschner et al., 2001; Martinez del Castillo et al., 2022; Schmied et al., 2023). Generally, the main effect of drought was observed in hydraulic traits, with the hydraulic diameter of the smallest cells (Dh min) being clearly negatively affected solely by drought. These findings align with well-established physiological responses to water limitation, where trees prioritize hydraulic safety by narrowing conduits to reduce embolism risk (Anderegg et al., 2016; Arend et al., 2022). However, the effect of drought on the other hydraulic traits is partially masked by the effect of masting, generating in

the mean and max hydraulic diameter (Dh and Dh max), specific hydraulic conductivity (Ks) and

coefficient of variation of the lumen area (CV LA) wide error bars and overlapping responses that lead

to non-significant comparison between drought and masting effects (Fig **3**). This lack of specific

response to drought by a certain array of traits might be explained by the fact that the sampling site is

located in the Mecklenburg Lake District, which constitutes a moist environment where microclimate

is significantly influenced by lakes (Scharnweber et al., 2011). Moreover, the conspicuous tree height

(34 - 42.9 m) also suggests a fertile soil and a generally favorable growing condition.

By studying masting in combination with drought, we were able to disentangle the specific impact of

masting on quantitative wood anatomy, providing novel insights into their interaction. This

comparison revealed that masting influenced anatomical, while drought alone did not show

statistically significant changes compared to the control group (MnDn). The broader anatomical

signature of masting is represented by reductions in tree-ring width (TRW), red intensity (RI), mean and

maximum hydraulic diameter (Dh and Dh max), lumen area and number of parenchyma cells (PC LA

and PC N), coupled with increases in vessel density (VD) and anatomical wood density (AWD). These

traits and the nature of their responses suggested a complex mechanism of general downregulation of

growth and investment in xylem tissue during masting, while reflecting a compensatory strategy to

maintain hydraulic function and mechanical stability under resource-constrained conditions.

Evidence suggests, in fact, that ecological mechanisms such as the ones driving masting effects on

growth, should not be reduced to simple carbon-mediated 'trade-off' (Mund et al., 2020), as they also

involve structural adjustments tied to whole-tree hydraulic architecture. Previous work on *Fagus*

sylvatica indicates that tree size and crown architecture better explain vessel diameter variation,

rather than general climatic factors (Kröner et al., 2024; Prislan et al., 2018). In this context, our

findings align with observations that mast years are associated with reduced shoot length and fewer

leaves per shoot (Han et al., 2011; Müller-Haubold et al., 2015), pointing towards the link between

reduced leaf area and lower conductive demand (decrease in Dh).

Notably, this study is among the first to explore such a broad suite of wood anatomical traits in

relation to masting, including a specific focus on axial parenchyma characteristics (Plavcová et al.,

2024; Słupianek et al., 2021). Parenchyma cells are primarily involved in nutrient storage and

mobilization functions (Tomasella et al., 2020), and while a reduction in their number (PC N) was

expected due to narrower tree rings and known trade-offs with hydraulic traits (Zheng and Martínez-

Cabrera, 2013), we also found a significant decrease in parenchyma lumen area (PC LA). This finding

is particularly interesting considering the existing literature reporting high genetic control and low

plasticity of wood anatomical features (Chhetri et al., 2020; Hajek et al., 2016). A plausible

explanation is that during mast years, when non-structural carbohydrates (NSC) used for

reproduction are predominantly sourced from current-season assimilates (Hoch, 2005; Ichie et al.,

2013), there is a reduced reliance on stored reserves. As a result, there may be limited carbon

investment into storage tissue development, which further contributes to explain the concurrent

reduction in both the number and lumen area of parenchyma cells.

Red intensity (RI), a proxy for lignin content in tree rings, showed opposite effects for masting than for

drought (Fig. **3**) (Resente et al., in review). This distinction is ecologically meaningful as lignin supports

both structural defense and hydraulic safety (Polo et al., 2020). The RI reduction during masting aligns

with reported carbon trade-offs in reproductive years, where lignin biosynthesis may be deprioritized

to allocate resources toward flowering and seed production (Han and Kabeya, 2017; Monks and Kelly,

2006; Redmond et al., 2019). Conversely, drought-induced RI increase mirrors global patterns of

enhanced lignification under water stress to reinforce xylem against embolism (Crivellaro et al., 2022;

Lima et al., 2018; Resente and Crivellaro, 2025). These opposing RI trends empirically validate

theoretical models predicting resource allocation shifts between reproduction and stress resilience

(Bogdziewicz, 2022).

The potential of wood anatomy for masting reconstruction

Previous attempts to reconstruct ecological events such as masting have relied solely on tree-ring

width (TRW) as anatomical proxy. For these reason existing reconstruction studies, either rely on the

combination of TRW and climatic data, or, in the case of Araucaria, employ the dioecious

characteristic of the species to enable the separation of masting from the climate signal (Drobyshev

et al., 2014; Hacket-Pain et al., 2024; Mundo et al., 2021). Given that masting influences forest

regeneration, carbon cycling, and wildlife dynamics (Bogdziewicz et al., 2021), accurate

reconstructions are crucial for understanding long-term ecosystem responses to environmental

change. Our findings demonstrate that although TRW carries substantial predictive power, it is

outperformed by models incorporating quantitative wood anatomical traits (QWA) or models

including a combination of QWA and TRW. To our knowledge, this is one of the first studies to

establish a direct link between masting events reconstruction and a wide suite of QWA traits, beside

Rodríguez-Ramírez et al. (2019) which focused on the hydraulic structure. Particularly, the QWA-only

model performed as well as the combined model, confirming that QWA traits contain substantial

independent information relevant to masting prediction. Among the QWA traits, the number of

parenchyma cells (PC N) and red intensity (RI) emerged as particularly strong predictors (Fig. 4). For

PC N, this can be attributed to the expected collinearity between PC N and TRW, since narrower tree

rings are generally associated with a lower number of cells (Babushkina et al., 2021). For RI, the

opposite effects of drought and masting are relevant; drought increased while masting decreased RI.

This provides a strong basis for robustly identifying mast years, which have previously been

challenging due to common response of TRW to these two drivers (Drobyshev et al., 2014). The strong

predictive responses of both PC N and RI validated our analytical approach to isolating the effects of masting; while traits such as VD, AWD, and Dhmax showed comparatively weaker predictive power, suggesting a less distinctive response to masting events.

When the same random forest modeling approach was applied to drought prediction, model performance dropped significantly across all metrics, particularly in relation to ROC AUC, which indicates the general reliability of the model in classifying the event as drought or non-drought. This suggests that traits selected to maximize the effect of masting on wood anatomy do not capture drought-specific signatures with the same reliability. The moderate performance predicting drought may be largely driven by the discriminative role of RI, which in the previous analysis (Fig. **3**) and in the ICE plots (Fig. **5**) showed a distinct opposite response to drought than to masting. Nonetheless, the generally lower precision and ROC AUC values indicate that anatomical traits selected better reflect the coordinated, resource-driven shifts associated with reproductive events. The ICE plots further support this interpretation. Traits such as vessel density (VD), red intensity (RI), and parenchyma cell number PCN show clear, directional relationships with masting probability, reflecting their strong influence on prediction. In contrast, ICE plots for drought prediction display flatter, more diffuse patterns, indicating that variability in these traits does not translate into consistent predictive signals for drought years. This suggests that while the same anatomical dataset may contain some signal of drought impact as one third of the mast years were also drought years, it lacks the distinct, coordinated anatomical signature seen in response to masting. Further support for the general applicability of our approach comes from the finding that individual tree identity, despite the low sample size, had negligible influence on model performance, suggesting that the predictive framework could be reliably extended across different trees and potentially across sites, which constitutes an important step toward developing scalable, species-wide reconstructions of masting dynamics.

Together, these findings show that QWA traits not only improve predictive accuracy but also provide insight into the physiological adjustments that trees undergo under biotic and abiotic pressures. Moreover, in line with previous dendrochronological reconstructions (Drobyshev et al., 2014; Mundo et al., 2021), our binary mast/non-mast approach lays the groundwork for future studies aiming to reconstruct continuous seed production—a crucial step toward detecting long-term trends and potential masting patterns.

Conclusions

In this study we present the first clear link between masting, the synchronized and variable seed
production, and structural changes in the wood anatomy of European beech.

Our results show that mast years are associated with a general downregulation of growth and storage
investment, evident in narrower rings, smaller hydraulic diameters, reduced lignin content, and fewer
and smaller parenchyma cells, coupled with increased vessel density and anatomical wood density.
These patterns suggest a complex mechanism of carbon reallocation towards reproduction, while
reflecting a compensatory strategy to maintain hydraulic function and mechanical stability under
resource-constrained conditions. Notably, red intensity – a proxy reflecting lignin content in tree rings
– was the only trait to show opposite responses to masting and drought, emphasizing the distinct
ecological strategies trees adopt under reproductive versus abiotic stress.

By capturing these specific wood anatomical adjustments, QWA traits emerged as strong predictors
of masting. The classification model based on QWA performed better than a model including TRW
alone, highlighting the advantage of the intra annual approach, and equally to a combined model of
QWA and TRW. Moreover, the most explanatory traits for predicting masting events mirrored in
intensity and nature the results found on the effect of masting in wood anatomy. The minimal effect of
individual tree identity further supports the possibility of generalization of our approach, promoting
species-wide reconstructions based on anatomical data. Importantly, the same anatomical traits that
reliably predicted masting, failed to accurately capture drought years in our dataset. This discrepancy
further reinforces that the anatomical signature observed is tightly linked to reproductive effort rather
than overlapping environmental stress. In addition, the study site presents particularly favorable
conditions for tree growth, therefore, we recommend replicating the study in a more drought-sensitive
location.

These findings underscore that quantitative wood anatomy provides a robust, species-wide tool for
reconstructing and predicting mast years in European beech, bridging the current gap on reliable
reconstruction methodologies, and therefore enabling more accurate predictions of future tree
reproduction trajectories.

Acknowledgements

563 DA and AP were supported by the Agritech National Research Center and the European Union Next-
564 Generation EU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2,
565 INVESTIMENTO 1.4 – D.D. 1032 June 17, 2022, CN00000022), Spoke 4, Task 4.1.3

Competing interests

The authors have no competing interests to declare.

Author contributions

All authors contributed equally to the study. Conceptualization: GR, MW, AC, AP, and DA. Design: GR and MW. Data Collection: GR, EF and FM. Analysis: GR, EF, FM, AHP, DA and AP. Original draft: GR, DA, AHP. Review & Editing: GR, AC, EF, AP, FM, MW, AHP, RM, and DA. Supervision: DA, RM, AHP, AP, and AC.

Data availability

The data that support the findings of this study are available from the corresponding author, GR, upon request.

References

Anderegg, W.R.L., Klein, T., Bartlett, M., Sack, L., Pellegrini, A.F.A., Choat, B., Jansen, S., 2016. Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. Proc. Natl. Acad. Sci. 113, 5024–5029. https://doi.org/10.1073/pnas.1525678113

Arend, M., Link, R.M., Zahnd, C., Hoch, G., Schuldt, B., Kahmen, A., 2022. Lack of hydraulic recovery as a cause of post-drought foliage reduction and canopy decline in European beech. New Phytol. 234, 1195–1205. https://doi.org/10.1111/nph.18065

Arnič, D., Gričar, J., Jevšenak, J., Božič, G., von Arx, G., Prislan, P., 2021. Different Wood Anatomical and Growth Responses in European Beech (Fagus sylvatica L.) at Three Forest Sites in Slovenia. Front. Plant Sci. 12. https://doi.org/10.3389/fpls.2021.669229

Ascoli, D., Hacket-Pain, A., Pearse, I.S., Vacchiano, G., Corti, S., Davini, P., 2021. Modes of climate variability bridge proximate and evolutionary mechanisms of masting. Philos. Trans. R. Soc. B Biol. Sci. 376, 20200380. https://doi.org/10.1098/rstb.2020.0380

Ascoli, D., Vacchiano, G., Turco, M., Conedera, M., Drobyshev, I., Maringer, J., Motta, R., Hacket-Pain, A., 2017. Inter-annual and decadal changes in teleconnections drive continental-scale synchronization of tree reproduction. Nat. Commun. 8, 2205. https://doi.org/10.1038/s41467-017-02348-9

Babushkina, E.A., Dergunov, D.R., Belokopytova, L.V., Zhirnova, D.F., Upadhyay, K.K., Tripathi, S.K., Zharkov, M.S., Vaganov, E.A., 2021. Non-linear Response to Cell Number Revealed and Eliminated From Long-Term Tracheid Measurements of Scots Pine in Southern Siberia. Front. Plant Sci. 12, 719796. https://doi.org/10.3389/fpls.2021.719796

Bajocco, S., Ferrara, C., Bascietto, M., Alivernini, A., Chirichella, R., Cutini, A., Chianucci, F., 2021. Characterizing the climatic niche of mast seeding in beech: Evidences of trade-offs between vegetation growth and seed production. Ecol. Indic. 121, 107139. https://doi.org/10.1016/j.ecolind.2020.107139

Beguería, S., Vicente-Serrano, S.M., 2023. SPEI: Calculation of the Standardized Precipitation-Evapotranspiration Index.

Bogdziewicz, M., 2022. How will global change affect plant reproduction? A framework for mast seeding trends. New Phytol. 234, 14–20. https://doi.org/10.1111/nph.17682

Bogdziewicz, M., Hacket-Pain, A., Ascoli, D., Szymkowiak, J., 2021. Environmental variation drives continental-scale synchrony of European beech reproduction. Ecology 102, e03384. https://doi.org/10.1002/ecy.3384

Bogdziewicz, M., Journé, V., Hacket-Pain, A., Szymkowiak, J., 2023. Mechanisms driving interspecific variation in regional synchrony of trees reproduction. Ecol. Lett. 26, 754–764. https://doi.org/10.1111/ele.14187

Brumme, R., Ahrends, B., Block, J., Schulz, C., Meesenburg, H., Klinck, U., Wagner, M., Khanna, P.K., 2021. Cycling and retention of nitrogen in European beech (*Fagus sylvatica* L.) ecosystems under elevated fructification frequency. Biogeosciences 18, 3763–3779. https://doi.org/10.5194/bg-18-3763-2021

Bunn, A.G., 2010. Statistical and visual crossdating in R using the dplR library. Dendrochronologia 28, 251–258. https://doi.org/10.1016/j.dendro.2009.12.001

Casalicchio, G., Molnar, C., Schratz, P., 2025. iml: Interpretable Machine Learning.

Chhetri, H.B., Furches, A., Macaya-Sanz, D., Walker, A.R., Kainer, D., Jones, P., Harman-Ware, A.E., Tschaplinski, T.J., Jacobson, D., Tuskan, G.A., DiFazio, S.P., 2020. Genome-Wide Association

621   Study of Wood Anatomical and Morphological Traits in Populus trichocarpa. Front. Plant Sci.
622   11. https://doi.org/10.3389/fpls.2020.545748

623 Cook, E.R., Kairiukstis, L.A. (Eds.), 1990. Methods of Dendrochronology. Springer Netherlands,
624   Dordrecht. https://doi.org/10.1007/978-94-015-7879-0

625 Crivellaro, A., Piermattei, A., Dolezal, J., Dupree, P., Büntgen, U., 2022. Biogeographic implication of
626   temperature-induced plant cell wall lignification. Commun. Biol. 5, 767.
627   https://doi.org/10.1038/s42003-022-03732-y

628 Ding, H., Pretzsch, H., Schütze, G., Rötzer, T., 2017. Size-dependence of tree growth response to
629   drought for Norway spruce and European beech individuals in monospecific and mixed-
630   species stands. Plant Biol. 19, 709–719. https://doi.org/10.1111/plb.12596

631 Drobyshev, I., Niklasson, M., Mazerolle, M.J., Bergeron, Y., 2014. Reconstruction of a 253-year long
632   mast record of European beech reveals its association with large scale temperature variability
633   and no long-term trend in mast frequencies. Agric. For. Meteorol. 192–193, 9–17.
634   https://doi.org/10.1016/j.agrformet.2014.02.010

635 Foest, J.J., Bogdziewicz, M., Pesendorfer, M.B., Ascoli, D., Cutini, A., Nussbaumer, A., Verstraeten, A.,
636   Beudert, B., Chianucci, F., Mezzavilla, F., Gratzer, G., Kunstler, G., Meesenburg, H., Wagner,
637   M., Mund, M., Cools, N., Vacek, S., Schmidt, W., Vacek, Z., Hacket-Pain, A., 2024. Widespread
638   breakdown in masting in European beech due to rising summer temperatures. Glob. Change
639   Biol. 30, e17307. https://doi.org/10.1111/gcb.17307

640 Gallego Zamorano, J., Hokkanen, T., Lehikoinen, A., 2018. Climate-driven synchrony in seed
641   production of masting deciduous and conifer tree species. J. Plant Ecol. 11, 180–188.
642   https://doi.org/10.1093/jpe/rtw117

643 Hacket-Pain, A., Ascoli, D., Berretti, R., Mencuccini, M., Motta, R., Nola, P., Piussi, P., Ruffinatto, F.,
644   Vacchiano, G., 2019. Temperature and masting control Norway spruce growth, but with high
645   individual tree variability. For. Ecol. Manag. 438, 142–150.
646   https://doi.org/10.1016/j.foreco.2019.02.014

647 Hacket-Pain, A., Foest, J.J., Pearse, I.S., LaMontagne, J.M., Koenig, W.D., Vacchiano, G., Bogdziewicz,
648   M., Caignard, T., Celebias, P., van Dormolen, J., Fernández-Martínez, M., Moris, J.V.,
649   Palaghianu, C., Pesendorfer, M., Satake, A., Schermer, E., Tanentzap, A.J., Thomas, P.A.,
650   Vecchio, D., Wion, A.P., Wohlgemuth, T., Xue, T., Abernethy, K., Aravena Acuña, M.-C., Daniel
651   Barrera, M., Barton, J.H., Boutin, S., Bush, E.R., Donoso Calderón, S., Carevic, F.S., de
652   Castilho, C.V., Manuel Cellini, J., Chapman, C.A., Chapman, H., Chianucci, F., da Costa, P.,
653   Croisé, L., Cutini, A., Dantzer, B., Justin DeRose, R., Dikangadissi, J.-T., Dimoto, E., da
654   Fonseca, F.L., Gallo, L., Gratzer, G., Greene, D.F., Hadad, M.A., Herrera, A.H., Jeffery, K.J.,
655   Johnstone, J.F., Kalbitzer, U., Kantorowicz, W., Klimas, C.A., Lageard, J.G.A., Lane, J., Lapin, K.,
656   Ledwoń, M., Leeper, A.C., Vanessa Lencinas, M., Lira-Guedes, A.C., Lordon, M.C., Marchelli,
657   P., Marino, S., Schmidt Van Marle, H., McAdam, A.G., Momont, L.R.W., Nicolas, M., de Oliveira
658   Wadt, L.H., Panahi, P., Martínez Pastur, G., Patterson, T., Luis Peri, P., Piechnik, Ł.,
659   Pourhashemi, M., Espinoza Quezada, C., Roig, F.A., Peña Rojas, K., Micaela Rosas, Y.,
660   Schueler, S., Seget, B., Soler, R., Steele, M.A., Toro-Manríquez, M., Tutin, C.E.G.,
661   Ukizintambara, T., White, L., Yadok, B., Willis, J.L., Zolles, A., Żywiec, M., Ascoli, D., 2022.
662   MASTREE+: Time-series of plant reproductive effort from six continents. Glob. Change Biol. 28,
663   3066–3082. https://doi.org/10.1111/gcb.16130

664 Hacket-Pain, A., Roig, F.A., Ascoli, D., LeQuesne, C., Hadad, M., 2024. Reconstruction of Araucaria
665   araucana cone production reveals warming intensifies regionally synchronized masting.
666   Ecosphere 15, e70087. https://doi.org/10.1002/ecs2.70087

667 Hacket-Pain, A.J., Friend, A.D., Lageard, J.G.A., Thomas, P.A., 2015. The influence of masting
668   phenomenon on growth–climate relationships in trees: explaining the influence of previous

summers' climate on ring width. Tree Physiol. 35, 319–330.
https://doi.org/10.1093/treephys/tpv007

Hadad, M.A., Roig, F.A., Arco Molina, J.G., Hacket-Pain, A., 2021. Growth of male and female
Araucaria araucana trees respond differently to regional mast events, creating sex-specific
patterns in their tree-ring chronologies. Ecol. Indic. 122, 107245.
https://doi.org/10.1016/j.ecolind.2020.107245

Hajek, P., Kurjak, D., von Wühlisch, G., Delzon, S., Schuldt, B., 2016. Intraspecific Variation in Wood
Anatomical, Hydraulic, and Foliar Traits in Ten European Beech Provenances Differing in
Growth Yield. Front. Plant Sci. 7, 791. https://doi.org/10.3389/fpls.2016.00791

Han, Q., Kabeya, D., 2017. Recent developments in understanding mast seeding in relation to
dynamics of carbon and nitrogen resources in temperate trees. Ecol. Res. 32, 771–778.
https://doi.org/10.1007/s11284-017-1494-8

Han, Q., Kabeya, D., Hoch, G., 2011. Leaf traits, shoot growth and seed production in mature Fagus
sylvatica trees after 8 years of CO_2 enrichment. Ann. Bot. 107, 1405–1411.
https://doi.org/10.1093/aob/mcr082

Hartig, F., Lohse, L., leite, M. de S., 2024. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level
/ Mixed) Regression Models.

Hoch, G., 2005. Fruit-bearing branchlets are carbon autonomous in mature broad-leaved temperate
forest trees. Plant Cell Environ. 28, 651–659. https://doi.org/10.1111/j.1365-
3040.2004.01311.x

Ichie, T., Igarashi, S., Yoshida, S., Kenzo, T., Masaki, T., Tayasu, I., 2013. Are stored carbohydrates
necessary for seed production in temperate deciduous trees? J. Ecol. 101, 525–531.
https://doi.org/10.1111/1365-2745.12038

Kelly, D., Sork, V.L., 2002. Mast Seeding in Perennial Plants: Why, How, Where? Annu. Rev. Ecol. Evol.
Syst. 33, 427–447. https://doi.org/10.1146/annurev.ecolsys.33.020602.095433

Koenig, W.D., Knops, J.M.H., Carmen, W.J., 2020. Can mast history be inferred from radial growth? A
test using five species of California oaks. For. Ecol. Manag. 472, 118233.
https://doi.org/10.1016/j.foreco.2020.118233

Kollmann, F.F.P., Côté, W.A., 1968. Principles of Wood Science and Technology: I Solid Wood.
Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-87928-9

Kröner, K., Larysch, E., Schindler, Z., Obladen, N., Frey, J., Stangler, D.F., Seifert, T., 2024. Influence of
crown morphology and branch architecture on tree radial growth of drought-affected *Fagus
sylvatica* L. For. Ecosyst. 11, 100237. https://doi.org/10.1016/j.fecs.2024.100237

Leuschner, C., Backes, K., Hertel, D., Schipka, F., Schmitt, U., Terborg, O., Runge, M., 2001. Drought
responses at leaf, stem and fine root levels of competitive *Fagus sylvatica* L. and *Quercus
petraea* (Matt.) Liebl. trees in dry and wet years. For. Ecol. Manag. 149, 33–46.
https://doi.org/10.1016/S0378-1127(00)00543-0

Lima, T.R.A., Carvalho, E.C.D., Martins, F.R., Oliveira, R.S., Miranda, R.S., Müller, C.S., Pereira, L.,
Bittencourt, P.R.L., Sobczak, J.C.M.S.M., Gomes-Filho, E., Costa, R.C., Araújo, F.S., 2018.
Lignin composition is related to xylem embolism resistance and leaf life span in trees in a
tropical semiarid climate. New Phytol. 219, 1252–1262. https://doi.org/10.1111/nph.15211

Lüdecke, D., Ben-Shachar, M.S., Patil, I., Waggoner, P., Makowski, D., 2021. performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. J. Open Source Softw.
6, 3139. https://doi.org/10.21105/joss.03139

Makowski, D., Lüdecke, D., Ben-Shachar, M.S., Patil, I., Thériault, R., 2025. modelbased: Estimation of
Model-Based Predictions, Contrasts and Means.

Martinez del Castillo, E., Zang, C.S., Buras, A., Hacket-Pain, A., Esper, J., Serrano-Notivoli, R., Hartl,
C., Weigel, R., Klesse, S., Resco de Dios, V., Scharnweber, T., Dorado-Liñán, I., van der
Maaten-Theunissen, M., van der Maaten, E., Jump, A., Mikac, S., Banzragch, B.-E., Beck, W.,
Cavin, L., Claessens, H., Čada, V., Čufar, K., Dulamsuren, C., Gričar, J., Gil-Pelegrín, E., Janda,
P., Kazimirovic, M., Kreyling, J., Latte, N., Leuschner, C., Longares, L.A., Menzel, A., Merela, M.,
Motta, R., Muffler, L., Nola, P., Petritan, A.M., Petritan, I.C., Prislan, P., Rubio-Cuadrado, Á.,
Rydval, M., Stajić, B., Svoboda, M., Toromani, E., Trotsiuk, V., Wilmking, M., Zlatanov, T., de
Luis, M., 2022. Climate-change-driven growth decline of European beech forests. Commun.
Biol. 5, 1–9. https://doi.org/10.1038/s42003-022-03107-3

McKee, T.B., Doesken, N.J., Kleist, J., 1993. The relationship of drought frequency and duration to time
scales InProceedings of the Eighth Conference on Applied Climatology, 179–184.

Michaud, T.J., Pearse, I.S., Kauserud, H., Andrew, C.J., Kennedy, P.G., 2024. Mast seeding in European
beech (Fagus sylvatica L.) is associated with reduced fungal sporocarp production and
community diversity. Ecol. Lett. 27, e14460. https://doi.org/10.1111/ele.14460

Monks, A., Kelly, D., 2006. Testing the resource-matching hypothesis in the mast seeding tree
Nothofagus truncata (Fagaceae). Austral Ecol. 31, 366–375. https://doi.org/10.1111/j.1442-
9993.2006.01565.x

Müller-Haubold, H., Hertel, D., Leuschner, C., 2015. Climatic Drivers of Mast Fruiting in European
Beech and Resulting C and N Allocation Shifts. Ecosystems 18, 1083–1100.
https://doi.org/10.1007/s10021-015-9885-6

Mund, M., Herbst, M., Knohl, A., Matthäus, B., Schumacher, J., Schall, P., Siebicke, L., Tamrakar, R.,
Ammer, C., 2020. It is not just a 'trade-off': indications for sink- and source-limitation to
vegetative and regenerative growth in an old-growth beech forest. New Phytol. 226, 111–125.
https://doi.org/10.1111/nph.16408

Mundo, I.A., Sanguinetti, J., Kitzberger, T., 2021. Multi-centennial phase-locking between reproduction
of a South American conifer and large-scale drivers of climate. Nat. Plants 7, 1560–1570.
https://doi.org/10.1038/s41477-021-01038-1

Nussbaumer, A., Gessler, A., Benham, S., de Cinti, B., Etzold, S., Ingerslev, M., Jacob, F., Lebourgeois,
F., Levanic, T., Marjanović, H., Nicolas, M., Ostrogović Sever, M.Z., Priwitzer, T., Rautio, P.,
Roskams, P., Sanders, T.G.M., Schmitt, M., Šrámek, V., Thimonier, A., Ukonmaanaho, L.,
Verstraeten, A., Vesterdal, L., Wagner, M., Waldner, P., Rigling, A., 2021. Contrasting Resource
Dynamics in Mast Years for European Beech and Oak—A Continental Scale Analysis. Front.
For. Glob. Change 4. https://doi.org/10.3389/ffgc.2021.689836

Ostfeld, R.S., Keesing, F., Ostfeld, R.S., Keesing, F., 2000. Pulsed resources and community dynamics
of consumers in terrestrial ecosystems. Trends Ecol. Evol. 15, 232–237.
https://doi.org/10.1016/S0169-5347(00)01862-0

Pearse, I.S., Koenig, W.D., Kelly, D., 2016. Mechanisms of mast seeding: resources, weather, cues,
and selection. New Phytol. 212, 546–562. https://doi.org/10.1111/nph.14114

Pearse, I.S., Wion, A.P., Gonzalez, A.D., Pesendorfer, M.B., 2021. Understanding mast seeding for
conservation and land management. Philos. Trans. R. Soc. B Biol. Sci. 376, 20200383.
https://doi.org/10.1098/rstb.2020.0383

Piovesan, G., Biondi, F., Filippo, A.D., Alessandrini, A., Maugeri, M., 2008. Drought-driven growth
reduction in old beech (Fagus sylvatica L.) forests of the central Apennines, Italy. Glob.
Change Biol. 14, 1265–1281. https://doi.org/10.1111/j.1365-2486.2008.01570.x

Plavcová, L., Jandová, V., Altman, J., Liancourt, P., Korznikov, K., Doležal, J., 2024. Variations in wood
anatomy in Afrotropical trees with a particular emphasis on radial and axial parenchyma. Ann.
Bot. 134, 151–162. https://doi.org/10.1093/aob/mcae049

Polo, C.C., Pereira, L., Mazzafera, P., Flores-Borges, D.N.A., Mayer, J.L.S., Guizar-Sicairos, M., Holler, M., Barsi-Andreeta, M., Westfahl, H., Meneau, F., 2020. Correlations between lignin content and structural robustness in plants revealed by X-ray ptychography. Sci. Rep. 10, 6023. https://doi.org/10.1038/s41598-020-63093-6

Prebble, J.G., Dunbar, G., van den Bos, V., Li, X., Vandergoes, M., Richardson, S., Horgan, H., Holt, K.A., Howarth, J., Levy, R., 2022. Toward a novel multi-century archive of tree mast using pollen from lake sediments. The Holocene 32, 1184–1192. https://doi.org/10.1177/09596836221114292

Prislan, P., Čufar, K., De Luis, M., Gričar, J., 2018. Precipitation is not limiting for xylem formation dynamics and vessel development in European beech from two temperate forest sites. Tree Physiol. 38, 186–197. https://doi.org/10.1093/treephys/tpx167

Redmond, M.D., Davis, T.S., Ferrenberg, S., Wion, A.P., 2019. Resource allocation trade-offs in a mast-seeding conifer: piñon pine prioritizes reproduction over defence. AoB PLANTS 11, plz070. https://doi.org/10.1093/aobpla/plz070

Resente, G., Crivellaro, A., 2025. Environmental Impacts on Plant Cell Wall Lignification. J. Bioresour. Bioprod. 10, 4–6. https://doi.org/10.1016/j.jobab.2024.11.001

Resente, G., Di Fabio, A., Scharnweber, T., Gillert, A., Crivellaro, A., Anadon-Rosell, A., Trouillier, M., Kreyling, J., Wilmking, M., 2024. The importance of variance and microsite conditions for growth and hydraulic responses following long-term rewetting in pedunculate oak wood. Trees 38, 1161–1175. https://doi.org/10.1007/s00468-024-02543-4

Resente, G., García-Hidalgo, M., Piermattei, A., Maimone, F., Ascoli, D., Wilmking, M., Crivellaro, A., in review. Exploring Red Intensity Trends in Double-Stained Wood Thin Sections. https://doi.org/10.2139/ssrn.5169865

Resente, G., Gillert, A., Trouillier, M., Anadon-Rosell, A., Peters, R.L., von Arx, G., von Lukas, U., Wilmking, M., 2021. Mask, Train, Repeat! Artificial Intelligence for Quantitative Wood Anatomy. Front. Plant Sci. 12. https://doi.org/10.3389/fpls.2021.767400

Rodríguez-Ramírez, E.C., Terrazas, T., Luna-Vega, I., 2019. The influence of climate on the masting behavior of Mexican beech: growth rings and xylem anatomy. Trees 33, 23–35. https://doi.org/10.1007/s00468-018-1755-3

Scharnweber, T., Manthey, M., Criegee, C., Bauwe, A., Schröder, C., Wilmking, M., 2011. Drought matters – Declining precipitation influences growth of *Fagus sylvatica* L. and *Quercus robur* L. in north-eastern Germany. For. Ecol. Manag. 262, 947–961. https://doi.org/10.1016/j.foreco.2011.05.026

Schmied, G., Pretzsch, H., Ambs, D., Uhl, E., Schmucker, J., Fäth, J., Biber, P., Hoffmann, Y.-D., Šeho, M., Mellert, K.H., Hilmers, T., 2023. Rapid beech decline under recurrent drought stress: Individual neighborhood structure and soil properties matter. For. Ecol. Manag. 545, 121305. https://doi.org/10.1016/j.foreco.2023.121305

Selås, V., Piovesan, G., Adams, J.M., Bernabei, M., 2002. Climatic factors controlling reproduction and growth of Norway spruce in southern Norway. Can. J. For. Res. 32, 217–225. https://doi.org/10.1139/x01-192

Skvareninova, J., Sitko, R., Vido, J., Snopková, Z., Skvarenina, J., 2024. Phenological response of European beech (Fagus sylvatica L.) to climate change in the Western Carpathian climatic-geographical zones. Front. Plant Sci. 15. https://doi.org/10.3389/fpls.2024.1242695

Słupianek, A., Dolzblasz, A., Sokołowska, K., 2021. Xylem Parenchyma—Role and Relevance in Wood Functioning in Trees. Plants 10, 1247. https://doi.org/10.3390/plants10061247

Stolz, J., Forkel, M., van der Maaten, E., Martin, J., van der Maaten-Theunissen, M., 2023. Through
eagle eyes—the potential of satellite-derived LAI time series to estimate masting events and
tree-ring width of European beech. Reg. Environ. Change 23, 74.
https://doi.org/10.1007/s10113-023-02068-5

Tomasella, M., Petrussa, E., Petruzzellis, F., Nardini, A., Casolo, V., 2020. The Possible Role of Non-
Structural Carbohydrates in the Regulation of Tree Hydraulics. Int. J. Mol. Sci. 21, 144.
https://doi.org/10.3390/ijms21010144

Vacchiano, G., Hacket-Pain, A., Turco, M., Motta, R., Maringer, J., Conedera, M., Drobyshev, I., Ascoli,
D., 2017. Spatial patterns and broad-scale weather cues of beech mast seeding in Europe.
New Phytol. 215, 595–608. https://doi.org/10.1111/nph.14600

Wion, A.P., Pearse, I.S., Broxson, M., Redmond, M.D., 2025. Mast hindcasts reveal pervasive effects of
extreme drought on a foundational conifer species. New Phytol. 246, 450–460.
https://doi.org/10.1111/nph.20321

Zheng, J., Martínez-Cabrera, H.I., 2013. Wood anatomical correlates with theoretical conductivity and
wood density across China: Evolutionary evidence of the functional differentiation of axial and
radial parenchyma. Ann. Bot. 112, 927–935. https://doi.org/10.1093/aob/mct153

Supplementary information legend

Figure S1. R-bar evaluation performed on data from the complete set of traits detrended with four

detrending method (cubic spline with a 10 years' time window, 20 years, 30 years and first order

difference)

Figure S2. Correlation matrix between all the traits detrended with both cubic spline 10 years' time

window and first order difference methods.

Figure S3. Estimated marginal means of masting events vs drought events computed for every

parameter considered in the analysis detrended with first order difference.

Figure S4. Performance evaluation and variable importance of Random Forest models in

predicting drought behavior.

Table S1. Field collected data, together with derived data from dendrochronological analyses.

Table S2. Masting time series (1970-2021) Id number of each column is composed of "site

id"_"NUTS1"_"source"

Table S3. Models' diagnostic tests

Table S4. Estimated marginal means metric with contrast applied to the TRW trait. P value

adjusted via the Benjamini-Yekutieli method.

Table S5. P-value results for significance for each comparison (Mn vs My vs Dn vs Dn) for both

detrending methods