Sedative choice alters *Klebsiella pneumoniae* lung pathogenesis and dissemination

David Mains[1†*], Ella R. Rotman[2§*], Deanna K. Aman[3*], Denise A. Ludvik[4], Anirudh Desikan[4||], Acadia A. Kocher[2#], An N.H. Tran[3], Mark J. Mandel[2,4], Nancy E. Freitag[1,3]

[1]Department of Microbiology and Immunology, University of Illinois at Chicago, Chicago, IL 60612

[2]Department of Microbiology-Immunology, Northwestern University Feinberg School of Medicine, Chicago, IL 60611

[3]Department of Pharmaceutical Sciences, University of Illinois, Chicago, IL 60611

[4]Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, Madison, WI 53706

* Equal Contribution

[†] Current Address: Corporate Research and Development, Solventum, Maplewood, MN 55144.

[§] Current Address: Department of Microbiology, University of Chicago, Chicago, IL 60637

[||] Current Address: Department of Biomedical Engineering, University of Illinois at Chicago, Chicago, IL 60607

[#] Current Address: Division of Molecular Genetics and Oncode Institute, Netherlands Cancer Institute, Amsterdam, The Netherlands

Short title: *K. pneumoniae* virulence influenced by anesthesia exposure

Keywords: *Klebsiella pneumoniae*, transposon insertion sequencing (INSeq/Tn-seq), pneumonia, propofol, ketamine, virulence

Correspondence to:

Mark J. Mandel
University of Wisconsin-Madison
Department of Medical Microbiology and Immunology
1550 Linden Drive, MSB 5155
Madison, WI 53706
Phone: (608) 261-1170
Email: mmandel@wisc.edu

Nancy E. Freitag
Department of Pharmaceutical Sciences
University of Illinois
Chicago, IL 60611
Phone: (312) 355-4903
Email: nfreitag@uic.edu

ABSTRACT (224 Words)

Klebsiella pneumoniae make up 85% of carbapenem-resistant Enterobacteriaceae
(CRE), bacteria that have become an urgent threat to public health. K. pneumoniae is largely
transmitted in healthcare settings, where inpatients and outpatients are often anesthetized with
the widely-used anesthetic induction agent propofol. Recent evidence obtained from rodent
infection models indicates that propofol exposure can dramatically increase host susceptibility to
microbial infections. Given that intensive care patients who are at a greater risk for K.
pneumoniae lung infections are often given propofol during their hospitalization, we investigated
the outcome of K. pneumoniae infections in mice briefly sedated with either propofol or
ketamine/xylazine as control. Propofol-sedated mice experienced more rapid dissemination
from the lungs to secondary sites of infection and their lungs exhibited more severe pathology.
Based on these observations, we investigated bacterial factors involved in infection and
dissemination in mice with propofol or ketamine/xylazine sedation using a high throughput
insertion sequencing (INSeq) approach. We identified numerous novel potential virulence
factors together with previously identified gene products, confirming the validity of our screen.
We further characterized a mutant lacking the phospholipid retrograde trafficking chaperone
MlaC and found that the degree of mutant attenuation was dependent upon sedation method.
These results highlight the importance of sedative choice when studying hospital-acquired
microbial infections and suggest that sedation can influence outcome of K. pneumoniae
infection and dissemination in animal models.

IMPORTANCE

Host sedation by either propofol or ketamine exposure differentially impacted the
severity of K. pneumoniae lung infection following intranasal inoculation of mice. While propofol-
sedated mice exhibited increased lung pathology, some bacterial mutants, such as those
lacking the MlaC gene product associated with the maintenance of inner and outer membrane

lipid asymmetry, exhibited more severe attenuation following propofol sedation versus

ketamine/xylazine. Given the dominating use of propofol in health care settings for the induction

and maintenance of anesthesia, procedural sedation, and sedation for intensive care patients,

this work provides important perspective as to how the choice of an anesthetic agent may

impact the outcome of healthcare-associated infections.

INTRODUCTION

Healthcare-associated infections (HAIs) pose the threat of increased morbidity for many

patients while serving as a major cause of mortality for intensive care patients. The anesthetic

induction agent propofol is widely used in critical care settings and is a common choice for light

sedation in the treatment for critically ill patients (1). Furthermore, patients requiring mechanical

ventilation are frequently administered sedatives (2) and are at increased risk for a variety of

complications including pneumonia, collectively referred to as ventilator-associated events

(VAEs) (3–6). The recent global pandemic of COVID-19, caused by SARS-CoV-2, has

highlighted risks of ventilator treatment (7–11). Mechanical ventilation increases the risk of

aspirating commensal bacteria into the lungs, and use of sedation during ventilation is

associated with an increased hazards ratio for VAEs (5–6).

The general anesthetic propofol has grown to become the predominant intravenous

anesthetic in clinical use throughout the world, lauded for its rapid onset and offset of sedation,

relatively mild side effects, and flexibility in both the induction and maintenance of anesthesia

(12, 13). While there have been multiple reports on propofol's immunomodulatory effects *in vitro*

(13–24), *in vivo* evidence has only recently begun to be reported. Klompas et al. correlated

propofol sedation in mechanically-ventilated patients with a significantly increased risk of VAEs,

including pneumonia, compared to patients not receiving propofol (5). Visvabharathy et al.

reported that acute propofol sedation dramatically increased susceptibility to both *Listeria*

monocytogenes and *Staphylococcus aureus* bloodstream infections in a mouse model and

additionally led to more severe pathology at primary sites of infection (25, 26). This susceptibility

correlated with a reduced capacity to effectively recruit mature immune effector cells to sites of

infection following propofol sedation. These propofol-associated influences on host immunity are

especially important to consider given that propofol is administered widely in both inpatient and

outpatient procedures and suggested to us that propofol could have a wider impact beyond the

intravenously administered gram-positive pathogens we had previously examined.

To explore the potential impact of propofol sedation on the outcome of respiratory

infections, we have made use of a mouse model of lung infection with the opportunistic

pathogen *Klebsiella pneumoniae*. *K. pneumoniae* is a gram-negative extracellular bacterium

that is ubiquitous in the water and soil of the environment and also colonizes the nasopharynx

and intestinal tract of humans and animals (27). It accounts for 3-8% of all nosocomial bacterial

infections including pneumonia, gram-negative bacteremia, and urinary tract infections (28). *K.*

pneumoniae is the second most common cause of gram-negative bacteremia in the United

States with 12-50% of *K. pneumoniae* bloodstream infections originating from the lungs (28–31).

A significantly worse outcome of bacteremia is associated with *K pneumoniae* infections from a

respiratory source (29). *K. pneumoniae* is a growing threat worldwide as a lethal nosocomial

pathogen that is rapidly acquiring antimicrobial resistance (32).

Using an intranasal model of *K pneumoniae* infection in mice, we compared the impact

of propofol versus ketamine/xylazine sedation on bacterial colonization of the lung, lung

pathology, and the dissemination of bacteria to distal organs. We further investigated whether

sedative choice influenced the requirement for distinct *K pneumoniae* virulence determinants

using a whole genome transposon insertion sequencing (INSeq) approach.

MATERIALS AND METHODS

Bacterial media and strains. *K. pneumoniae* and *Escherichia coli* strains used in this study are listed in **Table 1**. Strains were grown at 37 °C in lysogeny broth (LB; per liter, 10 g Bacto-tryptone [BD], 5 g yeast extract [BD], 10 g NaCl [Sigma-Aldrich]). When cells were grown in the presence of hygromycin, low-salt LB was used (i.e., 5 g NaCl per liter). When necessary, antibiotics (Gold Biotechnology) were added to the media at the following concentrations: rifampicin 30 µg ml^{-1}, hygromycin 100 µg ml^{-1}, kanamycin 50 µg ml^{-1}, and carbenicillin 100 µg ml^{-1}. When necessary, the following compounds were added to the media to the indicated final concentrations: sodium salicylate (Sigma-Aldrich): 2 mM; diaminopimelate (DAP; Sigma-Aldrich): 0.3 mM; arabinose (Gold Biotechnology): 100 mM. Growth media were solidified with 1.5% agar as needed. Standard molecular biology techniques were used to introduce plasmids into *E. coli* (i.e. electroporation) and into *K. pneumoniae* (i.e., electroporation or conjugation).

Animal infections. All animal procedures were approved by the University of Illinois Chicago Animal Care Committee and were conducted in the Biological Resources Laboratory. *K. pneumoniae* strains were cultured overnight with shaking (37°C, 180 rpm) in LB broth. The following morning, strains were subcultured 1:50 in fresh LB and incubated with shaking for one hour to an OD_{600} of 0.2-0.4. Optical density was adjusted to 0.2 OD_{600} with LB broth and bacteria were recovered by centrifugation, washed, and diluted to the desired CFU in sterile 1X phosphate-buffered saline (PBS) for infection. Strains were kept on ice until immediately before infection. Animals were sedated via tail vein injection with ketamine (20 mg/kg; Zoetis, Kalamazoo, MI) and xylazine (4 mg/kg; Akorn, Decatur, IL) in Intralipid as control or propofol (20 mg/kg; Zoetis) in Intralipid. For non-competitive infections, approximately 2.5 x 10^2 colony forming units (for propofol sedated mice) or 2.5 x 10^3 colony forming units (ketamine/xylazine sedated mice) of each strain was administered via intranasal route into 8- to 10-week old Swiss Webster mice (Envigo, Madison, WI) and for competitive infections, approximately 5 x 10^3 CFU of WT and a mutant were premixed and administered via intranasal route for a total 1 x 10^4 CFU

dose. For non-competitive infections, mice were sacrificed after 24 or 48 hours. Lungs, livers,

and spleens were isolated, homogenized, and plated in triplicate on LB agar containing

rifampicin for enumeration of bacterial burdens in each organ. For competitive infections, mice

were sacrificed after 24 hours and lungs were homogenized and plated in triplicate on LB agar

containing rifampicin and on LB agar containing rifampicin and kanamycin.

Histology. For histological analysis, mice were sacrificed individually via CO_2 asphyxiation.

Lungs were inflated with PBS at a flow rate no greater than 200 µl/second using a 3 ml syringe

with a 22 g needle inserted into the trachea. Once lungs were fully inflated and prior to removal

of the needle, the trachea was tied off with a surgeon's knot of suture thread. Lungs were

placed in 30 ml of neutral buffered formalin for 48 hours, then transferred into 70% ethanol.

Paraffin-embedded sections were prepared by the UIC Research Histology and Tissue Imaging

core facility. Five micron-thick sections were stained with hematoxylin and eosin (H&E) and

mounted. Digital scans of H&E-stained lung sections were captured on a Keyence digital

microscope and analyzed using ImageJ software.

H&E histopathological analyses. Mouse lung images standardized by containment of only the

left lobe and cranial and caudal portions of the right lobe were analyzed on ImageJ software by

an examiner blinded to sedative treatment. Total lesion area was quantified by the sum of

basophilic areas of inflammatory cell infiltration as a fraction of total lung area per image.

Number or lesions and average legion size composing the total lesion area were also

enumerated per whole lung slice.

Immunofluorescence. For immunofluorescent analysis, mice were sacrificed via isoflurane.

The lung vasculature was perfused with 10 ml ice cold PBS containing 50U/ml heparin to

remove blood and lungs were infused with one milliliter of neutral buffered formalin solution

(Millipore Sigma). Lungs were placed in 20 ml of neutral buffered formalin for 48 hours, then

transferred into 70% ethanol. Unstained, paraffin-embedded sections of 5µm thickness were

prepared by the UIC Research Histology and Tissue Imaging core facility. Paraffin was

dissolved in xylenes and sections were rehydrated in graded ethanol washes, followed by

washing in tris-buffered saline (TBS, pH 7.6). Antigen retrieval was carried out in 10 mM sodium

citrate buffer with 0.05% Tween-20 (pH 6.0) in a pressure cooker on high temperature for fifteen

minutes. Slides were blocked with Background Buster (Innovex, Richmond, CA). Primary

antibody was *K. pneumoniae* rabbit polyclonal (ThermoFisher, Waltham, MA) and secondary

antibody was goat anti-rabbit-FITC (Invitrogen, Carlsbad, CA). Slides were mounted with

ProLong Gold antifade with DAPI (ThermoFisher), imaged using a Zeiss Axio Imager A2 upright

microscope (Carl Zeiss, Thornwood, NY), and analyzed on Zen 2012 (Carl Zeiss) imaging

software.

Generation of KPPR1 transposon mutant library. KPPR1 and MJM2382 were grown

overnight in LB and LB carb DAP, respectively, at 37 °C with aeration. Saturated cultures were

diluted 1:80 in 3 ml and grown for 2 hours and 40 minutes at 37 °C. 2.5 ml of *E. coli* and 5 ml of

K. pneumoniae were pelleted and combined in 500 µl LB, which was spotted in fifty 10-µl spots

on LB. After 3.5 hours at 37 °C, each spot was swabbed into 500 µl LB, vortexed, and 50 µl was

spread onto LB plates containing kanamycin and salicylic acid for overnight incubation at 37 °C

(the presence of salicylic acid reduces the amount of capsule produced). An estimated 50,000

colonies were swabbed into 25 ml LB mixed with 12.5 ml 50% glycerol on ice and frozen at -

80˚C with a final density of 3×10^{11} CFU/ml.

INSeq analysis. Genomic DNA was extracted using a Maxwell 16 instrument (Promega,

Madison, WI). Transposon insertion junctions were amplified from three input samples and six

output samples according to the previously published protocol (33). Samples were single-end

(SE50) sequenced with an Illumina HiSeq 2500 platform (Illumina, San Diego, CA) at the Tufts

University Genomics Core Facility in Boston, MA with 45 million reads generated. FASTQ files

were first trimmed to the chromosomal sequence flanking each transposon insertion with

pyinseq v0.2 (https://github.com/mjmlab/pyinseq). Files were subsequently analyzed either with

the full pyinseq v0.2 pipeline to obtain cpm (normalized counts-per-million) data for each

replicate or with TRANSIT 3.8 (34) using the `resampling` algorithm against the KPPR1 genome

sequence CP009208.1 (35) with parameters -a -iN 5 -iC 10. Sequencing data were deposited at

NCBI SRA under accessions SRR6480865-SRR6480873 as detailed in **Table S1**. A table of

pyinseq analyzed values for each KPPR1 gene is posted as **Table S2**. Significant results from

the TRANSIT output are listed in **Table S3** and **Table S4**.

Generation of *K. pneumoniae* gene deletion strains. Deletions were made in the KPPR1

background using Lambda Red recombination (36). Knockout cassettes for each gene were

generated using the "KO" primers listed in **Table S3** to amplify the KanR cassette from pKD4

with homologous ends to chromosomal location of each gene. KPPR1 was electroporated with

pACBSR-Hyg to create strain MJM2698. This strain was electroporated with each of the

knockout cassettes. Strains with Kan resistance were selected and then patched onto low salt-

LB-Hyg plates to confirm the loss of pACBSR-Hyg. Kan-resistant and Hyg-sensitive isolates

were verified with PCR using the verification (V) primers listed in **Table S5**. The PCR product

was submitted for Sanger sequencing at the Northwestern University Sanger Sequencing Core

to confirm the integrity of the junctions where the knockout cassette replaced the gene of

interest. To remove the KanR cassette, strains were electroporated with pFLP-Hyg and

acquisition of the plasmid was selected on low salt-LB-Hyg media. Candidates were incubated

at 42 $^{\circ}$C overnight and then screened for loss of Kan resistance. Colonies that lost Kan

resistance were passaged for several days on LB plates and then screened for Hyg sensitivity

231 to confirm loss of pFLP-Hyg. PCR was used to confirm removal of the Kan cassette using the

232 verification (V) primers listed in **Table S5**.

234 **Construction of _K. pneumoniae_ complementation strains.** Complementation of deleted

235 genes was accomplished by integrating wild-type alleles of the gene at the presumed neutral

236 *att*Tn*7* chromosomal site using a site-specific Tn*7* transposon (**Figure S2**). Published vector

237 pJMP1185 expresses a Cas9 system with a constitutive promoter driving expression of *mRFP*

238 as a control (37). We first removed the *cas9* gene from the transposon to create pAD12.

239 Primers AD_254 and AD_255 were used to amplify pJMP1185 without the *cas9* gene. The

240 plasmid was ligated back together with T4 ligase and transformed into WM3064 *E. coli* cells.

241 Candidates were selected on LB-Kan plates, plasmid insertion was verified with PCR using

242 primers AD_191 and AD_192, and the correct sequence was confirmed with Oxford Nanopore

243 sequencing (Plasmidsaurus). To generate complement vectors, the *mRFP* gene was replaced

244 with one of the genes of interest to generate pAD14, pAD16, pAD18, or completely removed to

245 generate an empty vector control (pAD12b). Primers AD_258 and AD_259 were used to amplify

246 pAD12. The reaction was PCR purified and DpnI digested. Genes were amplified with either

247 AD_264/AD_265, AD_272/AD_273, or AD_276/AD_277 and then cloned into the backbone

248 using the NEB Hi-Fi assembly kit. For the empty vector control, primers DAT_535/DAT_536

249 were used to amplify the backbone and the Q5 site-directed mutagenesis kit (NEB) was used

250 for assembly. For the complementation plasmids and empty vector control, the assembly mix

251 was transformed into WM3064 *E. coli* cells and candidates were selected on LB-Kan. The

252 resulting plasmids were verified with Oxford Nanopore sequencing (Plasmidsaurus). Plasmids

253 were transformed into relevant *K. pneumoniae* strains using conjugation. Candidates were

254 selected using Kan resistance and verified using PCR (DAT_529/DAT_530) and Oxford

255 Nanopore sequencing (Plasmidsaurus).

SDS/EDTA susceptibility. To assess doubling time, strains were grown in LB broth at 37°C

with shaking for 3 h, with samples taken every 30 min. Serial dilutions were spot plated to

enumerate viable CFU and the doubling time was calculated from these points. To assess

growth on SDS/EDTA, cultures were grown to saturation in LB, serially diluted, and 5 µl was

spotted on LB plates or LB plates containing 0.1% SDS/1.5 mM EDTA. Plates were incubated

overnight at 37˚C and imaged with a Nikon D810 digital camera.

Statistical Analysis: For mouse experiments that compared multiple groups, bacterial burden

data was log10 transformed and analyzed via the Kruskal Wallis test with Dunn's post-hoc test.

Data graphing and analysis was conducted on GraphPad Prism software.

RESULTS

Sedation with the anesthetic propofol leads to increased lung colonization with *K.*

pneumoniae. Previous investigations into the impact of propofol on bloodstream infections

using the gram-positive pathogens *L. monocytogenes* and *S. aureus* revealed significant

alterations in bacterial colonization, growth, and pathology in target organs (25). Given the

prevalence of *K. pneumoniae* acquisition in health-care settings where propofol is widely used,

and to examine the impact of propofol on a model gram-negative pathogen, we assessed

whether propofol affected lung infection of *K. pneumoniae* via an intra-nasal inoculation route.

Female Swiss Webster mice were briefly sedated (5-10 minutes) with either intravenous

propofol or intravenous ketamine/xylazine and intranasally infected with $3x10^4$ CFU KPPR1.

Bacterial burdens were assessed at 24 and 48 hours post infection in the lung, liver, and spleen.

Brief propofol sedation had no measurable impact on lung burdens 24 hours post-infection, but

burdens were increased nearly 100-fold by 48 hours compared to ketamine/xylazine-sedated

controls (**Figure 1**). We also assessed the impact on dissemination from the lungs and again

found that while secondary organs had comparable burdens at 24 hours, average burdens in

both livers and spleens at 48 hours were more than 1,000-fold higher than ketamine/xylazine

sedated controls (**Figure 1**). We conclude that the increased susceptibility of murine hosts to

infection observed following propofol sedation is therefore broad enough to impact multiple

routes of infection with both gram-positive and gram-negative bacteria.

Propofol exacerbates lung pathology associated with *K. pneumoniae* infection. Previous

research using propofol sedation prior to infection identified gross changes in pathology at

primary sites of infection (25, 26). When examining lung histology, we observed that by 48

hours post-infection severe damage was evident in lungs regardless of sedation agent (**Figure

2A-B**), Quantification of these regions showed that propofol-sedated mice contained

significantly larger lung lesions while the total number of focal lesions did not differ based on

sedative agent (**Figure 2D-F**). Notably, only propofol-sedated mouse lungs featured large areas

of de-cellularized or denuded tissue containing little identifiable structure and only sporadic

immune infiltration. Immunofluorescent examination of these regions revealed that *K.*

pneumoniae had completely overrun local containment and appeared to have developed a

replicative niche (**Figure 2G-H**). It is likely that these bacteria-packed regions underlie the

disparate burdens observed between sedation groups.

**Transposon insertion sequencing identifies *K. pneumoniae* genes that are required for

outgrowth in the lung in a sedative-independent manner.** Genome-wide approaches for the

discovery of virulence factors required for *in vivo* growth have been applied in *K. pneumoniae*

previously with considerable success (38–43). In addition to examining the impact of propofol on

K. pneumoniae lung infection, we also sought to investigate what impact propofol might have on

the *K. pneumoniae* global virulence repertoire required for successful growth in the lung.

Transposon insertion sequencing (INSeq/Tn-seq) is a powerful, high resolution approach to

analyze whole genomes by coupling Mariner transposon mutagenesis with high-throughput

sequencing to rapidly and comprehensively quantify the relative fitness of large pools of

transposon mutants (44–46). We applied this approach by mutagenizing *K. pneumoniae* strain

KPPR1, sedating mice with either propofol or ketamine/xylazine, followed by the infection of

mice via the intranasal route. We generated a 50,000 mutant transposon library in KPPR1 using

the plasmid pJNW684 (46). Hits were observed across the chromosome with an average of

approximately 9.1 insertions per kb. Three propofol-sedated and three ketamine/xylazine

control-sedated mice were intranasally infected with the library at a dose of 6 x 10^5 CFU for 24

hours, after which the lungs were harvested and genomic DNA was extracted. The 24 h time

point was selected to identify mutants with fitness defects in advance of the lethality differences

that begin to occur at 48 h with the use of a higher CFU dose and so that the analysis was not

complicated by the high proportion of lethality at the later time point. The higher CFU dose of 6 x

10^5 CFU was selected to increase mutant representation within the library for the subsequent

INSeq analyses.

We analyzed the output data to examine genes that displayed fitness defects under both

sedatives. Using the software TRANSIT (34), we identified 217 genes for which mutants

exhibited significant depletion (and 2 genes for which mutants exhibited enrichment) under both

sedative conditions (**Fig. 3**; **Tables S3, S4**). The depleted mutants match factors identified in

previous transposon insertion sequencing studies, including *aroA, copA, hisA, ilvC, leuA, leuB,*

serA, serB, trpE, and *tyrA* (Fig. 3) (43, 47). Identification of conserved responses across the two

sedatives, and identification of responses consistent with published studies supports the quality

of the INSeq analysis and indicates that targeting of these pathways is likely to yield benefits to

the extent that propofol responses are conserved in human patients.

Identification of *K. pneumoniae* mutants that exhibit sedative-dependent responses. We

next asked whether the INSeq data had resolution to identify genes that exhibited differential

responses to the two sedatives. Examination of **Figure 3** revealed variation in the response to

the sedatives, even in mutants that were abundant in the input counts. A direct analysis of

mutants recovered from the two output conditions (i.e., propofol-anesthetized mice compared to

ketamine/xylazine-anesthetized mice) did not yield genes with significant enrichment or

depletion when the TRANSIT statistical test was applied. We reasoned that this was due to the

variation among transposon counts from the animal-recovered samples—in contrast to the

analysis in **Figure 3** in which variation from the mouse output treatment was compared to the

largely invariant input control. Therefore, we scanned the INSeq data for genes that exhibited

greater depletion under one sedative. Mutants in the genes *virK* (VK055_0094/RS00455)*,*

VK055_1993 (RS09970), and *glnB* (VK055_4623/RS23395) were more substantially depleted

following ketamine/xylazine sedation, while the genes *VK055_1398, VK055_3638 (RS18420),*

VK055_3462 (RS17525), and *fepC* (VK055_1930/ RS09665) were likely to be more important

following propofol sedation (**Figure S1**). Additionally, we chose to investigate *mlaC*

(VK055_3875) as it was identified as having a strong defect regardless of sedative choice and

was representative of most of the *mla* genes that encode a phospholipid transport system

(**Figure S3**).

To assess whether our approach could identify bona fide sedative-dependent mutants,

we generated deletion mutants and tested each individually in competition with the wild-type

parent. Competition infections were performed using a 1:1 mixture of KPPR1 and single deletion

mutants. Some of the gene candidates exhibited sedative-dependent phenotypes consistent

with the trends in the original transposon mutant selection (**Figure S1**). This was observed for

the *ΔglnB and ΔVK055_1993* mutants that exhibited competitive defects only in the

ketamine/xylazine treatment*,* and for *ΔVK055_1398* that demonstrated a defect only in propofol.

The *ΔmlaC* mutant exhibited a competition defect in both conditions as predicted. Trends for the

ΔfepC mutant also matched the INSeq data with slight defects under both conditions and more

pronounced effect in the propofol-sedated host (**Figure S1**). However, other mutants exhibited

phenotypes in the direct competition assays that differed from the trends observed in the INSeq

data. The *ΔVK055_3462* mutant did not exhibit a direct competition phenotype in either

condition. The *ΔVK055_3638* mutant yielded a mild and shared competitive defect in both

sedatives. Unexpectedly, the ΔvirK mutant exhibited a propofol defect in the direct competition,

in contrast to the ketamine/xylazine defect in the INSeq pooled competition experiment (**Figure**

S1). Although distinct from the result from INSeq, this outcome provides an additional mutant to

use as a probe for host function under propofol. In summary, despite weak support for sedative-

dependent mutant behavior in the INSeq data set, approximately half of the mutants that we

examined validated with expected phenotypes and provide useful tools to probe the differential

anesthesia response.

Conserved virulence factors can exhibit sedative-dependent responses during mono-

infection. We next wished to ask whether genes that were broadly required for *K. pneumoniae*

colonization were similarly required under propofol sedation as under ketamine/xylazine. For

this question, we focused on the *mlaC* mutant described above. We took advantage of the

known detergent sensitivity of *mla* mutants in *E. coli* (48) to further characterize the Δ*mlaC*

mutant. The *K. pneumoniae* Δ*mlaC* mutant was found to exhibit sensitivity to SDS and EDTA, a

phenotype that could be reversed upon complementation with the wild-type *mlaC* expressed at

the chromosomal *att*Tn*7* site (**Figure 5A, Figure S2**).

Given that propofol sedation appeared to significantly enhance mouse susceptibility to *K.*

pneumoniae infection, we identified an infectious dose for propofol sedated animals that

enabled mouse survival beyond 48 hours so as to provide a more extended time window for

assessment of potential mutant phenotypes. An intranasal dose of 250 CFU of KPPR1 was

found to establish robust and reproducible levels of infection despite being 10 to 100-fold lower

than doses routinely reported in the literature (**Figure 5**) (49–51). This dose was optimized for

intravenous propofol sedation as mice infected with a 10-fold higher inoculum at 2,500 CFU

could not meet humane criteria for the study duration of 48 hours. For the comparison of

propofol sedated animals with single Δ*mlaC* infections under ketamine sedation, a ten-fold

increase to the inoculum was required for animals sedated with ketamine to detect any

difference between attenuated mutant strains and WT as animals rapidly cleared lower doses of

WT *K. pneumoniae* (**Figure 5**). The requirement for *mlaC* was validated for mono-infections

under both sedative conditions and could be fully complemented by the introduction of *mlaC* at

a neutral chromosomal locus (**Figure 5, S2**). While the requirement for a 10-fold lower CFU

inoculum in propofol-sedated animals (given the increased susceptibility of these animals to *K.*

pneumoniae infection) complicates a direct comparison with ketamine/xylazine sedated mouse

infections, the Δ*mlaC* mutant is clearly highly attenuated in propofol-sedated mice with lung

burdens comparable to ketamine-sedated controls (**Figure 5**). For example, mice infected with

WT *K. pneumoniae* following propofol or ketamine/xylazine sedation exhibited comparable

numbers of bacterial CFU in the lung despite the 10-fold lower dose following ketamine

sedation. While the WT bacterial numbers in the lung are comparable for both groups, the

Δ*mlaC* mutant is recovered in numbers that are 100-1000 fold lower following propofol sedation

(**Figure 5**). Overall, our results support the requirement for specific *K. pneumoniae* virulence

factors to establish lung infections regardless of the type of sedative used during mouse

intranasal inoculation.

DISCUSSION

Previous work identified the significant impact of brief propofol exposure on the ability of

an infected host to combat bacterial bloodstream infections with either *L. monocytogenes* or *S.*

aureus, both gram-positive bacterial pathogens (25, 26). Here, we have demonstrated that

propofol also increases a murine host's susceptibility to both a new route of infection (intranasal)

and to a gram-negative pathogen, *K. pneumoniae*. Propofol sedation increased lung burdens

approximately 100-fold compared to ketamine/xylazine-sedated mice and furthermore increased

dissemination to secondary organs approximately 1,000-fold at 48 hours post-infection (**Figure**

1). We further identified that propofol sedation resulted in significantly larger areas of denuded

or destroyed lung tissue (**Figure 2**), likely forming a replicative niche for rapidly dividing *K.*

pneumoniae in vivo and possibly accounting for the increased lung burdens measured at 48

hours post-infection. Serious disseminated infections also resulted from a low dose inoculum

(200 CFU) indicating the increased susceptibility of propofol sedated mice to *K. pneumoniae*

lung infection (**Figure 5**). These data clearly demonstrate that propofol negatively impacts the

host's ability to address *K. pneumoniae* lung infection and subsequent bacteremia. Whether

these changes in host immune responses resemble those previously reported for bacterial

bloodstream infections, which included dramatic reductions in the populations of mature

macrophages (25, 26) has not yet been determined. Lung histology indicates abundant immune

cell infiltration into the infected lungs with evident differences in the resultant pathology (**Figure**

2).

425   Genome-wide mutagenic approaches such as signature-tagged mutagenesis (STM) and

transposon mutagenesis have previously been applied as *K. pneumoniae in vivo* virulence

screens and have identified dozens of putative virulence factors, which have been reviewed by

Paczosa and Mecsas (30). The screen described herein yielded novel gene products by

considering sedative exposure as a key variable in subsequent lung outgrowth. By comparing

outputs from infected lungs to their inputs, we were able to quantify the relative requirement of

numerous genes following either ketamine/xylazine or propofol sedation. This approach enabled

us to identify putative novel virulence determinants and begin to define the broad impact that

sedation has on pathogen fitness toward lung colonization. The choice of ketamine/xylazine for

comparison was based both on the common use of this anesthetic/sedative in animal

experiments, and because the drugs were previously shown to have minimal effects on the

control of bloodstream bacterial infections (25, 26). Interestingly, clear effects on host infection

were also evident following ketamine/xylazine sedation, suggesting that sedative use in general

may impact host immunity, albeit perhaps to different degrees. This finding is consistent with a

recent report from Gage et al indicating that ketamine sedation negatively impacted bacterial

clearance and lung tissue pathology using a *Pseudomonas aeruginosa* pneumonia model (52).

Based on the results of the INSeq analysis and previously reported contributions of the

mla locus to bacterial virulence (2), the *K. pneumoniae mlaC* mutant was selected for further

characterization. The Mla (maintenance of outer membrane lipid asymmetry) pathway has been

well characterized in *E. coli* (48). Mla comprises a transport system that shuttles phospholipids

from the outer membrane to the inner membrane (48). MlaFEDB forms an ABC transporter,

while MlaA is a lipoprotein attached to the outer membrane and MlaC is the periplasmic shuttle

between the two. Loss of *mlaC* resulted in a significant defect under both sedative conditions

(**Figure 5**). A direct comparison between propofol and ketamine/xylazine sedated animals for

mutant phenotypes was somewhat constrained by the need to use increased CFU to establish

infection in ketamine/xylazine sedated mice. Virulence defects similar to those observed for the

mlaC mutant would be predicted for *mlaA*, *mlaD*, *mlaE*, and *mlaF* based on their representation

in the INSeq screen (**Figure S3**). In contrast, the loss of *mlaB* does not appear to carry a

significant defect based on the INSeq results, a reasonable finding considering that MlaB is a

minor subunit in the transporter. *mlaA* has been recently confirmed as a virulence requirement

for *K. pneumoniae* lung infections in mice (53). The perturbation of membrane lipid homeostatic

mechanism is therefore likely a critical factor in the resistance of *K. pneumoniae* to host

antimicrobial strategies, and this host defense may be somehow enhanced by propofol

exposure, resulting in enhanced clearance of *mlaC* mutants.

Our dataset suggests numerous genes that are potentially differentially required

depending on host anesthetic exposure. Further validation and characterization of these gene

products is needed to provide clarity to these findings, and an expansion of the INSeq approach

in the anesthetized animals would likely provide valuable additional candidates to pursue. This

prediction is based on our finding of multiple interesting hits from our preliminary that were

validated despite the INSeq approach not being sufficiently powered to reveal global gene

requirements in propofol- versus ketamine/xylazine-sedated animals. Our results support a

growing body of evidence that the choice of sedative used for anesthesia can profoundly

influence the course of bacterial infection, our work goes on to show that the virulence gene

repertoire required for infection may differ dependent upon sedative choice. This finding has

important implications for the selection of potential new drug targets for *K. pneumoniae*

infections; the most effective targets are likely to be those gene products that are required

regardless of the type of sedation that may be used. Taken together, these results reveal that

propofol sedation leads to the exacerbation of lung infection in a murine model of experimental

pneumonia caused by the bacterial pathogen *K. pneumoniae.* Future studies aimed towards

identifying colonization and virulence factors required under multiple sedation conditions should

clarify which aspects of host immunity are negatively impacted by sedative exposure.

ACKNOWLEDGMENTS

We thank Alan Hauser and Nengding (Julie) Wang for plasmid pJNW684 and use of the

Promega Maxwell instrument, Jason Peters for pJMP1185 and advice on modifying the plasmid,

and Emanuel Burgos for computational assistance. We thank Mercy Obarewon for assistance

with histological analyses. This work was supported by the Chicago Biomedical Consortium

Catalyst Award (to M.J.M. and N.E.F.), NIH grant R21AI166098 (to N.E.F. and M.J.M.), and NIH

grant R35GM148385 (to M.J.M.). Its contents are solely the responsibility of the authors and do

not necessarily represent the official views of the funding source.

485
486   **Table 1. Bacterial strain and plasmid list.**

Strain	Relevant Genotype	Reference
K. pneumoniae		
MJM2383	KPPR1	(35)
MJM2402	Transposon mutant library in KPPR1	This study
MJM2698	MJM2383 pACBSR-Hyg	This study
MJM2793	KPPR1 Δ*VK055_3462*::*kan*	This study
MJM2794	KPPR1 Δ*VK055_1993*::*kan*	This study

MJM2795	KPPR1 Δ*virK*(*VK055_0094*)::*kan*	This study
MJM2796	KPPR1 Δ*fepC*(*VK055_1930*)::*kan*	This study
MJM2797	KPPR1 Δ*mlaC*(*VK055_3875*)::*kan*	This study
MJM2784	KPPR1 Δ*glnB*(*VK055_4623*)::*kan*	This study
MJM2816	KPPR1 Δ*VK055_3638*::*kan*	This study
MJM3001	KPPR1 Δ*VK055_1398*::*kan*	This study
MJM5804	KPPR1 Δ*mlaC*	This study
MJM6053	KPPR1 Δ*mlaC att*Tn*7*::*mlaC*	This study
MJM6260	KPPR1 *att*Tn*7*::*kan*	This study
MJM6264	KPPR1 Δ*mlaC att*Tn*7*::*kan*	This study
E. coli		
MJM537	DH5α λpir	(54)
MJM1424 (β3914)	F- RP4-2-Tc::Mu Δ*dapA*::(erm-pir) *gyrA*462 zei-298::Tn10 (ErmR KanR TetR)	(55)
MJM2382	MJM1424 pJNW684	(46)
MJM534	CC118 λpir / pEVS104	(56)
MJM1649	BW25141 / pKD4	(57)
MJM2532	MJM1424 pACBSR-Hyg	This study
MJM2369	pFLP-Hyg	(36)
MJM4078	WM6026 dap-, pir+, attTn7::acrIIA4, recA1, dapA::pir / pJMP1039	(58)
MJM4080	WM6026 dap-, pir+, attTn7::acrIIA4, recA1, dapA::pir / pJMP1185	(58)
MJM5331 (WM3064)	*thrB1004 pro thi rpsL hsdS lacZ*ΔM15 RP4-1360 Δ(*araBAD*)567 Δ*dapA1341*::[*erm pir*]	(59)
MJM5713	WM3064 λpir / pAD12	This study
MJM5851	WM3064 λpir / pAD18	This study
MJM6259	WM3064 λpir / pAD12b	This study
Plasmids	Genotype/Description	
pJNW684	Transposon vector	(46)
pKD4	*FRT-kan-FRT* template	(57)
pACBSR-Hyg	Lambda-Red genes	(36)
pFLP-Hyg	FLP recombinase	(36)
pJMP1039	Helper for transposon on pJMP1185	(58)
pJMP1185	Transposon vector containing *cas9* system	(58)
pAD12	pJMP1185 without *cas9*	This study
pAD12b	pAD12 without *mRFP*	This study
pAD18	pAD12 with *mlaC*	This study

489
490
Supplemental Tables S1-S5:
• **Table S1. NCBI SRA Accession numbers.** Data deposition for the INSeq samples.
• **Table S2. INSeq data analysis for all samples.** Shown are counts-per-million (cpm) per
gene for each sample from the pyinseq pipeline.
• **Table S3. Shared depleted mutants.** Genes for which mutants showed a significant
depletion under both sedatives, from the TRANSIT pipeline.
• **Table S4. Shared enriched mutants.** Genes for which mutants showed a significant
enrichment under both sedatives, from the TRANSIT pipeline.
• **Table S5. Primer list.** Primers used for <u>P</u>CR during construction (P), or for <u>v</u>erification by
PCR and/or Sanger sequencing to verify insertion/orientation.
501

502
FIGURE LEGENDS

Figure 1. Brief propofol sedation increases *Klebsiella* lung burdens and dissemination to

secondary tissues. Swiss Webster mice were sedated with either ketamine/xylazine (closed

circles) or propofol (open circles) and infected with 3×10^4 CFU intra-nasally and lung, liver, and

spleen burdens were assessed at 24 and 48 hours post infection. Statistics were calculated

using the Mann Whitney U test (*, $p \le 0.05$; **, $p \le 0.01$).

509
Figure 2. Histological analysis of *K. pneumoniae* lung pathology 48 hours post-infection.

Swiss Webster female mice were infected with 1×10^2 CFU KPPR1. Representative H&E images

of *Kp* infected mouse whole lung tissue 48H post infection after propofol (A), ketamine/xylazine

(B) sedation, or uninfected control (C). (D) Quantification of total lesioned area per lung section

(n=5 per treatment). (E) Area of individual lesion sizes. (F) Quantification of number of lesions in

whole lung sections from left and right (cranial and caudal) lung lobes (n=5 per treatment). (G-

H) Representative H&E (G) and Immunofluorescent staining (H) of propofol-sedated mouse

lungs with the propofol specific histopathological feature of acellular denuded regions packed

with *K. pneumoniae*. Black bar represents 100µm. Statistics were calculated using an unpaired,

two-tailed Student's t-test (*, $p \le 0.05$; ****, p<.0001).

Figure 3: Transposon insertion sequencing results for all genes following intranasal

infection in propofol-sedated mice and ketamine/xylazine-sedated mice. Each *K.*

pneumoniae gene that exhibited transposon insertions in our input pool (as defined in our

analysis) is represented twice in the plot, showing the performance of mutants in the gene

during infection in propofol-sedated mice vs input (circles) and in ketamine/xylazine-sedated

mice vs input (triangles). These data are plotted against the the representation of those mutants

in the input pool. Genes that are to the right of the graph have a higher representation of

mutants in the input pool and tend to produce more robust data in the INSeq analysis (e.g.,

larger genes with more available sites for transposon insertion, and genes for which transposon

insertion does not impair growth in culture). Genes for which mutants showed a significant

depletion (blue, n=217) or enrichment (red, n=2) under both sedatives are highlighted. Select

genes are labeled, with a full listing of the highlighted genes listed in Tables S3 and S4,

respectively.

Figure 4. Competitive infections with in-frame deletion mutants. Competitive indices of

each mutant against WT 24 hours post-infection (n=4 per group). Mice were infected with

approximately 5×10^3 CFU of WT and a mutant strain that were premixed and administered via

intranasal route for a total 1×10^4 CFU dose. Statistics on lung titers were calculated using an

unpaired, two-tailed Student's T-test. As noted in the manuscript text, depletion under INSeq

was statistically significant under both sedatives for *mlaC* but was not significant for mutants

under individual sedatives; the INSeq designations shown are based on the trends in Figure S1.

Figure 5. *mlaC* mutant demonstrates defects in a sedative-dependent manner in

monoculture infections. (A) Saturated cultures were serially diluted and spotted onto specified

media types, then grown at 37°C overnight. The growth defect of the *mlaC* mutant on

SDS/EDTA is able to be complemented. (B/C) Female Swiss Webster mice (n=8-9 per group)

were ketamine/xylazine sedated or propofol-sedated and infected with 2500 CFU

(ketamine/xylazine sedated mice) or 250 CFU (propofol sedated mice) intranasally. The

increased inoculum size for ketamine/xylazine sedated animals was necessary as mice cleared

the infections at the lower (250 CFU) inoculum dose. Organs were harvested at 48 h post-

infection for determination of bacterial CFU. Statistics were calculated using a two-tailed Mann-

Whitney U test (*, $p \leq 0.05$; **, $p \leq 0.01$) and significance is compared to KPPR1 under the

same sedative.

REFERENCES

1. Patel SB, Kress JP. 2012. Sedation and analgesia in the mechanically ventilated patient. Am J Respir Crit Care Med 185:486–497.

2. Caroff DA, Szumita PM, Klompas M. 2016. The relationship between sedatives, sedative strategy, and healthcare-associated infection: A systematic review. Infect Control Hosp Epidemiol 37:1234–1242.

3. Klompas M. 2019. Ventilator-associated events: What they are and what they are not. Respir Care 64:953–961.

4. Klompas M, Li L, Szumita P, Kleinman K, Murphy MV, CDC Prevention Epicenters Program. 2016. Associations between different sedatives and ventilator-associated events, length of stay, and mortality in patients who were mechanically ventilated. Chest 149:1373–1379.

5. de Irala Estévez J, Fernández-Crehuet Navajas R, Díaz Molina C, Martínez de la Concha D, Salcedo Leal I, Masa Calles J. 1997. Risk factors for pneumonia, bacteremia, and urinary tract infection in an intensive care unit. Med Clin (Barc) 109:733–737.

6. Velly L, Gayat E, Quintard H, Weiss E, De Jong A, Cuvillon P, Audibert G, Amour J, Beaussier M, Biais M, Bloc S, Bonnet MP, Bouzat P, Brezac G, Dahyot-Fizelier C, Dahmani S, de Queiroz M, Di Maria S, Ecoffey C, Futier E, Geeraerts T, Jaber H, Heyer L, Hoteit R, Joannes-Boyau O, Kern D, Langeron O, Lasocki S, Launey Y, le Saché F, Lukaszewicz AC, Maurice-Szamburski A, Mayeur N, Michel F, Minville V, Mirek S, Montravers P, Morau E, Muller L, Muret J, Nouette-Gaulain K, Orban JC, Orliaguet G, Perrigault PF, Plantet F, Pottecher J, Quesnel C, Reubrecht V, Rozec B, Tavernier B, Veber B, Veyckmans F, Charbonneau H, Constant I, Frasca D, Fischer M-O, Huraux C, Blet A, Garnier M. 2020. Guidelines: Anaesthesia in the context of COVID-19 pandemic. Anaesth Crit Care Pain Med 39:395–415.

7. Yamamoto K. 2020. Risk of propofol use for sedation in COVID-19 patient. Anaesthesiol Intensive Ther 52:354–355.

8. Hirota K, Lambert DG. 2020. Propofol and SARS-CoV-2 infection. Br J Anaesth 125:e475–e476.

9. Babu VK, Rojas P, Perez Del Nogal G, Garcia-Fernandez A. 2023. Unusual case of propofol-related infusion syndrome complicating severe COVID-19 ARDS. BMJ Case Rep 16:e249456.

10. Koutsogiannaki S, Bu W, Maisat W, Manzor M, Zhang Z, Ohto U, Eckenhoff RG, Yuki K. 2022. Propofol directly binds to and inhibits TLR7. FASEB J 36:e22481.

11. Vasileiou I, Xanthos T, Koudouna E, Perrea D, Klonaris C, Katsargyris A, Papadimitriou L. 2009. Propofol: a review of its non-anaesthetic effects. Eur J Pharmacol 605:1–8.

597   12.   Darnobid JA. 2015. The pharmacology of total intravenous anesthesia. Int Anesthesiol
598   Clin 53:13–27.

599   13.   Mikawa K, Akamatsu H, Nishina K, Shiga M, Maekawa N, Obara H, Niwa Y. 1998.
600   Propofol inhibits human neutrophil functions. Anesth Analg 87:695–700.

601   14.   Chen R-M, Wu C-H, Chang H-C, Wu G-J, Lin Y-L, Sheu J-R, Chen T-L. 2003. Propofol
602   suppresses macrophage functions and modulates mitochondrial membrane potential
603   and cellular adenosine triphosphate synthesis. Anesthesiology 98:1178–1185.

604   15.   Fourcade O, Simon M-F, Litt L, Samii K, Chap H. 2004. Propofol inhibits human platelet
605   aggregation induced by proinflammatory lipid mediators. Anesth Analg 99:393–398.

606   16.   Hsing C-H, Lin M-C, Choi P-C, Huang W-C, Kai J-I, Tsai C-C, Cheng Y-L, Hsieh C-Y,
607   Wang C-Y, Chang Y-P, Chen Y-H, Chen C-L, Lin C-F. 2011. Anesthetic propofol
608   reduces endotoxic inflammation by inhibiting reactive oxygen species-regulated
609   Akt/IKKβ/NF-κB signaling. PLoS One 6:e17598.

610   17.   Jawan B, Kao Y-H, Goto S, Pan M-C, Lin Y-C, Hsu L-W, Nakano T, Lai C-Y, Sun C-K,
611   Cheng Y-F, Tai M-H, Eng H-L, Wang C-S, Huang C-J, Lin C-R, Chen C-L. 2008.
612   Propofol pretreatment attenuates LPS-induced granulocyte-macrophage colony-
613   stimulating factor production in cultured hepatocytes by suppressing MAPK/ERK activity
614   and NF-κB translocation. Toxicol Appl Pharmacol 229:362–373.

615   18.   Inada T, Kubo K, Ueshima H, Shingu K. 2011. Intravenous anesthetic propofol
616   suppresses prostaglandin E2 production in murine dendritic cells. J Immunotoxicol
617   8:359–366.

618   19.   Meng T, Yu J, Lei Z, Wu J, Wang S, Bo Q, Zhang X, Ma Z, Yu J. 2013. Propofol reduces
619   lipopolysaccharide-induced, NADPH oxidase (NOX 2) mediated TNF- α and IL-6
620   production in macrophages. Clin Dev Immunol 2013:325481.

621   20.   Ren X, Lv F, Fang B, Liu S, Lv H, He G, Ma H, Cao Y, Wang Y. 2014. Anesthetic agent
622   propofol inhibits myeloid differentiation factor 88-dependent and independent signaling
623   and mitigates lipopolysaccharide-mediated reactive oxygen species production in human
624   neutrophils in vitro. Eur J Pharmacol 744:164–172.

625   21.   Inada T, Hirota K, Shingu K. 2015. Intravenous anesthetic propofol suppresses
626   prostaglandin E2 and cysteinyl leukotriene production and reduces edema formation in
627   arachidonic acid-induced ear inflammation. J Immunotoxicol 12:261–265.

628   22.   Li S, Wang C-X, Liu N-Z, Liu P. 2015. Anti-inflammatory effects of propofol on
629   lipopolysaccharides-treated rat hepatic Kupffer cells. Cell Biochem Biophys 71:845–850.

630   23.   Okuno T, Koutsogiannaki S, Ohba M, Chamberlain M, Bu W, Lin F-Y, Eckenhoff RG,
631   Yokomizo T, Yuki K. 2017. Intravenous anesthetic propofol binds to 5-lipoxygenase and
632   attenuates leukotriene B4 production. FASEB J 31:1584–1594.

24.   Meier A, Chien J, Hobohm L, Patras KA, Nizet V, Corriden R. 2019. Inhibition of human

neutrophil extracellular trap (NET) production by propofol and lipid emulsion. Front
Pharmacol 10:323.

25.   Visvabharathy L, Freitag NE. 2017. Propofol sedation exacerbates kidney pathology and
dissemination of bacteria during *Staphylococcus aureus* bloodstream infections. Infect
Immun 85:10.1128/iai.00097-17.

26.   Visvabharathy L, Xayarath B, Weinberg G, Shilling RA, Freitag NE. 2015. Propofol
increases host susceptibility to microbial infection by reducing subpopulations of mature
immune effector cells at sites of infection. PLoS One 10:e0138043.

27.   Dai P, Hu D. 2022. The making of hypervirulent *Klebsiella pneumoniae*. J Clin Lab Anal
36:e24743.

28.   Ashurst JV, Dawson A. 2025. Klebsiella pneumonia. StatPearls. StatPearls Publishing,
Treasure Island (FL).

29.   Al-Hasan MN, Lahr BD, Eckel-Passow JE, Baddour LM. 2010. Epidemiology and
outcome of *Klebsiella* species bloodstream infection: a population-based study. Mayo
Clin Proc 85:139–144.

30.   Paczosa MK, Mecsas J. 2016. *Klebsiella pneumoniae*: Going on the Offense with a
Strong Defense. Microbiol Mol Biol Rev 80:629–661.

31.   Holmes CL, Wilcox AE, Forsyth V, Smith SN, Moricz BS, Unverdorben LV, Mason S, Wu
W, Zhao L, Mobley HLT, Bachman MA. 2023. *Klebsiella pneumoniae* causes bacteremia
using factors that mediate tissue-specific fitness and resistance to oxidative stress. PLoS
Pathog 19:e1011233.

32.   Wyres KL, Holt KE. 2016. *Klebsiella pneumoniae* Population Genomics and
Antimicrobial-Resistant Clones. Trends Microbiol 24:944–956.

33.   Goodman AL, Wu M, Gordon JI. 2011. Identifying microbial fitness determinants by
insertion sequencing using genome-wide transposon mutant libraries. Nat Protoc
6:1969–1980.

34.   DeJesus MA, Ambadipudi C, Baker R, Sassetti C, Ioerger TR. 2015. TRANSIT--A
Software Tool for Himar1 TnSeq Analysis. PLoS Comput Biol 11:e1004401.

35.   Broberg CA, Wu W, Cavalcoli JD, Miller VL, Bachman MA. 2014. Complete genome
sequence of *Klebsiella pneumoniae* strain ATCC 43816 KPPR1, a rifampin-resistant
mutant commonly used in animal, genetic, and molecular biology studies. Genome
Announc 2:e00924-14.

36.   Huang T-W, Lam I, Chang H-Y, Tsai S-F, Palsson BO, Charusanti P. 2014. Capsule
deletion via a λ-Red knockout system perturbs biofilm formation and fimbriae expression
in *Klebsiella pneumoniae* MGH 78578. BMC Res Notes 7:13.

37.   Banta AB, Ward RD, Tran JS, Bacon EE, Peters JM. 2020. Programmable gene

knockdown in diverse bacteria using mobile-CRISPRi. Curr Protoc Microbiol 59:e130.

38.   Maroncle N, Balestrino D, Rich C, Forestier C. 2002. Identification of *Klebsiella*
pneumoniae genes involved in intestinal colonization and adhesion using signature-
tagged mutagenesis. Infect Immun 70:4729–4734.

39.   Struve C, Forestier C, Krogfelt KA. 2003. Application of a novel multi-screening
signature-tagged mutagenesis assay for identification of *Klebsiella pneumoniae* genes
essential in colonization and infection. Microbiology 149:167–176.

40.   Lawlor MS, Hsu J, Rick PD, Miller VL. 2005. Identification of *Klebsiella pneumoniae*
virulence determinants using an intranasal infection model. Mol Microbiol 58:1054–1073.

41.   Lau HY, Clegg S, Moore TA. 2007. Identification of *Klebsiella pneumoniae* genes
uniquely expressed in a strain virulent using a murine model of bacterial pneumonia.
Microb Pathog 42:148–155.

42.   Tu Y-C, Lu M-C, Chiang M-K, Huang S-P, Peng H-L, Chang H-Y, Jan M-S, Lai Y-C.
2009. Genetic requirements for *Klebsiella pneumoniae*-induced liver abscess in an oral
infection model. Infect Immun 77:2657–2671.

43.   Silver RJ, Paczosa MK, McCabe AL, Balada-Llasat J-M, Baleja JD, Mecsas J. 2019.
Amino acid biosynthetic pathways are required for *Klebsiella pneumoniae* growth in
immunocompromised lungs and are druggable targets during infection. Antimicrob
Agents Chemother 63:e02674-18.

44.   Goodman AL, McNulty NP, Zhao Y, Leip D, Mitra RD, Lozupone CA, Knight R, Gordon
JI. 2009. Identifying genetic determinants needed to establish a human gut symbiont in
its habitat. Cell Host Microbe 6:279–289.

45.   Brooks JF 2nd, Gyllborg MC, Cronin DC, Quillin SJ, Mallama CA, Foxall R, Whistler C,
Goodman AL, Mandel MJ. 2014. Global discovery of colonization determinants in the
squid symbiont *Vibrio fischeri*. Proc Natl Acad Sci U S A 111:17284–17289.

46.   Wang N, Ozer EA, Mandel MJ, Hauser AR. 2014. Genome-wide identification of
Acinetobacter baumannii genes necessary for persistence in the lung. mBio 5:e01163-
14.

47.   Bachman MA, Breen P, Deornellas V, Mu Q, Zhao L, Wu W, Cavalcoli JD, Mobley HLT.
2015. Genome-wide identification of *Klebsiella pneumoniae* fitness genes during lung
infection. MBio 6:e00775.

48.   Malinverni JC, Silhavy TJ. 2009. An ABC transport system that maintains lipid
asymmetry in the gram-negative outer membrane. Proc Natl Acad Sci U S A 106:8009–
8014.

49.   Bachman MA, Oyler JE, Burns SH, Caza M, Lépine F, Dozois CM, Weiser JN. 2011.
Klebsiella pneumoniae yersiniabactin promotes respiratory tract infection through
evasion of lipocalin 2. Infect Immun 79:3309–3316.

50. Vornhagen J, Sun Y, Breen P, Forsyth V, Zhao L, Mobley HLT, Bachman MA. 2019. The
Klebsiella pneumoniae citrate synthase gene, *gltA*, influences site specific fitness during
infection. PLoS Pathog 15:e1008010.

51. Walker KA, Miner TA, Palacios M, Trzilova D, Frederick DR, Broberg CA, Sepúlveda VE,
Quinn JD, Miller VL. 2019. A *Klebsiella pneumoniae* regulatory mutant has reduced
capsule expression but retains hypermucoviscosity. MBio 10.

52. Gage H, Hannah SM, Hancock B, Cornax I, Munguia J, Olson J, Bjånes E, Zurich R,
Hoffman A, Askarian F, Tong K, Liu L, Nizet V, Meier A. 2025. Anesthetic choice impacts
mortality and bacterial clearance in a murine experimental pneumonia model. BMC
Infect Dis 25:1–10.

53. Wu K, Lin X, Lu Y, Dong R, Jiang H, Svensson SL, Zheng J, Shen N, Camilli A, Chao Y.
2024. RNA interactome of hypervirulent *Klebsiella pneumoniae* reveals a small RNA
inhibitor of capsular mucoviscosity and virulence. Nat Commun 15:6946.

54. Brooks JF 2nd, Gyllborg MC, Kocher AA, Markey LEH, Mandel MJ. 2015. TfoX-based
genetic mapping identifies *Vibrio fischeri* strain-level differences and reveals a common
lineage of laboratory strains. J Bacteriol 197:1065–1074.

55. Le Roux F, Binesse J, Saulnier D, Mazel D. 2007. Construction of a *Vibrio splendidus*
mutant lacking the metalloprotease gene *vsm* by use of a novel counterselectable
suicide vector. Appl Environ Microbiol 73:777–784.

56. Stabb EV, Ruby EG. 2002. RP4-based plasmids for conjugation between *Escherichia
coli* and members of the *Vibrionaceae*. Methods Enzymol 358:413–426.

57. Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640–6645.

58. Peters JM, Koo B-M, Patino R, Heussler GE, Hearne CC, Qu J, Inclan YF, Hawkins JS,
Lu CHS, Silvis MR, Harden MM, Osadnik H, Peters JE, Engel JN, Dutton RJ, Grossman
AD, Gross CA, Rosenberg OS. 2019. Enabling genetic analysis of diverse bacteria with
Mobile-CRISPRi. Nat Microbiol 4:244–250.

59. Wang P, Yu Z, Li B, Cai X, Zeng Z, Chen X, Wang X. 2015. Development of an efficient
conjugation-based genetic manipulation system for *Pseudoalteromonas*. Microb Cell
Fact 14:11.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5