Title: Gut microbe-derived lactic acid optimizes host energy metabolism during starvation

Authors: Jason William Millington[1,2,*], Jamie Alcira Lopez[2,3,*], Amin M. Sajjadian[1,2],

Robert J. Scheffler[4,5], Brian C. DeFelice[6], William Basil Ludington[4,5], Benjamin H. Good[3,6],

Lucy Erin O'Brien[1,6,†], Kerwyn Casey Huang[2,6,7,†]

Affiliations:

[1]Department of Molecular and Cellular Physiology and Institute of Stem Cell Biology

and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA

94305

[2]Department of Bioengineering, Stanford University, Stanford, CA 94305

[3]Department of Applied Physics, Stanford University, Stanford, CA 94305

[4]Department of Embryology, Carnegie Institution for Science, Baltimore, MD 21218

[5]Department of Biology, Johns Hopkins University, Baltimore, MD 21218

[6]Chan Zuckerberg Biohub, San Francisco, CA 94158

[7]Department of Microbiology and Immunology, Stanford University School of

Medicine, Stanford, CA 94305

[*]These authors contributed equally

21   †Correspondence: kchuang@stanford.edu and lucye@stanford.edu

23   *Keywords*: *Drosophila melanogaster*; host-microbe interactions; *Lactiplantibacillus*

24   *plantarum*; gut microbiome; metabolism; oxidative phosphorylation

Abstract

Gut microbes convert dietary compounds into an array of metabolites that can directly provide energy to their host and indirectly impact host metabolism as systemic endocrine signals. Here, we show that gut microbe-derived metabolites can extend *Drosophila melanogaster* survival during starvation, despite minimal alteration of dietary energy intake. Combining survival assays with mathematical modeling and untargeted metabolomics, we identify a single, dominant mediator of starvation resilience: lactic acid produced by the commensal bacterium *Lactiplantibacillus plantarum*. We discover that the basis of starvation resilience is not catabolism of lactic acid using lactate dehydrogenase, but rather increased dietary energy yield through lactic acid-driven promotion of oxidative phosphorylation. Our findings emphasize the role of the microbiome as a source of endocrine cues coordinating host metabolism and underscore the potential of microbiome-derived metabolites as therapeutic molecules for manipulating metabolic health and preventing disease.

Introduction

Gut microbes have emerged as a key regulator of host energy metabolism (Fujisaka et al., 2023; Liu et al., 2022; Turnbaugh and Gordon, 2009; Turnbaugh et al., 2006). In response to host dietary intake, they utilize ingested nutrients for growth, increasing microbial biomass and producing an array of diverse metabolites as byproducts. These processes result in many nutrient-driven host-microbe interactions, including competition for nutrient resources, cross-feeding of microbe-derived metabolites that are taken up by the host to generate energy, and production of signaling molecules that coordinate changes to host metabolism in a manner evocative of endocrine regulation through hormones (Evans et al., 2013; Clarke et al., 2014; Neuman et al., 2015; Reghupaty et al., 2024). Each aspect of these interactions has important consequences for regulating host energy metabolism, and intestinal microbiome dysbiosis has been associated with metabolic disease (Agus et al., 2021).

An extreme example of adaptive regulation of energy metabolism is starvation resilience (Agrawal et a., 2016). Metabolic flexibility to endure unpredictable cycles of feast-famine is a desirable trait for many animal species in the wild, and persistent global prevalence of human malnourishment highlights its relevance to public health. Intriguingly, gut microbes have been successfully deployed as probiotics to treat malnourishment in animal models (Consuegra et al., 2020; Shin et al., 2011; Storelli et al., 2011; Storelli et al., 2018; Schwarzer et al., 2016; Schwarzer et al., 2023). However, the underlying mechanisms through which microbes promote starvation resilience remain largely unknown.

The fruit fly, *Drosophila melanogaster*, is a well-established model for the study of energy metabolism regulation during starvation (Chatterjee and Perrimon, 2021; Min et al., 2007), particularly with regards to host-microbe interactions. In *Drosophila*, *Lactiplantibacillus* and *Acetobacter* species dominate the microbiome (Broderick and Lemaitre, 2012; Pais et al., 2018; Aranda-Díaz et al., 2020; Wong et al., 2011; Dodge et al., 2023) and have been implicated in regulating host energy metabolism and lifespan (Brummel et al., 2004; Fast et al., 2018; Keebaugh et al., 2019; Douglas, 2018; Henriques et al., 2020; Leitão-Gonçalves et al., 2017; Gould et al., 2018; Storelli et al., 2011; Storelli et al., 2018; Shin et al., 2011). Microbiome composition correlates with starvation resilience in *Drosophila* (Keebaugh et al., 2018; Judd et al., 2018; Broderick et al., 2014), potentially occurring via the landscape of microbe-derived metabolites that may act directly as nutrients or indirectly as endocrine signals. However, although a few candidate microbe-derived metabolites have been proposed to impact host energy metabolism (Arnoldini et al., 2024; Fagundes et al., 2024; De Vadder et al., 2014), we lack systematic approaches to identify microbe-derived regulators of energy metabolism. Thus, the

79 immense, yet poorly characterized, chemical diversity produced by the microbiome
80 provides a rich ground for discovering regulation mechanisms.
81
82 Here, we develop an experimental platform and mathematical modeling framework to
83 systematically interrogate the impact of molecules on *Drosophila* energy metabolism
84 during starvation. We apply this method to the metabolites produced by *Drosophila* gut
85 microbiome commensals and find that these metabolites can substantially extend
86 lifespan during starvation despite minimal alteration to dietary caloric content. We
87 identify that *Lactiplantibacillus plantarum*-derived lactic acid is necessary and sufficient
88 for starvation survival extension, and show that lactic acid extends survival via
89 modulation of dietary energy yield in a manner largely independent of the metabolic
90 conversion of lactic acid to pyruvate. We hypothesize that this mechanism acts instead
91 via lactic acid-mediated regulation of oxidative phosphorylation. Consistent with our
92 hypothesis, we show that pharmacological inhibition of oxidative phosphorylation
93 abolishes the lactic acid-mediated starvation resilience. By demonstrating that a single
94 microbial metabolite globally remodels host energy metabolism for starvation
95 resilience, our results highlight potential roles of the microbiome as a source of
96 endocrine signals coordinating metabolic regulation.

Results

Microbe-derived metabolites prolong **Drosophila** *starvation survival*

To investigate how nutrient-based host-microbe interactions contribute to host energy metabolism in *Drosophila*, we developed an assay to screen for altered host starvation survival based on feeding flies microbial spent media (Fig. 1A). In flies near starvation, we found that gut microbiome load and composition were significantly altered relative to fed conditions (Fig. S1A-C). To amplify the host impact of microbe-derived metabolites, we utilized a chemically defined medium (CDM) designed for growth of fly gut microbes (Aumiller et al., 2023). CDM captures two key aspects necessary for our screening approach: (1) CDM permits the growth of all fly gut commensal bacteria and results in high levels of microbial metabolite production, and (2) CDM is a near-starvation baseline condition for fly survival. By itself, 1X CDM supports a marginal fly lifespan of ~1 week (Fig. 1C), versus 2-3 days in complete starvation (STV) conditions and ~2-3 months under *ad libitum* conditions (Piper et al., 2005). To control for potential media batch effects, all assays were performed with the relevant controls including STV and/or CDM conditions.

We used this assay to characterize the contribution to host starvation survival of two abundant *Drosophila* gut commensals isolated from wild flies: *Lactiplantibacillus plantarum* (hereafter *Lp*) and *Acetobacter pasteurianus* (*Ap*) (Obadia et al., 2017; Gould et al., 2018). By filtering stationary phase bacteria cultures, we generated *Lp*- and *Ap*-derived cell-free spent media, from which we produced solid substrates that we fed to adult wild-type *Oregon-R* mated female flies and quantified survival (Fig. 1B,C). Our null expectation was that host-microbe resource competition would result in reduced fly survival on spent media compared to CDM alone due to bacterial consumption of resources that would otherwise have been available to the fly. Indeed, flies fed the spent medium from an *LpAp* coculture (*LpAp*co) had a mean survival time of 5 days, slightly shorter than on fresh CDM (Fig. 1C). However, we found that flies fed 1X spent medium from either *Lp* or *Ap* lived significantly longer than those fed fresh CDM (mean survival of 14 days and 10.5 days, respectively, versus 6 days on CDM; Fig. 1C), indicating that microbe-derived metabolites can positively impact starvation resilience. *Lp*- and *Ap*-spent media prolonged survival without altering fly food intake or taste preference (Fig. S2A-D), active energy expenditure (Fig. S2E-G), or egg production (Fig. S2H). Altogether, these results argue against changes in food intake or active energy expenditure mediating prolonged survival and instead suggest that other aspects of energy metabolism are impacted by microbe-derived metabolites.

Energy balance models suggest that microbe-derived metabolites prolong survival beyond increasing dietary energy content

A multitude of factors could be responsible for the observed lifespan extension in spent
media, including changes in nutritional content, fly behavior, and/or fly metabolism. To
disentangle these potential mechanisms mediating differential survival, we utilized
mathematical models to interpret survival data through a framework based on changes
in energy balance. This framework considers organisms from a thermodynamic
perspective and has been widely applied to starvation and body mass dynamics across
vertebrate and invertebrate species (Hall, 2012; Hall, 2017; Hall and Guo, 2017; Sokolova
et al., 2012; Cesur et al., 2023; Mauritsson and Jonsson, 2023). In these models, survival
during starvation is determined by the initial amount of stored energy and the rates of
energy intake and expenditure. Once energy stores are consumed, organismal death
follows (Fig. 2A). Since the observed distributions of survival times were unimodal, we
focused on the mean and variance of fly survival times across feeding conditions.

Our model considers a population of flies, each with a time-dependent total energy
store $E_i(t)$ such that death occurs deterministically when $E_i = 0$. For simplicity, we
assume that variability in fly survival emerges from random variation across the
population in the initial value of E_i at the time starvation commences (t=0), which we
assume has mean μ_E and variance σ_E^2. Each fly expends energy at rate ϕ_O and intakes
energy at a rate ϕ_I, which is dependent on the energy extracted from the fly diet.
Starvation commences when intake drops below expenditure ($\phi_I < \phi_O$). For simplicity,
we assume here that these rates are constant across the population and over time for a
given diet (see Methods for an analysis of variable rates in energy input/output across
the population).

The mean and standard deviation of survival times is dictated by the mean and
standard deviation in initial values of E_i, scaled by $1/(\phi_O - \phi_I)$ (Methods):

$$\mu_\tau = \frac{\mu_E}{\phi_O - \phi_I}, \ \sigma_\tau = \frac{\sigma_E}{\phi_O - \phi_I}. \quad (1)$$

When the fly is near starvation ($\phi_I \sim \phi_O$), small changes in dietary energy extraction or
energy expenditure can lead to superlinear sensitivity of survival times to energy
metabolism changes (Methods). The similar scaling of μ_τ and σ_τ with respect to
$(\phi_I - \phi_O)$ provides two complementary tests for the model. Note that Eq. 1 predicts a
non-physical infinite lifespan when $\phi_O = \phi_I$; in practice, lifespan will saturate in
nutrient-replete conditions, eventually leading to a reduction in sensitivity to energy
metabolism changes (Methods).

We first sought to test whether it could successfully predict the quantitative dependence
of fly survival dynamics on caloric intake on fresh CDM. We varied caloric intake (ϕ_I)
by feeding flies CDM at concentrations ranging from 0.5X-4X, which resulted in two
regimes of behavior for the mean and standard deviation of survival time (μ_τ and σ_τ,

respectively): between 0.5X-1.5X CDM (low ϕ_I, the energy limitation regime), μ_τ and σ_τ
increased superlinearly with increasing caloric value (Fig. 2B-D). For 2X-4X CDM (high
ϕ_I), μ_τ and σ_τ saturated (Fig. 2C,D), presumably because energy demands are largely
rescued by nutrients present in the CDM (Fig. 2B). These data indicate that the
transition to the saturated regime occurs above 1X CDM concentration, hence 1X CDM
is a useful baseline condition due to the high lifespan sensitivity around this level of
energy intake. Moreover, the agreement between our data and the predictions of our
energy balance model across varying caloric intake for low ϕ_I (Fig. 2E,F) suggest that
fly survival on low levels of CDM is driven by simple energy balance, establishing a
baseline from which to interrogate the mechanisms underlying extended survival in
spent-media conditions.
187
We next asked whether the observed effects of spent media (Fig. 1C) are consistent with
straightforward changes in caloric value. As microbes grow in CDM, nutrients are
consumed and converted to biomass and microbe-derived metabolites. While we do not
know the identity of all microbe-derived metabolites, we can infer the approximate
change in the energy available to the fly. Most of the energy content of CDM (as
estimated by the Gibbs free energy of combustion) comes from glucose and amino
acids, both of which are accessible to the fly (Supplementary Text). Hence, it is unlikely
that microbial metabolism increases survival by increasing energy content; in fact, as
microbes grow, they utilize some of the energy present in the media (Supplementary
Text), lowering the total available energy in the media relative to fresh CDM. Thus, the
observation that spent media from *Lp* and *Ap* monocultures extend fly survival rather
than shorten it (Fig. 1C) suggests that microbe-derived metabolites alter the way
Drosophila extracts, expends, or stores energy. Based on the expectations of our energy
balance model, we concluded that the small observed differences in activity, food intake
volume, and egg production are quantitatively insufficient to explain the lifespan
extension in flies fed *Lp*-spent media (Fig. S2I,J).
204
Next, we sought to determine whether our energy balance model captures the survival
dynamics of flies fed spent media from *Lp*, *Ap*, or *LpAp*[co] at different initial caloric
values. Our analysis of CDM alone suggests that if survival is indeed controlled by
energy balance, *Lp-* and *Ap*-spent media should have minimal effects relative to CDM
alone at higher concentrations in the saturated regime. To test this prediction, we
measured fly survival at 2X caloric density for flies fed CDM, or *Lp-*, *Ap-*, or *LpAp*[co]-
spent media. Flies fed 2X *Lp-* and *Ap*-spent media had a mean survival time of ~26 days
and 24 days, respectively. These survival dynamics are only marginally longer than the
~23 day mean survival of flies fed 2X CDM (Fig. 2G, S3), suggesting that the effect of
microbe-derived metabolites on survival is strongest in the starvation regime. However,
flies fed 2X *LpAp*[co] spent medium had a significantly shorter mean survival time of ~17

days compared with CDM-fed flies (Fig. 2G, S3), equivalent to between 1X and 1.5X
CDM (Fig. 2B). This observation suggests that the co-culture depletes the media of fly-
accessible nutrients; the larger effect relative to 1X $LpAp^{co}$ spent medium (Fig. 1C)
highlights the superlinearity of survival in the starvation regime below 1.5X relative
caloric value (Fig. 2C). Thus, by applying energy balance models, we find that changes
in energy metabolism (and not changes in fly behaviors such as food intake) likely
underly the starvation resilience of flies fed Lp-spent medium.
223
Prolonged starvation survival is mediated by* Lp-*derived lactic acid
We next sought to identify the microbe-derived metabolite(s) that promote fly
starvation survival. To this end, we analyzed the chemical content of CDM and of Lp-,
Ap-, and $LpAp^{co}$-spent media via liquid chromatography with tandem mass
spectrometry (LC-MS, Fig. 3A). We identified ~1500 metabolomic features post-filtering
(mostly unannotated), with distinct metabolomic profiles across media (Fig. 3B,
Methods). One enriched metabolite in Lp-spent medium was lactic acid, as expected
given the metabolic capabilities of Lp, which was undetectable in CDM and Ap-spent
medium (Fig. 3C). Lactic acid was still present in $LpAp^{co}$-spent medium, but at
significantly lower abundance (Fig. 3C). While other metabolites were also enriched in
spent media, we initially focused on lactic acid as it is the primary metabolic product of
Lp fermentation and Lp-spent medium provided the largest lifespan extension (Fig. 1C).
236
Given the pronounced enrichment of lactic acid in Lp-spent medium, we sought to
determine the importance of Lp-derived lactic acid on fly starvation survival. We
utilized $Lp^{\Delta Ldh}$, a strain incapable of lactic acid production due to deletion of lactate
dehydrogenase (Ldh), which catalyzes the terminal fermentation of pyruvate to produce
lactic acid. Flies fed $Lp^{\Delta Ldh}$-spent medium had significantly shorter survival times than
spent medium from the parent Lp^{WCFS1} strain (Fig. 3D) and similar to that of CDM-fed
controls. This finding establishes Lp-derived lactic acid as necessary for increased
starvation survival, and that $Lp^{\Delta Ldh}$ does not substantially deplete CDM of nutrients that
benefit fly survival.
246
To examine whether supplementation of fresh CDM with lactic acid alone could extend
starvation survival, we diluted CDM two-fold and added DL-lactic acid (hereafter,
"lactic acid") to create near-isocaloric, lactic acid-supplemented CDM (0.5X CDM + 62.5
mM lactic acid). We found that lactic acid supplementation was sufficient to
significantly extend starvation survival compared to CDM-fed flies (Fig. 3E). As with
Lp-spent media, the effects of lactic acid were dependent on caloric value: at twice the
calories (1X CDM + 125 mM lactic acid), survival was unaffected compared to 2X CDM
(Fig. S4A,B). Thus, lactic acid is sufficient to extend survival in the starvation regime but
has no effect at higher caloric values as lifespan saturates. We further investigated

whether the deprotonated form of lactate (conjugate base) was sufficient for this effect.
We found directly feeding lactate provided no nutritional value, consistent with our
prediction that lactic acid dominates in spent media (Fig. S5A-D). However, as we do
not know which form predominates intracellularly and thus do not rule out a role for
intracellular lactate mediating this effect.

Lactic acid prolongs starvation survival through an interaction with glucose
metabolism
Given that *Lp*-spent medium extends starvation survival in a manner dependent on
lactic acid but independent of alterations to food intake and active energy expenditure
(Fig. S2D,E), we next considered alternate mechanisms of lactic acid action. One
possibility is a lactic acid-dependent increase in energy extraction from dietary or stored
nutrients (effectively increasing ϕ_I or μ_E). To determine the relative bioavailability of
lactic acid for energy production, we fed flies isocaloric solutions of either lactic acid or
glucose (the main carbon source in CDM). If lactic acid is more bioavailable than
glucose, then the lactic acid diet should promote longer starvation survival. However,
such an effect was not observed experimentally: isocaloric diets of glucose (1X D-
glucose, hereafter glucose) or lactic acid (1X lactic acid) led to similar survival lifespans
(Fig. S4C,D). Furthermore, doubling the concentration of lactic acid (2X lactic acid) had
no effect on starvation survival compared to 1X lactic acid (Fig. S4C,D). On the other
hand, higher concentrations of glucose increased starvation survival (Fig. S4C,D),
consistent with an energy balance mechanism. These findings exclude increased
bioavailability of lactic acid as the mechanism through which it prolongs starvation
survival.

An alternative hypothesis is that lactic acid prolongs starvation survival by interacting
with metabolism of other nutrients such as glucose. Supporting this possibility, we
found that flies fed a 1:1 mixture of glucose and lactic acid at 1X caloric value survived
significantly longer than flies fed either 1X glucose or 1X lactic acid alone (Fig. 3F;
S4C,D). Since all three diets were isocaloric, the survival extension produced by the
combined mixture suggests that lactic acid in the presence of glucose enables increased
dietary energy extraction.

Given the lack of support for a direct energy balance mechanism, we next considered a
signaling role for lactic acid that impacts energy metabolism. This hypothesis is
consistent with known links between lactate and signaling through G-protein coupled
receptor-81 (GPR81) that impacts lipolysis and energy metabolism in mammals (Liu et
al., 2009; Certo et al., 2022). In contrast to nutrients, which are utilized rapidly to impact
energy metabolism, signaling molecules can mediate long-term metabolic effects. We
hypothesized that if lactic acid has a signaling role, then priming via an initial interval

of lactic acid feeding could potentiate increased survival during subsequent feeding
with glucose. To test this hypothesis, we fed flies lactic acid for 5 days before
transferring them to an isocaloric glucose diet (without lactic acid) on which flies were
maintained until death. We also performed the reciprocal treatment of 5 days of glucose
feeding followed by transfer to lactic acid, as well as a control condition of flies
maintained on a mixture of glucose and lactic acid. We found that flies fed lactic acid
prior to glucose feeding survived significantly longer than flies transferred from glucose
to lactic acid (Fig. 3G). Moreover, the survival dynamics post-transfer from lactic acid to
glucose were quantitatively similar to those for the glucose and lactic acid mixture (Fig.
3G). These findings suggest that lactic acid has a priming effect on glucose metabolism
that extends starvation survival even after lactic acid is removed, consistent with a
signaling role for lactic acid.
308
Prolonged starvation survival does not require lactic acid breakdown
309
If lactic acid impacts starvation survival via signaling, we expected that this effect
would be independent of lactic acid metabolism. Bacterial Ldh (Fig. 3D), which is
functionally and structurally divergent from metazoan Ldh, nearly exclusively
produces lactate from pyruvate. In contrast, in metazoans, lactate is metabolized to
pyruvate in a reversible reaction catalyzed by Ldh, regenerating NAD^+ in the process
(Katsyuba et al., 2020). Pyruvate is then shuttled to the mitochondria to generate ATP
and fuel further glycolysis (Rabinowitz and Enerbäck, 2020). To determine the
importance of host lactic acid metabolism, we examined the effect of lactic acid on
starvation survival in flies whose *Ldh* activity was reduced through ubiquitous, RNAi-
mediated knockdown of *Ldh* in adulthood (*tubulin-GAL4, tubulin-GAL80^{ts}*). Flies were
fed isocaloric media containing either glucose, lactic acid, or a mixture of glucose and
lactic acid, and their survival was compared to control flies. If *Ldh* activity is necessary
for the mixture of glucose and lactic acid to prolong survival, then we expected *Ldh-*
RNAi flies to survive for a shorter interval than control flies. However, we found that in
both the knockdown and control genotypes, the mixture of glucose and lactic acid
prolonged survival compared to glucose only (Fig. S6A-C). The lactic acid-only medium
also prolonged survival of *Ldh-RNAi* flies but not control flies, suggesting RNAi-specific
effects that may be related to a decrease in lactic acid-induced stress. Together, these
data suggest that Ldh is not limiting for lactic acid to promote starvation survival in the
presence of glucose, consistent with a signaling role for lactic acid.
330
Lactic acid delays stored energy utilization
331
How can a signaling mechanism facilitate increased survival without higher food intake
or reduced energy expenditure? Given our priming data, we hypothesized that lactic
acid mediates an increase in food energy yield; specifically, that ingestion of glucose
and lactic acid together allows the fly to extract more energy than ingestion of only

lactic acid or only glucose alone. In this case, feeding with glucose and lactic acid
together should allow flies to mobilize their stores of body fat more slowly. To test this
prediction, we investigated the rate of stored body fat breakdown by measuring levels
of whole-body triglycerides, the main energy storage molecule in *Drosophila* and other
higher organisms (Walther et al., 2017; Walther and Farese, 2012). During complete
nutrient starvation, flies exhibited rapid triglyceride mobilization (Fig. 4A), as expected.
However, flies fed a mixture of glucose and lactic acid delayed their fat mobilization
relative to flies fed only glucose or lactic acid (Fig. 4A,B). These data are consistent with
a mechanism in which lactic acid mediates an increase in dietary energy yield.
345
Lactic acid prolongs starvation survival by potentiating OXPHOS
In the context of our energy balance model, increased energy yield corresponds to an
effective increase in ϕ_I. To generate hypotheses regarding the mechanisms responsible
for such an increase, we sought to generate the simplest model of *Drosophila* energy
metabolism consistent with three observations: (1) a mixture of lactic acid and glucose
promotes longer survival than glucose or lactic acid alone, (2) the survival benefit of the
mixture saturates with increasing caloric content, and (3) the survival benefit of the
mixture is largely host Ldh independent. Additionally, the model should exhibit
regimes in which glucose and lactic acid alone provide similar survival benefits (Fig.
3F). We focused on pharmacokinetic-pharmacodynamic (PKPD) models, a framework
with established efficacy for predicting metabolic outputs and the impact of effector
molecules in complex organisms (Bonate, 2011).
358
Our model (Methods) considers two compartments: the gut and the rest of the body
cavity (referred to as the central compartment) (Fig. 4C). The diet contains a mixture of
glucose and lactic acid, which flows through the gut and is absorbed into the central
compartment. Once absorbed, components undergo a highly simplified set of energy
metabolism reactions. Glucose undergoes glycolysis, producing 2 pyruvate and 2 ATP.
Lactic acid is reversibly converted to pyruvate, and each pyruvate can be processed via
oxidative phosphorylation (hereafter OXPHOS), which produces 17 ATP. Each
component is cleared from the central compartment, by excretion from the body or
other component-specific metabolic reactions. To connect this PKPD model to the
energy balance model, we assume that energy extraction ϕ_I from the diet is
proportional to ATP production.
370
371   We found that in the absence of active regulation of glycolysis and OXPHOS, the
372   dietary energy flux ϕ_I is predicted by the PK model to be a linear function of dietary
373   lactic acid and glucose concentrations. Thus, this form of the model does not capture the
374   nonlinear behavior we observed (Fig. 4D). We arrived at similar conclusions using a
375   genome-scale model of *Drosophila* metabolism that considers ~8,000 reactions and

factors such as redox balance (Fig S9A; Methods). These findings are consistent with our empirical results implying a signaling mechanism whereby lactic acid modulates energy metabolism.

To evaluate whether signaling mechanisms could account for our key experimental results, we focused on regulation of OXPHOS, which is where the most energy can be harvested and impacts both lactic acid and glucose energy yield. OXPHOS regulation is also in line with a previous *in vitro* study showing that lactic acid can promote oxidative phosphorylation in HepG2 cells and allow a higher yield of energy from glucose (Cai et al., 2023). Indeed, we found that a regulatory scheme in which the ratio of gut lactic acid to glucose regulates OXPHOS can recapitulate all key behaviors if (1) there is a basal level of OXPHOS activity in lactic acid or glucose diets, and (2) there is an optimal ratio of lactic acid to glucose (Fig. 4E). Furthermore, simulations across increasing levels of Ldh knockdown efficiency produced similar results (Fig. 4F), indicating that an OXPHOS regulatory scheme can be largely Ldh independent. Our proposed regulatory scheme is consistent with several possible mechanisms, such as gene expression altering OXPHOS enzyme levels or post-translational modification of OXPHOS enzymes. Nonmonotonic changes in the per-pyruvate ATP yield of OXPHOS would produce similar effects. Other nonlinear OXPHOS regulatory schemes, such as an optimal concentration of lactic acid rather than an optimal ratio of lactic acid to glucose concentrations, are also potentially consistent with our results.

This model predicts that survival of flies fed a mixture of lactic acid and glucose would be adversely impacted by inhibition of OXPHOS. To test this prediction, we inhibited OXPHOS pharmacologically using metformin, which inhibits OXPHOS by blocking complex I of the electron transport chain (Owen et al., 2000), and assessed the impact on starvation survival. In vehicle-treated control flies, we observed that flies fed a mixture of glucose and lactic acid survived significantly longer than glucose or lactic acid fed flies alone (Fig. 4G,H; S7). In contrast, flies fed the mixture in conjunction with metformin treatment did not survive longer than glucose-fed flies alone and their survival was significantly shorter than vehicle-treated controls (Fig. 4G,H; S7). Our observation that metformin abolishes extended survival in flies fed the mixture of glucose and lactic acid supports the model that lactic acid impacts starvation survival via OXPHOS regulation.

Discussion

Here, we showed that under starvation conditions, lactic acid generated by the gut commensal *Lp* extends host survival by enhancing dietary energy yield via OXPHOS. By developing and applying a method for screening and interrogating the impact of candidate effectors on energy metabolism in *Drosophila*, we uncovered a class of host-microbiome interactions in which a microbial metabolite alters host energy metabolism by modulating the energy yield from nutrient sources. We propose that the gut microbiome, a rich source of metabolic effectors, effectively serves as a cross-kingdom endocrine regulator of host energy metabolism.

Our experiments involved diets pre-conditioned by microbes, a context in which the microbes initially have exclusive access to dietary components. While this pre-conditioning mimics microbially fermented foods, it is unclear to what extent gut microbes chemically modify dietary components *in situ*. As a case study, we examined the lactic acid flux produced by *Lp* in non-starving *Drosophila*. By combining data on fly gut bacterial population sizes, gut transit times, and food intake measurements, we calculated that the flux of lactic acid produced by the fly gut microbiome is similar in magnitude to the flux of carbon compounds from food intake (Supplementary Text). This suggests that *in situ* bacterial metabolism can indeed generate physiologically relevant fluxes of effector metabolites. Note that owing to the smaller bacterial populations in starved flies (Fig. S1), these fluxes may be less relevant in severe starvation conditions. Our calculation is also consistent with results from other multicellular organisms. In humans, it is well-established that microbial metabolism of dietary compounds can produce physiologically relevant fluxes of fermentation products, supplying up to 10% of incoming energy (Arnoldini et al., 2024). Indeed, work on microbiome drug metabolism has shown that microbial modification of even non-food ingested compounds can substantially impact host physiology in mice (Zimmermann et al., 2024; Javdan et al., 2020) and humans (Lindenbaum et al., 1981). Thus, *in situ* chemical modification of ingested compounds may influence organismal physiology across the tree of life.

Our results also reveal a counter-intuitive relationship between dietary energy yield and organismal fitness. The role of diet in human energy balance can be largely captured by assuming that the energy yield is near the thermodynamic limit (i.e., the free energy of combustion) (Merrill and Watt 1955, Chow and Hall, 2008) and humans rarely excrete non-CO_2 carbonaceous waste (Rabinowitz and Enerbäck, 2020). However, we found that even under starvation, flies did not always fully utilize the energy contained in their diet, with flies fed 1X CDM leaving at least ~45% of the dietary energy unutilized (calculated from the increase in energy intake required for *Lp-*

mediated survival benefit; Fig. 1C). This finding contrasts with naïve expectations for
behavior during starvation, when an organism might be expected to be especially
energetically efficient to prolong survival. Why then does *Drosophila* underutilize
dietary energy during starvation? One possibility is a trade-off between energy yield
and other fitness-relevant phenotypes. For example, in single-celled organisms such as
bacteria, a trade-off between energy yield and overall energy influx rate has been well
characterized (De Groot et al., 2020), whereby it can be optimal to only partially oxidize
substrates to increase total substrate flux and overall energy flux. Similar trade-offs have
also been observed in cancer (Warburg, 1956) and yeast cells (Postma et al., 1989).
However, it is unlikely that this trade-off between substrate flux and yield explains the
behavior of *Drosophila* since food intake volume (Fig. S2D), and thus overall substrate
flux, does not change between diets. Other forms of trade-offs, such as with fertility,
could underlie this behavior. For example, prolonged periods of starvation can lead to
infertility in *Drosophila* (Churchill et al., 2019; Terashima and Bownes, 2004), decoupling
lifespan extension during starvation from reproductive fitness. In addition to nutrient
yield trade-offs, an alternative explanation is that the unmodified CDM diet is far from
typical *Drosophila* diets and that this under-yielding does not often occur in nature. The
diet of wild *Drosophila* consists largely of rotten fruit rich in *Lactiplantibacilli*, *Acetobacter*,
and yeasts amongst a host of other microbial species producing microbial metabolites,
including lactic acid. Thus, a fly in the wild may typically encounter mixtures of glucose
and lactic acid in its food source, rather than the pure glucose found in CDM. Further
characterization of phenotypes in relation to energy yield across hosts could shed light
on these fundamental evolutionary questions.
473
Finally, while our study focuses on one interaction between lactic acid and glucose
metabolism, the experimental and mathematical tools we developed represent a broad
platform for detecting and dissecting the role of energy metabolism effectors more
generally. The metagenome of intestinal microbial communities encodes a vast potential
for microbe-derived metabolite production (Qin et al., 2010), many of which can be
detected in host circulation (Lai et al., 2021; Quinn et al., 2020) and have potential
(largely uncharacterized) roles as signaling molecules, akin to the complex hormonal
milieu produced by different cell types in an endocrine organ. Our findings highlight
the scope for discovery of novel metabolic effectors among microbe-derived
metabolites, potentially paralleling the discovery of hormone-based treatments for
metabolic disease from mechanistic studies of circulating regulators of energy
metabolism such as GLP-1 and analogs such as semaglutide (Drucker et al., 2017;
Baggio and Drucker, 2007; Drucker and Holst, 2023; Wewer Albrechtsen et al., 2023).
One could similarly use energy balance models to guide experiments pinpointing the
aspects of energy metabolism altered by any diet and leverage untargeted
metabolomics to identify putative effector molecules. Beyond detecting microbiome-

derived effectors, our methods can be broadly applied to any putative effectors that can
be fed to flies as a solid agar. Indeed, this approach is a means to screen drug candidates
in vivo, many of which interact with the microbiome (Javdan et al., 2020; Zimmermann
et al., 2019), exemplified by our demonstration that metformin, a drug known to alter
human energy metabolism, impacts lactic acid signaling. Furthermore, given the
increasing prevalence of metabolic disease in humans (Liu et al., 2025), it will be
informative to leverage this platform to identify potential protective mechanisms of
both microbe-derived and xenobiotic metabolites on energy metabolism in *Drosophila*
models of metabolic disease.

Methods

Drosophila husbandry

Mated adult female flies 5-7 days post eclosion were used for all experiments. All flies were synchronized on diets during the L1 stage and raised at a density of 50 larvae per 10 mL food at 25 °C on standard cornmeal, molasses, yeast medium. Adult flies were maintained at a density of 10 flies per vial. All flies used for experiments were conventionally reared (not germ free).

Drosophila strains

We used the following fly strains from the Bloomington *Drosophila* Stock Center: *Oregon-R* (#5), *w^{1118}* (#3605), *Ldh^{16}* (#94698), *tubulin-GAL4* (#5138), *tubulin-GAL80^{ts}* (#7017), and *UAS-Ldh-RNAi* (#33640). All relevant genotypes are listed in the figure legends.

Bacterial culturing

To prepare spent media, *Lp* and *Ap* were grown from frozen stocks into colonies on De Man-Rogosa-Sharpe (MRS) agar plates and individual colonies were grown in liquid MRS medium for 48 h. These cultures were diluted 1:200 in a 50:50 mixture of MRS and CDM and grown for 24 h. For *LpAp* co-cultures, equal volumes of *Lp* and *Ap* MRS cultures were combined and diluted 1:200 in a 50:50 mixture of MRS and CDM. These cultures were then diluted 1:200 in CDM, grown for 24 h, diluted again 1:200, and grown for 48 h. Fresh CDM medium (as a control) and 48 h CDM cultures were pelleted via centrifugation at $4500g$ for 5 min. For 2X media, concentrated CDM was used to culture bacteria. The supernatants were filter sterilized using 0.22 μm filters (Thermo Scientific, 725-2520) to generate CDM control supernatant or cell-free spent media. To create a solid medium for fly ingestion, we mixed CDM supernatant or spent media 1:1 with 2% agar to create a 1% CDM or spent medium agar that was fed to flies. CDM was made according to published protocol (Aumiller et al., 2023). The following bacteria strains were used in this study: *Lp* (Obadia et al., 2017), *Ap* (Gould et al., 2018), *Lp^{WCFS1}* (Hayward and Davis, 1956), *Lp^{ΔLdh}* (Ferain et al., 1996). All culturing was performed at 30 °C with shaking.

Colony forming units

To enumerate bacterial density during starvation, flies were initially inoculated with *Lp* and *Ap*. Specifically, flies were allowed to mate for 2 days post eclosion before transfer to media containing a cocktail of antibiotics (neomycin (Fisher Scientific, BP266295), 0.1 mg mL^{-1}; ampicillin (Sigma-Aldrich, A9518), 0.1 mg mL^{-1}; vancomycin (MP Biomedicals, 02195540.2), 0.1 mg mL^{-1}; metronidazole (MP Biomedicals, 0215571025), 0.5 mg mL^{-1}) for 3 days. Flies were then transferred to sterile media inoculated with 100 μL of *LpAp* co-culture for 2 days. Flies were then transferred to 1% agar media before collection at

539   various time points post starvation. Flies were surface sterilized using ethanol before
540   homogenization using 0.5 mm silica beads (BioSpec Products, 11079105z) with shaking
541   (Omni International, BeadRuptor 12). Colonies were grown via serial dilution plating of
542   fly homogenate (Dodge and Ludington, 2023) and quantified using the ImageJ plugin
543   Count-On-It (Dodge and Ludington, 2023).
544
Survival assay
Flies were reared as described above and transferred in groups of 10 per biological
replicate to experimental media 5-7 days post eclosion. Survival was quantified at 12 h
intervals by enumerating dead flies. All experiments were performed at 25 °C, except
for temperature-sensitive experiments, which were performed at 29 °C, and vials
remained in incubators until all flies were dead. Mean survival was calculated at each
time point. The following chemicals were used to feed flies for survival assays: DL-lactic
acid (Sigma-Aldrich, L6661), D-glucose (Fisher Scientific, D16-3), agar (Sigma-Aldrich,
A1296), and metformin (Sigma-Aldrich, PHR1084). Each biological replicate represents
one vial of 10 flies.
555
Taste preference
Taste preference was determined using a two-choice assay developed in a previous
study (Stanley et al., 2021). In brief, groups of 10 mated female flies 5-7 days post
eclosion were starved for 12 h. Flies were then transferred to vials containing six 10 µL
drops of tastant with alternating red (0.5 mg mL^{-1} Amaranth Red, Sigma-Aldrich,
A1016) and blue color (0.125 mg mL^{-1} Erioglaucine Blue, Sigma-Aldrich, 861146). Each
drop contained a 1X concentration of CDM supernatant or spent media tastant or water.
To control for color preference, we alternated each tastant in color across experimental
vials to ensure that half of all replicates per group had each tastant in red and half in
blue. Flies were fed for 2 h at 25 °C, before freezing at -20 °C. Abdomen color was then
manually scored as red, blue, purple, or none using a dissection scope. Preference index
was calculated as ((# flies with tastant 1 color)-(# flies with tastant 2 color))/(# flies of
any color). Vials with less than 30% of flies scored as any color were not included. For
longitudinal measurements of taste preference during starvation, flies were collected at
indicated times post starvation and assayed as above.
571
Food intake
Food intake was quantified using the <u>ca</u>pillary <u>fee</u>der assay (CAFÉ) according to
previous studies (Ja et al., 2007; Stanley et al., 2021). In brief, vials containing 10 mated
female flies 5-7 days post eclosion were starved for 4 h. Flies were then transferred to
vials whose plugs had two accessible capillary tubes (Drummond Scientific, 2-000-020)
containing liquid media as indicated. To more easily visualize liquid volume in
capillary tubes, each liquid media contained 0.125 mg mL^{-1} Erioglaucine blue dye. To

control for liquid evaporation, an empty vial containing capillary tubes but no flies was
included. Evaporation volume was subtracted from each value before analysis. All
assays were performed at 25 °C.
582
Activity monitor
Activity and sleep were quantified as previously described (Vaughen et al., 2022).
Briefly, 5-7 day post-eclosion mated female flies were loaded singly into capillary tubes
compatible with the *Drosophila* activity monitor (DAM, Trikinetics) (Pfeiffenberger et al.,
2010). Each capillary tube (Trikinetics, PPT5x65) contained the indicated media. Activity
bouts were collected for 2 days with sleep defined as 5 min of inactivity. Flies with
protracted inactivity were determined to have perished and were omitted from
analyses. Analysis was performed using the web-based application ShinyDAM
(Cichewicz and Hirsh, 2018).
592
593   *Fecundity*
594   The number of eggs laid was enumerated as previously described (Millington et al.,
595   2021). In brief, 5- to 7-day old mated female flies were transferred onto indicated media
596   and the number of eggs laid over 24 h was quantified per group of 10 flies.
597
Triglyceride quantification
Triglycerides were quantified as previously described (Wat et al., 2020). In short, 5-7 day
post-eclosion mated female flies were collected at indicated time points in groups of five
and homogenized in 200 µL of 0.1% Tween in PBS using 1 mm glass beads (Sigma-
Aldrich, Z250473) with shaking (Omni International, BeadRuptor 12). Triglyceride
concentration was determined using a colorimetric assay as previously described
(Tennessen et al., 2014). Triglyceride concentration was normalized to body weight
(Mettler Toledo, MS204S) to calculate percent body fat.
606
Mass spectrometry
Liquid chromatography with tandem mass spectrometry was performed as previously
described (Ho et al., 2024). Briefly, spent media were generated and stored at -80 °C.
Samples were split into two 20 µL aliquots and dispensed into 96-well polypropylene
plates (Greiner Bio-One, 655161) on ice. To avoid biases associated with repeated freeze-
thaw cycles, samples were kept on ice and analysis was performed upon their first
thawing. As a quality control step, an additional pooled sample was generated by
mixing 5 µL of each sample; this mixed sample was injected at various times throughout
LC/MS analysis to check instrument performance. All samples were analyzed in both
ionization modes. Sample order was randomly determined for injection to prevent bias
in data collection. Blanks consisting of water in place of fresh CDM supernatant or spent
media were prepared by performing a parallel extraction and ran as additional quality

controls. Chromatographic conditions were modified to shorten the analysis time.
Specifically, a shorter ACQUITY BEH Amide 130 Å, 1.7 µm, 2.1 mm x 50 mm (Waters,
Milford MA) column was used. Chromatographic gradient was shortened as such: 0-0.5
min 100% B, shifting to 70% B at 1.95 min, 40% B at 2.55 min, returning to 100% B at 3.15
min and held until 3.8 min. All other analytical conditions were as described in (Ho et
al., 2024).
625

Metabolomics analysis
Untargeted feature extraction was performed with MS-Dial v. 4.60 (Tsugawa et al.,2015
& Tsugawa et al. 2020). Mass accuracy tolerances were set to 0.01 Da and 0.015 Da for
MS1 and MS2, respectively. Peak height threshold was set to 50k. Lactic acid was
annotated based on matching accurate mass, fragmentation spectra, and retention time
with an authenticated standard. All reported features were present in 100% of replicates
from at least one condition and had a maximum peak height at least five fold greater
than the average of the water blanks. Fold change of a given feature was computed as
the ratio of biological replicate means. To prevent division by zero when computing
fold changes, we added a pseudocount equal to the lowest nonzero feature area across
all samples. For principal component analyses (PCA), we first preprocessed the data by
removing features that were constant across all samples of interest and then
standardizing all features by subtracting the feature mean and dividing by its standard
deviation. PCA was performed using the *PCA* function within scikit-learn using the full
exact singular value decomposition solver. Details of statistical testing of normalized
peak areas are given in the figure legends.
642

643   *Survival curve data analyses*
644   The survival curve $S(\tau)$ is related to the distribution of death times $P(\tau)$ by
645

646
$$P(\tau) = -\frac{dS(\tau)}{d\tau}. \tag{S1}$$

648   Using this relation, the mean and variance of survival times can be estimated from $S(\tau)$
649   as

650
$$\mu_\tau = \int_0^\infty S(\tau)d\tau, \tag{S2}$$

651
$$\sigma_\tau^2 = 2\int_0^\infty \tau S(\tau)d\tau - \left(\int_0^\infty S(\tau)d\tau\right)^2. \tag{S3}$$

653   We use Simpson's method to numerically integrate the empirical survival curves in
654   each vial. In vials in which the flies died very rapidly, we found that the estimator for
655   variance can occasionally result in a slightly negative variance. In such cases, we set the

656 variance to zero. Descriptions of the statistical testing methods used for survival curve
657 moments are given in the figure legends.
658
659 *Generalized fly energy balance model*
660 Here, we present a general form of the simplified energy balance model shown in the
661 main text. In this section, we introduce the model and show how it can be reduced to
662 the expressions in the main text. In the following sections, we explore more complex
663 variations of the model.
664
Drawing upon energy balance models developed for humans (Speakman and
Westerterp, 2012; Chow and Hall, 2008), we model a population of flies, indexed by i,
that each have total stored energy $E_i(t)$ encompassing fat and other energy storage
molecules. For simplicity, we treat all energy storage forms as equivalent. We denote
the initial stored energy of a fly as $E_i(t = 0) = E_{i,0}$. Initially, the population has
$\text{mean}(E_{i,0}) = \mu_E$ and $\text{var}(E_{i,0}) = \sigma_E^2$. Each fly expends energy at rate $\phi_{O,i}(t)$ and intakes
energy at rate $\phi_{I,i}(t)$. The dynamics governing the energy balance of an individual fly
are therefore

$$\frac{dE_i}{dt} = \phi_{I,i}(t) - \phi_{O,i}(t). \tag{S4}$$

674
675 The solution to Eq. S4 is

$$E_i(t) = E_{i,0} + \int_0^t \phi_{I,i}(\zeta)\, d\zeta - \int_0^t \phi_{O,i}(\zeta)\, d\zeta. \tag{S5}$$

677 We assume that a fly dies of starvation once its energy storage reaches zero at a
678 starvation time τ_i, $E_i(\tau_i) = 0$.
679
680 To reach the minimal null model shown in the main text, we make two simplifications:
681 assuming (1) constant energy intake and expenditure, and (2) the same energy intake
682 and expenditure for all flies in the population, such that $\phi_{I,i}(t) = \phi_I$ and $\phi_{O,i}(t) = \phi_O$.
683 With these assumptions, the solution is $E_i(t) = E_{i,0} + (\phi_I - \phi_O)t$ and the starvation time
684 of an individual fly τ_i is

$$\tau_i = \frac{E_{i,0}}{\phi_O - \phi_I}. \tag{S6}$$

686 The model predicts that the survival time will be linear as a function of stored energy, a
687 prediction that approximately holds in more complex human energy balance models
688 (Speakman and Westerterp, 2012). Both the mean and variance are based on linear
689 operators (i.e., $\text{mean}(aX) = a\,\text{mean}(X)$ and $\text{var}(aX) = a^2\,\text{var}(X)$), and thus estimates of
690 the moments of τ_i given μ_E and σ_E^2 are

$$\mu_\tau = \frac{\mu_E}{\phi_O - \phi_I}, \tag{S7}$$

$$692 \qquad \sigma_\tau^2 = \frac{\sigma_E^2}{(\phi_O - \phi_I)^2} \Rightarrow \sigma_\tau = \frac{\sigma_E}{\phi_O - \phi_I}. \qquad \text{(S8)}$$

694   From Eq. S7, we can derive a variable transformation in which there is a linear
695   relationship between inverse mean starvation time and the energy intake ϕ_I (and
696   similarly for starvation standard deviation in Eq. S8):

$$697 \qquad \frac{1}{\mu_\tau} = \frac{\phi_O}{\mu_E} - \frac{\phi_I}{\mu_E}. \qquad \text{(S9)}$$

These transformed moments are the most suitable variables for plotting comparisons of
the energy balance model to experimental data. We plot a transformed version of Fig.
2C,E in Fig. S8, which shows the agreement between model and experiment.

To relate the above expressions to food energy density u, we assume a constant,
nutrient-independent inflow of food Q, such that $\phi_I = Qu$. In this simplified model, we
expect a linear relationship between inverse mean survival time, inverse survival time
standard deviation, and food energy density.

This mathematical model indicates that the sensitivity of the starvation assay to
changing dietary energy can be superlinear under certain conditions. This
superlinearity is apparent in the derivative of the mean survival time with respect to
input energy:

$$711 \qquad \frac{\partial \mu_\tau}{\partial \phi_I} = \frac{\mu_E}{(\phi_O - \phi_I)^2}. \qquad \text{(S10)}$$

This derivative diverges as the input energy flux approaches ϕ_O. Note that the
sensitivity will decrease sharply if lifespan plateaus at a finite value.

Energy balance models with finite lifespan
The model in the previous section predicts that fly lifespan will be infinite once the fly's
energy needs are met. In reality, lifespan will plateau at some finite value due to other
factors, limiting the linear relationship between lifespan and food energy density to
what we term the "starvation regime". We introduce two models that produce finite
lifespan as examples, but our analyses in this manuscript do not depend on the
underlying mechanism for the lifespan plateau.

First, consider a case where the fly's ability to extract energy from food is limiting (e.g.,
due to absorption limitations in the gastrointestinal tract): that is, $\phi_I \propto f(u)$, where $f(\cdot)$
is a saturating function. In this case, both the mean survival time and survival time
standard deviation will plateau at some maximum value, mimicking what is seen in
experiments.

Another possibility is that lifespan becomes limited by a second factor that is depleted
during starvation (e.g., vitamins or nitrogen). The storage level of this compound in fly
i, which we call X_i, should obey similar dynamics to the energy storage:

$$\frac{dX_i}{dt} = \pi_{I,i}(t) - \pi_{O,i}(t), \tag{S11}$$

where π_I and π_O represent expenditure and intake, respectively, of the second limiting
factor. In this case, lifespan will be dictated by which factor is depleted first. As energy
intake is increased, lifespan will smoothly reach a plateau as the other factor becomes
limiting. We expect a different variance relationship from the limited energy extraction
model as the variance will shift from being controlled by energy storage variance to the
storage variance of the second limiting factor.

Note that outside of the starvation regime we consider in this work, the relationship
between energy intake and lifespan is not straightforwardly controlled by simple
energy balance, with reduced caloric intake associated with longer lifespan in well-fed
flies (Min et al., 2007).

Energy balance model with population-varying energy intake and expenditure
We have thus far assumed that there is only substantial variation in the initial energy
storage of the flies. However, variation in energy expenditure and intake is also
possible. To determine how such variation would influence survival time, we start from
Eq. S5 and consider variation in the net rate of energy expenditure $\Delta\phi_i = \phi_{O,i} - \phi_{I,i}$,
with $\text{mean}(\Delta\phi_i) = \mu_{\Delta\phi}$ and $\text{var}(\Delta\phi_i) = \sigma^2_{\Delta\phi}$. Thus, we aim to determine the value of
$\text{mean}(\tau_i) = \text{mean}\left(\frac{E_{i,0}}{\Delta\phi_i}\right)$. To approximate this quantity, we perform a second-order
Taylor expansion for a bivariate function $f(X,Y)$ about the point $(X,Y) = (a,b)$:

$$f(X,Y) \approx f(a,b) + f_X(a,b)(X-a) + f_Y(a,b)(X-b) + \frac{f_{XX}(a,b)(X-a)^2}{2}$$

$$+ \frac{f_{YY}(a,b)(Y-b)^2}{2} + f_{XY}(a,b)(X-a)(Y-b). \tag{S12}$$

757 where f_X and f_{XX} denote the first and second derivative, respectively, of f with respect
758 to X. In our case, $f(E_{i,0}, \Delta\phi_i) = E_{i,0}/\Delta\phi_i$ and we expand about μ_E and $\mu_{\Delta\phi}$. Substituting
759 in these values and applying the expectation operator to both sides of Eq. S12 yields

$$\text{mean}(\tau_i) = \text{mean}\left(\frac{E_{i,0}}{\Delta\phi_i}\right) \approx \frac{\mu_E}{\mu_{\Delta\phi}} - \frac{\text{cov}(E_{i,0}, \Delta\phi_i)}{(\mu_{\Delta\phi})^2} + \frac{\mu_E \text{var}(\Delta\phi_i)}{(\mu_{\Delta\phi})^3} \tag{S13}$$

$$\text{mean}(\tau_i) \approx \frac{\mu_E}{\mu_{\Delta\phi}}\left(1 + \text{CV}^2_{\Delta\phi}\right) - \frac{\text{cov}(E_{i,0}, \Delta\phi_i)}{(\mu_{\Delta\phi})^2} \tag{S14}$$

Thus, variation and covariation in $\Delta\phi_i$ alter the mean only via higher-order terms. Since

τ_i is linearly dependent on $E_{i,0}$, variation in $E_{i,0}$ does not alter the mean of τ_i.

Fit of energy balance model to survival data

To fit energy balance models to our experimental data, we used a two-stage approach.

First, we fit the non-saturating version of the model to experiments with 0X-2X CDM, as

our data indicated that these conditions are outside the lifespan saturation regime. The

estimated moments from each vial were treated as independent replicates. We first fit

Eq. S7 to the mean survival times. Since the three parameters only appear in ratios, e.g.,

μ_E/ϕ_O and μ_E/ϕ_I, we arbitrarily set $\mu_E = 1$ and estimated ϕ_O and ϕ_I. We performed this

fitting using the MATLAB function *fit*, which minimizes the sum of squared errors.

Confidence intervals were estimated using the MATLAB function *confint*. We then fit

the variance data using Eq. S9, using estimates of ϕ_O and ϕ_I from the mean survival

time fit as fixed parameters and σ_E as a fitting parameter.

We found that the simple energy balance model fit the mean survival time data well for

flies fed CDM or a pure glucose diet, with adjusted r^2 values of 0.99 and 0.78,

respectively. The standard deviation of survival time was also well explained, with

adjusted r^2 values 0.85 and 0.72. As expected, the naïve energy balance models poorly

explained the *Lp*- and *LpAp*^{co}-spent media data, with both fits exhibiting negative

adjusted r^2 values.

To estimate the parameter changes required to achieve a given change in survival time

(e.g., the increase in energy intake ϕ_I required to produce the increase in survival

between CDM and *Lp*-spent medium), we used Eq. S7 in conjunction with parameters

from the above fitting procedure. As an example, we analyze the comparison between

1X CDM versus 1X *Lp*-spent medium. We begin with the parameters estimated using

the CDM data: ϕ_O^{CDM}, ϕ_I^{CDM}, and μ_E^{CDM}. Given that the ratio of mean survival times

between these two conditions is $\mu_\tau^{Lp}/\mu_\tau^{\text{CDM}}$, we wish to determine the changes in the

energy metabolism parameters (i.e., the values of ϕ_O^{Lp}, ϕ_I^{Lp}, and μ_E^{Lp}) required to produce

this change. We analyzed each parameter independently while holding the other two

constant. The mean stored energy is

794
$$\mu_E^{Lp} = \mu_E^{\text{CDM}}\left(\frac{\mu_\tau^{Lp}}{\mu_\tau^{\text{CDM}}}\right), \tag{S15}$$

795   the outgoing energy flux is

796
$$\phi_O^{Lp} = \frac{\mu_\tau^{\text{CDM}}}{\mu_\tau^{Lp}}\left(\phi_O^{\text{CDM}} - \phi_I^{\text{CDM}}\right) + \phi_I^{\text{CDM}}, \tag{S16}$$

797   and the incoming energy flux is

$$\phi_I^{Lp} = \frac{\mu_\tau^{\mathrm{CDM}}}{\mu_\tau^{Lp}} \left(\phi_I^{\mathrm{CDM}} - \phi_O^{\mathrm{CDM}} \right) + \phi_O^{\mathrm{CDM}}. \tag{S17}$$

Linear pharmacokinetic (PK) model of glucose and lactate metabolism in Drosophila
The goal of this analysis is to identify a minimal set of assumptions needed to
recapitulate three key experimental observations: (1) a mixture of lactic acid and glucose
can promote longer survival than glucose or lactic acid alone, (2) the survival benefit of
the mixture saturates with increasing caloric content, and (3) these behaviors are not
affected by strong but partial *Ldh* knockdown. Additionally, the model should exhibit
regimes in which lactic acid alone promotes similar mean survival as glucose alone.
Note that given the vast space of possible models, our goal is to identify whether a
model is consistent with our data, not whether it is the only model that can explain the
data.

We begin with a two-compartment pharmacokinetic model that considers the gut as one
compartment and the rest of the fly body as another compartment. The glycolysis-
OXPHOS reaction scheme is simplified and inspired by a previous study (Jacquet and
Stéphanou, 2023). The two primary reactions we consider are:

$$\text{Glucose} + 2\,\text{NAD}^+ \rightarrow 2\,\text{Pyruvate} + 2\,\text{ATP} + 2\,\text{NADH}, \tag{S18}$$
$$\text{Pyruvate} + \text{NADH} + 3\text{O}_2 \rightarrow 17\,\text{ATP} + \text{NAD}^+, \tag{S19}$$

corresponding to the net reactions of glycolysis and OXPHOS, respectively. We also
consider the activity of Ldh:

$$\text{Lactate} + \text{NAD}^+ \rightarrow \text{Pyruvate} + \text{NADH} + \text{H}^+. \tag{S20}$$

We assume all reactions are linear for analytic tractability. We do not consider the
energetic contribution of fat, focusing on food energy extraction, which appears to
supply the majority of the fly's energy in 1X nutrient conditions. Here, we do not
explicitly model redox (e.g., NAD^+/NADH ratio) and assume that the fly is able to
maintain this ratio to enable metabolism; we later will consider explicit redox dynamics.
We also assume that concentration of ADP, phosphate, oxygen, protons, carbon dioxide,
and water do not control the rate of these reactions and thus we do not explicitly model
their concentrations.

We model the gut as a well-mixed compartment with constant and equal inflow and
outflow rate δ. $1/\delta$ reflects the passage time of the fly gut. The concentration of glucose
and lactic acid in the diet is $[\text{G}]_{\mathrm{diet}}$ and $[\text{L}]_{\mathrm{diet}}$, respectively. The concentration in the gut
lumen is $[\text{G}]_{\mathrm{gut}}$ and $[\text{L}]_{\mathrm{gut}}$, respectively. Absorption of gut glucose and lactic acid occurs
linearly at rates $k_{\mathrm{a,G}}$ and $k_{\mathrm{a,L}}$, respectively. In the body, lactic acid can be converted to
pyruvate by lactate dehydrogenase at rate $k_{\mathrm{Ldh-f}}$, and pyruvate is converted to lactate at

rate $k_{\text{Ldh-r}}$. Pyruvate can be used for OXPHOS at rate k_{OXPHOS}, generating $Y_{\text{OXPHOS}} =$
17 ATP per pyruvate. Two pyruvates and $Y_{\text{glycolysis}} = 2$ ATP per glucose are produced by
glycolysis at rate $k_{\text{glycolysis}}$. Glucose, lactate, and pyruvate are cleared from the body or
used for other metabolic reactions at clearance rates CL_σ. We assume the incoming
energy flux ϕ_I is proportional to the ATP generated by glycolysis and OXPHOS. The
dynamics are thus

$$\frac{d[\text{G}]_{\text{gut}}}{dt} = \delta[\text{G}]_{\text{diet}} - k_{\text{a,G}}[\text{G}]_{\text{gut}} - \delta[\text{G}]_{\text{gut}}, \tag{S21}$$

$$\frac{d[\text{L}]_{\text{gut}}}{dt} = \delta[\text{L}]_{\text{diet}} - k_{\text{a,L}}[\text{L}]_{\text{gut}} - \delta[\text{L}]_{\text{gut}}, \tag{S22}$$

$$\frac{d[\text{G}]}{dt} = k_{\text{a,G}}[\text{G}]_{\text{gut}} - k_{\text{glycolysis}}[\text{G}] - \text{CL}_\text{G}[\text{G}], \tag{S23}$$

$$\frac{d[\text{L}]}{dt} = k_{\text{a,L}}[\text{L}]_{\text{gut}} - k_{\text{Ldh-f}}[\text{L}] + k_{\text{Ldh-r}}[\text{P}] - \text{CL}_\text{L}[\text{L}], \tag{S24}$$

$$\frac{d[\text{P}]}{dt} = 2k_{\text{glycolysis}}[\text{G}] - k_{\text{OXPHOS}}[\text{P}] + k_{\text{Ldh-f}}[\text{L}] - k_{\text{Ldh-r}}[\text{P}] - \text{CL}_\text{P}[\text{P}], \tag{S25}$$

$$\phi_I \propto \frac{d[\text{ATP}]}{dt} = Y_{\text{glycolysis}}k_{\text{glycolysis}}[\text{G}] + Y_{\text{OXPHOS}}k_{\text{OXPHOS}}[\text{P}]. \tag{S26}$$

849 *Steady-state behavior of the linear lactic acid and glucose PK model*
850 We sought to determine whether the above linear model could reproduce the starvation
851 dynamics seen in our assays. Assuming that the fly's internal reaction network reaches
852 steady state on a timescale faster than starvation, the energy influx from diet will be

$$\phi_I \propto Y_{\text{glycolysis}}k_{\text{glycolysis}}[\text{G}]^* + Y_{\text{OXPHOS}}k_{\text{OXPHOS}}[\text{P}]^*, \tag{S27}$$

854 where $[\text{G}]^*$ and $[\text{P}]^*$ denote the steady-state concentrations of glucose and pyruvate,
855 respectively, in the central compartment. Since this model is linear, we can solve for
856 these quantities by setting the left-hand-side of Eq. S21-S26 to zero:

$$[\text{G}]^* = [\text{G}]_{\text{diet}}\left(\frac{\delta k_{\text{a,G}}}{(k_{\text{a,G}} + \delta)(k_{\text{glycolysis}} + \text{CL}_\text{G})}\right), \tag{S28}$$

$$[\text{P}]^* = \frac{[\text{G}]_{\text{diet}}A_1 + [\text{L}]_{\text{diet}}A_2}{A_3}, \tag{S29}$$

$$A_1 = 2\delta k_{\text{a,G}}(k_{\text{a,L}} + \delta)k_{\text{glycolysis}}(\text{CL}_\text{L} + k_{\text{Ldh-f}}), \tag{S30}$$

$$A_2 = \delta k_{\text{a,L}}(k_{\text{a,G}} + \delta)(\text{CL}_\text{G} + k_{\text{glycolysis}})k_{\text{Ldh-f}}, \tag{S31}$$

$$A_3 = (k_{\text{a,G}} + \delta)(k_{\text{a,L}} + \delta)(\text{CL}_\text{G} + k_{\text{glycolysis}})[k_{\text{Ldh-f}}(\text{CL}_\text{P} + k_{\text{OXPHOS}}) + \text{CL}_\text{L}(\text{CL}_\text{P} + k_{\text{Ldh-r}}$$
$$+ k_{\text{OXPHOS}})]. \tag{S32}$$

Note that the energy flux is a linear function of the input dietary compositions $[\text{G}]_{\text{diet}}$
and $[\text{L}]_{\text{diet}}$. In our experiments, we impose a linear relationship between these two
compositions to maintain approximately equal caloric density ($\Delta[\text{G}]_{\text{diet}} = -2\Delta[\text{L}]_{\text{diet}}$).
Thus, this model is not capable of reproducing the nonlinear trends we see in our assay

in which a mixture of glucose and lactic acid prolongs lifespan more than diets of either
compound alone.
870
Genome-scale metabolic model of glucose and lactic acid metabolism
The linear model analyzed above is a substantial simplification of *Drosophila*
metabolism, which involves thousands of reactions and factors such as redox balance.
Before we turned our attention to models with active regulation of metabolic activity,
we first sought to determine whether a more realistic model of metabolism could
explain our experimental results. To do this, we analyzed iDrosophila1, a genome-scale
metabolic model (GEM) of *Drosophila* metabolism (Cesur et al., 2023). The goal of a GEM
is to represent the complete set of metabolic reactions that can occur within an
organism, typically constructed by referencing the organism's annotated genome.
GEMs can then be used to model metabolism using flux-balance analysis (FBA), which
predicts the metabolic fluxes in an organism given an objective function such as
biomass production. There are significant differences between the GEM model
formalism and the PK model formalism. While the PK model specifies a set of reaction
kinetics and estimates the steady-state chemical and energy fluxes given a diet, FBA-
based modeling finds a flux configuration maximizing an objective function, ignoring
reaction kinetics. Another notable difference is that iDrosophila1 models whole-
organism growth, considering factors such as amino acids that are not explicitly
accounted for in our energy-based PK model. However, this reliance on non-energetic
factors is represented in the version of the energy balance model with a second limiting
factor discussed in a previous section. Despite these differences, iDrosophila1 still
provides a useful benchmark for our simple PK model. If our basic assumptions about
energy metabolism are correct, the two models should produce similar behavior despite
the vast differences in complexity.
894
As a baseline diet for the iDrosophila1 analysis, we used metabolite intake values
calculated for the Holidic defined diet (Piper et al., 2014; Schönborn et al., 2019). The
primary carbon source in this diet is sucrose, a disaccharide composed of glucose and
fructose. To maintain a similar dietary energy composition, we replaced one unit of
sucrose flux with two units of glucose flux. Similarly, one unit of sucrose flux was
replaced with four units of lactic acid flux. We altered diet composition by changing the
maximum uptake rates of these carbon sources. Note that while our experiments
indicated that that lactic acid and lactate are not equivalent, with only lactic acid
extending starvation survival, iDrosophila1 only considers L-lactate. L-lactate supports
similar growth as other carbon sources in iDrosophila1.
905
In Fig. S9, we show the predicted growth rates from iDrosophila1 supplied the Holidic
diet with mixtures of glucose and L-lactate as the primary carbon sources. We used the

gurobi solver in the MATLAB version of the COBRA toolbox to solve for the optimal
flux configuration and growth rate. The "1X" condition corresponds to the maximum
carbon flux in the original Holidic diet, while "2X" corresponds to double that flux.
iDrosophila1 predicted similar behavior as the linear PK model, with a linear
relationship between glucose fraction and growth. Similar to the PK model, glucose
resulted in a slight growth advantage over L-lactate. Consistent with our energy
balance models, iDrosophila1 predicted saturation of growth rate as overall carbon flux
was increased: the 2X carbon condition did not provide double the growth rate in the
1X condition. This saturation occurs as growth under the Holidic diet is limited by
essential amino acids, not carbon or energy influx (Schönborn et al., 2019).
918
919 From this analysis, we conclude that our experimental results cannot be
920 straightforwardly explained by a more realistic model of the *Drosophila* metabolic
921 network. Despite only encoding a few reactions, our linear PK model performs similarly
922 to a GEM encoding 8,230 reactions.
923
Pharmacokinetic-pharmacodynamic model of Drosophila energy metabolism incorporating
OXPHOS regulation
Our analysis of the linear PK model and the iDrosophila model strongly indicates that
some form of diet-dependent regulation is required to produce the trends seen in our
experiment. Thus, we explored a pharmacokinetic-pharmacodynamic model in which
dietary components can alter the regulation of energy metabolism. We considered a
Gaussian function in which the ratio of glucose and lactic acid within the gut
compartment modulates OXPHOS activity:

$$k_{\text{OXPHOS}} = k_{\text{OXPHOS}}^{\max} \left(f_{\text{basal}} + (1 - f_{\text{basal}}) \exp \left[\frac{\left(\frac{[G]_{\text{gut}}}{[L]_{\text{gut}}} - r \right)^2}{2w^2} \right] \right), \qquad (S33)$$

933 where $k_{\text{OXPHOS}}^{\max}$ is the maximum OXPHOS rate, f_{basal} is the fraction of OXPHOS activity
934 that is basally active, r is the ratio of gut glucose to lactic acid at which OXPHOS
935 activity is maximized, and w is a parameter controlling the width of the regulation
936 function. This function produces a bell curve response, with OXPHOS activity
937 maximized at $\frac{[G]_{\text{gut}}}{[L]_{\text{gut}}} = r$ and minimized when only glucose or lactic acid are supplied.
938
Given that our goal is a qualitative recapitulation of the phenomena seen in our
experimental starvation assays, we did not attempt to quantitatively parameterize the
model based on our experiments. Owing to the large number of parameters in this
model, such parameterization likely would require an intractable number of additional
experiments. Instead, to analyze the model, we numerically integrated the ODEs until

all variables reached a steady state using MATLAB's *ode15s* integrator. We found that
this model can qualitatively recapitulate the three key phenomena observed in our
starvation assays (Fig. 4E). In the simulations shown in the main text, we assumed that
nutrient absorption is rapid relative to gut dilution and that Ldh operates primarily in
the forward direction (i.e., generating pyruvate). Table S1 contains a detailed list of all
parameters. For the model representing Ldh knockdown, we reduced the value of both
the forward and reverse Ldh reaction coefficients by the specified knockdown fraction.

Note that multiple regulatory schemes could produce similar behavior to this model.
For example, OXPHOS regulation instead being dependent on the internal
compartment glucose ratio $[G]/[L]$ would produce similar behavior. Non-monotonic
regulation in response to only lactic acid levels could also produce similar effects. It is
also possible that regulation of other enzymes within the pathway could produce non-
monotonicity in survival. For example, in Eq. S28-S32 the portion of energy flux from
OXPHOS is dependent on the forward and reverse rate constants of Ldh. Thus, a
regulatory function modulating these two rates may exist that can reproduce the non-
monotonicity in survival. However, such a regulatory scheme is less parsimonious than
the proposed OXPHOS regulatory scheme and inconsistent with prior *in vitro* work
finding that glucose and lactic acid mixtures regulate OXPHOS (Cai et al., 2023). A
scheme in which Ldh is the key regulatory target would also likely be inconsistent with
the results of our Ldh knockdown and metformin experiments.

The above PKPD model, while it captures the survival dynamics of flies fed glucose and
lactic acid diets, cannot capture the results of the priming experiment as it assumes that
regulation is instantaneous. To capture the observed priming results, one could
incorporate concepts from indirect response pharmacodynamic models, in which an
effector modulates an intermediate factor that then impacts the phenotype of interest
(Sharma and Jusko, 1998).

Analysis of fat breakdown data
We analyzed the fat breakdown time series using multiple fitting approaches. We first
estimated the time to 50% fat breakdown by fitting a piecewise linear model to the time
series. We identified the time interval in which the mean remaining body fat percentage
crossed 50%. In all time series, this crossing occurred only once. Once we identified this
interval, we fit a linear model $y = \beta_1 t + \beta_0$, where t is time and y is the percentage of
remaining body fat, using MATLAB's *fit* function and computed parameter confidence
intervals using the *confint* function. From these parameters, we estimated the time at
which the body fat decreased to 50% of its initial value as $t_{50} = (50 - \beta_0)/\beta_1$. To

estimate the confidence interval of this depletion time, we used the standard error

propagation formula: $\sigma_{t_{50}}^2 = \sigma_{\beta_0}^2 \left(\frac{\partial t_{50}}{\partial \beta_0}\right)^2 + \sigma_{\beta_1}^2 \left(\frac{\partial t_{50}}{\partial \beta_1}\right)^2$.

To obtain the average rate of fat depletion, we fit a linear model to the entire time series
using the methods specified above. We fixed the intercept to $\beta_0 = 100$ and estimate the
depletion rate β_0. From this linear fit, we computed an additional estimate of the t_{50}.
For estimating the confidence interval of this estimate of t_{50}, we set $\sigma_{\beta_0}^2 = 0$. We also fit
an exponential model, $y = \beta_0 e^{-\beta_1 t}$, where again $\beta_0 = 100$, using the same MATLAB
functions as with the linear model. The exponential model exhibited a somewhat lower
adjusted r^2 than the linear model.

While the results varied slightly across these three analysis methods, all supported the
same qualitative conclusion: starved flies exhibited the fastest fat breakdown, flies fed
pure glucose or lactic acid exhibited an intermediate rate, and flies fed a mixture of
glucose and lactic acid exhibited the slowest fat breakdown.

Modeling acid/base equilibria in fly diets
Our experiments indicated a substantial difference between the effects of lactate/acetate
versus lactic acid/acetic acid, with sodium lactate/acetate providing minimal lifespan
extension. To understand the composition of diets initially containing the weak acid
form of these compounds, we use standard acid/base modeling approaches. Consider a
mixture of two weak acids, HA and HB, with logarithmic dissociation constants $pK_{a,A}$
and $pK_{a,B}$ and initial concentrations $[HA]_0$ and $[HB]_0$, respectively. At equilibrium, a
fraction of these acids will dissociate into their conjugate base with concentrations $[A^-]$
and $[B^-]$. Neglecting the small initial concentration of protons present in water, the final
proton concentration will be $[H^+] = [A^-] + [B^-]$. The remaining concentration of weak
acid can be found by mass balance given the conjugate base concentrations. At
equilibrium, the concentrations will be

1010
$$K_{a,A} = \frac{[A^-]([A^-] + [B^-])}{[HA]_0 - [A^-]},$$
(S33)

1011
$$K_{a,B} = \frac{[B^-]([A^-] + [B^-])}{[HB]_0 - [B^-]}.$$
(S34)

1012 For a given mixture of lactic acid ($pK_a = 3.86$) and acetic acid ($pK_a = 4.76$), we
1013 numerically solved these equations for $[A^-]$ and $[B^-]$ for a range of dietary
1014 compositions using MATLAB's *fsolve* function (Fig. S5C, D). For both single acid and
1015 mixed acid diets, the weak acid form of the compound is predicted to dominate at
1016 equilibrium.

Supplementary Tables

Parameter	Name	Value
δ	Gut dilution rate	0.5
$[G]_{\text{diet}}$	Dietary glucose concentration	Varies
$[L]_{\text{diet}}$	Dietary lactic acid concentration	Varies
$k_{a,G}$	Glucose absorption rate	100
$k_{a,L}$	Lactic acid absorption rate	100
$k_{\text{glycolysis}}$	Glycolysis rate	1.5
$k_{\text{Ldh}-f}$	Forward Ldh rate	1
$k_{\text{Ldh}-r}$	Reverse Ldh rate	0.01
k_{OXPHOS}	OXPHOS rate	0.1, varies in regulation model
CL_G	Glucose clearance rate	0.1
CL_L	Lactic acid clearance rate	0.1
CL_P	Pyruvate clearance rate	0.1
$Y_{\text{glycolysis}}$	Per-glucose ATP yield of glycolysis	2
Y_{OXPHOS}	Per-pyruvate ATP yield of OXPHOS	17
$k_{\text{OXPHOS}}^{\max}$	Maximum OXPHOS rate	10
f_{basal}	Basal fraction of OXPHOS activity	0.01
r	Optimal ratio of glucose to lactic acid	0.5
w	Width of OXPHOS regulation function	0.13

Table S1: Parameters used in the PKPD model simulations shown in Fig. 4D-F. All rate
parameters have units of inverse time. These parameters are not a quantitative fit to the
reaction kinetics of metabolism in *Drosophila*, but rather represent a set of parameters
that provide a qualitative recapitulation of our data. The linear PK model simulated
with these parameters produced qualitatively similar trends to those predicted by the
iDrosophila1 genome-scale metabolic model (Cesur et al., 2023).
1025

Supplementary Text

Estimation of the maximal caloric content of CDM and lactic acid/glucose diets

Here, we outline our approach for estimating the maximum available energy densities of the diets used in our experiments. We compute the Gibbs free energy of combustion (ΔG) of dietary components using eQuilibrator (Beber et al., 2022), a calculator for thermodynamic analysis of biochemical reaction systems. This Gibbs free energy represents the maximum possible energy yield from complete oxidation of the dietary component. We calculated these free energies at pH 7, pMg 3, ionic strength 0.25 M, and reactant concentrations of 1 mM. The dominant source of energy and carbon in CDM is glucose, which has $\Delta G = -2911$ kJ/mol. Lactic acid fermentation converts a molecule of glucose into two molecules of lactic acid, which each have $\Delta G = -1347$ kJ/mol. Thus, complete fermentation of all glucose in CDM would result in only a ~7% loss in energy. As the spent media are filter sterilized before being fed to flies, we neglected the energy that the fly could gain from digestion of bacterial cells. This 7% difference in energy cannot explain the survival effects of *Lp*-spent medium nor the difference in survival between the diets composed of lactic acid or glucose alone. To mimic the outcome of lactic acid fermentation, we constructed our lactic acid and glucose diets by exchanging glucose with lactic acid at a 1:2 ratio. We refer to the resulting diets as "near isocaloric" since the differences are small relative to the nonlinear glucose-lactic acid interaction we observe.

Note that the increased energy yield we find in the *Lp*-spent medium condition indicates that the actual energy yield of *Drosophila* from 1X CDM diets is substantially lower than the maximum estimated here. Indeed, our model estimates that in the 1X CDM condition, at least ~45% of the theoretically available dietary energy is left unutilized.

Estimation of in situ Drosophila *microbiome metabolic fluxes*

First, we determined a relationship between *Lp* growth and lactate production. A prior study found that in MRS medium *Lp* grows to a density of $\sim 2 \times 10^9$ CFU/mL and produces ~200 mM of D+L-lactate (Fig. 2A and C) (Aranda-Díaz et al., 2020). These values lead to lactate yield estimate of $Y = 10^{-13}$ mol/CFU. As lactic acid fermentation is a central metabolic reaction, we assume the yield does not vary substantially between growth conditions.

Next, we estimated the typical population size and growth rate of *Lp* within non-starved *Drosophila.* In our experiments, we found that the steady-state *Lp* population, ρ_{LP}, within well-fed flies was $\sim 5 \times 10^5$ CFU/fly (Fig. S1). Assuming that the dominant source of death within the fly gut is dilution out of the gut, the steady-state growth rate

of Lp must match the rate of dilution. The transit time of a fly gut is ~1 h (Fig. 7D, von Frieling et al., 2020), corresponding to a dilution rate of $\delta \sim 24$ day^{-1}. Thus, the estimated per-fly lactate flux is $\rho_{\text{LP}} Y \delta = 1.2 \times 10^{-6}$ mol/day, corresponding to a carbon atom flux of 3.6×10^{-6} mol/day.

Finally, we determined whether the above flux is significant relative to the incoming flux of carbon compounds from a typical diet. Our CAFÉ assay data indicated that flies consume ~1.5 μL/day of food across diets. Well-fed flies consume a diet in which the primary carbon sources are yellow cornmeal (61.3 g/L) and molasses (75.2 mL/L). Yellow cornmeal is 77% carbohydrates by mass (FoodData Central ID: 169697, USDA, 2025). Molasses is 74.7% carbohydrates by mass and has a specific gravity of 1.33 (FoodData Central ID: 168820, USDA, 2025). The mass density of molasses in the fly food is therefore 101.5 g/L. Thus, an estimate of the total carbohydrate mass density of the food is 122 g/L. The carbohydrates in these two carbon sources are mostly composed of simple sugars or polymers of simple sugars. Assuming all sugars have a composition similar to glucose, this diet contains a glucose-equivalent concentration of 677 mM, corresponding to an input glucose flux of 1×10^{-6} mol/day and a carbon atom flux of 6×10^{-6} mol/day. The estimated Lp-derived lactic acid carbon atom flux of 3.6×10^{-6} mol/day is on the same order of magnitude as the dietary carbon flux, indicating that Lp cells can potentially produce physiologically relevant fluxes of lactic acid *in situ*.

Acknowledgements

We thank Dr. François Leulier for sharing the Lp^{WCFS1} and $Lp^{\Delta Ldh}$ strains with kind
permission from Dr. Pascal Hols (Ferain et al., 1996). We thank Dr. John Vaughen and
Dr. Thomas Clandinin for permitting use of the DAM and help with analyzing data. We
are grateful to M. Cesur, K. Patil, R. Porter, B. Fu, and J. Sonnenburg and members of
the O'Brien, Huang, Good, and Sonnenburg labs for helpful discussions. We are
thankful for Dr. L. Wat's kind sharing of a cartoon fly illustration. Stocks obtained from
the Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study.
We thank the TRiP at Harvard Medical School (NIH/NIGMS R01-GM084947) for
providing transgenic RNAi fly stocks used in this study. We acknowledge critical
resources and information provided by FlyBase (Öztürk-Çolak et al., 2024). FlyBase is
supported by a grant from the National Human Genome Research Institute at the U.S.
National Institutes of Health (U41 HG000739) and by the British Medical Research
Council (MR/N030117/1). The authors acknowledge support from a Helen Hay Whitney
Postdoctoral Fellowship (to J.W.M.), a Stanford PRISM Baker Fellowship (to J.A.L.), the
Bio-X Summer Research Experience for Undergraduates program (to A.M.S.), NSF
Awards IOS 2032985 and CAREER 2144342 (to W.B.L.), CZI Grant DAF2020-217689 (to
L.E.O. and K.C.H.), NIH Awards R35GM141885 (to L.E.O.) and R35GM146949 (to
B.H.G.), and NSF Award EF-2125383 (to. K.C.H.). B.H.G., L.E.O., and K.C.H. are Chan-
Zuckerberg Biohub Investigators.

Data availability

All codes and processed data used to generate figures are available at
https://github.com/jamie-alc-lopez/fly_energy_metabolism.

Figures

Figure 1: Microbe-derived metabolites impact *Drosophila* starvation survival.

    A) Schematic of *Drosophila* survival assay in flies fed chemically defined media (CDM) supernatant versus bacterial spent media and expected outcomes and interpretations. Spent media from stationary phase (48 h) cultures were obtained from fly gut commensals (*Lp* and *Ap*) grown in monoculture and co-culture by centrifugation and filtering before mixing with agar to make a solid substrate for fly survival quantification.

    B) Survival is determined by nutrient-driven interactions impacting energy yield, with the potential for both increased survival from beneficial metabolites and shortened survival from e.g., nutrient competition.

    C) Mean survival was significantly higher in *Ore-R* mated females fed *Lp*- and *Ap*-spent media compared to CDM ($p < 0.0001$ for both comparisons; one-way ANOVA followed by Dunnett's multiple comparison test). $N = 7$ biological replicates, $n = 70$ flies per condition. Mean survival was not significantly different between *Ore-R* mated females fed $LpAp^{co}$-spent medium compared to flies fed CDM supernatant ($p = 0.3$; one-way ANOVA followed by Dunnett's multiple comparison test). $N = 7$ biological replicates, $n = 70$ flies per condition. ****: $p < 0.0001$; ns: not significant; error bars indicate 1 SEM.

Figure 2: Microbe-derived metabolites cause survival to deviate from simple energy balance models in starvation regimes with substantial energy limitation.

A) Schematic of energy balance model used to interpret *Drosophila* survival data as described by Eq. 1. This model considers a population of flies with mean initial energy storage μ_E and each fly expends energy at flux ϕ_O and intakes energy at flux ϕ_I.

B) Mean survival increased with increasing CDM concentration.

C,D) Mean survival (C) and variance (D) increased superlinearly between 0-1.5X CDM, and saturate between 2-4X CDM.

E,F) Energy balance model predictions for mean survival (E) and variance (F) as a function of increasing relative caloric value. Survival is predicted to be superlinear in the energy limitation regime and to saturate when energy demands for normal lifespan are met.

G) The percent difference in mean survival relative to the control was higher at lower caloric values than at high caloric values for *Lp*- and *Ap*-spent media. Error bars indicate 1 SEM.

1149

Figure 3: *Lp*-derived lactic acid is necessary and sufficient for starvation survival and acts via an interaction with glucose metabolism.

A) Mean fold change of LC-MS signal intensity from spent media relative to fresh CDM supernatant for each species across all metabolomic features, annotated or otherwise. *N*=3 biological replicates.

B) Principal component analysis of fresh CDM supernatant and spent media from monocultures and co-culture.

C) Lactic acid levels in fresh CDM supernatant and spent media from monocultures and co-culture. Lactic acid was significantly more concentrated in *Lp*- compared with *LpAp*^{co}-spent medium ($p = 0.0002$; one-way ANOVA followed by Tukey HSD test). $N = 3$ biological replicates.

D) Mean survival was significantly lower in *Ore-R* mated females fed spent medium from the *Lp*$^{\Delta Ldh}$ mutant or CDM supernatant compared with flies fed spent medium from the parent strain *Lp*^{WCFS1} ($p < 0.0001$ for both comparisons; one-way ANOVA followed by Tukey HSD test). Mean survival was not significantly different between *Ore-R* mated females fed *Lp*$^{\Delta Ldh}$-spent medium compared to flies fed CDM supernatant ($p = 0.51$; one-way ANOVA followed by Tukey HSD test). $N = 7\text{-}9$ biological replicates, $n = 70\text{-}90$ flies per condition.

E) Mean survival was significantly higher in *Ore-R* mated females fed 1X 50% CDM + 50% lactic acid compared to flies fed 1X CDM ($p < 0.0001$; Student's *t* test). $N = 8\text{-}10$ biological replicates, $n = 80\text{-}100$ flies per condition.

F) Mean survival was significantly higher in *Ore-R* mated females fed 1X 50% glucose + 50% lactic acid compared to flies fed 1X glucose or 1X lactic acid ($p < 0.0001$ for both comparisons; one-way ANOVA followed by Tukey HSD test). Mean survival was not significantly different between *Ore-R* mated females fed 1X glucose compared to flies fed 1X lactic acid ($p = 0.98$; one-way ANOVA followed by Tukey HSD test). $N = 5$ biological replicates, $n = 50$ flies per condition.

G) Mean survival was significantly higher in *Ore-R* mated females fed 1X lactic acid for 5 days and then switched to 1X glucose after 5 days of lactic acid feeding compared to flies with the opposite feeding regimen (1X glucose followed by 1X lactic acid) or flies maintained on 1X 50% glucose + 50% lactic acid mixture ($p < 0.0001$ for both comparisons; one-way ANOVA followed by Tukey HSD test). Mean survival was not significantly different between *Ore-R* mated females first fed 1X lactic acid followed by 1X glucose compared to flies maintained on the 1X 50% glucose + 50% lactic acid mixture ($p = 0.83$; one-way ANOVA followed by Tukey HSD test). $N = 10$ biological replicates, $n = 100$ flies per condition. ****: $p < 0.0001$; ns: not significant; error bars indicate 1 SEM.

Figure 4: Lactic acid impacts starvation survival via regulation of oxidative phosphorylation (OXPHOS).

A) Percentage body fat in the form of triglycerides over 5 days of starvation in *Ore-R* mated females fed 1X glucose, 1X lactic acid, or 1X 50% glucose + 50% lactic acid. $N = 4$ biological replicates, $n = 20$ flies per condition.

B) The rate of triglyceride mobilization as determined by a linear fit was significantly slower in flies fed 1X 50% glucose + 50% lactic acid compared to flies fed 1X glucose or 1X lactic acid ($p = 0.017$ and 0.012, respectively; one-way ANOVA followed by Tukey HSD test). The rate of triglyceride mobilization was not significantly different between *Ore-R* mated females fed 1X glucose compared to flies fed 1X lactic acid ($p = 0.97$; one-way ANOVA followed by Tukey HSD test).

C) Schematic of pharmacokinetic-pharmacodynamic (PKPD) model used to interpret survival data. This model considers the fate of glucose and lactic acid in *Drosophila* using a simplified representation of central carbon metabolism pathways.

D) In the absence of regulation, PKPD model predicts equivalent mean survival between flies fed any ratio of glucose and lactic acid at either low or high caloric values.

E) PKPD model with regulation of OXPHOS dependent on the ratio of lactic acid to glucose levels predicts increased mean survival at intermediate ratios. The increase in survival is highest at low caloric values.

F) PKPD model with OXPHOS regulation predicts increased mean survival across a range of Ldh knockdown efficiencies. The magnitude of increased survival due to lactic acid addition to a glucose diet is highest with full Ldh expression.

G,H) Mean survival was significantly lower in *Ore-R* mated females treated with 25 mM metformin and fed 1X 50% glucose + 50% lactic acid compared to vehicle-treated flies with the same diet ($p < 0.0001$; two-way ANOVA followed by Sidak's multiple comparison test). Metformin treatment did not affect mean survival for complete starvation in *Ore-R* mated females ($p > 0.99$; two-way ANOVA followed by Sidak's multiple comparison test). Mean survival was significantly higher in *Ore-R* mated females treated with 25 mM metformin and fed 1X glucose compared to vehicle-treated flies fed 1X glucose ($p < 0.0001$; two-way ANOVA followed by Sidak's multiple comparison test). Mean survival was significantly lower in *Ore-R* mated females treated with 25 mM metformin and fed 1X lactic acid compared to vehicle-treated flies fed 1X lactic acid ($p < 0.0001$; two-way ANOVA followed by Sidak's multiple comparison test). $N = 10$ biological replicates, $n = 100$ flies per condition. *: $p < 0.05$, ****: $p < 0.0001$; ns: not significant; error bars indicate 1 SEM.

1228 **Supplemental Figures**

1229

1230

1231 **Figure S1: Microbial load decreases during starvation and composition shifts to**
1232 **increased frequency of the *Lactiplantibacillus* genus.**

A) Bacterial load in the *Drosophila* gut as measured by colony forming units on agar
selective for *Lactiplantibacillus plantarum* (*Lp*) or *Acetobacter pasteurianus* (*Ap*)
growth over 24 h of starvation. Microbe load decreased for both species.

B) Microbiome composition measured as the relative abundance of *Lp* and *Ap* shifts
toward increased *Lp* frequency.

C) Percentage of flies with any culturable microbes of either species decreased
during starvation. More *Lp* cells were culturable at 24 h of starvation than *Ap*
cells. $N = 40$ biological replicates.

Figure S2: Microbe-derived metabolites do not impact starvation survival via differential food intake, active energy expenditure, or fecundity.

A) Taste preference in *Ore-R* mated females as quantified by a dye-based, two-choice feeding assay demonstrated that fresh CDM supernatant, *Lp*-spent medium, and *Ap*-spent medium are appetitive to flies. $N = 10$ biological replicates, $n = 100$ flies per condition.

B) Taste preference in *Ore-R* mated females between CDM supernatant and *Lp*-spent medium was neutral, however flies preferred CDM supernatant or *Lp*-spent medium compared to *Ap*-spent medium. $N = 10\text{-}18$ biological replicates, $n = 100\text{-}180$ flies per condition.

C) Taste preference in *Ore-R* mated females between fresh CDM supernatant and *Lp*-spent medium remained neutral during prolonged starvation. $N = 8\text{-}10$ biological replicates, $n = 80\text{-}100$ flies per condition.

D) Food intake did not change in *Ore-R* mated females between flies fed fresh CDM supernatant, *Lp*-spent medium, and *Ap*-spent medium.

E) Locomotor activity in *Ore-R* mated females as measured using the *Drosophila* activity monitor (DAM) over 48 h of feeding with fresh CDM supernatant, *Lp*-spent medium, and *LpAp*^{co}-spent medium.

F) Time spent sleeping in *Ore-R* mated females measured using DAM over 48 h of feeding with fresh CDM supernatant, *Lp*-spent medium, and *LpAp*^{co}-spent medium.

G) Daily activity was similar among *Ore-R* mated females fed fresh CDM supernatant, *Lp*-spent medium, and *LpAp*^{co}-spent medium. $N = 24\text{-}28$ biological replicates.

H) The number of eggs laid in 24 h by *Ore-R* mated females fed with CDM supernatant, *Lp*-spent medium, and *Ap*-spent medium was similar. $N = 6$ biological replicates, $n = 60$ flies per condition.

I) Energy balance model prediction for the change in food intake necessary to account for the observed survival through a change in energy intake.

J) Energy balance model prediction for the change in activity necessary to account for the observed survival through a change in energy output.

1273

Figure S3: Caloric value affects how spent media impact starvation survival.
Mean survival was significantly higher in *Ore-R* mated females fed *Lp*-spent medium
compared to CDM supernatant ($p = 0.043$; one-way ANOVA followed by Dunnett's
multiple comparison test). Mean survival was not significantly different between *Ore-R*
mated females fed *Ap*-spent medium compared to CDM ($p = 0.59$; one-way ANOVA
followed by Dunnett's multiple comparison test). Mean survival was significantly
shorter in *Ore-R* mated females fed *LpAp*^{co}-spent medium compared to CDM ($p = 0.024$;
one-way ANOVA followed by Dunnett's multiple comparison test). $N = 3$ biological
replicates, $n = 30$ flies per condition. *: $p < 0.05$; ns: not significant; error bars indicate 1
SEM.

Figure S4: Caloric value affects how lactic acid impacts fly survival.

A) Mean survival was not significantly different in *Ore-R* mated females fed 2X 50% CDM + 50% lactic acid compared to flies fed 2X CDM ($p = 0.41$; Student's t test). $N = 4\text{-}10$ biological replicates, $n = 40\text{-}100$ flies per condition.

B) The percent difference in mean survival relative to the control was higher at lower caloric values in the strong energy limitation regime compared with high caloric values in the survival saturation regime for flies fed CDM versus flies fed CDM supplemented with lactic acid.

C) Mean survival was significantly higher in *Ore-R* mated females fed 2X glucose compared to flies fed 1X glucose ($p < 0.0001$; two-way ANOVA followed by Sidak's multiple comparison test). Mean survival was not significantly different between *Ore-R* mated females fed 2X lactic acid compared to flies fed 1X lactic acid or between flies fed 2X 50% glucose + 50% lactic acid compared to flies fed 1X 50% glucose + 50% lactic acid ($p = 0.99$ and 0.18, respectively; two-way ANOVA followed by Sidak's multiple comparison test). $N = 7\text{-}8$ biological replicates, $n = 70\text{-}80$ flies per condition.

D) The percent difference in mean survival relative to the control was higher at lower caloric values in the strong energy limitation regime compared with high caloric values in the survival saturation regime for flies fed glucose + lactic acid versus control diet flies. ****: $p < 0.0001$; ns: not significant; error bars indicate 1 SEM.

Figure S5: Neither dietary lactate nor neutralized lactic acid extend fly survival.

A) Mean survival was not significantly different in *Ore-R* mated females fed 100 mM sodium lactate or 100 mM sodium acetate compared to completely starved flies (p = 0.87 and 0.37, respectively; one-way ANOVA followed by Dunnett's multiple comparison test). N = 3 biological replicates, n = 30 flies per condition.

B) Mean survival was not significantly different in *Ore-R* mated females fed 1X lactic acid corrected to pH 7 with dropwise sodium hydroxide compared to completely starved flies (p = 0.80; Student's t test). N = 3-5 biological replicates, n = 30-50 flies per condition.

C) Fraction of lactic acid and acetic acid predicted to exist in the weak acid form across varying acid to glucose ratios in single acid solutions.

D) Fraction of lactic acid and acetic acid predicted to exist in the weak acid form across varying acid to glucose ratios in mixed acid solutions.

1320

Figure S6: Whole-body Ldh is dispensable for lactic acid-mediated effects on survival.

A) Mean survival was significantly higher in $tub^{ts} > +$ mated females (genotype: w^{1118} ;; *tub-GAL4, tub-GAL80^{ts}/+*) fed 1X 50% glucose + 50% lactic acid compared to flies fed 1X glucose or 1X lactic acid ($p < 0.0001$ for both comparisons; two-way ANOVA followed by Tukey HSD test). Mean survival was not significantly different between $tub^{ts} > +$ mated females fed 1X glucose compared to flies fed 1X lactic acid ($p = 0.86$; two-way ANOVA followed by Tukey HSD test).

B) Mean survival was significantly higher in $tub^{ts} > Ldh$-*RNAi* mated females (genotype: w^{1118} ;; *tub-GAL4, tub-GAL80^{ts}/UAS-Ldh-RNAi*) fed 1X 50% glucose + 50% lactic acid compared to 1X glucose but not 1X lactic acid fed flies ($p = 0.0003$ and 0.98, respectively; two-way ANOVA followed by Tukey HSD test).

C) Mean survival was not significantly different between $tub^{ts} > +$ and $tub^{ts} > Ldh$-*RNAi* mated females in either starved or 1X glucose feeding conditions ($p = 0.39$ and 0.61, respectively; two-way ANOVA followed by Tukey HSD test). Mean survival was significantly increased in $tub^{ts} > Ldh$-*RNAi* compared to $tub^{ts} > +$ mated females fed 1X lactic acid ($p = 0.0005$; two-way ANOVA followed by Tukey HSD test). Mean survival was significantly shorter in $tub^{ts} > Ldh$-*RNAi* compared to $tub^{ts} > +$ mated females fed 1X 50% glucose + 50% lactic acid ($p = 0.0027$; two-way ANOVA followed by Tukey HSD test). $N = 10$ biological replicates, $n = 100$ flies per condition. **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$; ns: not significant; error bars indicate 1 SEM.

1343

Figure S7: Metformin impacts survival in flies fed glucose and lactic acid.
Survival was significantly longer in *Ore-R* mated females treated with 25 mM
metformin and fed 1X glucose compared to vehicle-treated flies fed 1X glucose,
potentially due to a titrating effect on energy yield from OXPHOS. Survival was
significantly reduced in *Ore-R* mated females treated with 25 mM metformin and fed 1X
lactic acid compared to vehicle-treated flies fed 1X lactic acid. $N = 10$ biological
replicates, $n = 100$ flies per condition.

Figure S8: Transformed survival moments of CDM dilution series compared to energy balance model

A) Inverse mean survival is predicted to scale linearly between 0-1.5X CDM and to saturate between 2-4X CDM.

B) Energy balance model predictions for inverse mean survival as a function of increasing relative caloric value. Inverse mean survival is predicted to scale linearly in the energy limitation regime and to saturate when energy demands for normal lifespan are met.

1361

Figure S9: Genome-scale metabolic model of *Drosophila* metabolism (iDrosophila1)

predicts a linear relationship between diet composition and overall metabolic flux.

Predicted growth rate, a normalized representation of metabolic flux of ATP and other

essential compounds, in the iDrosophila1 model supplied with mixtures of only glucose

and L-lactate. Diet composition was modified by changing the maximum allowable

input flux of these two carbon sources. Aside from input fluxes of sucrose (set to zero),

glucose, and L-lactate, the fly diet composition is identical to the Holidic diet condition

from previous studies (Schönborn et al., 2019).

References

Agrawal, N., Delanoue, R., Mauri, A., Basco, D., Pasco, M., Thorens, B., Léopold, P., 2016. The Drosophila TNF Eiger Is an Adipokine that Acts on Insulin-Producing Cells to Mediate Nutrient Response. Cell Metabolism 23, 675–684. https://doi.org/10.1016/j.cmet.2016.03.003

Agriculture, U.S.D. of, 2025. FoodData Central.

Agus, A., Clément, K., Sokol, H., 2021. Gut microbiota-derived metabolites as central regulators in metabolic disorders. https://doi.org/10.1136/gutjnl-2020-323071

Aranda-Díaz, A., Obadia, B., Dodge, R., Thomsen, T., Hallberg, Z.F., Güvener, Z.T., Ludington, W.B., Huang, K.C., 2020. Bacterial interspecies interactions modulate pH-mediated antibiotic tolerance. eLife 9, e51493. https://doi.org/10.7554/eLife.51493

Arnoldini, M., Sharma, R., Moresi, C., Chure, G., Chabbey, J., Slack, E., Cremer, J., 2024. Quantifying the daily harvest of fermentation products from the human gut microbiota. https://doi.org/10.1101/2024.01.05.573977

Baggio, L.L., Drucker, D.J., 2007. Biology of Incretins: GLP-1 and GIP. Gastroenterology 132, 2131–2157. https://doi.org/10.1053/j.gastro.2007.03.054

Beber, M.E., Gollub, M.G., Mozaffari, D., Shebek, K.M., Flamholz, A.I., Milo, R., Noor, E., 2022. eQuilibrator 3.0: a database solution for thermodynamic constant estimation. Nucleic Acids Research 50, D603–D609. https://doi.org/10.1093/nar/gkab1106

Bonate, P.L., 2011. Pharmacokinetic-Pharmacodynamic Modeling and Simulation. Springer US, Boston, MA. https://doi.org/10.1007/978-1-4419-9485-1

Broderick, N.A., Lemaitre, B., 2012. Gut-associated microbes of Drosophila melanogaster. Gut Microbes 3, 307–321. https://doi.org/10.4161/gmic.19896

Brooks, G.A., 2018. The Science and Translation of Lactate Shuttle Theory. Cell Metab 27, 757–785. https://doi.org/10.1016/j.cmet.2018.03.008

Brummel, T., Ching, A., Seroude, L., Simon, A.F., Benzer, S., 2004. Drosophila lifespan enhancement by exogenous bacteria. Proceedings of the National Academy of Sciences 101, 12974–12979. https://doi.org/10.1073/pnas.0405207101

Cai, X., Ng, C.P., Jones, O., Fung, T.S., Ryu, K.W., Li, D., Thompson, C.B., 2023. Lactate activates the mitochondrial electron transport chain independently of its metabolism. Molecular Cell 83, 3904-3920.e7. https://doi.org/10.1016/j.molcel.2023.09.034

Certo, M., Llibre, A., Lee, W., Mauro, C., 2022. Understanding lactate sensing and signalling. Trends in Endocrinology & Metabolism 33, 722–735. https://doi.org/10.1016/j.tem.2022.07.004

Cesur, M.F., Basile, A., Patil, K.R., Çakır, T., 2023. A new metabolic model of *Drosophila melanogaster* and the integrative analysis of Parkinson's disease. Life Science Alliance 6, e202201695. https://doi.org/10.26508/lsa.202201695

Chatterjee, N., Perrimon, N., 2021. What fuels the fly: Energy metabolism in Drosophila and its application to the study of obesity and diabetes. Science Advances 7, eabg4336. https://doi.org/10.1126/sciadv.abg4336

Chow, C.C., Hall, K.D., 2008. The Dynamics of Human Body Weight Change. PLoS Computational Biology 4, e1000045. https://doi.org/10.1371/journal.pcbi.1000045

Churchill, E.R., Dytham, C., Thom, M.D.F., 2019. Differing effects of age and starvation on reproductive performance in Drosophila melanogaster. Sci Rep 9, 2167. https://doi.org/10.1038/s41598-019-38843-w

Cichewicz, K., Hirsh, J., 2018. ShinyR-DAM: a program analyzing Drosophila activity, sleep and circadian rhythms. Commun Biol 1, 1–5. https://doi.org/10.1038/s42003-018-0031-9

Clarke, G., Stilling, R.M., Kennedy, P.J., Stanton, C., Cryan, J.F., Dinan, T.G., 2014. Minireview: Gut Microbiota: The Neglected Endocrine Organ. Molecular Endocrinology 28, 1221–1238. https://doi.org/10.1210/me.2014-1108

De Groot, D.H., Lischke, J., Muolo, R., Planqué, R., Bruggeman, F.J., Teusink, B., 2020. The common message of constraint-based optimization approaches: overflow metabolism is caused by two growth-limiting constraints. Cellular and Molecular Life Sciences 77, 441–453. https://doi.org/10.1007/s00018-019-03380-2

De Vadder, F., Kovatcheva-Datchary, P., Goncalves, D., Vinera, J., Zitoun, C., Duchampt, A., Bäckhed, F., Mithieux, G., 2014. Microbiota-Generated Metabolites Promote Metabolic Benefits via Gut-Brain Neural Circuits. Cell 156, 84–96. https://doi.org/10.1016/j.cell.2013.12.016

Dodge, R., Jones, E.W., Zhu, H., Obadia, B., Martinez, D.J., Wang, C., Aranda-Díaz, A., Aumiller, K., Liu, Z., Voltolini, M., Brodie, E.L., Huang, K.C., Carlson, J.M., Sivak, D.A., Spradling, A.C., Ludington, W.B., 2023. A symbiotic physical niche in Drosophila melanogaster regulates stable association of a multi-species gut microbiota. Nat Commun 14, 1557. https://doi.org/10.1038/s41467-023-36942-x

Douglas, A.E., 2018. Contradictory Results in Microbiome Science Exemplified by Recent Drosophila Research. mBio 9, 10.1128/mbio.01758-18. https://doi.org/10.1128/mbio.01758-18

Drucker, D.J., Habener, J.F., Holst, J.J., 2017. Discovery, characterization, and clinical development of the glucagon-like peptides. J Clin Invest 127, 4217–4227. https://doi.org/10.1172/JCI97233

Drucker, D.J., Holst, J.J., 2023. The expanding incretin universe: from basic biology to clinical translation. Diabetologia 66, 1765–1779. https://doi.org/10.1007/s00125-023-05906-7

Evans, J.M., Morris, L.S., Marchesi, J.R., 2013. The gut microbiome: the role of a virtual organ in the endocrinology of the host. https://doi.org/10.1530/JOE-13-0131

Fagundes, R.R., Belt, S.C., Bakker, B.M., Dijkstra, G., Harmsen, H.J.M., Faber, K.N., 2024. Beyond butyrate: microbial fiber metabolism supporting colonic epithelial homeostasis. Trends in Microbiology 32, 178–189. https://doi.org/10.1016/j.tim.2023.07.014

Fast, D., Duggal, A., Foley, E., 2018. Monoassociation with Lactobacillus plantarum Disrupts Intestinal Homeostasis in Adult Drosophila melanogaster. mBio 9, 10.1128/mbio.01114-18. https://doi.org/10.1128/mbio.01114-18

Ferain, T., Hobbs, J.N., Richardson, J., Bernard, N., Garmyn, D., Hols, P., Allen, N.E., Delcour, J., 1996. Knockout of the two ldh genes has a major impact on peptidoglycan precursor synthesis in Lactobacillus plantarum. J Bacteriol 178, 5431–5437. https://doi.org/10.1128/jb.178.18.5431-5437.1996

Fujisaka, S., Watanabe, Y., Tobe, K., 2023. The gut microbiome: a core regulator of metabolism. https://doi.org/10.1530/JOE-22-0111

Gould, A.L., Zhang, V., Lamberti, L., Jones, E.W., Obadia, B., Korasidis, N., Gavryushkin, A., Carlson, J.M., Beerenwinkel, N., Ludington, W.B., 2018. Microbiome interactions shape host fitness. Proceedings of the National Academy of Sciences 115, E11951–E11960. https://doi.org/10.1073/pnas.1809349115

Hall, K.D., 2017. A review of the carbohydrate–insulin model of obesity. Eur J Clin Nutr 71, 323–326. https://doi.org/10.1038/ejcn.2016.260

Hall, K.D., 2012. Metabolism of mice and men: mathematical modeling of body weight dynamics. Current Opinion in Clinical Nutrition & Metabolic Care 15, 418. https://doi.org/10.1097/MCO.0b013e3283561150

Hall, K.D., Guo, J., 2017. Obesity Energetics: Body Weight Regulation and the Effects of Diet Composition. Gastroenterology 152, 1718-1727.e3. https://doi.org/10.1053/j.gastro.2017.01.052

Hayward, A.C., Davis, G.H.G., 1956. The isolation and classification of Lactobacillus strains from Italian saliva samples. Br. Dent. J. 43–46.

Henriques, S.F., Dhakan, D.B., Serra, L., Francisco, A.P., Carvalho-Santos, Z., Baltazar, C., Elias, A.P., Anjos, M., Zhang, T., Maddocks, O.D.K., Ribeiro, C., 2020. Metabolic cross-feeding in imbalanced diets allows gut microbes to improve reproduction and alter host behaviour. Nat Commun 11, 4236. https://doi.org/10.1038/s41467-020-18049-9

Ho, P.-Y., Nguyen, T.H., Sanchez, J.M., DeFelice, B.C., Huang, K.C., 2024. Resource competition predicts assembly of gut bacterial communities in vitro. Nat Microbiol 9, 1036–1048. https://doi.org/10.1038/s41564-024-01625-w

Ja, W.W., Carvalho, G.B., Mak, E.M., de la Rosa, N.N., Fang, A.Y., Liong, J.C., Brummel, T., Benzer, S., 2007. Prandiology of Drosophila and the CAFE assay. Proceedings of

the National Academy of Sciences 104, 8253–8256.
https://doi.org/10.1073/pnas.0702726104

Jacquet, P., Stéphanou, A., 2023. A reduced model of cell metabolism to revisit the glycolysis-OXPHOS relationship in the deregulated tumor microenvironment. Journal of Theoretical Biology 562, 111434. https://doi.org/10.1016/j.jtbi.2023.111434

Javdan, B., Lopez, J.G., Chankhamjon, P., Lee, Y.-C.J., Hull, R., Wu, Q., Wang, X., Chatterjee, S., Donia, M.S., 2020. Personalized Mapping of Drug Metabolism by the Human Gut Microbiome. Cell 181, 1661-1679.e22. https://doi.org/10.1016/j.cell.2020.05.001

Judd, A.M., Matthews, M.K., Hughes, R., Veloz, M., Sexton, C.E., Chaston, J.M., 2018. Bacterial Methionine Metabolism Genes Influence Drosophila melanogaster Starvation Resistance. Applied and Environmental Microbiology 84, e00662-18. https://doi.org/10.1128/AEM.00662-18

Katsyuba, E., Romani, M., Hofer, D., Auwerx, J., 2020. NAD+ homeostasis in health and disease. Nat Metab 2, 9–31. https://doi.org/10.1038/s42255-019-0161-5

Keebaugh, E.S., Yamada, R., Obadia, B., Ludington, W.B., Ja, W.W., 2018. Microbial Quantity Impacts Drosophila Nutrition, Development, and Lifespan. iScience 4, 247–259. https://doi.org/10.1016/j.isci.2018.06.004

Keebaugh, E.S., Yamada, R., Ja, W.W., 2019. The Nutritional Environment Influences the Impact of Microbes on Drosophila melanogaster Life Span. mBio 10, 10.1128/mbio.00885-19. https://doi.org/10.1128/mbio.00885-19

Lai, Y., Liu, C.-W., Yang, Y., Hsiao, Y.-C., Ru, H., Lu, K., 2021. High-coverage metabolomics uncovers microbiota-driven biochemical landscape of interorgan transport and gut-brain communication in mice. Nat Commun 12, 6000. https://doi.org/10.1038/s41467-021-26209-8

Lindenbaum, J., Tse-Eng, D., Butler, V.P., Rund, D.G., 1981. Urinary excretion of reduced metabolites of digoxin. The American Journal of Medicine 71, 67–74. https://doi.org/10.1016/0002-9343(81)90260-6

Leitão-Gonçalves, R., Carvalho-Santos, Z., Francisco, A.P., Fioreze, G.T., Anjos, M., Baltazar, C., Elias, A.P., Itskov, P.M., Piper, M.D.W., Ribeiro, C., 2017. Commensal bacteria and essential amino acids control food choice behavior and reproduction. PLOS Biology 15, e2000862. https://doi.org/10.1371/journal.pbio.2000862

Liu, C., Wu, J., Zhu, J., Kuei, C., Yu, J., Shelton, J., Sutton, S.W., Li, X., Yun, S.J., Mirzadegan, T., Mazur, C., Kamme, F., Lovenberg, T.W., 2009. Lactate Inhibits Lipolysis in Fat Cells through Activation of an Orphan G-protein-coupled Receptor, GPR81*. Journal of Biological Chemistry 284, 2811–2822. https://doi.org/10.1074/jbc.M806409200

Liu, H., Wang, S., Wang, J., Guo, X., Song, Y., Fu, K., Gao, Z., Liu, D., He, W., Yang, L.-L., 2025. Energy metabolism in health and diseases. Sig Transduct Target Ther 10, 1–71. https://doi.org/10.1038/s41392-025-02141-x

Liu, J., Tan, Y., Cheng, H., Zhang, D., Feng, W., Peng, C., 2022. Functions of Gut
Microbiota Metabolites, Current Status and Future Perspectives. Aging and disease
13, 1106–1126. https://doi.org/10.14336/AD.2022.0104

Mauritsson, K., Jonsson, T., 2023. A new flexible model for maintenance and feeding
expenses that improves description of individual growth in insects. Sci Rep 13,
16751. https://doi.org/10.1038/s41598-023-43743-1

Merrill, A.L., Watt, B.K., 1955. Energy Value of Foods. Human Nutrition Research
Branch, Agricultural Research Service, U. S. Department of Agriculture.

Millington, J.W., Brownrigg, G.P., Chao, C., Sun, Z., Basner-Collins, P.J., Wat, L.W.,
Hudry, B., Miguel-Aliaga, I., Rideout, E.J., 2021. Female-biased upregulation of
insulin pathway activity mediates the sex difference in Drosophila body size
plasticity. eLife 10, e58341. https://doi.org/10.7554/eLife.58341

Min, K., Flatt, T., Kulaots, I., Tatar, M., 2007. Counting calories in Drosophila diet
restriction. Experimental Gerontology 42, 247–251.
https://doi.org/10.1016/j.exger.2006.10.009

Neuman, H., Debelius, J.W., Knight, R., Koren, O., 2015. Microbial endocrinology: the
interplay between the microbiota and the endocrine system. FEMS Microbiology
Reviews 39, 509–521. https://doi.org/10.1093/femsre/fuu010

Obadia, B., Güvener, Z.T., Zhang, V., Ceja-Navarro, J.A., Brodie, E.L., Ja, W.W.,
Ludington, W.B., 2017. Probabilistic Invasion Underlies Natural Gut Microbiome
Stability. Current Biology 27, 1999-2006.e8. https://doi.org/10.1016/j.cub.2017.05.034

Owen, M.R., Doran, E., Halestrap, A.P., 2000. Evidence that metformin exerts its anti-
diabetic effects through inhibition of complex 1 of the mitochondrial respiratory
chain. Biochem J 348, 607–614.

Öztürk-Çolak, A., Marygold, S.J., Antonazzo, G., Attrill, H., Goutte-Gattat, D., Jenkins,
V.K., Matthews, B.B., Millburn, G., dos Santos, G., Tabone, C.J., FlyBase
Consortium, 2024. FlyBase: updates to the Drosophila genes and genomes
database. Genetics 227, iyad211. https://doi.org/10.1093/genetics/iyad211

Pais, I.S., Valente, R.S., Sporniak, M., Teixeira, L., 2018. Drosophila melanogaster
establishes a species-specific mutualistic interaction with stable gut-colonizing
bacteria. PLOS Biology 16, e2005710. https://doi.org/10.1371/journal.pbio.2005710

Pfeiffenberger, C., Lear, B.C., Keegan, K.P., Allada, R., 2010. Locomotor Activity Level
Monitoring Using the *Drosophila* Activity Monitoring (DAM) System: Figure 1.
Cold Spring Harb Protoc 2010, pdb.prot5518. https://doi.org/10.1101/pdb.prot5518

Piper, M.D.W., Blanc, E., Leitão-Gonçalves, R., Yang, M., He, X., Linford, N.J.,
Hoddinott, M.P., Hopfen, C., Soultoukis, G.A., Niemeyer, C., Kerr, F., Pletcher,
S.D., Ribeiro, C., Partridge, L., 2014. A holidic medium for Drosophila
melanogaster. Nat Methods 11, 100–105. https://doi.org/10.1038/nmeth.2731

Piper, M.D.W., Skorupa, D., Partridge, L., 2005. Diet, metabolism and lifespan in *Drosophila*. Experimental Gerontology, Metabolism, Aging and Longevity 40, 857–862. https://doi.org/10.1016/j.exger.2005.06.013

Postma, E., Verduyn, C., Scheffers, W.A., Van Dijken, J.P., 1989. Enzymic analysis of the crabtree effect in glucose-limited chemostat cultures of Saccharomyces cerevisiae. Applied and Environmental Microbiology 55, 468–477. https://doi.org/10.1128/aem.55.2.468-477.1989

Qin, J., Li, R., Raes, J., Arumugam, M., Burgdorf, K.S., Manichanh, C., Nielsen, T., Pons, N., Levenez, F., Yamada, T., Mende, D.R., Li, J., Xu, J., Li, Shaochuan, Li, D., Cao, J., Wang, B., Liang, H., Zheng, H., Xie, Y., Tap, J., Lepage, P., Bertalan, M., Batto, J.-M., Hansen, T., Le Paslier, D., Linneberg, A., Nielsen, H.B., Pelletier, E., Renault, P., Sicheritz-Ponten, T., Turner, K., Zhu, H., Yu, C., Li, Shengting, Jian, M., Zhou, Y., Li, Y., Zhang, X., Li, Songgang, Qin, N., Yang, H., Wang, Jian, Brunak, S., Doré, J., Guarner, F., Kristiansen, K., Pedersen, O., Parkhill, J., Weissenbach, J., Bork, P., Ehrlich, S.D., Wang, Jun, 2010. A human gut microbial gene catalogue established by metagenomic sequencing. Nature 464, 59–65. https://doi.org/10.1038/nature08821

Quinn, R.A., Melnik, A.V., Vrbanac, A., Fu, T., Patras, K.A., Christy, M.P., Bodai, Z., Belda-Ferre, P., Tripathi, A., Chung, L.K., Downes, M., Welch, R.D., Quinn, M., Humphrey, G., Panitchpakdi, M., Weldon, K.C., Aksenov, A., da Silva, R., Avila-Pacheco, J., Clish, C., Bae, S., Mallick, H., Franzosa, E.A., Lloyd-Price, J., Bussell, R., Thron, T., Nelson, A.T., Wang, M., Leszczynski, E., Vargas, F., Gauglitz, J.M., Meehan, M.J., Gentry, E., Arthur, T.D., Komor, A.C., Poulsen, O., Boland, B.S., Chang, J.T., Sandborn, W.J., Lim, M., Garg, N., Lumeng, J.C., Xavier, R.J., Kazmierczak, B.I., Jain, R., Egan, M., Rhee, K.E., Ferguson, D., Raffatellu, M., Vlamakis, H., Haddad, G.G., Siegel, D., Huttenhower, C., Mazmanian, S.K., Evans, R.M., Nizet, V., Knight, R., Dorrestein, P.C., 2020. Global chemical effects of the microbiome include new bile-acid conjugations. Nature 579, 123–129. https://doi.org/10.1038/s41586-020-2047-9

Rabinowitz, J.D., Enerbäck, S., 2020. Lactate: the ugly duckling of energy metabolism. Nat Metab 2, 566–571. https://doi.org/10.1038/s42255-020-0243-4

Reghupaty, S.C., Dall, N.R., Svensson, K.J., 2024. Hallmarks of the metabolic secretome. Trends in Endocrinology & Metabolism 35, 49–61. https://doi.org/10.1016/j.tem.2023.09.006

Rimal, S., Sang, J., Poudel, S., Thakur, D., Montell, C., Lee, Y., 2019. Mechanism of Acetic Acid Gustatory Repulsion in Drosophila. Cell Reports 26, 1432-1442.e4. https://doi.org/10.1016/j.celrep.2019.01.042

Sainz, F., Navarro, D., Mateo, E., Torija, M.J., Mas, A., 2016. Comparison of d-gluconic acid production in selected strains of acetic acid bacteria. International Journal of Food Microbiology 222, 40–47. https://doi.org/10.1016/j.ijfoodmicro.2016.01.015

Schönborn, J.W., Jehrke, L., Mettler-Altmann, T., Beller, M., 2019. FlySilico: Flux balance modeling of Drosophila larval growth and resource allocation. Scientific Reports 9, 17156. https://doi.org/10.1038/s41598-019-53532-4

Schwarzer, M., Gautam, U.K., Makki, K., Lambert, A., Brabec, T., Joly, A., Šrůtková, D., Poinsot, P., Novotná, T., Geoffroy, S., Courtin, P., Hermanová, P.P., Matos, R.C., Landry, J.J.M., Gérard, C., Bulteau, A.-L., Hudcovic, T., Kozáková, H., Filipp, D., Chapot-Chartier, M.-P., Šinkora, M., Peretti, N., Boneca, I.G., Chamaillard, M., Vidal, H., De Vadder, F., Leulier, F., 2023. Microbe-mediated intestinal NOD2 stimulation improves linear growth of undernourished infant mice. Science 379, 826–833. https://doi.org/10.1126/science.ade9767

Schwarzer, M., Makki, K., Storelli, G., Machuca-Gayet, I., Srutkova, D., Hermanova, P., Martino, M.E., Balmand, S., Hudcovic, T., Heddi, A., Rieusset, J., Kozakova, H., Vidal, H., Leulier, F., 2016. Lactobacillus plantarum strain maintains growth of infant mice during chronic undernutrition. Science 351, 854–857. https://doi.org/10.1126/science.aad8588

Sharma, A., Jusko, W.J., 1998. Characteristics of indirect pharmacodynamic models and applications to clinical drug responses. British Journal of Clinical Pharmacology 45, 229–239. https://doi.org/10.1046/j.1365-2125.1998.00676.x

Shin, S.C., Kim, S.-H., You, H., Kim, B., Kim, A.C., Lee, K.-A., Yoon, J.-H., Ryu, J.-H., Lee, W.-J., 2011. Drosophila Microbiome Modulates Host Developmental and Metabolic Homeostasis via Insulin Signaling. Science 334, 670–674. https://doi.org/10.1126/science.1212782

Sokolova, I.M., Frederich, M., Bagwe, R., Lannig, G., Sukhotin, A.A., 2012. Energy homeostasis as an integrative tool for assessing limits of environmental stress tolerance in aquatic invertebrates. Marine Environmental Research 79, 1–15. https://doi.org/10.1016/j.marenvres.2012.04.003

Speakman, J.R., Westerterp, K.R., 2012. A mathematical model of weight loss under total starvation: evidence against the thrifty-gene hypothesis. Disease Models & Mechanisms dmm.010009. https://doi.org/10.1242/dmm.010009

Stanley, M., Ghosh, B., Weiss, Z.F., Christiaanse, J., Gordon, M.D., 2021. Mechanisms of lactic acid gustatory attraction in Drosophila. Current Biology 31, 3525-3537.e6. https://doi.org/10.1016/j.cub.2021.06.005

Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J., Leulier, F., 2011. *Lactobacillus plantarum* Promotes *Drosophila* Systemic Growth by Modulating Hormonal Signals through TOR-Dependent Nutrient Sensing. Cell Metabolism 14, 403–414. https://doi.org/10.1016/j.cmet.2011.07.012

Storelli, G., Strigini, M., Grenier, T., Bozonnet, L., Schwarzer, M., Daniel, C., Matos, R., Leulier, F., 2018. *Drosophila* Perpetuates Nutritional Mutualism by Promoting the Fitness of Its Intestinal Symbiont *Lactobacillus plantarum*. Cell Metabolism 27, 362-377.e8. https://doi.org/10.1016/j.cmet.2017.11.011

Tennessen, J.M., Barry, W.E., Cox, J., Thummel, C.S., 2014. Methods for studying metabolism in *Drosophila*. Methods, Drosophila developmental biology methods 68, 105–115. https://doi.org/10.1016/j.ymeth.2014.02.034

Terashima, J., Bownes, M., 2004. Translating Available Food Into the Number of Eggs Laid by Drosophila melanogaster. Genetics 167, 1711–1719. https://doi.org/10.1534/genetics.103.024323

Turnbaugh, P.J., Gordon, J.I., 2009. The core gut microbiome, energy balance and obesity. The Journal of Physiology 587, 4153–4158. https://doi.org/10.1113/jphysiol.2009.174136

Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R., Gordon, J.I., 2006. An obesity-associated gut microbiome with increased capacity for energy harvest. Nature 444, 1027–1031. https://doi.org/10.1038/nature05414

Video: Fast Colony Forming Unit Counting in 96-Well Plate Format Applied to the Drosophila Microbiome [WWW Document], n.d. URL https://www.jove.com/v/64298/fast-colony-forming-unit-counting-96-well-plate-format-applied-to (accessed 5.9.25).

Von Frieling, J., Faisal, M.N., Sporn, F., Pfefferkorn, R., Nolte, S.S., Sommer, F., Rosenstiel, P., Roeder, T., 2020. A high-fat diet induces a microbiota-dependent increase in stem cell activity in the Drosophila intestine. PLOS Genetics 16, e1008789. https://doi.org/10.1371/journal.pgen.1008789

Walther, T.C., Chung, J., Jr, R.V.F., 2017. Lipid Droplet Biogenesis. Annual Review of Cell and Developmental Biology 33, 491–510. https://doi.org/10.1146/annurev-cellbio-100616-060608

Walther, T.C., Jr, R.V.F., 2012. Lipid Droplets and Cellular Lipid Metabolism. Annual Review of Biochemistry 81, 687–714. https://doi.org/10.1146/annurev-biochem-061009-102430

Warburg, O., 1956. On the Origin of Cancer Cells 123.

Wat, L.W., Chao, C., Bartlett, R., Buchanan, J.L., Millington, J.W., Chih, H.J., Chowdhury, Z.S., Biswas, P., Huang, V., Shin, L.J., Wang, L.C., Gauthier, M.-P.L., Barone, M.C., Montooth, K.L., Welte, M.A., Rideout, E.J., 2020. A role for triglyceride lipase brummer in the regulation of sex differences in Drosophila fat storage and breakdown. PLOS Biology 18, e3000595. https://doi.org/10.1371/journal.pbio.3000595

Wewer Albrechtsen, N.J., Holst, J.J., Cherrington, A.D., Finan, B., Gluud, L.L., Dean, E.D., Campbell, J.E., Bloom, S.R., Tan, T.M.-M., Knop, F.K., Müller, T.D., 2023. 100 years of glucagon and 100 more. Diabetologia 66, 1378–1394. https://doi.org/10.1007/s00125-023-05947-y

Wong, C.N.A., Ng, P., Douglas, A.E., 2011. Low-diversity bacterial community in the gut of the fruitfly Drosophila melanogaster. Environmental Microbiology 13, 1889–1900. https://doi.org/10.1111/j.1462-2920.2011.02511.x

1684   Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R., Goodman, A.L., 2019.
1685   Mapping human microbiome drug metabolism by gut bacteria and their genes.
1686   Nature 570, 462–467. https://doi.org/10.1038/s41586-019-1291-3
1687

Figure 1. Microbe-derived metabolites impact *Drosophila* starvation survival

Figure 2: Microbe-derived metabolites cause survival to deviate from simple energy balance models in starvation regimes with substantial energy limitation.

Figure 3: *Lp*-derived lactic acid is necessary and sufficient for starvation survival and acts via an interaction with glucose metabolism.

Figure 4. Lactic acid impacts starvation survival via regulation of oxidative phosphorylation (OXPHOS)

Figure S1: Microbial load decreases during starvation and composition shifts to increased frequency of the *Lactiplantibacillus* genus.

Figure S2. Microbe-derived metabolites do not impact starvation survival via differential food intake, active energy expenditure, or fecundity

Figure S3: Caloric value affects how spent media impact starvation survival.

Figure S4: Caloric value affects how lactic acid impacts fly survival.

Figure S5: Neither dietary lactate nor neutralized lactic acid extend fly survival.

Figure S6. Whole-body Ldh is dispensible for lactic acid mediated effects on survival

Figure S7. Metformin impacts survival in flies fed glucose and lactic acid

Figure S8. Transformed survival moments of CDM dilution series compared to energy balance model

Figure S9. Genome-scale metabolic model of *Drosophila* metabolism (iDrosophila) predicts a linear relationship between diet composition and overall metabolic flux