Chromosome-length genome assembly of *Uta stansburiana* and gene expression data

reveal fast pace-of-life comes with environmental stability

Sam R. Fellows[1]*, Raúl Araya-Donoso[2,3], Elizabeth Dávalos-Dehullu[3,4], Ruqayya Khan[5],

David Weisz[5], Olga Dudchenko[5,6], Erez Lieberman Aiden[5,6], Adrián Munguía-Vega[3,7,8], Greer

A. Dolby[3,9]*, Danielle L. Edwards[10]

[1] School of Natural Sciences, University of California Merced, California, USA

[2] School of Life Sciences, Arizona State University, Arizona, USA

[3] Baja Geogenomics Consortium, www.bajageogenomics.org

[4] Centro de Ciencias Genómicas, Programa de Biología de Sistemas, UNAM, Cuernavaca,

Morelos, Mexico

[5] The Center for Genome Architecture, Department of Molecular and Human Genetics,

Baylor College of Medicine, USA

[6] The Center for Theoretical Biological Physics, Rice University, USA

[7] School of Natural Resources and the Environment, University of Arizona, Arizona, USA

[8] Applied Genomics Lab, La Paz, Baja California Sur, Mexico

[9] Department of Biology, University of Alabama at Birmingham, Alabama, USA

[10] Research Institute for the Environment & Livelihoods, Charles Darwin University,

Northern Territory, Australia

Keywords: chromosome-level genome assembly, pace-of-life syndrome, *Uta stansburiana*,

Baja California, RNA-seq, seasonality

Summary: This study presents a high-quality reference genome paired with an RNA-seq

dataset collected to explore pace-of-life life history variation within the common side-

blotched lizard, *Uta stansburiana,* an emerging model species for studies of polymorphism,

adaptation, and reproduction. The authors paired long-read and Hi-C sequencing to generate

a chromosome-length genome assembly, annotated genes, and analyzed patterns of

differential gene expression between populations in different environments to reveal the

molecular basis for shifts in pace-of-life. These data and resources will provide a valuable

tool for evolutionary biologists, herpetologists, and geneticists interested in life history

evolution, vertebrate adaptation, and the maintenance of polymorphism.

Correspondence:

SRF, sam.r.fellows@gmail.com

GAD, gdolby@uab.edu

Abstract: *Uta stansburiana* are an emerging model system for studying sexual selection,

polymorphism, and the evolution of pace-of-life syndromes (POLS) whose distribution

covers variable environments and a wide latitudinal gradient. POLS are suites of traits

causing variation of life history along a slow maturing-fast maturing continuum. We present

a high-quality chromosome-level reference genome for *U. stansburiana* and pair it with

RNA-seq gene expression data to demonstrate, for the first time, the molecular basis for

pace-of-life differences between locations with higher and lower climate seasonality and

sexual size dimorphism (SSD). Our assembly is 2.1 Gbp, has scaffold N50 of 320 Mbp,

includes 104 scaffolds, and has an L50 of 3. The assembly comprises six macrochromosomes

and 11 microchromosomes. We annotated 20,350 genes for the assembly and found a repeat

element composition of 49.23%, similar to work in other phrynosomatid lizards. RNA-seq

data demonstrate differential expression in genes associated with pace-of-life divergence

including those related to stress, sexual reproduction, and cell proliferation/carcinogenesis

between distinctive environments. Our results provide the first differential gene expression

evidence of environmentally mediated pace-of-life processes related to different degrees of

SSD in *U. stansburiana* and demonstrate the utility of RNA-seq gene expression data in

detecting POLS.

1. <u>**Introduction**</u>

The pace-of-life syndrome (hereafter, POLS; Ricklefs and Wikelski 2002) hypothesis

invokes balancing selection to explain how life history variation is maintained within and among

populations (Royauté et al. 2018). The POLS hypothesis posits that life history, behavioral, and

physiological (especially metabolism, immunity, and stress responses) traits coevolve along a

"slow-fast" continuum which can create life history variation within or among populations

(Ricklefs and Wikelski 2002; Réale et al. 2007, 2010). Fast POLSs are often characterized by

fast metabolic rates, high stress responses, rapid development and maturation rates, bold

exploratory behaviors or high aggressiveness, high reproductive output, and are short-lived

(Réale et al. 2010; Dammhahn et al. 2018; Montiglio et al. 2018; but see also Royauté et al.

2018). Differences amongst POLS are likely exacerbated by variation in population density, e.g.,

lineages with high variation in population density tend to have the starkest differences in POLS

(Wright et al 2018). This suggests that the role of the environment in the formation and

maintenance of POLS is foundational (Dammhahn et al. 2018; Vasilieva 2022).

As genomic and gene expression data become more readily available, it becomes easier

to test and generate hypotheses regarding behavior (Hu et al. 2022), ecology (Xue et al. 2024),

niche (Mellor et al. 2025), and life history (Narum et al. 2018). Immonen et al (2018) reviewed

existing POLS hypothesis literature and identified diverse molecular pathways which, when

differentially expressed, may identify POLS differences including stress, immunity, metabolism,

life span, and sexual signals among others. To date, gene expression evidence for POLS has only

been identified in house mice (Prabh et al. 2023) where faster pace-of-life is associated with

genes related to metabolism, resulting in shorter generation time and more exploratory behavior.

However, on a molecular level stress brought on by a "fast" pace-of-life may shorten telomeres,

results in cellular damage due to oxidative stress, and often increases cancer risk (Giraudeau et

al. 2019; Ujvari et al. 2022). This provides a link to environmental conditions causing oxidative

stress, which can decrease telomere length or generally lead to decreased self-maintenance of

cell structures, apoptosis, higher risk of cancer, cell cycle arrest, or other senescent processes

(Young 2018) and presents an opportunity to use molecular tools to detect POLS. Despite this,

studies connecting the molecular mechanisms underlying POLS-related genes to such

phenotypes are rare. Linking key pace-of-life processes to gene expression differences in genes

associated with life history, behavior, physiology, and across environments using high-quality

reference genomes presents an unprecedented opportunity to detect evidence for the POLS

hypothesis in wild settings.

Common side-blotched lizards (*Uta stansburiana*, Baird and Girard 1852) are small,

wide-spread phrynosomatids native to the southwestern United States and northwestern Mexico,

whose range encompasses approximately 25 degrees latitude and 22 degrees longitude (Figure 1;

Stebbins 2003). Side-blotched lizards are small-bodied (up to 57 g and 60 mm snout-vent length)

and typically found amongst rocky scrub. They comprise a widespread species complex with

broad variation in morphological and life history characteristics across the range including

differences in sexual size dimorphism (SSD; Corl et al. 2012; Chelini et al. 2021), dorsal pattern

(McKinney 1971), genetically-determined mating tactics and associated throat color

polymorphism (Sinervo and Lively 1996; Sinervo et al. 2000; Corl et al. 2010), longevity (Zani

and Stein 2018; Smith et al. 2019), and reproductive investment (Alonzo and Sinervo 2001;

Smith et al. 2019). Such variation is highly indicative of pace-of-life processes operating as an

evolutionary mechanism within this species complex (Wright et al. 2019). Further, such patterns

are relevant to studies of sexual selection (Schlippe Justicia et al. 2024) and speciation (Corl et

al. 2012), so recovering the transcriptomic differences associated with these patterns will provide

a needed empirical assessment of POLS in an ideal, emerging model system.

Side-blotched lizards demonstrate strong life history trade-offs with respect to latitude

across their range. Individuals from the southern portions of the species distribution in the

southern United States and northern Mexico tend to be shorter-lived (up to two years, but usually

one or less; Wilson 1991; Smith et al. 2019) and lay many smaller eggs relative to longer-lived

individuals in the northern portions of the species range (i.e., up to 7 years in Oregon; Zani and

Stein 2018) which lay fewer, larger eggs (Smith et al 2019). This transition in longevity exists

alongside a range-wide shift from populations in the southern range with trimorphic throat colors

to populations in the northern distribution that are monomorphic orange-throated (Corl et al.

2010), mediated by differences in seasonality and annual precipitation (Chelini et al. 2021).

There have been suggestions that these dynamics are primarily underlain by sexual selection

dynamics and population densities given evolutionary correlations between SSD and the degree

of polymorphism (Corl et al. 2010b). However, recent evidence suggests that environments

independently generate variation in polymorphism and SSD, and the variation in the latter occurs

through sexually dimorphic growth rates in side-blotched lizards in alternate environmental

conditions (Chelini et al. 2021). Sexually dimorphic growth (and variation in its degree) that

correlates evolutionarily to alternate environmental conditions in different parts of the side-

blotched lizard range indicate alternate sex-specific optima in POLS, and such variation may

provide the mechanism connecting variation in SSD and polymorphism (Wright et al. 2019;

Arnqvist and Rowe 2023) by altering sexual selection dynamics to reinforce divergence among

populations in this system (Butlin and Smadja 2018). Furthermore, environmental conditions are

known to influence metabolic and immunity trait differences associated with alternate grown

rates in *U. stansburiana* (Smith et al. 2017) consistent with the POLS hypothesis (Dammhahn et

al. 2018).

We test for evidence of environmentally-mediated POLS differences in *U. stansburiana*

using a chromosome-level reference genome and annotation, paired with gene expression data

collected north and south of the Vizcaíno Desert, a major biogeographic structuring feature for

many species (Dolby et al. 2015), at the center of the Baja California peninsula. All side-blotched

lizard populations in this region are trimorphic for throat color, removing this as a confounding

variable, while environmental conditions such as temperature, precipitation seasonality, and

vegetation composition vary (Araya☐Donoso et al. 2024). Northern populations incur strong

seasonal variation in temperature between extremes, while the majority of rainfall occurs in the

winter. Southern populations encounter higher, more consistent temperatures through the year

with precipitation falling during the summer driven by the North American Monsoons (Figure

1C). These climatic differences lead to different environmental dynamics for *U. stansburiana*,

which is predominantly annual in Baja California. Northern environments are seasonally

disparate with relatively lower productivity given mismatches between lizard activity in the

spring/summer and rainfall, where southern environments are comparatively climatically stable

throughout the lifetime of *U. stansburiana* with precipitation coinciding with times when lizard

activity is at its peak. These differences in alignment between peak environmental productivity

and activity, and climatic consistency which influences growing season length likely impact life

history variation relevant to divergence in pace-of-life processes. For instance, environmental

seasonality is known to underlie variation in environmentally-mediated SSD and differences in

growth in side-blotched lizards (Chelini et al. 2021). Growing season differences between the

northern and southern Baja California peninsula areas we sampled may exacerbate POLS

differences (Cabezas-Cartes et al. 2018), and lead to variation in SSD among these regions

(Chelini et al. 2021).

In this study we present a high-quality chromosome-level reference genome for the

common side-blotched lizard (*Uta stansburiana*). This is the first genome assembly for the genus

Uta and the seventh chromosome-level published phrynosomatid genome, enabling

comprehensive comparative genomic analyses. The genome assembly is chromosome-scale (L90

= 8, N50 = 320Mb), has a diploid number 2n = 34 consistent with the hypothesized ancestral

karyotype of the family phrynosomatidae (Leaché and Sites, Jr. 2009), and reveals many

microchromosomes consistent with previous karyotyping work (Pennock et al. (1968, 1969); but

see Pinto et al. (2023)). We demonstrate the utility of this genome through analysis of RNA-seq

data collected from wild lizards in alternate environments. Our approach explores how life

history features (i.e., morphs and sexes) and environmental divergence (with impacts for

population density, SSD) associate with the molecular mechanisms that underlie pace-of-life

differences to develop hypotheses about the evolution of POLS within *U. stansburiana*. We

expect POLS divergence to present as differential expression in genes related to metabolism,

immune function, senescence, stress, regulatory/transcription pathways, and osmotic regulation.

We then test for transcriptomic evidence of POLS operating as a function of environmental

differences between northern and southern populations on the Baja California peninsula

(Hämäläinen et al. 2021; Dolby et al. 2022).

2. Materials and Methods

2.1 Sample collection: Genome

Fresh liver tissue was collected from an adult male *Uta stansburiana nevadensis*

collected by Dr. Pete Zani from Wrights Point, Harney County, OR, USA (permit ODFW024-

20; Fig. 1a). This site comprises a large population of *U. stansburiana nevadensis* living on a

lava flow surrounded by the Malheur wetlands and has been the subject of an 18-year mark-

recapture study. *U. stansburiana nevadensis* found at this site are known to live for up to seven

years (Zani and Stein 2018). The lizard was euthanized following IACUC protocol (University

of California Merced AUP21-0001) and liver tissue was immediately removed. Half of the tissue

was frozen and stored in 100% ethanol at -80° C, and the other half was used in immediate DNA

extraction. The remains of the individual were formalin-fixed and prepared for museum donation

(to be deposited at the Museum of Vertebrate Zoology, DLE3124).

Genomic DNA was extracted using a modified buffered lysis and detergent salting-out

protocol from Moore et al (2004) and Macdonald et al (2011). We assessed sample purity using a

NanoDrop 1.0 and quantified samples using the Qubit 2.0 high sensitivity kit. The extracted

DNA was sent to the University of California Davis DNA Technologies and Expression Analysis

core and sequenced using two Pacific Biosciences Sequel II SMRT-cells run in CCS mode. The

remaining frozen liver tissue was sent to DNA Zoo (dnazoo.org) for *in situ* Hi-C library

preparation and sequencing (Rao et al. 2014; Dudchenko et al. 2017, 2018).

2.2 Sample collection: RNA-seq

Adult *U. stansburiana elegans* were sampled by visual encounter survey and lassoing at

seven sites on the Baja California peninsula from northern and southern regions (Collecting

permit: 13439/19 DGVS Mexico, IACUC Protocol Arizona State University 20-1737R; IACUC

Protocol University of Arizona 20-627 PHS-CDC import permit 20220221-0651A; FWS import

2022CX2908360; Figure 1b) known to be genetically distinct lineages (Upton and Murphy

1997; Hollingsworth 1999). Sampling was undertaken at the same set of northern and southern

sites in the fall and spring during daylight hours. These sampling sites are in the ranges of two

reciprocally monophyletic groups located North and South of the center of the Peninsula (Upton

and Murphy 1997; Hollingsworth 1999; Araya-Donoso et al. 2024). Capture locations were

flagged with tape and morphological measurements were taken of the individuals, including

body size (snout-vent length, SVL; Table S1), and throat color photos. Tail tissue samples were

collected by autotomization and preserved in RNAlater on wet ice; tail tissue can be sampled

with minimal-to-no harm to individuals and is vascularized. As such, it is suitable for the scope

of these questions because of the wide array of RNAs found in even small reptilian blood

samples (Waits et al. 2020). A broader SVL dataset was used to calculate SSD as described in

Chelini et al. (2021) to show that northern sites (mean SSD = 1.04) were sexually monomorphic

in body size while southern sites were male-biased (mean SSD = 1.14) and these two regions

were significantly different in their degree of SSD (paired *t*-test; df = 2; *t*-stat = -9.04, p ≤ 0.05).

SVL data were collected from three populations in the northern and southern regions during the

spring breeding season only, using 2-8 individuals per sex, to estimate average adult male and

female SVL. We selected 33 individuals (14 male, 19 female) across seven sites (9 individuals

across three sites in the north, 24 individuals across four sites in the south) and roughly equal

throat colors (7OO, 4BB, 4YY, 4OB, 1BY, 2OY, 11 unassessable; Sinervo et al. 2001) for RNA

sequencing. The full metadata are available in Table S2. Preserved tail tips were sent to the Yale

Center for Genome Analysis for RNA isolation, poly-A mRNA library preparation, and

sequencing on the Illumina NovaSeq 6000 platform with 150 bp PE reads.

2.3 Bioinformatics:

2.3.1 Genome assembly

We prepared the draft genome assembly using hifiasm v0.16.1-r375 (Cheng et al. 2021)

and dropped k-mers that occurred more than 10 times the homozygous read coverage to improve

resolution in repetitive areas, but otherwise assembled the data under default parameters. Hi-C

reads were aligned to the hifiasm draft genome assembly using Juicer (Durand et al. 2016). The

alignments were then used to scaffold the draft to chromosome-length following Dudchenko et al

(2017, 2018) using the 3D-DNA pipeline and Juicebox Assembly Tools.

2.3.2 Genome annotation

We generated an annotation for the chromosome-length genome assembly of *U.*

stansburiana as follows. We renamed all HiC-scaffolds to simply "scaffold," and sorted the

chromosome-length scaffolds by decreasing length resulting in shuffling HiC-scaffold_14

through HiC-scaffold_17 (for a full list connecting HiC-scaffold names to the renamed scaffold

names, see Table S13). On the genome assembly, repeats were identified with RepeatModeler

v2.0.1 (Smit et al. 2015a) and soft-masked with RepeatMasker v4.1.1 (Smit et al. 2015b) Then,

we ran multiple iterations of Maker v3.01.03 (Campbell et al. 2014) to annotate the genome. A

first round of Maker was run to map and align evidence, which included the transcripts and

protein-coding sequences from the phrynosomatids *Urosaurus nigricaudus* (Davalos-Dehullu et

al. 2023) and *Phrynosoma platyrhinos* (Koochekian et al. 2022). Then, we ran two rounds of *ab*

initio gene model prediction using SNAP v2006-07-28 (Korf 2004) and Augustus v3.4.0 (Stanke

et al. 2006). After each round of Maker, we recorded the Annotation Edit Distance (AED) and

assessed the annotation completeness by using BUSCO v5.4.2 (Simão et al. 2015) on the

predicted transcripts obtained from Maker using the eukaryote and sauropsid gene datasets.

2.3.3 RNA-seq bioinformatics and differential expression analysis:

The RNA-seq data were assembled, quality-assessed, trimmed, aligned, and quantified

under the nf-core/rnaseq pipeline v3.14.0 (Patel et al. 2023), one of the nf-core workflows

(Ewels et al. 2020) which uses reproducible software from Bioconda (Grüning et al. 2018) and

BioContainers (Da Veiga Leprevost et al. 2017) projects. We ran this pipeline using Trim Galore

v0.6.7 (Krueger et al. 2021) to remove reads below 20 bp or with Q-values below 35, STAR

v2.7.9a (Dobin et al. 2013) to align reads to the reference genome, and quantified reads using

salmon v1.10.1 (Patro et al. 2017) under default parameters. All other processes in the pipeline

were run under default parameters.

We used the nf-core/rnaseq output files and tximport v1.32.0 (Soneson et al. 2015) to

prepare a matrix of non-normalized read counts for DESeq2 v1.44.0 (Love et al. 2014). We

conducted two different differential expression models: 1) a group-means parameterization of the

binary variables region (north and south), sex (male and female), and season (spring and fall)

with the categories north, female, and fall set as reference levels for contrasts, and 2) a group-

means parameterization of putatively homozygous throat colors (Sinervo et al. 2000, 2001)

against the other homozygous throat colors without respect to other variables. We extracted

contrasts for region (controlling for season and sex), sex (controlling for sex and region), orange

throats (controlling for blue and yellow throats), blue throats (controlling for orange and yellow

throats), and yellow throats (controlling for orange and blue throats). We calculated log2-fold

changes between significantly differentially expressed genes (Bonferroni-corrected $p < 0.025$

because we tested two differential expression models).

2.4 Gene Ontology term analyses and functional enrichment:

We used a two-pronged approach to determine which biological functions were

associated with high levels of differential expression between northern and southern populations,

controlling for season. First, we used g:Profiler v.e111_eg58_p18_f463989d (Raudvere et al.

2019) on the full list of differentially-expressed genes to determine major enriched biological

processes, KEGG pathways, and transcription factors (Tables S3-S7). For this analysis, we used

Mus musculus as reference because of its completeness in the database. We then visualized

enriched biological processes via semantic clustering as a treemap using REVIGO v.1.8.1

(Supek et al. 2011; Fig. S6, S7).

To supplement these results, we used STRING Interaction Database (Szklarczyk et al.

2023) to generate a network to visualize known interactions among the final gene list. Because of

the completeness of its functional annotation, we used human as the reference, and we used

medium confidence (0.4) as the minimum threshold. We then analyzed the gene networks in

Cytoscape (Shannon et al. 2003) to identify subnetworks and "hub" genes with high degrees of

connectivity.

2.5 Synteny evolution in Phrynosomatidae

We investigated synteny of protein-coding genes in *U. stansburiana* by comparative

analysis with other available phrynosomatid chromosome-level assemblies. We compared *U.*

stansburiana against *Phrynosoma platyrhinos* (Koochekian et al. 2022), *Sceloporus undulatus*

(Westfall et al. 2021), *Urosaurus nigricaudus* (Davalos-Dehullu et al. 2023) and the dactyloid

lizard *Anolis carolinensis* (Sampson et al. 2025; Gambón-Deza 2023) using GENESPACE

v1.3.1 (Lovell et al. 2022). Because the *A. carolinensis* assembly did not explicitly include

microchromosomes, we considered synteny across all scaffolds in all assemblies if they

contained syntenic blocks with a minimum of five hits.

3. Results

3.1 Genome assembly and annotation

PacBio HiFi long read sequencing resulted in 55.9 Gb of consensus sequences

comprising 204 contigs. The contig N50 was 93 Mb, and the longest contig was 170 Mb. The Hi-

C guided genome assembly for *U. stansburiana* (Table 1) was 2.10 Gb total length spanning 17

chromosome-length scaffolds (N50: 320 Mb, L90: 8) and high completeness (BUSCO

eukaryotic completeness of 99.6%). The chromosome-length scaffolds in the final DNA Zoo

assembly were arranged in descending order except for scaffold_14 (for associated HiC_scaffold

numbers, see Table S13) likely representing the sex chromosome based on a pseudoautosomal-

compatible region. The following scaffold, scaffold_18, is potentially a Y chromosome for the

same reason. Further analysis is required to definitively identify these as sex chromosomes.

Consistent with previous karyotype analysis (Pennock et al. 1968), the final assembly comprises

six macrochromosomes ranging from 426 Mb to 146 Mb and 11 microchromosomes ranging

from 38 Mb to 16 Mb (Fig. S8, S9). Roughly 50% of the genome corresponded to repeat

elements, comparable to the other phrynosomatids (Table 2). A total of 20,350 protein-coding

genes were annotated with an average gene length of 20,561 bp, corresponding to a 92.6%

completeness of all eukaryote BUSCO (either complete or fragmented; Table 2). Of the 49.29%

of the genome that comprises repeat elements (Table 3), the most abundant were unclassified

(31.17%) followed by LINEs (15.44%). Over 2/3 of the LINEs (10.26% of the genome)

belonged to the L2/CR1/Rex families (Table 3). RepeatModeler did not detect any SINEs, which

may account for the relatively high (31.71%) proportion of unclassified repeats.

3.2 Synteny evolution in phrynosomatid lizards

In comparing genomic synteny in *U. stansburiana* to other phrynosomatids, we found a

high level of synteny conservation in the six macrochromosomes with some short, inverted

regions (Figure 2). We identified several rearrangements in the microchromosomes, especially

scaffold_8, scaffold_12, scaffold_13, scaffold_15, scaffold_16, and scaffold_17. The likely X

chromosome, scaffold_14 showed substantial conserved synteny with other identified X

chromosomes (scaffold 14 in *Urosaurus nigricaudus*, scaffold 10 in *Sceloporus undulatus*,

scaffold 9 in *Phrynosoma platyrhinos*, and the X chromosome in *Anolis carolinensis*), again

suggesting that this scaffold represents the X chromosome in *U. stansburiana.*

3.3 RNA-seq bioinformatics

RNA sequencing resulted in 2.4 billion 101 bp paired-end reads, ranging from 24.2

million to 46.8 million raw reads per sample. 73.5% of raw reads had Q-scores of 35 or higher,

while 92.8% of raw reads had Q-scores of 30 or higher. Raw read GC content ranged from 40-

45% per sample. After trimming and filtering, average sequence length was 92 bp, ranged from

20-100 bp, and resulted in 2.35 billion paired-end reads. On average, we retained 35.6 million

paired-end reads per sample on average ranging from 24.2 million to 46.8 million raw reads per

sample. 74.9% of read-pairs uniquely mapped to the genome and 2.7% were multiply mapped;

only uniquely mapped transcripts were retained. The RNA-seq data comprised reads mapping to

19,970 genes (i.e., those with non-zero counts) in total, but averaged 14,830.5 genes per sample

and ranged from 12,823 to 16,317 genes per sample.

3.4 Differential expression by sex

DESeq2 analysis yielded 9 genes differentially expressed (Bonferroni-corrected adjusted

p-value <0.025, Table S3) between female (reference level) and male lizards controlling for

region and season, all of which had functional gene annotations. Moderated log2-fold change

values for gene annotations ranged from 1.50 (*gstm4*) to -1.40 (*shfl*). G:Profiler returned two

enriched biological process GO terms (Table S8): *long-chain fatty acid metabolic process* ($p =$

1.51×10^{-2}), and *xenobiotic catabolic process* ($p = 2.19 \times 10^{-2}$). Six enriched KEGG terms were

returned: *drug metabolism cytochrome P450* ($p = 2.92 \times 10^{-2}$), *glutathione metabolism* ($p = 3.00$

$\times 10^{-2}$), *metabolism of xenobiotics by cytochrome P450* ($p = 3.17 \times 10^{-2}$), *platinum drug*

resistance ($p = 3.70 \times 10^{-2}$), *chemical carcinogenesis-DNA adducts* ($p = 4.08 \times 10^{-2}$), and *drug*

metabolism- other enzymes ($p = 4.69 \times 10^{-2}$). The STRING interaction network contained only a

single edge, between *gstm4* and *gstm1*, did not contain significantly more interactions than

expected (p = 0.332), and presented one additional enriched biological process (*nitrobenzene*

metabolic process, FDR = 0.04) and one additional protein domain and feature (*glutathione s-*

transferase, Mu class, FDR = 0.03).

3.5 Differential expression by throat color

DESeq2 analysis yielded one statistically significantly differentially expressed gene in

orange-throated lizards compared to blue- or yellow-throated lizards (Bonferroni-corrected

adjusted p-value 0.025), which was associated with an annotation (*slc12a7*; Table S4); this had a

large moderated log2-fold change value of 7.21. g:Profiler returned a single enriched KEGG

term, *collecting duct acid secretion* ($p = 0.05$). Because there was a single enriched gene, further

exploration of GO terms was not pursued. Comparison between blue throated lizards and yellow-

and orange-throated lizards found nine differentially expressed genes, all of which had functional

annotations. Log2-fold change values for annotated genes ranged from -4.80 (*phpt*) to -6.73 x 10^{-7}

^{7} (*adam20*). g:Profiler did not return any biological process terms, but found one enriched KEGG

term (Table S10): *GnRH signaling pathway* ($p = 4.68$ x 10^{-2}). The STRING network did not

contain any edges, and did not yield any significant enrichments. Finally, the comparison

between yellow-throated and blue- and orange-throated lizards yielded two differentially

expressed genes both associated with annotations. Log2-fold change values ranged from 2.78

(*plcd1*) to 3.88 (*cd74*). g:Profiler returned one enriched biological process term: *positive*

regulation of small molecule metabolic process ($p = 4.98$ x 10^{-2}) and no enriched KEGG

processes.

3.6 Differential expression by region, controlling for sex and season

DESeq2 analysis revealed 163 annotated genes differentially expressed (Bonferroni-

corrected adjusted p-value < 0.025, Figure 3) between southern and northern lizards accounting

for sex and season. Log2-fold change values for annotated genes ranged from 11.43 (*adam20*) to

-9.40 (*taar1*) (Figure 4, Table S7). g:Profiler results returned two statistically enriched biological

processes: *regulation of response to stimulus* ($p = 7.51$ x 10^{-3}) and *regulation of multicellular*

organism process ($p = 2.90$ x 10^{-2}) (Table S12, Fig. S6). There were four enriched KEGG terms:

endocrine resistance ($p = 6.03$ x 10^{-3}), *parathyroid hormone synthesis, secretion, and action* ($p =$

1.38 x 10^{-3}), *chemical carcinogenesis- receptor activation* ($p = 2.47$ x 10^{-2}), and *hepatocellular*

carcinoma ($p = 2.88$ x 10^{-2}). The STRING interaction network had 153 nodes, an average

clustering coefficient of 0.35, and the network did not have significantly more interactions than

expected by chance ($p = 0.09$). Network analysis yielded four hub genes with ≥ 10 connected

edges: *egfr* (18), *ccnd1* (14), *fos* (14), and *kras* (10).

3.6.1 Endocrine pathways

Two of the four enriched KEGG pathways in the data by region were related to endocrine

signaling, specifically *endocrine resistance* (hsa01522), *parathyroid synthesis, secretion, action*

(hsa04928). Eight enriched genes were associated with this pathway in the STRING interaction

network: *prkcb* (log2-fold change 7.93), *adcy7* (log2-fold change 6.15), *egr1* (log2-fold change

3.22), *egfr* (log2-fold change 1.19), *fos* (log2-fold change -1.59), and *casr* (log2-fold change -

1.67). Endocrine resistance pathways (hsa01522) affect cell cycle and apoptosis; seven enriched

genes are associated with this pathway in the STRING interaction network: *adcy7* (log2-fold

change 6.15), *egfr* (log2-fold change 1.19), *dll1* (log2-fold change 1.17), *kras* (log2-fold change -

0.65), *ccnd1* (log2-fold change -0.097), *fos* (log2-fold change -1.59), and *ncor1* (log2-fold

change -1.88); g:Profiler did not identify *kras* as enriched for this pathway. Endocrine resistance

and secretion are components of metabolism, cell proliferation, and sexual maturation which are

predicted to occur with POLS differences.

3.6.2 Proto-oncogenes and cancer pathways

One of the four enriched KEGG pathways (*hepatocellular carcinoma*) and a major

keyword (proto-oncogene) in the analysis between regions were related to cancer or proto-

oncogenes. The hepatocellular carcinoma KEGG pathway (hsa05225) describes several key

signaling processes (genetic and epigenetic) that occur throughout liver cancer related to

chromatin remodeling and oxidative stress. Nine enriched genes are associated with this

pathway: *prkcb* (log2-fold change 9.60), *prkca* (log2-fold change 1.16), *smarcd3* (log2-fold

change 1.66), *egfr* (log2-fold change 1.34), *kras* (log2-fold change -0.71), *ccnd1* (log2-fold

change -1.07), *gsto1* (log2-fold change -1.22), *mgst1* (log2-fold change -1.25), and *dvl1* (log2-

fold change -2.18). Chemical carcinogenesis-receptor activation (hsa05207) was identified as an

enriched KEGG pathway in g:Profiler and describes cell signaling pathways which may control

the pace at which cells replicate. g:Profiler identified eight enriched genes associated with this

pathway: *fos* (log2-fold change -1.59), *gsto1* (log2-fold change -1.22), *mgst1* (log2-fold change -

1.25), *ccnd1* (log2-fold change -1.07), *dll1* (log2-fold change 1.17), *egfr* (log2-fold change 1.34),

adcy7 (log2-fold change 6.15), and *prkcb* (log2-fold change 9.60).

4. Discussion

Pace-of-life traits may be influenced by environmental processes that are likely to impact

sexual selection dynamics, yet little is known about the functional molecular mechanisms giving

rise to such variation within or between species. Side-blotched lizards are an emerging model

system exhibiting variation in several key traits that are known to predict POLS divergence;

variation in SSD, longevity, polymorphism, population density, sexual signaling, immune

response and growth, and other traits are all known to vary among populations experiencing

alternate environmental conditions. Our study provides a high-quality annotated reference

genome for *Uta stansburiana* as a resource for future experimental and wild population studies

in the processes which underlie the variation and adaptability across this widely distributed

species and a template for studying the molecular mechanisms underlying POLS variation in

other species. Our results support POLS divergence being mostly associated with differences in

environmental stability across the lifetime and identify brain development and function, cell

signaling and circadian rhythm, fertility, immunity, homeostasis, metal ion binding, and

metabolism as key mechanisms underlying POLS variation.

4.1 Genome features and comparative genomics

This is the first genome assembly for the genus *Uta*, and the seventh chromosome-level published phrynosomatid genome, enabling comprehensive comparative genomic analyses. The genome assembly is chromosome-scale (L90 = 8, N50 = 320 Mb), and reveals a large number of microchromosomes (first described as such by (Pennock et al. (1968, 1969); but see Pinto et al. (2023)) implying substantial chromosome fission or fusion events relative to closely related lizards. This is consistent with the hypothesized ancestral karyotype of 2n = 34 for phrynosomatidae (Leaché and Sites, Jr. 2009). We identified the 17 longest scaffolds as likely chromosomes of *U. stansburiana nevadensis*. These comprise six macrochromosomes and 11 microchromosomes with a diploid count of 2n = 34. This is consistent with previous karyotype analysis in *Uta sp.* (Pennock et al. 1968) and the hypothesized ancestral karyotype of all phrynosomatids (2n = 34; Leaché and Sites, Jr. 2009). Further analysis is needed to confirm the identity of the sex chromosomes, but preliminary observations of pseudoautosomal regions suggest that scaffold_14 is the X chromosome, consistent with synteny analysis, and that scaffold_18 is the Y.

At the macrochromosome level, the *U. stansburiana* genome shows high synteny and small regions of inversions relative to the other phrynosomatid genomes, consistent with previous findings in Iguania (Davalos-Dehullu et al. 2023). The microchromosomes also exhibit highly conserved syntenic blocks between these lizards, however, they also show clear evidence of substantial chromosome fission/fusion among phrynosomatids (Figure 4). In particular, scaffold_8 and scaffold_12 contain syntenic blocks that have undergone substantial fission/fusion. Finally, our hypothesized X chromosome (scaffold_14) was highly syntenic with

identified X chromosomes in other phrynosomatids, including the closely related *Urosaurus*

nigricaudus.

In general, microchromosomes contain conserved syntenic blocks of slowly evolving

genes while undergoing dynamic fission/fusion (Srikulnath et al. 2021), and the consequences of

these microchromosomes and their fission or fusion depend on their content. Increasing evidence

shows that microchromosomes could house genes under selection (Schield et al. 2019) and that

the syntenic lability provided by microchromosomes could allow for flexibility in local

adaptation via unique linkage disequilibrium regimes (Perry et al. 2021). This could provide

substantial capacity for local adaptation in side-blotched lizards, consistent with their large range

and highly disparate abiotic environments. However, previous karyotyping work in the genus

Uta suggests that this may not be a hard rule; Pennock et al. (1968, 1969) found the same diploid

number of chromosomes in all *Uta* species analyzed including insular taxa which presumably

have very distinct local adaptations and recombination regimes. The consequences of

microchromosome fission/fusion in side-blotched lizards relative to other phrynosomatid lizards

will depend on the genic content of the microchromosomes.

4.2 Sexual differentiation in gene expression and POLS

While there are limited reasons to expect that male and female lizards would experience

alternate POLS, many studies have detected large sexual differences in gene expression in other

taxa (e.g., Lipshutz et al. 2025). Our gene expression data allowed us to assess the relative

contribution of sex, throat color polymorphism, and environmental divergence among regions

relative to predictions under the POLS hypothesis. Results demonstrate minimal differential

expression between male and female *U. stansburiana*. Among differentially expressed genes, the

main enriched processes deal with glutathione metabolism with associated genes having

moderated log2-fold change values between 1.04 (*gstm1*) and 1.50 (*gstm4*). Glutathione

production protects cells from stress (Maher 2005) which relates to sexually-specific POLS, and

its production is increased in males indicating that male lizards are under higher stress. Our

results indicate that there may be limited mechanisms for sex-specific POLS in this species. It is

noteworthy to mention that other iguanian reptiles (*Anolis carolinensis,* Marin et al.

2017; *Sceloporus malachitichus,* Lisachov et al. 2021; the closely-related *Urosaurus*

nigricaudus, Davalos-Dehullu et al. 2023) show evidence of dosage compensation systems,

where males overexpress their single copy of the X chromosome to achieve expression levels of

female XX. If present in side-blotched lizards, this could contribute to the low number of sex-

biased expression observed. This also could be because genes expressed differentially by sex

could vary among tissue type (Oliva et al. 2020) and we did not sample gonad tissue (Xu et al.

2020). However, *U. stansburiana* are predominantly sexually monomorphic for size across their

range (Chelini et al. 2021) even while both sexes are polymorphic in throat color, suggesting

limited sex differences. Finally, our tail samples were vascularized and included blood which

allowed for detection of hormonal and reproductive gene expression differences in other

comparisons.

4.3 Differential gene expression, polymorphism, and POLS

There have been some suggestions that alternate reproductive tactics may lead to

divergence in POLS between morphs in polymorphic species (e.g., Friesen et al. 2017). Contrasts

among throat colors, indicative of alternative reproductive tactics, returned few DEGs, few

enriched GO terms, and no STRING interactions. The only DEG in orange-throated lizards was

slc12a7, which produces a solute carrier protein maintaining renal homeostasis in the presence of

potassium and chloride ions in humans. Relative to blue-throated lizards, orange- and yellow-

throated lizards over-expressed genes related to gonadotropin releasing hormone pathways,

however, moderated log2-fold change values for these DEGs were near zero. Blue-throated

lizards did not over-express any genes relative to the other morphs, and the two DEGs with high

moderated log2-fold change values (*phpt1*, a gene producing proteins involved in histidine

phosphorylation, and *ppp4r3b*, which is involved in carbohydrate metabolism and DNA repair).

Finally, yellow-throated lizards over-expressed genes involved in small molecule metabolic

processes; *cd74* is a component of the major histocompatibility complex (MHC) and plays a role

in regulation of prostaglandins (regulators of inflammation and fever response), while *plcd1* is

involved in inositol triphosphate synthesis which mobilizes calcium ions from cellular stores.

Such few DEGs among color morphs could be explained by the fact that we did not

sample tissue which might reveal differences in epidermal pigment or structural color. Previous

studies have suggested that there may be difference among morphs within *U. stansburiana*

(Hazard et al. 2019) and other polymorphic lizards (Friesen et al. 2017) in physiological traits

associated with POLS variation. We find limited evidence of that, since only blue and yellow

morphs show variation in one or two molecular pathways associated with metabolism, aging,

and/or immunity. We note that the throat color analysis did not include the correction for region,

season, or sex, meaning that while this should be a morph-specific expression signal, it could be

driven by uneven sample sizes across other variables or small sample sizes within variables.

4.4 Environmental influences on gene expression and POLS

Our analyses of gene expression among environmentally divergent northern and southern

regions revealed environmental stability across the lifetime experienced by populations as a

predominant force in differential expression. We find significant differences in POLS-related

gene pathways associated with brain development and cognition, temporal regulation, signaling,

homeostasis, genome stability, sperm production, immune function, and ion binding. Southern

populations overexpressed (log2-fold change > 2) eight genes related to brain development and

function. These genes were associated with neuron maturation (*fbxo41*), aggregation (*sspo*),

synapse assembly (*iqsec2*), synapse function (*slc9a6*, *chrdl1*, *fbxo41*), and post-synaptic

organization (*arhgap39*). Genes related to dopamine receptor binding (*clic6*) and GABA-based

neurotransmitter signaling (*gabra5*) implicated in learning and behavioral fear responses were

also enriched. In contrast, northern populations over-expressed (log2-fold change < -2) two

related genes. These genes were related to neurotransmitter response and release (*kcnh7*) or

reception (*taar1*) rather than neurogenesis.

Side-blotched lizards in populations with low predation and high sexual signaling

intensity tend to have larger ventral posterior amygdalae (LaDage et al. 2022) and increased

aggression and spatial processing needs, including in more complex spatial settings. This results

in increased neurogenesis (LaDage et al. 2013, 2016). Our data indicate that southern region

populations show male-biased SSD which is consistent with faster male growth and longer

growing seasons (Chelini et al. 2021) which could lead to increased reproductive rates. Side-

blotched lizard populations with increased reproductive rates often have increased rates of

predation (Parker and Pianka 1975). Over-expression of genes related to neurogenesis in the

southern populations could be an indication that southern lizards are signaling at higher rates in

spite of predation risk consistent with a faster POLS and is also consistent with observations that

most adult male lizards are removed from populations in southern portions of the species' range

by predators within their first breeding (Edwards D, pers. obs.).

Despite evidence for increased neurogenesis in side-blotched lizards, there exists mixed

evidence for cognitive performance as a component of POLS. Goulet et al. (2018) found that

skinks with fast POLS were more exploratory but slower learners, while De Meester et al. (2022)

found no link between cognitive ability and ecology in lacertids. This could be because increased

neurogenesis in side-blotched lizards is related to social behavior rather than memory and

learning (e.g., Mettke☐Hofmann 2014). This is supported by the over-expression of *gabra5* in

southern populations. *gabra5* is both implicated in elevated stress responses and decreased

learning in knockout studies and decreased short-term memory when up-regulated with drug

treatment in mice (Tan et al. 2011; Syding et al. 2023). Over-expression of genes related to

neurogenesis in southern populations is consistent with a fast POLS and male-biased SSD, but

these results should be validated with careful assessment of population density, sexual signaling

intensity, and cognitive ability in these populations.

Southern region populations of *U. stansburiana* showed higher expression (moderated lfc

> 2) of genes related to transcriptional regulation (*egr1*, *prkcb*, and *pprc1*), genome repair (*egr1*,

ppp4r3b, and *hmbox1*), and cellular and hormonal signaling especially in homeostasis (*prkcb*,

pprc1, *aoc1*, and *glp2r*). Many of these genes have multiple roles in these functions and may be

thematically linked by upregulation of *egr1*. *Egr1* is a multi-functional gene which regulates the

biosynthesis of luteinizing hormone and mediates metabolism, reproduction, cell cycle regulation

and DNA damage repair, and several core circadian clock genes (*bmal1*, *nr1d1*, and *per2*).

Network analysis did not identify it as a major hub gene given our specified threshold of node

degree > 10, but it was the sixth most connected DEG (seven connections). In northern

populations, we found over-expression of four genes related to metabolism and homeostasis

(*aox3*, *slc2a9*, *faah*, and *cdipt*). Transcriptional regulation, genome stability, and homeostasis are

foundationally associated with a slower pace of life. These patterns suggest that differential

expression of these genes in the south could emphasize regulatory complexity and flexibility,

consistent with a faster POLS, while northern populations prioritize metabolic stability and

homeostasis, consistent with a slower pace-of-life.

Lucas and French (2012) demonstrated POLS shifts in *U. stansburiana* when

environmental or urban stressors led to increased stress, decreased life maintenance processes,

and increased reproductive output. We identified two enriched KEGG pathways related to

endocrine processes and two related to cell cycle regulation associated with differential

expression between southern and northern regions. Parathyroid synthesis and activation increases

serum calcium and decreases bone density in other iguanid lizards (e.g., *Dipsosaurus dorsalis*

and *Sceloporus grammicus*; McWhinnie and Cortelyou 1968). Four of the six genes related to

this pathway were over-expressed in southern populations (*egfr*, *egr1*, *adcy7*, and *prkcb*).

Parathyroid hormone (PTH) synthesis, secretion, and action (hsa04928) is responsible for

regulating calcium and phosphorus ions. In cases of restricted dietary calcium, parathyroid

hormone will stimulate bone turnover and prevent loss of phosphate ions in the kidney. In

iguanian lizards, injection of PTH causes acute hypercalcemia and sometimes

hyperphosphatemia (McWhinnie and Cortelyou 1968) and may elevate in response to metabolic

or nutritional challenges and stress. This may reflect a deficit in resource allocation or increase in

some stressor in southern populations resulting in quicker use of energy reserves. For example,

predation rates might be higher in the south (but see Wilson 1991), higher population density or

long periods between typical hurricanes (Parker 1974) could limit resources and increase stress

(e.g., Malisch et al. 2020), or simply longer active seasons could result in higher generational

turnover rates. Elevated reproduction, high competition for resources, and increased stress are all

concordant with a faster POLS.

Cell signaling and proliferation, and cancer as a consequence, are functional components

of growth rates and aging which vary among POLS (Ujvari et al. 2022). The KEGG term

endocrine resistance refers to cell cycle resistance to hormonal signaling, typically through

cancer research in humans (i.e., Ribeiro and Freiman 2014). Three of the six genes related to this

pathway were over-expressed in the south (*egfr*, *dll1*, and *adcy7*). Mixed expression of endocrine

resistance genes is somewhat non-specific and may be due to many different processes such as

decreased need for endocrine resistance, efficient cell proliferation, etc. Understanding which of

these processes is responsible for generating this pattern will require detailed understanding of

signaling pathways resulting in cell proliferation (or lack thereof), and consequences for POLS

will depend on the effects of the DEGs.

Differences in longevity may also be associated with anticancer strategies (Tissier et al.

2022, Ujvari et al. 2022). The two enriched KEGG pathways related to cell cycle regulation were

hepatocellular carcinoma and *chemical carcinogenesis- receptor activation*. Like *endocrine*

resistance, these KEGG terms are primarily known through cancer research in humans. Three of

the seven genes involved in the *hepatocellular carcinoma* pathway were over-expressed in the

south (*egfr*, *prkcb*, and *smarcd3*). Four genes have functions specific to cell proliferation or

carcinogenesis- *smarcd3* is a tumor suppressor gene, *gsto1* and *mgst1* are involved in antioxidant

response, and *dvl1* is a cell proliferation regulator. The fact that a tumor suppressor gene is over-

expressed in the south while antioxidant and cell proliferation control genes are under-expressed

suggests that growth may be continuous and rapid in southern populations and is consistent with

other results pointing toward environmental variation influencing differences in POLS between

the regions.

The *chemical carcinogenesis- receptor activation* pathway had four of eight genes over-

expressed in the south (*egfr*, *dll1*, *adcy7*, and *prkcb*). This KEGG pathway describes several cell

receptor pathways which respond to chemical carcinogens to result in proliferation,

chemoresistance, and apoptosis. GSTO1 and MGST1 are also involved in this pathway. The role

of differential expression of genes related to cell proliferation and cancer pathways in *U.*

stansburiana is likely dependent on the function of the individual genes implicated. Specifically,

differences in biotic or abiotic conditions likely create alternate selection for POLS, echoing

previous evidence for environmental drivers of POLS (e.g., Williams et al. 2010; Hämäläinen et

al. 2021).

If, as we predict, southern region populations have higher densities and longer growing

and activity seasons, POLS variation in reproductive rates could present as differential

expression in molecular control of sexual function and fertility. Four genes over-expressed

(moderated lfc > 2) in the southern population were directly related to sperm production or male

fertility. *Adam20*, *topaz1*, and *piwil1* are directly involved in sperm maturation and

spermatogenesis. *Adam20* is the single-most differentially expressed gene in the dataset

(moderated log2 fold-change = 11.43). *Piwil1* which binds and suppresses TE-derived piwi

interfering RNAs (piRNAs) to maintain germline integrity and is associated with infertility of

certain crosses in canids (Stalker et al. 2016); it is thought to be a highly-conserved gene among

amniotes (Lim et al. 2013). Elevated expression of a piRNA gene and genes related to

spermatogenesis, especially those often differentially expressed in the testis, suggests that

southern lizards are over-producing sperm relative to northern males even when controlling for

seasonal effects. Over-production of sperm in southern populations after controlling for season

and regardless of throat color suggests that life history trade-offs are related to environmental

conditions rather than alternative mating strategies, and possibly related to longer mating seasons

in the south. Direct measurement of sperm production across the range of the species would

better resolve this issue.

Northern populations over-expressed two calcium-binding genes, *wdr49* and *calr*, and

one magnesium-binding gene, *ttll11*. *Calr* may be involved in oocyte maturation and the

prevention of polyspermy, possibly implicating increased investment in offspring rather than

total reproductive output. Southern populations expressed four calcium-binding genes, *pamr1*,

zbtb8os, *cpne4*, and *cdhr4*, one magnesium-binding gene, *adcy7*, and one zinc-binding gene,

otud7b. While these genes generally have diverse functions, the greater number of calcium ion

binding genes expressed in southern lizards suggest that they have elevated stress hormone levels

(Park et al. 2017) which can dampen immunocompetence (Hudson et al. 2020) and require

physiological compensation. Taken together, these results suggest that northern region

populations invest more in reproduction within a limited season, while southern region

populations operate under higher levels of stress because of a faster POLS.

Side-blotched lizard populations with higher parasite load tend to have higher

immunocompetence (Spence et al. 2017), and ectoparasite load was much higher in the south (G.

Dolby, pers. comm.). Side-blotched lizard populations which more heavily invest in reproduction

tend to heal wounds faster and have higher immunocompetence at the expense of body condition

during the reproductive season (Smith et al. 2019), a sign of a fast POLS. Both northern and

southern region populations differentially expressed genes related to immune function and ion

binding, but the connections between these categories and among the individual genes was

idiosyncratic. Immune genes expressed higher in the north are *zdhhc11* and *clec5a*, while

southern region populations over-expressed *adcy7*, *otud7b*, *tmem45b*, and *msmp*. In general,

zdhhc11 and *clec5a* are components of the innate immune system involved in viral infections,

while genes over-expressed in the south have a more general immune function. This may reflect

that southern lizards are exposed to a greater diversity of pathogens (Clark and Bradford 1969;

Quillfeldt et al. 2018), consistent with our hypothesis that side-blotched lizards in the south may

be at higher density. These results suggest that northern region lizards are subject to fewer

diseases while southern region lizards require a wider variety of immune responses to pathogens

possibly due to higher prevalence of disease and a higher population density because of a faster

POLS.

5. Conclusions

The molecular mechanisms controlling variation in POLS are relatively understudied. We

used the high-quality chromosome-level reference genome presented here alongside gene

expression data to test whether populations living under different environmental conditions vary

with respect to POLS for the first time in a wild reptile. In a system which demonstrates range-

wide shifts in longevity, polymorphic throat color and reproductive tactics, we find few genes

related to sex- or throat color-based expression differences and limited to no evidence that sex or

ARTs contribute to POLS differences. Our results indicate that environmental variation is the

predominant force resulting in POLS variation. We show that expression differences across a

latitudinal environment gradient are largely related to homeostasis in the face of resource stress,

sexual function, and cell proliferation control. This points to variation in environments as

mediating growth and reproduction opportunities, thereby affecting population dynamics and

driving the evolution of longevity which all influence POLS variation in this emerging model

system. We hypothesize that this, in turn, may drive sexual selection dynamics and therefore

variation in polymorphism to reinforce population divergence (Butlin and Smadja 2018). Future

research should consider *U. stansburiana* to be a model system for studying the evolution of

POLS, especially for understanding the relationships environmental variation and differences in

alternative reproductive tactics, growth, reproduction, aging, senescence, telomere length, cancer

suppression, and resource allocation.

6. Acknowledgments

664   We thank S. Baty and A. Biddy for help with bioinformatics and analysis, D. Ardell for

665 insight into modeling, P. Zani for providing the specimen from which we generated the genome,

666 B. Carlson and N. Whutthituntisil for feedback on the manuscript, O. Nguyen and L. Froenicke

667 for help with long-read sequencing, K. Kusumi and D. Denardo for helpful discussions, and C.

668 Pedraza-Marrón and M. Dawson for their advice and support.

7. Data Availability Statement

Raw Hi-C data are available on SRA (BioProject PRJNA512907, BioSample

SAMN46443096). An interactive Hi-C contact map for the final assembly is available at

https://www.dnazoo.org/post/a-top-notch-blotch,

https://www.dnazoo.org/assemblies/uta_stansburiana. The assembled and annotated genome will

be available on NCBI, submission number SUB15340363. The raw long-read and RNA-seq data

are available on SRA (long-read data are under submission number SUB15364625, and RNA-

seq data are under submission number SUB15372950; both are under BioProject

PRJNA1268585).

8. Funding

This project was funded by NSF EAR #2305608 to G.A.D.; NSF EAR # 1925771 to

A.M.V.; NSF DBI- 2021795, NSF PHY-2210291, and NIH CEGS RM1HG011016-01A1 to

E.L.A., R.A.D. was supported by the doctoral fellowship 72200094 (ANID, Chile); E.D-D. was

supported from CONAHCYT fellowship 162588; D.L.E. and S.R.F. acknowledge funding

support from the University of California, Merced. The Hi-C guided assembly was done in

association with the DNA Zoo consortium (dnazoo.org), which acknowledges support from

Illumina, IBM, and Pawsey Supercomputing Center.

9. Contributions

SRF, DLE, AMV & GAD conceptualized and designed the study; SRF, RAD, EDD,

AMV & OD collected tissues and data; SRF, RAD, RK & DW undertook analyses on collected

data; SRF, DLE, AMV & GAD interpreted results; DLE, AMV & GAD provided resources

necessary to execute the study; DLE, AMV, GAD & ELA acquired funding for the study; DLE

& GAD supervised student researchers and oversaw administration for the work, SRF & DLE

wrote the paper; All authors reviewed, edited and approved the paper.

10. References

Alonzo S.H., Sinervo B. 2001. Mate choice games, context-dependent good genes, and genetic
cycles in the side-blotched lizard, Uta stansburiana. Behavioral Ecology and
Sociobiology. 49:176–186.

Araya☐Donoso R., Biddy A., Munguía☐Vega A., Lira☐Noriega A., Dolby G.A. 2024. Habitat
quality or quantity? Niche marginality across 21 plants and animals suggests differential
responses between highland and lowland species to past climatic changes. Ecography.
2024:e07391.

Arnqvist G., Rowe L. 2023. Ecology, the pace☐of☐life, epistatic selection and the maintenance
of genetic variation in life☐history genes. Molecular Ecology. 32:4713–4724.

Baird S.F., Girard C. 1852. Characteristics of some new reptiles in the Museum of the
Smithsonian Institution. Proc. Acad. Nat. Sci. Philadelphia. 6:68–70.

Butlin R.K., Smadja C.M. 2018. Coupling, Reinforcement, and Speciation. The American
Naturalist. 191:155–172.

Cabezas-Cartes F., Mariela Boretto J., Ruth Ibargüengoytía N. 2018. Effects of Climate and
Latitude on Age at Maturity and Longevity of Lizards Studied by Skeletochronology.
Integrative and Comparative Biology.

Campbell M.S., Holt C., Moore B., Yandell M. 2014. Genome Annotation and Curation Using
MAKER and MAKER□P. CP in Bioinformatics. 48.

Chelini M.C., Brock K., Yeager J., Edwards D.L. 2021. Environmental drivers of sexual
dimorphism in a lizard with alternative mating strategies. J of Evolutionary Biology.
34:1241–1255.

Cheng H., Concepcion G.T., Feng X., Zhang H., Li H. 2021. Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. Nat Methods. 18:170–175.

Clark G.W., Bradford J. 1969. Blood Parasites of Some Reptiles of the Pacific Northwest. The
Journal of Protozoology. 16:578–581.

Corl A., Davis A.R., Kuchta S.R., Sinervo B. 2010. Selective loss of polymorphic mating types is
associated with rapid phenotypic evolution during morphic speciation. Proceedings of the
National Academy of Sciences. 107:4254–4259.

Corl A., Lancaster L.T., Sinervo B. 2012. Rapid Formation of Reproductive Isolation between
Two Populations of Side-Blotched Lizards, Uta stansburiana. Copeia. 2012:593–602.

Da Veiga Leprevost F., Grüning B.A., Alves Aflitos S., Röst H.L., Uszkoreit J., Barsnes H.,
Vaudel M., Moreno P., Gatto L., Weber J., Bai M., Jimenez R.C., Sachsenberg T.,
Pfeuffer J., Vera Alvarez R., Griss J., Nesvizhskii A.I., Perez-Riverol Y. 2017.
BioContainers: an open-source and community-driven framework for software
standardization. Bioinformatics. 33:2580–2582.

Dammhahn M., Dingemanse N.J., Niemelä P.T., Réale D. 2018. Pace-of-life syndromes: a
framework for the adaptive integration of behaviour, physiology and life history. Behav
Ecol Sociobiol. 72:62, s00265-018-2473-y.

Davalos-Dehullu E., Baty S.M., Fisher R.N., Scott P.A., Dolby G.A., Munguia-Vega A., Cortez
D. 2023. Chromosome-Level Genome Assembly of the Blacktail Brush Lizard,
Urosaurus nigricaudus , Reveals Dosage Compensation in an Endemic Lizard. Genome
Biology and Evolution. 15:evad210.

De Meester G., Van Linden L., Torfs J., Pafilis P., Šunje E., Steenssens D., Zulčić T., Sassalos A.,
Van Damme R. 2022. Learning with lacertids: Studying the link between ecology and
cognition within a comparative framework. Evolution. 76:2531–2552.

Dobin A., Davis C.A., Schlesinger F., Drenkow J., Zaleski C., Jha S., Batut P., Chaisson M.,
Gingeras T.R. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29:15–
21.

Dolby G.A., Bennett S.E.K., Dorsey R.J., Stokes M.F., Riddle B.R., Lira-Noriega A., Munguia-
Vega A., Wilder B.T. 2022. Integrating Earth–life systems: a geogenomic approach.
Trends in Ecology & Evolution. 37:371–384.

Dolby G.A., Bennett S.E.K., Lira-Noriega A., Wilder B.T., Munguía-Vega A. 2015. Assessing
the Geological and Climatic Forcing of Biodiversity and Evolution Surrounding the Gulf
of California. jsw. 57:391–455.

Dudchenko O., Batra S.S., Omer A.D., Nyquist S.K., Hoeger M., Durand N.C., Shamim M.S.,
Machol I., Lander E.S., Aiden A.P., Aiden E.L. 2017. De novo assembly of the *Aedes*
aegypti genome using Hi-C yields chromosome-length scaffolds. Science. 356:92–95.

Dudchenko O., Pham M., Lui C., Batra S.S., Hoeger M., Nyquist S.K., Durand N.C., Shamim
M.S., Machol I., Erskine W., Aiden E.L., Kaur P. 2018. Hi-C yields chromosome-length
scaffolds for a legume genome, *Trifolium subterraneum*. .

Durand N.C., Shamim M.S., Machol I., Rao S.S.P., Huntley M.H., Lander E.S., Aiden E.L. 2016.
Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments.
Cell Systems. 3:95–98.

Ewels P.A., Peltzer A., Fillinger S., Patel H., Alneberg J., Wilm A., Garcia M.U., Di Tommaso P.,
Nahnsen S. 2020. The nf-core framework for community-curated bioinformatics
pipelines. Nat Biotechnol. 38:271–278.

Fick S.E., Hijmans R.J. 2017. WorldClim 2: new 1 km spatial resolution climate surfaces for
global land areas. Int. J. Climatol. 37:4302–4315.

Friesen C.R., Johansson R., Olsson M. 2017. Morph specific metabolic rate and the timing of
reproductive senescence in a color polymorphic dragon. J Exp Zool Pt A. 327:433–443.

Gambón-Deza F. 2023. A new T Cell Receptor in Squamata Reptiles. bioRxiv.

Giraudeau M., Angelier F., Sepp T. 2019. Do Telomeres Influence Pace of Life Strategies in
Response to Environmental Conditions Over a Lifetime and Between Generations?
BioEssays. 41:1800162.

Goulet C.T., Michelangeli M., Chung M., Riley J.L., Wong B.B.M., Thompson M.B., Chapple
D.G. 2018. Evaluating cognition and thermal physiology as components of the pace-of-
life syndrome. Evol Ecol. 32:469–488.

Grüning B., Dale R., Sjödin A., Chapman B.A., Rowe J., Tomkins-Tinch C.H., Valieris R.,
Köster J. 2018. Bioconda: sustainable and comprehensive software distribution for the
life sciences. Nat Methods. 15:475–476.

Hämäläinen A.M., Guenther A., Patrick S.C., Schuett W. 2021. Environmental effects on the
covariation among pace of life traits. Ethology. 127:32–44.

Hazard L.C., Nagy K.A., Miles D.B., Svensson E.I., Costa D., Sinervo B. 2019. Integration of
Genotype, Physiological Performance, and Survival in a Lizard (*Uta stansburiana*) with
Alternative Mating Strategies. Physiological and Biochemical Zoology. 92:303–315.

Hollingsworth B.D. 1999. The Molecular Systematics of the Side-blotched Lizards (IGUANIA:
PHRYNOSOMATIDAE: UTA). .

Hu C.K., York R.A., Metz H.C., Bedford N.L., Fraser H.B., Hoekstra H.E. 2022. cis-Regulatory
changes in locomotor genes are associated with the evolution of burrowing behavior. Cell
Reports. 38:110360.

Hudson S.B., Lidgard A.D., French S.S. 2020. Glucocorticoids, energy metabolites, and
immunity vary across allostatic states for plateau side☐blotched lizards (*Uta*
stansburiana uniformis) residing in a heterogeneous thermal environment. J Exp Zool Pt
A. 333:732–743.

Immonen E., Hämäläinen A., Schuett W., Tarka M. 2018. Evolution of sex-specific pace-of-life
syndromes: genetic architecture and physiological mechanisms. Behav Ecol Sociobiol.
72:60.

Koochekian N., Ascanio A., Farleigh K., Card D.C., Schield D.R., Castoe T.A., Jezkova T. 2022.
A chromosome-level genome assembly and annotation of the desert horned lizard,
Phrynosoma platyrhinos , provides insight into chromosomal rearrangements among
reptiles. GigaScience. 11:giab098.

Korf I. 2004. Gene finding in novel genomes. BMC Bioinformatics. 5:59.

Krueger F., James F., Ewels P., Afyounian E., Schuster-Boeckler B. 2021. Trim Galore. .

LaDage L.D., Maged R.M., Forney M.V., Roth T.C., Sinervo B., Pravosudov V.V. 2013.
Interaction between territoriality, spatial environment, and hippocampal neurogenesis in
male side-blotched lizards. Behavioral Neuroscience. 127:555–565.

LaDage L.D., Roth T.C., Sinervo B., Pravosudov V.V. 2016. Environmental experiences
influence cortical volume in territorial and nonterritorial side-blotched lizards, Uta
stansburiana. Animal Behaviour. 115:11–18.

LaDage L.D., Yu T., Zani P.A. 2022. Higher Rate of Male Sexual Displays Correlates with
Larger Ventral Posterior Amygdala Volume and Neuron Soma Volume in Wild-Caught
Common Side-Blotched Lizards, **Uta stansburiana**. Brain Behav Evol. 97:298–308.

Leaché A.D., Sites, Jr. J.W. 2009. Chromosome Evolution and Diversification in North American
Spiny Lizards (Genus *Sceloporus*). Cytogenet Genome Res. 127:166–181.

Lim S.L., Tsend-Ayush E., Kortschak R.D., Jacob R., Ricciardelli C., Oehler M.K., Grützner F.
2013. Conservation and Expression of PIWI-Interacting RNA Pathway Genes in Male
and Female Adult Gonad of Amniotes1. Biology of Reproduction. 89.

811 Lipshutz S.E., Hibbins M.S., Bentz A.B., Buechlein A.M., Empson T.A., George E.M., Hauber
812    M.E., Rusch D.B., Schelsky W.M., Thomas Q.K., Torneo S.J., Turner A.M., Wolf S.E.,
813    Woodruff M.J., Hahn M.W., Rosvall K.A. 2025. Repeated behavioural evolution is
814    associated with convergence of gene expression in cavity-nesting songbirds. Nat Ecol
815    Evol. 9:845–856.

816 Lisachov A.P., Tishakova K.V., Romanenko S.A., Molodtseva A.S., Prokopov D.Yu., Pereira
817    J.C., Ferguson-Smith M.A., Borodin P.M., Trifonov V.A. 2021. Whole-chromosome
818    fusions in the karyotype evolution of *Sceloporus* (Iguania, Reptilia) are more frequent in
819    sex chromosomes than autosomes. Phil. Trans. R. Soc. B. 376:20200099.

820 Love M.I., Huber W., Anders S. 2014. Moderated estimation of fold change and dispersion for
821    RNA-seq data with DESeq2. Genome Biol. 15:550.

822 Lovell J.T., Sreedasyam A., Schranz M.E., Wilson M., Carlson J.W., Harkess A., Emms D.,
823    Goodstein D.M., Schmutz J. 2022. GENESPACE tracks regions of interest and gene copy
824    number variation across multiple genomes. eLife. 11:e78526.

825 MacDonald A.J., Sarre S.D., FitzSimmons N.N., Aitken N. 2011. Determining microsatellite
826    genotyping reliability and mutation detection ability: an approach using small-pool PCR
827    from sperm DNA. Mol Genet Genomics. 285:1–18.

828 Maher P. 2005. The effects of stress and aging on glutathione metabolism. Ageing Research
829    Reviews. 4:288–314.

830 Malisch J.L., Garland T., Claggett L., Stevenson L., Kohl E.A., John-Alder H.B. 2020. Living on
831    the edge: Glucocorticoid physiology in desert iguanas (Dipsosaurus dorsalis) is predicted
832    by distance from an anthropogenic disturbance, body condition, and population density.
833    General and Comparative Endocrinology. 294:113468.

834 Marin R., Cortez D., Lamanna F., Pradeepa M.M., Leushkin E., Julien P., Liechti A., Halbert J.,
835    Brüning T., Mössinger K., Trefzer T., Conrad C., Kerver H.N., Wade J., Tschopp P.,
836    Kaessmann H. 2017. Convergent origination of a *Drosophila* -like dosage compensation
837    mechanism in a reptile lineage. Genome Res. 27:1974–1987.

838 McKinney C.O. 1971. An Analysis of Zones of Intergradation in the Side-Blotched Lizard, Uta
839    stansburiana (Sauria: Iguanidae). Copeia. 1971:596.

840 McWhinnie D.J., Cortelyou J.R. 1968. Influence of parathyroid extract on blood and urine
841    mineral levels in iguanid lizards. General and Comparative Endocrinology. 11:78–87.

842 Mellor N.J., Webster T.H., Byrne H., Williams A.S., Edwards T., DeNardo D.F., Wilson M.A.,
843    Kusumi K., Dolby G.A. 2025. Divergence in Regulatory Regions and Gene Duplications
844    May Underlie Chronobiological Adaptation in Desert Tortoises. Molecular Ecology.
845    34:e17600.

846 Mettke☐Hofmann C. 2014. Cognitive ecology: ecological factors, life☐styles, and cognition.
847    WIRES Cognitive Science. 5:345–360.

848   Montiglio P.-O., Dammhahn M., Dubuc Messier G., Réale D. 2018. The pace-of-life syndrome
849   revisited: the role of ecological conditions and natural history on the slow-fast
850   continuum. Behav Ecol Sociobiol. 72:116.

851   Moore E., Arnscheidt A., Ger K. 2004. Simplified protocols for the preparation of genomic DNA
852   from bacterial cultures. Molecular Microbial Ecology Manual. Kluwer Academic
853   Publishers. p. 3–18.

854   Narum S.R., Di Genova A., Micheletti S.J., Maass A. 2018. Genomic variation underlying
855   complex life-history traits revealed by genome sequencing in Chinook salmon. Proc. R.
856   Soc. B. 285:20180935.

857   Oliva M., Muñoz-Aguirre M., Kim-Hellmuth S., Wucher V., Gewirtz A.D.H., Cotter D.J.,
858   Parsana P., Kasela S., Balliu B., Viñuela A., Castel S.E., Mohammadi P., Aguet F., Zou Y.,
859   Khramtsova E.A., Skol A.D., Garrido-Martín D., Reverter F., Brown A., Evans P.,
860   Gamazon E.R., Payne A., Bonazzola R., Barbeira A.N., Hamel A.R., Martinez-Perez A.,
861   Soria J.M., GTEx Consortium, Pierce B.L., Stephens M., Eskin E., Dermitzakis E.T.,
862   Segrè A.V., Im H.K., Engelhardt B.E., Ardlie K.G., Montgomery S.B., Battle A.J.,
863   Lappalainen T., Guigó R., Stranger B.E., Aguet F., Anand S., Ardlie K.G., Gabriel S.,
864   Getz G.A., Graubert A., Hadley K., Handsaker R.E., Huang K.H., Kashin S., Li X.,
865   MacArthur D.G., Meier S.R., Nedzel J.L., Nguyen D.T., Segrè A.V., Todres E., Balliu B.,
866   Barbeira A.N., Battle A., Bonazzola R., Brown A., Brown C.D., Castel S.E., Conrad D.F.,
867   Cotter D.J., Cox N., Das S., De Goede O.M., Dermitzakis E.T., Einson J., Engelhardt
868   B.E., Eskin E., Eulalio T.Y., Ferraro N.M., Flynn E.D., Fresard L., Gamazon E.R.,
869   Garrido-Martín D., Gay N.R., Gloudemans M.J., Guigó R., Hame A.R., He Y., Hoffman
870   P.J., Hormozdiari F., Hou L., Im H.K., Jo B., Kasela S., Kellis M., Kim-Hellmuth S.,
871   Kwong A., Lappalainen T., Li X., Liang Y., Mangul S., Mohammadi P., Montgomery
872   S.B., Muñoz-Aguirre M., Nachun D.C., Nobel A.B., Oliva M., Park Y., Park Y., Parsana
873   P., Rao A.S., Reverter F., Rouhana J.M., Sabatti C., Saha A., Stephens M., Stranger B.E.,
874   Strober B.J., Teran N.A., Viñuela A., Wang G., Wen X., Wright F., Wucher V., Zou Y.,
875   Ferreira P.G., Li G., Melé M., Yeger-Lotem E., Barcus M.E., Bradbury D., Krubit T.,
876   McLean J.A., Qi L., Robinson K., Roche N.V., Smith A.M., Sobin L., Tabor D.E., Undale
877   A., Bridge J., Brigham L.E., Foster B.A., Gillard B.M., Hasz R., Hunter M., Johns C.,
878   Johnson M., Karasik E., Kopen G., Leinweber W.F., McDonald A., Moser M.T., Myer K.,
879   Ramsey K.D., Roe B., Shad S., Thomas J.A., Walters G., Washington M., Wheeler J.,
880   Jewell S.D., Rohrer D.C., Valley D.R., Davis D.A., Mash D.C., Branton P.A., Barker
881   L.K., Gardiner H.M., Mosavel M., Siminoff L.A., Flicek P., Haeussler M., Juettemann T.,
882   Kent W.J., Lee C.M., Powell C.C., Rosenbloom K.R., Ruffier M., Sheppard D., Taylor
883   K., Trevanion S.J., Zerbino D.R., Abell N.S., Akey J., Chen L., Demanelis K., Doherty
884   J.A., Feinberg A.P., Hansen K.D., Hickey P.F., Jasmine F., Jiang L., Kaul R., Kibriya
885   M.G., Li J.B., Li Q., Lin S., Linder S.E., Pierce B.L., Rizzardi L.F., Skol A.D., Smith
886   K.S., Snyder M., Stamatoyannopoulos J., Tang H., Wang M., Carithers L.J., Guan P.,
887   Koester S.E., Little A.R., Moore H.M., Nierras C.R., Rao A.K., Vaught J.B., Volpi S.
888   2020. The impact of sex on gene expression across human tissues. Science.
889   369:eaba3066.

Park W., Rengaraj D., Kil D.-Y., Kim H., Lee H.-K., Song K.-D. 2017. RNA-seq analysis of the
kidneys of broiler chickens fed diets containing different concentrations of calcium. Sci
Rep. 7:11740.

Parker W.S. 1974. Home Range, Growth, and Population Density of Uta stansburiana in Arizona.
Journal of Herpetology. 8:135.

Parker W.S., Pianka E.R. 1975. Comparative Ecology of Populations of the Lizard Uta
stansburiana. Copeia. 1975:615.

Patel H., Ewels P., Peltzer A., Botvinnik O., Sturm G., Moreno D., Vemuri P., Garcia M.U.,
silviamorins, Pantano L., Binzer-Panchal M., bot nf-core, Syme R., Zepper M., Kelly G.,
Hanssen F., Yates J.A.F., Cheshire C., rfenouil, Espinosa-Carrasco J., marchoeppner,
Miller E., Talbot A., Zhou P., Guinchard S., Hörtenhuber M., Gabernet G., Mertes C.,
Straub D., Tommaso P.D. 2023. nf-core/rnaseq: nf-core/rnaseq v3.12.0 - Osmium
Octopus. .

Patro R., Duggal G., Love M.I., Irizarry R.A., Kingsford C. 2017. Salmon provides fast and bias-
aware quantification of transcript expression. Nat Methods. 14:417–419.

Pennock L.A., Tinkle, D W, Shaw, M W. 1969. Minute Y chromosome in the lizard genus Uta
(family Iguanidae). Cytogenetics. 8:9–19.

Pennock LewisA., Tinkle DonaldW., Shaw MargeryW. 1968. Chromosome number in the lizard
genus Uta (family Iguanidae). Chromosoma. 24.

Perry B.W., Schield D.R., Adams R.H., Castoe T.A. 2021. Microchromosomes Exhibit Distinct
Features of Vertebrate Chromosome Structure and Function with Underappreciated
Ramifications for Genome Evolution. Molecular Biology and Evolution. 38:904–910.

Pinto B.J., Gamble T., Smith C.H., Wilson M.A. 2023. A lizard is never late: Squamate genomics
as a recent catalyst for understanding sex chromosome and microchromosome evolution.
Journal of Heredity. 114:445–458.

Prabh N., Linnenbrink M., Jovicic M., Guenther A. 2023. Fast adjustment of pace of life and
risk taking to changes in food quality by altered gene expression in house mice. Ecology
Letters. 26:99–110.

Quillfeldt P., Romeike T., Masello J.F., Reiner G., Willems H., Bedolla-Guzmán Y. 2018.
Molecular survey of coccidian infections of the side-blotched lizard *Uta stansburiana* on
San Benito Oeste Island, Mexico. Parasite. 25:43.

Rao S.S.P., Huntley M.H., Durand N.C., Stamenova E.K., Bochkov I.D., Robinson J.T., Sanborn
A.L., Machol I., Omer A.D., Lander E.S., Aiden E.L. 2014. A 3D Map of the Human
Genome at Kilobase Resolution Reveals Principles of Chromatin Looping. Cell.
159:1665–1680.

Raudvere U., Kolberg L., Kuzmin I., Arak T., Adler P., Peterson H., Vilo J. 2019. g:Profiler: a
web server for functional enrichment analysis and conversions of gene lists (2019
update). Nucleic Acids Research. 47:W191–W198.

Réale D., Garant D., Humphries M.M., Bergeron P., Careau V., Montiglio P.-O. 2010. Personality
and the emergence of the pace-of-life syndrome concept at the population level. Phil.
Trans. R. Soc. B. 365:4051–4063.

Réale D., Reader S.M., Sol D., McDougall P.T., Dingemanse N.J. 2007. Integrating animal
temperament within ecology and evolution. Biological Reviews. 82:291–318.

Ribeiro J.R., Freiman R.N. 2014. Estrogen signaling crosstalk: Implications for endocrine
resistance in ovarian cancer. The Journal of Steroid Biochemistry and Molecular Biology.
143:160–173.

Ricklefs R.E., Wikelski M. 2002. The physiology/life-history nexus. Trends in Ecology &
Evolution. 17:462–468.

Royauté R., Berdal M.A., Garrison C.R., Dochtermann N.A. 2018. Paceless life? A meta-
analysis of the pace-of-life syndrome hypothesis. Behav Ecol Sociobiol. 72:64.

Sampson J.M., Morrissey K.A., Mikolajova K.J., Zimmerly K.M., Gemmell N.J., Gardner M.G.,
Bertozzi T., Miller R.D. 2025. Squamate reptiles may have compensated for the lack of
γδTCR with a duplication of the TRB locus. Front. Immunol. 15:1524471.

Schield D.R., Card D.C., Hales N.R., Perry B.W., Pasquesi G.M., Blackmon H., Adams R.H.,
Corbin A.B., Smith C.F., Ramesh B., Demuth J.P., Betrán E., Tollis M., Meik J.M.,
Mackessy S.P., Castoe T.A. 2019. The origins and evolution of chromosomes, dosage
compensation, and mechanisms underlying venom regulation in snakes. Genome Res.
29:590–601.

Schlippe Justicia L., Mayer M., Lorioux-Chevalier U., Dittrich C., Rojas B., Chouteau M. 2024.
Intraspecific divergence of sexual size dimorphism and reproductive strategies in a
polytypic poison frog. Evol Ecol. 38:121–139.

Shannon P., Markiel A., Ozier O., Baliga N.S., Wang J.T., Ramage D., Amin N., Schwikowski
B., Ideker T. 2003. Cytoscape: A Software Environment for Integrated Models of
Biomolecular Interaction Networks. Genome Res. 13:2498–2504.

Simão F.A., Waterhouse R.M., Ioannidis P., Kriventseva E.V., Zdobnov E.M. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 31:3210–3212.

Sinervo B., Bleay C., Adamopoulou C. 2001. SOCIAL CAUSES OF CORRELATIONAL
SELECTION AND THE RESOLUTION OF A HERITABLE THROAT COLOR
POLYMORPHISM IN A LIZARD. Evolution. 55:2040–2052.

Sinervo B., Lively C.M. 1996. The rock–paper–scissors game and the evolution of alternative
male strategies. Nature. 380:240–243.

Sinervo B., Svensson E., Comendant T. 2000. Density cycles and an offspring quantity and
quality game driven by natural selection. Nature. 406:985–988.

Smit A.F.A., Hubley R., Green P. 2015a. RepeatModeler. .

Smit A.F.A., Hubley R., Green P. 2015b. RepeatMasker. .

Smith G.D., Zani P.A., French S.S. 2019. Life☐history differences across latitude in common
side☐blotched lizards (*Uta stansburiana*). Ecology and Evolution. 9:5743–5751.

Soneson C., Love M.I., Robinson M.D. 2015. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. F1000Res. 4:1521.

Spence A.R., Durso A.M., Smith G.D., Skinner H.M., French S.S. 2017. Physiological Correlates
of Multiple Parasitic Infections in Side-Blotched Lizards. Physiological and Biochemical
Zoology. 90:321–327.

Srikulnath K., Ahmad S.F., Singchat W., Panthum T. 2021. Why Do Some Vertebrates Have
Microchromosomes? Cells. 10:2182.

Stalker L., Russell S.J., Co C., Foster R.A., LaMarre J. 2016. PIWIL1 Is Expressed in the Canine
Testis, Increases with Sexual Maturity, and Binds Small RNAs1. Biology of
Reproduction. 94.

Stanke M., Keller O., Gunduz I., Hayes A., Waack S., Morgenstern B. 2006. AUGUSTUS: ab
initio prediction of alternative transcripts. Nucleic Acids Research. 34:W435–W439.

Stebbins R.C. 2003. A Field Guide to Western Reptiles and Amphibians. Boston: Houghton
Mifflin Company.

Supek F., Bošnjak M., Škunca N., Šmuc T. 2011. REVIGO Summarizes and Visualizes Long
Lists of Gene Ontology Terms. PLoS ONE. 6:e21800.

Syding L.A., Kubik-Zahorodna A., Reguera D.P., Nickl P., Hruskova B., Kralikova M.,
Kopkanova J., Novosadova V., Kasparek P., Prochazka J., Rozman J., Turecek R.,
Sedlacek R. 2023. Ablation of Gabra5 Influences Corticosterone Levels and Anxiety-like
Behavior in Mice. Genes. 14:285.

Szklarczyk D., Kirsch R., Koutrouli M., Nastou K., Mehryary F., Hachilif R., Gable A.L., Fang
T., Doncheva N.T., Pyysalo S., Bork P., Jensen L.J., von Mering C. 2023. The STRING
database in 2023: protein–protein association networks and functional enrichment
analyses for any sequenced genome of interest. Nucleic Acids Research. 51:D638–D646.

Tan S., Rudd J.A., Yew D.T. 2011. Gene Expression Changes in GABAA Receptors and
Cognition Following Chronic Ketamine Administration in Mice. PLoS ONE. 6:e21328.

Ujvari B., Raven N., Madsen T., Klaassen M., Dujon A.M., Schultz A.G., Nunney L., Lemaître
J., Giraudeau M., Thomas F. 2022. Telomeres, the loop tying cancer to organismal
life□histories. Molecular Ecology. 31:6273–6285.

Upton D.E., Murphy R.W. 1997. Phylogeny of the Side-Blotched Lizards (Phrynosomatidae:Uta)
Based on mtDNA Sequences: Support for a Midpeninsular Seaway in Baja California.
Molecular Phylogenetics and Evolution. 8:104–113.

Vasilieva N.A. 2022. Pace-of-Life Syndrome (POLS): Evolution of the Concept. Biol Bull Russ
Acad Sci. 49:750–762.

Waits D.S., Simpson D.Y., Sparkman A.M., Bronikowski A.M., Schwartz T.S. 2020. The utility
of reptile blood transcriptomes in molecular ecology. Molecular Ecology Resources.
20:308–317.

Williams J.B., Miller R.A., Harper J.M., Wiersma P. 2010. Functional Linkages for the Pace of
Life, Life-history, and Environment in Birds. Integrative and Comparative Biology.
50:855–868.

Wilson B.S. 1991. Latitudinal Variation in Activity Season Mortality Rates of the Lizard Uta
Stansburiana. Ecological Monographs. 61:393–414.

Wright J., Bolstad G.H., Araya□Ajoy Y.G., Dingemanse N.J. 2019. Life□history evolution
under fluctuating density□dependent selection and the adaptive alignment of
pace□of□life syndromes. Biological Reviews. 94:230–247.

Xu C., Dolby G.A., Drake K.K., Esque T.C., Kusumi K. 2020. Immune and sex-biased gene
expression in the threatened Mojave desert tortoise, Gopherus agassizii. PLoS ONE.
15:e0238202.

Xue X., Thompson A.R., Adams B.J. 2024. An Antarctic worm and its soil ecosystem: A review
of an emerging research program in ecological genomics. Applied Soil Ecology.
193:105110.

Young A.J. 2018. The role of telomeres in the mechanisms and evolution of life-history trade-
offs and ageing. Phil. Trans. R. Soc. B. 373:20160452.

Zani P.A., Stein S.J. 2018. Field and laboratory responses to drought by Common Side-blotched
Lizards (Uta stansburiana). Journal of Arid Environments. 154:15–23.

1023

1024

1025

1026

1027 **Table 1.** Genome assembly statistics for *U. stansburiana* with three other phrynosomatid

1028 reference genomes for comparison.

	Species			
	Uta stansburiana	*Urosaurus nigricaudus*	*Sceloporus undulatus*	*Phrynosoma platyrhinos*
Genome version	UCM_Usta_1	ASU_Unig_1	SceUnd_v1.1	MUOH_PhPlat_1.1
Assembly length (Gbp)	2.1	1.87	1.91	1.9
N50 (Mbp)	320	279	276	273
L50 (n°)	3	3	3	3
Largest scaffold (bp)	426,646,591	380,405,096		
Number scaffolds	104	327		
Eukaryote BUSCO Assembly (%)	99.6	99.2	99.6	98.8

1029

1030 **Table 2.** Genome annotation statistics for *U. stansburiana* with three other phrynosomatid

1031 reference genomes for comparison.

	Species			
	Uta stansburiana	*Urosaurus nigricaudus*	*Sceloporus undulatus*	*Phrynosoma platyrhinos*
Genome version	UCM_Usta_1	ASU_Unig_1	SceUnd_v1.1	MUOH_PhPlat_1.1
% Repeats	49.23	45.42	41.33	44.45
N° genes	20,350	17,902	15,472	17,864
Average gene length (bp)	20,560.80	18,191.50	22,998.8	21,769.8
Eukaryote BUSCO Annotation (%)	C + F: 92.6	C + F: 94.9	C + F: 96.1	C + F: 92.5
Reference	Dudchenko et al., 2017; Dudchenko et al., 2018; This study	Dávalos-Dehullu et al., 2023	Westfall et al. 2021	Koochekian et al., 2022

1032

1033 **Table 3.** Repeat element statistics for *U. stansburiana* genome from RepeatModeler v2.0.1 (Smit

1034 et al., 2015a) and RepeatMasker (Smit et al., 2015b). Total interspersed repeats is 49.29%.

1035 Percentages are relative to the total genome length.

Retroelements (17.32%)	LINEs	15.44%	L2/CR1/Rex	10.26%
			R2/R4/NeSL	0.01%
			RTE/Bov-B	2.61%
			L1/CIN4	1.82%
	SINEs	0%		
	Penelope	0.59%		
	LTR elements	1.88%	Ty1/Copia	0.27%
			Gypsy/DIRS1	1.03%
			Retroviral	0.15%
DNA transposons (0.27%)	hobo-Activator	0.11%		
	Tc1-IS630-Pogo	0.11%		
	PiggyBac	0.02%		
	Tourist/Harbinger	0.03%		
Unclassified (31.71%)				
Simple repeats (1.58%)				
Low complexity (0.18%)				

1036

1037

Figure 1. A) Map of full range of *Uta stansburiana* (IUCN Redlist) with collection locality of
specimen from which the reference genome was sequenced (DLE 3124) and the study area in
Baja California, MX is highlighted. **B)** Photo of the individual male *Uta stansburiana nevadensis*
(DLE 3124) from whom the reference genome was generated (photo courtesy Dr. Pete Zani). **C)**
Map of samples collected for RNA with percentage of average total annual precipitation
occurring in the winter (December-February; Fick and Hijmans 2017). Samples represent 33
specimens collected across seven sites. Sites were divided into those north (3 populations, 9
samples) and south (4 populations, 24 samples) of the Vizcaíno desert. **D)** Differences in
temperature seasonality (standard deviation of monthly temperatures x 100, BIO4; Fick and
Hijmans 2017) between northern and southern sites.

1048

Chromosomes scaled by gene rank order

Figure 2. Riparian plot of genomic syntenic blocks in four phrynosomatid reference genomes (*Phrynosoma platirhinos*, *Sceloporus undulatus*, *Urosaurus nigricaudus*, and *Uta stansburiana*) and one dactyloid (*Anolis carolinensis*) reference genome. Chromosome-level scaffolds have been re-named to reflect the position of each scaffold when sorted by length and appended with the first letter or first two letters of the genus. Suspected or identified X chromosomes are noted.

1057
1058 **Figure 3. A)** Heatmap of log2-transformed count data for genes with differential expression by
1059 region, controlling for season and sex. Samples are labeled by region, season, and sex. **B)**
1060 Principal components analysis of regular log-transformed count data. PC2 and PC3 are shown
1061 here to better illustrate the distinctions between groups; see Fig S4 for PC1 and PC2. Ellipses
1062 enclose the centroids of the northern and southern samples.
1063

1064

Figure 4. Volcano plot of genes differentially expressed between northern and southern groups,
controlling for seasonal differences. Positive log2 fold change values correspond to genes
differentially over-expressed in the southern group, while negative log2 fold change values
correspond to genes differentially under-expressed in the southern group. ADAM20 is not plotted
here, but see Fig. S5.