SMURF Reconstructs Single-Cells from Visium HD Data to Reveal Zonation of Transcriptional Programs in the Intestine

Juanru Guo[1,2], Simona Sarafinovska[1,5], Ryan A. Hagenson[2,3], Mark C. Valentine[2,4], Joseph D. Dougherty[1,5,6*], Robi D. Mitra[1,2*] and Brian D. Muegge[2,7,8*]

[1]Department of Genetics, Washington University in St. Louis School of Medicine, Saint Louis, 63110, MO, USA.
[2]Edison Family Center for Genome Sciences and Systems Biology, Washington University in St. Louis School of Medicine, Saint Louis, 63110, MO, USA.
[3]Division of Biology and Biomedical Sciences, Washington University in St. Louis School of Medicine, Saint Louis, 63110, MO, USA.
[4]Department of Obstetrics and Gynecology, Washington University in St. Louis School of Medicine, Saint Louis, 63110, MO, USA.
[5]Department of Psychiatry, Washington University in St. Louis School of Medicine, Saint Louis, 63110, MO, USA.
[6]Intellectual and Developmental Disabilities Research Center, Washington University School of Medicine, Saint Louis, 63108, MO, USA.
[7]Department of Medicine,VA Medical Center, Saint Louis, 63106, MO, USA.
[8]Department of Medicine,Washington University in St. Louis School of Medicine, Saint Louis, 63110, MO, USA.

*Corresponding author(s). E-mail(s): jdougherty@wustl.edu; rmitra@wustl.edu; mueggeb@wustl.edu;
Contributing authors: g.juanru@wustl.edu;

Abstract

High-resolution spatial transcriptomics demands new computational tools to accurately assign transcripts to individual cells. We developed SMURF (Segmentation and Manifold UnRolling Framework), a novel soft-segmentation algorithm that maps mRNAs from barcoded capture spots to nearby nuclei. SMURF also "unrolls" cells in complex tissue structures, placing them on standard Cartesian coordinates, thus empowering studies of cellular organization in tissues. Using SMURF, we identified the zonation of gene expression programs along the maturing intestinal villus and the transcription factors that regulate them. We investigated how functionally important proximal-to-distal gene expression gradients in the intestine are specified. We discovered that gene expression gradients along the proximal-distal axis accumulate in the villus tip and that tip gene expression is reprogrammed by environmental signals in the lumen, suggesting environmental inputs are a major determinant of regional transcriptional identity. These results demonstrate SMURF provides a powerful toolbox for analysis of regional transcriptional programs along tissue manifolds.

Introduction

Biology is poised to address a fundamental challenge: describing the molecular features of all cell types in the body at single-cell resolution. However, a full understanding of how cells function can't be achieved without considering how they exist within their microenvironments. Spatial transcriptomics (ST) addresses this challenge by allowing researchers to measure cellular gene expression while preserving spatial context.

High-resolution ST is particularly valuable for the study of organs such as the small intestine, which has a complex and highly organized architecture that is critical for its function. The intestine is a long tube made up of millions of repeating crypt-villus units [1]. Each crypt-villus unit is highly organized and contains multiple cell types that collectively regulate nutrient absorption, motility, and host defense. Cells in the crypt-villus units are specialized to carry out region-specific functions. Despite intense research, defining the factors that shape the spatial organization of the intestine remains a challenge. Important studies using Laser Capture Microdissection (LCM) or early generations of ST technologies revealed interesting transcriptional dynamics along the crypt-to-villus [2, 3, 4, 5, 6] or longitudinal axes [7] but were somewhat limited because they lacked single-cell resolution.

The Visium HD platform dramatically improves the resolution of capture ST methods but brings new analytical challenges. First, there is no native relationship between the tiled oligonucleotides and overlying cells. As a result, many of the spots aren't aligned with cells and thus contain zero or few transcripts. Second, transcripts from a single cell may be spread across multiple adjacent spots, and conversely, a single spot may contain transcripts from multiple adjacent cells. Existing high-resolution segmentation algorithms typically assume that each capture spot contains mRNA from only a single cell [8, 9]. Furthermore, we lack tools to analyze cell organization and gradients of gene expression in tissue substructures, which often have complex geometries, complicating this task. Finally, these substructures often deform during sample preparation and processing, obscuring their native spatial arrangement. Thus, there is a need for new computational approaches that accurately segment and deconvolve mRNA in capture spots to single cells and of regional transcriptional programs along tissue substructures.

To address these challenges, we developed the Segmentation and Manifold UnRolling Framework (SMURF) to extract Visium HD gene expression at single-cell resolution. We introduce the concept of "soft-segmentation", where mRNA counts within a capture spot may be assigned to a single cell or partitioned and assigned to multiple nearby cells in a fashion that optimizes the cosine similarity of each cell to the gene expression cluster to which it belongs. SMURF maximizes the mRNA counts assigned to each cell and achieves precise cell clustering. SMURF also introduces a framework to unroll complex tissue structures in two-dimensional histologic sections using manifold theory. This enables the analysis of gene expression as a function of spatial location along two different axes in the tissue. We used SMURF to reveal the spatial organization of gene expression programs in the intestine along the crypt-to-villus axis. We also identified the transcription factor networks that regulate these programs. Finally, we used SMURF to investigate a fundamental question in the development of the small intestine: are proximal-to-distal gradients in gene expression predetermined, or are they established in response to signals in the luminal environment? We found that the gene programs varying along the proximal-to-distal axis are concentrated in the villus tip. This suggested the hypothesis that signals in the lumen shape proximal-to-distal gradients in gene expression, which we confirmed

using a dataset where a segment of the distal small intestine is surgically moved to a proximal location in an adult mouse. This work demonstrates that SMURF accurately segments Visium HD expression data, is highly sensitive for detecting rare cell types, and can extract novel biology from ST datasets.

Results

Soft-segmentation for spatial transcriptomics

Visium HD capture spots do not align with cellular boundaries. Some spots collect mRNA from multiple cells, and conversely, many cells have transcripts that are distributed across multiple spots. This makes it difficult to reliably assign transcripts to individual cells. SMURF overcomes this limitation using a novel three-step "soft segmentation" approach.

In the first step, SMURF identifies nuclei from H&E or DAPI/IF images using established segmentation methods [10, 11, 12, 13]. It then constructs a gene expression matrix by assigning mRNA counts from capture spots directly overlying each nucleus. Nuclei are then clustered by expression similarity using Scanpy [14, 15, 16]. Simultaneously, SMURF builds a nearest-neighbor network based on nuclear positions to later help assign surrounding spots to nearby nuclei (Figure 1a).

Because nuclei occupy only a small fraction of total cell volume, the initial single-nucleus gene expression matrix is sparse, with low UMI counts [17]. Thus, the second step of soft-segmentation expands the nucleus-anchored gene expression by incrementally assigning nearby spots to nuclei through concentric expansion. Nearby spots are added only if they increase the cosine similarity between the expression profiles of the nucleus and its cluster. This process is repeated iteratively for each nucleus until gene expression clustering stabilizes using normalized mutual information (NMI) [18]. At this step, SMURF allows flexible assignment: a spot can be linked to one nucleus, several nuclei, or none (Figure 1b).

The third and final step of SMURF's soft-segmentation is count deconvolution for spots shared by multiple cells. SMURF deconvolves individual mRNA counts from capture spots assigned to multiple cells, using a combination of quadratic programming and deep learning [19, 20] (Methods, Figure 1c).

SMURF improves single-cell reconstruction of Visium HD data

We benchmarked SMURF's soft segmentation algorithm using ST data from the mouse brain and distal small intestine (the ileum) [21, 22]. We compared SMURF to three commonly used methods (Figure 2a&b). The 8 μm method, the default from 10x Genomics, aggregates 16 2-μm-spots into 8×8 μm^2 "cells" to increase UMI counts for clustering and differential expression. It does not consider nuclear boundaries, often assigning transcripts to "cells" that lack nuclei, only partially cover a nucleus, or span multiple nuclei (Figure 2a&b, 2nd column). Bin2cell, a recently described method for single-cell segmentation, performs better in the mouse brain but often merges multiple cells or misses nuclei entirely in the densely packed small intestine [9] (Figure 2a&b,3 rd column). The Nuclei segmentation method only uses spots directly overlapping nuclei (Figure 2a&b, 4th column); this results in a sparse expression matrix, as it excludes many transcripts from the surrounding cytoplasm. In contrast, SMURF achieves reliable single-cell resolution while accurately assigning cytoplasmic transcripts, outperforming existing methods (Figure 2a&b, last column).

We next quantified segmentation accuracy by measuring the percentage of cells lacking a nucleus and the percentage containing multiple nuclei for each method

(Figure 2c). SMURF was the most accurate in both the brain and small intestine data. Only 0.31% of SMURF cells in the mouse brain lacked a nucleus, and only 1.44% contained multiple nuclei. This was significantly better than Bin2Cell or the $8\,\mu$m method. SMURF also outperformed these methods in the densely populated small intestine, with the lowest rates of both nucleus-less (3.69%) and multi-nucleus cells (11.77%, Figure 2c, right panels), demonstrating that SMURF more accurately assigns transcripts to cells at single-cell resolution than existing methods. We also found that SMURF's cell-level normalization corrects a severe count imbalance artifact observed in transcript-sparse regions when using the 8 μm method (Supplementary Notes, Extended Data Figure 1a). Compared to the nuclei segmentation approach, SMURF assigned far more UMIs to each cell, and the average cell size was much bigger (Figure 2d). Together, these results demonstrate that SMURF can accurately assign Visium HD counts to single cells, with substantially more counts per cell than nuclei segmentation alone.

SMURF improves cell type identification and capture of rare cell types

We hypothesized that SMURF's higher transcript assignment per cell would lead to more accurate cell-type identification and improved estimation of their relative abundances. To test this, we used cell-type proportions from a probe-based ST method as a reference standard [23]. Using Kullback–Leibler (KL) divergence [24], we quantified how closely SMURF and nuclei segmentation (without expansion) reproduced the probe-based ST-derived proportions. SMURF showed substantially lower KL divergence (0.1 vs. 0.738), indicating a more accurate representation of cell-type abundances (Extended Data Figure 1b).

Comparable ground-truth data for the small intestine are limited, so we clustered and annotated cell types independently for nuclei segmentation without expansion and SMURF. Both approaches detected enterocytes, the dominant epithelial population (Figure 2e). However, only SMURF reliably identified rarer epithelial cell types, such as tuft cells and heterogeneous enteroendocrine cells (EECs), as well as smooth muscle cells from the surrounding muscle layer. These cell types were not captured by nuclei segmentation, potentially due to irregular nuclear morphology, low nuclear transcript abundance, or mRNA polarization—factors known to affect these cell populations.

Taken together, our results show that SMURF accurately assigns transcripts to individual cells with single-cell resolution, retaining transcripts from both the nucleus and cytoplasm. Compared to nuclei segmentation, SMURF recovers more transcripts per cell and improves the identification of known cell types in Visium HD data. SMURF's increased count depth enhances statistical power for downstream analyses and enables more reliable detection of rare or morphologically irregular cell types.

SMURF applies manifold detection to unroll ST data from complex tissue structures

After assigning transcripts to single cells, we next wanted to place the cells on a shared set of coordinates that reflect their *in vivo* locations in complex tissue structures. To do this, we developed a novel manifold detection approach that projects cells onto a coordinate system aligned with the structure of interest (Methods). Manifold detection is a mathematical concept that assigns local Euclidean coordinates to small areas of a complex shape. SMURF uses cell-resolved transcriptomes to identify

the tissue structures (e.g. layers in the cortex or the muscularis mucosa of the intestine) and define a manifold. SMURF then "unrolls" this manifold, projecting the cells onto standard Cartesian axes. This functionality empowers studies of cellular organization and regional specification in the coordinate system most interpretable for that tissue.

We evaluated SMURF's manifold detection by analyzing the mouse brain cortex. We first identified a one-dimensional manifold path representing the spatial distribution of cells in the cortical basement membrane using a locally linear embedding algorithm [25, 26] (Figure 3a). We then extended the manifold to two dimensions to include all cortical cells, assigning each an (x, y) coordinate based on its position relative to the basement membrane (Figure 3b&c). This effectively projected the cortex onto a rectilinear coordinate system—"unrolling" the tissue for spatial analysis.

To illustrate how SMURF's manifold algorithm can reveal regional patterning of cell types and gene expression, we divided the unrolled y-axis of the cortex into five bins spanning from the cortical base to the surface. We then plotted the expression of known cortical layer markers across these bins as a heatmap [27] (Figure 3d). Layer-specific gene expression aligned precisely with SMURF-assigned locations: markers of layers 2–4 neurons, such as Cux1 and Pou3f2, were enriched nearer the cortical surface, while Bcl11b and Foxp2, markers of layers 5 and 6, were expressed near the base.

Next, we assigned every cortical cell a cell-type identity based on its transcriptional profile, using the Allen Brain Cell Atlas as a reference [28, 29]. As expected, the cell types stratified along the y-axis, with Layer 2 and 3 neurons at the cortical surface and Layer 6 neurons at the base (Figure 3e). Cortical thickness increased moving laterally on the x-axis, consistent with previous anatomical studies [30, 31]. Non-neuronal cell types also projected according to their known anatomic distributions [32, 33] (Extended Data Figure 2). Taken together, these results demonstrate that SMURF can accurately "unroll" a manifold of cells onto rectilinear coordinates while preserving patterns of gene expression and cellular organization.

SMURF accurately unrolls complex and discontinuous tissue

We next deployed SMURF to better understand regional transcriptional programs in the mouse small intestine. Intestinal gene expression, cell type distributions, and functions vary along both the proximal-to-distal (longitudinal) and crypt-to-villus (radial) axes. We reasoned that SMURF's methods to identify gene expression in individual intestinal cells and then map those cells to a common proximal-to-distal and crypt-to-villus coordinate system could provide new insight into the molecular logic that controls spatial organization in the gut, along with the cell-cell interactions shaping those gradients.

We analyzed Visium HD data from healthy mouse ileum (distal small intestine) prepared in a "Swiss roll" configuration [22] (Figure 3f). We soft-segmented over 40,000 cells and applied SMURF's generalized unrolling tool to reconstruct the tissue's original longitudinal layout (Figure 3g). Pseudobulked epithelial gene expression of the Visium data along the x-axis correlated with a recently published scRNA-seq atlas of the mouse small intestine [7] (Extended Data Figure 3b). SMURF accurately localized regional markers, including proximally enriched Lactase (Lct) and distally enriched Ileal bile acid transporter (Ibat) [7] (Figure 3h). These results demonstrate that SMURF successfully unrolls complex histological sections, restoring the tissue's native spatial context.

We next assigned cells to coordinates along the crypt-to-villus axis. The dataset contains over 1,000 villi, and this analysis posed several challenges (Figure 3i). Villi

often bend and are not perpendicular to the muscularis base. In addition, some villi are partially severed or detached during sectioning. To address these issues, we developed an iterative method inspired by dynamic programming. We identified sequential layers of cells to identify a corrected y-axis for each villus, normalized to maximum villus height (Methods, Figure 3k, Extended Data Figure 3c, Extended Data Figure 4). Cell types were detected in the expected positions, including neuronal and muscle cells at the base of muscularis propria, stem and Paneth cells in the crypt, proliferative transit-amplifying (TA) cells above the stem cell zone, and differentiated epithelial types along the villus (Figure 3i). Marker genes aligned appropriately along this axis, including the stem cell marker Olfm4 at the crypt base, the TA marker Krt19 above the stem cell zone, enterocytic Fabp6 along the villus, and Ada at the villus tip (Extended Data Figure 3d-e). Interestingly, some EECs and tuft cells appeared near the base of the TA zone, suggesting they escaped the typical upward migration of differentiated cells. While retrograde migration has been observed for EECs, it has not been described for tuft cells [34, 35].

Together, these results demonstrate that SMURF accurately defines manifolds along complex tissue structures and projects cells onto rectilinear coordinates. Our methods can be applied to subregions that are torn, bent, and/or missing to preserve the *in vivo* spatial arrangements of different cell types.

SMURF identifies the spatial trajectory of intestinal development

Stem cells at the crypt base continuously give rise to rapidly dividing TA cells, which differentiate into specialized digestive and barrier functions as they migrate up the villus [36] (Figure 4a). We hypothesized that SMURF's projection of complex tissue structures onto rectilinear coordinates could be used to quantitatively examine the relationship between intestinal epithelial cells' developmental state and their spatial position on the crypt-villus axis. We focused on the developmental trajectories of the most abundant epithelial cell type, enterocytes, from stem and TA cells. Pseudotime trajectories were measured using diffusion maps and visualized with partition-based graph abstraction (PAGA) and force-directed graph layouts [37, 38, 39] (Figure 4b). Pseudotime trajectory captured the expected transition from stem and Paneth cells to mature villus cells (left and middle panels). SMURF-assigned y-axis position followed a similar distribution (right panel), and was significantly correlated with pseudotime ($corr = 0.79$, $p < 10^{-10}$; Figure 4c). These findings demonstrate SMURF can recapitulate the spatial-temporal coupling of cell differentiation along the crypt-villus axis solely from spatial location.

SMURF defines novel ileal zonation gene programs

We next sought to identify the gene programs that change along the crypt-villus axis during epithelial cell maturation. We binned villi into 6 zones spanning the crypt to villus tip based on the corrected y-axis positions (Figure 4d). We pseudobulked gene expression from the stem/Paneth, TA, and enterocyte clusters for each zone. Normalized gene expression was highly zonated across the villi (Figure 4e). We identified 1,631 genes that were significantly zonated by differential expression analysis (Methods, Supplementary Table 1). We refined this list further to a set of "landmark" genes highly specific for each zone by algorithmic and manual curation (Supplemental Note, Extended Data Figure 5).

KEGG pathway enrichment analysis revealed specialized transcriptional programs in each zone (Figure 4f). Zone 1 contained pathways regulating stem cells, including DNA replication and Wnt signaling, while TA cells in Zone 2 were enriched

in pathways involved in the cell cycle and RNA synthesis. For example, the canonical stem cell markers Lgr5 and Mcm2 were localized to Zone 1, while proliferative genes that mark TA cells, including Mki67, Nusap1, and Pcna, are more abundant in Zone 2 (Figure 4g). Enterocytes at the villus base in Zones 3 and 4 express programs for carbohydrate and vitamin metabolism, exemplified by sucrase isomaltase (Sis) and the B12-intrinsic factor receptor Cubulin (Cubn) (Figure 4h). As enterocytes moved up the villus, they acquired functional enrichment for protein and vitamin metabolism in Zone 5, such as the brush border peptidase Dpp4. This maturation progressed to pathways for lipid metabolism in tip cells of Zone 6, including Fabp1 and Apoc2, the cofactor for lipoprotein lipase (Figure 4h). This functional progression is concordant with prior studies in the jejunum [2], but ST provides more power to identify zonated genes with better resolution along the crypt-villus axis.

SMURF defines novel secretory cell zonation programs

We next turned our attention to crypt-to-villus zonation in less abundant secretory epithelial cell types. Goblet cells, tuft cells, and the heterogeneous EECs are individually scattered throughout the villus and can't be isolated by LCM or conventional spatial analysis with large bin sizes. Using the same six zones, SMURF analysis also identified significant zonation along the crypt-to-villus axis for each secretory cell type (Figure 4i-k). For instance, a recent study of the human small intestine reported that tuft cell markers were zonated towards the bottom of the villus [4]. By contrast, in the mouse ST data, tuft markers are variably distributed. The tuft marker Kit was localized to the base, the essential transcription factor Pou2f3 was evenly distributed, and immune and chemosensory receptors, including Il17rb, Sucnr1, and Gnat3, were more abundant at the middle or upper villus [40, 41, 42, 43] (Figure 4j). Similarly, hormone expression by EECs was highly zonated. Crypts were enriched for cells expressing Substance P (Tac1) and GLP-1 (Gcg), while cells expressing Serotonin (Tph1), Neurotensin (Nts), and Peptide YY (Pyy) are detected only in villus zones (Figure 4k). These results are consistent with prior immunohistochemical studies [44, 45]. While villus morphogen gradients, including bone morphogenetic proteins (BMPs) have been implicated in Nts and Pyy emergence, the factors that limit Tac1 and Gcg expression in the crypt remain unknown.

Zonated transcription factor regulatory networks along the villus

The network of transcription factors (TF) regulating zonated gene expression in the mature villus remains poorly defined. Using SCENIC, we identified active TFs and their targets in each zone based on expression profiles from stem/Paneth, TA, and enterocyte cells [46]. We detected 65 TFs with significant gene regulon score in at least one of the six zones (≥ 0.05, Table S3, Supplementary Notes). The analysis confirmed previously reported zonation of TFs, including the stem/Paneth regulating TF Sox9 in Zone 1 and immediately early TFs Jun and Fos at the villus tip [2, 47, 48] (Figure 5a). Notably, our analysis revealed previously unappreciated zonation of several TF regulons. For example, the Vitamin D receptor (Vdr), the bile acid receptor Nr1h4 (also known as Fxr), and the peroxisome proliferator-activated receptor delta (Ppard) exhibited peak regulon scores at the base of the villus, despite their luminal ligands being more abundant near the villus tip. We next asked if TF expression on the crypt-villus axis mirrors the zonation of their gene targets (Figure 5b). The majority of TFs were most highly expressed where their regulons were active. In contrast, the ileal TF Gata6 was expressed in the crypt (Zones 1-2), but its regulon

scores peaked in the villus and tip (Zone 4-6) [49]. This suggests Gata6 function may depend on post-transcriptional regulation or interaction with other zonated factors.

Cooperative TF binding is a well-established mechanism for regulating gene expression. We therefore mapped TF interaction networks within each zone (Extended Data Figure 6). In Zone 2, a highly interconnected network of 7 TFs emerged, anchored by Elf3 and Klf5 (Figure 5c). Elf3 mediates Wnt-independent B-catenin activation in colon cancer, and Klf5 is an essential regulator of intestinal barrier integrity and stemness [50, 51, 52, 53]. Other members of this network, including Ehf, Nono, Bclaf1, and Vezf1, are not well studied in the intestine. Genes with enriched expression in Zone 2 were significantly more likely to be targets of multiple TFs in this network (p-value $= 8 \times 10^{-7}$, Methods, Figure 5c). Genes co-targeted by two or more TFs were enriched for RNA biogenesis and mitochondrial metabolism, core TA cell functions. Thus, we identified a novel cooperative TF network controlling TA cell proliferation.

At the villus tip, we identified a distinct TF network, including AP-1 complex members Jun and Fos, previously linked to tip identity, and novel TFs Irf7, Mxd1, and Klf4 [4, 2, 54, 55]. As in Zone 2, tip-zonated genes were over-represented among those targeted by multiple TFs (p-value $< 10^{-10}$, Methods, Figure 5d). These genes are involved in amino acid, lipid, and vitamin metabolism, as well as host defense, suggesting that AP-1 complex and environmental sensors such as Irf7 cooperate to drive tip-specific functions.

Regionally specific gene programs accumulate in the epithelial villus tip

The intestine is functionally specialized along its proximal-distal axis, yet the molecular mechanisms that maintain regional identity remain unclear. To address this, we divided villi from the proximal and distal ileum by x-axis coordinates and performed differential expression analysis on pseudobulked counts from each villus zone. Surprisingly, more genes were differentially expressed in upper villus zones than at the base [7, 56, 57] (Figure 6a). Our ST data are derived from the mouse ileum. To test the idea that villus bottom zonation is preserved across regions while villus top zonation is regionally specific, we plotted markers of jejunal crypt-villus zonation identified previously by LCM [2] (Extended Figure 7a&b). The genes identified as bottom markers in the jejunum were also expressed predominantly in Zones 2 and 3 of the ileum. In contrast, the jejunal villus top markers were not segregated at the top of the ileum, supporting the idea that villus tips have regionally specific transcriptional programs. Functional enrichment showed that proximal ileal villus tips upregulate metabolic pathways for cholesterol and pyruvate metabolism, including lipid transporters Apoa4 and Apoc3, consistent with the proximal ileum's role in absorbing excess dietary fats [58] (Figure 6b,d). In contrast, distal ileal tips are enriched for bile secretion and steroid biosynthesis, such as the beta-alanine transporter Slc6a14 [59] (Figure 6c,d).

These data suggest that proximal–distal cellular identity is not dictated solely by stem cell-intrinsic transcriptional programs. Although intestinal stem cells encode much of their progeny's regional identity, recent evidence challenges this view [7, 60, 61]. We therefore hypothesized that luminal gradients of environmental signals, including dietary components, xenobiotics, and microbes, interact with intestinal TF networks to establish regional identity.

To test this, we examined a small bowel resection (SBR) model, in which 50% of the small intestine is surgically removed and the remaining segments are joined,

exposing the distal ileum to proximal luminal contents [62] (Figure 6e). Our hypothesis predicts that this exposure would induce proximal transcriptional programs at distal villi tips. We re-analyzed previously reported scRNA-seq from SBR ileal tissue and sham-operated controls, identified differentially expressed genes, and mapped these genes to spatial zones in healthy ileum using our ST dataset. As predicted, genes upregulated after resection were significantly enriched in the tip zonated genes (empirical chi-square test, $p = 9.99 \times 10^{-6}$), Figure 6f). None of the induced genes were expressed in stem and TA cells, indicating that most transcriptional changes occur downstream of any stem cell reprogramming. As reported, the SBR upregulated genes were enriched for proximal lipid metabolism pathways [62]. Together, these results support our hypothesis that environmental cues, rather than stem cell-intrinsic programs alone, shape regional identity. However, we cannot exclude additional effects from epithelial proliferation or hormonal signaling triggered by surgery.

Discussion

Visium HD and other high-resolution, capture-based ST platforms have created an urgent need for algorithms that can accurately assign transcripts in densely tiled capture spots to individual cells. Here, we showed that SMURF's novel "soft-segmentation" method more accurately and comprehensively assigns mRNAs to cells, segments cells with a higher resolution, and more reliably detects moderate and rare cell types than existing methods. SMURF also "unrolls" tissue substructures, even when parts of them are damaged or missing, to place cells on Cartesian coordinates allowing for quantitative analysis of the *in vivo* spatial organization of cell types and transcriptional programs.

We applied SMURF to analyze Visium HD data from the small intestine and gained new insights into the spatial organization of this organ. Prior work in this area has revealed important principles, but was constrained by coarse resolution or the use of labor-intensive methods [2, 3, 4, 6, 63]. Using SMURF, we identified cell-type-specific zonated gene expression programs and novel TF regulatory networks along the crypt-villus axis, many of which were not previously identified. We also used SMURF to investigate a fundamental question about small intestine development: are proximal-to-distal gradients in gene expression predetermined or established in response to environmental signals? We found that regional gene expression differences are largely restricted to villus tip cells and can be reprogrammed by relocating the distal intestine to a proximal environment, demonstrating that proximal–distal patterning in adult tissue is shaped more by environmental cues than by intrinsic stem cell programming.

SMURF, in its current form, also has some limitations. First, SMURF uses H&E or DAPI/IF-stained images to detect nuclei, so it cannot analyze datasets lacking tissue images. Second, when assigning transcripts from capture spots near cell boundaries, SMURF cannot always resolve cells that share a high degree of expression similarity and are closely adjacent, such as the stem and Paneth cells in the intestinal crypt, although it still outperforms other methods in this regard. Finally, ST datasets, including associated histology images, can be extremely large, making the analysis computationally demanding. SMURF's full implementation of employs a deep-learning algorithm for deconvolution, which requires GPUs to run efficiently [64]. To make SMURF more widely accessible, we have also developed a "lite" version of SMURF, which can run on most workstations (Supplementary Notes).

As ST continues to evolve, new platforms will utilize even higher resolution capture spots and profile larger numbers of cells. The full potential of these technologies will only be achieved with approaches like SMURF's that consider all capture spots and have the ability to assign mRNAs in the same capture spot to different cells. SMURF's framework is inherently extensible to emerging technologies for three-dimensional (3D) tissue ST. Finally, SMURF's unrolling algorithm could be extended to other spatial technologies, such as protein-based assays or image-based tools.

Our study highlights SMURF's utility for uncovering principles of tissue organization during normal development. An important future direction will be to deploy this approach to study disease models at single-cell resolution. Many intestinal disorders, including adaptation to massive SBR and epithelial injury in inflammatory bowel disease, disrupt normal crypt–villus architecture. SMURF offers a framework to dissect how multiple cell types interact in these contexts, including "4D" studies of evolving responses over time. We have also demonstrated SMURF's utility in other tissues (Supplementary Notes) and anticipate that SMURF will provide insight into other spatially complex organs such as the kidney, liver, and lung.

Methods

Soft Segmentation Analysis

Introduction

In this section, we provide a detailed description of the soft-segmentation method implemented in SMURF. Cells are denoted by the index i, genes by j, and spots by k. Since cells are typically associated with exactly one nucleus, we denote both a cell and its corresponding nucleus by i, emphasizing their one-to-one correspondence. The inputs to the soft segmentation algorithm consist of: (1) the $2\,\mu$m spot gene expression matrix S; (2) the H&E or DAPI/IF tissue image Img^o; and (3) the "spot to pixel" data frame.

We define the $2\,\mu$m spot gene expression matrix S to have dimensions $K_{\text{spots}} \times J_{\text{genes}}$, where K_{spots} represents the number of spots and J_{genes} the number of genes, with each element $S_{kj} \geq 0$. Each pixel in the original image Img^o represents a grayscale value (0–255) for DAPI/IF images or a 1×3 RGB array for H&E stained images. The matrix S is linked to the processed image Img via the "spot to pixel" data frame. Note that some pixels in Img^o are not covered by spots, and each spot encompasses multiple pixels; hence, we focus on the area covered by spots, defined as Img. We denote by Img_k the set of pixels corresponding to spot k.

SMURF requires users to generate their own nuclei segmentation results. To facilitate this, we provide tutorials for DAPI/IF images (using Cellpose [10, 65]) and H&E-stained images (using StarDist [11, 12, 13]); users may also choose their preferred segmentation tools. The final nuclei segmentation output, denoted as $Nuclei\_img$, is a two-dimensional matrix with the same dimensions as Img. In $Nuclei\_img$, regions without nuclei are assigned a value of 0, whereas regions containing nuclei receive unique numerical identifiers.

Given the substantial volume of data and resulting outputs, SMURF defines a new Python object, the Spatial Object (SO), initially constructed from the three inputs. The SO records the nuclei segmentation results, detailed information for each cell, and visualization outputs. For clarity, we refer to the components of SO by their object names rather than describing each individual element of SO here.

The soft-segmentation in SMURF involves three main steps: nuclei analysis, cell expansion, and count deconvolution for shared spots. For the first step, a nucleus-by-gene matrix is constructed and clustered to provide initial cell type annotations, and a nearest neighbor network is constructed to estimate approximate cell size; second, cells are expanded by iteratively assigning adjacent spots based on their similarity to the cell's cluster identity, with periodic reclustering to refine assignments; finally, overlapping spots that are shared by multiple cells are resolved through optimization methods (quadratic programming and deep learning), ensuring accurate gene expression profiles for each cell. Finally, SMURF outputs a spatial single-cell `AnnData` object containing the gene expression matrix G ($I_{\text{cells}} \times J_{\text{genes}}$). Each cell is annotated with its gene expression profile, spatial coordinates, preliminary cluster assignment, and the cosine similarity to its assigned cluster.

Step one: Nuclei Analysis

The nuclei analysis includes mainly three parts: building the nuclei gene expression array, clustering nuclei, and estimating the potential sizes of cells based on location. It takes in the 2μm spot gene expression matrix S and nuclei segmentation results $Nuclei\_img$ from the image.

Before the analysis, we first filter the $2\,\mu$m spot gene expression matrix S based on the criterion

$$\sum_{k=1}^{n_{\text{spots}}} S_{kj} \geq \text{mincounts},$$

with the default threshold set at mincounts $= 1000$. We denote the resulting filtered matrix as S'.

The `Nuclei_img` contains pixel-level data, and each $2\,\mu$m spot corresponds to approximately 50 pixels. Consequently, many nuclei overlap multiple capture spots and, conversely, some capture spots are shared by multiple nuclei. We define $pct_{ik} =$ (number of pixels in nucleus i of spot k)/Img_k, with $0 \leq pct_{ik} \leq 1$, to denote the percentage of pixels in spot k that overlap the nucleus of cell i. For each nucleus i, we record all spot IDs for which the fraction of overlapping pixels is at least a specified minimum (the default fraction is $1/6$). The transcripts captured in all spots that meet this criterion for nucleus i are assigned to that nucleus. Using this initial assignment of transcripts to cells, we create the gene count matrix $X^{(0)}$, defined as:

$$X_{ij}^{(0)} = \sum_{k=1}^{n_{\text{spots}}} \mathbf{1}(pct_{ik} \geq 1/6) \times S'_{kj}$$

where $X^{(0)}$ is an $I_{\text{cells}} \times J_{\text{genes\_filtered}}$ matrix and $X_{ij}^{(0)}$ indicates the count expression of gene j in nucleus i.

In the next part, we filter out nuclei with few transcripts assigned to them, since extremely low counts could be due to errors like waterdrops on the capture surface and influence the accuracy of further analysis. We then analyze matrix $X^{(0)}$ using the standard Scanpy [14, 15] workflow, including filtering out genes that are not expressed in some minimum number of cells, count normalization, dimensionality reduction via principal component analysis (PCA), and Leiden clustering. Through this approach, we obtain initial clustering results $C^{(0)}$ based on the $X^{(0)}$, where $C^{(0)}$ is a $1 * n_{cells}$ array that assigns each nucleus into one of $n^{(0)}$ clusters based on gene expression profile. These initial cluster assignments are based solely on transcripts assigned to nuclei from overlapping capture spots.

Finally, we estimate the potential number of spots belonging to a cell based on its cell density. We calculate the center of each nucleus by averaging the locations of the identified spots associated with that nucleus. We employ the k-nearest neighbors (KNN) algorithm [66], linking each nucleus to its three closest neighbors. For each nucleus i, we record $size_i$ as the number of spots occupied by the nucleus and the distances d_{io} (for $o = 1, 2, 3$) from the center of nucleus i to the centers of its three nearest neighbors. Assuming that the nuclei are approximately circular, we calculate d'_i as:

$$d'_i = \frac{1}{3} \sum_{o=1}^{3} d_{io} - 2\sqrt{\frac{size_i}{\pi}},$$

which represents the average distance between nearby cells minus twice the estimated radius (calculated from the cell size), thus approximating the maximum extent of the spots.

Step two: Cell Expansion

Having assigned transcripts to nuclei from all overlapping capture spots, we next sought to assign transcripts from nearby, non-overlapping capture spots, effectively

expanding these nuclei into cells. During this nuclei expansion phase, we use an alternating optimization strategy to expand the nuclei, which generates robust cell gene profiles and reliable clustering results. In essence, we iteratively assign adjacent spots to cells based on whether their addition increases the cell's similarity to the cluster to which it is assigned. We perform periodic reclustering to refine the assignments.

Here, we use the iteration variable r (starting from $r = 1$) to record each iteration and define the initial nuclei results as the starting point ($r = 0$). For example, as previously introduced, $X^{(0)}$ represents the nuclei-by-gene expression matrix, $C^{(0)}$ denotes the nuclei clustering results.

In the beginning of each iteration r, we calculate the average expression of each cluster, denoted by $w^{(r-1)}$, which has dimensions $n^{(r-1)} \times n_{\text{genes\_filtered}}$. For each cluster $c = 1, 2, \ldots, n_c^{(r-1)}$ and for each gene j, the average expression is defined as

$$w_{cj}^{(r-1)} = \frac{1}{\|C_c^{(r-1)}\|} \sum_{i \in C_c^{(r-1)}} X_{ij}^{(r-1)},$$

where $C_c^{(r-1)}$ denotes the set of cells belonging to cluster c in $C^{(r-1)}$ and $\|C_c^{(r-1)}\|$ represents the number of cells in this set. Then, we normalize the gene expression by

$$\hat{w}_c^{(r-1)} = \frac{w_c^{(r-1)}}{\|w_c^{(r-1)}\|}.$$

Additionally, we calculate

$$\bar{d}_c^{(r-1)} = \frac{1}{\|C_c^{(r-1)}\|} \sum_{i \in C_c^{(r-1)}} d_i',$$

to decide the maximum expansion of spots for each nuclei. By default, we use

$$\tilde{d}_c^{(r-1)} = \max\left(1, \min\left(2, \lceil \bar{d}_c^{(r-1)} \rceil + 1\right)\right).$$

In the second stage, our goal is to derive the matrix $X^{(r)}$ ($n_{\text{cells}} \times n_{\text{genes\_filtered}}$) so that cells within the same cluster exhibit high cosine similarity. We denote the gene expression vector for cell i in the current iteration as $X_i^{(r)}$ and let $cell_i^r$ represent the spots assigned to cell i that were not previously associated with nucleus i in iteration r. The objective function is defined as follows:

$$
\begin{aligned}
\Phi_1(X^{(r)}) &= \sum_{c=1}^{n_c^{(r-1)}} \sum_{i \in C_c^{(r-1)}} \cos_{\text{sim}}(X_i^{(r)}, \hat{w}_c^{(r-1)}) \\
&= \sum_{c=1}^{n_c^{(r-1)}} \sum_{i \in C_c^{(r-1)}} \cos_{\text{sim}}\left(X_i^{(0)} + \sum_{k \in \text{cell}_i^r} S_{kj}', \hat{w}_c^{(r-1)}\right)
\end{aligned}
\tag{1}
$$

Here, our objective is to determine $cell_i^r$ for $i = 1, 2, \ldots, n_{cells}$. For each spot k included in $cell_i^r$, the distance from the spot to the nucleus must be less than $\tilde{d}_c^{(r-1)}$, and the spots within a cell must form a connected region.

To achieve this within limited memory resources, we employ a greedy algorithm that expands the spots one-by-one in a specified order. A spot is added if its inclusion improves the objective function (1), and expansion in a particular direction ceases if the previously added spot is excluded.

In the next phase of iteration r, we analyze $X^{(r)}$ following the Scanpy tutorial as described earlier, thereby obtaining the clustering information $C^{(r)}$. To compare the clustering outcomes between iterations r and $r-1$, we calculate the normalized mutual information score (NMI) using the formula:

$$\text{NMI}(C^{(r)}, C^{(r-1)}) = \frac{2 \times I(C^{(r)}; \, C^{(r-1)})}{H(C^{(r)}) + H(C^{(r-1)})}$$

where $I(C^{(r)}; \, C^{(r-1)})$ denotes the mutual information between the clustering iterations, defined as:

$$I(C^{(r)}; \, C^{(r-1)}) = \sum_{u \in C^{(r)}} \sum_{v \in C^{(r-1)}} P(u,v) \log \left(\frac{P(u,v)}{P(u)P(v)} \right),$$

$H(C^{(r)})$ and $H(C^{(r-1)})$ represent the Shannon entropies of the respective clustering iterations, defined as:

$$H(C^{(r)}) = -\sum_{u \in C^{(r)}} P(u) \log P(u).$$

This calculation quantifies the consistency of clustering results between iterations r and $r-1$, with higher NMI values indicating greater similarity. Typically, the NMI increases as the iterations progress, and by default, the process terminates when there is no improvement in the NMI between two consecutive iterations.

Step three: Count Deconvolution for Shared Spots

The previous step assigns transcripts from spots that do not overlap nuclei to cells; however, transcripts from a single spot may be assigned to multiple cells. In this step, we deconvolve counts from shared spots so that every transcript is assigned to at most one cell. We denote these assignments as $r = *$. At this stage, we isolate spots that are exclusively associated with a single cell and sum their gene counts to obtain X^{final_0}. Next, we focus on the ambiguous spots, which can be categorized into two types: those shared by cells from different clusters, and those shared by cells within the same cluster. We first address the former by transforming situations where multiple clusters share spots into scenarios resembling the latter case. Finally, we apply deep learning optimization to the objective function (Equation 1) to determine the proportion of each cell cluster present within the shared spots.

For the first scenario, we simplify the problem to a case where cells from different clusters, denoted by the set u_k, share a single spot k, with each cluster's contribution represented by the weight matrix $\hat{w}_c^*$. Due to the resolution of Visium HD, u_k typically contains 2 or 3 clusters. We then employ sequential least-squares programming and define the equation as:

$$\sum_{c \in u_k} \text{pct}_{ck} \cdot \hat{w}_c^* = S_k' / \left(\sum_{j}^{n_{genes}} S_{kj}' \right),$$

where $\sum_{c \in u_k} \text{pct}_{ck} = 1$ ensures that the proportions sum up to unity and S_k' represents the gene expression array for spot k. After obtaining the values of pct_{ck},

we employ a binomial model to determine the precise count via random number generation. We then define X^{final_1} as $\sum_{k \in cell_i \bigcap (\text{i is only cell of c in k }) \bigcap (c \in u_k)} pct_{ck} * S'_k$.

Here, each specific cell i contains spots with predetermined area percentages. However, there remains a subset of spots v_k that are shared by at least two cells within the same cluster. We refine the objective function (2) to incorporate both the already assigned spots and those yet to be assigned, as follows:

$$\Phi_2(X) = \sum_{c}^{n_c^*} \sum_{i \in C_c^*} \cos_{\text{sim}} \left(X_i^{final_0} + X_i^{final_1} + \sum_{k \in v_k} pct_{ik} \cdot S'_k, \hat{w}_c^* \right) \qquad (2)$$

pct_{ik} satisfies the condition:

$$\sum_{i \in (\text{cells partially in spot k})} pct_{ik} = 1.$$

The maximization of Equation (2) is achieved through gradient-based optimization using the PyTorch library. Subsequently, we employ a binomial model to precisely determine counts via random number generation. Moreover, since each cell comprises a spot and its associated proportion, we revert to the unfiltered S to recover the gene expression count matrix for all genes.

Unrolling Analysis

Many tissues possess intricate biological structures that are challenging to study, especially since some tissues can become deformed on slides, obscuring their native organization. SMURF's soft segmentation algorithm identifies distinct clusters of cells that include spatial information. Using these clusters to define a manifold, we propose a framework to unroll and elucidate the underlying biological architecture of tissue substructures.

We first select a specific cluster from the soft-segmentation output to guide the unrolling process. Because the relevant cluster may not perfectly correspond to the boundary of a tissue substructure, the set of cell coordinates can be augmented by user-defined points and by inserting the midpoints between existing cells. Cells that are isolated or located at the periphery are filtered out if their average distance to the nearest neighboring cells exceeds a specified threshold. Next, we apply Locally Linear Embedding (LLE) to preserve local neighborhood structures, yielding a new x-axis represented by the matrix L (of dimensions $n_l \times 2$), where n_l denotes the number of selected cells. The LLE algorithm comprises several key steps: neighborhood selection, weight matrix construction, and low-dimensional embedding calculation. For each data point l_i in L, its local neighborhood is identified using the k-nearest neighbors method:

$$N(i) = \{l_{j_1}, l_{j_2}, \ldots, l_{j_k}\}$$

where $N(i)$ denotes the neighborhood of the data point l_i. For each data point l_i, we compute weights p_{ij} that best approximate l_i using its neighbors $N(i)$, with the objective to minimize the reconstruction error:

$$E(P) = \sum_{i=1}^{n} \left\| l_i - \sum_{j \in N(i)} p_{ij} l_j \right\|^2$$

620 subject to the constraint $\sum_{j \in N(i)} p_{ij} = 1$, which guarantees that the weights sum to
621 one and maintain local linearity. Once the weights p_{ij} are determined, the algorithm
622 seeks a low-dimensional embedding q_i for each l_i that preserves the local geome-
623 try defined by these weights. The embedding is then optimized by minimizing the
624 following cost function:

$$\Phi'(Q) = \sum_{i=1}^{n} \left\| q_i - \sum_{j \in N(i)} p_{ij} q_j \right\|^2$$

625 After obtaining Q, we reorder L according to the magnitude of q_i for each element
626 in L. To present a smooth result with consistent distances between units on the x-
627 axis, we first smooth the points L' $((n_l - n_{avg} + 1) \times 2)$ by averaging the closest n_{avg}
628 points:

$$l'_i = \frac{\sum_{j=i}^{i+n_{avg}-1} l_i}{n_{avg}}, \quad i = 1, \ldots, (n_l - n_{avg} + 1)$$

629 Next, we use interpolation functions to create an x-axis such that the actual distance
630 between each unit point is consistent. We first calculate the Euclidean distances d_i
631 between two nearby units in L'. The cumulative distance D_i is the total distance from
632 the first point to the i-th point. Specifically, $D_0 = 0$, and for $i = 1, 2, \ldots, (n_l - n_{avg})$,
633 it is given by:

$$D_i = D_{i-1} + d_{i-1}$$

634 The total distance $D_{n_l - n_{avg}}$ represents the distance from the starting point to the
635 last point. A cubic interpolation function f is established for the coordinates of L'.
636 We define a new sequence of distances containing $n_l - n_{avg}$ points evenly distributed
637 points D'_j, where:

$$D'_j = j \cdot \frac{D_{n_l - n_{avg}}}{n_l - n_{avg}}, \quad \text{for} \quad j = 0, 1, 2, \ldots, n_l - n_{avg}$$

The new coordinates $f(D'_j)$ are computed using the interpolation functions for each
D'_j, forming new points ℓ''_j. Thus, we create evenly and smoothly distributed points
for the x-axis.

Depending on the tissue type, the y-axis is constructed either based on absolute
distances or on cell connectivity. In the first case, for each cell, we identify its nearest
unit on the x-axis and calculate the Euclidean distance from the cell to that unit,
assigning this distance as the y-coordinate. In the second case, cells that are in
close proximity are considered connected. For each cell, we determine the shortest
connected path to the x-axis, define the y-layer as the number of cells along that
path, and compute the y-coordinate by summing the distances along this path.

However, given the limited computational resources and the vast number of cells,
calculating the shortest path for each cell individually is time-consuming. Moreover,
the available paths are normally restricted to a limited set of directions. To over-
come these challenges, we progressively generate y-layers by iteration. We begin by
constructing the first layer using an iterative approach. For each unit on the x-axis,
we identify the 18 closest cells within a specified distance and label them as Layer
1. Then, for each cell in Layer 1, we find the nearest neighboring cells among the
remaining cells within the limited distance. To ensure that the cells are captured in
the correct direction, we use a central reference point and require that the direction
from the center to the original cell is consistent with the direction from the center

to the Layer 1 cell. If a cell is selected multiple times by different Layer 1 cells, we retain the path with the shortest total length and record that path as the shortest one, assigning the corresponding x-axis coordinate to that cell. This process continues by iterations: in each iteration, the next layer is generated following the same procedure until no qualified cells are left.

Mouse Brain data Analysis from 10x Genomics[21]

Nuclei Segmentation

SMURF requires nuclei segmentation results as input. In this study, we utilized Squidpy (v1.4.1) [67] with Stardist [13] (v0.8.5) to perform nuclei segmentation on mouse brain H&E data. We first used SMURF to get the exact spot-occupied area of the H&E plot. Since a small portion of the H&E staining in this data was damaged and exhibited abnormally dark red regions, we converted these areas to a pink hue similar to that of the blank regions. Subsequently, we divided the entire image into 20 blocks with a small overlapping area between adjacent sections. For each block, the "2D_versatile_he" model from Stardist was applied, and the segmentation results were then combined. To account for split cells at the block boundaries, we ensured that the overlapping regions were preserved and only retained cells that were completely contained within a block. These refined nuclei segmentation results were used as input for the SMURF analysis of the mouse brain data.

SMURF Analysis

We input the nuclei segmentation data along with the original output from 10x Genomics and followed the SMURF protocol to generate the final gene expression matrix along with primary clustering results. For the unwrapping of the mouse cortex, we first selected the cluster corresponding to the basement membrane of the cortex, then filtered out isolated cells within this cluster and used the filtered cells to build the x-axis. We used the Euclidean distance to define the y-axis and selected a center point to determine the orientation. Next, we constructed both the x-axis and y-axis based on the spatial coordinates of the cells. The y-axis was evenly divided into five sections, and we subsequently plotted a heatmap of gene expression markers that delineated the distinct layers of the cortex [27, 30].

Cell type annotation

We uploaded the complete gene expression dataset directly from the SMURF output to MapMyCells (RRID: SCR_024672) to align cells with the 10X Whole Mouse Brain (CCN20230722) reference [29]. We then selected layer-specific neurons with subclass bootstrapping probabilities greater than 0.90 and plotted these cells along the new axes, with colors indicating layer information. Similarly, we filtered for specific non-neuronal cell types with subclass bootstrapping probabilities exceeding 0.95 and visualized them on the same coordinate system.

Mouse Small Intestine Data Analysis from 10x Genomics [22]

Nuclei Segmentation

SMURF requires nuclei segmentation results as input. In this study, we utilized Squidpy (v1.4.1) [67] with Stardist (v0.8.5) [13] to perform nuclei segmentation on mouse small intestine H&E data. We first used SMURF to determine the exact

spot-occupied areas of the H&E image. The entire image was then divided into 36 blocks with a small overlapping area between adjacent sections. For each block, the "2D_versatile_he" model from Stardist was applied, and the segmentation results were subsequently combined. To account for split cells at block boundaries, we preserved the overlapping regions and retained only those cells that were completely contained within a block. These refined nuclei segmentation results were used as input for the SMURF analysis of the small intestine data.

SMURF analysis

We followed the SMURF protocol to obtain the final gene expression matrix along with primary clustering results. For the unrolling analysis, we manually removed regions where the small intestine was inadvertently cut or damaged at both the beginning and the end. We then selected the clusters corresponding to the muscularis externa and myenteric plexus to define the x-axis, and filtered out isolated cells within these clusters. Finally, we adhered to the SMURF pipeline to define the y-axis layer by layer.

Single-Cell Downstream Analysis

We followed the Scanpy (v. 1.10.0) [14] pipeline to analyze the raw counts generated by SMURF. Initial filtering removed low-count cells based on SMURF's cluster assignments: for cells in Peyer's patches we required total counts to be at least smaller of 50 and 5 % of that cluster's total counts, and for all other clusters we required total counts to be at least the smaller of 50 and 10 % of that cluster's total counts. We then excluded cells with 2000 or more detected genes, 2000 or more total counts, a mitochondrial gene fraction of 25 % or higher, or a cell size of 45 or greater. These quality-control thresholds ensured the removal of low-quality or abnormal cells, enabling robust downstream analysis.

We then normalized the cells by scaling each cell's total counts to 1000 and performed log-normalization to stabilize variance across cells. Next, we retained the top 4000 variable genes and two key genes (Pyy and Sct). The data were scaled such that the maximum expression value for each gene was set to 10. We performed PCA to reduce the dimensionality to 40 components, followed by the construction of a nearest-neighbor graph using 25 neighbors. For visualization, we applied UMAP, and cell clustering was conducted using the Leiden algorithm with a resolution of 3 [68, 16]. Differential expression analysis was carried out using the default method of Scanpy (scanpy.tl.rank_genes_groups, t-test) [14]. We annotate the cell types by gene markers according to reference data and knowledge with the help of location information [7, 69, 70].

To detect the crypt–villus boundary, we first performed a rough division based on a normalized distance threshold of 9, temporarily assigning cells with lower values to the crypt group and those with higher values to the villus group. We then conducted differential gene expression analysis between these two groups, selecting genes with an adjusted p-value below 1×10^{-5}. Using these genes, we built a random forest classifier by XGBoost (v 2.1.1) [71] to distinguish between crypt and villus cells based on their expression profiles. Recognizing that the crypt–villus continuum is a continuous structure, we further examined the local neighborhood of each cell. We assumed that the distribution of normalized y-axis positions for the crypt and villus groups follows a mixed Gaussian distribution and used the resulting boundary to precisely define the crypt–villus interface at each location.

Pseudotime Analysis

We used Scanpy (v1.10.0) [14] to perform pseudotime analysis on a subset of cells, including Paneth and stem cells, TA cells, and enterocytes. First, we computed diffusion maps to capture the underlying cellular trajectories. One cell was randomly selected from the Paneth and stem cells group as the putative starting point, and pseudotime was computed using the diffusion pseudotime function (sc.tl.dpt) [37]. To better visualize the structure of this cell subset, we employed PAGA (sc.tl.paga) [38] and force-directed graph drawing (sc.tl.draw_graph) [39] to generate a layout on which the pseudotime values were mapped.

We then selected cells that were not part of the highly concentrated Peyer's patches and normalized both the pseudotime values and the normalized corrected y-axis to a scale from 0 to 1. Using pseudotime (0-1) as the independent variable and the normalized corrected y-axis (0-1) as the dependent variable, we performed linear regression analysis and computed the Pearson correlation coefficient to assess the strength and significance of their relationship.

Zonation Analysis

We selected only intact complete villi, including tip enterocytes for zonation analysis. The intestine was divided into small sections along the x-axis, and we applied a Kolmogorov–Smirnov test to evaluate the distribution of tip enterocytes in each section, following a uniform distribution. To ensure comprehensive coverage, multiple candidate sections from each area that passed the test were retained, and the final selection was refined through manual curation. We retained Paneth and stem cells, TA cells, and enterocytes. We then subdivided the villi into six distinct zones according to their normalized, corrected y-axis coordinates, ensuring that each zone contained an equivalent number of cells. Gene count were pseudobulked in each zone. and only genes with counts greater than 10 were retained (12,482 genes in total). We normalized the average gene expression of each gene across the zones to a maximum expression of 1. To identify the genes that were significantly zonated, we used pairwise differential expression analysis of each zone against all other zones using Scanpy (scanpy.tl.rank_genes_groups, t-test) [14]. We considered a gene as significantly zonated if its absolute log2-fold-change (LFC) was greater than 1.5 and the adjusted p-value was below 1×10^{-8} compared to all other zones.

KEGG Analysis

We used gprofiler-official (v1.0.0) [72] to analyze KEGG pathway enrichment for each zone. KEGG pathways were retained if any zone exhibited an adjusted p-value below 1×10^{-8} and an LFC of 1.5. We generated bubble plots of significant KEGG pathways across all zones, where the bubble size indicates the fraction of genes from that pathway that were significantly enriched in the zone, and the bubble color represents the corresponding p-value.

Gene Expression by Zone

We used the y-axis boundaries identified in zonation analysis for all plots of cell-type-specific gene expression by zones. Cells in the designated cluster were split by zone and pseudobulked. Plots show the average log expression of the gene in each zone.

We then sought to capture proximal–distal differential genes across different zones. Cells were first divided along the x-axis, with the first half designated as the

proximal group and the second half as the distal group. Only the subset of Paneth and stem cells, TA cells, and enterocytes was used, with each zone containing an approximately equal number of cells. For each zone, mitochondrial genes were filtered out, and only genes expressed in at least 30 cells were retained. Differential expression analysis was performed using an LFC threshold of 1.5 and an adjusted p-value cutoff of 0.001. The number of differential genes was then normalized by dividing by the total number of genes remaining after filtration. Finally, we generated plots of the differential gene count and the fraction of differential genes versus zones. KEGG pathways of the proximal and distal zone 6 genes were calculated if they exhibited an adjusted p-value below 0.001 and an LFC of 1.5. Line-plot trajectories were drawn for important marker genes across the zonation axis. Because stem, Paneth, and TA cells are predominantly confined to Zones 1–2, we displayed these three populations together with enterocytes for direct comparison. For every other lineage, the plot was restricted to cells of that specific type.

Transcription Factor Analysis

We selected Paneth cells, stem cells, TA cells, and enterocytes — each annotated with villus-zone information — from the previous section, and then followed the pySCENIC protocol (https://github.com/aertslab/SCENICprotocol) by applying `pyscenic` to infer TF–target gene regulatory networks and to identify and score regulons significantly enriched in our dataset by comparison with established reference databases [46].

A total of 192 regulons have been called by the results. For each regulon, we computed its mean activity score across the six spatial zones. We then visualized the normalized activity, scaled such that the maximum score for every regulon across the six zones equals 1, for 19 representative regulons in a heatmap. We generated violin plots showing the distribution of the corrected y-axis positions (μm) of the constituent genes for each regulon.

We selected zone-specific regulons (LFC > 0.3 and adjusted p-value $< 1 \times 10^{-10}$) and plotted the TF–target network for all six zones. For Zone 2, we built a TF–target network for the seven regulons whose activities differed significantly across zones by differential t-test from Scanpy. In the graph, large blue nodes denote the TF genes, small green nodes mark non-zonated targets, red nodes mark zonated targets, and nodes in additional colours highlight zonated targets enriched for specific functional categories. To probe zonation enrichment, we applied Fisher's exact tests to compare genes targeted by at least two TFs with all remaining targets, and to compare each TF — comparing its unique targets with the pool of all other target genes separately. We completed the same analysis for Zone 6 with five differential regulons.

Single-cell Analysis of Seiler et al [62]

We downloaded the dataset from GEO (GSE130113), which contains raw counts from three sham and two SBR replicates. We first filtered cells by requiring a minimum of 200 genes per cell, and filtered genes by retaining only those expressed in at least 3 cells. The data were then normalized to a total of 10,000 counts per cell and log-normalized. We selected highly variable genes, scaled the data, performed PCA, and applied Harmony for batch correction. UMAP was used for dimensionality reduction, and clustering was performed using the Leiden algorithm [68, 16]. Differential expression analysis was carried out using a t-test. Although the differential results were not exactly identical to those reported in the original paper, they were extremely similar—the top five genes were exactly the same, and all the top 10 genes

reported in the paper appeared as the top 11 in our analysis. Finally, we selected differential genes using a threshold of adjusted p-value less than 1×10^{-50} and an absolute LFC larger than 1, resulting in a total of 77 genes significantly differentially expressed in the SBR group.

We then mapped the differential genes to the spatial dataset, and 72 of them were expressed in the spatial data. Next, we sought to determine the specific zone in which each gene was predominantly expressed by defining a gene as expressed in a particular zone if its LFC in that zone was greater than 1.5 and its adjusted p-value was less than 1×10^{-8}. Finally, we plotted the zones in which all SBR differential genes were expressed in the spatial dataset.

Comparison Analysis for Results from Different Methods

We compared the results of $8\,\mu$m spot data, Bin2cell results, nuclei segmentation results, and SMURF results. For each dataset, we used the direct output of 4×4 spots (corresponding to an $8\,\mu$m $\times$ $8\,\mu$m spot size) as the $8\,\mu$m spot data. For Bin2cell, we followed the instructions provided in the Bin2cell GitHub repository [73]. Nuclei segmentation results were obtained using Squidpy (v1.4.1) [67] with Stardist (v0.8.5) [13].

In the analysis of the percentage of cells covered by nuclei, we encountered the issue that a single spot might encompass several nuclei, leading to multiple nuclei being assigned to a cell, an outcome that is not accurate. To address this, we defined a cell as covered by a nucleus only if at least 30% of the pixels in the cell region are occupied by that nucleus.

For the definition of nuclei data in the Mouse Brain dataset, we considered a spot to belong to a nucleus if at least 90% of the spot's counts were within that nucleus. We then uploaded the raw counts matrix of nuclei, SMURF, and Xenium results from 10x [23] to MapMyCells [28, 29] by selecting the "10x Whole Mouse Brain (CCN20230722)" as the reference dataset. Using the Hierarchical Mapping strategy available in MapMyCells, we obtained the cell type results for the three datasets. We calculated the KL divergence according to the "class_name" from the MapMyCells results. In the mouse small intestine dataset, we applied the same segmentation and clustering pipeline as provided by 10x Genomics [74].

Data Availability

The public mouse brain Visium HD and mouse small intestine datasets are available on the 10x Genomics website (mouse brain Visium HD, mouse intestine). Public mouse brain 10x Xenium probe-based data can be accessed on the 10x Genomics website (fresh-frozen mouse brain replicates). The public SBR experiment dataset from Seiler et al (2019). is available from GEO under accession number GSE130113. The ISH staining for gene Slc1a2 was obtained from the Allen Brain Atlas (https://mouse.brain-map.org/gene/show/20273).

Code Availability

The SMURF package is available at https://github.com/The-Mitra-Lab/SMURF. The code to reproduce the results presented in this study can be accessed at https://github.com/The-Mitra-Lab/SMURF_figures. The documentation for SMURF is available at https://the-mitra-lab.github.io/SMURF/.

Funding

This work has been funded by grants from National Institute of Mental Health (RF1MH117070, RF1MH126723, R01DA056829, and RM1MH138313 to J.D.D. and R.D.M.; R01MH124808 to J.D.D.; R01DE032865 to R.D.M.). S.S. was supported in part by the Autism Science Foundation (22-007). B.D.M. was supported by the Department of Veterans Affairs, Veterans Health Administration, Biomedical Laboratory Research and Development (1IK2BX004909) and the Washington University Digestive Disease Research Core Center (P30DK052574).

This manuscript is the result of funding in whole or in part by the National Institutes of Health (NIH). It is subject to the NIH Public Access Policy. Through acceptance of this federal funding, NIH has been given a right to make this manuscript publicly available in PubMed Central upon the Official Date of Publication, as defined by NIH.

Acknowledgements

We thank Hao Fan for discussions that helped shape this project, and Jennifer Hayden for scientific illustrations. The contents of this manuscript do not represent the views of the Department of Veterans Affairs or the United States Government.

Figures

a

H&E / DAPI Staining

① Nuclei Segmentation

Create Nearest Neighbor Network

Assign Nuclei to Clusters

③ Deconvolution

Transcripts: Spots x Genes

Transcripts: Nuclei x Genes

UMAP

Cluster Nuclei

② Expand Nuclei to Cells using Cosine Similarity

Reclustering Cells

b Cell Expansion

Nucleus 1
Nucleus 2

Assigned by Nucleus 1
Assigned by Nucleus 2
Assigned by Nuclei 1 & 2
Rejected by Nuclei 1 & 2

c Count Deconvolution for Shared Spots

Fig. 1: **SMURF uses soft-segmentation to reconstruct single-cell transcriptomes.** (a) Schematic of SMURF's soft-segmentation algorithm. SMURF segments nuclei from the image data, assigns transcripts to nuclei from overlapping capture spots and clusters the resulting nuclei-by-genes matrix (Step 1). The algorithm then iteratively expands cell boundaries to assign transcripts from capture spots that do not overlap nuclei to cells (Step 2). Finally, SMURF leverages deep learning to deconvolve spots assigned to multiple cells during the expansion step (Step 3). (b) Example of capture spots assignment to cells. Capture spots can be assigned to zero, one, or multiple nuclei during cell expansion (Step 2), depending on their proximity and expression similarity. (c) Count deconvolution for shared spots. After the cell expansion step, capture spots may be shared between multiple cells (purple squares in the first two panels). Transcripts from shared capture spots are divided among cells using a quadratic programming or deep learning algorithm (see Methods).

Fig. 2: **SMURF accurately assigns transcripts to cells in mouse brain and mouse small intestine data.** (a,b) Representative images illustrating how SMURF and other ST segmentation methods aggregate Visium HD capture spots to define cells in the mouse brain (top row) and mouse small intestine (bottom row). Capture spots are shaded according to their assigned cell. SMURF uniquely manages to both assign only one nucleus per cell and analyze captures spots under cell bodies. (c) Percentage of cells without an associated nucleus or containing multiple nuclei in the mouse brain (left panel) and mouse small intestine (right panel) using different ST segmentation strategies. (d) Median number of UMIs assigned to cells, total cells detected, and median cell size using SMURF or nuclei-based segmentation. (e) Cell type detection in mouse small intestine using clusters generated by SMURF or nuclei-segmentation. Expression of marker genes for four intestinal cell types is plotted across clusters. SMURF but not nuclei segmentation has clusters that clearly correspond to each cell type. Abbreviations: UMI, Unique Molecular Identifier.

Fig. 3: **Manifold unrolling of Visium HD data in cortex and small intestine.**
(a) Selection of the X-axis contour in the mouse cortex based on positions of cells in
the basement membrane cluster. (b) Overlay of cortical manifold onto mouse brain
section, colored by X-axis coordinates (µm). (c) Overlay of cortical manifold onto
mouse brain section, colored by Y-axis coordinates (µm). (d) Heatmap of marker
genes for cortical layers across different Y-bins of the mouse brain manifold. (e)
Distribution of cortical layer-specific neurons in the unrolled rectilinear coordinate
system. (f) Schematic illustration of the mouse small intestine and its "Swiss roll"
histological orientation. (g) Overlay of mouse small intestine data, colored by unrolled
X-axis coordinates based on cells in the muscle layer (µm). (h) Expression of selected
proximal and distal enriched genes in the original ST image and histogram plot of
those genes on the unrolled X-axis of the mouse small intestine. (*Lct, Slc10a2(Ibat)*).
(i) Illustration of the crypt-villus axis in the mouse small intestine. (j) Schematic
of four common villus structures encountered during construction of the Y-axis for
mouse small intestine data. (k) Overlay of intestinal manifold in selected villi with
cells colored by normalized, corrected Y-axis. (l) Cell type distribution along the
corrected Y-axis. Red dashed line indicates the approximate boundary between crypt
and villus regions.

Fig. 4: **Temporal and functional zonation of transcriptional programs on the ileal crypt-villus axis.** (a) Schematic illustrating the epithelial differentiation trajectory in intestinal villi from proliferative stem cells (bottom) to differentiated enterocytes (top). (b) Concordance of pseudotime trajectory and corrected y-axis position during enterocyte maturation. PAGA embedding of intestinal stem and Paneth cells, TA, and enterocytes, colored by pseudotime position (left panel), cell type (center panel), and corrected y-axis position (right panel). (c) Scatter plot comparing the normalized pseudotime progression score and the normalized corrected Y position. Each point represents a single cell. The dashed red line indicates a linear fit (Pearson correlation = 0.79). (d) Example of the discrete binning of cells of the intestinal villus into 6 zones along the y-axis. (e) Heatmap of normalized gene expression across zones. Gene expression in all stem and Paneth cells, TA, and enterocytes is included. The heatmap displays normalized expression of each gene (columns) in each zone (rows). (f) Bubble plot displaying KEGG pathway enrichment of significantly zonated genes. (g-j) Zonation profiles of selected genes in enterocytes (g), goblet cells (h), tuft cells (i), and EEC (j).

Fig. 5: **Spatially resolved transcription-factor programs along the crypt-villus axis.** (a) Heatmap of normalized regulon activity scores for 20 key transcription factors (TFs) across the six villus zones. Regulon scores calculated with Scenic. (b) Violin plots of number of cells expressing a given TFs, plotted against the unrolled corrected Y-axis coordinate (μm). Black bars indicate the median y-axis position of TF-expressing cells and the inter-quartile range; violin width reflects local cell density.

(c) Integrated TF target networks for Zone 2. TFs are depicted in blue circles. Lines connect TFs to their gene targets. Selected target genes are color-coded by their predominant functional category. *: 500 of the 1362 Elf3 targets are shown for legibility.

(d) Integrated TF target networks for Zone 6. Coloring as in panel c.

Fig. 6: **Regionally specific gene programs along the proximal-distal intestinal axis accumulate in the villus tip.** (a) Number (top) and fraction (bottom) of genes that are differentially expressed in each zone, comparing proximal and distal regions of villi. Only genes expressed in more than 30 cells are considered. (b) Expression of proximal-tip enriched genes (*Apoa4* and *Apoc3*). Data shows expression in the original ST data, as well as expression along the unwrapped x and y axes. (c) Expression of distal-tip enriched gene (*Slc6a14*. Data shows expression in the original ST data, as well as expression along the unwrapped x and y axes. (d) KEGG pathway enrichment for proximal and distal tip-associated genes. (e) Schematic of small bowel resection and scRNA-seq dataset (Seiler et al [62]). (f) Tip cells, but not cells at the base of crypts, are reprogrammed one week after small bowel resection. For each zone, the number and identity of differentially expressed genes from the single-cell data are shown.

909 # Extended Data Figures

Extended Data Figure 1: **Soft segmentation with SMURF** (a) Illustration of the data imbalance issue when the standard normalized 8 μm Spot method is used. The spatial expression pattern of Slc1a2 is discordant between Visium HD mouse brain data analyzed by the 8 μm Spot method and ISH data from the Allen Brain Atlas. In contrast, when analyzed by SMURF, there is concordance. (b) Cell type proportions annotated by MapMyCells in probe-based mouse brain data, comparing nuclei results and SMURF results. KL divergence values quantify differences in cell type distributions. Lower KL divergence indicates better concordance.

Extended Data Figure 2: Distribution of cortical layer-specific non-neuronal cells in the unrolled rectilinear coordinate system

Extended Data Figure 3: **Mouse small intestine analysis.** (a) UMAP embedding of intestinal cells, colored by annotated cell types. (b) Correlation between single-cell transcriptomes from Zwick et al. [7] and spatial transcriptomic profiles across intestinal sections. (c) Overlay of selected mouse small intestine data with cells colored by coordinate value along the Y-axis at different stages of "unrolling": raw layer positioning, corrected Y (μm), and normalized corrected Y. (d) Marker gene expression visualized in the selected unrolled intestinal tissue (*Olfm4*, *Krt19*, *Fabp2*, and *Ada*). (e) Normalized corrected Y-axis distribution of cells expressing selected genes with red dashed lines indicating the approximate boundary between crypt and villus regions.

Extended Data Figure 4: **Different Distances Examples Mapped on Real H&E Plot.** (a) Layer. (b) Corrected Distance. (c) Normalized Corrected Distance. (d) Zone.

Extended Data Figure 5: **Highly zonated enterocyte genes.** Statistically enriched genes were identified in each zone. These lists were manually inspected to select the most illustrative examples of zonated genes. Heatmaps depict normalized gene expression for Zone 1 (a), Zone 2 (b), Zone 3 (c), Zone 4 (d), Zone 5 (e), and Zone 6 (f).

Extended Data Figure 6: Transcription-factor (TF) interaction plots in each zone. Regulons are shown only when the $\log_2$ fold change exceeds 0.3 and the adjusted p-value is below 1×10^{-10}.

Extended Data Figure 7: **Heatmap of villus tip and bottom marker genes from Moor et al. [2] in ileal SMURF dataset.** (a) Heatmap of villus with tip marker genes from Moor et al. (b) Heatmap of villus with bottom marker genes from Moor et al.

References

[1] R.F. Hagemann, C.P. Sigdestad, and S. Lesher. A quantitative description of the intestinal epithelium of the mouse. *American Journal of Anatomy*, 129(1):41–52, 1970.

[2] Andreas E Moor, Yotam Harnik, Shani Ben-Moshe, Efi E Massasa, Milena Rozenberg, Raya Eilam, Keren Bahar Halpern, and Shalev Itzkovitz. Spatial reconstruction of single enterocytes uncovers broad zonation along the intestinal villus axis. *Cell*, 175(4):1156–1167, 2018.

[3] Rita Manco, Inna Averbukh, Ziv Porat, Keren Bahar Halpern, Ido Amit, and Shalev Itzkovitz. Clump sequencing exposes the spatial expression programs of intestinal secretory cells. *Nature communications*, 12(1):3074, 2021.

[4] Yotam Harnik, Lisa Buchauer, Shani Ben-Moshe, Inna Averbukh, Yishai Levin, Alon Savidor, Raya Eilam, Andreas E Moor, and Shalev Itzkovitz. Spatial discordances between mrnas and proteins in the intestinal epithelium. *Nature metabolism*, 3(12):1680–1693, 2021.

[5] Toufic Mayassi, Chenhao Li, Åsa Segerstolpe, Eric M Brown, Rebecca Weisberg, Toru Nakata, Hiroshi Yano, Paula Herbst, David Artis, Daniel B Graham, et al. Spatially restricted immune and microbiota-driven adaptation of the gut. *Nature*, pages 1–10, 2024.

[6] Altay Yuzeir, David Alejandro Bejarano, Stephan Grein, Jan Hasenauer, Andreas Schlitzer, and Jiangyan Yu. Intestline: a shiny-based application to map the rolled intestinal tissue onto a line. *Bioinformatics*, 39(4):btad140, 2023.

[7] Rachel K Zwick, Petr Kasparek, Brisa Palikuqi, Sara Viragova, Laura Weichselbaum, Christopher S McGinnis, Kara L McKinley, Asoka Rathnayake, Dedeepya Vaka, Vinh Nguyen, et al. Epithelial zonation along the mouse and human small intestine defines five discrete metabolic domains. *Nature Cell Biology*, 26(2):250–262, 2024.

[8] Hao Chen, Dongshunyi Li, and Ziv Bar-Joseph. Scs: cell segmentation for high-resolution spatial transcriptomics. *Nature methods*, 20(8):1237–1243, 2023.

[9] Krzysztof Polański, Raquel Bartolomé-Casado, Ioannis Sarropoulos, Chuan Xu, Nick England, Frode L Jahnsen, Sarah A Teichmann, and Nadav Yayon. Bin2cell reconstructs cells from high resolution visium hd data. *Bioinformatics*, 40(9):btae546, 2024.

[10] Carsen Stringer, Tim Wang, Michalis Michaelos, and Marius Pachitariu. Cellpose: a generalist algorithm for cellular segmentation. *Nature methods*, 18(1):100–106, 2021.

[11] Uwe Schmidt, Martin Weigert, Coleman Broaddus, and Gene Myers. Cell detection with star-convex polygons. In *Medical Image Computing and Computer Assisted Intervention - MICCAI 2018 - 21st International Conference, Granada, Spain, September 16-20, 2018, Proceedings, Part II*, pages 265–273, 2018.

[12] Martin Weigert, Uwe Schmidt, Robert Haase, Ko Sugawara, and Gene Myers. Star-convex polyhedra for 3d object detection and segmentation in microscopy. In *The IEEE Winter Conference on Applications of Computer Vision (WACV)*, March 2020.

[13] Martin Weigert and Uwe Schmidt. Nuclei instance segmentation and classification in histopathology images with stardist. In *The IEEE International Symposium on Biomedical Imaging Challenges (ISBIC)*, 2022.

[14] F Alexander Wolf, Philipp Angerer, and Fabian J Theis. Scanpy: large-scale single-cell gene expression data analysis. *Genome biology*, 19:1–5, 2018.

[15] Lukas Heumos, Anna C Schaar, Christopher Lance, Anastasia Litinetskaya, Felix Drost, Luke Zappia, Malte D Lücken, Daniel C Strobl, Juan Henao, Fabiola Curion, et al. Best practices for single-cell analysis across modalities. *Nature Reviews Genetics*, 24(8):550–572, 2023.

[16] Vincent A Traag, Ludo Waltman, and Nees Jan Van Eck. From louvain to leiden: guaranteeing well-connected communities. *Scientific reports*, 9(1):1–12, 2019.

[17] John T Chamberlin, Younghee Lee, Gabor T Marth, and Aaron R Quinlan. Differences in molecular sampling and data processing explain variation among single-cell and single-nucleus rna-seq experiments. *Genome Research*, 34(2):179–188, 2024.

[18] Aaron F McDaid, Derek Greene, and Neil Hurley. Normalized mutual information to evaluate overlapping community finding algorithms. *arXiv preprint arXiv:1110.2515*, 2011.

[19] Stephen Boyd and Lieven Vandenberghe. *Convex Optimization*. Cambridge University Press, Cambridge, UK, 2004.

[20] Tianlong Chen, Xiaohan Chen, Wuyang Chen, Howard Heaton, Jialin Liu, Zhangyang Wang, and Wotao Yin. Learning to optimize: A primer and a benchmark. *Journal of Machine Learning Research*, 23(189):1–59, 2022.

[21] 10x Genomics. Visium hd cytassist gene expression libraries of mouse brain he. https://www.10xgenomics.com/datasets/visium-hd-cytassist-gene-expression-libraries-of-mouse-brain-he, 2024. Accessed: 2025-01-01.

[22] 10x Genomics. Visium hd cytassist gene expression libraries of mouse intestine. https://www.10xgenomics.com/datasets/visium-hd-cytassist-gene-expression-libraries-of-mouse-intestine, 2024. Accessed: 2025-01-01.

[23] 10x Genomics. Fresh frozen mouse brain replicates: In situ gene expression dataset (3 replicates). https://www.10xgenomics.com/datasets/fresh-frozen-mouse-brain-replicates-1-standard, 2023. Analyzed using Xenium Onboard Analysis 1.0.2; dataset comprises 3 replicates; licensed under CC BY 4.0. Accessed: 16 February 2025.

[24] Solomon Kullback and Richard A Leibler. On information and sufficiency. *The annals of mathematical statistics*, 22(1):79–86, 1951.

[25] Lingling Xu, Abhijit Nirwane, and Yao Yao. Basement membrane and blood–brain barrier. *Stroke and vascular neurology*, 4(2), 2019.

[26] Sam T. Roweis and Lawrence K. Saul. Nonlinear dimensionality reduction by locally linear embedding. *Science*, 290(5500):2323–2326, 2000.

[27] Bosiljka Tasic, Vilas Menon, Thuc Nghi Nguyen, Tae Kyung Kim, Tim Jarsky, Zizhen Yao, Boaz Levi, Lucas T Gray, Staci A Sorensen, Tim Dolbeare, et al. Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nature neuroscience*, 19(2):335–346, 2016.

[28] Allen Institute for Brain Science. Mapmycells (rrid:scr_024672). https://portal.brain-map.org/atlases-and-data/bkp/mapmycells. Accessed: 16 February 2025.

[29] Zizhen Yao, Cindy TJ van Velthoven, Michael Kunst, Meng Zhang, Delissa McMillen, Changkyu Lee, Won Jung, Jeff Goldy, Aliya Abdelhak, Matthew Aitken, et al. A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain. *Nature*, 624(7991):317–332, 2023.

[30] Pasko Rakic. Evolution of the neocortex: a perspective from developmental biology. *Nature Reviews Neuroscience*, 10(10):724–735, 2009.

[31] Maria Antonietta Tosches and Gilles Laurent. Evolution of neuronal identity in the cerebral cortex. *Current opinion in neurobiology*, 56:199–208, 2019.

[32] Antonio Pinto-Duarte, Terrence J Sejnowski, and Shyam Srinivasan. A conserved principle of glial organization in the paleocortex and neocortex. 2025.

[33] Riikka Pietilä, Francesca Del Gaudio, Liqun He, Elisa Vázquez-Liébanas, Michael Vanlandewijck, Lars Muhl, Giuseppe Mocci, Katrine D Bjørnholm, Caroline Lindblad, Alexander Fletcher-Sandersjöö, et al. Molecular anatomy of adult mouse leptomeninges. *Neuron*, 111(23):3745–3764, 2023.

[34] Matthew Bjerknes and Hazel Cheng. The stem-cell zone of the small intestinal epithelium. iii. evidence from columnar, enteroendocrine, and mucous cells in the adult mouse. *American Journal of Anatomy*, 160(1):77–91, 1981.

[35] Yoshitatsu Sei, Xinping Lu, Alice Liou, Xilin Zhao, and Stephen A Wank. A stem cell marker-expressing subset of enteroendocrine cells resides at the crypt base in the small intestine. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, 300(2):G345–G356, 2011.

[36] Nicholas Barker, Johanna H van Es, Johannes Kuipers, Päivi Kujala, Marijke van den Born, M Cozijnsen, A Haegebarth, Joost Korving, H Begthel, Peter J Peters, et al. Identification of stem cells in the small intestine and colon by marker gene lgr5. *Nature*, 449(7165):1003–1007, 2007.

[37] Laleh Haghverdi, Florian Buettner, and Fabian J Theis. Diffusion maps for high-dimensional single-cell analysis of differentiation data. *Bioinformatics*, 31(18):2989–2998, 2015.

[38] F Alexander Wolf, Fiona K Hamey, Mireya Plass, Jordi Solana, Joakim S Dahlin, Berthold Göttgens, Nikolaus Rajewsky, Lukas Simon, and Fabian J Theis. Paga: graph abstraction reconciles clustering with trajectory inference through a topology preserving map of single cells. *Genome biology*, 20:1–9, 2019.

[39] Mathieu Jacomy, Tommaso Venturini, Sebastien Heymann, and Mathieu Bastian. Forceatlas2, a continuous graph layout algorithm for handy network visualization designed for the gephi software. *PloS one*, 9(6):e98679, 2014.

[40] Lulu Huang, Jochem H Bernink, Amir Giladi, Daniel Krueger, Gijs JF van Son, Maarten H Geurts, Georg Busslinger, Lin Lin, Harry Begthel, Maurice Zandvliet, et al. Tuft cells act as regenerative stem cells in the human intestine. *Nature*, 634(8035):929–935, 2024.

[41] Michael R Howitt, Sydney Lavoie, Monia Michaud, Arthur M Blum, Sara V Tran, Joel V Weinstock, Carey Ann Gallini, Kevin Redding, Robert F Margolskee, Lisa C Osborne, et al. Tuft cells, taste-chemosensory cells, orchestrate parasite type 2 immunity in the gut. *Science*, 351(6279):1329–1333, 2016.

[42] Christoph Schneider, Claire E O'Leary, Jakob von Moltke, Hong-Erh Liang, Qi Yan Ang, Peter J Turnbaugh, Sridhar Radhakrishnan, Michael Pellizzon, Averil Ma, and Richard M Locksley. A metabolite-triggered tuft cell-ilc2 circuit drives small intestinal remodeling. *Cell*, 174(2):271–284, 2018.

[43] Julie Bas, Philippe Jay, and François Gerbe. Intestinal tuft cells: Sentinels, what else? In *Seminars in cell & developmental biology*, volume 150, pages 35–42. Elsevier, 2023.

[44] Keven A Roth and Jeffrey I Gordon. Spatial differentiation of the intestinal epithelium: analysis of enteroendocrine cells containing immunoreactive serotonin, secretin, and substance p in normal and transgenic mice. *Proceedings of the National Academy of Sciences*, 87(16):6408–6412, 1990.

[45] Joep Beumer, Benedetta Artegiani, Yorick Post, Frank Reimann, Fiona Gribble, Thuc Nghi Nguyen, Hongkui Zeng, Maaike Van den Born, Johan H Van Es, and Hans Clevers. Enteroendocrine cells switch hormone expression along the crypt-to-villus bmp signalling gradient. *Nature cell biology*, 20(8):909–916, 2018.

[46] Sara Aibar, Carlos B. González-Blas, Thomas Moerman, Vân A. Huynh-Thu, Hana Imrichova, Geert Hulselmans, Florian Rambow, José C. Marine, Pierre Geurts, and Stein Aerts. SCENIC: single-cell regulatory network inference and clustering. *Nature Methods*, 14(11):1083–1086, 2017.

[47] Philippe Blache, Marc Van De Wetering, Isabelle Duluc, Claire Domon, Philippe Berta, Jean-Noël Freund, Hans Clevers, and Philippe Jay. Sox9 is an intestine crypt transcription factor, is regulated by the wnt pathway, and represses the cdx2 and muc2 genes. *Journal of Cell Biology*, 166(1):37–47, 2004.

[48] Eric J Formeister, Ayn L Sionas, David K Lorance, Carey L Barkley, Ginny H Lee, and Scott T Magness. Distinct sox9 levels differentially mark stem/progenitor populations and enteroendocrine cells of the small intestine epithelium. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, 296(5):G1108–G1118, 2009.

[49] Emily M Walker, Cayla A Thompson, and Michele A Battle. Gata4 and gata6 regulate intestinal epithelial cytodifferentiation during development. *Developmental biology*, 392(2):283–294, 2014.

[50] Ian Y Luk, Camilla M Reehorst, and John M Mariadason. Elf3, elf5, ehf and spdef transcription factors in tissue homeostasis and cancer. *Molecules*, 23(9):2191, 2018.

[51] JL Wang, ZF Chen, HM Chen, MY Wang, X Kong, YC Wang, TT Sun, J Hong, W Zou, J Xu, et al. Elf3 drives β-catenin transactivation and associates with poor prognosis in colorectal cancer. *Cell death & disease*, 5(5):e1263–e1263, 2014.

[52] Kristin N Bell and Noah F Shroyer. Krüpple-like factor 5 is required for proper maintenance of adult intestinal crypt cellular proliferation. *Digestive diseases and sciences*, 60:86–100, 2015.

[53] Chang-Kyung Kim, Madhurima Saxena, Kasmika Maharjan, Jane J Song, Kenneth R Shroyer, Agnieszka B Bialkowska, Ramesh A Shivdasani, and Vincent W Yang. Krüppel-like factor 5 regulates stemness, lineage specification, and regeneration of intestinal epithelial stem cells. *Cellular and Molecular Gastroenterology and Hepatology*, 9(4):587–609, 2020.

[54] Keren Bahar Halpern, Hassan Massalha, Rachel K Zwick, Andreas E Moor, David Castillo-Azofeifa, Milena Rozenberg, Lydia Farack, Adi Egozi, Dan R Miller, Inna Averbukh, et al. Lgr5+ telocytes are a signaling source at the intestinal villus tip. *Nature communications*, 11(1):1936, 2020.

[55] Joep Beumer, Jens Puschhof, Fjodor Yousef Yengej, Lianzheng Zhao, Adriana Martinez-Silgado, Marloes Blotenburg, Harry Begthel, Charelle Boot, Alexander van Oudenaarden, Ye-Guang Chen, et al. Bmp gradient along the intestinal villus axis controls zonated enterocyte and goblet cell states. *Cell Reports*, 38(9), 2022.

[56] Adrianna K San Roman and Ramesh A Shivdasani. Boundaries, junctions and transitions in the gastrointestinal tract. *Experimental cell research*, 317(19):2711–2718, 2011.

[57] Cayla A Thompson, Ann DeLaForest, and Michele A Battle. Patterning the gastrointestinal epithelium to confer regional-specific functions. *Developmental biology*, 435(2):97–108, 2018.

[58] Alyssa S Zembroski, Theresa D'Aquila, and Kimberly K Buhman. Characterization of cytoplasmic lipid droplets in each region of the small intestine of lean and diet-induced obese mice in response to dietary fat. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, 321(1):G75–G86, 2021.

[59] Catriona MH Anderson, Vadivel Ganapathy, and David T Thwaites. Human solute carrier slc6a14 is the β-alanine carrier. *The Journal of physiology*, 586(17):4061–4067, 2008.

[60] Sabine Middendorp, Kerstin Schneeberger, Caroline L Wiegerinck, Michal Mokry, Ronald DL Akkerman, Simone Wijngaarden, Hans Clevers, and Edward ES Nieuwenhuis. Adult stem cells in the small intestine are intrinsically programmed with their location-specific function. *Stem cells*, 32(5):1083–1091, 2014.

[61] Ozge Kayisoglu, Franziska Weiss, Carolin Niklas, Isabella Pierotti, Malvika Pompaiah, Nina Wallaschek, Christoph-Thomas Germer, Armin Wiegering, and Sina Bartfeld. Location-specific cell identity rather than exposure to gi microbiota defines many innate immune signalling cascades in the gut epithelium. *Gut*, 70(4):687–697, 2021.

[62] Kristen M Seiler, Sarah E Waye, Wenjun Kong, Kenji Kamimoto, Adam Bajinting, William H Goo, Emily J Onufer, Cathleen Courtney, Jun Guo, Brad W Warner, et al. Single-cell analysis reveals regional reprogramming during adaptation to massive small bowel resection in mice. *Cellular and molecular gastroenterology and hepatology*, 8(3):407–426, 2019.

[63] Sara M Parigi, Ludvig Larsson, Srustidhar Das, Ricardo O Ramirez Flores, Annika Frede, Kumar P Tripathi, Oscar E Diaz, Katja Selin, Rodrigo A Morales, Xinxin Luo, et al. The spatial transcriptomic landscape of the healing mouse intestine following damage. *Nature communications*, 13(1):828, 2022.

[64] Adam Paszke, Sam Gross, Francisco Massa, Adam Lerer, James Bradbury, Gregory Chanan, Trevor Killeen, Zeming Lin, Natalia Gimelshein, Luca Antiga, et al. Pytorch: An imperative style, high-performance deep learning library. *Advances in neural information processing systems*, 32, 2019.

[65] Carsen Stringer and Marius Pachitariu. Cellpose3: one-click image restoration for improved cellular segmentation. *Nature Methods*, pages 1–8, 2025.

[66] Thomas Cover and Peter Hart. Nearest neighbor pattern classification. *IEEE transactions on information theory*, 13(1):21–27, 1967.

[67] Giovanni Palla, Hannah Spitzer, Michal Klein, David Fischer, Anna Christina Schaar, Louis Benedikt Kuemmerle, Sergei Rybakov, Ignacio L Ibarra, Olle Holmberg, Isaac Virshup, et al. Squidpy: a scalable framework for spatial omics analysis. *Nature methods*, 19(2):171–178, 2022.

[68] Etienne Becht, Leland McInnes, John Healy, Charles-Antoine Dutertre, Immanuel WH Kwok, Lai Guan Ng, Florent Ginhoux, and Evan W Newell. Dimensionality reduction for visualizing single-cell data using umap. *Nature biotechnology*, 37(1):38–44, 2019.

[69] Cynthia Arroyo Portilla, Romain Fenouil, Camille Wagner, Cécilia Luciani, Margaux Lagier, Clément Da Silva, Fanny Hidalgo-Villeda, Lionel Spinelli, Mathieu Fallet, Julie Tomas, et al. Peyer's patch phagocytes acquire specific transcriptional programs that influence their maturation and activation profiles. *Mucosal Immunology*, 16(4):527–547, 2023.

[70] Eugene Drokhlyansky, Christopher S Smillie, Nicholas Van Wittenberghe, Maria Ericsson, Gabriel K Griffin, Gokcen Eraslan, Danielle Dionne, Michael S Cuoco, Max N Goder-Reiser, Tatyana Sharova, et al. The human and mouse enteric nervous system at single-cell resolution. *Cell*, 182(6):1606–1622, 2020.

[71] Tianqi Chen and Carlos Guestrin. Xgboost: A scalable tree boosting system. In *Proceedings of the 22nd acm sigkdd international conference on knowledge discovery and data mining*, pages 785–794, 2016.

[72] Uku Raudvere, Liis Kolberg, Ivan Kuzmin, Tambet Arak, Priit Adler, Hedi Peterson, and Jaak Vilo. g: Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic acids research*, 47(W1):W191–W198, 2019.

[73] Teichlab. bin2cell. https://github.com/Teichlab/bin2cell, n.d. Accessed: September 14, 2024.

[74] 10x Genomics. Segmentation for visium hd. https://www.10xgenomics.com/analysis-guides/segmentation-visium-hd. Analysis Guide. Accessed: 16 February 2025.