Detection of a hybrid PrPfr state in the dark reversion of a bathy phytochrome indicates inter-dimer allostery.

Sayan Prodhan[1], Petra Mészáros[1], Szabolcs Bódizs[1], Yalin Zhou[1], Micha Maj[2], and Sebastian Westenhoff[1,*]

[1]Department of Chemistry - BMC, Biochemistry, Uppsala University, 75123 Uppsala, Sweden
[2]Department of Chemistry—Ångström Laboratory, Uppsala University, Uppsala, Sweden
[*]Corresponding author: sebastian.westenhoff@kemi.uu.se

ABSTRACT

Phytochromes are photosensor proteins which detect light in plants, fungi, and bacteria. They pho-
toswitch between red light absorbing (Pr) and far-red light absorbing (Pfr) states, however, ther-
mal reversion in the dark is an equally important factor in controlling their signaling levels. Phy-
tochromes are generally dimeric proteins, and mixed PrPfr states are therefore possible. These
states have been implied in the dark reversion studies of plant phytochromes, but not in bacterial
phytochromes. Here, we investigate the dark reversion kinetics of the 'bathy' phytochrome from *P.*
***aeruginosa* (*Pa*BphP) using UV-Vis absorption spectroscopy. A single set of time-resolved spectra**
does not conclusively reveal the presence of a mixed PrPfr state, as both a direct $\mathrm{Pr} \to \mathrm{Pfr}$ model or
a sequential $\mathrm{Pr} \to \mathrm{PrPfr} \to \mathrm{Pfr}$ model fit the spectral kinetics. However, a systematic analysis of dark
reversion kinetics with varying Pr/Pfr ratios can only be satisfactorily fit by the sequential model,
which indicates the presence of an intermediate PrPfr state. A newly designed monomeric vari-
ant of *Pa*BphP provides strong support for this interpretation. Temperature-dependent kinetics
revealed similarly low activation energies for the dark reversion processes of both proteins, consis-
tent with a previously proposed keto-enol tautomerization preceding dark reversion. Interestingly,
our results suggest allosteric regulation of dark reversion across the dimer, which we propose to
be a contributing factor in phytochrome signaling.

INTRODUCTION

Many organisms rely on photoreception to adapt to changing environmental conditions[1,2]. Among vari-
ous light-sensing systems, phytochromes are red light-detecting photoreceptors that regulate physiolog-
ical responses in plants, fungi, and bacteria[3–5]. In plants, they control essential processes like seed germi-
nation, chloroplast development, and flowering[6–8], while in bacteria, they regulate pigmentation, motil-
ity, and biofilm formation[9–11]. Therefore, the phytochrome photocycle is extensively studied for biotech-
nological applications, as well as for the engineering of optogenetic tools and near-infrared fluorescent
markers used in microscopy and cell biology[12,13].

Phytochromes are typically homodimeric proteins with a modular structure[14]. Each monomer consists

of an N-terminal photosensory core module (PCM), responsible for light sensing, and a C-terminal output

module (OPM), which mediates signal transduction. In some cases, additional N-terminal extensions may

also contribute to the effector function of the protein. The PCM is composed of three functionally distinct

domains: PAS (Per-ARNT-Sim)[15], GAF (cGMP phosphodiesterase / adenylate cyclase / FhlA)[16], and PHY

(phytochrome-specific domain), and contains a bilin cofactor that serves as the chromophore[17].

Phytochromes interconvert between two structurally distinct states: the red light-absorbing Pr state (

$\lambda_{max} \approx 700$ nm, for bacterial phytochromes) and the far-red-light-absorbing Pfr state ($\lambda_{max} \approx 750$ nm).

These states are associated with distinct biochemical properties and, consequently, different signaling

activities[18,19]. Based on the relative free energies of these states, phytochromes are classified as either

canonical, the most common type with a thermally stable Pr state, or reverse-activated (bathy), in which

the Pfr state is thermodynamically favored (Fig.1(a))[20–22]. In the absence of light, phytochromes ther-

mally revert to their resting state.

Canonical and bathy phytochromes have been extensively studied using spectroscopy and structural bi-

ology methods. Most of these studies focus on understanding the sequence of events during photoac-

tivation[23–28]. They show that upon photoactivation, the chromophore undergoes isomerization around

the C15=C16 bond, followed by a series of structural changes through intermediate states (i.e., Lumi-F/R

and Meta-F/R).[29–31]. Crystal, NMR, and cryo-EM structures of phytochromes in both the Pr and Pfr states

have been solved[32–37]. Of particular interest are two recent crystal and cryo-EM structures, which reveal

phytochromes in mixed Pr/Pfr heterodimers[38,39]. However, it remains unclear how these intermediate

states are formed. The dark reversion has been studied in both plant and bacterial phytochromes[19,25,28,29,35,37,40–56]

with studies on plant phytochromes providing the most detailed insights, including differences in activa-

tion energy across various temperature ranges[40], the presence of PrPfr hybrid states[54], and temperature-

dependent kinetics that allow phytochromes to function as temperature sensors[19,50]. Plants also have

multiple isoforms of phytochromes, which contribute to varying dark reversion rates, allowing for more

complex adaptation to different environmental conditions[43].

In bacterial phytochromes, the dark reversion process is less well understood. While dark reversion is com-

monly used as a standard assay in structural and biochemical studies of phytochromes, only a few de-

tailed studies have been conducted[47,51,52,57–59]. However, a hybrid PrPfr state has not been observed as

part of the dark reversion process in bacterial phytochromes yet.

In this study, we systematically investigate the dark reversion kinetics of a bathy phytochrome from *Pseu-*

domonas aeruginosa (*Pa*BphP). By comparing the wild-type full-length dimer (FL-*Pa*BphP-D) with a

newly designed monomeric variant (PSM-*Pa*BphP-M), we provide clear evidence for the presence of PrPfr

states as intermediates in the dark reversion process for the wild-type protein. This finding suggests a

novel allosteric mechanism between the two protomers.

Figure 1: The phytochrome photoswitches between Pr and Pfr states, and chromatography confirms the oligomeric states of the two protein samples. (a) Conformational differences in the biliverdin structure distinguish the Pfr ($\approx 750nm$) and Pr ($\approx 700nm$) states, which are reflected in their steady-state absorption spectra.(b) Preparatory SEC analysis of FL-*Pa*BphP-D and PSM-*Pa*BphP-M revealing their oligomerization state. The shaded region indicates the fraction collected fractions for FL-*Pa*BphP-D. For PSM-*Pa*BphP-M, the collection zone was around the peak maximum. (c) Analytical SEC of FL-*Pa*BphP-D in Pr and Pfr state confirms that the protein is predominatly dimeric. The slight shift in the retention time between the states likely reflects conformational changes.

MATERIALS AND METHODS

Recombinant protein production

The plasmid used for the recombinant expression of the full-length *P. aeruginosa* phytochrome contained the *Pa*BphP insert (GenBank: AAG07504.1) with a C-terminal His_6-tag and a pET28a(+) backbone (Novagen). The plasmid used for the recombinant expression of the *P. aeruginosa* phytochrome PAS-GAF-PHY mutant fragment contained the *Pa*BphP insert (GenBank: AAG07504.1) with a C-terminal His6-tag and a pET24a(+) plasmid backbone (GenScript). To facilitate biliverdin incorporation during expression, all the phytochrome constructs were co-expressed with a heme oxygenase plasmid, kindly provided by Prof. Janne A. Ihalainen. The plasmids were transformed into BL21(DE3) *Escherichia coli* strain for expression.

Cells were grown in LB medium supplemented with 50 μg/mL kanamycin and 34.5 μg/mL chloramphenicol at $37°C$ until reaching an OD_{600} of 0.8. Expression of *Pa*BphP and heme oxygenase was induced by the addition of 1 mM IPTG and 1 mM 5-aminolevulinic acid. Protein expression was carried out overnight at 18°C.

Protein purification

Cells were re-suspended in 10 mL of lysis buffer per gram of pellets (50 mM TrisHCl, pH 8.0, 150 mM NaCl, 5% glycerol), supplemented with one tablet of EDTA-free protease inhibitor (cOmplete Protease Inhibitor Cocktail, Roche) and 10 U/mL DNase I. Lysis was carried out by sonication on ice at 70% amplitude

Figure 2: Design and structural prediction of truncated monomeric mutant *Pa*BphP construct. (a) Locations of I294E and V298E mutation sites in the wild-type, truncated structure, highlighting their spatial proximity. (b) AlphaFold prediction of the wild-type truncated sequence as a dimer, showing proper dimerization and a conformation comparable to the crystal structure. (c) AlphaFold prediction of the I294E/V298E mutant in its monomeric form, showing proper folding of the monomeric unit.

for 5 min, with 5 s pulses on and 10 s pulses off. The supernatant was collected after centrifugation at

20,000g for 35 min (JA 14.50 rotor, Beckman) and was briefly incubated with a molar excess of biliverdin

hydrochloride.ă The lysate was purified by Ni-NTA affinity chromatography using a HisTrap HP 5 mL col-

umn (Cytiva). The bound protein was washed with wash buffer I (50 mM Tris, 50 mM NaCl, pH 8.0) and

wash buffer II (50 mM Tris, 1 M NaCl, pH 8.0), and eluted with elution buffer (50 mM Tris, 50 mM NaCl,

300 mM imidazole, pH 8.0). This was followed by size-exclusion chromatography using a HiLoad 16/600

Superdex 200 pg column (Cytiva) in wash buffer I. The purified protein was then flash-frozen for storage.

Design of monomeric variant

The mutant plasmid was designed using only the photosensory module (PSM) of the protein, excluding

the histidine kinase module, to promote monomerization with minimal mutational changes. Mutation

sites were selected based on amino acid properties and spatial proximity (within 4 Å). I294E and V298E

mutations were introduced into the wild-type sequence to replace hydrophobic residues with charged

glutamate residues (Fig. 2(a)). The mutant plasmid was generated using GenScripts site-directed muta-

genesis service.

UV-Vis absorption spectroscopy

Steady-state UVVis absorption measurements were performed using a Shimadzu UV-1900i spectropho-

tometer equipped with LabSolutions UVVis software (Version 1.13). For dark reversion kinetics, 100 se-

quential UVVis spectra scans were recorded at 45-second intervals with a delay of 4 seconds between

diode illumination of the phytochrome samples and the start of each measurement. For time-dependent

diode illumination experiments, samples were exposed to a 780 nm light-emitting diode (LED) for dura-

tions ranging from 6 to 60 seconds. Temperature-dependent measurements were conducted by irradiat-

ing the phytochrome samples with a 780 nm LED for 120 seconds at temperatures ranging from 10 °C

to 35 °C. To obtain a fully uniform Pr state, the sample was illuminated with a 780 nm LED for 120 sec-

onds. To obtain a fully uniform Pfr state, the sample was kept in darkness for 3 hours.

102 Calculation of Pr/Pfr fraction in diode illuminated phytochrome

103   To determine the relative amounts of Pr and Pfr in the illuminated samples, we first acquired absorp-

104   tion spectra for the pure Pr and pure Pfr states. The pure Pfr spectrum was recorded after incubating the

105   sample in the dark for approximately three hours. The pure Pr spectrum was recorded after illuminating

106   the sample with a 780 nm LED for 2 minutes. The concentrations of Pr and Pfr in the illuminated sample

107   were determined by fitting the initial spectrum obtained after illumination to the following equation:

$$I(\lambda) = X \cdot I_{Pfr}(\lambda) + (1 - X) \cdot I_{Pr}(\lambda) \tag{1}$$

108   Here, $I, I_{Pfr}(\lambda)$, and $I_{Pr}(\lambda)$ are the intensity of experimental, pure Pfr and pure Pr spectra respectively. X

109   and (1-X) represent the fractions of Pfr and Pr, respectively. We assume that no sample degradation hap-

110   pens during the dark reversion process.

111 RESULTS

112 Design of monomeric variant

To study the dark reversion kinetics as a function of the oligomeric state of PaBphP, we prepared a trun-

cated, monomeric variant of the protein. Mutations I294E and V298E were introduced into the photo-

sensory module (PSM) to disrupt dimer formation by weakening hydrophobic interactions and introduc-

ing electrostatic repulsion (Fig. 2(a)). This mutational strategy was motivated by previous studies on the

canonical phytochrome from *Deinococcus radiodurans*, which showed that glutamate substitutions

at F145, L311, and L314 positions in *Dr*BphP effectively disrupted dimerization and produced stable

monomers.[53,60] Our AlphaFold predictions supported that these mutations would disrupt the dimer

interface while maintaining proper monomer folding. Structural models of both the wild-type and mu-

tant truncated proteins were generated using the GenBank sequence AAG07504.1, in both dimeric and

monomeric forms. While the wild-type sequence was reliably predicted to form a stable dimer (Fig. 2(b)),

AlphaFold did not predict a stable dimeric structure for the mutant variant. However, it successfully pre-

dicted a stable monomeric fold for the mutant (Fig. 2(c)), indicating that the introduced mutations dis-

rupted dimerization without compromising the native folding.

126 Size-exclusion chromatography

127   To confirm the oligomeric states of the full-length protein (FL-*Pa*BphP-D) and the truncated mutant

128   (PSM-*Pa*BphP-M), we performed size-exclusion chromatography (SEC). Preparative SEC (Fig. 1(b)) showed

Figure 3: The sequential and direct model cannot be distinguished by fitting the dark reversion kinetics of FL-*Pa*BphP-D and PSM-*Pa*BphP-M. UV-Vis spectra recorded over time during dark reversion for (a) FL-*Pa*BphP-D, and (b) PSM-*Pa*BphP-M. (c) Global fitting of dark reversion kinetics for FL-*Pa*BphP-D at 700 nm and 750 nm using a direct sequential model. Solid lines represent the global fit with a direct model, while dashed lines show the fit using a sequential model. The temperature for this experiment is maintained at 30 °C.

that FL-*Pa*BphP-D (blue trace) predominantly forms dimers, indicated by a major peak at lower elution

volumes (68-72 mL). A minor peak at 78 mL suggests a small monomeric population. The shaded re-

gion marks the fraction collected for further analysis. In contrast, PSM-*Pa*BphP-M (red trace) showed

a single peak around 88 mL, consistent with monomeric species. Analytical SEC of the FL-*Pa*BphP-D

fraction, performed both in the dark and under continuous 780 nm illumination, confirmed the dimeric

state in both Pr and Pfr forms, with elution volumes of 1.24-1.26 mL. A small shoulder at 1.5 mL may

correspond to a minor monomeric fraction (Fig. 1(c)). A slight shift in the dimer peak between states likely

reflects conformational differences between Pr and Pfr that influence the proteins shape. We conclude

that PSM-*Pa*BphP-M is monomeric, whereas the wild-type FL-*Pa*BphP-D is predominantly dimeric (>90%).

Dark reversion

To measure dark reversion kinetics, all phytochrome samples were illuminated for 120 s with LED light

at 780 nm to populate the Pr state. Fig. 3(a,b) shows the time-dependent spectral evolution from the

initially photoconverted Pr state back to the Pfr state. Under these conditions, FL-*Pa*BphP-D and PSM-

*Pa*BphP-M display qualitatively similar spectral features over time.

The dark reversion kinetics of the protein is analyzed using two kinetic models to elucidate the underly-

ing mechanism: a direct transition and a sequential model involving an intermediate state. We first de-

scribe the direct transition model:

$$\text{PrPr} \xrightarrow{k_1} \text{PfrPfr} \tag{2}$$

146   Here, k_1 is the transition rate constant from PrPr to PfrPfr. The analytic solution for fitting the kinetic data

147   is

$$c_1(\lambda, t) = c_0^{(1)}(\lambda) \cdot \exp(-k_1 \cdot t) \tag{3}$$

148   where $c_0^{(1)}(\lambda)$ and $c_1(\lambda, t)$ are the initial and time-dependent concentrations of the PrPr state, respectively.

149   The sequential model incorporates the presence of an intermediate state (PrPfr) between the initial and

150   final species of the dark reversion process. This model is more plausible than the direct model for the

151 dimeric system, provided that the intermediate is sufficiently long-lived to be experimentally detectable.

152 The model is described by:

$$PrPr \xrightarrow{k_1} PrPfr \xrightarrow{k_2} PfrPfr \tag{4}$$

153 Here k_1 and k_2 are the dark reversion rate constant from PrPr to PrPfr, and from PrPfr to PfrPfr respec-

154 tively. The solutions of the sequential model used for the global fitting are

$$c_1(\lambda, t) = c_0^{(1)}(\lambda) \cdot \exp(-k_1 \cdot t), \tag{5}$$

$$c_2(\lambda, t) = c_0^{(1)}(\lambda) \cdot \left(\frac{k_1}{k_2 - k_1} \right) \cdot \exp(-k_1 \cdot t) + \left(c_0^{(2)}(\lambda) - c_0^{(1)}(\lambda) \cdot \left(\frac{k_1}{k_2 - k_1} \right) \right) \cdot \exp(-k_2 \cdot t), \tag{6}$$

156 and

$$c_3(\lambda, t) = 1 - c_1(\lambda, t) - c_2(\lambda, t), \tag{7}$$

where $c_0^{(1)}(\lambda), c_0^{(2)}(\lambda)$ are the initial concentration of PrPr and PrPfr species. $c_1(\lambda, t), c_2(\lambda, t)$, and $c_3(\lambda, t)$ rep-

resent the time-dependent concentration of the PrPr, PrPfr and PfrPfr forms, respectively. We perform

global fitting of the time-dependent dark reversion spectra using these kinetic models.

In the first experiment, dark reversion kinetics is measured starting from a state with a maximal Pr pop-

ulation. Global fitting of the kinetic data for FL-PaBphP-D is performed using both the direct (Equation

3) and sequential (Equations 5-7) models. The resulting kinetic fits at 700 and 750 nm are shown in Fig.

3(c). Although the sequential model yields a slightly better fit than the direct model, this difference is not

sufficient to clearly favor one model over the other.

To differentiate between the models, we investigated the dependence of the refined rate constants on

the initial photostationary state. To achieve varying initial photostationary states, we altered the duration

of the illumination at 780 nm from 6 to 60 seconds. This variation in illumination time modified the ini-

tial fraction of the Pr state at the onset of the dark reversion measurement (Fig. 4(a,b)). The resulting pho-

tostationary states are defined as the fraction of Pr relative to the total phytochrome population, as given

by[44]:

$$\varphi = \frac{Pr}{P_{\text{total}}}. \tag{8}$$

171 φ was determined by fitting the spectra with the pure spectra for Pr and Pfr (section 2.5). In terms of the

172 sequential model, the initial relative concentrations of PrPr, PrPfr, and PfrPfr species depend on φ and

173 can be calculated assuming a stochastic occupancy of the hybrid state as[41]

$$PrPr = \varphi^2 \tag{9}$$

$$PrPfr = 2\,(1 - \varphi)\,\varphi, \tag{10}$$

$$PfrPfr = (1 - \varphi)^2. \tag{11}$$

The relative concentrations of PrPr, PrPfr, and PfrPfr, needed for the global fit, are calculated using Equa-

tion 9-11 and shown in Fig. 4(c).

After global fitting of all data sets, we now analyze the rate-refined constants as a function of the initial

photostationary states. Using the direct transition model (Equation 2-3) the rate constants (k_1) remain in-

dependent of the initial photostationary states for PSM-PaBphP-M, as shown in Fig. 4(d). This is expected

as the rate constant of dark reversion should not change with the population of the initial state. In con-

trast, for the FL-PaBphP-D, the refined rates vary significantly with φ (Fig. 4(d)). This dependence indi-

cates that the direct transition model does not adequately describe the reaction, as true rate constants

should be independent of initial concentrations. When examining the rates of the sequential model, re-

fined for spectra starting from different photostationary states, we find that k_1 and k_2 remain constant for

$\varphi > 0.5$ (Fig. 4(e) and Table 1). Notably, k_1 at the lowest excitation fraction ($\varphi = 0.3$) appears as an out-

lier, which is likely due to the low abundance of PrPr under this illumination condition, which limits the

reliable determination of k_1. This analysis strongly supports the applicability of the sequential model to

the dark reversion of the dimeric protein, as the rate constants are independent of the initial population

distribution. The monomeric variant further confirms this, as the direct model adequately describes the

dark reversion across all φ values.

We extended our investigation by probing the temperature dependence of dark reversion kinetics. Spec-

tra for both proteins were recorded across a temperature range of $10°C$ to $35°C$. Each measurement was

initiated from pure Pr (PrPr) state at maximum photoexcitation. The initial spectra recorded immediately

after illumination are consistent across all temperatures, confirming a uniform starting population of the

Pr state. To extract the temperature-dependent rate constants, we globally fit the kinetic data using the

sequential model (Equation 4) for FL-PaBphP-D and the direct model (Equation 2) for PSM-PaBphP-M.

The resulting temperature dependence of the rate constants is shown in Fig.- 5a,b and tabulated in Ta-

ble 2. For both proteins, the kinetic rate constants increase significantly with temperature, indicating that

thermal energy facilitates dark reversion. To quantify this effect, the activation energy (E_a) for each kinetic

step was determined using the Arrhenius equation. For FL-PaBphP-D, the activation energies for k_1 and

k_2 are 2.44 and 5.11 kJ/mol, respectively. These values lead to the unexpected conclusion that the pho-

tochemical state of the sister protomer in the dimer influences the activation energy of the reaction. In

contrast, PSM-PaBphP-M shows a significantly higher activation energy of 15.21 kJ/mol for k_1, suggest-

ing that even the quaternary structure of the protein impacts the activation energy.

Figure 4: **Preparation of different initial concentrations of Pr and Pfr and global kinetic fits. The temperature for this experiment is maintained at 30 °C. Spectra support the sequential model for the wild-type protein and reveal the existence of the hybrid PrPfr state.** Dark reversion spectra at various photostationary states for (a) PSM-*Pa*BphP-M and (b) FL-*Pa*BphP-D. (c) Variation in the starting fractions of PrPr, PrPfr, and PfrPfr states with varying Pr concentration for the FL-*Pa*BphP-D. (d) Variation of the dark reversion rate constants, obtained from global fitting with the direct step model for FL-*Pa*BphP-D and PSM-*Pa*BphP-M. (e) Variation of the fast (k_1) and slow (k_2) dark reversion rate constants with Pr fraction for FL-*Pa*BphP-D. (f) Variation of simulated relative concentration of PrPr, PrPfr, and PfrPfr over time during dark reversion.

Our kinetic analysis shows that the transition rate (k_1) from PrPr to PrPfr is about 2-3 times faster than

the transition rate (k_2) from PrPfr to PfrPfr. This suggests that a significant portion of the PrPfr interme-

diate state is formed during the intermediate phase of dark reversion in FL-*Pa*BphP-D (Fig. 4(f)). More

importantly, this shows that substantial allosteric interactions may occur between the two protomers in

the dimer, as the dark reversion time differs depending on whether both monomers are in the Pr state

or if one is in the Pfr state. This points to an as-yet undescribed interaction between the two protomers in

the dimer.

DISCUSSION

Our kinetic analysis of *Pa*BphP provides the first direct demonstration of a PrPfr heterodimer interme-

diate during the thermal reversion of bacterial phytochromes. By analyzing kinetic models in a series of

experiments with varying initial concentrations of Pr and Pfr, we show that a two-step mechanism accu-

rately describes the dark reversion pathway. Only this model yields rate constants that are independent

of the initial Pr:Pfr ratio

This finding mirrors observations made for plant phytochromes, where dark reversion proceeds via a PrPfr

heterodimer before returning to the Pr state.[41,44] *Pa*BphP being a bathy phytochrome exhibits a similar

sequential interconversion pathway, but in the reverse direction, from Pr to Pfr.

To our knowledge and somewhat surprisingly, earlier studies on bacterial phytochromes have not con-

sidered the possibility of a PrPfr intermediate state[22,28,37,52,57,61 35,45–47,51,58]. This may be due to the fact

Figure 5: Arrhenius analysis shows notably low activation energies for the dark reversion of _Pa_BphP.
Logarithmic variation of (a) fast (k_1) and slow (k_2) dark reversion rate constants with the inverse of temperature for FL-_Pa_BphP-D and (b) logarithmic variation of fast (k_1) dark reversion rate constants with the inverse of temperature for PSM-_Pa_BphP-M. Activation energy is calculated using Arrhenius equation ($k = Ae^{-\frac{E_a}{kT}}$)

that, in canonical phytochromes, the initial state is typically a mixture of Pr and Pfr, making it difficult

to assign a specific dark recovery pathway. In contrast, bathy phytochromes allow for the generation of

nearly pure Pr states through far-red illumination, enabling the identification of a PrPfr intermediate state.

It is established that the Pfr form of _Pa_BphP represents the biochemical "off" state.[37,52] Huber _et al._ qual-

itatively correlated biochemical kinase activity, measured via radionucleotide assays, with the decaying

Pr/Pfr ratio during dark reversion.[52] However, their study did not consider the potential presence of a

mixed PrPfr heterodimer. This raises the question of whether the PrPfr state is also biochemically active

and whether it might even substitute for PrPr as the functional state. Notably, in plant phytochromes, the

hybrid PrPfr state is known to play a distinct functional role compared to the pure PrPr or PfrPfr states.[41,56]

A definitive assignment of the PrPfr states function would require a quantitative analysis of the radionu-

cleotide assays, a challenging task, as acknowledged by Huber _et al.._. Nonetheless, their assays show high

biochemical activity after prolonged far-red illumination,[52], a condition in which the PrPr population is

significantly higher than that of PrPfr (see Fig. 4(c) at $\varphi = 1$). We believe that PrPr is indeed kinase-active.

However, this line of reasoning does not rule out the possibility that the PrPfr state also exhibits biochem-

ical activity.

Table 1: Photostationary-state-dependent dark reversion kinetic rates for FL-_Pa_BphP-D construct

φ	FL-_Pa_BphP-D	
	k_1 (s^{-1})	k_2 (s^{-1})
0.30	0.011	0.0011
0.55	0.0058	0.0013
0.64	0.0056	0.0013
0.70	0.0058	0.0014
0.80	0.0051	0.0015
0.85	0.0052	0.0015
0.92	0.0055	0.0015
0.98	0.0056	0.0015

237   Our temperature-dependent measurements reveal activation energies that are significantly lower than

238   those typically reported for dark reversion in plant phytochromes. In studies on the effect of temperature

Table 2: Temperature dependent rate constants for FL-*Pa*BphP-D and PSM-*Pa*BphP-M construct

Temperature	FL-*Pa*BphP-D		PSM-*Pa*BphP-M
(°C)	k_1 (s^{-1})	k_2 (s^{-1})	k_1 (s^{-1})
10	0.0062	0.00168	0.0009
15	0.0064	0.00175	0.0011
20	0.0066	0.00179	0.0013
25	0.0065	0.00191	0.0014
30	0.0069	0.00191	0.0015
35	0.0067	0.0021	0.0016

on the dark reversion properties of *Arabidopsis thaliana* phyA, activation enthalpies of 92.9 and 74.5

kJ·mol^{-1} were reported for PfrPfr homodimers and PrPfr heterodimers, respectively.[42] Another study

found that all four isoforms showed strong temperature dependence, with activation energies of 104, 62,

163, and 75.3 kJ·mol^{-1} for PhyA, PhyB, PhyC, and PhyE, respectively.[43] Furthermore, phosphorylation

has been shown to modulate the dark reversion rate.[55]

The unusually low activation barriers observed for the *Pa*BphP protein are difficult to explain solely by

a steric model of C15=C16 double bond isomerization. An alternative chemical mechanism may be in-

volved. In fact, ketoenol tautomerization of the D-ring in its Z-isomerized (Pr) state has been proposed to

precede isomerization of the C15C16 bond.[51] The enol form lowers the bond order of the C15C16 bond,

which would significantly reduce the energy of the transition state and could account for the unusually

low activation energy observed in our measurements.

Interestingly, we observe distinct rates and activation energies for the two steps of dark reversion (Fig. 5),

suggesting that the state of the sister monomer within the dimer influences the chromophores reversion

reaction. This strongly indicates an allosteric interaction. Supporting this, the monomeric variant, which

lacks a dimer interface, exhibits an even higher activation energy for dark reversion (Fig. 5b). A similar

observation has been made for plant phytochromes, suggesting that the two-step mechanism is tightly

regulated by the protein and likely optimized through evolution[42]

So far, allostery in bacterial phytochromes has primarily been discussed in the context of linking chro-

mophore photoactivation to the regulation of biochemical output. However, the allosteric interaction ob-

served here happens between the sister protomers within the dimer, suggesting the existence of a pre-

viously unidentified signaling pathway between the two protein subunits. Recent NMR investigations of

the *D. radiodurans* phytochrome support this idea, revealing chemical shift changes that propagate from

the chromophore to the dimer interface within the PAS/GAF domains[62]. Further structural studies are

needed to uncover the molecular basis of this signaling route.

CONCLUSION

In this study, we have successfully demonstrated the transient accumulation of a mixed PrPfr hybrid state

in the bacterial phytochrome *Pa*BphP during the dark reversion process. Our kinetic analysis indicates

that both subunits within the dimeric phytochrome can exist in different photoisomeric states simultaneously during dark reversion, and that the hybrid state is transiently populated at high concentration. We also observe that a monomeric variant of *Pa*BphP follows a direct Pr-to-Pfr reaction scheme, which further supports the assignment of the PrPfr intermediate as a feature arising from the dimeric nature of the wild-type protein. Furthermore, the temperature-dependent kinetics reveal low activation energies for the dark reversion processes in both the dimeric and monomeric PaBphP, consistent with a previously proposed keto-enol tautomerization as a crucial step preceding dark reversion. The distinct kinetic behavior observed in the dimeric *Pa*BphP, specifically, the transient accumulation of the PrPfr hybrid state and the difference in activation energy, suggests the presence of allosteric regulation between the two subunits. We propose that this plays an important role in controlling the signaling output of bacterial phytochromes.

ACKNOWLEDGEMENTS

The author acknowledge Göran Gustaffson foundations and Carl-Trygger Foundation for supporting this work. MM acknowledges the Swedish Research Council (Grant No. VR 2020-05403), the Swedish Society for Medical Research (S20-0156), and Harald and Greta Jeanssons Foundation Grant (J2021-0114).

REFERENCES

[1] P. Hegemann, *annurev.arplant*, 2008, **59**, 167–189.

[2] A. Möglich, X. Yang, R. A. Ayers and K. Moffat, *annurev.arplant*, 2010, **61**, 21–47.

[3] N. C. Rockwell, Y.-S. Su and J. C. Lagarias, *annurev.arplant*, 2006, **57**, 837–858.

[4] H. Wang and H. Wang, *Mol. Plant.*, 2015, **8**, 540–551.

[5] T. Lamparter, N. KrauSS and P. Scheerer, *Photochem. Photobiol.*, 2017, **93**, 642–655.

[6] J. J. Casal and R. A. Sánchez, *Seed Sci. Res*, 1998, **8**, 317329.

[7] E. Monte, J. M. Tepperman, B. Al-Sady, K. A. Kaczorowski, J. M. Alonso, J. R. Ecker, X. Li, Y. Zhang and P. H. Quail, *Proc. Natl. Acad. Sci. U.S.A*, 2004, **101**, 16091–16098.

[8] M. Endo, Y. Tanigawa, T. Murakami, T. Araki and A. Nagatani, *Proc. Natl. Acad. Sci. U.S.A*, 2013, **110**, 18017–18022.

[9] J. A. Kyndt, T. E. Meyer and M. A. Cusanovich, *Photochem. Photobiol. Sci*, 2004, **3**, 519–530.

[10] L. Wu, R. S. McGrane and G. A. Beattie, *MBio*, 2013, **4**, 10–1128.

[11] R. K. Verma, A. Biswas, A. Kakkar, S. K. Lomada, B. B. Pradhan and S. Chatterjee, *Cell Rep*, 2020, **32**, year.

[12] K. G. Chernov, T. A. Redchuk, E. S. Omelina and V. V. Verkhusha, *Chem Rev*, 2017, **117**, 6423–6446.

[13] A. A. Kaberniuk, A. A. Shemetov and V. V. Verkhusha, *Nat. Methods*, 2016, **13**, 591–597.

[14] A. T. Ulijasz and R. D. Vierstra, *Curr. Opin. Plant Biol.*, 2011, **14**, 498–506.

[15] A. Möglich, R. A. Ayers and K. Moffat, *Structure*, 2009, **17**, 1282–1294.

[16] L. Aravind and C. P. Ponting, *Trends Biochem. Sci.*, 1997, **22**, 458–459.

[17] L.-O. Essen, J. Mailliet and J. Hughes, *Proc. Natl. Acad. Sci. U.S.A*, 2008, **105**, 14709–14714.

[18] E. S. Burgie and R. D. Vierstra, *Plant Cell*, 2014, **26**, 4568–4583.

[19] M. Legris, C. Klose, E. S. Burgie, C. C. R. Rojas, M. Neme, A. Hiltbrunner, P. A. Wigge, E. Schäfer, R. D. Vierstra and J. J. Casal, *Science*, 2016, **354**, 897–900.

[20] G. Rottwinkel, I. Oberpichler and T. Lamparter, *J. Bacteriol. Res*, 2010, **192**, 5124–5133.

[21] F. Velázquez Escobar, D. Buhrke, N. Michael, L. Sauthof, S. Wilkening, N. N. Tavraz, J. Salewski, N. Frankenberg-Dinkel, M. A. Mroginski, P. Scheerer *et al.*, *Photochem. Photobiol*, 2017, **93**, 724–732.

[22] M. F. López, M. Dahl, F. V. Escobar, H. R. Bonomi, A. Kraskov, N. Michael, M. A. Mroginski, P. Scheerer and P. Hildebrandt, *Phys. Chem. Chem. Phys.*, 2022, **24**, 11967–11978.

[23] F. Velázquez Escobar, D. Buhrke, N. Michael, L. Sauthof, S. Wilkening, N. N. Tavraz, J. Salewski, N. Frankenberg-Dinkel, M. A. Mroginski, P. Scheerer *et al.*, *Photochem. Photobiol*, 2017, **93**, 724–732.

[24] S. Westenhoff, P. Meszaros and M. Schmidt, *Curr. Opin. Struct. Biol.*, 2022, **77**, 102481.

[25] A. Björling, O. Berntsson, H. Lehtivuori, H. Takala, A. J. Hughes, M. Panman, M. Hoernke, S. Niebling, L. Henry, R. Henning, I. Kosheleva, V. Chukharev, N. V. Tkachenko, A. Menzel, G. Newby, D. Khakhulin, M. Wulff, J. A. Ihalainen and S. Westenhoff, *Sci. Adv.*, 2016, **2**, e1600920.

[26] M. Chenchiliyan, J. Kübel, S. A. Ooi, G. Salvadori, B. Mennucci, S. Westenhoff and M. Maj, *J. Chem. Phys.*, 2023, **158**, 085103.

[27] S. G. Sokolovski, E. A. Zherebtsov, R. K. Kar, D. Golonka, R. Stabel, N. B. Chichkov, A. Gorodetsky, I. Schapiro, A. Möglich and E. U. Rafailov, *Biophys. J.*, 2021, **120**, 964–974.

[28] Y. Yang, T. Stensitzki, L. Sauthof, A. Schmidt, P. Piwowarski, F. Velazquez Escobar, N. Michael, A. D. Nguyen, M. Szczepek, F. N. Brünig, R. R. Netz, M. A. Mroginski, S. Adam, F. Bartl, I. Schapiro, P. Hildebrandt, P. Scheerer and K. Heyne, *Nat. Chem.*, 2022, **14**, 823–830.

[29] H. Takala, P. Edlund, J. A. Ihalainen and S. Westenhoff, *Photochem. Photobiol. Sci.*, 2020, **19**, 1488–1510.

[30] X. Yang, Z. Ren, J. Kuk and K. Moffat, *Nature*, 2011, **479**, 428–432.

[31] X. Yang, J. Kuk and K. Moffat, *Proc. Natl. Acad. Sci. U.S.A*, 2009, **106**, 15639–15644.

[32] J. R. Wagner, J. S. Brunzelle, K. T. Forest and R. D. Vierstra, *Nature*, 2005, **438**, 325–331.

[33] L.-O. Essen, J. Mailliet and J. Hughes, *Proceedings of the National Academy of Sciences*, 2008, **105**, 14709–14714.

[34] X. Yang, J. Kuk and K. Moffat, *PNAS*, 2008, **105**, 14715–14720.

[35] W. Y. Wahlgren, E. Claesson, I. Tuure, S. Trillo-Muyo, S. Bódizs, J. A. Ihalainen, H. Takala and S. Westenhoff, *Nat. Commun*, 2022, **13**, 7673.

[36] H. Li, E. S. Burgie, Z. T. K. Gannam, H. Li and R. D. Vierstra, *Nature*, 2022, **604**, 127–133.

[37] S. Bódizs, P. Mészáros, L. Grunewald, H. Takala and S. Westenhoff, *Structure*, 2024, **32**, 1952–1962.e3.

[38] D. Buhrke, G. Gourinchas, M. Müller, N. Michael, P. Hildebrandt and A. Winkler, *J. Biol. Chem.*, 2020, **295**, 539–551.

[39] T. N. Malla, C. Hernandez, S. Muniyappan, D. Menendez, D. Bizhga, J. H. Mendez, P. Schwander, E. A. Stojkovi and M. Schmidt, *Sci. Adv*, 2024, **10**, eadq0653.

[40] E. Schäfer and W. Schmidt, *Planta*, 1974, **116**, 257–266.

[41] J. BROCKMANN, S. RIEBLE, N. KAZARINOVA-FUKSHANSKY, M. SEYFRIED and E. SCHÄFER, *Plant Cell Environ.*, 1987, **10**, 105–111.

[42] L. Hennig and E. Schäfer, *J. Biol. Chem.*, 2001, **276**, 7913–7918.

[43] E. S. Burgie, Z. T. K. Gannam, K. E. McLoughlin, C. D. Sherman, A. S. Holehouse, R. J. Stankey and R. D. Vierstra, *Proc. Natl. Acad. Sci. U.S.A*, 2021, **118**, e2105649118.

[44] W. Schmidt and E. Schäfer, *Planta*, 1974, **116**, 267–272.

[45] C. Huber, M. Strack, I. SchultheiSS, J. Pielage, X. Mechler, J. Hornbogen, R. Diller and N. Frankenberg-Dinkel, *J. Biol. Chem.*, 2024, **300**, 107148.

[46] A. Kraskov, A. D. Nguyen, J. Goerling, D. Buhrke, F. Velazquez Escobar, M. Fernandez Lopez, N. Michael, L. Sauthof, A. Schmidt, P. Piwowarski, Y. Yang, T. Stensitzki, S. Adam, F. Bartl, I. Schapiro, K. Heyne, F. Siebert, P. Scheerer, M. A. Mroginski and P. Hildebrandt, *Biochemistry*, 2020, **59**, 1023–1037.

[47] G. Merga, M. F. Lopez, P. Fischer, P. Piwowarski, . Nogacz, A. Kraskov, D. Buhrke, F. V. Escobar, N. Michael, F. Siebert, P. Scheerer, F. Bartl and P. Hildebrandt, *Phys. Chem. Chem. Phys.*, 2021, **23**, 18197–18205.

[48] H. Takala, H. Lehtivuori, H. Hammarén, V. P. Hytönen and J. A. Ihalainen, *Biochemistry*, 2014, **53**, 7076–7085.

[49] E. S. Burgie, Z. T. K. Gannam, K. E. McLoughlin, C. D. Sherman, A. S. Holehouse, R. J. Stankey and R. D. Vierstra, *PNAS*, 2021, **118**, e2105649118.

[50] J.-H. Jung, M. Domijan, C. Klose, S. Biswas, D. Ezer, M. Gao, A. K. Khattak, M. S. Box, V. Charoensawan, S. Cortijo, M. Kumar, A. Grant, J. C. W. Locke, E. Schäfer, K. E. Jaeger and P. A. Wigge, *Science*, 2016, **354**, 886–889.

[51] F. Velazquez Escobar, P. Piwowarski, J. Salewski, N. Michael, M. Fernandez Lopez, A. Rupp, B. M. Qureshi, P. Scheerer, F. Bartl, N. Frankenberg-Dinkel, F. Siebert, M. Andrea Mroginski and P. Hildebrandt, *Nat. Chem.*, 2015, **7**, 423–430.

[52] C. Huber, M. Strack, I. SchultheiSS, J. Pielage, X. Mechler, J. Hornbogen, R. Diller and N. Frankenberg-Dinkel, *J. Biol. Chem.*, 2024, **300**, 107148.

[53] H. Takala, A. Björling, M. Linna, S. Westenhoff and J. A. Ihalainen, *J. Biol. Chem.*, 2015, **290**, 16383–16392.

[54] W. Schmidt and E. Schäfer, *Planta*, 1974, **116**, 267–272.

[55] M. Medzihradszky, J. Bindics, . Ádám, A. Viczián, . Klement, S. Lorrain, P. Gyula, Z. Mérai, C. Fankhauser, K. F. Medzihradszky, T. Kunkel, E. SchÄfer and F. Nagy, *The Plant Cell*, 2013, **25**, 535–544.

[56] C. Klose, F. Venezia, A. Hussong, S. Kircher, E. Schäfer and C. Fleck, *Nat. Plants*, 2015, **1**, 15090.

[57] H. Takala, A. Björling, M. Linna, S. Westenhoff and J. A. Ihalainen, *J. Biol. Chem.*, 2015, **290**, 16383–16392.

[58] K. D. Piatkevich, F. V. Subach and V. V. Verkhusha, *Nat. Commun.*, 2013, **4**, 2153.

[59] A. Kraskov, A. D. Nguyen, J. Goerling, D. Buhrke, F. Velazquez Escobar, M. Fernandez Lopez, N. Michael, L. Sauthof, A. Schmidt, P. Piwowarski, Y. Yang, T. Stensitzki, S. Adam, F. Bartl, I. Schapiro, K. Heyne, F. Siebert, P. Scheerer, M. A. Mroginski and P. Hildebrandt, *Biochemistry*, 2020, **59**, 1023–1037.

[60] M. E. Auldridge, K. A. Satyshur, D. M. Anstrom and K. T. Forest, *J. Biol. Chem.*, 2012, **287**, 7000–7009.

[61] R. Tasler, T. Moises and N. Frankenberg-Dinkel, *The FEBS Journal*, 2005, **272**, 1927–1936.

[62] L. Isaksson, E. Gustavsson, C. Persson, U. Brath, L. Vrhovac, G. Karlsson, V. Orekhov and S. Westenhoff, *Structure*, 2020, **29**, 151–160.e3.