1 **CRISPRi-based functional genomic screening identifies genes essential for**
2 **CH$_4$-dependent growth in the methanotroph *Methylococcus capsulatus* Bath.**
3

4 Jessica M. Henard, Spencer A. Lee, Yao-Chuan Yu, Danyang Shao, Rajeev K. Azad, and Calvin
5 A. Henard*.
6
7 Department of Biological Sciences and BioDiscovery Institute, University of North Texas,
8 Denton, TX, USA.
9
10 *Correspondence to Calvin.Henard@unt.edu

11
12 Running title: Methanotroph CRISPRi
13
14 Keywords: methanotroph, CRISPRi, methane monooxygenase, RubisCO, functional genomics

Abstract

Aerobic methanotrophic bacteria are the primary organisms that consume atmospheric methane (CH_4) and have potential to mitigate the climate-active gas. However, a limited understanding of the genetic determinants of methanotrophy hinders the development of biotechnologies leveraging these unique microbes. Here, we developed and optimized a CRISPR interference (CRISPRi) system to enable functional genomic screening in methanotrophic bacteria. We built a genome-wide single guide RNA (sgRNA) library in the industrial methanotroph, *Methylococcus capsulatus*, consisting of ~45,000 unique sgRNAs mediating inducible, CRISPRi-dependent transcriptional repression. A selective screen during growth on CH_4 identified 233 genes whose transcription repression resulted in a fitness defect and repression of 13 genes associated with a fitness advantage. Enrichment analysis of the 233 putative essential genes linked many of the encoded proteins with critical cellular processes like ribosome biosynthesis, translation, transcription, and other central biosynthetic metabolism, highlighting the utility of CRISPRi for functional genetic screening in methanotrophs. The CRISPRi screen uncovered that 50 ppb Co^{2+} in methanotroph Nitrate Mineral Salts medium is growth inhibitory and decreasing Co^{2+} to < 10 ppb improved biomass productivity from CH_4. Collectively, our results show that the CRISPRi system and sgRNA library developed here can be used for facile gene-function analyses and genomic screening to identify novel genetic determinants of methanotrophy and isolation of improved methanotroph biocatalysts.

Introduction

Aerobic methanotrophic bacteria, methanotrophs, play an essential role in balancing atmospheric carbon by oxidizing methane (CH_4) to multi-carbon molecules (methanotroph biomass) and carbon dioxide (CO_2). These ubiquitous microbes limit the escape of CH_4 to the atmosphere via biofiltration in terrestrial and aquatic systems[1]. Methanotrophs can also be leveraged for the bioconversion of abundant CH_4-rich gas streams from oil and natural gas extraction sites, coal mines, agricultural waste, landfills, and wastewater treatment facilities, and have the potential to capture rising CH_4 greenhouse gas and mitigate climate change[2]. However, significant advances in methanotroph biology and engineering for effective CH_4 capture and conversion are needed to realize cost-effective solutions using methanotrophs.

Methanotrophs belong to a broad group of methylotrophic microbes that use organic single carbon (C_1) molecules as sources of carbon and/or energy, which, in addition to CH_4, includes methanol (CH_3OH), formate (CH_2O), and/or methylamine. Unique to methanotrophs is the CH_4 monooxygenase (MMO) enzyme that oxidizes the strong C-H bond of CH_4 to generate CH_3OH[3]. Methanotroph C_1 metabolic enzymes and pathways for CH_3OH assimilation to biomass and its dissimilation to CO_2 generally overlap with those in methylotrophs[4]. These include the ribulose monophosphate and serine cycles used by Gammaproteobacterial methanotrophs and Alphaproteobacterial methanotrophs for C_1 assimilation, respectively[5–7]. In addition, the methanotroph dissimilatory metabolic processes that couple C_1 oxidation to the electron transport chain for ATP production are coincident with those in other methylotrophs[89]. These central CH_4 metabolic pathways have been biochemically validated using radiolabeled isotopes in a subset of methanotrophs[5,7,10] and the availability of several complete methanotroph genomes and comparative genomics have provided critical insight into methanotroph CH_4 metabolism[11–18]. However, many genes that may be required for methanotrophy, especially those outside of the core C_1 metabolic gene clusters, are difficult to identify due to their strict CH_4-dependent cultivation.

Early research using chemical and insertional transposon mutagenesis of the CH_3OH-utilizing methylotrophic bacterium *Methylobacterium extorquens* AM1 identified many genes required for methylotrophic growth, including those encoding the enzymes needed for methanol oxidation, formaldehyde oxidation, formate oxidation, and assimilation via the serine cycle[6,19–26]. Additionally, transposon mutagenesis led to the identification of genes required for transport and utilization of rare earth elements[27] and transcriptional regulators of the C_1 metabolic pathways[20,28]. Notably, the elucidation of genes required for methylotrophy was enabled by leveraging their facultative capacity to utilize alternative multi-carbon sources (i.e. succinate) supplied during library generation[28]. In contrast, the vast majority of methanotrophs are obligate CH_4-utilizers; thus, mutations in essential genes required for growth on CH_4 would be lost from the mutagenized population. There are only two reports of successful chemical mutagenesis[29] and one report of transposon insertional mutagenesis[30] in methanotrophs, underscoring barriers

76 applying forward genetic techniques to CH_4-oxidizing bacteria and significant knowledge gaps
77 related thereto.
78
Clustered regularly interspaced short palindromic repeats (CRISPR) interference (CRISPRi) is a
genetic tool that has been leveraged for functional genetic screens in diverse bacteria[31–39].
CRISPRi uses the nuclease-deficient "dead" CRISPR-associated (Cas) variant dCas9 and a
single-guide RNA (sgRNA) for the sequence-specific repression of gene expression[40]. This
technology can be used to target single genes or scaled for genome-wide screening via massively
parallel oligonucleotide synthesis of pooled sgRNAs. Functional genomic screens using
CRISPRi are coupled with next generation sequencing to quantify the depletion or enrichment of
sgRNAs (sgRNA differential fitness) during selective bacterial cultivation[41]. Notably,
transcription repression by CRISPRi can be tightly regulated using inducible promoters to drive
dCas9 and/or sgRNA expression; thus, in contrast to transposon mutagenesis, CRISPRi can be
applied to conditionally evaluate the effect of repressing essential gene expression on bacterial
growth. Recent developments of methanotroph CRISPR genome editing tools indicate that Cas9
and sgRNAs are functional in Gammaproteobacterial and Alphaproteobacterial
methanotrophs[31,42–46] paving the way for CRISPRi and forward genetic approaches in
methanotrophs that have eluded these bacteria for decades.
94
Here, we modified a previously developed methanotroph CRISPR-Cas9 genome editing system[44]
to facilitate CRISPRi and demonstrate its utility for repressing gene transcription in the
phylogenetically diverse methanotrophs *Methylococcus capsulatus* Bath and *Methylosinus*
trichosporium OB3b. We performed a genome-scale CRISPRi screen using 45,798 unique
sgRNAs targeting all annotated genes in the *M. capsulatus* Bath genome to investigate gene-
function relationships during CH_4 cultivation. We identified 233 genes whose transcription
repression resulted in significantly decreased fitness and repression of 13 genes associated with a
fitness advantage. sgRNAs with decreased fitness scores targeted genes encoding proteins
involved in critical cellular processes like ribosome biosynthesis, translation, transcription, and
central biosynthetic metabolism, underscoring the utility of CRISPRi for the identification of
predicted and novel essential genes. The CRISPRi system and sgRNA library developed here can
be used for facile gene-function analyses and genomic screening to identify methanotroph
genetic determinants under an array of screening conditions. Further, these CRISPRi screening
methodologies can also be applied to high-throughput engineering approaches for isolation of
improved methanotroph biocatalysts.

Materials and Methods
Bacterial strains and cultivation. Bacterial strains used in this study are shown in Table 1. *E.*
coli strains were cultured in lysogeny broth (Lennox) with 25 µg/mL kanamycin, 50 µg/mL
spectinomycin, or 10 µg/mL gentamicin for transformant selection. *Methylococcus capsulatus*
Bath and *Methylosinus trichosporium* OB3b frozen stocks were streaked onto nitrate mineral
salts (NMS) solid medium and maintained in stainless steel gas chambers supplied with 20%
CH_4 in the gas phase at 37°C or 30°C, respectively, as previously described[44]. Methanotroph
strains were passaged weekly for 4 weeks (maximum passage = 5) after which time a new
culture from frozen stock was initiated. Plasmids were transferred to methanotrophs via
biparental mating by spreading equivalent biomass of S17-1λ *E. coli* and recipient methanotroph
biomass on NMS mating agar mating plates and incubating in a 20% CH_4 atmosphere for 24 h as
previously described [44]. After mating, methanotroph transformants harboring plasmids were
selected on NMS medium containing 50 µg/mL spectinomycin (pJHCi and derivatives), 10
µg/mL gentamicin (pDS1), and/or 50 µg/mL kanamycin (pJHint). Methanotrophs were cultured
in 150 mL vials containing 10 mL of NMS medium at 37°C (*M. capsulatus*) or 30°C (*M.*
trichosporium) at 200 rpm orbital shaking. After inoculation with plate-derived biomass to OD_{600}
0.01, vials were crimped with gray butyl stoppers to create gas-tight seals followed by
CH_4 addition to the headspace via syringe to reach a final CH_4 concentration of 20% in air (v/v).
Where applicable, sgRNA and/or dCas9 expression were induced with 0.2 µg/mL
anhydrotetracycline (aTc). High resolution bacterial growth was measured every 20 seconds by
Cell Growth Quantifier (CGQ) optical sensors (Scientific Bioprocessing), and the data is
presented as the mean arbitrary light backscatter units as directly detected by the sensors.
132
CRISPRi plasmid construction and sgRNA exchange. Primers and other synthetic DNA used in
this study are shown in Table 2. To construct a broad-host-range, single plasmid CRISPRi
system, we amplified a P_{mxaF}-sgRNA with an mRFP spacer sequence from the pgRNA plasmid[44]
using primers oCAH267/289 and ligated it with pCAH01[SpR]::Cas9[44] linearized with primers
oCAH230/231 via isothermal assembly using HiFi Gibson Assembly Master Mix (New England
Biolabs). We then amplified the dCas9 gene from plasmid pdCas9-bacteria[40] with
oCAH952/1507 and assembled it with the above plasmid backbone amplified with oCAH4/3,
exchanging the Cas9 gene with dCas9 to generate plasmid pJHCi. The aTc-inducible suicide
plasmid pJHint was constructed via isothermal assembly of the following six DNA fragments: 1)
pUC origin of replication from pCAH01 amplified with oCAH403/404; 2) 1kb upstream
homology arm amplified with oCAH405/406 from *M. capsulatus* genomic DNA; 3) P_{tet}
promoter/operator amplified from pCAH01 with oCAH407/408 ; 4) *tetR*-kn[R] cassette from
pCAH01 amplified with oCAH409/410; 5) 1kb downstream homology arm amplified with
oCAH411/412 from *M. capsulatus* genomic DNA; and 6) *oriT* origin of transfer amplified from
pCAH01 with oCAH413/414. pJHint was then amplified with primers oCAH3/4 and assembled
with the dCas9 gene fragment to generate plasmid pJHintdCas9. pJHintCas9 was transferred to
M. capsulatus Bath via conjugation and transformants with the P_{tet}-dCas9 module integrated into

an intergenic site between MCA_tRNA-Ile and MCA_tRNA-Pro were selected on NMS solid
medium containing 25 µg/mL kanamycin. Successful integration was confirmed via PCR using
primers oCAH447/448 that flank the integration site. For sgRNA expression, we initially
amplified the mRFP targeting sgRNA from pJHCi with primers oCAH1083/1081 and cloned it
into pCAH01 amplified without a ribosomal binding site with primers oCAH4/119 to generate
p01sgRNA^{mrfp}. Other aTc-inducible sgRNA expression plasmids were generated by exchanging
the spacer sequence via isothermal assembly of a single-stranded oligo containing a unique
spacer and the pCAH01sgRNA^{mrfp} amplified with primers oCAH1190/119.

Fluorescence measurements. pJHCi was transferred to *M. capsulatus* Bath or *M. trichosporium*
OB3b harboring the pDS1 mRFP reporter plasmid via conjugation. Alternatively, pDS1 and
p01sgRNA^{mrfp} were iteratively transferred to the *M. capsulatus* Bath strain with chromosomally
integrated P$_{tet}$-dCas9. Overnight seed cultures of these strains were subcultured to $OD_{600} = 0.01$
in liquid medium containing appropriate antibiotics with or without aTc induction and cultivated
for 72h to ~ OD_{600} 2.0. 200 µL culture was transferred to a 96-well microplate, and mRFP1
fluorescence (ex$_{532nm}$, em$_{588nm}$, gain = 80) and optical density (A$_{600nm}$) was measured with a
BioTek Synergy Mx microplate reader. The data were normalized by dividing raw fluorescence
reads by the optical density. mRFP expression was also monitored via fluorescence microscopy
by transferring 5 µL *M. capsulatus* logarithmic growth phase cultures with or without CRISPRi
induction to a glass slide and mounted with a cover slip. RFP (ex575-640nm, em629nm) images
were acquired using a Zeiss AXIO Imager M2 fluorescence microscope with 100X (oil) Plan-
Apochromat (HC DIC) objective and processed with ZEN software.

Construction of pLSQ and cbbL PAM mutant. The *M. capsulatus* Bath *cbbLSQ* operon
(MCA2743-2746) including 150bp upstream of *cbbL* presumed to contain the *cbbL* promoter
was amplified from purified genomic DNA using primers oCAH1629 and oCAH1750. The
amplicon was assembled with pDS1 amplified with primers oCAH1007 and oCAH16 to generate
pLSQ. This plasmid was then linearized with primers oCAH1751 and oCAH16 and assembled
with a synthetic fragment containing a *cbbL*:c.57C>T protospacer adjacent motif (PAM)
mutation to generate pLSQ^{ΔPAM}. The pCAH01sgRNA^{cbbL} plasmid was transferred *to M.*
capsulatus Bath harboring the pLSQ or pLSQ^{ΔPAM} strains and growth was assessed with or
without aTc induction using CGQ optical sensors.

Design, synthesis, and construction of the M. capsulatus genome-wide sgRNA library. Synbio
Technologies designed and synthesized the sgRNA oligo pool following design rules to identify
15 PAM and target sequences per the predicted 3,034 open reading frames in the NC_002977.6
RefSeq genome and to preferentially target the non-template DNA strand within the first 50% of
the 5' end of the target gene. The sgRNA library consisted of 45,798 sgRNA oligos (120 bp)
with each containing a unique spacer sequence and 5' and 3'homologous sequence for facile
cloning into the p01sgRNA plasmid. The library also included 400 non-targeting sgRNAs with

randomly generated spacers. The pooled sgRNA library was amplified via PCR using primers
oCAH1416/1417 and Q5 polymerase (New England Biolabs) following the manufacturers
recommended reaction conditions for 20 cycles to limit biased amplification. The amplicons
were assembled with p01sgRNA^{mrfp} amplified with oCAH1190/119 via isothermal assembly and
transformed into high-efficiency DH10b competent cells (New England Biolabs). Prior to plating
on selective medium, the competent cells were serially diluted to determine transformant colony
forming units (CFU/mL). 1.02×10^7 DH10b transformants were obtained after transformation,
three orders of magnitude greater than the synthesized library size. Twenty antibiotic-resistant
transformants were screened via PCR and Sanger sequencing to determine p01sgRNA^{mrfp}
background, which was < 5 %. The transformants were then pooled and the library miniprepped
using the Monarch Plasmid Miniprep Kit (New England Biolabs). The mixed p01sgRNA library
was then transferred to chemically competent S17 *E. coli* obtaining 1.15×10^6 S17
transformants, which were conjugated with *M. capsulatus* Bath strain with integrated P$_{tet}$-dCas9
to generate the methanotroph CRISPRi library. *M. capsulatus* Bath transformants were
quantified via serial dilution plating and pooled for genomic DNA extraction and long-term -
80°C storage. 6.8×10^7 transconjugants were obtained, 1000X the amount of unique sgRNAs in
the synthesized library. Library composition was determined via Oxford Nanopore Technologies
amplicon sequencing as described below.
208
Oxford Nanopore Technologies sequencing and bioinformatic analysis. The sgRNA library
was cultivated in liquid NMS with or without CRISPRi induction. After 72 h of selection,
genomic DNA was extracted from an OD = 2 equivalent cell pellet using the Zymo DNA
extraction kit following the manufacturers protocol. 50 ng pooled sgRNA library was used as a
template to amplify a ~250 bp amplicon containing the sgRNA sequence with Q5 polymerase
using primers oCAH1416/1417 and the same reaction conditions as above (20 cycles). The
amplicons were barcoded with the native barcoding kit 24 V14 following the manufacturers
protocol (Oxford Nanopore Technologies, ONT) and sequenced using a MinION sequencing
device with high accuracy basecalling. Fastq sequences were then trimmed and filtered using
Geneious Prime software. The 20 bp sgRNA target sequences were extracted from the trimmed
and filtered sequence reads using a custom Python script to process ONT sequence data stored as
compressed .fasta.gz files. The script performs a two-step search for the constant sgRNA
sequences (TATTTTACCACTCCCTATCAGTGAT, TTTTAGAGCTAGAAATAGCAAGTT
or their reverse complements ATCACTGATAGGGAGTGGTAAAATA,
AACTTGCTATTTCTAGCTCTAAAAC) flanking the target sequence and extracts the adjacent
5' or 3' 20 bp target sequence from the read dependent on the identified query sequence/read
strand orientation. The script utilizes pairwise alignment to refine the search results, ensuring
accurate extraction of relevant sequences. After extraction, the sgRNA target sequences were
aligned to either a custom reference library containing the 45,798 sgRNA library sequences or
the *M. capsulatus* Bath genome (NC_002977.6); mapped reads were quantified and normalized
as sgRNA target sequence per million (TPM); and the fitness score (sgRNA target sequence

differential abundance between induced and uninduced samples) was determined following a
DEseq2 workflow using Geneious software. Enrichment analysis of the most differentially
abundant sgRNA targets was performed using ShinyGo 0.82 using the STRING database[47].
233
Co^{2+}-deplete cultivation, gas chromatography and biomass productivity calculations. Wild-
type *M. capsulatus* solid medium-derived biomass was inoculated into acid-washed serum vials
containing 10 mL NMS medium with decreasing $CoCl_2$ in the trace elements to $OD_{600} = 0.01$ and
growth was measured with CGQ optical sensors. All medium components were made with
HPLC-grade water in acid-washed glassware. The CH_4 in the headspace was determined 10 min
after gas addition and every 24 h during methanotroph cultivation using an SRI 8610c gas
chromatograph with TCD and FID detectors (SRI Instruments). CH_4 was quantified as %
atmospheric composition by comparison to known standards and converted to weight using the
known headspace volume and specific gravity (0.622 mg/mL at 37°C). Dry cell weight (DCW)
was determined by converting OD_{600} to DCW using a formula empirically determined by
comparing OD_{600} measurements to colony forming units during serum vial cultivation ($OD_{600} = 1$
is equivalent to 0.254 ± 0.0249 g DCW/L).
246
247 ***Illustrations.*** Data were graphed using GraphPad Prism 10 software. Circos plots were generated
248 using Circa (https://circa.omgenomics.com/). Workflow schematics were created with
249 BioRender.com.
250
251 ***Statistical analysis.*** Statistical analysis was performed using GraphPad Prism 10 software. The
252 data between two groups were analyzed using unpaired t tests. Data were considered statistically
253 significant when $p \leq 0.05$.

Results and Discussion

Construction and validation of methanotroph CRISPRi tools. We previously developed a
two-plasmid, broad-host-range CRISPR-Cas9 system that enabled *M. capsulatus* genome editing
[44]. This two-plasmid system consisted of a 1) pBBR-based plasmid with the *M. capsulatus*
constitutive PQQ-dependent methanol dehydrogenase P_{mxaF} promoter driving single guide RNA
(sgRNA) expression, and a compatible 2) IncP-based plasmid, pCAH01, with
anhydrotetracycline-inducible Cas9 expression (pCAH01^{SpR}::*cas9*)[44]. To construct a single
plasmid CRISPRi system, we moved the P_{mxaF}-sgRNA module to pCAH01^{SpR}::*cas9* and replaced
the *cas9* gene with the dead cas9 (*dcas9*) gene encoding the nuclease-deficient dCas9 variant
(D10A, H840A) creating pJHCi (Figure 1a). The ability of the CRISPRi system to repress
transcription in the Gammaproteobacterial methanotroph *M. capsulatus* Bath and
Alphaproteobacterial methanotroph *Methylosinus trichosporium* OB3b was tested using strains
expressing an mRFP reporter (Figure 1b)[48]. pJHCi with a sgRNA targeting the non-template
strand of the mRFP open reading frame (sgRNA^{mRFP}) was transferred to the methanotroph
reporter strains via conjugation. The CRISPRi system reduced *M. capsulatus* and *M.
trichosporium* mRFP fluorescence by 6.3-fold and 2.3-fold, respectively, after 72 h of bacterial
cultivation in liquid NMS with CH_4 as the carbon source (Figure 1c). The decreased efficiency of
the CRISPRi system in *M. trichosporium* may be related to differences in sgRNA and/or dCas9
abundance as promoters derived from *Gammaproteobacteria*, like those driving transcription of
the sgRNA (*M. capsulatus* P_{mxa}) and dCas9 (P_{tet}), exhibit lower activity in *M. trichosporium*
compared to *M. capsulatus*[48].

Figure 1. Construction and validation of a methanotroph CRISPRi system. a) Schematic overview of
converting a two-plasmid CRISPR genome editing system into a single-plasmid CRISPRi system for targeted
repression of gene expression. b and c) The mRFP fluorescent reporter was used as a read-out of CRISPRi
functionality in Gammaproteobacterial methanotroph *Methylococcus capsulatus* Bath and Alphaproteobacterial
methanotroph *Methylosinus trichosporium* OB3b with (+) or without (-) dCas9 induction by anhydrotetracycline
(aTc). A representative image of cell pellets is shown. Data in c represent the mean ± SEM from two independent
experiments (n=3-5). ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$.

The pJHCi CRISPRi system limited the production of fluorescent protein in the absence of dCas9 induction in both methanotrophs (Figure 1c), suggesting that low dCas9 expression from leaky P_{tet} promoter gene transcription in the context of high, constitutive sgRNA expression is sufficient for transcription repression. The leaky functionality of the CRISPRi system is not desirable for functional genomic applications where sgRNA expression could cause fitness defects during strain or library generation. Thus, to develop a system with more tightly regulated CRISPRi repression, we designed a configuration wherein both dCas9 and the sgRNA are inducible (Figure 2a). mRFP fluorescence in this strain was similar to a control strain with the mRFP reporter plasmid alone (Figure 2b), supporting that the CRISPRi system did not inhibit mRFP transcription in the absence of aTc induction. In contrast, simultaneous induction of dCas9 and sgRNA^{mRFP} led to a 3.3-fold decrease in mRFP fluorescence (Figure 2b). Fluorescence microscopy was used to assess CRISPRi efficiency in individual cells. As shown in Figure 2c, CRISPRi repressed mRFP fluorescence in most of the bacterial population, but some cells exhibited high mRFP fluorescence indicating a small percentage of the population may "escape" CRISPRi repression.

Figure 2. A CRISPRi system with inducible dCas9 and sgRNA enables fine-tuned transcription repression in *M. capsulatus*. a) The pJHCi single plasmid system exhibited leaky functionality, so an *M. capsulatus* Bath strain was generated with an aTc-inducible dCas9 cassette integrated into the chromosome and plasmid with aTc-inducible sgRNA expression. b) mRFP fluorescence with or without aTc induction compared to the pDS1 mRFP reporter plasmid control strain showing tightly regulated CRISPRi gene transcription repression. c) Fluorescence microscopy of *M. capsulatus* with or without CRISPRi induction for mRFP repression. Representative images from biological replicates are shown. Data in b represent the mean ± SEM from two independent experiments (n=3-5). ns-not significant, *** p ≤ 0.001.

CRISPRi-mediated transcription repression of *M. capsulatus* RubisCO causes growth inhibition. We previously showed that the RubisCO enzyme encoded by *M. capsulatus* is essential for its growth on CH_4[49]. As such, we targeted the *cbbL* gene encoding the RubisCO large subunit to test the ability of the CRISPRi system to repress transcription from the *M. capsulatus* chromosome. Transconjugants could only be obtained in the absence of sgRNA[cbbL] and dCas9 induction (Figure 3a), supporting the essentiality of RubisCO for *M. capsulatu*s growth[49]. Transconjugants isolated in the absence of induction were subsequently cultivated in liquid NMS with or without the aTc inducer, and high-resolution growth analysis showed that CRISPRi-repression of *cbbL* transcription significantly inhibited bacterial growth (Figure 3b). Notably, the induced culture began to expand after five days and exhibited typical logarithmic growth kinetics thereafter (Figure 3b). The aTc inducer has a 17 h half-life in minimal medium at pH 7 and 37°C [50], so we hypothesized that the delayed growth observed in induced samples may be due to ineffective aTc concentrations after five days cultivation at 37°C. However, we did not observe a growth defect when the induced, outgrown sgRNA[cbbL] culture was diluted into fresh induction medium (data not shown), indicating that the CRISPRi system is no longer effective in this expanded population. Known mutations in the dCas9 gene[51] or others in the sgRNA, P*tet-tetO* promoter, or the *tetR* gene that would render the CRISPRi system nonfunctional were not identified in five "escapee" colonies via Sanger sequencing (data not shown), so the underlying mechanism of resistance to CRISPRi repression in these strains remains elusive. Given the essentiality of RubisCO for *M. capsulatus* growth, it is likely there is a strong selective pressure to overcome *cbbL* transcription repression in these cells.

Figure 3. CRISPRi-mediated transcription repression of the genes encoding the essential ribulose 1,5 bisphosphate carboxylase/oxygenase (RubisCO) enzyme limits *M. capsulatus* growth. Growth of *M. capsulatus* with (+) or without (-) induction of dCas9 and an sgRNA targeting the *cbbL* gene encoding the large subunit of RubisCO on solid NMS medium (a) or in liquid NMS medium in a serum vial over time (b) with a 20% CH_4 in air atmosphere. c) A complementing plasmid expressing the *cbbLSQ* operon (pLSQ) was constructed and the protospacer adjacent motif (PAM) site in the *cbbL* target sequence (MCA2743) was mutated (c.57C>T). d) Growth of strains carrying the pLSQ or pLSQ$^{\Delta PAM}$ compared to an isogenic background control strain with or without CRISPRi induction. Data in b and d represent the means of two independent high-resolution cultivation experiments (n=6).

To further test the specificity of the CRISPRi system, we constructed a plasmid containing the *cbbLSQ* operon under control of its native promoter region (pLSQ) and another plasmid with a mutated *cbbL* target protospacer adjacent motif (PAM) (MCA2743:c.57C>T, pLSQ$^{\Delta PAM}$) expected to prevent dCas9 DNA binding[52][53] (Figure 3c). CRISPRi targeting of *cbbL* in the pLSQ$^{\Delta PAM}$ strain did not inhibit growth (Figure 3d), supporting that expression of RubisCO from this plasmid was not inhibited and able to complement chromosomal *cbbLSQ* transcription repression. In contrast, CRISPRi-mediated repression of *cbbL* transcription in the strain with pLSQ inhibited bacterial growth, but not to the extent of that observed in the strain lacking the plasmid (Figure 3d). The pLSQ plasmid has a pBBR-based replicon, which is maintained in *M. capsulatus* at ~50 copies/cell[48]. As such, the difference in growth inhibition observed between the pSLQ and control strain may be correlated to an increase in target sites in cells harboring pSLQ, thereby limiting CRISPRi transcription repression efficiency compared to targeting a single chromosomal site.

CRISPRi functional genomic screening identifies genes essential for CH$_4$-dependent growth. To evaluate the utility of our CRISPRi system for functional genomic screening, we designed and synthesized a pooled sgRNA library consisting of 45,798 sgRNAs, ~15 sgRNAs per *M. capsulatus* protein-coding genes (3,034 annotated in the NC_002977.6 RefSeq genome) and 400 non-targeting control sgRNAs (supplemental Table S1-S2). Eighteen open-reading frames had less than 15 sgRNAs in the library and zero sgRNA target sequences were identified for 3 genes (MCA0050, MCA0307, MCA1517, supplemental Table S3). The workflow for cloning and screening the sgRNA library is shown in Figure 4a. We determined the *M. capsulatus* library coverage by amplifying the sgRNAs from genomic DNA isolated from 6.8 x 10^7 pooled transconjugants. The sgRNA amplicons were sequenced using an Oxford Nanopore Technologies MinION device; 680,425 reads with QC 12+ scores were trimmed and filtered using Geneious software followed by extraction of the 20 bp sgRNA targeting sequences from the processed sequences using a custom script. In total, 680,304 extracted sgRNA spacers were aligned to the synthesized genome-wide CRISPRi sgRNA library reference with 45,554 of the 45,798 synthesized sgRNAs detected (99.5% sgRNA library coverage, supplemental Table S4) with at least one sgRNA per predicted DNA coding sequence (except the three genes originally not included in the synthesized library, supplemental Table S5).

The methanotroph genome-wide sgRNA library was cultivated in serum vials with an atmosphere of 20% CH$_4$ in air with and without CRISPRi induction to identify genes essential for *M. capsulatus* growth. We observed a minor growth defect of CRISPRi induced cultures compared to uninduced cultures, but logarithmic growth kinetics and entry into stationary phase were similar between the conditions (Figure 4b). The sgRNA library compositions in CRISPRi induced and noninduced samples 72 h after selection were quantified by sgRNA amplicon sequencing and mapping to both the synthesized sgRNA library and the *M. capsulatus* genome (supplemental Tables S6-S13, read counts). We identified between 61.1-91.8% of the synthesized sgRNAs in the sample sequences, with increased coverage strongly correlated to sequencing depth (supplemental Figure S1, $R^2 = 0.93$). Although complete library coverage was not obtained in the screening samples with the sequencing depth achieved here, all gene targets were represented in the sequence data with 3,027/3,034 (99.8%) coding DNA sequences with ≥10 guides represented (supplemental Tables S6-S13). Comparing the sgRNA library composition between CRISPRi induced vs noninduced samples showed that 1,019 sgRNAs were

depleted (log2 fitness score ≤ 1, $p ≤ 0.001$) and 59 sgRNAs were enriched (log2 fitness score ≤
1, $p ≤ 0.001$) after 72 h of selection, presumably corresponding to a fitness defect or advantage
phenotype associated with transcription repression of the sgRNA target gene (Figure 4c and
Table S14). Enrichment analysis of 429 genes with sgRNA log2 fitness scores ≤ -3 and
probability values ≤ 0.001 indicated that these genes encode proteins associated with essential
cellular processes like translation, transcription, ribosome biosynthesis, and amino acid and
nucleotide biosynthesis (Figure 4d and Table S15). Enrichment analysis of the genes whose
transcription repression resulted in a fitness advantage did not enrich any pathways using the
available ShinyGO databases. Notably, 99.5% (398/400) of the non-targeting control sgRNAs
did not have significant differential fitness scores (log2 ≤-1 or ≥1, $p ≤ 0.001$) between the two
conditions (supplemental Table S14).

396
Figure 4. A CRISPRi genome-wide screen identifies genes essential for CH₄-dependent growth of *M.*
capsulatus. a) 45,798 sgRNAs were synthesized via massively parallel oligonucleotide synthesis and cloned into the

inducible plasmid pCAH01 to generate a cloned sgRNA library. The sgRNA library was transferred to an *M.*
capsulatus Bath strain with a chromosomally integrated P_{tet}-dCas9 cassette via biparental mating. b) The library was
cultivated with or without sgRNA and dCas9 induction in NMS medium in serum vials with a 20% CH_4 in air
atmosphere. After selection, the sgRNA composition and abundances were determined via Oxford Nanopore
Technologies amplicon sequencing and bioinformatic analysis to quantify sgRNA fitness. c) Circos plot of the
methanotroph sgRNA library raw reads (outer ring) mapped to the 45,798 synthesized sgRNA oligos (rainbow ring
with non-targeting sgRNAs in blue). Differential sgRNA fitness scores determined via DeSeq2 between CRISPRi
induced (n=4) and uninduced (n=4) samples during selection/cultivation with CH_4 as the sole carbon and energy
source (4th ring, blue dots log2 >1; red dots log2 <1). sgRNAs depleted in the screen (negative fitness scores)
correspond to genes with transcription repression causing a growth defect where those enriched (positive fitness
scores) correspond to genes with transcription repression causing a growth advantage. sgRNAs with statistically
significant ($p \leq 0.01$) differential fitness scores are indicated (blue squares, 3rd ring). d) ShinyGo enrichment analysis
of the 429 genes with sgRNA log2 fitness scores $\leq$ -3 and $p \leq 0.001$. e) CH_4 assimilatory and dissimilatory
metabolic pathways predicted by sgRNA fitness scores. The mean sgRNA fitness scores are indicated in parentheses
next to the gene loci that encode metabolic enzymes. 3PG, 3-phosphoglycerate; G3P, glyceraldehyde 3-phosphate;
F1,6P, fructose 1,6-bisphosphate; F6P, fructose 6-phosphate; E4P, erythrose 4-phosphate; Xu5P, xylulose 5-
phosphate; S7P, sedoheptulose 7-phosphate; R5P, ribose 5-phosphate; Ru5P, ribulose 5-phosphate; R1,5P, ribulose
1,5-bisphosphate; G6P, glucose-6-phosphate; 6PG, 6-phosphogluconate; KDPG, 2-keto-3-deoxy-6-
phosphogluconate; Pyr, pyruvate; H4F=CH2, methylene-tetrahydrofolate; H4MPT=CH2, methylene-
tetrahydromethanopterin; H4F≡CH, methenyl-tetrahydrofolate; H4MPT≡CH, methenyl-tetrahydromethanopterin;
H4F-CHO, formyl-tetrahydrofolate; H4MPT-CHO, formyl-tetrahydromethanopterin. Scores with $p \geq 0.01$ are
italicized. Data in b represent the means of two independent high-resolution cultivation experiments (n=6).
421
Several sgRNAs targeting known central CH_4 metabolic enzyme genes had significant negative
fitness scores (Figure 4e and supplemental Table S14), including those targeting the particulate
methane monooxygenase operons (MCA1796-98 and MCA2853-55), the calcium- and PQQ-
dependent methanol dehydrogenase (MCA0779), and the RuMP cycle operons (MCA3040-45
and 3046-50). As expected based on our initial CRISPRi experiments (Figure 3), sgRNAs
targeting the *cbbLSQ* operon (MCA2743-46) were among those with the lowest fitness scores.
Further, the sgRNAs targeting the Entner-Doudoroff pathway *edd* and *eda* genes (MCA0037-38)
were significantly depleted in the screen, consistent with biochemical data and genome-scale
models predicting this pathway to be essential and the primary route for pyruvate biosynthesis in
M. capsulatus Bath[5,54,55]. Also, the depletion of sgRNAs targeting the tetrahydromethanopterin
formaldehyde dissimilatory pathway corroborate flux balance model predictions that this is the
primary route used by *M. capsulatus* for formaldehyde conversion to formate[54,55]. Notably,
sgRNAs targeting genes encoding serine cycle enzymes showed no differential fitness during
growth on CH_4.
436
Many sgRNAs targeting the same gene or genes within operons exhibited similar fitness scores,
highlighting the potential of the CRISPRi library and screening pipeline to identify functional
gene targets (supplemental Table S14). For example, sgRNAs targeting ribosomal protein operon
genes MCA2368, MCA2371, MCA2372, and MCA2374 were consistently depleted under
CRISPRi induction conditions with log2 fitness scores of -5.47, -5.57, -5.70, and -6.37,
respectively, while sgRNAs targeting the cobalt transporter and cobalamin biosynthetic operon
genes MCA2300, MCA2301, and MCA2302 were significantly enriched with log2 fitness scores
of 3.10, 2.70 and 3.65, respectively. However, significant fitness score variances between
sgRNAs targeting the same gene were observed, suggesting CRISPRi effectiveness in repressing
transcription can vary depending on the sgRNA target sequence. Further evaluation of the
individual *pmoCAB*- and *cbbLSQ*-targeting sgRNAs showed highly variable fitness scores,
especially those targeting, *pmoC*, *pmoA*, and the *cbbL* genes (supplemental Figure S2a). To

confirm this variability, we tested the ability of three *pmoC*-targeting sgRNAs with variable
fitness scores (sgRNA^{pmoC1}:log2 = 0.34; sgRNA^{pmoC2}: log2 = -1.11*;* sgRNA^{pmoC3}: log2 = -2.10) to
inhibit *M. capsulatus* growth. The extent of bacterial growth inhibition was correlated to the
respective sgRNA fitness scores (supplemental Figure S2b). The high-GC content of the *M.*
capsulatus genome increases the possibility of sgRNA secondary structure formation, which may
limit sgRNA targeting efficiency or render some nonfunctional [56,57].
We next mapped the sgRNA target sequences from CRISPRi induced and noninduced cultures
to the *M. capsulatus* Bath genome to determine overall sgRNA/gene fitness. Given the size of
the sgRNA library and multiple sgRNAs per ORF, only those genes with overall positive or
negative sgRNA fitness scores were expected to be identified in the analysis. We identified 233
genes whose transcription repression resulted in a fitness defect (sgRNAs depleted in the induced
samples, log2 ≤ -1, $p ≤ 0.001$) and repression of 13 genes associated with a fitness advantage
(sgRNAs enriched in the induced samples, log2 ≥ 1, $p ≤ 0.001$) (supplemental Figure S3a and
supplemental Table S16). Enrichment analysis of the 233 genes with significant sgRNA log2
fitness scores ≤ -1 linked the genes to the essential processes of translation, transcription,
ribosome biosynthesis, and amino acid and nucleotide biosynthesis (supplemental Figure S3b
and supplemental Table S17), consistent with the analysis using the sgRNA reference library.
The *edd* gene encoding the ED pathway 6-phosphogluconate dehydratase had the most negative
differential fitness score (log2 = -3.9, $p = 1.63e^{-71}$, Table 3). Several putative essential genes are
involved in antioxidant defenses, Fe-S cluster assembly/repair, and disulfide reduction, including
superoxide dismutase (MCA2458, log2 = -3.2), a putative Rnf electron transport complex
(MCA2893-2899, log2 = -3.1), ferredoxin (MCA2470, log2 = -3.0), and thioredoxin disulfide
reductase (MCA1694, log2 = -2.6). These genes may be involved in balancing redox
homeostasis during CH_4 oxidation[58,59] and represent gene targets for downstream gene-function
analysis. sgRNAs targeting genes in the cobalamin biosynthetic operon (MCA2290-MCA2302)
had consistent positive fitness scores and MCA2302, encoding a putative CbtB subunit of a
cobalt transporter, was identified as significantly enriched (log2 = 1.26, $p = 3.9e^{-11}$) when
mapping all sgRNAs to the *M. capsulatus* genome (Table 3 and supplemental Table S16). As
expected, based on the variable fitness scores for sgRNAs targeting the pMMO and RubisCO
operons (supplemental Figure S2), only the *cbbS* (encoding the RubisCO small subunit) and
cbbQ (encoding a putative RubisCO activase) genes with consistent sgRNA scores were
associated with negative fitness scores when mapping to the genome. Interestingly, another
putative RubisCO activase gene (MCA2754, log2 = -3.1, $p = 2.4e^{-22}$) may be essential for *M.*
capsulatus growth on CH_4. Evaluating the RubisCO activases under differential growth
conditions may provide insight into their roles in methanotrophic growth of *M. capsulatus*.
484
To validate our screening results, we individually targeted five genes with negative fitness scores
involved in *M. capsulatus* central metabolism (*edd*, MCA0037, encoding phosphogluconate
dehydratase of the Entner-Doudoroff pathway); DNA replication (*dnaA*, MCA3033, encoding
the DNA replication initiator), redox metabolism (*fdx*, MCA2470, encoding a ferredoxin;
ribosome biogenesis (r*psB*, MCA0567, encoding the S2 protein component of the ribosome 30S
subunit); and amino acid biosynthesis (*asnB*, MCA2127, encoding the asparagine synthase); and
one gene with a positive fitness score (*cbtB*, MCA2302, encoding a putative cobalt
transporter)(Table 3). Consistent with the screening results, high-resolution growth analysis of
M. capsulatus strains expressing individual sgRNAs displayed significantly inhibited (*edd, dnaA,*
fdx, rpsB, asnB) or improved (*cbtB*) growth compared to uninduced controls or a non-targeting

sgRNA (Figure 5a-f, Figure 6a). Like that observed when targeting the *cbbL* gene (Figure 3),
growth-inhibited strains began to expand after several days of selective pressure. These data
validate that the CRISPRi system is effective for functional genomic screening and identification
of genes required for CH$_4$-dependent growth.
499

500
Figure 5. CRISPRi targeted transcription repression of putative essential genes inhibits CH$_4$-dependent
growth. a-f) High resolution growth curves of strains with sgRNAs targeting genes identified in the genome-wide
screen as essential for CH$_4$-dependent growth with (blue line) or without (purple line) CRISPRi induction in NMS
medium in serum vials with 20% CH$_4$ in air atmosphere. A strain expressing an sgRNA targeting mRFP was
included as a control (a). Genes represent those encoding enzymes involved in amino acid biosynthesis (b, *asnB*,
MCA2127); DNA replication (c, *dnaA*, MCA3033); CH$_4$ assimilation (d, *edd*, MCA0037); redox homeostasis (e,
fdx, MCA2470); and ribosome biosynthesis (f, *rpsB*, MCA0567). Data represent the mean of two independent high-
resolution cultivation experiments (n=6).
509

Cobalt is an essential trace element; however, excess can limit bacterial growth by competing
with iron and through the generation of reactive oxygen species [60,61]. Based on the growth
improvement observed when transcription of the CbtB cobalt transporter was inhibited (Figure
7a), we hypothesized that the trace amount of cobalt present in our NMS growth medium (1 µM
CoCl$_2$) was cytotoxic. Consistent with this, *M. capsulatus* cultivated in NMS with decreasing
Co^{2+} concentrations displayed a dose-dependent growth improvement, primarily linked to
reduced lag phase (Figure 6b). *M. capsulatus* exhibited a 1.4-fold improvement in biomass yield
within 48 h in NMS medium with 0.01 µM CoCl$_2$ compared to standard NMS (Figure 6c, 0.65
vs 0.48 g DCW/g CH$_4$, p = 0.006), indicating higher CH$_4$ flux to biomass under these conditions.
The improved methanotroph growth kinetics in the optimized growth medium equates to a 4.7-
fold improvement in methanotroph biomass productivity (Figure 6d). Methanotroph biomass (i.e.
single cell protein) is currently generated industrially as a valuable CH$_4$-derived product used in
the production of animal feed and agricultural fertilizer. Technoeconomic analyses have
identified methanotroph productivity as a primary cost driver of methane bioconversion
processes[62,63]; thus the optimized medium formulation identified here would be expected to
significantly decrease single cell protein production costs.

Figure 6. CRISPRi screen-guided growth medium optimization improves methanotroph biomass productivity from CH₄. High resolution growth curves of a) a strain with an sgRNA targeting the *cbtB* gene with (blue line) or without (purple line) CRISPRi induction or b) wild-type *M. capsulatus* in NMS medium with decreasing CoCl₂ in the trace elements. c) CH₄ consumption (solid lines) measured by gas chromatography and biomass accumulation (dashed lines) between wild-type cells cultured in NMS medium with high (1 µM CoCl₂, blue symbols) or low (0.01 µ CoCl₂, brown symbols) in serum vials with 20% CH₄ in air atmosphere. d) *M. capsulatus* biomass productivity in high or low CoCl₂ NMS medium at 48h. Data in a and b represent the mean of two independent high-resolution cultivation experiments (n=6). Data in c and d represent the mean ± SEM from two independent experiments (n=6-10). **** $p \leq 0.0001$.

Conclusions

Collectively, our results demonstrate that the CRISPRi system developed here can be used for targeted transcription repression in phylogenetically diverse methanotrophic bacteria. Selective screening of a genome-wide sgRNA library during CH₄-dependent cultivation identified predicted and novel essential genes required for methanotrophic growth. The screen provides genetic evidence that the Entner-Doudoroff pathway is essential for CH₄ assimilation and the tetrahydromethanopterin pathway is the primary route for CH₄ dissimilation in *M. capsulatus*. Additionally, the decreased fitness of sgRNAs targeting the ribulose bisphosphate carboxylase/oxygenase (RubisCO) and RubisCO activase genes reiterates the essentiality of CO₂ assimilation for *M. capsulatus* growth on CH₄. Notably, our investigations led to the identification of an optimized growth medium for improved methanotroph biomass productivity. The CRISPRi system and associated datasets herein will serve as valuable resources for hypothesis-driven inquiry into genes required for methanotrophy or alternative gene-function analyses. Additionally, these CRISPRi tools can be coupled with other synthetic biology approaches to optimize methanotrophic strains for carbon- and energy-efficient CH₄ capture and conversion.

References

1.   Guerrero-Cruz, S. *et al.* Methanotrophs: discoveries, environmental relevance, and a
perspective on current and future applications. *Front. Microbiol.* **12**, 678057 (2021).

2.   Lidstrom, M. E. Direct methane removal from air by aerobic methanotrophs. *Cold Spring*
Harb. Perspect. Biol. **16**, (2024).

3.   Tucci, F. J. & Rosenzweig, A. C. Direct methane oxidation by copper- and iron-
dependent methane monooxygenases. *Chem. Rev.* **124**, 1288–1320 (2024).

4.   Kalyuzhnaya, M. G., Gomez, O. A. & Murrell, J. C. The Methane-Oxidizing Bacteria
(Methanotrophs). in *Taxonomy, Genomics and Ecophysiology of Hydrocarbon-*
Degrading Microbes (ed. McGenity, T. J.) 1–34 (Springer International Publishing,
2019). doi:10.1007/978-3-319-60053-6_10-1.

5.   Strom, T., Ferenci, T. & Quayle, J. R. The carbon assimilation pathways of
Methylococcus capsulatus, *Pseudomonas methanica* and *Methylosinus trichosporium*
(OB3b) during growth on methane. *Biochem. J.* **144**, 465–476 (1974).

6.   Heptinstall, J. & Quayle, J. R. Pathways leading to and from serine during growth of
Pseudomonas AM1 on C1 compounds or succinate. *Biochem. J.* **117**, 563–572 (1970).

7.   Lawrence, A. J., Kemp, M. B. & Quayle, J. R. Synthesis of cell constituents by methane-
grown *Methylococcus capsulatus* and *Methanomonas methanooxidans*. *Biochem. J.* **116**,
631–639 (1970).

8.   Chistoserdova, L., Vorholt, J. A., Thauer, R. K. & Lidstrom, M. E. C1 transfer enzymes
and coenzymes linking methylotrophic bacteria and methanogenic Archaea. *Science* **281**,
99–102 (1998).

9.   Vorholt, J. A. Cofactor-dependent pathways of formaldehyde oxidation in methylotrophic
bacteria. *Arch. Microbiol.* **178**, 239–249 (2002).

10.   Kalyuzhnaya, M. G. *et al.* Highly efficient methane biocatalysis revealed in a
methanotrophic bacterium. *Nat. Commun.* **4**, 2785 (2013).

11.   Vuilleumier, S. *et al.* Genome sequence of the haloalkaliphilic methanotrophic bacterium
Methylomicrobium alcaliphilum 20Z. *J. Bacteriol.* **194**, 551–552 (2012).

12.   Mateos-Rivera, A., Islam, T., Marshall, I. P. G., Schreiber, L. & Øvreås, L. High-quality
draft genome of the methanotroph *Methylovulum psychrotolerans* Str. HV10-M2 isolated
from plant material at a high-altitude environment. *Stand. Genomic Sci.* **13**, 10 (2018).

13.   Miroshnikov, K. K. *et al.* Draft genome sequence of *Methylocapsa palsarum* NE2T, an
obligate methanotroph from subarctic soil. *Genome Announc.* **5**, (2017).

14.   Kalyuzhnaya, M. G. *et al.* Draft genome sequences of gammaproteobacterial
methanotrophs isolated from lake Washington sediment. *Genome Announc.* **3**, (2015).

15.   Hamilton, R. *et al.* Draft genomes of gammaproteobacterial methanotrophs isolated from
terrestrial ecosystems. *Genome Announc.* **3**, (2015).

16.   Orata, F. D., Kits, K. D. & Stein, L. Y. Complete Genome Sequence of *Methylomonas*
denitrificans Strain FJG1, an Obligate Aerobic Methanotroph That Can Couple Methane
Oxidation with Denitrification. *Genome Announc.* **6**, (2018).

17.   Oshkin, I. Y. *et al.* Complete Genome Sequence of *Methylococcus capsulatus* MIR, a
Methanotroph Capable of Growth on Methanol. *Microbiol. Resour. Announc.* **11**,
e0054222 (2022).

18.   Stein, L. Y. *et al.* Genome sequence of the obligate methanotroph *Methylosinus*
trichosporium strain OB3b. *J. Bacteriol.* **192**, 6497–6498 (2010).

19.   Van Dien, S. J. *et al.* Reconstruction of C(3) and C(4) metabolism in *Methylobacterium*
extorquens AM1 using transposon mutagenesis. *Microbiology (Reading, Engl)* **149**, 601–
609 (2003).

20.   Marx, C. J., O'Brien, B. N., Breezee, J. & Lidstrom, M. E. Novel methylotrophy genes of
Methylobacterium extorquens AM1 identified by using transposon mutagenesis including
a putative dihydromethanopterin reductase. *J. Bacteriol.* **185**, 669–673 (2003).

21.   Lee, K. E., Stone, S., Goodwin, P. M. & Holloway, B. W. Characterization of transposon
insertion mutants of *Methylobacterium extorquens* AM1 (*Methylobacterium* strain AM1)
which are defective in methanol oxidation. *Microbiology* (1991).

22.   Chistoserdova, L. Metabolism of Formaldehyde in *M. extorquens* AM1: Molecular
genetic analysis and mutant characterization. in *Microbial growth on C1 compounds* (eds.
Lidstrom, M. E. & Tabita, F. R.) 16–24 (Springer Netherlands, 1996). doi:10.1007/978-
94-009-0213-8_4.

23.   Vorholt, J. A., Marx, C. J., Lidstrom, M. E. & Thauer, R. K. Novel formaldehyde-
activating enzyme in *Methylobacterium extorquens* AM1 required for growth on
methanol. *J. Bacteriol.* **182**, 6645–6650 (2000).

24.   Nunn, D. N. & Lidstrom, M. E. Phenotypic characterization of 10 methanol oxidation
mutant classes in *Methylobacterium* sp. strain AM1. *J. Bacteriol.* **166**, 591–597 (1986).

25.   Springer, A. L., Chou, H. H., Fan, W. H., Lee, E. & Lidstrom, M. E. Methanol oxidation
mutants in *Methylobacterium extorquens* AM1: identification of new genetic
complementation groups. *Microbiology (Reading, Engl)* **141 (Pt 11)**, 2985–2993 (1995).

26.   Whitta, S., Sinclair, M. I. & Holloway, B. W. Transposon Mutagenesis in
Methylobacterium AM1 (Pseudomonas AM1). *Microbiology* (1985).

27.   Roszczenko-Jasińska, P. *et al.* Gene products and processes contributing to lanthanide
homeostasis and methanol metabolism in *Methylorubrum extorquens* AM1. *Sci. Rep.* **10**,
12663 (2020).

28.   Ochsner, A. M. *et al.* Transposon sequencing uncovers an essential regulatory function of
phosphoribulokinase for methylotrophy. *Curr. Biol.* **27**, 2579-2588.e6 (2017).

29.   Groom, J. D., Ford, S. M., Pesesky, M. W. & Lidstrom, M. E. A Mutagenic Screen
Identifies a TonB-Dependent Receptor Required for the Lanthanide Metal Switch in the
Type I Methanotroph *Methylotuvimicrobium buryatense* 5GB1C. *J. Bacteriol.* **201**,
(2019).

30.   Sharpe, P. L. *et al.* Use of transposon promoter-probe vectors in the metabolic
engineering of the obligate methanotroph *Methylomonas* sp. strain 16a for enhanced C40
carotenoid synthesis. *Appl. Environ. Microbiol.* **73**, 1721–1728 (2007).

31.   Cheng, M., Pei, D., He, L., Fei, Q. & Yan, X. Cre/lox-Mediated CRISPRi Library
Reveals Core Genome of a Type I Methanotroph *Methylotuvimicrobium buryatense*
5GB1C. *Appl. Environ. Microbiol.* **89**, e0188322 (2023).

32.   Hawkins, J. S., Wong, S., Peters, J. M., Almeida, R. & Qi, L. S. Targeted transcriptional
repression in bacteria using CRISPR interference (CRISPRi). *Methods Mol. Biol.* **1311**,
349–362 (2015).

33.   de Bakker, V., Liu, X., Bravo, A. M. & Veening, J.-W. CRISPRi-seq for genome-wide
fitness quantification in bacteria. *Nat. Protoc.* **17**, 252–281 (2022).

34.   Hawkins, J. S. *et al.* Mismatch-CRISPRi Reveals the Co-varying Expression-Fitness
Relationships of Essential Genes in *Escherichia coli* and *Bacillus subtilis*. *Cell Syst.* **11**,
523-535.e9 (2020).

35.   Yao, L. *et al.* Pooled CRISPRi screening of the cyanobacterium *Synechocystis* sp PCC
6803 for enhanced industrial phenotypes. *Nat. Commun.* **11**, 1666 (2020).

36.   Shin, J. *et al.* Genome-wide CRISPRi screen identifies enhanced autolithotrophic
phenotypes in acetogenic bacterium *Eubacterium limosum*. *Proc Natl Acad Sci USA* **120**,
e2216244120 (2023).

37.   Mo, X.-H. *et al.* Establishment of CRISPR interference in *Methylorubrum extorquens*
and application of rapidly mining a new phytoene desaturase involved in carotenoid
biosynthesis. *Appl. Microbiol. Biotechnol.* **104**, 4515–4532 (2020).

38.   Wang, T. *et al.* Pooled CRISPR interference screening enables genome-scale functional
653   genomics study in bacteria with superior performance. *Nat. Commun.* **9**, 2475 (2018).

654   39.   Jiang, W., Oikonomou, P. & Tavazoie, S. Comprehensive Genome-wide Perturbations
655   via CRISPR Adaptation Reveal Complex Genetics of Antibiotic Sensitivity. *Cell* **180**,
656   1002-1017.e31 (2020).

657   40.   Qi, L. S. *et al.* Repurposing CRISPR as an RNA-guided platform for sequence-specific
658   control of gene expression. *Cell* **152**, 1173–1183 (2013).

659   41.   Enright, A. L., Heelan, W. J., Ward, R. D. & Peters, J. M. CRISPRi functional genomics
660   in bacteria and its application to medical and industrial research. *Microbiol. Mol. Biol.*
661   *Rev.* **88**, e0017022 (2024).

662   42.   Rumah, B. L. *et al.* In vivo genome editing in type I and II methanotrophs using a
663   CRISPR/Cas9 system. *ACS Synth. Biol.* **12**, 544–554 (2023).

664   43.   Jeong, J. *et al.* A highly efficient and versatile genetic engineering toolkit for a
665   methanotroph-based biorefinery. *Chemical Engineering Journal* **453**, 139911 (2023).

666   44.   Tapscott, T., Guarnieri, M. T. & Henard, C. A. Development of a CRISPR/Cas9 System
667   for *Methylococcus capsulatus* In Vivo Gene Editing. *Appl. Environ. Microbiol.* **85**,
668   (2019).

669   45.   Nath, S., Henard, J. M. & Henard, C. A. Optimized tools and methods for methanotroph
670   genome editing. *Methods Mol. Biol.* **2489**, 421–434 (2022).

671   46.   Emelianov, G. *et al.* Engineered *Methylococcus capsulatus* Bath for efficient methane
672   conversion to isoprene. *Bioresour. Technol.* **393**, 130098 (2024).

673   47.   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals
674   and plants. *Bioinformatics* **36**, 2628–2629 (2020).

675   48.   Bhat, E. H. *et al.* Construction of a broad-host-range Anderson promoter series and
676   particulate methane monooxygenase promoter variants expand the methanotroph genetic
677   toolbox. *Synthetic and Systems Biotechnology* **9**, 250–258 (2024).

678   49.   Henard, C. A. *et al.* Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase (RubisCO) Is
679   Essential for Growth of the Methanotroph *Methylococcus capsulatus* Strain Bath. *Appl.*
680   *Environ. Microbiol.* **87**, e0088121 (2021).

681   50.   Politi, N. *et al.* Half-life measurements of chemical inducers for recombinant gene
682   expression. *J. Biol. Eng.* **8**, 5 (2014).

683   51.   Zhao, H. *et al.* Depletion of Undecaprenyl Pyrophosphate Phosphatases Disrupts Cell
684   Envelope Biogenesis in *Bacillus subtilis*. *J. Bacteriol.* **198**, 2925–2935 (2016).

52.   Sternberg, S. H., Redding, S., Jinek, M., Greene, E. C. & Doudna, J. A. DNA
interrogation by the CRISPR RNA-guided endonuclease Cas9. *Nature* **507**, 62–67 (2014).

53.   Wu, X. *et al.* Genome-wide binding of the CRISPR endonuclease Cas9 in mammalian
cells. *Nat. Biotechnol.* **32**, 670–676 (2014).

54.   Gupta, A. *et al.* Genome-scale metabolic reconstruction and metabolic versatility of an
obligate methanotroph *Methylococcus capsulatus* str. Bath. *PeerJ* **7**, e6685 (2019).

55.   Lieven, C. *et al.* A Genome-Scale Metabolic Model for *Methylococcus capsulatus* (Bath)
Suggests Reduced Efficiency Electron Transfer to the Particulate Methane
Monooxygenase. *Front. Microbiol.* **9**, 2947 (2018).

56.   Riesenberg, S., Helmbrecht, N., Kanis, P., Maricic, T. & Pääbo, S. Improved gRNA
secondary structures allow editing of target sites resistant to CRISPR-Cas9 cleavage. *Nat.*
Commun. **13**, 489 (2022).

57.   Moreb, E. A. & Lynch, M. D. A Meta-Analysis of gRNA Library Screens Enables an
Improved Understanding of the Impact of gRNA Folding and Structural Stability on
CRISPR-Cas9 Activity. *The CRISPR Journal* **5**, 146–154 (2022).

58.   Jouanneau, Y., Jeong, H. S., Hugo, N., Meyer, C. & Willison, J. C. Overexpression in
Escherichia coli of the *rnf* genes from *Rhodobacter capsulatus*--characterization of two
membrane-bound iron-sulfur proteins. *Eur. J. Biochem.* **251**, 54–64 (1998).

59.   Alleman, A. B. & Peters, J. W. Mechanisms for Generating Low Potential Electrons
across the Metabolic Diversity of Nitrogen-Fixing Bacteria. *Appl. Environ. Microbiol.* **89**,
e0037823 (2023).

60.   Thorgersen, M. P. & Downs, D. M. Cobalt targets multiple metabolic processes in
Salmonella enterica. *J. Bacteriol.* **189**, 7774–7781 (2007).

61.   Leonard, S. *et al.* Cobalt-mediated generation of reactive oxygen species and its possible
mechanism. *J. Inorg. Biochem.* **70**, 239–244 (1998).

62.   Fei, Q. *et al.* Biological valorization of natural gas for the production of lactic acid:
techno-economic analysis and life cycle assessment. *Biochem. Eng. J.* 107500 (2020)
doi:10.1016/j.bej.2020.107500.

63.   El Abbadi, S. H., Sherwin, E. D., Brandt, A. R., Luby, S. P. & Criddle, C. S. Displacing
fishmeal with protein derived from stranded methane. *Nat. Sustain.* **5**, 47–56 (2021).

Acknowledgements
This work was supported by a BioDiscovery Institute Seed Grant to CH and RA, the Department of Energy via contract with the MSI STEM Research & Development Consortium (MSRDC) to CH, and the National Science Foundation MCB award # 2225776 to CH. We would like to thank Tallon Coxe in the Azad laboratory for computational support with initial sgRNA library design.

Author Contributions
CH conceived the project. JH, SL, and YCY performed experiments. CH acquired funding, analyzed and graphed data, and provided overall guidance. CH wrote the manuscript. All others edited and approved the manuscript.

Data Availability
Plasmids will be available at Addgene upon publication. ONT sequencing data will be deposited at the NCBI Sequence Read Archive (SRA) prior to publication. Code for sgRNA target sequence extraction can be downloaded from the Henard laboratory GitHub database (https://github.com/calvinhenard/CRISPRi-library).

732
733
Figure 1. Construction and validation of a methanotroph CRISPRi system. a) Schematic
overview of converting a two-plasmid CRISPR genome editing system into a single-plasmid
CRISPRi system for targeted repression of gene expression. b and c) The mRFP fluorescent
reporter was used as a read-out of CRISPRi functionality in Gammaproteobacterial
methanotroph *Methylococcus capsulatus* Bath and Alphaproteobacterial methanotroph
Methylosinus trichosporium OB3b with (+) or without (-) dCas9 induction by
anhydrotetracycline (aTc). A representative image of cell pellets is shown. Data in c represent
the mean ± SEM from two independent experiments (n=3-5). ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$.

743
Figure 2. A CRISPRi system with inducible dCas9 and sgRNA enables fine-tuned
transcription repression in *M. capsulatus*. a) The pJHCi single plasmid system exhibited leaky
functionality, so an *M. capsulatus* Bath strain was generated with an aTc-inducible dCas9
cassette integrated into the chromosome and plasmid with aTc-inducible sgRNA expression. b)
mRFP fluorescence with or without aTc induction compared to the pDS1 mRFP reporter plasmid
control strain showing tightly regulated CRISPRi gene transcription repression. c) Fluorescence
microscopy of *M. capsulatus* with or without CRISPRi induction for mRFP repression.
Representative images from biological replicates are shown. Data in b represent the mean ± SEM
from two independent experiments (n=3-5). ns-not significant, *** $p \leq 0.001$.

753
Figure 3. CRISPRi-mediated transcription repression of the genes encoding the essential
ribulose 1,5 bisphosphate carboxylase/oxygenase (RubisCO) enzyme limit *M. capsulatus*
growth. Growth of *M. capsulatus* with (+) or without (-) induction of dCas9 and an sgRNA
targeting the *cbbL* gene encoding the large subunit of RubisCO on solid NMS medium (a) or in
liquid NMS medium in a serum vial over time (b) with a 20% CH_4 in air atmosphere. c) A
complementing plasmid expressing the *cbbLSQ* operon (pLSQ) was constructed and the
protospacer adjacent motif (PAM) site in the *cbbL* target sequence (MCA2743) was mutated
(c57C>T). d) Growth of strains carrying the pLSQ or pLSQ$^{\Delta PAM}$ compared to an isogenic
background control strain with or without CRISPRi induction. Data in b and d represent the
means of two independent high-resolution cultivation experiments (n=6).

764
Figure 4. A CRISPRi genome-wide screen identifies genes essential for CH₄-dependent
growth of *M. capsulatus*. a) 45,798 sgRNAs were synthesized via massively parallel
oligonucleotide synthesis and cloned into the inducible plasmid pCAH01 to generate a cloned
sgRNA library. The sgRNA library was transferred to an *M. capsulatus* Bath strain with a
chromosomally integrated P$_{tet}$-dCas9 cassette via biparental mating. b) The library was cultivated
with or without sgRNA and dCas9 induction in NMS medium in serum vials with a 20% CH₄ in
air atmosphere. After selection, the sgRNA composition and abundances were determined via
Oxford Nanopore Technologies amplicon sequencing and bioinformatic analysis to quantify

sgRNA fitness. c) Circos plot of the methanotroph sgRNA library raw reads (outer ring) mapped to the 45,798 synthesized sgRNA oligos (rainbow ring with non-targeting sgRNAs in blue). Differential sgRNA fitness scores determined via DeSeq2 between CRISPRi induced (n=4) and uninduced (n=4) samples during selection/cultivation with CH_4 as the sole carbon and energy source (4th ring, blue dots log2 >1; red dots log2 <1). sgRNAs depleted in the screen (negative fitness scores) correspond to genes with transcription repression causing a growth defect where those enriched (positive fitness scores) correspond to genes with transcription repression causing a growth advantage. sgRNAs with statistically significant ($p \leq 0.01$) differential fitness scores are indicated (blue squares, 3rd ring). d) ShinyGo enrichment analysis of the 429 genes with sgRNA log2 fitness scores $\leq$ -3 and $p \leq 0.001$. e) CH_4 assimilatory and dissimilatory metabolic pathways predicted by sgRNA fitness scores. The mean sgRNA fitness scores are indicated in parentheses next to the gene loci that encode metabolic enzymes. 3PG, 3-phosphoglycerate; G3P, glyceraldehyde 3-phosphate; F1,6P, fructose 1,6-bisphosphate; F6P, fructose 6-phosphate; E4P, erythrose 4-phosphate; Xu5P, xylulose 5-phosphate; S7P, sedoheptulose 7-phosphate; R5P, ribose 5-phosphate; Ru5P, ribulose 5-phosphate; R1,5P, ribulose 1,5-bisphosphate; G6P, glucose-6-phosphate; 6PG, 6-phosphogluconate; KDPG, 2-keto-3-deoxy-6-phosphogluconate; Pyr, pyruvate; $H_4F=CH_2$, methylene-tetrahydrofolate; $H_4MPT=CH_2$, methylene-tetrahydromethanopterin; $H_4F\equiv CH$, methenyl-tetrahydrofolate; $H_4MPT\equiv CH$, methenyl-tetrahydromethanopterin; H_4F-CHO, formyl-tetrahydrofolate; H_4MPT-CHO, formyl-tetrahydromethanopterin. Scores with $p \geq 0.01$ are italicized. Data in b represent the means of two independent high-resolution cultivation experiments (n=6).

Figure 5. CRISPRi targeted transcription repression of putative essential genes inhibits
CH$_4$-dependent growth. a-f) High resolution growth curves of strains with sgRNAs targeting
genes identified in the genome-wide screen as essential for CH$_4$-dependent growth with (blue
line) or without (purple line) CRISPRi induction in NMS medium in serum vials with 20% CH$_4$
in air atmosphere. A strain expressing an sgRNA targeting mRFP was included as a control (a).
Genes represent those encoding enzymes involved in amino acid biosynthesis (b, *asnB*,
MCA2127); DNA replication (c, *dnaA*, MCA3033); CH$_4$ assimilation (d, *edd*, MCA0037); redox
homeostasis (e, *fdx*, MCA2470); and ribosome biosynthesis (f, *rpsB*, MCA0567). Data represent
the mean of two independent high-resolution cultivation experiments (n=6).

Figure 6. CRISPRi screen-guided growth medium optimization improves methanotroph biomass productivity from CH$_4$. High resolution growth curves of a) a strain with an sgRNA targeting the *cbtB* gene with (blue line) or without (purple line) CRISPRi induction or b) wild-type *M. capsulatus* in NMS medium with decreasing CoCl$_2$ in the trace elements. c) CH$_4$ consumption (solid lines) measured by gas chromatography and biomass accumulation (dashed lines) between wild-type cells cultured in NMS medium with high (1 µM CoCl$_2$, blue symbols) or low (0.01 µ CoCl$_2$, brown symbols) in serum vials with 20% CH$_4$ in air atmosphere. d) *M. capsulatus* biomass productivity in high or low CoCl$_2$ NMS medium at 48h. Data in a and b represent the mean of two independent high-resolution cultivation experiments (n=6). Data in c and d represent the mean ± SEM from two independent experiments (n=6-10). **** $p \leq 0.0001$.

815 **Tables**

Table 1. Strains and Plasmids		
Name	**Genotype**	**Source**
Methylococcus capsulatus **Bath**	Wild-type	Lab stock (originating from Kalyuzhnaya lab)
CH2236	*M. capsulatus* Bath with pDS1	Bhat et al[48]
CH2484	*M. capsulatus* Bath with pDS1 and pJHCi	This study
CH2486	*M. capsulatus* Bath with pJH1 and pJHCi^{sfgfp}	This study
CH2342	*M. capsulatus* Bath with inducible dCas9 integrated into the chromosome	This study
CH2475	CH2342 with p01sgRNA^{mrfp}	This study
CH2473	CH2342 with p01sgRNA^{cbbL}	This study
CH2529	CH2342 with p01sgRNA^{gwlibrary}	This study
CH2537	CH2342 with p01sgRNA^{MCA2127}	This study
CH2477	CH2342 with p01sgRNA^{MCA3033}	This study
CH2535	CH2342 with p01sgRNA^{MCA0037}	This study
CH2476	CH2342 with p01sgRNA^{MCA2470}	This study
CH2536	CH2342 with p01sgRNA^{MCA0567}	This study
CH2598	CH2342 with p01sgRNA^{MCA2302}	This study
CH2474	CH2342 with p01sgRNA^{pmoC-1}	This study
CH2478	CH2342 with p01sgRNA^{pmoC-2}	This study
CH25104	CH2342 with p01sgRNA^{pmoC-3}	This study
Methylosinus trichosporium **OB3b**	Wild-type	[18]
CH2507	*M. trichosporium* OB3b with pDS1	Bhat et al[48]
CH2508	*M. trichosporium* OB3b with pDS1 and pJHCi	This study
Escherichia coli **Zymo 10B**	F- *mcrA* Δ(*mrr-hsdRMS-mcrBC*) Φ80*lacZ*ΔM15 Δ*lacX*74 *recA1 endA1 araD139* Δ(*ara leu*) 7697 *galU galK rpsL nupG*	Zymo Research
E. coli **S17-1**	Tp^r Sm^r *recA thi pro hsd*(r⁻m⁺)RP4-2-Tc::Mu::Km Tn7	ATCC 47055
Plasmids		
Name	**Description**	**Source**
pCAH01^{SpR}	IncP broad-host-range inducible expression plasmid	Addgene #128159[44]
pCAH01^{SpR}::Cas9	Inducible Cas9 expression	Addgene #128160[44]
pdCas9-bacteria	aTc-inducible expression of catalytically inactive	Addgene #44249[40]

	S. pyogenes Cas9 (dCas9)	
pDS1	pBBR1MCS-5 with BBa_J23119-mRFP reporter	Bhat et al[48]
pJHCi	pCAH01^{SpR}::Cas9 *cas9* gene replaced with *dcas9* and *M. capsulatus* P$_{mxa}$ driving expression of mRFP targeting sgRNA	This study
pJHint	*tetO*-TetR inducible suicide plasmid that integrates into the *M. capsulatus* genome between MCA and MCA.	This study
pJHint::dCas9	pJHint with dCas9 gene for chromosomal integration of inducible dCas9 cassette	This study
pLSQ	*M. capsulatus cbbLSQ* operon with native promoter cloned into pBMTL-2	This study
pL$^{PAMC>T}$SQ	pCbbLSQ with mutated PAM site	This study
p01sgRNA^{mrfp}	pCAH01 with sgRNA targeting mRFP ORF	This study
p01sgRNA^{cbbL}	pCAH01 with sgRNA targeting MCA2743 (*cbbL*) ORF	This study
p01sgRNA^{MCA2127}	pCAH01 with sgRNA targeting MCA2127 (*asnB*) ORF	This study
p01sgRNA^{MCA3033}	pCAH01 with sgRNA targeting MCA3033 (*dnaA*) ORF	This study
p01sgRNA^{MCA0037}	pCAH01 with sgRNA targeting MCA0037 (*edd*) ORF	This study
p01sgRNA^{MCA2470}	pCAH01 with sgRNA targeting MCA2470 (*fdx*) ORF	This study
p01sgRNA^{MCA0567}	pCAH01 with sgRNA targeting MCA0567 (*rpsB*) ORF	This study
p01sgRNA^{MCA2302}	pCAH01 with sgRNA targeting MCA2301 (*cbtB*) ORF	This study
p01sgRNA$^{MCA1798/2855}$	pCAH01 with sgRNA targeting MCA (*pmoC-1*) ORF	This study
p01sgRNA$^{MCA1798/2855}$	pCAH01 with sgRNA targeting MCA (*pmoC-2*) ORF	This study
p01sgRNA$^{MCA1798/2855}$	pCAH01 with sgRNA targeting MCA (*pmoC-3*) ORF	This study

Table 2. Primers and synthetic DNA fragments

Name	Sequence
Construction of pJHCi	
oCAH230 pCAH01^SpR::Cas9 F	GTTCTTTCCTGCGTTATCCC
oCAH231 pCAH01^SpR::Cas9 R	GTCTGACGCTCAGTGGAA
oCAH267 P$_{mxa}$-sgRNA F	CGTTCCACTGAGCGTCAGACGAGGTTCAGGCGAAACCG
oCAH268 P$_{mxa}$-sgRNA R	GGGATAACGCAGGAAAGAACAAAAAAAGCACCGACTCGGT
oCAH4 pCAH01 F	AAGCTTGACCTGTGAAGTG
oCAH3 pCAH01 R	TTCACTTTTCTCTATCACTGATAG
oCAH952 dCas9 F	gtgatagagaaaagtgaaATGGATAAGAAATACTCAATAGGCTTAGCT
oCAH1507 dCas9 R	cttcacaggtcaagcttTCAGTCACCTCCTAGCTGACTCAAAT
Construction of pJHint and M. capsulatus strain with chromosomal dCas9 integration	
oCAH403 part 1 F	gcagcggaaaagCCCGTAGAAAAGATCAAAGG
oCAH404 part 1 R	cggatcgtgcgcATGTGAGCAAAAGGCCAG
oCAH405 part 2 F	cttttgctcacatGCGCACGATCCGTATTCC
oCAH406 part 2 R	ttaggaattaatcATCTGTCGAAGCGTGCCG
oCAH407 part 3 F	gcttcgacagatGATTAATTCCTAATTTTTGTTGAC
oCAH408 part 3 R	aacgagacatcatACTCTTCCTTTTTCAATATTATTG
oCAH409 part 4 F	aaaaggaagagtATGATGTCTCGTTTAGATAAAAGTAAAGTGATTAACAG
oCAH410 part 4 R	caagggccttgcGCGCGCTCTACGAACTGC
oCAH411 part 5 F	tcgtagagcgcgcGCAAGGCCCTTGGCTCTTG
oCAH412 part 5 R	cttcgcggagctATCTCTCGCAAGACGGCG
oCAH413 part 6 F	tcttgcgagagatAGCTCCGCGAAGTCGCTC
oCAH414 part 6 F	tcttttctacgggCTTTTCCGCTGCATAACCCTG
oCAH447 F	GACGACGACCGTCACAAA
oCAH448 R	GTTTTCGTGATTCGCCGC
Construction of pLSQ and MCA2743(cbbL) PAM mutation	
oCAH1007 pDS1 F	GGATCCAAACTCGAGTAAG
oCAH16 pDS1 R	TAGCCGCTTATGTCTATTGCTG
oCAH1629 P$_{cbbL}$ F	cagcaatagacataagcggctaTTATGCTAATCGCCCTTCATCCG
oCAH1750 *cbbQ* R	gagatccttactcgagtttggatccTCAGAAAAACGTCGCTACGG
oCAH1751 pLSQ F	TGACCGACCTCGACTATTAC
cbbL:c.57C>T synthetic fragment	CGGTAAACCAGCAATAGACATAAGCGGCTATTATGCTAATCGCCCTTCATCCGGCGGTAGCAGGCAGGCATATCAGTATAATGCGCGCTTATTGTCGTGATTATGACAATTTAGTTCCGTTTATGGATATCAGCGCTTATAGTGCCTCGCGAAAAGGGATCGCCCCGATGCCGGCTAGGGAACGGCGGTTCATTGCCATAGCGCTGTTGCAAAACGAGTAGGAGAAAACCATGGCTGTCAAAACATACAACGCGGGCGTCAAGGAATACCGCGAAACCTACTGGGATCCGAACTACACTCCCGCCGACACCGATCTGCTGGCGGTCTTCAAGATCACCCCTCAGCCGGGTGTG

	CCGCGCGAAGAAGCCGCCGCCGCCGTGGCCGCGGAATCG TCGACCGGCACCTGGACCACCGTCTGGACCGACCTGCTG ACCGACCTCGACTATTACAAGGGCCGCG
sgRNA exchange and sgRNA library generation	
oCAH1083 F	taccactccctatcagtgatTGGAACCGTACTGGAACTGC
oCAH1081 R	cttcacaggtcaagcttAAAAAAAGCACCGACTCGGT
oCAH1190 01*sgRNA^{mrfp}* F	GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCT
oCAH119 01*sgRNA* R	ATCACTGATAGGGAGTGGTAAAATAACTCTATCAATGAT AGAG
oCAH1084 *sgRNA^{cbbL}* oligo	TACCACTCCCTATCAGTGATCATTGTCGGCGGGAGTGTAG TTCGTTTTAGAGCTAGAAATAGCAAGTT
oCAH1735 *sgRNA^{MCA0037}* oligo	TACCACTCCCTATCAGTGATTGAGATCGAGGTAAGCGGC GGTTTTAGAGCTAGAAATAGC
oCAH1736 *sgRNA^{MCA0567}* oligo	TACCACTCCCTATCAGTGATCATCTGACGCATGGTAACTG GTTTTAGAGCTAGAAATAGC
oCAH1737 *sgRNA^{MCA2127}* oligo	TACCACTCCCTATCAGTGATGGGCCAGGCGCTCGATCATG GTTTTAGAGCTAGAAATAGC
oCAH1738 *sgRNA^{MCA2470}* oligo	TACCACTCCCTATCAGTGATGCAGTTTTCGGTCACAACAA GTTTTAGAGCTAGAAATAGC
oCAH1739 *sgRNA^{MCA3033}* oligo	TACCACTCCCTATCAGTGATCAGGTGTTGAACTGTTGCGG GTTTTAGAGCTAGAAATAGC
oCAH1893 *sgRNA^{MCA2302}* oligo	TACCACTCCCTATCAGTGATCGACATGGACCGGCAAGGA AGTTTTAGAGCTAGAAATAGC
oCAH1082 *sgRNA^{pmoC1}* oligo	TACCACTCCCTATCAGTGATCGCCGCAGCTGCACCACCGT TTTAGAGCTAGAAATAGC
oCAH1740 *sgRNA^{pmoC2}* oligo	TACCACTCCCTATCAGTGATCCGCACCCACAGGTAGAAC AGTTTTAGAGCTAGAAATAGC
oCAH1905 *sgRNA^{pmoC3}* oligo	TACCACTCCCTATCAGTGATGTACAGGAAATTCATCCAGT GTTTTAGAGCTAGAAATAGC
oCAH1416 sgRNA^{library} F	GTTGACACTCTATCATTGATAGA
oCAH1417 sgRNA^{library} R	TGATAACGGACTAGCCTTATTTT

818   *Lowercase sequence are homology arms for isothermal assembly. Bold sequence is the sgRNA*
819   *target sequence*

820 **Table 3. Genes with greatest differential sgRNA fitness scores.**

Name	Locus tag	Differential Fitness Score (Log2)	Differential Fitness Score p-value
edd CDS	MCA0037	-3.89	1.6E-71
ispC CDS	MCA0573	-3.59	7.2E-19
sodB CDS	MCA2458	-3.22	2.3E-18
rsxD CDS	MCA2896	-3.21	3.4E-23
dnaA CDS	MCA3033	-3.14	7.0E-22
nitric oxide reductase activation protein NorD CDS	MCA2754	-3.10	2.4E-22
formylmethanofuran dehydrogenase subunit B CDS	MCA2860	-3.08	7.9E-24
rsxG CDS	MCA2897	-3.06	1.1E-49
ferredoxin family protein CDS	MCA2470	-3.02	2.7E-17
frr CDS	MCA0570	-3.01	7.9E-22
rpsB CDS	MCA0567	-2.88	4.8E-22
HAD-IB family hydrolase CDS	MCA2087	-2.87	1.7E-15
rsxA CDS	MCA2893	-2.84	1.7E-19
formylmethanofuran dehydrogenase subunit A CDS	MCA2859	-2.70	6.2E-14
rpsS CDS	MCA2368	-2.68	1.2E-42
cysG CDS	MCA2089	-2.63	2.3E-11
tsf CDS	MCA0568	-2.57	3.9E-14
trxB CDS	MCA1694	-2.56	2.7E-23
pyrH CDS	MCA0569	-2.53	1.8E-13
asnB CDS	MCA2127	-2.43	9.4E-13
infB CDS	MCA1315	-2.36	1.2E-31
sufD CDS	MCA0990	-2.35	5.5E-22
rplD CDS	MCA2371	-2.34	9.5E-21
rplC CDS	MCA2372	-2.28	2.3E-23
CbbQ/NirQ/NorQ/GpvN family protein CDS	MCA2746	-2.24	1.3E-16
OadG family protein CDS	MCA2481	-2.23	2.8E-18
glyQ CDS	MCA2049	-2.22	3.1E-27
ispG CDS	MCA2483	-2.19	9.0E-17
DNA-binding protein CDS	MCA2958	-2.18	1.6E-08
ABC transporter substrate-binding protein CDS	MCA1278	-2.15	9.8E-17
bifunctional cobalt-precorrin-7 (C(5))-methyltransferase/cobalt-precorrin-6B (C(15))-methyltransferase CDS	MCA2296	-2.14	1.0E-12
phosphoribosylaminoimidazolesuccinocarboxamide synthase CDS	MCA0021	-2.08	6.3E-17
calcium/proton transporter CDS	MCA0545	-2.05	1.3E-14
OmpA family protein CDS	MCA1819	1.00	1.5E-08
YihY family inner membrane protein CDS	MCA0711	1.05	1.8E-05
atpA CDS	MCA0010	1.08	8.1E-10
cas5e CDS	MCA0932	1.10	1.2E-07
glutathione S-transferase N-terminal domain-containing protein CDS	MCA1959	1.11	2.9E-07
hypothetical protein CDS	MCA1324	1.14	6.5E-07
phosphoribulokinase CDS	MCA3051	1.15	7.6E-11
NUDIX hydrolase CDS	MCA0333	1.15	7.6E-13
PhoH family protein CDS	MCA1458	1.20	6.2E-07
CbtB-domain containing protein CDS	MCA2302	1.26	6.1E-11
MucB/RseB C-terminal domain-containing protein CDS	MCA1469	1.41	1.2E-08
CbtA family protein CDS	MCA2301	1.61	1.3E-11
RseA family anti-sigma factor CDS	MCA1470	1.62	1.1E-06

822   **Supplemental Material**

823
Figure S1. The relationship between sequencing depth and sgRNA library coverage. Oxford
Nanopore Technology sequencing reads were processed to extract sgRNA target sequences that
were mapped to the synthesized genome-wide sgRNA library. Mapped sequence reads varied
between samples. Linear regression analysis suggests 413,960 mapped reads are needed to
achieve 100% sgRNA library inclusion in downstream analyses, which is ~10x sequencing
depth.

Figure S2. Some sgRNAs targeting the same gene have variable differential fitness scores.
Mapped sgRNA target sequences (black rectangles) and corresponding differential fitness score
within the *M. capsulatus* a) *pmoCAB* and b) *cbbLSQ* operon genes encoding the particulate
methane monooxygenase and ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO) and
associated RubisCO activase, respectively. c) High resolution CH_4-dependent growth curves of
strains with sgRNAs with differential fitness scores (*pmoC-1*, *pmoC-2*, *pmoC-3*) targeting the
pmoC gene with (blue line) or without (purple line) CRISPRi induction. The growth curves
represent the mean of 4-6 independent observations. The mean of the three uninduced sgRNA
controls is shown.

Figure S3. A CRISPRi genome-wide screen identifies genes essential for CH$_4$-dependent
growth of *M. capsulatus*: mapping to the *M. capsulatus* genome. a) Circos plot of the
methanotroph sgRNA library raw reads (outer ring) mapped to the NC_002977 *M. capsulatus*
reference genome (2nd ring, genes in forward direction are in dark grey and reverse direction in
light grey). Differential sgRNA fitness scores determined via DeSeq2 between CRISPRi induced
and uninduced samples during selection/cultivation with CH$_4$ as the sole carbon and energy
source (4th ring, blue dots log2 >1; red dots log2 <1). sgRNAs depleted in the screen (negative
fitness scores) correspond to genes with transcription repression causing a growth defect where
those enriched (positive fitness scores) correspond to genes with transcription repression causing
a growth advantage. sgRNAs with statistically significant ($p \leq 0.01$) differential fitness scores
are indicated (blue squares, 3rd ring). b) ShinyGo enrichment analysis of the 233 genes with
sgRNA log2 fitness scores $\leq$ -1 and $p \leq 0.001$.

854   Supplemental Tables S1-S17 can be found here:

855   https://docs.google.com/spreadsheets/d/1I1Mwm8o_4b5vcYRpeE11nL-bj-

856   knuRPL/edit?usp=drive_link&ouid=103067052205615496775&rtpof=true&sd=true