1    Interaction of two tightly linked loci in an adaptive genomic hotspot explains

2    exploration behavior in juvenile Atlantic salmon in freshwater.

3    Tutku Aykanat[1*], Jaakko Erkinaro[2]

4        1.   Organismal and Evolutionary Biology Research Program, Faculty of Biological and Environmental

5             Sciences, University of Helsinki, Finland.

6        2.   Natural Resources Institute Finland (Luke), Oulu, Finland

## Abstract

1)   Traits that are important for adaptation may exhibit genetic correlation due to pleiotropy or as a result of

linkage. Understanding the genetic architecture of such correlations is important for predicting the

selection response of populations. Exploration in fishes is a behavioral trait by which individuals may find

habitats with better foraging and growth opportunities that subsequently improve their fitness. In Atlantic

salmon (*Salmo salar*), for example, all individuals are originated in larger rivers that females can lay eggs,

but some juveniles migrate to small tributaries that provide favorable habitat patches. The increased

growth in these nursery streams may also facilitate earlier sexual maturation, implying a potential

correlation between exploration and maturation traits.

2)   In this study, by sampling juveniles from two wild populations in the large Teno River catchment in

northernmost Fennoscandia, we tested for genetic association between exploration behavior to nursery

streams across four SNPs that spans a 70 kb long genomic region with a major effect on age at maturity

variation. Three of these SNPs are missense mutations in the *vgll3* and *akap11* genes, and one SNP tags a

putative regulatory region with the strongest association with the age at maturity trait.

3)    We show that the exploration behavior was linked to the genomic region in one of the two studied

populations. However, the genetic association was substantially stronger in the missense SNP located in

the *akap11* gene, which is farthest away from the *vgll3* SNPs and previously ruled out as being linked to

the age at maturity. We also detected a significant interaction between SNPs at the *vgll3* gene and

*akap11*, indicating a complex genetic architecture underlying the trait variation.

4)    Our results suggest that exploration and age at maturity are co-inherited within the same haplotype block,

but we find no evidence for direct causality. The observed genetic interaction may indicate that these two

traits form a co-adapted trait complex that may be instrumental in local adaptation processes.

**Keywords:** Correlated traits, exploration, habitat shift, age at maturity, epistasis, pleiotropy, Atlantic salmon

## Introduction

Elucidating the genetic architecture underlying adaptive traits is essential for understanding the mechanistic

basis of natural selection in the wild. Genome-wide scans have been essential for locating genomic regions

associated with phenotypic variation, or elucidating adaptive genetic differences within and among

populations. However, assessing the adaptive landscape at a finer-scale resolution is challenging, especially in

natural populations. A genomic region may harbor multiple loci associated with trait variation, and depending

on the number of loci and the linkage disequilibrium between them, the predicted response to selection will

vary. Close genomic proximity, or close genetic linkage distances between adaptive genetic loci can constrain

adaptation (Connallon & Hall 2018), but may also be selected for to preserve co-adapted allele combinations in

the form of epistatic interaction (Yeaman 2013; Bank 2022). For example, the evolution of inversions reduces

recombination along the inversion and allows gene complexes to be preserved over large genomic distances

(Kirkpatrick 2010; Küpper *et al.* 2015). Such an epistatic structure has been suggested as an important form of

genetic architecture that facilitates the co-evolution of co-adapted trait complexes, such as behavior and

reproduction or behavior and life history (Küpper *et al.* 2015; Tuttle *et al.* 2016; Funk *et al.* 2021). Likewise,

population models suggest such non-recombining regions may be preserved even without a need for co-

adaptation or epistasis (Kirkpatrick & Barton 2006; Hoffmann & Rieseberg 2008), suggesting their potential

prevalence in the natural populations.

The loci that are in close proximity have inherently low recombination rate. Hence, similar to long non-

recombining genomic regions, they could also harbor co-adapted gene complexes or independent adaptive

loci. However, the prevalence of such mechanisms is not well understood in natural populations. For example,

it is challenging to distinguish whether multiple traits are genetically associated with independent loci within

the region (spurious pleiotropy) or due to true pleiotropy (linked to one causal genic polymorphism, e.g,

Solovieff *et al.* 2013). Distinguishing these two modes of inheritance has implications. In the evolutionary

sense, pleiotropic loci are harder to decouple than tightly linked independent loci (Chebib & Guillaume 2021)

hence are less responsive to evolutionary pressures. Furthermore, multivariate phenotypes associated with

tightly linked loci might indicate co-adaptation (between traits) within local demes (Yeaman & Whitlock 2011),

while phenotypes under a pleiotropic effect of a gene may have a developmental basis (Wagner, Pavlicev &

Cheverud 2007). Therefore, linking genetic architecture to ecological processes and adaptation requires

information not only about the phenotypes associated with the genomic region but also about the fine-scale

genetic association between phenotypes and polymorphisms within genomic regions.

The *vgll3* locus in Atlantic salmon (*Salmo salar*) has strong potential as a genomic tool to better understand the

adaptive dynamics in the wild associated with a small linked genomic region. The *vgll3* genomic region, which

spans approximately 70 kb and encompasses *vgll3* and *akap11* genes, was initially identified as a major

genomic region explaining age at maturity, a life history trait with substantial fitness effects (Ayllon *et al.* 2015;

Barson *et al.* 2015). Further research identified that variation in other traits was also linked with the genetic

variation in the region, such as precocious male maturation (Debes *et al.* 2021), repeat-spawning (Aykanat *et*

*al.* 2019), metabolism (Prokkola *et al.* 2022a; Prokkola *et al.* 2024), foraging behavior (Aykanat et al. 2024),

exploration and aggression (Bangura et al. 2022; Bangura et al. 2024), morphology (Aykanat *et al.* 2025) and

early life fitness (Aykanat *et al.* 2024). The region harbors a large number of single nucleotide polymorphisms

(SNP), as revealed by whole genome sequencing (e.g., Barson *et al.* 2015), but four SNPs remains as the most

important candidates for phenotype association. These includes two potentially significant functional missense

polymorphisms within the coding regions in the *vgll3* gene ($vgll3_{Met54Thr}$ and $vgll3_{Asn323Lys}$), one missense

polymorphism in the *akap11* gene ($akap11_{Val214Met}$), and a putatively regulatory SNP with the strongest

association with maturation age, so called $vgll3_{TOP}$ (Barson *et al.* 2015). However, in most of these studies, the

trait-gene association was not refined further. For example, in the studies that involved cross-breeding designs,

parental fish usually was selected to have a homogeneous *vgll3* haplotype block with full linkage among these

four important SNPs, primarily to increase inference power of additive genetic relation. In one exceptional

case, cross-breeding was designed to segregate the genetic variation between *vgll3* and *akap11* genes, which

allowed further refinement of the location of causal SNPs within the genomic region (Sinclair-Waters *et al.*

2022). The genetic variation across tightly linked loci can be also refined in wild populations by exploiting the

natural segregation within the region. For example, Aykanat *et al.* (2019) attempted to narrow down the

genetic variation in the *vgll3* region associated with the repeat-spawning phenotype along four important SNPs

in the region, but none of the four SNPs explained the phenotypic variation better than the others, indicating

insufficiency of power. Taken together, the presence of multiple adaptive traits and putatively functionally

important SNPs linked to the genomic region signify the importance of better characterizing the genetic

architectures. This will provide insights into the adaptive dynamics of the genomic region within and among

Atlantic salmon populations.

Exploration in fishes is an important behavioral trait in that enables populations to spread and occupy new

habitat patches and resources. It can reduce the negative effects of density dependence (Einum, Sundt-Hansen

& H. Nislow 2006), and improve population stability (Baker *et al.* 2025). In juvenile Atlantic salmon, exploration

within freshwater (before emigration to the ocean) has the potential to contribute to improved feeding

opportunities, better growth, and ultimately higher survival (Erkinaro & Niemelä 1995; Erkinaro & Erkinaro

1998; Johansen, Elliott & Klemetsen 2005a; Johansen, Elliott & Klemetsen 2005b). Such habitat shifts may also

affect individual life history patterns (Erkinaro *et al.* 1997) via improved feeding and growth opportunities (e.g.,

Udea *et al.* 2025), making the individuals more likely to exceed the maturation threshold (Metcalfe 1998;

Thorpe *et al.* 1998; Jonsson & Jonsson 2014). At the genetic level, exploration and covarying traits such as

metabolism and aggression have been linked with the major loci affecting age-at-maturity, *vgll3* (Bangura *et al.*

2022; Prokkola *et al.* 2022a; Bangura *et al.* 2024; Prokkola *et al.* 2024). For example, using a common garden

design in an aquarium setting, Bangura *et al.* (2024) showed that juvenile Atlantic salmon with the late-

maturing *vgll3*[LL] genotype (L referring to the allele associated with late maturation) were significantly more

active in the 30-minute period following their release into a novel environment than individuals with the early-

maturing *vgll3*[EE] genotype (E referring to the allele associated with early maturation). This indicates that

exploration and maturation might be influenced by the same genetic locus. However, in their cross design, the

*vgll3* haplotype block was in complete linkage, and the study could not further refine the genetic basis within

the genomic region.

Traditionally, Atlantic salmon juveniles are considered relatively site-attached and sedentary in their early years

in freshwater (e.g., Jonsson & Jonsson 2011). However, in some river systems, older age groups of juvenile

salmon (parr) disperse to more favorable habitat patches within freshwater systems, including lacustrine areas

and smaller tributaries (Erkinaro 1995; Erkinaro, Julkunen & Niemela 1998; Halvorsen & Jørgensen 2006).

These small tributaries are nursery streams that are not suitable for salmon reproduction but may provide

favorable growth and survival conditions for salmon parr by means of increased food availability and decreased

competition and predation (Erkinaro & Niemelä 1995; Erkinaro & Erkinaro 1998; Erkinaro, Julkunen & Niemela

1998; Johansen, Elliott & Klemetsen 2005b). The extent of exploratory behavior to nursery streams and its

adaptive importance may be influenced by various factors, including environmental conditions, genetic

predispositions, and population density (e.g., Erkinaro *et al.* 1997; Johansen, Erkinaro & Amundsen 2010). For

example, the use of nursery streams by juvenile Atlantic salmon have been observed especially in northern

areas with depauperate fish fauna that exhibit less competition and predation threat to juvenile salmon

(Erkinaro 1995; Erkinaro & Gibson 1997; Shustov, Baryshev & Belyakova 2012).

Measuring the active migration of juvenile Atlantic salmon to nursery streams is a good proxy for quantifying

exploration behavior in natural settings (Erkinaro, Julkunen & Niemela 1998) and may help us to quantify

genetic basis of the exploration across populations. Furthermore, such migratory behavior may be linked to

maturation age during adulthood: The relative proportion of early maturation is shown to be higher among

individuals that occupy nursery streams during the juvenile stage (Erkinaro *et al.* 1997), which may be

facilitated by improved growth opportunities and the likelihood of reaching the maturation threshold (e.g.,

Thorpe *et al.* 1998). Thus, a genetic covariation between exploration and maturation timing may be

hypothesized as the basis of this observation.

Here, we quantified the genetic basis of exploration to nursery streams in two wild Atlantic salmon populations

in the Teno River system in northernmost Fennoscandia. We first asked whether the *vgll3* genomic region is

associated with explorative behaviors to nursery streams. By further exploring the genetic architecture within

the *vgll3* haploblock, we asked if exploration and age-at-maturity traits are pleiotropic or associated with

different genetic loci, and if epistasis within the region may play a role in shaping trait variation.

## Material and Methods

*Study area and sampling*

The Teno River (Deatnu in Sámi, Tana in Norwegian) catchment is located in far-north Europe (68–70°N, 25–

27°E), runs northwardly between Finland and Norway, and drains into the Barents Sea. The Teno system

supports a number of Atlantic salmon populations with wide diversity in life history characteristics and genetic

structure within and among populations (Vähä *et al.* 2017; Erkinaro *et al.* 2019). Juvenile salmon samples used

in this study were originally collected for a series of studies characterizing the exploration behaviors of juvenile

salmon from the main stem spawning areas into small nursery streams in relation to environmental conditions

and fish age (Erkinaro & Niemelä 1995; Erkinaro, Julkunen & Niemela 1998). Samples were collected from two

populations that reproduce in the Teno main stem and Inarijoki (Anárjohka in Sámi and Norwegian), a large

headwater tributary of the Teno River. These two populations exhibit a low level of genetic divergence ($F_{ST}$ =

0.012, Vähä *et al.* 2008; Aykanat *et al.* 2015), but have substantial differences in their life histories, with the

former population having high proportions of large, later-maturing individuals, and the latter population

primarily consisting of younger individuals at maturity (Czorlich *et al.* 2018, Erkinaro *et al.* 2019). In the

spawning rivers, the Teno main stem and Inarijoki, where adult Atlantic salmon reproduce and females lay

eggs, juvenile salmon were captured using 600- to 900-V pulsed DC electrofishing equipment. Scales were

sampled (initially for age determination, but later used for DNA extraction), and after sampling, fish were

released unharmed at the site of capture.

Sampling sites in both spawning rivers were located in close proximity (up to 100 meters) to the outlets of two

nursery streams, Baððá and Guoldnájohka, which drain into the Teno main stem and Inarijoki, respectively

(Erkinaro, Julkunen & Niemela 1998). Juvenile salmon in the streams were captured with a counting fence

installed at the lowest parts of the streams, 20–50 m from the outlets, and as above, scales were sampled.

After sampling, fish were released in the direction in which they had been heading (*cf.*, Erkinaro, Julkunen &

Niemela 1998). The migrant fish were classified as smolt or parr based on external characteristics, especially

the coloration, visibility of parr marks, and size (Erkinaro & Niemelä 1995). Overall, we sampled and genotyped

785 individuals, of which 635 were used in the main analysis after filtering out IDs with missing genotype

information in focal SNPs (see below) or missing cohort year information (i.e., due to poor-quality scale

samples; Supplementary Table 1).

*DNA extraction, SNP genotyping by targeted sequencing*

DNA was extracted from the scales of the juvenile Atlantic salmon using QIAamp 96DNA QIAcube HTKit

(Qiagen) following the manufacturer's protocol. Individuals were genotyped by targeted sequencing using a

SNP panel consisting of 173 SNP markers using a GTSeq approach (Campbell, Harmon & Narum 2015) as

outlined in (Aykanat *et al.* 2020). The SNP panel contained four SNP from the *vgl3* genomic regions including

the SNP that is most highly associated with age at maturity ($vgll3_{TOP}$), two missense mutations in the *vgll3* gene

($vgll3_{Met54Thr}$ and $vgll3_{Asn323Lys}$), and one missense mutation in the *akap11* gene ($akap11_{Val214Met}$, Barson *et al.*

2015). These SNPs are located on chromosome 25 and are 0.8 kb ($vgll3_{Met54Thr}$), 2.9 kb ($vgll3_{Asn323Lys}$), 10.9 kb

($vgll3_{TOP}$), and 65.5 kb ($akap11_{Val214Met}$) downstream from the transcription start site of the *vgll3* gene. The SNP

panel also included *six6* locus, a SNP highly spatially divergent between Atlantic salmon populations (Barson *et*

*al.* 2015), and so-called outlier and baseline SNPs. The outlier module includes 53 SNPs highly differentiated

between the Inarijoki and Teno main stem populations, allowing for a more powerful assignment of the

population of origin between these two closely related populations. The baseline module included 136

putatively neutral markers in low linkage disequilibrium, distributed throughout the whole genome and

previously filtered to have a minor allele frequency larger than 0.05 and heterozygosity larger than 0.2 for

these two populations (Aykanat *et al.* 2016). Individuals missing genotype information in any of these SNPs

were excluded from the analysis. Bonferroni correction was applied when multiple SNPs were compared.

*Statistical analysis*

We first tested if the exploration trait is co-localized in the *vgll3* genomic region by comparing single SNP

association to exploration behavior, which may also address if exploration and age at maturity are pleiotropic

or associated with different loci within the region. We modeled this separately for the Teno main stem and

Inarijoki populations using a generalized linear mixed model with a binomial error structure as follows:

$Y_{tenoMS} = \mu + SNP + \varepsilon_{error}$ for Teno main stem,

$Y_{inari} = \mu + SNP + \varepsilon_{hatchyear} + \varepsilon_{error \, for}$ Inarijoki,

where Y is a Boolean response variable for individuals' explorative behavior, in which individuals sampled in the

main stem were assumed to be sedentary, and coded as "0", and individuals sampled in the streams, either

descending or ascending, were assumed to be explorative and coded as "1". SNP is the genotype score of SNPs,

coded as 0 when individuals are homozygous for the reference alleles, 1 when homozygous for the alternative

alleles, and 0.5 when heterozygous. For the Inarijoki population, $\varepsilon_{hatchyear}$ is the random variance associated

with the hatch year calculated as the year sampled minus the age of the fish. Neither hatch year nor sampling

year overlapped for the Teno main stem populations between nursery streams and main stem collections, so

for this population, the model did not include the random term $\varepsilon_{hatchyear}$. The model was implemented for four

focal SNPs in the vgll3 genomic region (*vglll3*$_{TOP}$, *vgll3*$_{Met54Thr}$, *vgll3*$_{Asn323Lys}$), *akap1*$_{Val214Met}$), and 169 SNPs in the

SNP panel, including *six6*, which is shown to be associated with age at maturity and metabolism (Sinclair-

Waters *et al.* 2020; Prokkola *et al.* 2022b; Prokkola *et al.* 2024). Using the same model structure as above, we

then implemented a second model to quantify the effect of epistasis within the genomic region by modeling

pairwise interactions of SNPs within the *vgll3* genomic regions. Finally, we tested for potential gene-by-

environment effects by examining if results were consistent across the two populations, i.e., Inarijoki and Teno

main stem, which are in close proximity and genetically similar, but with substantial differences in their life

histories.

All models were run using the glmmTMB package, version 1.1.5, (Brooks *et al.* 2017) in R software version 4.4.1

(R Core Team 2019), using binomial error structure. Marginal means for genotype effects and their confidence

intervals were calculated using the emmeans package, version 1.8.5 (Lenth, 2023). Model diagnostics were

performed using the DHARMa package, version 0.4.6 (Hartig 2022).

## Results

Overall, the genotyping success rate was generally low (85%, Supplementary Figure 1), mostly due to the small

amount of material available for DNA extraction. After removing individuals with no genotype success in any of

the focal SNPs (N=143) and those with no scale aging data (N=15, of which eight also had no genotype success),

635 individuals remained in the final dataset out of an initial 785.

In the Teno main stem population, exploration behavior toward the nursery stream, Baðđá, was not

significantly associated with any of the four SNPs in the *vgll3* genomic region, or any other SNPs in the panel

after the Bonferroni correction for multiple comparisons (Figure 1a). In the Inarijoki population, the only SNP

that reached the Bonferroni-corrected threshold was *akap11*$_{\text{Val214Met}}$ (Figure 1b). Specifically, the

*akap11*$_{\text{Val214Met}}^{\text{EE}}$ genotype was 7.36 times (95% CI: 3.22 – 17.26) more likely to explore the nursery stream,

Guoldnájohka, than the *akap11*$_{\text{Val214Met}}^{\text{LL}}$ allele ($p < 0.001$, Figure 2a). The *six6* loci did not exhibit a significant

association in any of the populations.

**Figure 1:** SNP - exploration association across 173 SNPs in Teno main stem (a) and Inarijoki (b) populations in Atlantic salmon. Focal SNPs are colored in red. Uncorrected (p = 0.05) and Bonferroni corrected p-value thresholds are indicated with dashed and continuous lines.

When pairwise interactions within the *vgll3* genomic region were modeled in the Teno main stem population, none of the models were more significant than the null model that did not contain genetic effects. For the Inarijoki population, the most parsimonious model explaining the exploration included the interaction of *akap11*$_{Val214Met}$ and *vgll3*$_{Asn323Lys}$, which was superior to the null model ($\Delta$AIC = 16.36), or the model that only included *akap11*$_{Val214Met}$ ($\Delta$AIC = 4.47), but not significantly better than the interaction between *akap11*$_{Val214Met}$ and two other SNPs in the *vgll3* genomic region ($\Delta$AIC < 2, Supplementary Figure 3). The effect was driven by a substantial decrease in exploration behavior associated with the genotype combination of *akap11*$_{Val214Met}^{EL}$ and

230   *vgll3*Asn323Lys$^{EL}$ (Figure 2b). As such, the model predicts that the odds of exploration for individuals with

231   *akap11*Val214Met$^{EL}$ and *vgll3*Asn323Lys$^{EL}$ were 4.13 (95% CI 2.24-7.84, *p*<0.001) and 3.10 (95% CI 1.34-7.15, *p*=0.041)

232   times less than for *akap11*Val214Met$^{EE}$ and *vgll3*Asn323Lys$^{EL}$ and *akap11*Val214Met$^{EL}$ and *vgll3*Asn323Lys$^{LL}$, respectively (Figure

233   2b)

**Figure 2: a)** Odds of exploration in Inarijoki Atlantic salmon as a function of *akap11*Val214Met SNP, and **b)** as a
function of *akap11*Val214Met and *vgll3*Asn323Lys interaction. Error bars indicate 95% CI of the marginal estimates, and
first and second numbers in parenthesis indicates the sample sizes in the nursery stream and the main stream,
respectively. Y axis is drawn in log scale.

The fish sampled in Guoldnájohka contained 83 individuals that had undergone smoltification (or were in the

process of becoming smolt), which is a suite of physiological, morphological, and behavioral changes in Atlantic

salmon juveniles for becoming ready for seaward migration (McCormick & Saunders 1987), which may have

altered their exploration behavior. However, when the smolts were excluded from the analysis, the results

were mostly qualitatively similar (Supplementary Figure 4). One notable difference was that a model with the

*akap11*Val214Met and *vgll3*Asn323Lys interaction was only marginally more significant than a model with

independent, additive effects of these SNPs (*p* = 0.085, ΔAIC = 0.93, Supplementary Figure 4). However, the

change in significance was not meaningfully different from an effect that might have resulted from a reduced

sample size in a random manner. As such, when 83 parr were randomly sampled and excluded from the

dataset (i.e., the same number as the smolts in the dataset) or 83 randomly selected parrs in the dataset were

used as a replacement for excluded smolts, the ΔAIC range of these models (each permuted 1000 times) was

similar to the ΔAIC values obtained in the original model (Supplementary Figure 5).

## Discussion

Factors shaping the genetic architecture of ecologically important traits are fundamentally important for

understanding adaptive dynamics in the wild. We have demonstrated that exploration behavior in juvenile

Atlantic salmon is linked to a genomic region on chromosome 25 in one of the two populations in the Teno

River. This region contains *vgll3* and *akap11* genes and is associated with a suite of traits involved in salmon

adaptation (reviewed in Mobley *et al.* 2021) including age at maturity (Barson *et al.* 2015). We have found that

a missense SNP in the *akap11* gene ($akap11_{Val214Met}$) was substantially linked to exploratory behavior, while

SNPs in the *vgll3* gene exhibited epistasis with this SNP, suggesting a complex genetic architecture in the

region. Furthermore, the genetic effect was only observed in one of the two studied populations, indicating the

importance of environment or population-specific factors in triggering gene-dependent exploration behavior.

Our results showed that the early maturation allele in the *akap11* gene was associated with increased

exploration to nursery streams in the Inarijoki population. This is consistent with the findings of Erkinaro *et al.*

(1997) who, by analyzing freshwater growth patterns of salmon, suggested that juveniles entering nursery

streams had a greater tendency to mature and return from their sea journey earlier than those that stayed in

the main stem for their juvenile years. Differences in growth opportunity in Atlantic salmon juveniles, which is

linked to efficient acquisition of resources, may be instrumental in exceeding the maturation threshold in

subsequent years (Metcalfe 1998; Thorpe *et al.* 1998; Jonsson & Jonsson 2014). As such, causality between

exploration and migrating to nursery streams may be suggested based on the increased likelihood of

individuals in the streams to reach the maturation threshold (e.g., Thorpe *et al.* 1998) upon increased growth

opportunities. On the other hand, the association was predominantly driven by genetic variation in the

*akap11*$_{Val214Met}$ SNP, which is about 70 kb downstream of the *vgll3* gene and exhibits a weaker association with

sea age at maturity (Barson *et al.* 2015). The *akap11*$_{Val214Met}$ SNP was also previously ruled out as the genetic

basis of early male maturation in Atlantic salmon in controlled settings (Sinclair-Waters *et al.* 2022). Combined,

while exploration behavior may facilitate earlier maturation in a context-dependent manner, our results do not

provide evidence to support a direct genetic causation between the two, but that a small region explains both

maturation and exploration behaviors, and that the two traits are genetically correlated by linkage.

A multitude of phenotypic traits may be linked within a genomic region via pleiotropy or linkage disequilibrium

(i.e., spurious pleiotropy, Solovieff *et al.* 2013). This makes it challenging to identify the phenotypic traits under

selection, i.e., by using genomic scans (Paaby & Rockman 2013; Solovieff *et al.* 2013). Our results indicate that

the *vgll3* genomic region contains multiple independent SNPs linked to adaptive traits, ruling out pleiotropy as

the basis of multi-trait association. The presence of multiple adaptive traits within a tightly linked genomic

region raises the question of whether the genetic co-variation between these loci harbors co-adapted trait

combinations. Linked architectures may facilitate adaptation by enabling co-inheritance of adapted gene

complexes (Mullon, Keller & Lehmann 2017; Oomen, Kuparinen & Hutchings 2020). For example, genomic

inversions have been shown to be important in maintaining co-adapted trait combinations, including migratory

strategies and dispersal (Mullon, Keller & Lehmann 2017; Wellenreuther & Bernatchez 2018; Rubenstein *et al.*

2019). Similarly, genetic variation in life-history migratory ecotypes in rainbow trout (*Oncorhynchus mykiss*,

Pearse *et al.* 2014) and Atlantic cod (*Gadus morhua*) have been shown to be associated with genetic markers in

a large inverted region (Kirubakaran *et al.* 2016). A tight linkage between adaptive traits might also be selected

for when locally adapted populations may have varying local fitness optima (for these traits), and exhibit high

gene flow between them (Yeaman & Whitlock 2011). In this case, tight linkage between locally adapted traits

would help in purging maladaptive genetic variants with a relatively reduced genetic load.

In a practical sense, our results also highlight the insufficiency of drawing conclusions when only one SNP is

used as a tag marker to represent the genetic makeup of tightly linked genomic regions. In common garden

cross-breeding designs, one SNP marker may be utilized to tag the whole region when exploring the trait-

genotype association of the region (e.g., Prokkola *et al.* 2022a; Bangura *et al.* 2024). While such designs provide

substantial statistical and cost-reduction benefits, they would not be able to discern associations across SNPs in

close proximity, and their interactions. For example, using a common garden setup, Bangura *et al.* (2024),

suggested *vgll3* is likely to play a role in exploration behaviors. However, in the cross-breeding design, parents

were chosen to be in complete linkage regarding the genetic variation in *akap11* and *vgll3* SNPs, thereby

deduction of SNP effects independently is not possible (see also, Bangura *et al.* 2022). In fact, when Sinclair-

Waters *et al.* (2022) refined the SNP association in the vgll3 genomic region, in relation to their effect on male

parr maturation, they found an opposite effect of SNPs in the *vgll3* and *akap11* gene, whereby the early

maturation allele in the *akap11* gene was indeed marginally associated with reduced maturation, further

signifying the importance of refining genetic associations when multiple candidate SNPs may underlie the

phenotypic variation.

Our study provides a further outlook on the dynamics of behavioral syndromes and correlated behaviors

(Höjesjö, Johnsson & Bohlin 2004; Garamszegi, Markó & Herczeg 2013; Mathot, Dingemanse & Nakagawa

2019; Axling *et al.* 2023). The classic paradigm suggests that dispersal is negatively correlated with dominance

and aggression, whereby dominant individuals monopolize favorable habitat patches, and force subordinate

individuals to displace in search of new habitats (Stamps 1994; Stamps 1999). However, in Atlantic salmon,

migration to nursery streams might be carried out by individuals holding higher positions in the dominance

hierarchy (Erkinaro & Niemelä 1995). Using seasonal growth in scale depositions, Erkinaro and Niemelä (1995)

estimated growth in previous seasons -a trait correlated with dominance and aggression in early life stages

(Cutts *et al.* 2005)-, and showed that faster growing individuals are more likely to migrate to the small nursery

streams. A similar tendency in individual growth patterns has also been identified in individuals shifting their

juvenile rearing habitat to lacustrine areas of the catchment (Erkinaro *et al.* 1998). A niche shift, i.e., by

migrating to nursery streams, may be hypothesized as a mechanism to overcome potential resource mismatch

in the original habitat. Therefore, we would predict that the EE allele in the *vgll3* genomic region is likely

associated with behavioral traits correlated with exploration in salmon, such as dominance.

On the other hand, common garden experiments in controlled conditions do not support our prediction. For

example, by quantifying novel environment exploration in a confined space (an aquarium with a 25 cm x 30 cm

surface area), Bangura *et al.* (2024) found that the *vgll3* L allele was associated with higher exploration in a

novel environment. Likewise, using a similar experimental setup Bangura *et al.* (2022) also found increased

aggression associated with the *vgll3* L allele. These results suggest a genetic covariation exists between

aggression and novel environmental exploration, but with an opposite direction of allelic effect to our study.

(Note that in the crosses used by Bangura *et al.* (2022) and Bangura *et al.* (2024) *akap11* and *vgll3* were in full

linkage, hence conclusions related to the genetic variation in the *vgll3* SNPs are equally valid for the *akap11*

SNP.) These contrasting results may reflect contextual differences between natural and common garden

settings. As such, a meta-analysis has shown that these two personality traits (aggression and exploration) only

correlate when there is a contextual overlap that underlies the similarity of trait expression (Garamszegi,

Markó & Herczeg 2013). Therefore, the contrasting conclusions in these two settings (natural vs. laboratory)

may stem from different functional physiological processes. Alternatively, strong environment (common

garden vs. natural setting) by genotype, or population by genotype interactions may explain the contrast. For

the latter, for example, the parental fish used in Bangura *et al.* (2022) and Bangura *et al.* (2024) are obtained

from a hatchery broodstock of salmon managed by the Natural Resources Institute Finland (LUKE), initially

originated from a Baltic Sea population (River Neva). Hence differences in the genetic background of fish may

also explain the discrepancies. In fact, the genetic association between SNPs in the *vgll3* genomic region and

exploration was significant only in the Inarijoki population, while no detectable effect was found in the Teno

main stem population, further emphasizing a potential role of population specific differences in the genetic

basis of trait variation. Our study illustrates the complex dynamics of correlated behaviors of Atlantic salmon

and their genetic background.

Finally, we demonstrated an epistasis between *akap11* and the *vgll3* gene. While the statistical power was

insufficient to which of the SNP in the *vgll3* gene are the most likely candidates, the presence of epistasis

suggests a complex genetic architecture within the genomic region. Epistasis contributes to the trait variance,

and help maintain additive genetic variation upon selection (Cheverud & Routman 1995). Further, the trait

complex between exploration and age at maturity may contribute to adaptation dynamics. For example,

optimal trait combinations (of traits co-inherited within a haplotype block) may be locally variable among

populations and may improve overall fitness compared to what would be predicted by only additive gene

action. Such an epistatic interaction would also facilitate purging maladapted genetic combinations (Yeaman &

Whitlock 2011).

In this study, we used historical field data that was not initially intended for a genetic study such as this one.

Hence, the design has some drawbacks. For example, the hatching years of individuals collected in the Teno

main stem and Baððá stream do not match, therefore we cannot rule out the possibility that the (lack of)

difference in genetic response is due to temporal changes in allele frequencies, even though the time span

between sampling periods is only a few years. Secondly, genotype calling failed for a substantial portion of

individuals due to the suboptimal amount of tissue remains in the storage envelopes. In addition, we employed

a targeted genotyping approach by scoring only the most likely candidate SNPs in the genomic regions. An

alternative approach to sequence the whole genomic region might provide finer resolution in trait association,

and help to better narrow down the causal region.

Movement patterns in the Salmonidae family are mostly studied in the context of the phenology of migration,

which is central to their life histories, has a heritable component, and strong fitness consequences (Niemelä *et*

*al.* 2006; Stewart, Middlemas & Youngson 2006). Many species exhibit a strong dichotomy of the trait within

and among populations, as evidenced by the presence of both resident and migratory forms in, e.g., brown

trout, rainbow/steelhead trout, and Arctic char (Jonsson & Jonsson 1993). Movement within the freshwater

habitat at the juvenile stage and its association with migration at later stages is much less explored. Our study

demonstrates a complex genetic interaction between exploration in the freshwater and age at maturity. This

may help to maintain co-adapted trait complexes (exploration and maturation) and may be instrumental in

local adaptation processes, fitness, and the long-term survival of populations.

## Acknowledgement

373 We thank Susanna Ukonaho and Meri Lindquist from the University of Turku for help with DNA extraction,

374 Matti Kylmäaho from Luke for scale aging, and Lars Paulin from the University of Helsinki for coordinating the

375 genotyping-by-sequencing assays. Funding came from the Research Council of Finland (353388 and 325964 to

376 T.A.). We thank to Craig Primmer for his communication during the making of the study.

## Conflict of Interest

379 The authors declare no competing interests.

## References

Axling, J., Vossen, L.E., Peterson, E. & Winberg, S. (2023) Boldness, activity, and aggression: Insights from a
large-scale study in Baltic salmon (Salmo salar L). *PLoS One,* **18,** e0287836.
10.1371/journal.pone.0287836.
Aykanat, T., Debes, P.V., Jansouz, S., Gueguen, L., House, A.H., Ruokolainen, A., Erkinaro, J., Pritchard, V.L.,
Primmer, C.R. & Bolstad, G.H. (2025) Large effect life-history genomic regions are associated with
functional morphological traits in Atlantic salmon. *G3 (Bethesda)*.

Aykanat, T., Johnston, S.E., Orell, P., Niemela, E., Erkinaro, J. & Primmer, C.R. (2015) Low but significant genetic differentiation underlies biologically meaningful phenotypic divergence in a large Atlantic salmon population. *Molecular Ecology,* **24,** 5158-5174. 10.1111/mec.13383.

Aykanat, T., Lindqvist, M., Pritchard, V.L. & Primmer, C.R. (2016) From population genomics to conservation and management: a workflow for targeted analysis of markers identified using genome-wide approaches in Atlantic salmon Salmo salar. *Journal of Fish Biology,* **89,** 2658-2679. 10.1111/jfb.13149.

Aykanat, T., McLennan, D., Metcalfe, N.B. & Prokkola, J.M. (2024) Early survival in Atlantic salmon is associated with parental genotypes at loci linked to timing of maturation. *Evolution,* **78,** 1441-1452. 10.1093/evolut/qpae072.

Aykanat, T., Ozerov, M., Vaha, J.P., Orell, P., Niemela, E., Erkinaro, J. & Primmer, C.R. (2019) Co-inheritance of sea age at maturity and iteroparity in the Atlantic salmon vgll3 genomic region. *Journal of Evolutionary Biology,* **32,** 343-355. 10.1111/jeb.13418.

Aykanat, T., Rasmussen, M., Ozerov, M., Niemela, E., Paulin, L., Vaha, J.P., Hindar, K., Wennevik, V., Pedersen, T., Svenning, M.A. & Primmer, C.R. (2020) Life-history genomic regions explain differences in Atlantic salmon marine diet specialization. *Journal of Animal Ecology,* **89,** 2677-2691. 10.1111/1365-2656.13324.

Ayllon, F., Kjaerner-Semb, E., Furmanek, T., Wennevik, V., Solberg, M.F., Dahle, G., Taranger, G.L., Glover, K.A., Almen, M.S., Rubin, C.J., Edvardsen, R.B. & Wargelius, A. (2015) The vgll3 Locus Controls Age at Maturity in Wild and Domesticated Atlantic Salmon (Salmo salar L.) Males. *PLoS Genetics,* **11,** e1005628. 10.1371/journal.pgen.1005628.

Baker, H.K., Obedzinski, M., Grantham, T.E. & Carlson, S.M. (2025) Variation in Salmon Migration Phenology Bolsters Population Stability but Is Threatened by Drought. *Ecology Letters,* **28,** e70081. 10.1111/ele.70081.

Bangura, P.B., Tiira, K., Aykanat, T., Niemela, P.T., Erkinaro, J., Liljestrom, P., Toikkanen, A. & Primmer, C.R. (2024) Sex-specific associations of the maturation locus vgll3 with exploratory behavior and boldness in Atlantic salmon juveniles. *Ecol Evol,* **14,** e11449. 10.1002/ece3.11449.

Bangura, P.B., Tiira, K., Niemela, P.T., Erkinaro, J., Liljestrom, P., Toikkanen, A. & Primmer, C.R. (2022) Linking vgll3 genotype and aggressive behaviour in juvenile Atlantic salmon (Salmo salar). *Journal of Fish Biology,* **100,** 1264-1271. 10.1111/jfb.15040.

Bank, C. (2022) Epistasis and Adaptation on Fitness Landscapes. *Annual Review of Ecology, Evolution, and Systematics,* **53,** 457-479. 10.1146/annurev-ecolsys-102320-112153.

Barson, N.J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G.H., Fiske, P., Jacq, C., Jensen, A.J., Johnston, S.E., Karlsson, S., Kent, M., Moen, T., Niemela, E., Nome, T., Naesje, T.F., Orell, P., Romakkaniemi, A., Saegrov, H., Urdal, K., Erkinaro, J., Lien, S. & Primmer, C.R. (2015) Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature,* **528,** 405-408. 10.1038/nature16062.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., Skaug, H.J., Machler, M. & Bolker, B.M. (2017) glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *R Journal,* **9,** 378-400. Doi 10.32614/Rj-2017-066.

Campbell, N.R., Harmon, S.A. & Narum, S.R. (2015) Genotyping-in-Thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. *Mol Ecol Resour,* **15,** 855-867. 10.1111/1755-0998.12357.

Chebib, J. & Guillaume, F. (2021) Pleiotropy or linkage? Their relative contributions to the genetic correlation of quantitative traits and detection by multitrait GWA studies. *Genetics,* **219**. 10.1093/genetics/iyab159.

Cheverud, J.M. & Routman, E.J. (1995) Epistasis and its contribution to genetic variance components. *Genetics,* **139,** 1455-1461. 10.1093/genetics/139.3.1455.

Connallon, T. & Hall, M.D. (2018) Genetic constraints on adaptation: a theoretical primer for the genomics era. *Annals of the New York Academy of Sciences,* **1422,** 65-87. 10.1111/nyas.13536.

Cutts, C.J., Brembs, B., Metcalfe, N.B. & Taylor, A.C. (2005) Prior residence, territory quality and life-history
strategies in juvenile Atlantic salmon (Salmo salar L.). *Journal of Fish Biology,* **55,** 784-794.
10.1111/j.1095-8649.1999.tb00717.x.
Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P. & Primmer, C.R. (2018) Rapid sex-specific evolution of age at
maturity is shaped by genetic architecture in Atlantic salmon. *Nat Ecol Evol*, **2**, 1800-1807.
10.1038/s41559-018-0681-5.
Debes, P.V., Piavchenko, N., Ruokolainen, A., Ovaskainen, O., Moustakas-Verho, J.E., Parre, N., Aykanat, T.,
Erkinaro, J. & Primmer, C.R. (2021) Polygenic and major-locus contributions to sexual maturation
timing in Atlantic salmon. *Molecular Ecology,* **30,** 4505-4519. 10.1111/mec.16062.
Einum, S., Sundt-Hansen, L. & H. Nislow, K. (2006) The partitioning of density-dependent dispersal, growth and
survival throughout ontogeny in a highly fecund organism. *Oikos,* **113,** 489-496. 10.1111/j.2006.0030-
1299.14806.x.
Erkinaro, H. & Erkinaro, J. (1998) Feeding of Atlantic salmon,Salmo salarL., parr in the subarctic River Teno and
three tributaries in northernmost Finland. *Ecology of Freshwater Fish,* **7,** 13-24. 10.1111/j.1600-
0633.1998.tb00168.x.
Erkinaro, J. (1995) The age structure and distribution of Atlantic salmon parr, Salmo salar L., in small tributaries
and main stems of the subarctic River Teno, northern Finland. *Ecology of Freshwater Fish,* **4,** 53-61.
10.1111/j.1600-0633.1995.tb00117.x.
Erkinaro, J., Czorlich, Y., Orell, P., Kuusela, J., Falkegård, M., Länsman, M., Pulkkinen, H., Primmer, C.R. &
Niemelä, E. (2019) Life history variation across four decades in a diverse population complex of Atlantic
salmon in a large subarctic river. *Canadian Journal of Fisheries and Aquatic Sciences,* **76,** 42-55.
10.1139/cjfas-2017-0343.
Erkinaro, J., Dempson, J.B., Julkunen, M. & Niemelä, E. (1997) Importance of ontogenetic habitat shifts to
juvenile output and life history of Atlantic salmon in a large subarctic river: an approach based on
analysis of scale characteristics. *Journal of Fish Biology,* **51,** 1174-1185. DOI 10.1111/j.1095-
8649.1997.tb01134.x.
Erkinaro, J. & Gibson, R.J. (1997) Interhabitat migration of juvenile Atlantic salmon in a Newfoundland river
system, Canada. *Journal of Fish Biology,* **51,** 373-388. 10.1111/j.1095-8649.1997.tb01673.x.
Erkinaro, J., Julkunen, M. & Niemela, E. (1998) Migration of juvenile Atlantic salmon Salmo salar in small
tributaries of the subarctic River Teno, northern Finland. *Aquaculture,* **168,** 105-119. Doi
10.1016/S0044-8486(98)00343-3.
Erkinaro, J. & Niemelä, E. (1995) Growth Differences between the Atlantic Salmon Parr, Salmo-Salar, of Nursery
Brooks and Natal Rivers in the River Teno Watercourse in Northern Finland. *Environmental Biology of
Fishes,* **42,** 277-287. Doi 10.1007/Bf00004921.
Erkinaro, J., Niemelä, E., Saari, A., Shustov, Y. & Jørgensen, L. (1998) Timing of habitat shift by Atlantic salmon
parr from fluvial to lacustrine habitat: analysis of age distribution, growth, and scale characteristics.
*Canadian Journal of Fisheries and Aquatic Sciences,* **55,** 2266-2273. 10.1139/f98-107.
Funk, E.R., Mason, N.A., Palsson, S., Albrecht, T., Johnson, J.A. & Taylor, S.A. (2021) A supergene underlies
linked variation in color and morphology in a Holarctic songbird. *Nat Commun,* **12,** 6833.
10.1038/s41467-021-27173-z.
Garamszegi, L.Z., Markó, G. & Herczeg, G. (2013) A meta-analysis of correlated behaviors with implications for
behavioral syndromes: relationships between particular behavioral traits. *Behavioral Ecology,* **24,** 1068-
1080. 10.1093/beheco/art033.
Halvorsen, M. & Jørgensen, L. (2006) Lake-use by juvenile Atlantic salmon (Salmo salarL.) and other salmonids
in northern Norway. *Ecology of Freshwater Fish,* **5,** 28-36. 10.1111/j.1600-0633.1996.tb00034.x.
Hartig F (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. R
package version 0.4.6.

Höjesjö, J., Johnsson, J.r. & Bohlin, T. (2004) Habitat complexity reduces the growth of aggressive and dominant
brown trout (Salmo trutta) relative to subordinates. *Behavioral Ecology and Sociobiology,* **56**.
10.1007/s00265-004-0784-7.
Hoffmann, A.A. & Rieseberg, L.H. (2008) Revisiting the Impact of Inversions in Evolution: From Population
Genetic Markers to Drivers of Adaptive Shifts and Speciation? *Annu Rev Ecol Evol Syst,* **39,** 21-42.
10.1146/annurev.ecolsys.39.110707.173532.
Johansen, M., Elliott, J.M. & Klemetsen, A. (2005a) A comparative study of juvenile salmon density in 20
streams throughout a very large river system in northern Norway. *Ecology of Freshwater Fish,* **14,** 96-
110. 10.1111/j.1600-0633.2004.00082.x.
Johansen, M., Elliott, J.M. & Klemetsen, A. (2005b) Relationships between juvenile salmon,Salmo salarL., and
invertebrate densities in the River Tana, Norway. *Ecology of Freshwater Fish,* **14,** 331-343.
10.1111/j.1600-0633.2005.00107.x.
Johansen, M., Erkinaro, J. & Amundsen, P.-A. (2010) The When, What and Where of Freshwater Feeding.
*Atlantic Salmon Ecology*, pp. 89-114. Wiley-Blackwell.
Jonsson, B. & Jonsson, N. (1993) Partial migration: niche shift versus sexual maturation in fishes. *Reviews in Fish*
*Biology and Fisheries,* **3,** 348-365. 10.1007/bf00043384.
Jonsson, B. & Jonsson, N. (2011) Ecology of Atlantic salmon and brown trout : habitat as a template for life
histories. Springer, Dordrecht.
Jonsson, B. & Jonsson, N. (2014) Early environment influences later performance in fishes. *Journal of Fish*
*Biology,* **85,** 151-188. 10.1111/jfb.12432.
Kirkpatrick, M. (2010) How and why chromosome inversions evolve. *PLoS Biology,* **8**.
10.1371/journal.pbio.1000501.
Kirkpatrick, M. & Barton, N. (2006) Chromosome inversions, local adaptation and speciation. *Genetics,* **173,**
419-434. 10.1534/genetics.105.047985.
Kirubakaran, T.G., Grove, H., Kent, M.P., Sandve, S.R., Baranski, M., Nome, T., De Rosa, M.C., Righino, B.,
Johansen, T., Ottera, H., Sonesson, A., Lien, S. & Andersen, O. (2016) Two adjacent inversions maintain
genomic differentiation between migratory and stationary ecotypes of Atlantic cod. *Molecular Ecology,*
**25,** 2130-2143. 10.1111/mec.13592.
Küpper, C., Stocks, M., Risse, J.E., dos Remedios, N., Farrell, L.L., McRae, S.B., Morgan, T.C., Karlionova, N.,
Pinchuk, P., Verkuil, Y.I., Kitaysky, A.S., Wingfield, J.C., Piersma, T., Zeng, K., Slate, J., Blaxter, M., Lank,
D.B. & Burke, T. (2015) A supergene determines highly divergent male reproductive morphs in the ruff.
*Nature Genetics,* **48,** 79-83. 10.1038/ng.3443.
Lenth R (2023). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.8.5.
Li, M.H. & Merila, J. (2011) Population differences in levels of linkage disequilibrium in the wild. *Molecular*
*Ecology,* **20,** 2916-2928. 10.1111/j.1365-294X.2011.05154.x.
Lillehammer A (1974) Studie av laks og ørretyngelens forflytning over elvebunnen og evertebratfaunaen i
temporært oversvømte områder. (Summary: Studies on the movement of salmon and trout fry and
invertebrates into flooded areas). *Fauna* 27: 69-73.
Mathot, K.J., Dingemanse, N.J. & Nakagawa, S. (2019) The covariance between metabolic rate and behaviour
varies across behaviours and thermal types: meta-analytic insights. *Biological Reviews of the*
*Cambridge Philosophical Society,* **94,** 1056-1074. 10.1111/brv.12491.
McCormick, S.D. & Saunders, R.L. (1987) Preparatory Physiological Adaptations for Marine Life of Salmonids:
Osmoregulation, Growth, and Metabolism.
Metcalfe, N.B. (1998) The interaction between behavior and physiology in determining life history patterns in
Atlantic salmon (Salmo salar). *Canadian Journal of Fisheries and Aquatic Sciences,* **55,** 93-103.
10.1139/d98-005.
Mobley, K.B., Aykanat, T., Czorlich, Y., House, A., Kurko, J., Miettinen, A., Moustakas-Verho, J., Salgado, A.,
Sinclair-Waters, M., Verta, J.-P. & Primmer, C.R. (2021) Maturation in Atlantic salmon (Salmo salar,

Salmonidae): a synthesis of ecological, genetic, and molecular processes. *Reviews in Fish Biology and*
*Fisheries,* **31,** 523-571. 10.1007/s11160-021-09656-w.
Mullon, C., Keller, L. & Lehmann, L. (2017) Social polymorphism is favoured by the co-evolution of dispersal
with social behaviour. *Nature Ecology & Evolution,* **2,** 132-140. 10.1038/s41559-017-0397-y.
Niemelä, E., Orell, P., Erkinaro, J., Dempson, J.B., BrØrs, S., Svenning, M.A. & Hassinen, E. (2006) Previously
spawned Atlantic salmon ascend a large subarctic river earlier than their maiden counterparts. *Journal*
*of Fish Biology,* **69,** 1151-1163. 10.1111/j.1095-8649.2006.01190.x.
Oomen, R.A., Kuparinen, A. & Hutchings, J.A. (2020) Consequences of Single-Locus and Tightly Linked Genomic
Architectures for Evolutionary Responses to Environmental Change. *Journal of Heredity,* **111,** 319-332.
10.1093/jhered/esaa020.
Paaby, A.B. & Rockman, M.V. (2013) The many faces of pleiotropy. *Trends in Genetics,* **29,** 66-73.
10.1016/j.tig.2012.10.010.
Pearse, D.E., Miller, M.R., Abadia-Cardoso, A. & Garza, J.C. (2014) Rapid parallel evolution of standing variation
in a single, complex, genomic region is associated with life history in steelhead/rainbow trout. *Proc Biol*
*Sci,* **281,** 20140012. 10.1098/rspb.2014.0012.
Prokkola, J.M., Asheim, E.R., Morozov, S., Bangura, P., Erkinaro, J., Ruokolainen, A., Primmer, C.R. & Aykanat, T.
(2022a) Genetic coupling of life-history and aerobic performance in Atlantic salmon. *Proc Biol Sci,* **289,**
20212500. 10.1098/rspb.2021.2500.
Prokkola, J.M., Chew, K.K., Anttila, K., Maamela, K.S., Yildiz, A., Asheim, E.R., Primmer, C.R. & Aykanat, T. (2024)
Tissue-specific metabolic enzyme levels covary with whole-animal metabolic rates and life-history loci
via epistatic effects. *Philos Trans R Soc Lond B Biol Sci,* **379,** 20220482. 10.1098/rstb.2022.0482.
Prokkola, J.M., Wagner, A., Koskinen, E., Debes, P.V., House, A., Åsheim, E.R., Primmer, C.R., Pirinen, E. &
Aykanat, T. (2022b) Adipose tissue mitochondrial respiration in Atlantic salmon: implications for sex-
dependent life-history variation. 10.1101/2022.07.25.501389.
R Core Team (2019) R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria.
Rubenstein, D.R., Ågren, J.A., Carbone, L., Elde, N.C., Hoekstra, H.E., Kapheim, K.M., Keller, L., Moreau, C.S.,
Toth, A.L., Yeaman, S. & Hofmann, H.A. (2019) Coevolution of Genome Architecture and Social
Behavior. *Trends in Ecology & Evolution,* **34,** 844-855. 10.1016/j.tree.2019.04.011.
Shustov, Y.A., Baryshev, I.A. & Belyakova, E.E. (2012) Specific features of the feeding of juvenile atlantic salmon
(Salmo salar L.) in the subarctic varzuga river and its small tributaries (Kola Peninsula). *Inland Water*
*Biology,* **5,** 288-292. 10.1134/s1995082912030121.
Sinclair-Waters, M., Odegard, J., Korsvoll, S.A., Moen, T., Lien, S., Primmer, C.R. & Barson, N.J. (2020) Beyond
large-effect loci: large-scale GWAS reveals a mixed large-effect and polygenic architecture for age at
maturity of Atlantic salmon. *Genetics Selection Evolution,* **52,** 9. 10.1186/s12711-020-0529-8.
Sinclair-Waters, M., Piavchenko, N., Ruokolainen, A., Aykanat, T., Erkinaro, J. & Primmer, C.R. (2022) Refining
the genomic location of single nucleotide polymorphism variation affecting Atlantic salmon maturation
timing at a key large-effect locus. *Molecular Ecology,* **31,** 562-570. 10.1111/mec.16256.
Solovieff, N., Cotsapas, C., Lee, P.H., Purcell, S.M. & Smoller, J.W. (2013) Pleiotropy in complex traits: challenges
and strategies. *Nature Reviews: Genetics,* **14,** 483-495. 10.1038/nrg3461.
Stamps, J. (1994) Territorial Behavior: Testing the Assumptions. pp. 173-232.
Stamps, J.A. (1999) Dominance and Territoriality. *Proc. 22 Int. Ornithol. Congr.,* (eds N.J. Adams & R.H. Slotow),
pp. 1592-1605. Durban.
Stewart, D.C., Middlemas, S.J. & Youngson, A.F. (2006) Population structuring in Atlantic salmon (Salmo salar):
evidence of genetic influence on the timing of smolt migration in sub-catchment stocks. *Ecology of*
*Freshwater Fish,* **15,** 552-558. 10.1111/j.1600-0633.2006.00197.x.

Thorpe, J.E., Mangel, M., Metcalfe, N.B. & Huntingford, F.A. (1998) Modelling the proximate basis of salmonid
577           life-history variation, with application to Atlantic salmon, Salmo salar L. *Evolutionary Ecology,* **12,** 581-
578           599. Doi 10.1023/A:1022351814644.
579     Tuttle, E.M., Bergland, A.O., Korody, M.L., Brewer, M.S., Newhouse, D.J., Minx, P., Stager, M., Betuel, A.,
580           Cheviron, Z.A., Warren, W.C., Gonser, R.A. & Balakrishnan, C.N. (2016) Divergence and Functional
581           Degradation of a Sex Chromosome-like Supergene. *Current Biology,* **26,** 344-350.
582           10.1016/j.cub.2015.11.069.
583     Ueda, R., Kanaiwa, M., Terui, A., Takimoto, G. & Sato, T. (2025) Seasonal timing of ecosystem linkage mediates
584           life-history variation in a salmonid fish population. *Ecology*, **106**, e70114. 10.1002/ecy.70114.
585     Vähä, J.-P., Erkinaro, J., Falkegård, M., Orell, P. & Niemelä, E. (2017) Genetic stock identification of Atlantic
586           salmon and its evaluation in a large population complex. *Canadian Journal of Fisheries and Aquatic
587           Sciences,* **74,** 327-338. 10.1139/cjfas-2015-0606.
588     Vähä, J.P., Erkinaro, J., Niemelä, E. & Primmer, C.R. (2008) Temporally stable genetic structure and low
589           migration in an Atlantic salmon population complex: implications for conservation and management.
590           *Evolutionary Applications,* **1,** 137-154. DOI 10.1111/j.1752-4571.2007.00007.x.
591     Wagner, G.P., Pavlicev, M. & Cheverud, J.M. (2007) The road to modularity. *Nature Reviews: Genetics,* **8,** 921-
592           931. 10.1038/nrg2267.
593     Wellenreuther, M. & Bernatchez, L. (2018) Eco-Evolutionary Genomics of Chromosomal Inversions. *Trends in
594           Ecology & Evolution,* **33,** 427-440. 10.1016/j.tree.2018.04.002.
595     Yeaman, S. (2013) Genomic rearrangements and the evolution of clusters of locally adaptive loci. *Proc Natl
596           Acad Sci U S A,* **110,** E1743-1751. 10.1073/pnas.1219381110.
597     Yeaman, S. & Whitlock, M.C. (2011) The genetic architecture of adaptation under migration-selection balance.
598           *Evolution,* **65,** 1897-1911. 10.1111/j.1558-5646.2011.01269.x.

599